






29,815/0

N.I. 0  
12





Digitized by the Internet Archive  
in 2017 with funding from  
Wellcome Library

[https://archive.org/details/b28771485\\_0001](https://archive.org/details/b28771485_0001)





















---

A  
MATHEMATICAL AND PHILOSOPHICAL  
DICTIONARY.

V O L. I.

---





A  
MATHEMATICAL AND PHILOSOPHICAL  
DICTIONARY:

CONTAINING  
*AN EXPLANATION OF THE TERMS, AND AN ACCOUNT OF THE SEVERAL SUBJECTS,*  
COMPRIZED UNDER THE HEADS  
MATHEMATICS, ASTRONOMY, AND PHILOSOPHY  
BOTH NATURAL AND EXPERIMENTAL:

WITH AN  
HISTORICAL ACCOUNT OF THE RISE, PROGRESS, AND PRESENT STATE OF THESE SCIENCES:

ALSO  
MEMOIRS OF THE LIVES AND WRITINGS OF THE MOST EMINENT AUTHORS,  
BOTH ANCIENT AND MODERN,

*WHO BY THEIR DISCOVERIES OR IMPROVEMENTS HAVE CONTRIBUTED TO THE ADVANCEMENT OF THEM.*

IN TWO VOLUMES.

WITH MANY CUTS AND COPPER-PLATES.

---

---

BY CHARLES HUTTON, LL.D.

F. R. S. OF LONDON AND EDINBURGH, AND OF THE PHILOSOPHICAL SOCIETIES OF HAARLEM AND AMERICA;  
AND PROFESSOR OF MATHEMATICS IN THE ROYAL MILITARY ACADEMY, WOOLWICH.

---

---

VOL. I.

LONDON:

PRINTED BY J. DAVIS,  
FOR J. JOHNSON, IN ST. PAUL'S CHURCH-YARD; AND G. G. AND J. ROBINSON,  
IN PATERNOSTER-ROW.

M.DCC.XCVI.





*[Faint, illegible text, likely bleed-through from the reverse side of the page.]*

*[Faint, illegible text, likely bleed-through from the reverse side of the page.]*

*[Faint, illegible text, likely bleed-through from the reverse side of the page.]*

---

# P R E F A C E.

---

AMONG the Dictionaries of Arts and Sciences which have been published, of late years, in various parts of Europe, it is matter of surprise that Philosophy and Mathematics should have been so far overlooked as not to be thought worthy of a separate Treatise, in this form. These Sciences constitute a large portion of the present stock of human knowledge, and have been usually considered as possessing a degree of importance to which few others are entitled; and yet we have hitherto had no distinct Lexicon, in which their constituent parts and technical terms have been explained, with that amplitude and precision, which the great improvements of the Moderns, as well as the rising dignity of the Subject, seem to demand.

THE only works of this kind in the English language, deserving of notice, are Harris's *Lexicon Technicum*, and Stone's *Mathematical Dictionary*; the former of which, though a valuable performance at the time it was written, is now become too dry and obsolete to be referred to with pleasure or satisfaction: and the latter, consisting only of one volume in 8vo, must be regarded merely as an unfinished sketch, or brief compendium, extremely limited in its plan, and necessarily deficient in useful information.

It became, therefore, the only resource of the Reader, in many cases where explanation was wanted, to have recourse to Chambers's Dictionary, in four large Volumes folio, or to the *Encyclopædia Britannica*, now in eighteen large volumes 4to, or the still more stupendous performance of the French Encyclopedists; and even here his expectations might be frequently disappointed. These great and useful works, aiming at a general comprehension of the whole circle of the Sciences, are sometimes very deficient in their descriptions of particular branches; it being almost impossible, in such extensive



extensive undertakings, to appreciate, with exactness, the due value of every article : They are, besides, so voluminous and heterogeneous in their nature, as to render a frequent reference to them extremely inconvenient ; and even if this were not the case, their high price puts them out of the reach of the generality of readers.

WITH a view to obviate these defects, the Public are here presented with a Dictionary of a moderate size and price, which is devoted solely to Philosophical and Mathematical subjects. It is a work for which materials have been collecting through a course of many years ; and is the result of great labour and reading. Not only most of the Encyclopedias already extant, and the various publications of the Learned Societies throughout Europe, have been carefully consulted, but also all the original works, of any reputation, which have hitherto appeared upon these subjects, from the earliest writers down to the present times.

FROM the latter of these sources, in particular, a considerable portion of information has been obtained, which the curious reader will find, in many cases, to be highly interesting and important. The History of Algebra, for instance, which is detailed at considerable length in the First Volume, under the head of that Article, will afford sufficient evidence to shew in what a superficial and partial way the inquiry has been hitherto investigated, even by professed writers on the subject ; the principal of whom are M. Montucla, our countryman the celebrated Dr. Wallis, and the Abbé De Gua, a late French author, who has pretended to correct the Doctor's errors and misrepresentations.

REGULAR historical details are in like manner given of the origin and progress of each of these Sciences, as well as of the inventions and improvements by which they have been gradually brought from their first rude beginnings to their present advanced state.

It is also to be observed, that besides the articles common to the generality of Dictionaries of this kind, an interesting Biographical Account is here introduced of the most celebrated Philosophers and Mathematicians, both ancient and modern ; among which will be found the Lives of many eminent characters, who have hitherto been either wholly overlooked, or very imperfectly recorded. Complete lists of their works are also subjoined to each Article, where they could be procured ; which cannot but prove highly acceptable to that class of readers, who are desirous of obtaining the most satisfactory information upon the subjects of their particular enquiries and pursuits. On the head of Biography however the Author has still to lament the want of many other respectable names which he was desirous to have added



to his list of authors, not having been able to procure any circumstantial accounts of their lives. He could have wished to have comprised in his list, the lives of all such public literary characters as the University Professors of Astronomy, Philosophy and Mathematics, as well as those of the other more respectable class of Authors on those Sciences. He will therefore thankfully receive the communication of any such memoirs from the hands of gentlemen possessed of them; as well as hints and information on such useful improvements in the sciences as may have been overlooked in this Dictionary, or any articles that may here have been imperfectly or incorrectly treated; that he may at some future time, by adding them to this work, render it still more complete and deserving the public notice.

As this work is an attempt to separate the words in the sciences of Astronomy, Mathematics, and Philosophy, from those of other arts or sciences, in several of which there are already separate Dictionaries; as in Chemistry, Geography, Music, Marine and Naval affairs, &c; words sometimes occurred which it was rather doubtful whether they could be considered as properly belonging to the present work or not; in which case many of such words have been here inserted. But such as appeared clearly and peculiarly to belong to any of those other subjects, have been either wholly omitted, or else have had a very short account only given of them. The readers of this work therefore, recollecting that it is not a General Dictionary of all the Arts and Sciences, will not expect to find all sorts of words and subjects here treated of; but such only as peculiarly appertain to the proper matter of the work. And therefore, although some few words may inadvertently have been omitted; yet when the Reader does not immediately find every word which he wishes to consult, he will not always consider them as omissions of the Author, but for the most part as relating to some other science foreign to this Dictionary.

In all cases where it could be conveniently done, the necessary figures and diagrams are inserted in the same page with the subjects which they are designed to elucidate; a method which will be found much more commodious than that of putting them in separate plates at the end of each volume, but, which has added very considerably to the expence of the undertaking: where the subjects are of such a nature that they could not be otherwise well represented, they are engraved on Copperplates.

As the whole of this work was written before it was put to the press, the Reader will find it of an equal and uniform nature and construction throughout; in which respect many publications of this kind are very defective, from the subjects being diffusely treated under the first letters of the alphabet, while articles  
of

of equal importance in the latter part are so much abridged as to be rendered almost useless, in order that the whole might be comprised in a limited number of sheets, according to proposals made before the works were composed. The present Dictionary having been completed without any of these unfavourable circumstances, will be found in most cases equally instructive and useful, and may be consulted with no less advantage by the Man of Science than the Student.



---

A

PHILOSOPHICAL and MATHEMATICAL

DICTIONARY.

---

A.

A B A

A B A

**A**BACIST, an Arithmetician. In this sense we find the word used by William of Malmesbury, in his *History de Gestis Anglorum*, written about the year 1150; where he shews that one Gerbert, a learned monk of France, who was afterwards made pope of Rome in the year 998 or 999, by the name of Silvester the 2d, was the first who got from the Saracens the abacus, and that he taught such rules concerning it, as the Abacists themselves could hardly understand.

ABACUS, in *Arithmetic*, an ancient instrument used by most nations for casting up accounts, or performing arithmetical calculations: it is by some derived from the Greek *αβαξ*, which signifies a cupboard or beaufet, perhaps from the similarity of the form of this instrument; and by others it is derived from the Phœnician *abak*, which signifies dust or powder, because it was said that this instrument was sometimes made of a square board or tablet, which was powdered over with fine sand or dust, in which were traced the figures or characters used in making calculations, which could thence be easily defaced, and the abacus refitted for use. But Lucas Pacioli, in the first part of his second distinction, thinks it is a corruption of Arabicus, by which he meant their Algorithm, or the method of numeral computation received from them.

We find this instrument for computation in use, under some variations, with most nations, as the Greeks, Romans, Germans, French, Chinese, &c.

The Grecian abacus was an oblong frame, over which were stretched several brass wires, strung with little ivory balls, like the beads of a necklace; by the various arrangements of which all kinds of computa-

tions were easily made. Mahudel, in *Hist. Acad. R. Inscr.* t. 3. p. 390.

The Roman Abacus was a little varied from the Grecian, having pins sliding in grooves, instead of strings or wires and beads. *Philos. Trans.* No. 180.

The Chinese Abacus, or Shwan-pan, like the Grecian, consists of several series of beads strung on brass wires, stretched from the top to the bottom of the instrument, and divided in the middle by a cross piece from side to side. In the upper space every string has two beads, which are each counted for 5; and in the lower space every string has five beads, of different values, the first being counted as 1, the second as 10, the third as 100, and so on, as with us. See SHWAN-PAN.


The Abacus chiefly used in European countries, is nearly upon the same principles, though the use of it is here more limited, because of the arbitrary and unequal divisions of money, weights, and measures, which, in China, are all divided in a tenfold proportion, like our scale of common numbers. This is made by drawing any number of parallel lines, like paper ruled for music, at such a distance as may be at least equal to twice the diameter of a calculus, or counter. Then the value of these lines, and of the spaces between them, increases, from the lowest to the highest, in a tenfold proportion. Thus, counters placed upon the first line, signify so many units or ones; on the second line 10's, on the third line 100's, on the fourth line 1000's, and so on: in like manner a counter placed in the first space, between the first and second line, denotes 5, in the second space 50, in the third space 500, in the fourth space 5000,

B

and



and so on. So that there are never more than four counters placed on any line, nor more than one placed in any space, this being of the same value as five counters on the next line below. So the counters on the Abacus, in the figure here below, express the number or sum 47382.



Besides the above instruments of computation, there have been several others invented by different persons; as *Napier's rods* or *bones*, described in his *Rabdologia*, which see under the word *NAPIER*; also the *Abacus Rhabdologicus*, a variation of *Napier's*, which is described in the first vol. of *Machines et Inventions approuvées par l'Academie Royale des Sciences*. An ingenious and general one was also invented by Mr. Gamaliel Smethurst, and is described in the *Philosophical Transactions*, vol. 46; where the inventor remarks that computations by it are much quicker and easier than by the pen, are less burthensome to the memory, and can be performed by blind persons, or in the dark as well as in the light. A very comprehensive instrument of this kind was also contrived by the late learned Dr. Nicholas Saunderson, by which he performed very intricate calculations: an account of it is prefixed to the first volume of his *Algebra*, and it is there by the editor called *Palpable Arithmetic*: which see.

**ABACUS, Pythagorean**, so denominated from its inventor, Pythagoras; a table of numbers, contrived for readily learning the principles of arithmetic; and was probably what we now call the multiplication-table.

**ABACUS, or ABACISCUS**, in *Architecture*, the upper part or member of the capital of a column; serving as a crowning both to the capital and to the whole column. Vitruvius informs us that the *Abacus* was originally intended to represent a square flat tile laid over an urn, or a basket; and the invention is ascribed to Calimachus, an ingenious statuary of Athens, who, it is said, adopted it on observing a small basket, covered with a tile, over the root of an *Acanthus* plant, which grew on the grave of a young lady; the plant shooting up, encompassed the basket all around, till meeting with the tile, it curled back in the form of scrolls: Calimachus passing by, took the hint, and immediately executed a capital on this plan; representing the tile by the *Abacus*, the leaves of the *acanthus* by the volutes or scrolls, and the basket by the vase or body of the capital. See *ACANTHUS*.

*Abacus* is also used by Scamozzi for a concave moulding in the capital of the Tuscan pedestal. And the word is used by Palladio for other members which he describes. Also, in the ancient architecture, the same term is used to denote certain compartments in the incrustation or lining of the walls of state-rooms, mosaic-pavements, and the like. There were *Abaci* of marble,

porphyry, jasper, alabaster, and even glass; variously shaped, as square, triangular, and such-like.

**ABACUS Logisticus** is a right angled triangle, whose sides, about the right angle, contain all the numbers from 1 to 60; and its area the products of each two of the opposite numbers. This is also called a *canon of sexagesimals*, and is no other than a multiplication-table carried to 60 both ways.

**ABACUS & Palmula**, in the Ancient Music, denote the machinery by which the strings of the polyplectra, or instruments of many strings, were struck, with a plectrum made of quills.

**ABACUS Harmonicus** is used by Kircher for the structure and disposition of the keys of a musical instrument, either to be touched with the hands or feet.

**ABACUS**, in *Geometry*, a table or slate upon which schemes or diagrams are drawn.

**ABAS**, a weight used in Persia for weighing pearls; and is an eighth part lighter than the European carat.

**ABASSI**, a silver coin current in Persia, deriving its name from Schaw Abbas II. King of Persia, and is worth near eighteen pence English money.

**ABATIS, or ABATTIS**, from the French *abattre*, to throw down, or beat down, in the Military Art, denotes a kind of retrenchment made by a quantity of whole trees cut down, and laid lengthways beside each other, the closer the better, having all their branches pointed towards the enemy, which prevents his approach, at the same time that the trunks serve as a breast-work before the men. The *Abattis* is a very useful work on most occasions, especially on sudden emergencies, when trees are near at hand; and has always been practised with considerable success, by the ablest commanders in all ages and nations.

**ABBREVIATE**; to abbreviate fractions in arithmetic and algebra, is to lessen proportionally their terms, or the numerator and denominator; which is performed by dividing those terms by any number or quantity, which will divide them without leaving a remainder. And when the terms cannot be any farther so divided, the fraction is said to be in its least terms.

$$\text{So } \frac{16}{24} = \frac{8}{12} = \frac{4}{6} = \frac{2}{3},$$

by dividing the terms continually by 2.

$$\text{And } \frac{294}{504} = \frac{147}{252} = \frac{49}{84} = \frac{7}{12},$$

by dividing by 2, 3, and 7.

$$\text{Also } \frac{3 \times 8 \times 5}{6} = \frac{8 \times 5}{2} = 4 \times 5 = 20,$$

by dividing by 3 and by 2.

$$\text{And } \frac{12abx^2}{4acx} = \frac{3bx}{c}, \text{ by dividing by } 4ax.$$

$$\text{And } \frac{ab^2 + b^2x}{ax + x^2} = \frac{b^2}{x}, \text{ by dividing by } a + x.$$


**ABBREVIATION**, of fractions, in Arithmetic and Algebra, the reducing them to lower terms.

**ABERRATION**, in *Astronomy*, an apparent motion of the celestial bodies, occasioned by the progressive motion of light, and the earth's annual motion in her orbit.

This effect may be explained and familiarized by the motion



motion of a line parallel to itself, much after the manner that the composition and resolution of forces are explained. If light have a progressive motion, let the



proportion of its velocity to that of the earth in her orbit, be as the line  $BC$  to the line  $AC$ ; then, by the composition of these two motions, the particle of light will seem to describe the line  $BA$  or  $DC$ , instead of its real course  $BC$ ; and will appear in the direction  $AB$  or  $CD$ , instead of its true direction  $CB$ . So that if  $AB$  represent a tube, carried with a parallel motion by an observer along the line  $AC$ , in the time that a particle of light would move over the space  $BC$ ,

the different places of the tube being  $AB, ab, cd, CD$ ; and when the eye, or end of the tube, is at  $A$ , let a particle of light enter the other end at  $B$ ; then when the tube is at  $ab$ , the particle of light will be at  $e$ , exactly in the axis of the tube; and when the tube is at  $cd$ , the particle of light will arrive at  $f$ , still in the axis of the tube; and lastly, when the tube arrives at  $CD$ , the particle of light will arrive at the eye or point  $C$ , and consequently will appear to come in the direction  $DC$  of the tube, instead of the true direction  $BC$ . And so on, one particle succeeding another, and forming a continued stream or ray of light in the apparent direction  $DC$ . So that the apparent angle made by the ray of light with the line  $AE$ , is the angle  $DCE$ , instead of the true angle  $BCE$ ; and the difference,  $BCD$  or  $ABC$ , is the quantity of the aberration.

M. de Maupertuis, in his *Elements of Geography*, gives also a familiar and ingenious idea of the aberration, in this manner: "It is thus," says he, "concerning the direction in which a gun must be pointed to strike a bird in its flight; instead of pointing it straight to the bird, the fowler will point a little before it, in the path of its flight, and that so much the more as the flight of the bird is more rapid, with respect to the flight of the shot." In this way of considering the matter, the flight of the bird represents the motion of the earth, or the line  $AC$ , in our scheme above, and the flight of the shot represents the motion of the ray of light, or the line  $BC$ .

Mr. Clairaut too, in the *Memoires of the Academy of Sciences* for the year 1746, illustrates this effect in a familiar way, by supposing drops of rain to fall rapidly and quickly after each other from a cloud, under which a person moves with a very narrow tube; in which case it is evident that the tube must have a certain inclination, in order that a drop which enters at the top, may fall freely through the axis of the tube, without touching the sides of it; which inclination must be more or less according to the velocity of the drops in respect to that of the tube: then the angle made by the direction of the tube and of the falling drops, is the aberration arising from the combination of those two motions.

This discovery, which is one of the brightest that have been made in the present age, we owe to the accuracy and ingenuity of the late Dr. Bradley, Astronomer Royal; to which he was occasionally led by the result

of some accurate observations which he had made with another view, namely, to determine the annual parallax of the fixed stars, or that which arises from the motion of the earth in its annual orbit about the sun.

The annual motion of the earth about the sun had been much doubted, and warmly contested. The defenders of that motion, among other proofs of the reality of it, conceived the idea of adducing an incontestable one from the annual parallax of the fixed stars, if the stars should be within such a distance, or if instruments and observations could be made with such accuracy, as to render that parallax sensible. And with this view various attempts have been made. Before the observations of M. Picard, made in 1672, it was the general opinion, that the stars did not change their position during the course of a year. Tycho Brahe and Ricciolus fancied that they had assured themselves of it from their observations; and from thence they concluded that the earth did not move round the sun, and that there was no annual parallax in the fixed stars. M. Picard, in the account of his *Voyage d'Uranibourg*, made in 1672, says that the pole star, at different times of the year, has certain variations which he had observed for about 10 years, and which amounted to about  $40''$  a year: from whence some who favoured the annual motion of the earth were led to conclude that these variations were the effect of the parallax of the earth's orbit. But it was impossible to explain it by that parallax; because this motion was in a manner contrary to what ought to follow only from the motion of the earth in her orbit.

In 1674 Dr. Hook published an account of observations which he said he had made in 1669, and by which he had found that the star  $\gamma$  Draconis was  $23'$  more northerly in July than in October: observations which, for the present, seemed to favour the opinion of the earth's motion, although it be now known that there could not be any truth or accuracy in them.

Flamsteed having observed the pole star with his mural quadrant, in 1689 and the following years, found that its declination was  $40''$  less in July than in December; which observations, although very just, were yet however improper for proving the annual parallax; and he recommended the making of an instrument of 15 or 20 feet radius, to be firmly fixed on a strong foundation, for deciding a doubt which was otherwise not soon likely to be brought to a conclusion.

In this state of uncertainty and doubt, then, Dr. Bradley, in conjunction with Mr. Samuel Molineux, in the year 1725, formed the project of verifying, by a series of new observations, those which Dr. Hook had communicated to the public almost 50 years before. And as it was his attempt that chiefly gave rise to this, so it was his method in making the observations, in some measure, that they followed; for they made choice of the same star, and their instrument was constructed upon nearly the same principles: but had it not greatly exceeded the former in exactness, they might still have continued in great uncertainty as to the parallax of the fixed stars. And this was chiefly owing to the accuracy of the ingenious Mr. George Graham, to whom the lovers of astronomy are also indebted for several other exact and convenient instruments.



The success then of the intended experiment, evidently depending very much on the accuracy of the instrument, that leading object was first to be well secured. Mr. Molineux's apparatus then having been completed, and fitted for observing, about the end of November 1725, on the third day of December following, the bright star in the head of Draco, marked  $\gamma$  by Bayer, was for the first time observed, as it passed near the zenith, and its situation carefully taken with the instrument. The like observations were made on the fifth, eleventh, and twelfth days of the same month; and there appearing no material difference in the place of the star, a farther repetition of them, at that season, seemed needless, it being a time of the year in which no sensible alteration of parallax, in this star, could soon be expected. It was therefore curiosity that chiefly urged Dr. Bradley, being then at Kew, where the instrument was fixed, to prepare for observing the star again on the 17th of the same month; when, having adjusted the instrument as usual, he perceived that it passed a little more southerly this day than it had done before. Not suspecting any other cause of this appearance, they first concluded that it was owing to the uncertainty of the observations, and that either this, or the foregoing, was not so exact as they had before supposed. For which reason they proposed to repeat the observation again, to determine from what cause this difference might proceed: and upon doing it, on the 20th of December, the doctor found that the star passed still more southerly than at the preceding observation. This sensible alteration surprised them the more, as it was the contrary way from what it would have been, had it proceeded from an annual parallax of the star. But being now pretty well satisfied, that it could not be entirely owing to the want of exactness in the observations, and having no notion of any thing else that could cause such an apparent motion as this in the star; they began to suspect that some change in the materials, or fabric of the instrument itself, might have occasioned it. Under these uncertainties they remained for some time; but being at length fully convinced, by several trials, of the great exactness of the instrument; and finding, by the gradual increase of the star's distance from the pole, that there must be some regular cause that produced it; they took care to examine very nicely, at the time of each observation, how much the variation was; till about the beginning of March 1726, the star was found to be 20'' more southerly than at the time of the first observation: it now indeed seemed to have arrived at its utmost limit southward, as in several trials, made about this time, no sensible difference was observed in its situation. By the middle of April it appeared to be returning back again towards the north; and about the beginning of June, it passed at the same distance from the zenith, as it had done in December, when it was first observed.

From the quick alteration in the declination of the star about this time, increasing about one second in three days, it was conjectured that it would now proceed northward, as it had before gone southward, of its present situation; and it happened accordingly; for the star continued to move northward till September following, when it became stationary again; being then near 20'' more northerly than in June, and upwards of

39'' more northerly than it had been in March. From September the star again returned towards the south, till, in December, it arrived at the same situation in which it had been observed twelve months before, allowing for the difference of declination on account of the precession of the equinox.

This was a sufficient proof that the instrument had not been the cause of this apparent motion of the star; and yet it seemed difficult to devise one that should be adequate to such an unusual effect. A nutation of the earth's axis was one of the first things that offered itself on this occasion; but it was soon found to be insufficient; for though it might have accounted for the change of declination in  $\gamma$  Draconis, yet it would not at the same time accord with the phenomena observed in the other stars, particularly in a small one almost opposite in right ascension to  $\gamma$  Draconis, and at about the same distance from the north pole of the equator: for though this star seemed to move the same way, as a nutation of the earth's axis would have made it; yet changing its declination but about half as much as  $\gamma$  Draconis in the same time, as appeared on comparing the observations of both made on the same days, at different seasons of the year, this plainly proved that the apparent motion of the star was not occasioned by a real nutation; since, had that been the case, the alteration in both stars would have been nearly equal.

The great regularity of the observations left no room to doubt, but that there was some uniform cause by which this unexpected motion was produced, and which did not depend on the uncertainty or variety of the seasons of the year. Upon comparing the observations with each other, it was discovered that, in both the stars above mentioned, the apparent difference of declination from the *maxima*, was always nearly proportional to the versed sine of the sun's distance from the equinoctial points. This was an inducement to think that the cause, whatever it was, had some relation to the sun's situation with respect to those points. But not being able to frame any hypothesis, sufficient to account for all the phenomena, and being very desirous to search a little farther into this matter, Dr. Bradley began to think of erecting an instrument for himself at Wanstead; that, having it always at hand, he might with the more ease and certainty enquire into the laws of this new motion. The consideration likewise of being able, by another instrument, to confirm the truth of the observations hitherto made with that of Mr. Molineux, was no small inducement to the undertaking; but the chief of all was, the opportunity he should thereby have of trying in what manner other stars should be affected by the same cause, whatever it might be. For Mr. Molineux's instrument being originally designed for observing  $\gamma$  Draconis, to try whether it had any sensible parallax, it was so contrived, as to be capable of but little alteration in its direction; not above seven or eight minutes of a degree: and there being but few stars, within half that distance from the zenith of Kew, bright enough to be well observed, he could not, with his instrument, thoroughly examine how this cause affected stars that were differently situated, with respect to the equinoctial and solstitial points of the ecliptic.

These considerations determined him; and by the contrivance and direction of the same ingenious person, Mr.



Mr. Graham, his instrument was fixed up the 19th of August 1727. As he had no convenient place where he could make use of so long a telescope as Mr. Molineux's, he contented himself with one of but little more than half the length, namely of 12 feet and a half, the other being 24 feet and a half long, judging from the experience he had already had, that this radius would be long enough to adjust the instrument to a sufficient degree of exactness: and he had no reason afterwards to change his opinion; for by all his trials he was very well satisfied, that when it was carefully rectified, its situation might be securely depended on to half a second. As the place where his instrument was hung, in some measure determined its radius; so did it also the length of the arc or limb, on which the divisions were made, to adjust it: for the arc could not conveniently be extended farther, than to reach to about  $6\frac{1}{2}$  degrees on each side of his zenith. This however was sufficient, as it gave him an opportunity of making choice of several stars, very different both in magnitude and situation: there being more than two hundred, inserted in the British Catalogue, that might be observed with it. He needed not indeed to have extended the limb so far, but that he was willing to take in *Capella*, the only star of the first magnitude that came so near his zenith.

His instrument being fixed, he immediately began to observe such stars as he judged most proper to give him any light into the cause of the motion already mentioned. There was a sufficient variety of small ones, and not less than twelve that he could observe through all seasons of the year, as they were bright enough to be seen in the day-time, when nearest the sun. He had not been long observing, before he perceived that the notion they had before entertained, that the stars were farthest north and south when the sun was near the equinoxes, was only true of those stars which are near the solstitial colure. And after continuing his observations a few months, he discovered what he then apprehended to be a general law observed by all the stars, namely, that each of them became stationary, or was farthest north or south, when it passed over his zenith at six of the clock, either in the evening or morning. He perceived also that whatever situation the stars were in, with respect to the cardinal points of the ecliptic, the apparent motion of every one of them tended the same way, when they passed his instrument about the same hour of the day or night; for they all moved southward when they passed in the day, and northward when in the night; so that each of them was farthest north, when it came in the evening about six of the clock, and farthest south when it came about six in the morning.

Though he afterwards discovered that the maxima, in most of these stars, do not happen exactly when they pass at those hours; yet, not being able at that time to prove the contrary, and supposing that they did, he endeavoured to find out what proportion the greatest alterations of declination, in different stars, bore to each other; it being very evident that they did not all change their declination equally. It has been before noticed, that it appeared from Mr. Molineux's observations, that  $\gamma$  *Draconis* changed its declination above twice as much as the before-mentioned small star that was nearly op-

posite to it; but examining the matter more nicely, he found that the greatest change in the declination of these stars, was as the sine of the latitude of each star respectively. This led him to suspect that there might be the like proportion between the *maxima* of other stars; but finding that the observations of some of them would not perfectly correspond with such an hypothesis, and not knowing whether the small difference he met with might not be owing to the uncertainty and error of the observations, he deferred the farther examination into the truth of this hypothesis, till he should be farther furnished with a series of observations made in all parts of the year; which would enable him not only to determine what errors the observations might be liable to, or how far they might safely be depended on; but also to judge, whether there had been any sensible change in the parts of the instrument itself.

When the year was completed, he began to examine and compare his observations; and having pretty well satisfied himself as to the general laws of the phenomena, he then endeavoured to find out the cause of them. He was already convinced that the apparent motion of the stars was not owing to a nutation of the earth's axis. The next that occurred to him, was an alteration in the direction of the plumb-line, by which the instrument was constantly adjusted; but this, upon trial, proved insufficient. Then he considered what refraction might do; but here also he met with no satisfaction. At last, through an amazing sagacity, he conjectured that all the phenomena hitherto mentioned, proceeded from the progressive motion of light, and the earth's annual motion in her orbit: for he perceived, that if light were propagated in time, the apparent place of a fixed object would not be the same when the eye is at rest, as when it is moving in any other direction but that of the line passing through the object and the eye; and that when the eye is moving in different directions, the apparent place of the object would be different.

He considered this matter in the following manner. He imagined *CA* to be a ray of light, falling perpendicularly upon the line *BD*: then, if the eye be at rest at *A*, the object must appear in the direction *AC*, whether light be propagated in time, or in an instant. But if the eye be moving from *B* towards *A*, and light be propagated in time, with a velocity that is to the velocity of the eye, as *AC* to *AB*; then, light moving from *C* to *A*, whilst the eye moves from *B* to *A*, that particle of it by which the object will be discerned, when the eye in its motion comes to *A*, is at *C* when the eye is at *B*. Joining the points *B*, *C*, he supposed the line *BC* to be a tube, inclined to the line *BD* in the angle *DBC*, and of such a diameter as to admit of but one particle of light: then it was easy to conceive, that the particle of light at *C*, by which the object must be seen when the eye arrives at *A*, would pass through the tube *BC*, so inclined to the line *BD*, and accompanying the eye in its motion from *B* to *A*; and that it would not come to





to the eye, placed behind such a tube, if it had any other inclination to the line B D. If, instead of supposing B C so small a tube, we conceive it to be the axis of a larger; then, for the same reason, the particle of light at C cannot pass through that axis, unless it be inclined to B D in the same angle D B C.

In the like manner, if the eye move the contrary way, from D towards A, with the same velocity; then the tube must be inclined in the angle B D C. Although therefore the true or real place of an object, be perpendicular to the line in which the eye is moving, yet the visible place will not be so; since that must doubtless be in the direction of the tube. But the difference between the true and apparent place, will be, *ceteris paribus*, greater or less, according to the different proportions between the velocity of light and that of the eye: so that if we could suppose light to be propagated in an instant, then there would be no difference between the real and visible place of an object, although the eye were in motion; for in that case, A C being infinite with respect to A B, the angle A C B, which is the difference between the true and visible place, vanishes. But if light be propagated in time, which was then allowed by most philosophers, then it is evident from the foregoing considerations, that there will always be a difference between the true and visible place of an object, except when the eye is moving either directly towards, or from the object. And in all cases, the sine of the difference between the true and visible place of the object, will be to the sine of the visible inclination of the object to the line in which the eye is moving, as the velocity of the eye, is to the velocity of light.

If light moved only 1000 times faster than the eye, and an object, supposed to be at an infinite distance, were really placed perpendicularly over the plane in which the eye is moving; it follows, from what has been said, that the apparent place of such object will always be inclined to that plane, in an angle of  $89^{\circ} 56\frac{1}{2}'$ ; so that it will constantly appear  $3\frac{1}{2}'$  from its true place, and will seem so much less inclined to the plane, that way towards which the eye tends. That is, if A C be to A B or A D, as 1000 to 1, the angle A B C will be  $89^{\circ} 56\frac{1}{2}'$ , and the angle A C B  $3\frac{1}{2}'$ , and B C D or 2 A C B will be  $7'$ , if the direction of the motion of the eye be contrary at one time to what it is at another.

If the earth revolve about the sun annually, and the velocity of light were to the velocity of the earth's motion in its orbit, as 1000 is to 1; then it is easy to conceive, that a star really placed in the pole of the ecliptic, would to an eye carried along with the earth, seem to change its place continually; and, neglecting the small difference on account of the earth's diurnal revolution on its axis, it would seem to describe a circle about that pole, every where distant from it by  $3\frac{1}{2}'$ . So that its longitude would be varied through all the points of the ecliptic every year, but its latitude would always remain the same. Its right ascension would also change, and its declination, according to the different situation of the sun in respect of the equinoctial points; and its apparent distance from the north pole of the equator, would be  $7'$  less at the autumnal, than at the vernal equinox.

The greatest alteration of the place of a star, in the pole of the ecliptic, or, which in effect amounts to the same, the proportion between the velocity of light and the earth's motion in its orbit, being known, it will not be difficult to find what would be the difference, on this account, between the true and apparent place of any other star at any time; and, on the contrary, the difference between the true and apparent place being given, the proportion between the velocity of light, and the earth's motion in her orbit, may be found.

After the history of this curious discovery, related by the author nearly in the terms above, he gives the results of a multitude of accurate observations, made on a great number of stars, at all seasons of the year. From all which observations, and the theory as related above, he found that every star, in consequence of the earth's motion in her orbit and the progressive motion of light, appears to describe a small ellipse in the heavens, the transverse axis of which is equal to the same quantity for every star, namely  $40''$  nearly; and that the conjugate axis of the ellipse, for different stars, varies in this proportion, namely, as the right sine of the star's latitude; that is, radius is to the sine of the star's latitude, as the transverse axis to the conjugate axis: and consequently a star in the pole of the ecliptic, its latitude being there  $90^{\circ}$ , whose sine is equal to the radius, will appear to describe a small circle about that pole as a centre, whose radius is equal to  $20''$ . He also gives the following law of the variation of the star's declination: if A denote the angle of position, or the angle at the star made by two great circles drawn from it through the poles of the ecliptic and equator, and B another angle, whose tangent is to the tangent of A, as radius is to the sine of the star's latitude; then B will be equal to the difference of longitude between the sun and the star, when the true and apparent declination of the star are the same. And if the sun's longitude in the ecliptic be reckoned from that point in which it is when this happens; then the difference between the true and apparent declination of the star, will be always as the sine of the sun's longitude from that point. It will also be found that the greatest difference of declination that can be between the true and apparent place of the star, will be to  $20''$ , the semitransverse axis of the ellipse, as the sine of A to the sine of B.

The author then shews, by the comparison of a number of observations made on different stars, that they exactly agree with the theory deduced from the progressive motion of light, and that consequently it is highly probable that such motion is the cause of those variations in the situation of the stars. From which he infers, that the parallax of the fixed stars is much smaller, than hath been hitherto supposed by those, who have pretended to deduce it from their observations. He thinks he may venture to say, that in the stars he had observed, the parallax does not amount to  $2''$ ; nay, that if it had amounted to  $1''$ , he should certainly have perceived it, in the great number of observations that he made, especially of  $\gamma$  Draconis; which agreeing with the hypothesis, without allowing any thing for parallax, nearly as well when the sun was in conjunction with, as in opposition to, this star, it seems very probable



bable that the parallax of it is not so much as one single second; and consequently that it is above 400000 times farther from us than the sun.

From the greatest variation in the place of the stars, namely  $40''$ , Dr. Bradley deduces the ratio of the velocity of light in comparison with that of the earth in her orbit. In the preceding figure,  $AC$  is to  $AB$ , as the velocity of light to that of the earth in her orbit, the angle  $ACB$  being equal to  $20''$ ; so that the ratio of those velocities is that of radius to the tangent of  $20''$ , or of radius to  $20''$ , since the tangent has no sensible difference from so small an arc: but the radius of a circle is equal to the arc of  $57^{\circ}\frac{3}{8}$  nearly, or equal to  $206260''$ ; therefore the velocity of light is to the velocity of the earth, as  $206260$  to  $20$ , or as  $10313$  to  $1$ .

And hence also the time in which light passes over the space from the sun to the earth, is easily deduced; for this time will be to one year, as  $AB$  or  $20''$  to  $360^{\circ}$  or the whole circle; that is,  $360^{\circ} : 20'' :: 365\frac{1}{4}$  days :  $8^m 7^s$ , namely, light will pass from the sun to the earth in the time of 8 minutes, 7 seconds; and this will be the same, whatever the distance of the sun is.

Dr. Bradley having annexed to his theory the rules or formulæ for computing the aberration of the fixed stars in declination and right ascension; these rules have been variously demonstrated, and reduced to other practical forms, by Mr. Clairaut in the *Memoirs of the Academy of Sciences* for 1737; by Mr. Simpson in his *Essays* in 1740; by M. Fontaine des Crues in 1744; and several other persons. The results of these rules are as follow: Every star appears to describe in the course of a year, by means of the aberration, a small ellipse, whose greater axis is  $40''$ , and the less axis, perpendicular to the ecliptic, is equal to  $40''$  multiplied by the sine of the star's latitude, the radius being  $1$ . The eastern extremity of the longer axis, marks the apparent place of the star, the day of the opposition; and the extremity of the less axis, which is farthest from the ecliptic, marks its situation three months after.

The greatest aberration in longitude, is equal to  $20''$  divided by the cosine of its latitude. And the aberration for any time, is equal to  $20''$  multiplied by the cosine of the elongation of the star found for the same time, and divided by the cosine of its latitude. This aberration is subtractive in the first and last quadrant of the argument, or of the difference between the longitudes of the sun and star; and additive in the second and third quadrants. The greatest aberration in latitude, is equal to  $20''$  multiplied by the sine of the star's latitude. And the aberration in latitude for any time, is equal to  $20''$  multiplied by the sine of the star's latitude, and multiplied also by the sine of the elongation. The aberration is subtractive before the opposition, and additive after it.

The greatest aberration in declination, is equal to  $20''$  multiplied by the sine of the angle of position  $A$ , and divided by the sine of  $B$  the difference of longitude between the sun and star when the aberration in declination is nothing. And the aberration in declination at any other time, will be equal to the greatest aberration multiplied by the sine of the difference between the sun's place at the given time and his place when the

aberration is nothing. Also the sine of the latitude of the star is to radius, as the tangent of  $A$  the angle of position at the star, is to the tangent of  $B$ , the difference of longitude between the sun and star when the aberration in declination is nothing. The greatest aberration in right-ascension, is equal to  $20''$  multiplied by the cosine of  $A$  the angle of position, and divided by the sine of  $C$  the difference in longitude between the sun and star when the aberration in right ascension is nothing. And the aberration in right-ascension at any other time, is equal to the greatest aberration multiplied by the sine of the difference between the sun's place at the given time, and his place when the aberration is nothing. Also the sine of the latitude of the star is to radius, as the cotangent of  $A$  the angle of position, to the tangent of  $C$ .

*ABERRATION of the Planets*, is equal to the geocentric motion of the planet, the space it appears to move as seen from the earth, during the time that light employs in passing from the planet to the earth. Thus, in the sun, the aberration in longitude is constantly  $20''$ , that being the space moved by the sun, or, which is the same thing, by the earth, in the time of  $8^m 7^s$ , which is the time in which light passes from the sun to the earth, as we have seen in the foregoing article. In like manner, knowing the distance of any planet from the earth, by proportion it will be, as the distance of the sun is to the distance of the planet, so is  $8^m 7^s$  to the time of light passing from the planet to the earth: then computing the planet's geocentric motion in this time, that will be the aberration of the planet, whether it be in longitude, latitude, right-ascension, or declination.

It is evident that the aberration will be greatest in the longitude, and very small in latitude, because the planets deviate very little from the plane of the ecliptic, or path of the earth; so that the aberration in the latitudes of the planets, is commonly neglected, as insensible; the greatest in Mercury being only  $4''\frac{1}{3}$ , and much less in the other planets. As to the aberrations in declination and right-ascension, they must depend on the situation of the planet in the zodiac. The aberration in longitude, being equal to the geocentric motion, will be more or less according as that motion is; it will therefore be least, or nothing at all, when the planet is stationary; and greatest in the superior planets Mars, Jupiter, Saturn, &c, when they are in opposition to the sun; but in the inferior planets Venus and Mercury, the aberration is greatest at the time of their superior conjunction. These maxima of aberration for the several planets, when their distance from the sun is least, are as below: viz, for

Saturn	-	-	-	$27''\cdot 0$
Jupiter	-	-	-	$29 \cdot 8$
Mars	-	-	-	$37 \cdot 8$
Venus	-	-	-	$43 \cdot 2$
Mercury	-	-	-	$59 \cdot 0$
The Moon	-	-	-	$\frac{2}{3}$

And between these numbers and nothing the aberrations of the planets, in longitude, vary according to their situations. But that of the sun varies not, being constantly  $20''$ , as has been before observed. And this may alter his declination by a quantity, which varies from  $0$  to near  $8''$ ; being greatest or  $8''$  about the equinoxes, and vanishing in the solstices.


The



The methods of computing these, and the formulas for all cases, are given by M. Clairaut in the Memoirs of the Academy of Sciences for the year 1746, and by M. Euler in the Berlin Memoirs, vol. 2, for 1746.

*Optic ABERRATION*, the deviation or dispersion of the rays of light, when reflected by a speculum, or refracted by a lens, by which they are prevented from meeting or uniting in the same point, called the geometrical focus, but are spread over a small space, and produce a confusion of images. Aberration is either lateral or longitudinal: the lateral aberration is measured by a perpendicular to the axis of the speculum or lens, drawn from the focus to meet the refracted or reflected ray: the longitudinal aberration is the distance, on the axis, between the focus and the point where the ray meets the axis. The aberrations are very amply treated in Smith's Complete System of Optics, in 2 volumes 4to.

There are two species of aberration, distinguished according to their different causes: the one arises from the figure of the speculum or lens, producing a geometrical dispersion of the rays, when these are perfectly equal in all respects; the other arises from the unequal refrangibility of the rays of light themselves; a discovery that was made by Sir Isaac Newton, and for this reason it is often called the Newtonian aberration. As to the former species of aberration, or that arising from the figure, it is well known that if rays issue from a point at a given distance; then they will be reflected into the other focus of an ellipse having the given luminous point for one focus, or directly from the other focus of an hyperbola; and will be variously dispersed by all other figures. But if the luminous point be infinitely distant, or, which is the same, the incident rays be parallel, then they will be reflected by a parabola into its focus, and variously dispersed by all other figures. But those figures are very difficult to make, and therefore curved specula are commonly made spherical, the figure of which is generated by the revolution of a circular arc, which produces an aberration of all rays, whether they are parallel or not, and therefore it has no accurate geometrical focus which is common to all the rays. Let  $BVF$  represent a concave spherical speculum, whose centre is  $C$ ; and let  $AB, EF$  be incident rays parallel to the axis  $CV$ . Because the angle of incidence is equal to the angle of reflection in



all cases, therefore if the radii  $CB, CF$  be drawn to the points of incidence, and thence  $BD$  making the angle  $CBD$  equal to the angle  $CBA$ , and  $FG$  making the angle  $CFG$  equal to the angle  $CFE$ ; then  $BD, FG$  will be the reflected rays, and  $D, G$ , the points where

they meet the axis. Hence it appears that the point of coincidence with the axis is equally distant from the point of incidence and the centre: for because the angle  $CBD$  is equal to the angle  $CBA$ , which is equal to the alternate angle  $BCD$ , therefore their opposite sides  $CD, DB$  are equal: and in like manner, in any other,  $GF$  is equal to  $GC$ . And hence it is evident that when  $B$  is indefinitely near the vertex  $V$ ; then  $D$  is in the middle of the radius  $CV$ ; and the nearer the incident ray is to the axis  $CV$ , the nearer will the reflected ray come to the middle point  $D$ ; and the contrary. So that the aberration  $DG$  of any ray  $EF$ , is always more and more, as the incident ray is farther from the axis, or the incident point  $F$  from the vertex  $V$ ; till when the distance  $VI$  is 60 degrees, then the reflected ray falls in the vertex  $V$ , making the aberration equal to the whole length  $DV$ . And this shews the reason why specula are made of a very small segment of a sphere, namely, that all their reflected rays may arrive very near the middle point or focus  $D$ , to produce an image the most distinct, by the least aberration of the rays. And in like manner for rays refracted through lenses.

In spherical lenses, Mr. Huygens has demonstrated that the aberration from the figure, in different lenses, is as follows.

1. In all plano-convex lenses, having their plane surface exposed to parallel rays, the longitudinal aberration of the extreme ray, or that remotest from the axis, is equal to  $\frac{2}{3}$  of the thickness of the lens.

2. In all plano-convex lenses, having their convex surface exposed to parallel rays, the longitudinal aberration of the extreme ray, is equal to  $\frac{7}{8}$  of the thickness of the lens. So that in this position of the same plano-convex lens, the aberration is but about one-fourth of that in the former; being to it only as 7 to 27.

3. In all double convex lenses of equal spheres, the aberration of the extreme ray, is equal to  $\frac{5}{8}$  of the thickness of the lens.

4. In a double convex lens, the radii of whose spheres are as 1 to 6, if the more convex surface be exposed to parallel rays, the aberration from the figure is less than in any other spherical lens; being no more than  $\frac{1}{4}$  of its thickness.

But the foregoing species of aberration, arising from the figure, is very small, and easily remedied, in comparison with the other, arising from the unequal refrangibility of the rays of light; insomuch that Sir Isaac Newton shews in his Optics, pa. 84 of the 8vo. edition, that if the object-glass of a telescope be plano-convex, the plane side being turned towards the object, and the diameter of the sphere, to which the convex side is ground, be 100 feet, the diameter of the aperture being 4 inches, and the ratio of the sine of incidence out of glass into air, be to that of refraction, as 20 to 31; then the diameter of the circle of aberrations will in this case be only  $\frac{261}{7200000}$  parts of an inch: while the diameter of the little circle, through which the same rays are scattered by unequal refrangibility, will be about the 55th part of the aperture of the object-glass, which here is 4 inches. And therefore the error arising from the spherical figure of the glass, is to the error arising from the different refrangibility of the rays, as  $\frac{261}{7200000}$  to  $\frac{4}{55}$ , that is as 1 to 5449.



So that it may seem strange that objects appear through telescopes so distinct as they do, considering that the error arising from the different refrangibility, is almost incomparably larger than that of the figure. Newton however solves the difficulty by observing that the rays, under their various aberrations, are not scattered uniformly over all the circular space, but collected infinitely more dense in the centre than in any other part of the circle; and that, in the way from the centre to the circumference, they grow more and more rare, so as at the circumference to become infinitely rare; and, by reason of their rarity, they are not strong enough to be visible, unless in the centre, and very near it.

In consequence of the discovery of the unequal refrangibility of light, and the apprehension that equal refractions must produce equal divergencies in every sort of medium, it was supposed that all spherical object-glasses of telescopes would be equally affected by the different refrangibility of light, in proportion to their aperture, of whatever materials they might be constructed: and therefore that the only improvement that could be made in refracting telescopes, was that of increasing their length. So that Sir Isaac Newton, and other persons after him, despairing of success in the use and fabric of lenses, directed their chief attention to the construction of reflecting telescopes.

However, about the year 1747, M. Euler applied himself to the subject of refraction; and pursued a hint suggested by Newton, for the design of making object-glasses with two lenses of glass inclosing water between them; hoping that, by constructing them of different materials, the refractions would balance one another, and so the usual aberration be prevented. Mr. John Dollond, an ingenious optician in London, minutely examined this scheme, and found that Mr. Euler's principles were not satisfactory. M. Clairaut likewise, whose attention had been excited to the same subject, concurred in opinion that Euler's speculations were more ingenious than useful. This controversy, which seemed to be of great importance in the science of optics, engaged also the attention of M. Klingenstierna of Sweden, who was led to make a careful examination of the 8th experiment in the second part of Newton's Optics, with the conclusions there drawn from it. The consequence was, that he found that the rays of light, in the circumstances there mentioned, did not lose their colour, as Sir Isaac had imagined. This hint of the Swedish philosopher led Mr. Dollond to re-examine the same experiment: and after several trials it appeared, that different substances caused the light to diverge very differently, in proportion to their general refractive powers. In the year 1757 therefore he procured wedges of different kinds of glass, and applied them together so that the refractions might be made in contrary directions, that he might discover whether the refraction and divergency of colour would vanish together. The result of his first trials encouraged him to persevere; for he discovered a difference far beyond his hopes in the qualities of different kinds of glass, with respect to their divergency of colours. The Venice glass and English crown glass were found to be nearly allied in this respect: the common English plate glass made the rays diverge more; and the English flint

glass most of all. But without enquiring into the cause of this difference, he proceeded to adapt wedges of crown glass, and of white flint glass, ground to different angles, to each other, so as to refract in different directions; till the refracted light was entirely free from colours. Having measured the refractions of each wedge, he found that the refraction of the white glass was to that of the crown glass, nearly as 2 to 3: and he hence concluded in general, that any two wedges made in this proportion, and applied together so as to refract in contrary directions, would refract the light without any aberration of the rays.

Mr. Dollond's next object was to make similar trials with spherical glasses of different materials, with the view of applying his discovery to the improvement of telescopes: and here he perceived that, to obtain a refraction of light in contrary directions, the one glass must be concave, and the other convex; and the latter, which was to refract the most, that the rays might converge to a real focus, he made of crown glass, the other of white flint glass. And as the refractions of spherical glasses are inversely as their focal distances, it was necessary that the focal distances of the two glasses should be inversely as the ratios of the refractions of the wedges; because that, being thus proportioned, every ray of light that passes through this compound glass, at any distance from its axis, will constantly be refracted, by the difference between two contrary refractions, in the proportion required; and therefore the different refrangibility of the light will be entirely removed.

But in the applications of this ingenious discovery to practice, Mr. Dollond met with many and great difficulties. At length, however, after many repeated trials, by a resolute perseverance, he succeeded so far as to construct refracting telescopes much superior to any that had hitherto been made; representing objects with great distinctness, and in their true colours:

Mr. Clairaut, who had interested himself from the beginning in this discovery, now endeavoured to ascertain the principles of Mr. Dollond's theory, and to lay down rules to facilitate the construction of these new telescopes. With this view he made several experiments, to determine the refractive power of different kinds of glass, and the proportions in which they separated the rays of light: and from these experiments he deduced several theorems of general use. M. D'Alembert made likewise a great variety of calculations to the same purpose; and he shewed how to correct the errors to which these telescopes are subject, sometimes by placing the object-glasses at a small distance from each other, and sometimes by using eye-glasses of different refractive powers. But though foreigners were hereby supplied with the most accurate calculations, they were very defective in practice. And the English telescopes, made, as they imagined, without any precise rule, were greatly superior to the best of their construction.

M. Euler, whose speculations had first given occasion to this important and useful enquiry, was very reluctant in admitting Mr. Dollond's improvements, because they militated against a pre-conceived theory of his own. At last however, after several altercations, being convinced of their reality and importance by M. Clair-



aut, he assented; and he soon after received farther satisfaction from the experiments of M. Zeiher, of Petersburg.

M. Zeiher shewed by experiments that it is the lead, in the composition of glass, which gives it this remarkable property, namely, that while the refraction of the mean rays is nearly the same, that of the extreme rays considerably differs. And, by increasing the lead, he produced a kind of glass, which occasioned a much greater separation of the extreme rays than that of the flint glass used by Mr. Dollond, and at the same time considerably increased the mean refraction. M. Zeiher, in the course of his experiments, made glass of minium and lead, with a mixture also of alkaline salts; and he found that this mixture greatly diminished the mean refraction, and yet made hardly any change in the dispersion: and he at length obtained a kind of glass greatly superior to the flint glass of Mr. Dollond for the construction of telescopes; as it occasioned three times as great a dispersion of the rays as the common glass, whilst the mean refraction was only as 1.61 to 1.


Other improvements were also made on the new or achromatic telescopes by the inventor Mr. John Dollond, and by his son Peter Dollond; which may be seen under the proper words. For various dissertations on the subject of the aberration of light, colours, and the figure of the glass, see *Philos. Trans.* vols. 35, 48, 50, 51, 52, 55, 60; *Memoirs of the Academy of Sciences of Paris*, for the years 1737, 1746, 1752, 1755, 1756, 1757, 1762, 1764, 1765, 1767, 1770; the *Berlin Ac.* 1746, 1762, 1766; *Swed. Mem.* vol. 16; *Com. Nov. Petropol.* 1762; M. Euler's *Dioptrics*; M. d'Alembert's *Opuscles Math.*; M. de Rochon *Opuscles*; &c, &c.

**ABRIDGING**, in *Algebra*, is the reducing a compound equation, or quantity, to a more simple form of expression. This is done either to save room, or the trouble of writing a number of symbols; or to simplify the expression, either to ease the memory, or to render the formula more easy and general.

So the equation  $x^3 - ax^2 + abx - abc = 0$ ; by putting  $p = a$ ,  $q = ab$ , and  $r = abc$ , becomes  $x^3 - px^2 + qx - r = 0$ .

And the equation  $x^2 + (a+b)x - \frac{ab}{c} = 0$ , by putting  $p = a+b$ , and  $q = \frac{ab}{c}$ , becomes  $x^2 + px - q = 0$ .


**ABSCISS**, **ABSCISSE**, or **ABSCISSA**, is a part or segment cut off a line, terminated at some certain point, by an ordinate to a curve; as AP or BP.



The absciss may either commence at the vertex of the curve, or at any other fixed point. And it may be taken either upon the axis or diameter of the curve, or upon any other line drawn in a given position.

Hence there are an infinite number of variable abscissas, terminated at the same fixed point at one end, the other end of them being at any point of the given line or diameter.

In the common parabola, each ordinate PQ has but



one absciss AP; in the ellipse or circle, the ordinate has two abscissas AP, BP lying on the opposite sides of it; and in the hyperbola the ordinate PQ has also two abscissas, but they lie both on the same side of it. That is, in general, a line of the second kind, or a curve of the first kind, may have two abscissas to each ordinate. But a line of the third order may have three abscissas to each ordinate; a line of the fourth order may have four; and so on.

The use of the abscissas is, in conjunction with the ordinates, to express the nature of the curves, either by some proportion or equation including the absciss and its ordinate, with some other fixed invariable line or lines. Every different curve has its own peculiar equation or property by which it is expressed, and different from all others: and that equation or expression is the same for every ordinate and its abscissas, whatever point of the curve be taken. So, in the circle, the square of any ordinate is equal to the rectangle of its two abscissas, or  $AP \cdot PB = PQ^2$ ; in the parabola, the square of the ordinate is equal to the rectangle of the absciss and a certain given line called the parameter; in the ellipse and hyperbola, the square of the ordinate is always in a certain constant proportion to the rectangle of the two abscissas, namely, as the square of the conjugate to the square of the transverse, or as the parameter is to the transverse axis; and so other properties in other curves.

When the natures or properties of curves are expressed by algebraic equations, any general absciss, as AP, is commonly denoted by the letter  $x$ , and the ordinate PQ by the letter  $y$ ; the other or constant lines being represented by other letters. Then the equations expressing the nature of these curves are as follow; namely, for the

circle -  $dx - x^2 = y^2$ , where  $d$  is the diameter AB;  
parabola -  $px = y^2$ , where  $p$  is the parameter;  
ellipse -  $t^2 : c^2 :: tx - a^2 : y^2$ , { where  $t$  is the transverse,  
hyperbola  $t^2 : c^2 :: tx + a^2 : y^2$ , { &  $c$  the conjugate axis.

**ABSIS**, **ABSIDES**. See **APSIS**, **APSIDES**.

**ABSOLUTE**



**ABSOLUTE EQUATION**, in *Astronomy*, is the sum of the optic and excentric equations. The apparent inequality of a planet's motion, arising from its not being equally distant from the earth at all times, is called its optic equation; and this would subsist even if the planet's real motion were uniform. The excentric inequality is caused by the planet's motion being not uniform. To illustrate this, conceive the sun to move, or to appear to move, in the circumference of a circle, in whose centre the earth is placed. It is manifest, that if the sun move uniformly in this circle, then he must appear to move uniformly to a spectator at the earth; and in this case there will be no optic nor excentric equation. But suppose the earth to be placed out of the centre of the circle; and then, though the sun's motion should be really uniform, it would not appear to be so, being seen from the earth; and in this case there would be an optic equation, without an excentric one. Imagine farther, the sun's orbit to be, not circular, but elliptical, and the earth in its focus: it will be full as evident that the sun cannot appear to have an uniform motion in such ellipse; so that his motion will then be subject to two equations; that is, the optic equation, and the excentric equation. See **EQUATION**, and **OPTIC INEQUALITY**.

**ABSOLUTE Number**, in *Algebra*, is that term or member of an equation that is completely known, and which is equal to all the other, or unknown terms, taken together; and is the same as what Vieta calls the *homogeneum comparationis*. So, of the equation  $x^2 + 16x - 36 = 0$ , or  $x^2 + 16x = 36$ , the absolute number, or known term, is 36.

**ABSOLUTE Gravity, Motion, Space, Time, &c.** See the respective substantives.

**ABSTRACT MATHEMATICS**, otherwise called pure mathematics, is that which treats of the properties of magnitude, figure, or quantity, absolutely and generally considered, without restriction to any species in particular: such as Arithmetic and Geometry. In this sense, abstract or pure mathematics, is opposed to mixed mathematics, in which simple and abstract properties, and the relations of quantities, primitively considered in pure mathematics, are applied to sensible objects; as in astronomy, hydrostatics, optics, &c.

**ABSTRACT Number**, is a number, or collection of units, considered in itself, without being applied to denote a collection of any particular and determinate things. So, for example, 3 is an abstract number, so far as it is not applied to something: but when we say 3 feet, or 3 persons, the 3 is no longer an abstract, but a concrete number.

**ABSURD**, or **ABSURDUM**, a term commonly used in demonstrating converse propositions; a mode of demonstration, in which the proposition intended is not proved in a direct manner, by principles before laid down; but it proves that the contrary is absurd or impossible; and so indirectly as it were proves the proposition itself. The 4th proposition in the first book of Euclid, is the first in which he makes use of this mode of proof; where he shews that if the extremities of two lines coincide, those lines will coincide in all their parts, otherwise they would inclose a space, which is absurd or contrary to the 10th axiom. Most converse

propositions are proved in this way, which mode of proof is called *reductio ad absurdum*.

**ABUNDANT Number**, in *Arithmetic*, is a number whose aliquot parts, added all together, make a sum which is greater than the number itself. Thus 12 is an abundant number, because its aliquot parts, namely 1, 2, 3, 4, 6, when added together, make 16, which is greater than the number 12 itself.

An abundant number is opposed to a deficient one, which is less than the sum of its aliquot parts taken together, as the number 14, whose aliquot parts 1, 2, 7, make no more than 10; and to a perfect number, which is exactly equal to the sum of all its aliquot parts, as the number 6, which is equal to the sum of 1, 2, 3, which are its aliquot parts.

**ACADEMICIAN**, a member of a society called an academy, instituted for the promotion of arts, sciences, or natural knowledge in general.

**ACADEMICS**, an ancient sect of philosophers, who followed the doctrine of Socrates and Plato, as to the uncertainty of knowledge, and the incomprehensibility of truth.

*Academic*, in this sense, amounts to much the same with *Platonist*; the difference between them being only in point of time. Those who embraced the system of Plato, among the ancients, were called *academici*, *academician* or *academic*; whereas those who did the same since the restoration of learning, have assumed the denomination of *Platonists*.

We usually reckon three sects of academics; though some make five. The ancient academy was that of which Plato was the chief.

Arcefilas, one of Plato's successors, introducing some alterations into the philosophy of this sect, founded what they call the second academy.

The establishment of the third, called also the new academy, is attributed to Lacydes, or rather to Carneades.

Some authors add a fourth, founded by Philo; and a fifth, by Antiochus, called the Antiochan, which tempered the ancient academy with Stoicism.

The ancient academy doubted of every thing; and carried this principle so far as to make it a doubt, whether or no they ought to doubt. It was a kind of a principle with them, never to be certain or satisfied of any thing; never to affirm or to deny any thing, either for true or false.

The new academy was somewhat more reasonable; they acknowledged several things for truths, but without attaching themselves to any with entire assurance. These philosophers had found that the ordinary commerce of life and society was inconsistent with the absolute and universal doubtfulness of the ancient academy: and yet it is evident that they looked upon things rather as probable, than as true and certain: by this amendment thinking to secure themselves from those absurdities into which the ancient academy had fallen.

**ACADEMIST**, the same as *Academician*.

**ACADEMY**, **ACADEMIA**, in *Antiquity*, a fine villa or pleasure house, in one of the suburbs of Athens, about a mile from the city; where Plato, and the wise men who followed him, held assemblies for disputes



and philosophical conference; which gave the name to the sect of Academics.

The house took its name, *Academy*, from one *Academus*, or *Ecademus*, a citizen of Athens, to whom it originally belonged: he lived in the time of Theseus; and here he used to have gymnastic sports or exercises.

The academy was farther improved by Cimon, and adorned with fountains, trees, shady walks, &c, for the convenience of the philosophers and men of learning, who here met to confer and dispute for their mutual improvement. It was surrounded with a wall by Hipparchus, the son of Pisistratus; and it was also used as the burying-place for illustrious persons, who had deserved well of the republic.

It was here that Plato taught his philosophy; and hence it was that all public places, destined for the assemblies of the learned and ingenious, have been since called *Academies*.

Sylla sacrificed the delicious walks and groves of the academy, which had been planted by Cimon, to the ravages of war; and employed those very trees in constructing machines to batter the walls of the city which they had adorned.

Cicero too had a villa, or country retirement, near Puzzuoli, which he called by the same name, *Academia*. Here he used to entertain his philosophical friends; and here it was that he composed his *Academical Questions*, and his books *De Naturâ Deorum*.

ACADEMY, among the moderns, denotes a regular society or company of learned persons, instituted under the protection of some prince, or other public authority, for the cultivation and improvement of arts or sciences.

Some authors confound Academy with University; but though much the same in Latin, they are very different things in English. An university is properly a body composed of graduates in the several faculties; of professors, who teach in the public schools; of regents or tutors, and students who learn under them, and aspire likewise to degrees. Whereas an academy is not intended to teach, or profess any art or science, but to improve it: it is not for novices to be instructed in, but for those that are more knowing; for persons of learning to confer in, and communicate their lights and discoveries to each other, for their mutual benefit and improvement.

The first modern academy we read of, was established by Charlemagne, by the advice of Alcuin, an English monk: it was composed of the chief geniuses of the court, the emperor himself being a member. In their academical conferences, every person was to give some account of the ancient authors he had read; and each one assumed the name of some ancient author, that pleased him most, or some celebrated person of antiquity. Alcuin, from whose letters we learn these particulars, took that of Flaccus, the surname of Horace; a young lord, named Augilbert, took that of Homer; Adelard, bishop of Corbie, was called Augustin; Reluse, bishop of Mentz, was Dametas; and the king himself, David.

Since the revival of learning in Europe, academies have multiplied greatly, most nations being furnished with several, and from their communications the chief

improvements have been made in the arts and sciences, and in cultivating natural knowledge. There are now academies for almost every art, or species of knowledge; but I shall give a short account only of those institutions of this kind, which regard the cultivation of subjects mathematical or philosophical, which are the proper and peculiar objects of our undertaking.

Italy abounds more in academies than all the world besides; there being enumerated by Jarckius not less than five hundred and fifty in all; and even to the amount of twenty-five in Milan itself. These are however mostly of a private and inferior nature; the consequence of their too great number.

The first academy of a philosophical kind was established at Naples, in the house of Baptista Porta, about the year 1560, under the name of *Academy Secretorum Naturæ*; being formed for the improvement of natural and mathematical knowledge. This was succeeded by the

ACADEMY of *Lyncei*, founded at Rome by prince Frederick Cesi, towards the end of the same century. It was rendered famous by the notable discoveries made by several of its members; among whom was the celebrated Galileo Galilei.

Several other academies contributed also to the advancement of the sciences; but it was by speculations rather than by repeated experiments on the phenomena of nature: such were the academy of Bessarion at Rome, and that of Laurence de Medicis at Florence, in the 15th century; and in the 16th were that of Inflammati at Padua, of Vegna Juoli at Rome, of Ortolani at Placentia, and of Umidi at Florence. The first of these studied fire and pyrotechnia, the second wine and vineyards, the third pot-herbs and gardens, the fourth water and hydraulics. To these may be added that of Venice, called *La Veneta*, and founded by Frederick Badoara, a noble Venetian; another in the same city; of which Campegio, bishop of Feltro, appears to have been the chief; also that of Cosenza, or *La Consentina*, of which Bernadin Telefio, Sertorio Quatromanni, Paulus Aquinas, Julio Cavalcanti, and Fabio Cicali, celebrated philosophers, were the chief members. The compositions of all these academies, of the 16th century, were good in their kind; but none of them comparable to those of the *Lyncei*.

ACADEMY of *del Cimento*, that is, of Experiments, arose at Florence, some years after the death of Torricelli, namely in the year 1657, under the protection of prince Leopold of Tuscany, afterwards cardinal de Medicis, and brother to the Grand Duke Ferdinand the Second. Galileo, Toricelli, Aggiunti, and Viviani had prepared the way for it: and some of its chief members were Paul del Buono, who in 1657 invented the instrument for trying the incompressibility of water, namely a thick globular shell of gold, having its cavity filled with water; then the globe being compressed by a strong screw, the water came through the pores of the gold rather than yield to the compression: also, Alphonfus Borelli, well known for his ingenious treatise *De Motu Animalium*, and other works; Candide del Buono, brother of Paul; Alexander Marsili, Vincent Viviani, Francis Rhedi, and the Count Laurence Magalotti, secretary of this academy, who published



lished a volume of their curious experiments in 1667, under the title of *Saggi di Naturali Esperienze*; a copy of which being presented to the Royal Society, it was translated into English by Mr. Waller, and published at London, in 4to, 1684: A curious collection of tracts, containing ingenious experiments on the pressure of the air, on the compressing of water, on cold, heat, ice, magnets, electricity, odours, the motion of sound, projectiles, light, &c. &c. But we have heard little or nothing more of the academy since that time. It may not be improper to observe here, that the Grand Duke Ferdinand, above mentioned, was no mean philosopher and chemist, and that he invented thermometers, of which the construction and use may be seen in the collection of the academy del Cimento.

ACADEMY *degl' Inquieti* at Bologna, incorporated afterwards into that *della traccia* in the same city, followed the example of that *del Cimento*. The members met at the house of the abbot Antonio Sampieri; and here Geminiano Montanari, one of the chief members, made excellent discourses on mathematical and philosophical subjects, some parts of which were published in 1667, under the title of *Pensieri Fisico-Matematici*. This academy afterwards met in an apartment of Eustachio Manfredi; and then in that of Jacob Sandri; but it arrived at its chief lustre while its assemblies were held in the palace Marfilli.

ACADEMY of *Rossano*, in the kingdom of Naples, called *La Societa Scientifica Rossanese degl' Incuriosi*, was founded about the year 1540, under the name of *Naviganti*; and was renewed under that of *Spensierati* by Camillo Tuscano, about the year 1600. It was then an academy of belles-lettres, but was afterwards transformed into an academy of sciences, on the solicitation of the learned abbot Don Giacinto Gimma; who, being made president under the title of promoter-general, in 1695, gave it a new set of regulations. He divided the academists into several classes, namely, grammarians, rhetoricians, poets, historians, philosophers, physicians, mathematicians, lawyers, and divines; with a separate class for cardinals and persons of quality. To be admitted a member, it was necessary that the candidate have degrees in some faculty. Members, in the beginning of their books, are not allowed to take the title of *academist* without a written permission from the president, which is not granted till the work has been examined by the censors of the academy. This permission is the highest honour the academy can confer; since they hereby, as it were, adopt the work, and engage to answer for it against any criticisms that may be made upon it. The president himself is not exempt from this law: and it is not permitted that any academist publish any thing against the writings of another, without leave obtained from the society.

There have been several other academies of sciences in Italy, but which have not subsisted long, for want of being supported by the princes. Such were at Naples that of the *Investiganti*, founded about the year 1679, by the marquis d'Arena, Don Andrea Conculletto; and that which, about the year 1698, met in the palace of Don Lewis della Cerda, the duke de Medina, and viceroy of Naples: at Rome, that of *Fisico-Matematici*, which in 1686 met in the house of

Signior Ciampini: at Verona, that of *Aletofili*, founded the same year by Signior Joseph Gazola, and which met in the house of the count Serenghi della Cucca: at Brescia, that of *Filefotici*, founded the same year for the cultivation of philosophy and mathematics, and terminated the year following: that of F. Francisco Lana, a jesuit of great skill in these sciences: and lastly that of *Fisico-Critici* at Sienna, founded in 1691, by Signior Peter Maria Gabrielli.

Some other academies, still subsisting in Italy, repair with advantage the loss of the former. One of the principal is the academy of *Filarmonici* at Verona, supported by the marquis Scipio Maffei, one of the most learned men in Italy; the members of which academy, though they cultivate the belles lettres, do not neglect the sciences. The academy of *Ricovrati* at Padua still subsists with reputation; in which, from time to time, learned discourses are held on philosophical subjects. The like may be said of the academy of the *Muti di Reggio*, at Modena. At Bologna is an academy of sciences, in a flourishing condition, known by the name of *The Institute of Bologna*; which was founded in 1712 by count Marsigli, for cultivating physics, mathematics, medicine, chemistry, and natural history. The history of it is written by M. de Limiers, from memoirs furnished by the founder himself. Among the new academies, the first place, after the Institute of Bologna, is given to that of the Countess Donna Clelio Grillo Boromeo, one of the most learned ladies of the age, to whom Signior Gimma dedicates his literary history of Italy. She had lately established an academy of experimental philosophy in her palace at Milan; of which Signior Vallisnieri was nominated president, and had already drawn up the regulations for it, though we do not find it has yet taken place. In the number of these academies may also be ranked the assembly of the learned, who of late years met at Venice in the house of Signior Cristino Martinelli, a noble Venetian, and a great patron of learning.

ACADEMIA *Cosmografica*, or that of the Argonauts, was instituted at Venice, at the instance of F. Coronelli, for the improvement of geography; the design being to procure exact maps, geographical, topographical, hydrographical, and ichnographical, of the celestial as well as terrestrial globe, and their several regions or parts, together with geographical, historical, and astronomical descriptions accommodated to them: to promote which purposes, the several members oblige themselves, by their subscription, to take one copy or more of each piece published under the direction of the academy; and to advance the money, or part of it, to defray the charge of publication. To this end there are three societies settled, namely at Venice, Paris, and Rome; the first under F. Moro, provincial of the Minorites of Hungary; the second under the abbot Laurence au Rue Payenne au Marais; the third under F. Ant. Baldigiani, jesuit, professor of mathematics in the Roman college; to whom those address themselves who are willing to engage in this design. The Argonauts number near 200 members in the different countries of Europe; and their device is the terraqueous globe, with the motto *Plus ultra*. All the globes, maps, and geographical writings of F. Coronelli have been published at the expence of this academy.



THE ACADEMY of *Artists*, or Impartial Academy, deserves to be mentioned on account of the extent of its plan, including universally all arts and sciences. It holds from time to time public meetings at Florence, where any person, whether academist or not, may read his works, in whatever form, language, or subject; the academy receiving all with the greatest impartiality.

In France there are many academies for the improvement of arts and sciences. F. Merfenne, it is said, gave the first idea of a philosophical academy in France, about the beginning of the seventeenth century, by the conferences of mathematicians and naturalists, held occasionally at his lodgings; at which Des Cartes, Gassendus, Hobbes, Roberval, Pascal, Blondel, and others, assisted. F. Merfenne proposed to each of them certain problems to examine, or certain experiments to be made. These private assemblies were succeeded by more public ones, formed by M. Monmort, and M. Thevenot, the celebrated traveller. The French example animated several Englishmen of rank and learning to erect a kind of philosophical academy at Oxford, towards the close of Cromwell's administration; which after the restoration was erected, by public authority, into a Royal Society: an account of which see under the word. The English example, in its turn, animated the French. In 1666 Louis XIV, assisted by the counsels of M. Colbert, founded an academy of sciences at Paris, called the

ACADEMIE Royale des Sciences, or Royal Academy of Sciences, for the improvement of philosophy, mathematics, chemistry, medicine, belles-lettres, &c. Among the principal members, at the commencement in 1666, were the respectable names of Carcavi, Huygens, Roberval, Frenicle, Auzout, Picard, Buot, Du Hamel the Secretary, and Mariotte. There was a perfect equality among all the members, and many of them received salaries from the king, as at present. By the rules of the academy, every class was to meet twice a week; the philosophers and geometricians were to meet, separately, every Wednesday, and then both together on the Saturday, in a room of the king's library, where the philosophical and mathematical books were kept: the history class was to meet on the Monday and Thursday in the room of the historical books: and the class of belles-lettres on the Tuesday and Friday: and on the first Thursday of every month all the classes met together, and by their secretaries made a mutual report of what had been transacted by each class during the preceding month.

In 1699, on the application of the president, the abbé Bignon, the academy received, under royal authority and protection, a new form and constitution; by the articles of which, the academy was to consist of four sorts of members, namely honorary, pensionary, associates, and eleves. The honorary class to consist of ten persons, and the other three classes of twenty persons each. The president to be chosen annually out of the honorary class, and the secretary and treasurer to be perpetual, and of the pensionary class. The meetings to be twice a week, on the Wednesday and Saturday; besides two public meetings in the year.

Of the pensionaries, or those who receive salaries, three to be geometricians, three astronomers, three mechanists, three anatomists, three botanists, and three

chemists, the other two being the secretary and treasurer. Of the twenty associates, of which twelve to be French, and eight might be foreigners, two were to cultivate geometry, two astronomy, two mechanics, two anatomy, two botany, and two chemistry. Of the twenty eleves, one to be attached to each pensionary, and to cultivate his peculiar branch of science. The pensionaries and their eleves to reside at Paris. No regulars nor religious to be admitted, except into the honorary class: nor any person to be admitted a pensioner who was not known by some considerable work, or some remarkable discovery.

In 1716 the Duke of Orleans, then regent of France, by the king's authority made some alteration in their constitution. The class of eleves was suppressed; and instead of them were instituted twelve adjuncts, two to each of the six classes of pensioners. The honorary members were increased to twelve: and a class of six free associates was made, who were not under the obligation of cultivating any particular branch of science, and in this class only could the regulars or religious be admitted. A president and vice-president to be appointed annually from the honorary class, and a director and sub-director annually from that of the pensioners. And no person to be allowed to make use of his quality of academician, in the title of any of his books that he published, unless such book were first approved by the academy.

The academy has for a device or motto, *Invenit & perficit*. And the meetings, which were formerly held in the king's library, have since the year 1699 been held in a fine hall of the old Louvre.

Finally, in the year 1785 the king confirmed, by letters patent, dated April 23, the establishment of the academy of sciences, making the following alterations, and adding classes of agriculture, natural history, mineralogy, and physics; incorporating the associates and adjuncts, and limiting to fix the members of each class, namely three pensioners and three associates; by which the former receive an increase of salary, and the latter approach nearer to becoming pensioners.

By the articles of this instrument it is ordained, that the academy shall consist of eight classes, namely, that of geometry, 2d astronomy, 3d mechanics, 4th general physics, 5th anatomy, 6th chemistry and metallurgy, 7th botany and agriculture, and 8th natural history and mineralogy. That each class shall remain irrevocably fixed at six members; namely, three pensioners and three associates, independent however of a perpetual secretary and treasurer, of twelve free-associates and of eight associate-strangers or foreigners, the same as before, except that the adjunct-geographer for the future be called the associate-geographer.

The classes at first to be filled by the following persons, namely, that of geometry by Messieurs de Borda, Jaurat, Vandermonde, as pensioners; and Messieurs Cousin, Meusnier, and Charles, as associates: that of astronomy by Messieurs le Monnier, de la Lande, and le Gentil, as pensioners; and Messieurs Messier, de Cassini, and Dagelet, as associates: that of mechanics by Messieurs l'abbé Bossut, l'abbé Rochon, and de la Place, as pensioners; and Messieurs Coulomb, le Gendre, and Perrier, as associates: that of general physics by Messieurs Leroy, Briffon, and Bailly,



Bailly, as pensioners; and Messieurs Monge, Mechain, and Quatremere, as associates: that of anatomy by Messieurs Daubluton, Tenon, and Portal, as pensioners; and Messieurs Sabatier, Vicq-d'azir, and Broussonet, as associates: that of chemistry and metallurgy by Messieurs Cadet, Lavoisier, and Beaume, as pensioners; and Messieurs Cornette, Bertholet, and Fourcroy, as associates: that of botany and agriculture by Messieurs Guettard, Foucheroux, and Adanson, as pensioners; and Messieurs de Jussieu, de la Marck, and Desfontaines, as associates: and that of natural history and mineralogy by Messieurs Desmarest, Saye, and l'abbé de Gua, as pensioners; and Messieurs Darcet, l'abbé Haui, and l'abbé Tessier, as associates. All names respectable in the common-wealth of letters; and from whom the world might expect still farther improvements in the several branches of science.

The late M. Rouille de Meilay, counsellor of the parliament of Paris, founded two prizes, the one of 2500 livres, the other of 2000 livres, which the academy distributed alternately every year: the subjects of the former prize respecting physical astronomy, and of the latter, navigation and commerce.

The world is highly indebted to this academy for the many valuable works they have executed, or published, both individually and as a body collectively, especially by their memoirs, making upwards of a hundred volumes in 4to, with the machines, indexes, &c. in which may be found most excellent compositions in every branch of science. They publish a volume of these memoirs every year, with the history of the academy, and eulogies of remarkable men lately deceased: also a general index to the volumes every ten years. An alteration was introduced into the volume for 1783, which it seems is to be continued in future, by omitting, in the history, the minutes or extracts from the registers, containing some preliminary account of the subjects of the memoirs; but still however retaining the eulogies of distinguished men, lately deceased.

M. l'abbé Rozier also has published in four 4to volumes, an excellent index of the contents of all the volumes, and the writings of all the members, from the beginning of their publications to the year 1770; with convenient blank spaces for continuing the articles in writing.

Their history also, to the year 1697, was written by M. Du Hamel; and after that time continued from year to year by M. Fontenelle, under the following titles, Du Hamel *Historiæ Regiæ Academiæ Scientiarum*, Paris, 4to. *Histoire de l'Académie Royale des Sciences, avec les Mémoires de Mathématique & de Physique, tirés des Registres de l'Académie*, Paris, 4to. *Histoire de l'Académie Royale des Sciences depuis son établissement en 1666, jusqu'en 1699, en 13 tomes*, 4to. A new history, from the institution of the academy, to the period from whence M. de Fontenelle commences, has been formed; with a series of the works published under the name of this academy, during the first interval.

Since the foregoing account was written, it is said the Academy has been suppressed and abolished, by the present convention of France.

Besides the academies in the capital, there are a great many in other parts of France. The *ACADEMIE Royale*, at Caen, was established by letters

patent in the year 1705; but it had its rise fifty years earlier in private conferences, held first in the house of M. de Brioux. M. de Segrais retiring to this city, to spend the rest of his days, restored and gave new lustre to their meetings. In 1707 M. Foucault, intendant of the generality of Caen, procured the king's letters patent for erecting them into a perpetual academy, of which M. Foucault was to be protector for the time, and the choice afterwards left to the members, the number of whom was fixed to thirty, chosen for this time by M. Foucault. But besides the thirty original members, leave was given to add six supernumerary members, from the ecclesiastical communities in that city.

At Toulouse is the *Académie des jeux floraux*, composed of forty persons, the oldest of the kingdom: besides an academy of sciences and belles-lettres, founded in 1750.

At Montpellier is the royal society of sciences, which since 1708 makes but one body with the royal academy of sciences at Paris.

There are also other academies at Bourdeaux, founded in 1703, at Soissons in 1674, at Marseilles in 1726, at Lyons in 1700, at Pau in Bearn in 1721, at Montauban in 1744, at Angers in 1685, at Amiens in 1750, at Villefranche in 1679, at Dijon in 1740, at Nîmes in 1682, at Besançon in 1752, at Chalons in 1775, at Rochelle in 1734, at Beziers in 1723, at Rouen in 1744, at Metz in 1760, at Arras in 1773, &c. The number of these academies is continually augmenting; and even in such towns as have no academies, the literati form themselves into literary societies, having nearly the same objects and pursuits.

In Germany and other parts of Europe, there are various academies of sciences, &c. The

*ACADEMIE Royale des Sciences & des Belles-Lettres* of Prussia, was founded at Berlin, in the year 1700, by Frederic I. king of Prussia, of which the famous M. Leibnitz was the first president, and its great promoter. The academy received a new form, and a new set of statutes in 1710; by which it was ordained, that the president shall be one of the counsellors of state; and that the members be divided into four classes; the first to cultivate physics, medicine, and chemistry; the second, mathematics, astronomy, and mechanics; the third, the German language, and the history of the country; and the fourth, oriental learning, particularly as it may concern the propagation of the gospel among infidels. That each class elect a director for themselves, who shall hold his post for life. That they meet in the castle called the New Marshal, the classes to meet in their turns, one each week. And that the members of any of the classes have free access into the assemblies of the rest. Several volumes of their transactions have been published in Latin, from time to time, under the title of *Miscellanea Berolinensia*.

In 1743 the late famous Frederic II. king of Prussia, made great alterations and improvements in the academy. Instead of a great lord or minister of state, who had usually presided over the academy, he wisely judged that office would be better filled by a man of letters; and he honoured the French academy of sciences by fixing upon one of its members for a president, namely M. Maupertuis, a distinguished character in the literary world, and whose conduct in improving the academy was a proof of the sound judgment of the king,



king, who at the same time made new regulations for the academy, and took the title of its Protector. From that time the transactions of the academy have been published, under the title of *Histoire de l'Academie Royale des Sciences et Belles Lettres à Berlin*, much in the manner of those of the French academy of sciences, and in the French language; and the volumes are now commonly published annually. Besides the ordinary private meetings of the academy, it has two public ones in the year, in January and May, at the latter of which is given a prize gold medal, of the value of 50 ducats, or about 20 guineas. The subject of the prize is successively physics, mathematics, metaphysics, and general literature. For the academy has this peculiar circumstance, that it embraces also metaphysics, logic, and morality; having one class particularly appropriated to these objects, called the class of Speculative Philosophy.

*Imperial ACADEMY of Petersburg.* This academy was projected by the Czar Peter I, commonly called Peter the Great, who in so many other instances also was instrumental in raising Russia from the state of barbarity in which it had been immersed for so many ages. Having visited France in 1717, and among other things informed himself of the advantages of an academy of arts and sciences, he resolved to establish one in his new capital, whither he had drawn by noble encouragements several learned strangers, and made other preparations, when his death prevented him from fully accomplishing that object, in the beginning of the year 1725. Those preparations and intentions however were carried into execution the same year, by the establishment of the academy, by his consort the czarina Catherine, who succeeded him. And soon after the academy composed the first volume of their works, published in 1728, under the title of *Commentarii Academiæ Scientiarum Imperialis Petropolitanae*; which they continued almost annually till 1746, the whole amounting to 14 volumes, which were published in Latin, and the subjects divided and classed under the following heads, namely mathematics, physics, history, and astronomy. Their device a tree bearing fruit not ripe, with the modest motto *paullatim*.

Most part of the strangers who composed this academy being dead, or having retired, it was rather in a languishing state at the beginning of the reign of the empress Elizabeth, when the count Rasomowski was happily appointed president, who was instrumental in recovering its vigour and labours. This empress renewed and altered its constitution, by letters patent dated July 24, 1747, giving it a new form and regulations. It consists of two chief parts, an academy, and a university, having regular professors in the several faculties, who read lectures as in our colleges. The ordinary assemblies are held twice a week, and public or solemn ones thrice in the year; in which an account is given of what has been done in the private ones. The academy has a noble building for their meetings, &c, with a good apparatus of instruments, a fine library, observatory, &c. Their first volume, after this renovation, was published for the years 1747 and 1748, and they have been since continued from year to year, now to the amount of near thirty volumes, under the title of *Novi Commentarii Academiæ Scientiarum Imperialis*

*Petropolitanae*. They are printed in the Latin language, and contain many excellent compositions in all the sciences, especially the mathematical papers of the late excellent M. L. Euler, which always made a considerable portion of every volume. The subjects are classed under heads in the following order, mathematics, physico-mathematics, physics, which include botany, anatomy, &c, and astronomy; the whole prefaced by historical extracts, or minutes, relating to each paper or memoir, after the manner of the volumes of the French academy; but wanting however the eulogies of deceased eminent men. Their device is a heap of ripe fruits piled on a table, with the motto *En addit fructus ætate recentes*.

*Imperial and Royal ACADEMY of Sciences and Belles Lettres, at Brussels.* This academy was founded in the year 1773; and several volumes of their memoirs have been published.

*Royal ACADEMY of Sciences*, at Stockholm, was instituted in 1739, and since that time it has published about sixty volumes of transactions, quarterly, in 8vo, in the Swedish language.

For an account of the Royal Society of London, and several other similar institutions, see the words *Journal*, *Society*, &c.

*American ACADEMY of Arts and Sciences*, was established in 1780 by the council and house of representatives in the province of Massachusetts Bay, for promoting the knowledge of the antiquities of America, and of the natural history of the country; for determining the uses to which its various natural productions may be applied; for encouraging medicinal discoveries, mathematical disquisitions, philosophical enquiries and experiments, astronomical, meteorological, and geographical observations, and improvements in agriculture, manufactures, and commerce; and, in short, for cultivating every art and science, which may tend to advance the interest, honour, dignity, and happiness, of a free, independent, and virtuous people. The members of this academy are never to be less than forty, nor more than two hundred.

ACADEMY is also used among us for a kind of collegiate school, or seminary; where youth are instructed in the liberal arts and sciences in a private way: now indeed it is used for all kinds of schools.

Frederic I, king of Prussia, established an academy at Berlin in 1703, for educating the young nobility of the court, suitable to their extraction. The expence of the students was very moderate, the king having undertaken to pay the extraordinaries. This illustrious school, which was then called the academy of princes, has now lost much of its first splendour.

The Romans had a kind of military academies established in all the cities of Italy, under the name of *Campi Martis*. Here the youth were admitted to be trained for war at the public expence. And the Greeks, besides academies of this kind, had military professors, called *Tactici*, who taught all the higher offices of war, &c.

We have two royal academies of this kind in England, the expences of which are defrayed by the government; the one at Woolwich, for the artillery and military engineers; and the other at Portsmouth, for the navy. The former was established by his late majesty.



majesty king George II, by warrants dated April the 30th and November the 18th, 1741, for instructing persons belonging to the military part of the ordnance, in the several branches of mathematics, fortification, &c, proper to qualify them for the service of artillery and the office of engineers. This institution is under the direction of the master-general and board of ordnance for the time being: and at first the lectures of the masters in the academy were attended by the practitioner-engineers, with the officers, serjeants, corporals, and private men of the artillery, besides the cadets. At present however none are educated there but the gentlemen cadets, to the number of 90 or 100, where they receive an education perhaps not to be obtained or purchased for money in any part of the world. The master-general of the ordnance is always captain of the cadets' company, and governor of the academy; under him are a lieutenant-governor, and an inspector of studies. The masters have been gradually increased, from two or three at first, now to the number of twelve, namely, a professor of mathematics, and two other mathematical masters, a professor of fortification, and an assistant, two drawing masters, two French masters, with masters for fencing, dancing, and chemistry. This institution is of the greatest consequence to the state, and it is hardly credible that so important an object should be accomplished at so trifling an expence. It is to be lamented however that the academy is fixed in so unhealthy a situation; that the lecture rooms and cadets' barracks are so small as to be insufficient for the purposes of the institution; and that the salaries of the professors and masters should be so inadequate to their labours, and the benefit of their services.

The Royal Naval Academy at Portsmouth was founded by George I, in 1722, for instructing young gentlemen in the sciences useful for navigation, to breed officers for the royal navy. The establishment is under the direction of the board of admiralty, who give salaries to two masters, by one of whom the students are boarded and lodged, the expence of which is defrayed by their own friends, nothing being supplied by the government but their education.

ACANTHUS, in *Architecture*, the leaves of a plant which forms the ornament of the capital of the Corinthian order. Vitruvius ascribes the use of it to the following accident. A young girl dying, her nurse was desirous of consecrating to her manes certain toys which she was fond of in her life-time; which the good woman carried in a little basket, covered with a square tile, and placed it among some green plants which grew on her grave. One of these, which happened to be the acanthus, as it grew up, invironed and in a manner embraced the basket; which Callimachus, a noted Greek sculptor, casting his eyes upon, from thence took the hint of this elegant ornament. See ABACUS.

ACCELERATED *Motion*, is that which receives fresh accessions of velocity; and the acceleration may be either equably or unequably: if the accessions of velocity be always equal in equal times, the motion is said to be equably or uniformly accelerated; but if the

accessions, in equal times, either increase or decrease, then the motion is unequably or variably accelerated.

Acceleration is directly opposite to retardation, which denotes a diminution of velocity.

ACCELERATION comes chiefly under consideration in physics, in the descent of heavy bodies, tending or falling towards the centre of the earth, by the force of gravity.

That bodies are accelerated in their natural descent, is evident both to the sight, and from observing that the greater height they fall from, the greater force they strike with, and the deeper impressions they make in soft substances.

The acceleration of falling bodies has been ascribed to various causes, by different philosophers. Some have attributed it to the pressure of the air downwards: the more a body descends, the longer and heavier, say they, must be the column of atmosphere incumbent upon it; to which they add, that the whole mass of fluid pressing by an infinity of right-lines all ultimately meeting in the earth's centre, such central point must support, as it were, the pressure of the whole mass; and that consequently the nearer a body approaches to it, the more must it receive of the pressure of a multitude of lines tending to unite in the central point.

Mr. Hobbes endeavours to account for this acceleration from a new impression of the cause which makes bodies fall; in which he is so far right. But then he as far mistakes, as to the cause of the fall, which he thinks is the air: at the same time, says he, that one particle of air ascends, another descends; for in consequence of the earth's motion being two-fold, that is circular and progressive, the air must at once both ascend and circulate; whence it follows, that a body falling in this medium, and receiving a new pressure every instant, must have its motion accelerated.

But to both these systems it may be answered, that the air is quite out of the question; for it is very evident that bodies fall, and in falling have their motion accelerated, in *vacuo*, as in open air, and even more than in the air, in as much as this opposes and somewhat retards their fall.

The Gassendists assign another reason for the acceleration: they pretend that there are continually issuing out of the earth certain attractive corpuscles, directed in an infinite number of rays; those, say they, ascend and then descend, in such sort that the nearer a body approaches to the earth's centre, the more of these attractive rays press upon it, in consequence of which its motion becomes more accelerated.

The peripatetics endeavour to explain the matter thus: the motion of heavy bodies downward, arises, say they, out of an intrinsic principle that causes a tendency in them to the centre, as the place appropriated to their element; where, when they can once arrive, they will be at perfect rest; and therefore, continue they, the nearer bodies approach to it, the more the velocity of their motion is increased: a notion too idle to merit confutation.

The Cartesians account for acceleration, by reiterated impulses of their *materia subtilis*, acting continually



on falling bodies, and propelling them downwards: a conceit equally unintelligible and absurd with the former.

But, leaving all such visionary causes of acceleration, and only admitting the existence of such a force as gravity, so evidently inherent in all bodies, without regard to what may be the cause of it, the whole mystery of motion will be cleared up. Consider gravity then, Galileo, only as a cause or force which acts constantly on heavy bodies; and it will be easy to conclude that the principle of gravitation, which determines bodies to descend, must by a necessary consequence accelerate them in falling.

A body then having once begun to descend, through the impulse of gravity; the state of descending is now, by Newton's first law of nature, become as it were natural to it; inasmuch that, were it left to itself, it would for ever continue to descend, even though the first cause of its descent should cease. But besides this determination to descend, impressed upon it by the first cause of motion, which would be sufficient to continue to infinity the degree of motion already begun, new impulses are continually superadded by the same cause; which continues to act upon the body already in motion, in the same manner as if it had remained at rest. There being then two causes of motion, acting both in the same direction; it necessarily follows, that the motion which they unitedly produce, must be more considerable than what either could produce separately. And as long as the velocity is thus continued, the same cause still subsisting to increase it more, the descent must of necessity be continually accelerated.

Supposing then that gravity, from whatever principle it arises, acts uniformly upon all bodies at the same distance from the centre of the earth: dividing the time which the heavy body takes up in falling to the earth, into indefinitely small equal parts; gravity will impel the body toward the centre of the earth, in the first indefinitely short instant of the descent. If after this we suppose the action of gravity to cease, the body will continue perpetually to advance uniformly toward the earth's centre, with an indefinitely small velocity, equal to that which resulted from the first impulse.


But then if we suppose that the action of gravity still continues the same after the first impulse; in the second instant, the body will receive a new impulse toward the earth, equal to that which it received in the first instant; and consequently its velocity will be doubled; in the third instant, it will be tripled; in the fourth, quadrupled; in the fifth, quintupled; and so on continually: for the impulse made in any preceding instant, is no ways altered by that which is made in the following one; but they are, on the contrary, always accumulated on each other.

So that the instants of time being supposed indefinitely small, and all equal, the velocity acquired by the falling body, will be, in every instant, proportional to the times from the beginning of the descent; and consequently the velocity will be proportional to the time in which it is produced. So that if a body, by this constant force, acquire a velocity of  $16\frac{1}{2}$  feet suppose in one second of time; it will acquire a velocity

of  $32\frac{1}{2}$  feet in two seconds,  $48\frac{1}{2}$  feet in 3 seconds,  $64\frac{1}{2}$  in 4 seconds, and so on. Nor ought it to seem strange that all bodies, small or large, acquire, by the force of gravity, the same velocity in the same time. For every equal particle of matter being endued with an equal impelling force, namely its gravity or weight, the sum of all the forces, in any compound mass of matter, will be proportional to the sum of all the weights, or quantities of matter to be moved; consequently, the forces and masses moved, being thus constantly increased in the same proportion, the velocities generated will be the same in all bodies, great or small. That is, a double force moves a double mass of matter, with the same velocity that the single force moves the single mass; and so on. Or otherwise, the whole compound mass falls all together with the same velocity, and in the same manner, as if its particles were not united, but as if each fell by itself, separated all from one another. And thus all being let go at once, they would fall together, just as if they were united into one mass.

The foregoing law of the descent of falling bodies, namely that the velocities are always proportional to the times of descent, as well as the following laws concerning the spaces passed over, &c, were first discovered and taught by the great Galileo, and that nearly in the following manner.

Because the constant velocity with which any body moves, or the space it passes over in a given time, as suppose one second, being multiplied by the time, or number of seconds it is in motion, expresses the space passed over in that time; and the area or space of a rectangular figure being denoted by the length multiplied by the breadth; therefore the space so run over, may be considered as a rectangle compounded of the time and velocity, that is a rectangle of which the time denotes the length, and the velocity the breadth. Suppose then A to be the heavy body which descends, and A B to denote the whole time of any descent; which




let be divided into a certain number of equal parts, denoting intervals or portions of the given time, as A C, C D, D E, &c. Imagine the body to descend, during the time expressed by the first of the divisions A C, with a certain uniform velocity arising from the force of gravity acting on it, which let be denoted by A F, the breadth of the rectangle C F; then the space run through during the time denoted by A C, with the velocity denoted by A F, will be expressed by the rectangular space C F.

Now the action of gravity having produced, in the first moment, the velocity A F, in the body, before at rest; in the first two moments it will produce the velocity C G, the double of the former; in the third moment, to the velocity C G will be added one degree more,



more, by which means will be produced the velocity  $DH$ , triple of the first; and so of the rest; so that during the whole time  $AB$ , the body will have acquired the velocity  $BK$ . Hence, taking the divisions of the line  $AB$  at pleasure; for example, the divisions  $AC$ ,  $CD$ , &c, for the times; the spaces run through during those times, will be as the areas or rectangles  $CF$ ,  $DG$ , &c; and so the space described by the moving body during the whole time  $AB$ , will be equal to all the rectangles, that is, equal to the whole indented space  $ABKIHGF$ . And thus it will happen if the increments of velocity be produced, as we may say, all at once, at the end of certain portions of finite time; for instance at  $C$ , at  $D$ , &c; so that the degree of motion remains the same to the instant that a new acceleration takes place.

By conceiving the divisions of time to be shorter, for example but half as long as the former, the indentures of the figure will be proportionably more contracted, and it will approach nearer to a triangle; and so much the nearer as the divisions of time are shorter: and if these be supposed infinitely small; that is, if increments of the velocity be supposed to be acquired continually, and at each indivisible particle of time, which is really the case, the rectangles so successively produced, will form a true triangle, as  $ABC$ ; the whole time  $AB$  consisting of minute portions  $A1$ ,



$12$ ,  $23$ , &c; and the area of the triangle  $ABC$ , of all the minute surfaces, or minute trapeziums, which answer to the divisions of the times; the area of the whole triangle  $ABC$ , denoting the space run through during the whole time  $AB$ ; and the area of any smaller triangle  $A7g$ , denoting the space run through during the corresponding time  $A7$ . But the triangles  $A1a$ ,  $A7g$ , &c, being similar, have their areas to each other as the squares of their like sides  $A1$ ,  $A7$ , &c; and consequently the spaces gone through, in any times counted from the beginning, are to each other as the squares of the times.

Hence, in any right-angled triangle, as  $ABC$ , the one side  $AB$  represents the time, the other side  $BC$  the velocity acquired in that time, and the area of the triangle the space described by the falling body.

From the preceding demonstration is also drawn this other general theorem in motions that are uniformly accelerated; namely, that a body descending with a uniformly accelerated motion, describes in the whole time of its descent, a space, which is exactly the half of that which it would describe uniformly in the same time, with the velocity it has acquired at the end of its accelerated fall. For it has been shewn that the whole space which the falling body has run through in the time  $AB$ , is represented by the triangle  $ABC$ , the last velocity being  $BC$ ; and the space which the

body would run through uniformly in the same time  $AB$ , constantly with the said greatest velocity  $BC$ , is represented by the rectangle  $ABCD$ ; but it is well known that the rectangle  $ABCD$  is double the triangle  $ABC$ ; and therefore the latter space run through, is double the former; that is, the space run through by the accelerated motion, is just half of that which the body would describe in the same time, moving uniformly with the velocity acquired at the end of its accelerated fall.

Hence then, from the foregoing considerations are deduced the following general laws of uniformly accelerated motions, namely,

1st. That the velocities acquired, are constantly proportional to the times; in a double time a double velocity, &c.

2d. That the spaces described in the whole times, each counted from the commencement of the motion, are proportional to the squares of the times, or to the squares of the velocities; that is, in twice the time, the body will describe 4 times the space; in thrice the time, it will describe 9 times the space; in quadruple the time, 16 times the space; and so on. In short, if the times are proportional

to the numbers - 1, 2, 3, 4, 5, &c,

the spaces will be as 1, 4, 9, 16, 25, &c,

which are the squares of the former. So that if a body, by the natural force of gravity, fall through the space of  $16\frac{1}{2}$  feet in the first second of time; then in the first two seconds of time it will fall through four times as much, or  $64\frac{1}{2}$  feet; in the first three seconds it will fall nine times as much, or  $144\frac{1}{2}$  feet; and so on. And as the spaces fallen through are as the squares of the times, or of the velocities; therefore the times, or the velocities, are proportional to the square roots of the spaces.

3d. The spaces described by falling bodies, in a series of equal instants or intervals of time, will be as the odd numbers - 1, 3, 5, 7, 9, &c, which are the differences of }  
the squares or whole spaces } 1, 4, 9, 16, 25, &c,  
that is, the body which has run through  $16\frac{1}{2}$  feet in the first second, will in the next second run through  $48\frac{1}{2}$  feet, in the third second  $80\frac{1}{2}$ , and so on.

4th. If the body fall through any space in any time, it acquires a velocity equal to double that space; that is, in an equal time, with the last velocity acquired, if uniformly continued, it would pass over just double the space. So if a body fall through  $16\frac{1}{2}$  feet in the first second of time, then it has acquired a velocity of  $32\frac{1}{2}$  feet in a second; that is, if the body move uniformly for one second, with the velocity acquired, it will pass over  $32\frac{1}{2}$  feet in this one second: and if in any time the body fall through 100 feet; then in another equal time, if it move uniformly with the velocity last acquired, it will pass over 200 feet. And so on.

But, as the method of demonstration used by Galileo, by means of infinitely small parts forming a regular triangle, is not approved of by many persons, the same laws may be otherwise demonstrated thus: let the whole time of a body's free descent be divided into any number of parts, calling each of these parts 1; and let  $a$  denote the velocity acquired at the end of the first



part of time; then will  $2a$ ,  $3a$ ,  $4a$ , &c, represent the velocities at the end of the 2d, 3d, 4th, &c, part of time, because the velocities are as the times; and for the same reason  $\frac{1}{2}a$ ,  $\frac{3}{2}a$ ,  $\frac{5}{2}a$ , &c; will be the velocities at the middle point of the first, second, third, &c, part of time. But now as the velocities increase uniformly, the space described in any one of these parts of time, may be considered as uniformly described with its middle velocity, or the velocity in the middle of that part of time; and therefore multiplying those mean velocities each by their common time 1, we have the same fractions  $\frac{1}{2}a$ ,  $\frac{3}{2}a$ ,  $\frac{5}{2}a$ , &c, for the spaces passed over in the successive parts of the time; that is, the space  $\frac{1}{2}a$  in the first time,  $\frac{3}{2}a$  in the second,  $\frac{5}{2}a$  in the third, and so on: then add these spaces successively to one another, and we obtain  $\frac{1}{2}a$ ,  $\frac{4}{2}a$ ,  $\frac{9}{2}a$ ,  $\frac{16}{2}a$ , &c, for the whole spaces described from the beginning of the motion to the end of the first, second, third, &c, portion of time; namely  $\frac{1}{2}a$  space in one time,  $\frac{4}{2}a$  in 2 times,  $\frac{9}{2}a$  in 3 times, and so on: and it is evident that these spaces are as the numbers 1, 4, 9, 16, &c, which are as the squares of the times.

And from this mode of demonstration, all the properties above-mentioned evidently flow; such as that the whole spaces -  $\frac{1}{2}a$ ,  $\frac{4}{2}a$ ,  $\frac{9}{2}a$ , &c, are as the squares of the times - 1, 2, 3, &c, that the separate spaces -  $\frac{1}{2}a$ ,  $\frac{3}{2}a$ ,  $\frac{5}{2}a$ , &c, described in the successive times, } 1, 3, 5, &c, are as the odd numbers - and that the velocity  $a$  acquired in any time 1, is double the space -  $\frac{1}{2}a$  described in the same time.

As the laws of acceleration are very important, I shall here insert the two following propositions, sent me by my learned friend Mr. Abram Robertson, of Christ Church College Oxford, in which those laws are demonstrated in a manner somewhat different.

“ PROPOSITION I.

If from the point P in the straight line A B, the points M, N begin to move at the same time, namely, M towards A with a motion uniformly retarded, and N from rest towards B with a motion uniformly accelerated; and if the velocity of M decreases as much as the velocity of N increases in the same time; then the space M N is generated by an uniform motion, equal to the velocity with which M begins to move.



For, by hypothesis, whatever is lost in the velocity of M by retardation, is added to the velocity of N by acceleration: the joint velocities, therefore, of M and N must always be equal. But it is by the joint velocities of M and N that the space M N is generated. Consequently M N is generated by an uniform motion, which is evidently equal to the velocity with which M begins to move.

“ PROPOSITION II.


If a point begins to move in the direction of a straight line, and continues to move in the same di-

rection with a velocity uniformly accelerated; the space passed over in any given time, will be equal to half the space passed over in the same time with the velocity with which the acceleration ends.

Let the point D begin to move from A towards B, along the straight line A B, with a motion uniformly accelerated; the space A D passed over, is equal to half the space which the point would pass over, in the same time with the acquired velocity at D.



Let the points M, N begin to move in the straight line G H, at the same time, with equal velocities uniformly accelerated; M beginning to move from G, and N from P; and at the same time that M comes to the point P, let N come to H. Then as M and N




move with equal velocities, uniformly accelerated, it is evident that the spaces, which they pass over in the same time, are equal to one another; consequently the space G P is equal to the space P H. Now as M begins to move from G with a velocity uniformly accelerated, it will arrive at P with an acquired velocity. Hence it is evident, if it be supposed to begin to move from P with this acquired velocity, and proceed toward G with a velocity uniformly retarded in the same degree that it was accelerated when it began to move from G, that it will pass over the same space G P in the same time. Wherefore, supposing the two points M, N to begin to move from P at the same time, namely the point M beginning to move with the acquired velocity mentioned above, and proceeding towards G with the velocity uniformly retarded, described above; and the point N as before with the velocity uniformly accelerated: then as the acceleration and retardation are uniform, they will be equal in equal spaces of time. Again, as M is retarded in the same degree that it was accelerated when it began to move from G, that is, in the same degree that N is accelerated, by the former prop. M N is generated by an uniform velocity. But when the point M arrives at G, its velocity becomes equal to 0 or nothing; and at the time that M arrives at G, N arrives at H with the acquired velocity. Wherefore, as the velocities of M and N taken jointly are equal, and consequently uniform, the space G H is passed over with the velocity of N at H, in the same time that P H is passed over by N beginning to move from P with a velocity uniformly accelerated to H. But P H is half of G H. “ Hence the prop. is manifest.”

And hence the other laws of the spaces, before-mentioned, easily follow.

Since the spaces descended are as the squares of the times, and the abscissas of a parabola are as the squares of the ordinates, therefore the relation of the times and spaces descended may be very well represented by the ordinates and abscissas of that figure. Thus if A B be the axis of the parabola A b d f h, and A C a tangent at





at the vertex perpendicular to the axis, divided into any number of equal parts  $Aa, ac, ce, &c$ , for the times; and if there be drawn  $ab, cd, ef, &c$ , parallel to the axis: hence if  $ab$  be the space descended in the time  $Aa$ , then  $cd$  will be the space descended in the time  $Ac$ , and  $ef$  the space de-

scended in the time  $Ae$ , and so on continually.

From the properties above-demonstrated, are derived the following practical formulas or theorems for use. Namely, if  $g$  denote the space passed over in the first second of time, by a body urged by any constant force, denoted by  $1$ , and  $t$  denote the time or number of seconds in which the body passes over any other space  $s$ , and  $v$  the velocity acquired at the end of that time; then from the foregoing laws we have  $v = 2gt$ , and  $s = gt^2$ ; and from these two equations result the following general formulas:

$$t = \frac{v}{2g} = \frac{2s}{v} = \sqrt{\frac{s}{g}},$$

$$v = 2gt = 2\sqrt{gs} = \frac{2s}{t},$$

$$s = gt^2 = \frac{vv}{4g} = \frac{tv}{2},$$

$$g = \frac{s}{t^2} = \frac{v}{2t} = \frac{vv}{4s}.$$

And here, when the constant force  $1$ , is the natural force of gravity, then the distance  $g$  descended in the first second, in the latitude of London, is  $16\frac{1}{2}$  feet: but if it be any other constant force, the value of  $g$  will be different, in proportion as the force is more or less.

The motion of an ascending body, or of one that is impelled upwards, is diminished or retarded by the same principle of gravity, acting in a contrary direction, after the same manner that a falling body is accelerated.

A body projected upwards, ascends until it has lost all its motion; which it does in the same space of time, that the body would have taken up in acquiring, by falling, a velocity equal to that with which the falling body began to be projected upwards. And consequently the heights to which bodies ascend, when projected upwards with different velocities, are to each other as the squares of those velocities.

**ACCELERATED Motion of Bodies on Inclined Planes.** The same general laws obtain here, as in bodies falling freely, or perpendicularly; namely, that the velocities are as the times, and the spaces descended down the planes as the squares of the times, or of the velocities. But those velocities are less, according to the sine of the plane's inclination; and the spaces less, according to the square of the sine. See *INCLINED Plane*.

**ACCELERATED Motion of Pendulums.** See *PENDULUM*.

**ACCELERATED Motion of Projectiles.** See *PROJECTILE*.


**ACCELERATED Motion of Compressed Bodies,** in ex-

panding or restoring themselves. See *DILATATION, COMPRESSION, and ELASTICITY*.

**ACCELERATING FORCE**, in Physics, is the force that accelerates the motion or velocity of bodies; and it is equal to, or expressed by, the quotient arising from the motive or absolute force, divided by the mass or weight of the body that is moved. In treating of physical considerations respecting forces, velocities, times, and spaces gone over, the first inquiry is the accelerating or accelerative force. This force is greater or less in proportion to the velocity it generates in the same time, and by this velocity it is measured. All accelerating forces are equal, and generate equal velocities, that have the motive forces directly proportional to the quantities of matter: so a double motive force will move a double quantity of matter with the same velocity, as also a triple motive force a triple quantity, a quadruple force a quadruple quantity, &c, all with the same velocity. And this is the reason why all bodies fall equally swift by the force of gravity; for the motive force is exactly proportional to their weight or mass. In general, the accelerating force is in the direct ratio of the motive force, and inverse ratio of the quantity of matter. When a body is let fall freely, to descend by the force of its natural gravity, it has been found by experiment that it falls through  $16\frac{1}{2}$  feet in one second of time, and requires a velocity of  $32\frac{1}{2}$  feet in that time: but if the quantity of matter be doubled, and the motive force remain the same as before, by connecting the falling body to another of equal weight by means of a thread, this other body being laid on a horizontal plane, and the falling body hanging down off the plane, and drawing the other equal body along the plane after it; then the accelerating force will be only half of what it was before, and the space fallen in one second will be only  $8\frac{1}{2}$  feet, and the velocity acquired  $16\frac{1}{2}$ : and if the quantity of matter be tripled, or the body drawn along the plane doubled; then the accelerating force will be only one-third of what it was at first, and the space descended in one second, and velocity acquired, each one-third of the first: and so on.

But accelerating forces are sometimes variable, as well as sometimes constant; and the variation may be either increasing or decreasing.

The nature of constant and variable accelerating forces, may be illustrated in the following manner. Let two weights  $W, w$ , be connected by a thread




passing over a pulley at  $A, B$ , or  $C$ ; and let the weight  $W$  descend perpendicularly down, while it draws the smaller weight  $w$  up the line  $AD$ , or  $BE$ , or  $CF$ , the first.



first being a straight inclined plane, and the other two curves, the one convex and the other concave to the perpendicular. Then the small weight  $w$  will always make some certain resistance to the free descent of the large weight  $W$ , and that resistance will be constantly the same in every part of the plane  $AD$ , the difficulty to draw it up being the same in every point of it, because every part of it has the same inclination to the horizon, or to the perpendicular; and consequently the accessions to the velocity of the descending weight  $W$ , will be always equal in equal times; that is, in this case  $W$  descends by a uniformly accelerating force. But in the two curves  $BE$ ,  $CF$ , the resistance or opposition of the small weight  $w$  will be constantly altering as it is drawn up the curves, because every part of them has a different inclination to the horizon, or to the perpendicular: in the former curve, the direction becomes more and more upright, or nearer perpendicular, as the small weight  $w$  ascends, and the opposition it makes to the descent of  $W$ , becomes more and more; and consequently the accessions to the velocity of  $W$  will be always less and less in equal times; that is,  $W$  descends by a decreasing accelerating force: but in the latter curve  $CF$ , as  $w$  ascends, the direction of the curve becomes less and less upright, and the opposition it makes to the descent of  $W$ , becomes always less and less; and consequently the accessions to the velocity of  $W$  will be always more and more in equal times; that is,  $W$  descends by an increasing accelerating force. So that although the velocity continually increases in all these cases, yet whilst it increases in a constant ratio to the times of motion, in the plane  $AD$ ; the velocity increases in a less ratio than the time it ascended up  $BE$ , and in a greater ratio than the time increases in the other curve  $CF$ .

Now the relations between the times and velocities in all these cases, may be very well represented by the relations between the abscisses and ordinates of certain lines. Thus let  $AB$  and  $AC$  be two straight lines,



making any angle  $BAC$ ; and  $AD$ ,  $AE$  two curves, the former concave, and the latter convex towards  $AB$ : divide  $AB$  into any parts  $Aa$ ,  $Ab$ , &c, representing the times of motion; and draw the perpendiculars  $acd$ ,  $bfg$ , &c, representing the velocities. Then in the right line  $AC$ , the ordinates  $ad$ ,  $bg$ , being as the abscisses  $Aa$ ,  $Ab$ , this represents the case of uniformly accelerated motion, in which the velocities are always as the times: but in the curve  $AD$ , the ordinates  $ac$ ,  $bf$  increase in a less ratio than the abscisses  $Aa$ ,  $Ab$ ; and therefore this represents the case of decreasing acceleration, in which the velocities increase

in a less ratio than the times: and in the other curve  $AE$ , the ordinates  $ae$ ,  $bb$  increase in a greater ratio than the abscisses; and therefore this represents the case of increasing acceleration, in which the velocities increase in a greater ratio than the times.

The several algebraic formulas or theorems, respecting the time, velocity, space, for constant accelerating forces, are delivered above, at the article *Accelerated Motion*, where the value of each circumstance is expressed in finite determinate quantities. But in the cases of variably accelerated motions, the formulas will require the help of the method of fluxions to express, not those general relations themselves, but the fluxions of them; and consequently, taking the fluents of those expressions, in particular cases, the relations of time, space, velocity, &c, are obtained.

Now if  $t$  denote the time in motion,

$v$  the velocity generated by any force,

$s$  the space passed over,

and  $2g$  the variable force at any part of the motion, or the velocity the force would generate in one second of time, if it should continue invariable, like the force of gravity, during that one second; and therefore the value of this velocity  $2g$ , will be in proportion to  $32\frac{1}{8}$  feet, as that variable force, is to 1 the force of gravity. Then because the force may be supposed constant during the indefinitely small time  $t$ , and that in uniform motions the spaces and velocities are proportional to the times, we from thence obtain these two general fundamental proportions,

$$v : s :: 1'' : t, \text{ or } s = t v;$$

$$2g : v :: 1'' : t, \text{ or } v = 2gt.$$

From which are derived the four formulas below, in which the value of each quantity is expressed in terms of the rest.

$$1. \quad t = \frac{s}{v} = \frac{v}{2g}.$$

$$2. \quad v = 2gt = \frac{2gs}{v}.$$

$$3. \quad s = vt = \frac{v^2}{2g}.$$

$$4. \quad 2g = \frac{v}{t} = \frac{v^2}{s}.$$

And these theorems equally hold good for the destruction of motion and velocity, by means of retarding forces, as for the generation of the same by means of accelerating forces.

ACCELERATION, in *Mechanics*, the increase of velocity in a moving body.

ACCELERATION. *Astron.* The Diurnal Acceleration of the fixed stars, is the time which the stars, in one diurnal revolution, anticipate the mean diurnal revolution of the sun; which is  $3^m 55^s \frac{2}{3}$  of mean time, or nearly  $3^m 56^s$ : that is, a star rises, or sets, or passes the meridian, about  $3^m 56^s$  sooner each day. This acceleration of the stars, which is only apparent in them, arises from the real retardation of the sun, owing to his apparent



rent motion in his orbit towards the east, which is about  $59' 8'' \frac{2}{5}$  of a degree every day. So that the star which passed the meridian yesterday at the same moment with the sun, is to-day about  $59' 8''$  past the meridian to the west, when the sun arrives at it; which will take him up about  $3^m 56^s$  of time to pass over; and therefore the star passes by  $3^m 56^s$  sooner than the sun each day, or anticipates his motion at that rate. The true quantity of this anticipation, or acceleration, is found by this proportion,  $360^\circ 59' 8'' \frac{1}{5} : 59' 8'' \frac{1}{5} :: 24 \text{ hours} : 3^m 55^s \frac{2}{5}$ , the fourth term of which is the acceleration.

The diurnal acceleration serves to regulate the lengths or vibration of pendulums. If I observe a fixed star set or pass behind a hill, steeple, or such like, when the pendulum marks for instance  $8^h 10^m$ ; and the next day, the eye being in the same place as before, the passage be at  $8^h 6^m 4^s$ ; I thence conclude that the pendulum is well regulated, or truly measures mean time.

**ACCELERATION of a Planet.** A planet is said to be accelerated in its motion, when its real diurnal motion exceeds its mean diurnal motion. And, on the other hand, the planet is said to be retarded in its motion, when the mean exceeds the real diurnal motion. This inequality arises from the change in the distance of the planet from the sun, which is continually varying; the planet moving always quicker in its orbit when nearer the sun, and slower when farther off.

**ACCELERATION of the Moon,** is a term used to express the increase of the moon's mean motion from the sun, compared with the diurnal motion of the earth; by which it appears that, from some uncertain cause, it is now a little quicker than it was formerly. Dr. Halley was led to the discovery, or suspicion, of this acceleration, by comparing the ancient eclipses observed at Babylon, &c, and those observed by Albategnius in the ninth century, with some of his own time; as may be seen in N<sup>o</sup> 218 of the Philosophical Transactions. He could not however ascertain the quantity of the acceleration, because the longitudes of Bagdat, Alexandria, and Aleppo, where the observations were made, had not been accurately determined. But since his time the longitude of Alexandria has been ascertained by Chazelles; and Babylon, according to Ptolemy's account, lies  $50'$  east of Alexandria. From these data, Mr. Dunthorne, vol. 46 Philof. Transactions, compared the recorded times of several ancient and modern eclipses, with the calculations of them by his own tables, and thereby verified the suspicion that had been started by Dr. Halley; for he found that the same tables gave the moon's place more backward than her true place in ancient eclipses, and more forward than her true place in later eclipses; and thence he justly inferred that her motion in ancient times was slower, and in later times quicker, than the tables give it.

Not content however with barely ascertaining the fact, he proceeded to determine, as well as the observations would allow, the quantity of the acceleration; and by means of the most authentic eclipse, of which any good account remains, observed at Babylon in the year 721 before Christ, he found that the observed beginning of this eclipse was about an hour and three quarters sooner than the beginning by the tables; and that therefore the moon's true place preceded her place by computation by about  $50'$  of a degree at that time.

Then admitting the acceleration to be uniform, and the aggregate of it as the square of the time, it will be at the rate of about  $10''$  in 100 years.

Dr. Long, vol. ii. p. 436 of his Astronomy, enumerates the following causes from some one or more of which the acceleration may arise. Either 1<sup>st</sup>, the annual and diurnal motion of the earth continuing the same, the moon is really carried about the earth with a greater velocity than formerly: or, 2<sup>dly</sup>, the diurnal motion of the earth, and the periodical revolution of the moon, continuing the same, the annual motion of the earth about the sun is retarded; which makes the sun's apparent motion in the ecliptic a little slower than formerly; and consequently the moon, in passing from any conjunction with the sun, takes up a less time before she again overtakes the sun, and forms a subsequent conjunction: in both these cases, the motion of the moon from the sun is really accelerated, and the synodical month actually shortened: or, 3<sup>dly</sup>, the annual motion of the earth, and the periodical revolution of the moon, continuing the same, the rotation of the earth upon its axis is a little retarded; in this case, days, hours, minutes, &c, by which all periods of time must be measured, appear of a longer duration; and consequently the synodical month will appear to be shortened, though it really contain the same quantity of absolute time as it always did. If the quantity of matter in the body of the sun be lessened, by the particles of light continually streaming from it, the motion of the earth about the sun may become slower: if the earth increases in bulk, the motion of the moon about the earth may thereby be quickened.

**ACCELERATIVE FORCE, &c,** the same as ACCELERATING.

**ACCESSIBLE,** something that may be approached, or to which we can come. In Surveying, it is such a place as will admit of having a distance or length of ground measured from it; or such a height or depth as can be measured by actually applying a proper instrument to it. For the means of doing which, see ALTIMETRY, LONGIMETRY, or HEIGHTS-AND-DISTANCES.

**ACCIDENS, ACCIDENT, Philof.**

*Per* ACCIDENS is a term often used among philosophers, to denote what does not follow from the nature of a thing, but from some accidental quality of it: in this sense it stands opposed to *per se*, which denotes the nature and essence of a thing. Thus, fire is said to burn *per se*, or considered as fire, and not *per accidens*; but a piece of iron, though red-hot, only burns *per accidens*, by a quality accidental to it, and not considered as iron.

**ACCIDENTS,** in *Astrology*, denote the most extraordinary occurrences in the course of a person's life, either good or bad: such as a remarkable instance of good fortune, a signal deliverance, a great sickness, &c.

**ACCIDENTAL,** something that partakes of the nature of an accident; or that is indifferent, or not essential to its subject.—Thus whiteness is accidental to marble, and sensible heat to iron.

**ACCIDENTAL Colours,** so called by M. Buffon, are those which depend on the affections of the eye, in contradistinction to such as belong to light itself.


The impressions made upon the eye, by looking steadfastly on objects of a particular colour, are various according



according to the single colour, or assemblage of colours, in the object; and they continue for some time after the eye is withdrawn, and give a false colouring to other objects that are viewed during their continuance. M. Buffon has endeavoured to trace the connection between these accidental colours, and those that are natural, in a variety of instances. M. d'Arcy contrived a machine for measuring the duration of those impressions on the eye; and from the result of several trials he inferred, that the effect of the action of light on the eye continued about eight thirds of a minute.

The subject has also been considered by M. de la Hire, and M. Aepinus, &c. See Mem. Acad. Paris 1743, and 1765; Nov. Com. Petrop. vol. 10; also Dr. Priestley's Hist. of Discoveries relating to Vision, pa. 631.

**ACCIDENTAL Point**, in Perspective, is the point in which a right line drawn from the eye, parallel to another right line, cuts the picture or perspective plane.



Let  $AB$  be the line given to be put into perspective,  $CFD$  the picture or perspective plane, and  $E$  the eye: draw  $EF$  parallel to  $AB$ ; so shall  $F$  be the accidental point of the line  $AB$ , and indeed of all lines parallel to it, since only one parallel to them, namely  $EF$ , can be drawn from the same point  $E$ : and in the accidental point concur or meet the representations of all the parallels to  $AB$ , when produced.

It is called the accidental point, to distinguish it from the principal point, or point of view, where a line drawn from the eye perpendicular to the perspective plane, meets this plane, and which is the accidental point to all lines that are perpendicular to the same plane.

**ACCIDENTAL Dignities**, and **Debilities**, in *Astrology*, are certain casual dispositions, and affections, of the planets, by which they are supposed to be either strengthened, or weakened, by being in such a house of the figure.

**ACCLIVITY**, the slope or steepness of a line or plane inclined to the horizon, taken upwards; in contradistinction to *declivity*, which is taken downwards. So the ascent of a hill, is an *acclivity*: the descent of the same, a *declivity*.

Some writers on fortification use *acclivity* for *talus*: though more commonly the word *talus* is used to denote the slope, whether in ascending or descending.

**ACCOMPANYMENT**, in *Music*, denotes either the different parts of a piece of music for the different instruments, or the instruments themselves which accompany a voice, to sustain it, as well as to make the music more full.

The Accompaniment is used in recitative, as well as in song; on the stage, as well as in the choir, &c.

The ancients had likewise their accompaniments on the theatre; and they had even different kinds of instruments to accompany the chorus, from those which accompanied the actors in the recitation.

The accompaniment among the moderns, is often a different part, or melody, from the song it accompanies. But it is disputed whether it was so among the ancients.

Organists sometimes apply the word to several pipes which they occasionally touch to accompany the treble; as the drone, the flute, &c.

**ACCOMPT**. See **ACCOUNT**.

**ACCORD**, according to the modern French music, is the union of two or more sounds heard at the same time, and forming together a regular harmony.

They divide Accords into *perfect* and *imperfect*; and again into *consonances* and *dissonances*.

Accord is more commonly called **CONCORD**, which see.

**ACCORD** is also spoken of the state of an instrument, when its fixed sounds have among themselves all the justness that they ought to have.

**ACCOUNT**, or **ACCOMPT**, in *Arithmetic*, &c, a calculation or computation of the number or order of certain things; as the computation of time, &c.

There are various ways of accounting; as, by enumeration, or telling one by one; or by the rules of arithmetic, addition, subtraction, &c.

**ACCOUNT**, in *Chronology*, is nearly synonymous with *style*. Thus, we say the English, the foreign, the Julian, the Gregorian, the Old, or the New account, or style.

We account time by years, months, &c; the Greeks accounted it by olympiads; the Romans, by indictions, lustres, &c.

**ACHERNER**, or **ACHARNER**, in *Astronomy*, a star of the first magnitude in the southern extremity of the constellation Eridanus, marked  $\alpha$  by Bayer. Its longitude for 1761,  $\propto 11^{\circ} 55' 1''$ ; and latitude south  $59^{\circ} 22' 4''$ .

**ACHILLES**, a name given by the schools to the principal argument alleged by each sect of philosophers in behalf of their system. In this sense we say this is his Achilles; that is, his master-proof: alluding to the strength and importance of the hero Achilles among the Greeks.

Zeno's argument against motion is peculiarly termed Achilles. That philosopher made a comparison between the swiftness of Achilles, and the slowness of a tortoise, pretending that a very swift animal could never overtake a slow one that was before it, and that therefore there is no such thing as motion: for, said he, if the tortoise were one mile before Achilles, and the motion of Achilles 100 times swifter than that of the tortoise, yet he would never overtake it; and for this reason, namely, that while Achilles runs over the mile, the tortoise will creep over one hundredth part of a mile, and will be so much the foremost; again while Achilles runs over this  $\frac{1}{100}$ th part, the tortoise will creep over the 100th part of that  $\frac{1}{100}$ th part, and will still be this last part the foremost; and so on continually, according to an infinite series of 100th parts: from which he concluded that the swifter could never overtake the slower in any finite time, but that they must go on approaching



proaching to infinity. But this sophism lay in their considering as an infinite time, the sum of the infinite series of small times in which Achilles could run over the infinite series of spaces,  $1 + \frac{1}{100} + \frac{1}{10000} + \frac{1}{1000000} + \dots$  &c, not knowing that the sum of this infinite series is equal to the quantity  $1\frac{1}{99}$  of a mile, and that therefore Achilles will overtake the tortoise when the latter has crawled over  $\frac{1}{99}$ th of a mile.

ACHROMATIC, in *Optics*, without colour; a term which, it seems, was first used by M. de la Lande, in his astronomy, to denote telescopes of a new invention, contrived to remedy aberrations and colours. See *Aberration* and *Telescope*.

ACHROMATIC TELESCOPE, a singular species of refracting telescope, said to be invented by the late Mr. John Dollond, optician to the king, and since improved by his son Mr. Peter Dollond, and others.

Every ray of light passing obliquely from a rarer into a denser medium, changes its direction towards the perpendicular; and every ray passing obliquely from a denser into a rarer medium, changes its direction from the perpendicular. This bending of the ray, caused by the change of its direction, is called its refraction; and the quality of light which subjects it to this refraction, is called its refrangibility. Every ray of light, before it is refracted, is white, though it consists of a number of component rays, each of which is of a different colour. As soon as it is refracted, it is separated into its component rays, which, from that time, proceed diverging from each other, like rays from a centre: and this divergency is caused by the different refrangibility of the component rays, in such sort, that the more the original or component ray is refracted, the more will the compound rays diverge when the light is refracted by one given medium only.

From hence it has been concluded, that any two different mediums that can be made to produce equal refractions, will necessarily produce equal divergencies: whence it should also follow, that equal and contrary refractions should not only destroy each other, but that the divergency of the colours caused by one refraction, should be corrected by the other; and that to produce refraction that would not be affected by the different refrangibility of light, is impossible.

But Mr. Dollond has proved, by many experiments, that these conclusions are not well founded; from which experiments it appeared, that a ray of light, after equal and contrary refractions, was still spread into component rays differently coloured: in other words, that two different mediums may cause equal refraction, but different divergency; and equal divergency, with different refraction. It follows therefore that refraction may be produced, which is not affected by the different refrangibility of light. In other words, that, if the mediums be different, different refractions may be produced, though at the same time the divergency caused by one refraction shall be exactly counteracted by the divergency caused by the other; and so an object may be seen through mediums which, together, cause the rays to converge, without appearing of different colours.

This is the foundation of Mr. Dollond's improvement of refracting telescopes. By subsequent experiments he found, that different sorts of glass differed greatly in their refractive qualities, with respect to the divergency of colours. He found that crown glass causes the least diver-

gency, and white flint the most, when they are wrought into forms that produce equal refractions. He ground a piece of white flint glass into a wedge, whose angle was about 25 degrees; and a piece of crown glass to another, whose angle was about 29 degrees; and these he found refracted nearly alike, but that their divergency of colours was very different.

He then ground several other pieces of crown glass to wedges of different angles, till he got one that was equal, in the divergency it produced, to that of a wedge of flint glass of 25 degrees; so that when they were put together, in such a manner as to refract in contrary directions, the refracted light was perfectly free from colour. Then measuring the fractions of each wedge, he found that that of the white flint glass, was to that of the crown glass, nearly as two to three. And hence any two wedges, made of these two substances, and in this proportion, would, when applied together so as to refract in contrary directions, refract the light without any effect arising from the different refrangibility of the component rays.

Therefore, to make two spherical glasses that refract the light in contrary directions, one must be concave, and the other convex; and as the rays, after passing through both, must meet in a focus, the excess of the refraction must be in the convex one: and as the convex is to refract most, it appears from the experiment that it must be made of crown glass; and as the concave is to refract least, it must be made of white flint.

And farther, as the refractions of spherical glasses are in an inverse ratio of their focal distances, it follows that the focal distances of the two glasses should be in the ratio of the refractions of the wedges; for, being thus proportioned, every ray of light that passes through this combined glass, at whatever distance from its axis, will constantly be refracted by the difference between two contrary refractions, in the proportion required; and therefore the effect of the different refrangibility of light will be prevented.

The removal of this impediment, however, produced another: for the two glasses, which were thus combined, being segments of very deep spheres, the aberrations from the spherical surfaces became so considerable, as greatly to disturb the distinctness of the image. Yet considering that the surfaces of spherical glasses admit of great variations, though the focal distance be limited, and that by these variations their aberration might be made more or less at pleasure; Mr. Dollond plainly saw that it was possible to make the aberrating of any two glasses equal; and that, as in this case the refractions of the two glasses were contrary to each other, and their aberrations being equal, these would destroy each other.

Thus he obtained a perfect theory of making object glasses, to the apertures of which he could hardly perceive any limits: for if the practice could come up to the theory, they must admit of apertures of great extent, and consequently bear great magnifying powers.

The difficulties of the practice are, however, still very considerable. For first, the focal distances, as well as the particular surfaces, must be proportioned with the utmost accuracy to the densities and refracting powers of the glasses, which vary even in the same sort of glass, when made at different times. Secondly, there are four surfaces to be wrought perfectly spherical.



However, Mr. Dollond could construct refracting telescopes upon these principles, with such apertures and magnifying powers, under limited lengths, as greatly exceed any that were produced before, in forming the images of objects bright, distinct, and uninfected with colours about the edges, through the whole extent of a very large field or compass of view; of which he has given abundant and undeniable testimony. See TELESCOPE.

There has lately appeared in the Gentleman's Magazine (1790, pa. 890) a paper on the refracting telescope, by an author who signs *Veritus*, in which the invention is ascribed to another person, not heretofore mentioned; in these words: "As the invention has been claimed by M. Euler, M. Klingenstierna, and some other foreigners, we ought, for the honour of England, to assert our right, and give the merit of the discovery to whom it is due; and therefore, without farther preface, I shall observe, that the inventor was Chester More Hall, Esq. of More-hall, in Essex, who, about 1729, as appears by his papers, considering the different humours of the eye, imagined they were placed so as to correct the different refrangibility of light. He then conceived, that if he could find substances having such properties as he supposed these humours might possess, he should be enabled to construct an object glass that would shew objects colourless. After many experiments he had the good fortune to find those properties in two different sorts of glass, and making them disperse the rays of light in different directions, he succeeded. About 1733 he completed several achromatic object glasses (though he did not give them this name), that bore an aperture of more than  $2\frac{3}{4}$  inches, though the focal length did not exceed 20 inches; one of which is now in the possession of the Rev. Mr. Smith, of Charlotte Street, Rathbone Place.

This glass has been examined by several gentlemen of eminence and scientific abilities, and found to possess the properties of the present achromatic glasses.

Mr. Hall used to employ the working opticians to grind his lenses; at the same time he finished them with the radii of the surfaces, not only to correct the different refrangibility of rays, but also the aberration arising from the spherical figure of the lenses. Old Mr. Bais, who at that time lived in Bridewell precinct, was one of these working opticians, from whom Mr. Hall's invention seems to have been obtained.

In the trial at Westminster-hall about the patent for making achromatic telescopes, Mr. Hall was allowed to be the inventor; but Lord Mansfield observed, that "It was not the person that locked up his invention in his scrutoire that ought to profit by a patent for such an invention, but he who brought it forth for the benefit of the public." This, perhaps, might be said with some degree of justice, as Mr. Hall was a gentleman of property, and did not look to any pecuniary advantage from his discovery; and, consequently, it is very probable that he might not have an intention to make it generally known at that time.

That Mr. Ayscough, optician on Ludgate Hill, was in possession of one of Mr. Hall's achromatic telescopes in 1754, is a fact which at this time will not be disputed."

**ACHRONICAL**, or *Achronycal*. See **ACRONYCHAL**.

**ACOUSTICS**. This term, in physico-mathematical meaning, signifies the doctrine of hearing, and the art of assisting that sense by means of speaking trumpets, hearing trumpets, whispering galleries, and such like. See **STENTROPHONIC TUBE**.

Sturmius, in his Elements of Universal Mechanics, treating of Acoustics, after examining into the nature of sounds, describes the several parts of the external and internal ear, and their several uses and connexions with each other; and from thence deduces the mechanism of hearing: and lastly, he treats of the means of adding an intensity of force to the voice and other sounds; and explains the nature of echoes, otacoustic tubes, and speaking trumpets. See **SOUND**, **EAR**, **VOICE**, and **ECHO**.

Dr. Hook, in the preface to his Micrography, asserts that the lowest whisper, by certain means, may be heard at the distance of a furlong; and that he knew a way by which it is easy to hear any one speak through a wall of three feet thick; also that by means of an extended wire, sound may be conveyed to a very great distance, almost in an instant.

**ACRE**, from the Saxon *æcre*, or German *acker*, a field, of the Latin *ager*. It is a measure of land, containing, by the ordinance for measuring land, made in the 33d and 34th of Edward I, 160 perches or square poles of land; that is, 16 in length and 10 in breadth, or in that proportion: and as the statute length of a pole is  $5\frac{1}{2}$  yards, or  $16\frac{1}{4}$  feet, therefore the acre will contain 4840 square yards, or 43560 square feet. The chain with which land is commonly measured, and which was invented by Gunter, is 4 poles or 22 yards in length; and therefore the acre is just 10 square chains; that is, 10 chains in length and one in breadth, or in that proportion. Farther, as a mile contains 1760 yards, or 80 chains in length, therefore the square mile contains 640 acres.

The acre, in surveying, is divided into 4 roods, and the rood is 40 perches.

The French acre, *arpent*, is equal to  $1\frac{1}{4}$  English acre;

The Strasburg contains about  $\frac{1}{2}$  an English acre;

The Welch acre contains about 2 English acres;

The Irish acre contains 1 ac. 2 r.  $19\frac{2}{3}\frac{7}{8}$  p. English.

Sir William Petty, in his Political Arithmetic, reckons that England contains 39 million acres: but Dr. Greve shews, in the Philos. Trans. N° 330, that England contains not less than 46 million acres. Whence he infers that England is above 46 times as large as the province of Holland, which it is said contains but about one million of acres.

By a statute of the 31st of Elizabeth, it is ordained, that if any man erect a cottage, he shall annex four acres of land to it.

**ACRONYCHAL**, or **ACRONYCAL**, in *Astronomy*, is said of a star or planet, when it is opposite to the sun. It is from the Greek *ακρονυχος*, the point or extremity of night, because the star rose at sun-set, or the beginning of night, and set at sun-rise, or the end of night; and so it shone all the night.

The acronychal is one of the three Greek poetic risings and settings of the stars; and stands distinguished from Cosmical and Heliacal. And by means of which, for want of accurate instruments, and other observations, they might regulate the length of their year.

**ACROTERIA**,



**ACROTHERIA**, or **ACROTHERS**, in *Architecture*, small pedestals, usually without bases, placed on pediments, and serving to support statues.

Those at the extremities ought to be half the height of the tympanum; and that in the middle, according to Vitruvius, one eighth part more.

**ACROTHERIA** also are sometimes used to signify figures, whether of stone or metal, placed as ornaments or crownings, on the tops of temples, or other buildings.

It is also sometimes used to denote those sharp pinnacles or spiry battlements, that stand in ranges about flat buildings, with rails and balustrades.

**ACTION**, in *Mechanics* or *Physics*, a term used to denote, sometimes the effort which some body or power exerts against another body or power, and sometimes it denotes the effects resulting from such effort.

The Cartesians resolve all physical action into metaphysical. Bodies, according to them, do not act on one another; the action comes all immediately from the Deity; the motions of bodies, which seem to be the cause, being only the occasions of it.

It is one of the laws of nature, that action and reaction are always equal, and contrary to each other in their directions.

Action is either instantaneous or continued; that is, either by collision or percussion, or by pressure. These two sorts of action are heterogeneous quantities, and are not comparable, the smallest action by percussion exceeding the greatest action of pressure, as the smallest surface exceeds the longest line, or as the smallest solid exceeds the largest surface: thus, a man by a small blow with a hammer, will drive a wedge below the greatest ship on the stocks, or under any other weight; that is, the smallest percussion overcomes the pressure of the greatest weight. These actions then cannot be measured the one by the other, but each must have a measure of its own kind, like as solids must be measured by solids, and surfaces by surfaces: time being concerned in the one, but not in the other.

If a body be urged at the same time by equal and contrary actions, it will remain at rest. But if one of these actions be greater than its opposite, motion will ensue towards the part least urged.

The actions of bodies upon each other, in a space that is carried uniformly forward; are the same as if the space were at rest; and any powers or forces that act upon all bodies, so as to produce equal velocities in them in the same, or in parallel right lines, have no effect on their mutual actions, or relative motions. Thus the motion of bodies on board of a ship that is carried uniformly forward, are performed in the same manner as if the ship was at rest. And the motion of the earth about its axis has no effect on the actions of bodies and agents at its surface, except in so far as it is not uniform and rectilinear. In general, the actions of bodies upon each other, depend not on their absolute, but relative motion.

*Quantity of ACTION*, in *Mechanics*, a name given by M. de Maupertuis, in the Memoirs of the Academy of Sciences of Paris for 1744, and in those of Berlin for 1746, to the continual product of the mass of a body, by the space which it runs through, and by its celerity. He lays it down as a general law, that in the

changes made in the state of a body, the quantity of action necessary to produce such change is the least possible. This principle he applies to the investigation of the laws of refraction, and even the laws of rest, as he calls them; that is, of the equilibrium or equipollency of pressures; and even to the modes of acting of the Supreme Being. In this way Maupertuis attempts to connect the metaphysics of final causes with the fundamental truths of mechanics; to shew the dependence of the collision of both elastic and hard bodies, upon one and the same law, which before had always been referred to separate laws; and to reduce the laws of motion, and those of equilibrium, to one and the same principle.

But this quantity of motion, of Maupertuis, which is defined to be the product of the mass, the space passed over, and the celerity, comes to the same thing as the mass multiplied by the square of the velocity, when the space passed over is equal to that by which the velocity is measured; and so the quantity of force will be proportional to the mass multiplied by the square of the velocity; since the space is measured by the velocity continued for a certain time.

In the same year that Maupertuis communicated the idea of his principle, professor Euler, in the supplement to his book, intitled *Methodus inveniendi lineas curvas maximi vel minimi proprietate gaudentes*, demonstrates, that in the trajectories which bodies describe by central forces, the velocity multiplied by what the foreign mathematicians call the element of the curve, always makes a minimum: which Maupertuis considered as an application of his principle to the motion of the planets.

It appears from Maupertuis's Memoir of 1744, that it was his reflections on the laws of refractions, that led him to the theorem above mentioned. The principle which Fermat, and after him Leibnitz, made use of, in accounting for the laws of refraction, is sufficiently known. Those mathematicians pretended, that a particle of light, in its passage from one point to another, through two mediums, in each of which it moves with a different velocity, must do it in the shortest time possible: and from this principle they have demonstrated geometrically, that the particle cannot go from the one point to the other in a right line; but being arrived at the surface that separates the two mediums, it must alter its direction in such a manner, that the sine of its incidence shall be to the sine of its refraction, as its velocity in the first medium is to its velocity in the second: whence they deduced the well known law of the constant ratio of those sines.

This explanation, though very ingenious, is liable to this pressing difficulty, namely, that the particle must approach towards the perpendicular, in that medium where its velocity is the least, and which consequently resists it the most: which seems contrary to all the mechanical explanations of the refraction of bodies, that have hitherto been advanced, and of the refraction of light in particular.

Sir Isaac Newton's way of accounting for it, is the most satisfactory of any that has hitherto been offered, and gives a clear reason for the constant ratio of the sines, by ascribing the refraction to the attractive force of the mediums; from which it follows, that the densest



mediums, whose attraction is the strongest, should cause the ray to approach the perpendicular; a fact confirmed by experiment. But the attraction of the medium could not cause the ray to approach towards the perpendicular, without increasing its velocity; as may easily be demonstrated. Thus then, according to Newton, the refraction must be towards the perpendicular, when the velocity is increased: contrary to the law of Fermat and Leibnitz.

Maupertuis has attempted to reconcile Newton's explanation with metaphysical principles. Instead of supposing, as the aforesaid gentlemen do, that a particle of light proceeds from one point to another in the shortest time possible; he contends that a particle of light passes from one point to another in such a manner, that the quantity of action shall be the least possible. This quantity of action, says he, is a real expence, in which nature is always frugal. In virtue of this philosophical principle he discovers, that not only the sines are in a constant ratio, but also that they are in the inverse ratio of the velocities, according to Newton's explanation, and not in the direct ratio, as had been pretended by Fermat and Leibnitz.

It is remarkable that, of the many philosophers who have written on refraction, none should have fallen upon so simple a manner of reconciling metaphysics with mechanics; since no more is necessary to that, than making a small alteration in the calculus founded upon Fermat's principle. Now according to that principle, the time, that is, the space divided by the velocity, should be a minimum; so that calling the space run through in the first medium  $S$ , with the velocity  $V$ , and the space run through in the second

medium  $s$ , with the velocity  $v$ , we shall have  $\frac{S}{V} + \frac{s}{v}$

= a minimum; that is to say,  $\frac{\dot{S}}{V} + \frac{\dot{s}}{v} = 0$ , or  $\frac{\dot{S}}{V} = -$

$\frac{\dot{s}}{v}$ . Now it is easy to perceive, that the sines of incidence and refraction are to each other, as  $\dot{S}$  to  $-\dot{s}$ ; whence it follows, that those sines are in the direct ratio of the velocities  $V, v$ ; which is exactly what Fermat makes it to be. But in order to have those sines to be in the inverse ratio of the velocities, it is only supposing  $V \dot{S} + v \dot{s} = 0$ ; which gives  $S V + s v$  or  $S \times V + s \times v$  a minimum: which is Maupertuis's principle.

In the Memoirs of the Academy of Berlin, above cited, may be seen all the other applications which Maupertuis has made of this principle. And whatever may be determined as to his metaphysical basis of it, as also to the idea he has annexed to the quantity of action, it will still hold good, that the product of the space by the velocity is a minimum in some of the most general laws of nature.

ACTIVE, the quality of an agent, or of communicating motion or action to some body. In this sense the word stands opposed to passive: thus we say an active cause, active principle, &c.

Sir Isaac Newton shews, that the quantity of motion in the world must be always decreasing, in consequence of the vis inertiae, &c. So that there is a necessity for

certain active principles to recruit it: such he takes the cause of gravity to be, and the cause of fermentation; adding, that we see but little motion in the universe, except what is owing to these active principles.


ACTIVITY, the virtue or faculty of acting. As the activity of an acid, a poison, &c: the activity of fire exceeds all imagination.

According to Sir Isaac Newton, bodies derive their activity from the principle of attraction.

*Sphere of ACTIVITY*, is the space which furrounds a body, as far as its efficacy or virtue extends to produce any sensible effect. Thus we say, the sphere of activity of a loadstone, of an electric body, &c.

ACUBENE, in *Astronomy*, the Arabic name of a star of the fourth magnitude, in the southern claw of Cancer, marked  $\alpha$  by Bayer. Its longitude for 1761,  $\Omega$   $10^{\circ} 18' 9''$ , south latitude  $5^{\circ} 5' 56''$ .

ACUTE, or sharp; a term opposed to obtuse. Thus, *Acute Angle*, in *Geometry*, is that which is less than a right angle; and is measured by less than  $90^{\circ}$ , or by less than a quadrant of a circle. As the angle A B C.



*Acute angled Triangle*, is that whose three angles are all acute; and is otherwise called an oxygenous triangle. As the triangle D E F.

*Acute-angled Cone*, is that whose opposite sides make an acute angle at the vertex, or whose axis, in a right cone, makes less than half a right angle with the side. As the cone G H I.

Pappus, in his *Mathematical Collections*, says, this name was given to such a cone by Euclid and the ancients, before the time of Apollonius. And they called an

*Acute-angled Section of a Cone*, an Ellipsis, which was made by a plane cutting both sides of an acute-angled cone: not knowing that such a section could be generated from any cone whatever, till it was shewn by Apollonius.

ACUTE, in *Music*, is understood of a tone, or sound, which is high, sharp, or shrill, in respect of some other: in which sense the word stands opposed to grave. And both these sounds are independent of loudness or force: so that the tone may be acute or high, without being loud; and loud without being high or acute. For both the affections of acute and grave, depend intirely on the quickness or slowness of the vibrations by which they are produced.

Sounds considered as grave and acute, that is, in the relation of gravity and acuteness, constitute what is called tune, the foundation of all harmony.

ADAGIO, in *Music*, one of the terms used by the Italians to express a degree or distinction of time.

Adagio denotes the slowest time except grave.

Sometimes the word is repeated, as *adagio, adagio*, to denote a still slower time than the former.

Adagio



Adagio also signifies a slow movement, when used substantively.

ADAMAS, in *Astrology*, a name given to the moon.

ADAR, in the Hebrew *Chronology*, is the 6th month of their civil year, but the 12th of their ecclesiastical year. It contains only 29 days; and it answers to our February; but sometimes entering into the month of March, according to the course of the moon.

ADDITION, the uniting or joining of two or more things together; or the finding of one quantity equal to two or more others taken together.

ADDITION, in *Arithmetic*, is the first of the four fundamental rules or operations of that science; and it consists in finding a number equal to several others taken together, or in finding the most simple expression of a number according to the established notation. The quantity so found equal to several others taken together, is named their sum.

The sign or character of addition is  $+$ , and is called *plus*. This character is set between the quantities to be added, to denote their sum: thus,  $3 + 6 = 9$ , that is, 3 plus 6 are equal to 9; and  $2 + 4 + 6 = 12$ , that is, 2 plus 4 plus 6 are equal to 12.

Simple numbers are either added as above; 6  
or else by placing them under one another, as 4  
in the margin, and adding them together, one 2  
after another, beginning at the bottom: thus 2 —  
and 4 make 6, and 6 make 12. 12 —

Compound numbers, or numbers consisting of more figures than one, are added, by first ranging the numbers in columns under each other, placing always the numbers of the same denomination under each other, that is, units under units, tens under tens, and so on; and then adding up each column separately, beginning at the right hand, setting down the sum of each column below it, unless it amount to ten or some number of tens, and in that case setting down only the overplus, and carrying one for each ten to the next column. Thus, to add 451 and 326,

$$\begin{array}{r} 451 \\ 326 \\ \hline \end{array} \left. \vphantom{\begin{array}{r} 451 \\ 326 \end{array}} \right\} \text{that is } \left\{ \begin{array}{l} 400 + 50 + 1 \\ 300 + 20 + 6 \end{array} \right.$$

$$\text{Sum } 777 = 700 + 70 + 7$$

Also to add the numbers  $329 + 1562 + 20347 + 712048$ ; set them down as in the margin, and beginning at the lowest number on the right hand, say 8 and 7 make 15, and 2 make 17, and 9 make 26; set down 6, and carry 2 to the next column, saying 2 and 4 make 6, and 4 make 10, and 6 make 16, and 2 make 18; set down 8, and carry 1, saying 1 and 3 make 4, and 5 make 9, and 3 make 12; set down 2, and carry 1, saying 1 and 2 make 3, and 1 make 4, which set down; then 1 and 2 make 3; and 7 is 7 to set down: so the sum of all together is 734286. Or it is the same as the sums of the columns set under one another, as in the margin, and then these added up in the same manner.

$$\begin{array}{r} 329 \\ 1562 \\ 20347 \\ 712048 \\ \hline 734286 \\ \hline 26 \\ 16 \\ 11 \\ 3 \\ 3 \\ 7 \\ \hline 734286 \end{array}$$

When a great number of separate sums or numbers are to be added, as in long accounts, it is easier to break or separate them into two or more parcels, which are added up severally, and then their sums added together for the total sum. And thus also the truth of the addition may be proved, by dividing the numbers into parcels different ways, as the totals must be the same in both cases when the operation is right.

Another method of proving addition was given by Dr. Wallis, in his *Arithmetic*, published 1657, by casting out the nines, which method of proof extends also to the other rules of arithmetic. The method is this: add the figures of each line of numbers together severally, casting out always 9 from the sums as they arise in so adding, adding the overplus to the next figure, and setting at the end of each line what is over the nine or nines; then do the same by the sum-total, as also by the former excesses of 9, so shall the last excesses be equal when the work is right. So the former example will be proved as below:

$$\begin{array}{r|l} 329 & 5 \\ 1562 & 5 \\ 20347 & 7 \\ 712048 & 4 \\ \hline 734286 & 3 \end{array} \begin{array}{l} \text{Excess of 9's} \\ \hline \end{array}$$

When the numbers are of different denominations; as pounds, shillings, and pence; or yards, feet, and inches; place the numbers of the same kind under one another, as pence under pence, shillings under shillings, &c; then add each column separately, and carry the overplus as before, from one column to another. As in the following examples:

l.	s.	d.		Yards.	Feet.	Inches.
271	12	3		271	10	3
94	14	7		36	2	7
42	5	10		14	2	5
<hr/>				<hr/>		
408	12	8	sums	323	0	3
<hr/>				<hr/>		

ADDITION of *Decimals*, is performed in the same manner as that of whole numbers, placing the numbers of the same denomination under each other, in which case the decimal separating points will range straight in one column; as in this example, to add together these numbers  $371.0496 + 25.213 + 1.704 + 924.61 + .0962$ .

$$\begin{array}{r} 371.0496 \\ 25.213 \\ 1.704 \\ 924.61 \\ .0962 \\ \hline \end{array}$$

The sum 1322.6728

ADDITION of *Vulgar Fractions*, is performed by bringing all the proposed fractions to a common denominator, if they have different ones, which is an indispensable preparation; then adding all the numerators together,



together, and placing their sum over the common denominator for the sum total required.

$$\text{So } \frac{3}{7} + \frac{2}{7} = \frac{5}{7}. \text{ And } \frac{1}{5} + \frac{2}{5} + \frac{3}{5} = \frac{6}{5} = 1\frac{1}{5}.$$

$$\text{And } \frac{1}{2} + \frac{2}{3} = \frac{3}{6} + \frac{4}{6} = \frac{7}{6} = 1\frac{1}{6}.$$

$$\text{And } \frac{5}{8} + \frac{2}{3} \text{ of } \frac{3}{4} = \frac{5}{8} + \frac{2}{8} = \frac{7}{8} = \frac{4}{3} = 1\frac{1}{3}.$$

$$\text{And } \frac{1}{2} \text{ of } \frac{3}{4} + \frac{2}{3} \text{ of } 5 = \frac{3}{8} + \frac{10}{3} = \frac{3}{24} + \frac{80}{24} = \frac{83}{24} = 3\frac{11}{24}.$$

**ADDITION in Algebra**, or the addition of indeterminate quantities, denoted by letters of the alphabet, is performed by connecting the quantities together by their proper signs, and uniting or reducing such as are susceptible of it, namely similar quantities, by adding their co-efficients together if the signs are the same, but subtracting them when different. Thus the quantity  $a$  added to the quantity  $b$ , makes  $a + b$ ; and  $a$  joined with  $-b$ , makes  $a - b$ ; also  $-a$  and  $-b$  make  $-a - b$ ; and  $3a$  and  $5a$  make  $3a + 5a$  or  $8a$ , by uniting the similar numbers 3 and 5 to make 8.

$$\text{Thus also } 3a + 2bc - 4a^2b + c^3$$

$$\text{added to } 2a - 2bc + 3a^2b + 2c^3$$

$$\text{makes } -5a \quad 0 \quad -a^2b + 3c^3$$

$$\text{Also } \frac{4a}{3c} + \frac{2a}{3c} = \frac{6a}{3c} = \frac{2a}{c};$$

$$\text{and } 3\sqrt{ac} + 5\sqrt{ac} - 1\frac{1}{2}\sqrt{ac} = 6\frac{1}{2}\sqrt{ac};$$

$$\text{and } a\sqrt{ac} + b\sqrt{ac} = a+b\sqrt{ac} \text{ or } (a+b)\sqrt{ac};$$

$$\text{and } 5\sqrt{a^2-x^2} + 7\sqrt{a^2-x^2} = 12\sqrt{a^2-x^2};$$

$$\text{and } 6\sqrt{3} + 4\sqrt{3} - 2\sqrt{3} = 8\sqrt{3};$$

$$\text{and } \frac{3a + 7\sqrt{2x}}{a+x} + \frac{7-3a\sqrt{2x}}{a+x} = \frac{14\sqrt{2x}}{a+x};$$

$$\text{and } \frac{8a\sqrt{ax}}{a+2c} + \frac{16c\sqrt{ax}}{a+2c} = \frac{8a+16c\sqrt{ax}}{a+2c} = 8\sqrt{ax};$$

$$\text{and } -\frac{3ax}{2c} - \frac{5ax}{2c} = -\frac{8ax}{2c} = -\frac{4ax}{c};$$

$$\text{and } -a\sqrt{x} - b\sqrt{x} = -(a+b)\sqrt{x};$$

$$\text{and } +3 - 2 = 1; \text{ and } -3 + 2 = -1;$$

$$\text{and } \frac{7ax}{b} - \frac{4ax}{c} = \frac{7acx}{bc} - \frac{4abx}{bc} = (7c-4b)\frac{ax}{bc}.$$

In the addition of surd or irrational quantities, they must be reduced to the same denomination, or to the same radical, if that can be done; then add or unite the rational parts, and subjoin the common surd. Otherwise connect them with their own signs.

$$\text{So } \sqrt{8} + \sqrt{18} = \sqrt{4 \times 2} + \sqrt{9 \times 2} = 2\sqrt{2}$$

$$+ 3\sqrt{2} = 5\sqrt{2} = \sqrt{50};$$

$$\text{and } \sqrt{12} + \sqrt{17} = \sqrt{4 \times 3} + \sqrt{9 \times 3} = 2\sqrt{3}$$

$$+ 3\sqrt{3} = 5\sqrt{3} = \sqrt{75};$$

$$\text{and } \sqrt{50} + \sqrt{75} = \sqrt{25 \times 2} + \sqrt{25 \times 3} = 5\sqrt{2}$$

$$+ 5\sqrt{3} = (\sqrt{2} + \sqrt{3}) 5;$$

but of  $\sqrt{5}$  and  $\sqrt{6}$  the sum is set down  $\sqrt{5} + \sqrt{6}$ , because the terms are incommensurable, and not reducible to a common surd.

**ADDITION of Logarithms.** See Logarithms.

**ADDITION of Ratios**, the same as composition of ratios; which see.

**ADDITIVE**, denotes something to be added to another, in contradistinction to something to be taken away or subtracted. So astronomers speak of additive equations, and geometricians of additive ratios.

**ADELARD**, or **ATHELARD**, was a learned monk of Bath, in England, who flourished about the year 1130, as appears by some manuscripts of his in Corpus Christi, and Trinity Colleges, Oxford. Vossius says he was universally learned in all the sciences of his time; and that, to acquire all sorts of knowledge, he travelled into France, Germany, Italy, Spain, Egypt, and Arabia. He wrote many books himself, and translated others from different languages; thus, he translated, from Arabic into Latin, Euclid's Elements, at a time before any Greek copies had been discovered; also Erichiafarum, upon the seven planets. He wrote a book on the seven liberal arts, another on the astrolabe, another on the causes of natural compositions, besides several on physics and on medicine.

Although Vossius refers to Oxford for some of these manuscripts, it would yet seem they were not to be found there in Wallis's time; for the Doctor, speaking of this author, and other English authors and travellers about the same age, says, "A particular account of these travels of Shelley and Morley was a while since to be seen in two prefaces to two manuscript books of theirs in the library of Corpus-Christi College in Oxford, but hath lately (by some unknown hand) been cut out, and carried away; which prefaces (one or both of them) did also make mention of the travels of Athelardus Bathoniensis, and are, to that purpose, cited by Vossius out of that manuscript copy. Whoever hath them, would do a kindness (by some way or other) to restore them, or at least a copy of them." Wallis's Algebra, pa. 6.

**ADELM**, **ALDHLMUS**, or **ALTHELMUS**, a learned Englishman, who flourished about the year 680. He was first abbot of Malmesbury, and afterward bishop of Shirburn. He died in the year 709, in the monastery of Malmesbury.

Adelm was the son of Kenred or Kenten, who was the brother of Ina, king of the West Saxons in England. Beside certain books in theology, he composed several on the mathematical sciences &c; as Arithmetic, and Astrology, and librum de philosophorum disciplinis. See Bede's History, lib. 5. cap. 19. He is also mentioned by Bale and William of Malmesbury.

**ADERAIMIN**, or **ALDERAIMIN**, the Arabic name of a star of the third magnitude, in the left shoulder of Cepheus, marked  $\alpha$  by Bayer. Its longitude for 1761,  $79^{\circ} 30' 8''$  north latitude  $68^{\circ} 56' 20''$ .

**ADFFECTED**, see **AFFECTED**.

**ADHESION**, **ADHERENCE**, in *Physics*, is the state of two bodies, joined or fastened together, whether by mutual attraction, the interposition of their own parts, or the impulse or pressure of external bodies. See **COHESION**.

Thus two hollow hemispheres, exhausted of air, are made to adhere firmly together by the pressure of the atmosphere on their convex or external surfaces; for if they are introduced into an exhausted receiver, they presently fall asunder. Also two very well polished planes



planes adhere firmly together, partly by the external pressure of the atmosphere, and partly by the attraction of their parts.

In No. 389 of the *Philos. Trans.* Dr. Desaguliers has given experiments of the adhesion of leaden bullets to each other: the cause of which he resolves into the principle of attraction.

M. Musschenbroeck, in his *Essai de Physique*, has given a great many remarks on the adhesion of bodies, and relates various experiments which he had made upon this matter, but chiefly relative to the resistance made by bodies to fracture, in virtue of the adhesion of their parts; which adhesion he ascribes principally to their mutual attraction. Common experiments prove the mutual adhesion of the parts of water to each other, as well as to the bodies they touch. The same may be said of the particles of air, on which M. Petit has a memoir among those of the Paris Academy of Sciences for the year 1731.

Some authors however are not willing to admit that the adhesion of the parts of water, or indeed of bodies in general, is to be attributed to the attraction of their parts; and they reason thus: suppose, say they, that attraction acts at any small distance, as for example to the distance of one-tenth of an inch, from a particle of water: and about this particle describe a circle whose radius is one-tenth of an inch: then the particle of water will be attracted only by the particles included within the circle; but as these particles act in contrary directions, their mutual effects must destroy one another, and there can be no attraction of the particle, since it will have no more tendency one way than another.

ADHIL, in *Astronomy*, a star, of the sixth magnitude, upon the garment of Andromeda, under the last star in her foot.

ADJACENT, whatever lies immediately by the side of another.

ADJACENT *Angle*, in *Geometry*, is said of an angle when it is immediately contiguous to another, so that they have both one common side. And the term is more particularly used when the two angles have not only one common side, but also when the other two sides form one continued right line.

ADJACENT *bodies*, in *Physics*, are understood of those that are near, or next to, some other body.

ADJUTAGE, or rather AJUTAGE; which see.

ADSCRIPTS, in *Trigonometry*, is used by some mathematicians, for the tangents of arcs. Vieta calls them also *prolines*.

ADVANCE-FOSSE, in *Fortification*, a ditch thrown round the esplanade or glacis of a place, to prevent its being surprised by the besiegers.

The ditch sometimes made in that part of the lines or retrenchments nearest the enemy, to prevent him from attacking them, is also called the advance-fosse.

The advance-fosse should always be full of water, otherwise it will serve to cover the enemy from the fire of the place, if he should become master of the fosse. Beyond the advance-fosse it is usual to construct lunettes, redouts, &c.

ADVENT, *Adventus*, in the Calendar, the time immediately preceding Christmas; and was anciently employed in pious preparation for the *adventus*, or coming on, of the feast of the Nativity.

Advent includes four Sundays, or weeks; commencing either with the Sunday which falls on St. Andrew's day, namely the 30th day of November, or the nearest Sunday to that day, either before or after.

ÆOLIPILE, *Æolipile*, in *Hydraulics*, a hollow ball of metal, with a very small hole or opening; chiefly used to shew the convertibility of water into elastic steam. The best way of fitting up this instrument, is with a very slender neck or pipe, to screw on and off, for the convenience of introducing the water into the inside; for by unscrewing the pipe, and immersing the ball in water, it readily fills, the hole being pretty large; and then the pipe is screwed on. But if the pipe do not screw off, its orifice is too small to force its way in against the included air; and therefore to expel most of the air, the ball is heated red hot, and suddenly plunged with its orifice into water, which will then rush in till the ball is about two-thirds filled with the water. The water having been introduced, the ball is set upon the fire, which gradually heats the contained water, and converts it into elastic steam, which rushes out by the pipe with great violence and noise; and thus continues till all the water is so discharged; though not with a constant and uniform blast, but by fits: and the stronger the fire is, the more elastic will the steam be, and the force of the blast. Care should be taken that the ball be not set upon a violent fire with very little water in it, and that the small pipe be not stopped with any thing; for in such case, the included elastic steam will suddenly burst the ball with a very dangerous explosion.

This instrument was known to the ancients, being mentioned by Vitruvius, lib. 1. cap. 6. It is also treated of, or mentioned, by several modern authors, as Descartes, in his *Meteor.* cap. 4; and Father Merfennus, in prop. 29 *Phædom. Pneumat.* uses it to weigh the air, by first weighing the instrument when red hot, and having no water in it; and afterwards weighing the same when it becomes cold. But the conclusion gained by this means, cannot be quite accurate, as there is supposed to be no air in the ball when it is red hot; whereas it is shewn by Varenus, in his *Geography*, cap. 19, sect. 6, prop. 10, that the air is rarefied but about 70 times; and consequently the weight obtained by the above process, will be about one-70th too small, or more or less according to the intensity of the heat.

In Italy it is said that the *Æolipile* is often used to cure smoky chimneys: for being hung over the fire, the blast arising from it carries up the loitering smoke along with it.

And some have imagined that the *æolipile* might be employed as bellows to blow up a fire, having the blast from the pipe directed into the fire: but experience would soon convince them of their mistake; for it would rather blow the fire out than up, as it is not air, but rarefied water, that is thus violently blown through the pipe.

ÆOLUS, in *Mechanics*, a small portable machine, not long since invented by Mr. Tidd, for refreshing and changing the air in rooms which are made too close.

The machine is adapted to supply the place of a square



square of glass in a sash-window, where it works with little or no noise, on the principle of the sails of a mill, or a smoke-jack; and thus admitting an agreeable quantity of air, at a convenient part of the room.

*Æolus's Harp*, or *Æolian Harp*, an instrument so named, from its producing an agreeable melody, merely by the action of the wind.

Neither the age nor inventor of this instrument are very well known. It is not mentioned by Merfennus in his *Harmonics*, where he describes most sorts of musical instruments: and yet the description and use of it was given soon after, by Kircher, in his book, *Magia Phenotactica & Phonurgia*.

The construction of this instrument is thus; let a box be made of as thin deal as possible, its length answering exactly to the width of the window in which it is to be placed; five or six inches deep, and seven or eight inches wide. Across the top, and near each end, glue on a bit of wainscot, about half an inch high, and a quarter of an inch thick, to serve as two bridges for the strings to be stretched over, by means of pins inserted into holes a little behind the bridges, nearer the ends, half the number being at one end, and half at the other end: these pins are like those of a harpsichord; and for their better support in the thin deal, a piece of beech of about an inch square, and length equal to the breadth of the box, is glewed on the inside of the lid, immediately under the place of the pins, the holes for receiving them being bored through this piece. It is strung with small catgut, or blue first fiddle strings, more or less at pleasure, on the outside and lengthways of the lid, fixing one end to one of the small pins, and twisting the other end about the opposite or stretching pin. A couple of sound-holes are cut in the lid; and the thinner this is, the better will be the performance.

When the strings are tuned unison, and the instrument placed, with the top or stringed side outwards, in the window to which it is fitted, the air blowing upon that window, the instrument will give a sound like a distant choir, increasing or decreasing according to the strength of the wind.

*ÆRA*, in Chronology, is the same as epoch, or epocha, and means a fixed point of time, from which to begin a computation of the years ensuing.

The word is sometimes also written *era* in ancient authors. Its origin is contested, though it is generally supposed that it had its rise in Spain. Some imagine that it is formed from *a. er. a.* the abbreviations of the words, *annus erat Augusti*, or from *a. e. r. a.* the initials of the words *annus erat regni Augusti*, because the Spaniards began their computation from the time that their country came under the dominion of Augustus. Others derive it from *as*, *brass*, the tribute money with which Augustus taxed the world. It is also said that *era* originally signified a number stamped on money to determine its current value. And that the ancients used *as* or *era* as an article, as we do the word *item*, to each particular of an account; and hence it came to stand for a sum or number itself.

*ÆRA* also means the way or mode of accounting time. Thus we say such a year of the Christian æra, &c.

*Spanish ÆRA*, otherwise called the year of Cæsar,

was introduced after the second division of the Roman provinces, between Augustus, Anthony, and Lepidus, in the 714th year of Rome, the 4676th year of the Julian period, and the 38th year before Christ. In the 447th year of this æra, the Alani, the Vandals, Suevi, &c, entered Spain. It is frequently mentioned in the Spanish affairs; their councils, and other public acts, being all dated according to it. Some say it was abolished under Peter IV, king of Arragon, in the year of Christ 1358, and the Christian æra introduced instead of it. But Mariana observes that it ceased in the year of Christ 1383, under John I, king of Castile. The like was afterwards done in Portugal.

*Christian ÆRA*. It is generally allowed by Chronologers, that the computation of time from the birth of Christ, was only introduced in the sixth century in the reign of Justinian; and it is commonly ascribed to Dionysius Exiguus. This æra came then into use in deeds, and such like; before which time either the olympiads, the year of Rome, or that of the reign of the emperors, was used for such purposes.

See an account of the other principal æras under the word Epoch.

*AERIAL Perspective*, is that which represents bodies diminished and weakened, in proportion to their distance from the eye.

*Ærial Perspective* chiefly respects the colours of objects, whose force and lustre it diminishes more or less, to make them appear as if more or less remote.

It is founded upon this, that the longer the column of air an object is seen through, the more feebly do the visual rays emitted from it affect the eye.

*AEROGRAPHY*, a description of the air, or atmosphere, its limits, dimensions, properties, &c.

*AEROLOGY*, the doctrine or science of the air, and its phænomena, its properties, good and bad qualities, &c. It is much the same with the foregoing word, *Aerography*.

*AEROMETRY*, *Aerometria*, the science of measuring the air, its powers and properties; comprehending not only the quantity of the air itself, as a fluid body, but also its pressure or weight, its elasticity, rarefaction, condensation, &c.

The term is not much used at present; this branch of natural philosophy being usually called pneumatics, which see. Wolfius, late professor of mathematics at Hall, having reduced several properties of the air to geometrical demonstrations, first published at Leipzig his *Elements of Aerometry*, in the German language, and afterwards more enlarged in Latin, which have since been inserted in his *Cursus Mathematicus*, in five volumes in 4to.

*AERONAUTICA*, the pretended art of sailing through the air, or atmosphere, in a vessel, sustained as a ship in the sea.

*AEROSTATICA*, is properly the doctrine of the weight, pressure, and balance of the air and atmosphere.

*AEROSTATION*, in its proper and primary sense, denotes the science of weights suspended in the air; but in the modern application of the term, it signifies the art of navigating or floating in the air, both as to the practice and principles of it. Hence also the machines which are employed for this purpose, are called  
*aerostats*,



*aerostats*, or *aerostatic* machines ; and which, on account of their round and bell-like shape, are otherwise called *air balloons*. Also *aeronaut* is the name given to the person who navigates or floats in the air by means of such machines.

*Principles of AEROSTATION.* The fundamental principles of this art have been long and generally known, as well as speculations on the theory of it ; but the successful application of them to practice seems to be altogether a modern discovery. These principles chiefly respect the weight or pressure, and elasticity of the air, with its specific gravity, and that of the other bodies to be raised or floated in it : the particular detail of which principles may be seen under the respective words in this dictionary. Suffice it therefore in this place to observe, that any body which is specifically, or bulk for bulk, lighter than the atmosphere, or air encompassing the earth, will be buoyed up by it, and ascend, like as wood, or a cork, or a blown bladder, ascends in water. And thus the body would continue to ascend to the top of the atmosphere, if the air were every where of the same density as at the surface of the earth. But as the air is compressible and elastic, its density decreases continually in ascending, on account of the diminished pressure of the superincumbent air, at the higher elevations above the earth ; and therefore the body will ascend only to such height where the air is of the same specific gravity with itself ; where the body will float, and move along with the wind or current of air, which it may meet with at that height. This body then is an aerostatic machine, of whatever form or nature it may be. And an air-balloon is a body of this kind, the whole mass of which, including its covering and contents, and the weights annexed to it, is of less weight than the same bulk of air in which it rises.

We know of no solid bodies however that are light enough thus to ascend and float in the atmosphere ; and therefore recourse must be had to some fluid or aeriform substance.

Among these, that which is called inflammable air is the most proper of any that have hitherto been discovered. It is very elastic, and from six to ten or eleven times lighter than common atmospheric air at the surface of the earth, according to the different methods of preparing it. If therefore a sufficient quantity of this kind of air be inclosed in any thin bag or covering, the weight of the two together will be less than the weight of the same bulk of common air ; and consequently this compound mass will rise in the atmosphere, and continue to ascend till it attain a height at which the atmosphere is of the same specific gravity as itself ; where it will remain or float with the current of air, as long as the inflammable air does not escape through the pores of its covering. And this is an inflammable air-balloon.

Another way is to make use of common air, rendered lighter by warming it, instead of the inflammable air. Heat, it is well known, rarefies and expands common air, and consequently lessens its specific gravity ; and the diminution of its weight is proportional to the heat applied. If therefore the air, inclosed in any kind of a bag or covering, be heated, and consequently dilated, to such a degree, that the excess of the weight of an equal bulk of common air, above the

weight of the heated air, be greater than the weight of the covering and its appendages, the whole compound mass will ascend in the atmosphere, till, by the diminished density of the surrounding air, the whole become of the same specific gravity with the air in which it floats ; where it will remain, till, by the cooling and condensation of the included air, it shall gradually contract and descend again, unless the heat is renewed or kept up. And such is a heated air-balloon, otherwise called a Montgolfier, from its inventor.

Now it has been discovered, by various experiments, that one degree of heat, according to the scale of Fahrenheit's thermometer, expands the air about one five-hundredth part ; and therefore that it will require about 500 degrees, or nearer 484 degrees of heat, to expand the air to just double its bulk. Which is a degree of heat far above what it is practicable to give it on such occasions. And therefore, in this respect, common air heated, is much inferior to inflammable air, in point of levity and usefulness for aerostatic machines.

Upon such principles then depends the construction of the two sorts of air-balloons. But before treating of this branch more particularly, it will be proper to give a short historical account of this late-discovered art.

*History of AEROSTATION.* Various schemes for rising in the air, and passing through it, have been devised and attempted, both by the ancients and moderns, and that upon different principles, and with various success. Of these, some attempts have been upon mechanical principles, or by virtue of the powers of mechanism : and such are conceived to be the instances related of the flying pigeon made by Archytas, the flying eagle and fly by Regiomontanus, and various others. Again, other projects have been formed for attaching wings to some part of the body, which were to be moved either by the hands or feet, by the help of mechanical powers ; so that striking the air with them, after the manner of the wings of a bird, the person might raise himself in the air, and transport himself through it, in imitation of that animal. But of these and various other devices of the like nature, a particular account will be given under the article *artificial flying*, as belonging rather to that species or principle of motion, than to our present subject of aerostation, which is properly the sailing or floating in the air by means of a machine rendered specifically lighter than that element, in imitation of aqueous navigation, or the sailing upon the water in a ship, or vessel, which is specifically lighter than the water.

The first rational account that we have upon record, for this sort of sailing, is perhaps that of our countryman Roger Bacon, who died in the year 1292. He not only affirms that the art is feasible, but assures us that he himself knew how to make an engine, in which a man sitting might be able to carry himself through the air like a bird ; and he farther affirms that there was another person who had tried it with success. And the secret it seems consisted in a couple of large thin shells, or hollow globes, of copper, exhausted of air ; so that the whole being thus rendered lighter than air, they would support a chair, on which a person might sit.

Bishop Wilkins too, who died in 1672, in several  
F of



of his works, makes mention of similar ideas being entertained by divers persons. "It is a pretty notion to this purpose, says he (in his *Discovery of a New World*, prop. 14), mentioned by Albertus de Saxonia, and out of him by Francis Mendoza, that the air is in some part of it navigable. And that upon this statick principle, any brass or iron vessel (suppose a kettle), whose substance is much heavier than that of the water; yet being filled with the lighter air, it will swim upon it, and not sink." And again, in his *De dalus*, chap. 6, "Scaliger conceives the framing of such volant automata to be very easy. *Volantis columbe machinulam, cujus autorem Archytam tradunt, vel facillime profiteri audeo.* Those ancient motions were thought to be contrived by the force of some included air: So *Gellius, Ita erat scilicet libramentis suspensum, & aura spiritus inclusa, atque occulta confitum, &c.* As if there had been some lamp, or other fire within it, which might produce such a forcible rarefaction, as should give a motion to the whole frame." From which it would seem that Bishop Wilkins had some confused notion of such a thing as a heated air-balloon.

Again F. Francisco Lana, in his *Prodroma*, printed in 1670, proposes the same method with that of Roger Bacon, as his own thought.

He considered that a hollow vessel, exhausted of air, would weigh less than when filled with that fluid; he also reasoned that, as the capacity of spherical vessels increases much faster than their surface, if there were two spherical vessels, of which the diameter of one is double the diameter of the other; then the capacity of the former will be equal to 8 times the capacity of the latter, but the surface of that only equal to 4 times the surface of this: and the one sphere have its diameter equal to triple the diameter of the other; then the capacity of the greater will be equal to 27 times the capacity of the less, while its surface is only 9 times greater: and so on, the capacities increasing as the cubes of the diameters, while the surfaces increase only as the squares of the same diameters. And from this mathematical principle, father Lana deduces, that it is possible to make a spherical vessel of any given matter, and thickness, and of such a size as, when emptied of air, it will be lighter than an equal bulk of that air, and consequently that it will ascend in that element, together with some additional weight attached to it. After stating these principles, father Lana computes that a round vessel of plate brass, 14 feet in diameter, weighing 3 ounces the square foot, will only weigh 1848 ounces; whereas a quantity of air of the same bulk will weigh 2155 $\frac{2}{3}$  ounces, allowing only one ounce to the cubic foot; so that the globe will not only be sustained in the air, but will also carry up a weight of 307 $\frac{2}{3}$  ounces: and by increasing the bulk of the globe, without increasing the thickness of the metal, he adds, a vessel might be made to carry a much greater weight.

Such then were the ingenious speculations of learned men, and the gradual approaches towards this art. But one thing more was yet wanting: although acquainted in some degree with the weight of any quantity of air, considered as a detached substance, it seems they were not aware of its great elasticity, and the universal pressure of the atmosphere; by which pressure, a globe of the dimensions above-described, and exhausted of its

air, would immediately be crushed inwards, for want of the equivalent internal counter pressure, to be sought for in some element, much lighter than common air, and yet nearly of equal pressure or elasticity with it; a property or circumstance attending common air when considerably heated. It is evident then that the schemes of ingenious men hitherto must have terminated in mere speculation; otherwise they could never have recorded schemes, which, on the first attempt to put in practice, must have manifested their own insufficiency, by an immediate failure of success: For instead of exhausting the vessel of air, it must either be filled with common air heated, or with some other equally elastic and lighter air. So that upon the whole it appears, that the art of traversing the air, is an invention of our own time; and the whole history of it is comprehended within a very short period.

The rarefaction and expansion of air by heat, is a property of it that has long been known, not only to philosophers, but even to the vulgar: by this means it is, that the smoke is continually carried up our chimneys; and the effect of heat upon air, is made very sensible by bringing a bladder, only partly full of air, near a fire; when the air presently expands with the heat, and distends the bladder so as almost to burst it: and so well are the common people acquainted with this effect, that it is the common practice of those who kick blown bladders about for foot-balls, to bring them from time to time to the fire, to restore the spring of the air, and distension of the ball, lost by the continual waste of that fluid through the sides of it.

But the great levity of inflammable air, is a very modern discovery. As to the inflammable property of this air itself, it had been long known to miners, and especially in coal mines, by the dreadful effects it sometimes produces by its explosions. Among them it is sometimes vulgarly called sulphur, but more properly the fire damp, or inflammable damp, to distinguish it from the choak damp, and other damps, a species of air sometimes found in deep wells and mines, and which does not explode nor take fire, but presently extinguishes candles, and suffocates the persons who may happen to go into it. But it seems that it was Mr. Cavendish who first discovered with exactness the specific gravity of inflammable air; and his experiments and observations upon it, are published in the 56th volume of the *Philosophical Transactions* for the year 1766. Soon after this discovery of Mr. Cavendish, it occurred to the ingenious Dr. Black of Edinburgh, that if a bladder, or other vessel, sufficiently light and thin, were filled with this air, it would form altogether a mass lighter than the same bulk of atmospheric air, and consequently that it would ascend in it.

This idea he mentioned in his chemical lectures in the year 1767 or 1768; and he farther proposed to exhibit the experiment, by filling the allantois of a calf with such air. The allantois however was not prepared just at the time when he was at that part of his lectures, and other avocations afterwards prevented his design: so that, considering it only as an amusing experiment, and being fully satisfied of the truth of so evident an effect, he contented himself with barely mentioning the experiment from time to time in his lectures. About the year 1777 or 1778 too it occurred to Mr. Cavallo, that



that it might be possible to construct a vessel, which, when filled with inflammable air, would ascend in the atmosphere: and there is no doubt but that similar ideas would occur to many other persons, of so evident a consequence of Mr. Cavendish's discovery.

But it seems to have been Mr. Cavallo who first actually attempted the experiment, in which however he succeeded no farther than in being able to raise soap bubbles of two or three inches diameter: a thing which had been done by children for their amusement time immemorial. These experiments Mr. Cavallo made in the beginning of the year 1782, and an account of them was read at a public meeting of the Royal Society on the 20th day of June of that year. From which it appears that he tried bladders and paper of various sorts. But the bladders, however thin they were made by scraping, &c, were still found too heavy to ascend in the atmosphere, when fully inflated with the inflammable air: and in using China paper, he found that this air passed through its pores, like water through a sieve. And having failed of success by blowing the same air into a thick solution of gum, thick varnishes, and oil paint, he was obliged to rest satisfied with soap-balls or bubbles, which, being filled with inflammable air, by dipping the end of a small glass tube, connected with a bladder containing the air, into a thick solution of soap, and gently compressing the bladder, ascended rapidly in the atmosphere, and broke against the ceiling of the room.

Here however it seems the matter might have rested, had it not been for experiments made in France soon after, by the two brothers Stephen and Joseph Montgolfier, upon principles suggested, not by the levity of inflammable air, which probably they had never heard of, but by that of smoke and clouds ascending in the atmosphere. These two brothers it seems were natives of Annonay, a town in the Vivarais, about 36 miles distant from Lyons; and that in their youth, Stephen, the elder, had assiduously studied the mathematics, but the other had applied himself more particularly to natural philosophy and chemistry. They were not intended for any particular way of business, but the death of a brother obliged them to put themselves at the head of a considerable paper manufactory at Annonay. In the intervals of time allowed by their business they amused themselves in several philosophical pursuits, and particularly with the experiments in aerostation, of which we are now to give some account. It would be perhaps impossible to know all the particular steps and ideas which finally produced this discovery: but it has been said that the real principle, upon which the effect of the aerostatic machine depends, was unknown even for a considerable time after its discovery: that M. Montgolfier attributed the effect of the machine, not to the rarefaction of the air, which is the true cause, but to a certain gas, specifically lighter than common air, which was supposed to be developed from burning substances, and which was commonly called Montgolfier's gas. Be this however as it may, it is well known that the two brothers began to think of the experiment of the aerostatic machine about the middle or the latter part of the year 1782. The natural ascension of the smoke and the clouds in the atmosphere, suggested the first idea; and to imitate those bodies,

or to inclose a cloud in a bag, and let the latter be lifted up by the buoyancy of the former, was the first project of those celebrated gentlemen.

Accordingly the first experiment was made at Avignon by Stephen, the elder brother, about the middle of November 1782. Having prepared a bag of fine silk, in the shape of a parallelopipedon, and of about 40 cubic feet in capacity, he applied burning paper to an aperture in the bottom, which rarefied the air, and thus formed a kind of cloud in the bag; and when it became sufficiently expanded, it ascended rapidly to the ceiling.

Soon afterwards the experiment was repeated with the same machine at Annonay, by the two brothers, in the open air; when the bag ascended to the height of about 70 feet. Encouraged by this success, they constructed another machine, of about 650 cubic feet capacity; which, when inflated as before, broke the cords which confined it, and after ascending rapidly to the height of about 600 feet, descended and fell on the adjoining ground. With another larger machine, of 37 feet diameter, they repeated the experiment on the 25th day of April, which answered exceedingly well: the machine had such force of ascension, that, breaking abruptly from its confinement of ropes, it rose to the height of more than 1000 feet, and then, being carried by the wind, descended and fell at a place about three quarters of a mile from the place of its ascension. The capacity of this machine was equal to above 23 thousand cubic feet, and, being nearly globular, when inflated, it measured 117 English feet in circumference. The covering was formed of linen, lined with paper; and its aperture at the bottom was fixed to a wooden frame, of about 4 feet square, or 16 feet in surface. When filled with vapour, which it was conjectured might be about half as heavy as common air, it was capable of lifting up about 490 pounds, besides its own weight, which, together with that of the wooden frame, was equal to about 500 pounds. With this same machine the next experiment was publicly performed at Annonay, on the 5th of June 1783, before a great multitude of spectators. The flaccid bag was suspended on a pole 35 feet high; straw and chopped wool were burned under the opening at the bottom; the vapour, or rather smoke, soon inflated the bag, so as to distend it in all its parts; and this enormous mass ascended in the air with such velocity, that in less than ten minutes it reached the height of above 6 thousand feet, when a breeze carried it in an horizontal direction to the distance of 7668 feet, or near a mile and a half, where it descended gently to the ground.

As soon as the news of this experiment reached Paris, the philosophers of that city, conceiving that a new species of gas, of about half the weight of common air, had been discovered by Messrs. Montgolfier; and knowing that the weight of inflammable air was but about the eighth or tenth part of the weight of common air, they justly concluded that inflammable air would answer the purpose of this experiment better than the gas of Montgolfier, and accordingly they resolved to make trial of it.

A subscription was opened by M. Faujas de St. Fond, towards defraying the expence of the experiment. A sufficient sum of money having soon been raised,



Messrs. Roberts were appointed to construct the machine, and M. Charles, professor of experimental philosophy, to superintend the work. After a considerable time spent, and surmounting many difficulties in obtaining a sufficient quantity of inflammable air, and searching out a substance light enough for the covering, they at length constructed a globe of the silk called lutestring, which was rendered impervious to the inclosed air by a varnish of elastic gum or caeutchouc, dissolved in some kind of spirit or essential oil. The diameter of this globe was about 13 feet; and it had only one aperture, like a bladder, to which a stop-cock was adapted: and the weight of this covering, when empty, together with that of the stop-cock, was 25 pounds.

On the 23d of August 1783, they began to fill the globe with inflammable air; but this, being their first attempt, was attended with many obstructions and disappointments, which took up two or three days to overcome.

At length however it was prepared for exhibition, and on the 27th it was carried from the Place des Victoires, where it had been prepared, to the Champs de Mars, a spacious open ground in the front of the Military School, where, after introducing some more inflammable air, and disengaging it from the cords by which it was held down, it rose, in less than two minutes, to the height of 3123 feet: the specific gravity of the balloon, when it went up, being 35 pounds less than that of common air. At that height the balloon entered a cloud, but soon appeared again; and at last it was lost among other clouds. After floating about in the air for about three quarters of an hour, it fell in a field about 15 miles from the place of its ascent; where, as we may easily imagine, it occasioned great amazement to the peasants who found it. Its fall was owing to a rent in the covering, probably occasioned by the superior elasticity of the inflammable air, over that of the rare part of the atmosphere to which it had ascended.

In consequence of this brilliant experiment, numberless small balloons were made, mostly of gold-beater's skin, from 6 and 9 to 18 or 20 inches diameter; their cheapness putting it in the power of almost every family to satisfy its curiosity relative to the new experiment; and in a few days time balloons were seen flying all about Paris, from whence they were soon after sent abroad.

Mr. Joseph Montgolfier repeated an experiment with a machine of his construction before the commissaries of the Academy of Sciences, on the 11th and 12th of September. The machine was about 74 feet high, and 43 feet in diameter; it was made of canvass, covered with paper both within and without, and weighed 1000 pounds. It was filled with rarefied air in 9 minutes, and in one trial the weight of eight men was not sufficient to keep it down. It was not suffered to go up, as it had been intended for exhibition before the Royal Family, a few days after. By the violence of the rain, however, which fell about this time, it was so much spoiled, that he thought proper to construct another for that purpose, in which he used so great dispatch, that it was completed in the short space of four days time. This machine was constructed of cloth made of linnen and cotton

thread, and painted with water colours both within and without. Its height was 60 feet, and diameter 43 feet. Having made the necessary preparation for inflating it, the operation was begun about one o'clock on the 19th of the same month, before the king and queen, the court, and the inhabitants of the place, as well as all the Parisians who could procure a conveyance to Versailles. The balloon was soon filled, and in eleven minutes after the commencement of the operation, the ropes being cut, it ascended, bearing up with it a wicker cage, containing a cock, a duck, and a sheep, the first animals that ever ascended into the atmosphere with an aerostatic machine. Its power of ascension, or the weight by which it was lighter than an equal bulk of common air, allowing for the animals and their cage, was 696 pounds. The balloon rose to the height of 1440 feet; and being driven by the wind for the space of eight minutes, it gradually descended in consequence of two large rents made in the covering by the wind, and fell in a wood at the distance of 10,200 feet, or about two miles from Versailles. The animals landed again as safe as when they went up, and the sheep was found feeding.

The success of this experiment induced M. Pilatre de Rozier, with a philosophical intrepidity which will be recorded with applause in the history of aerostation, to offer himself as the first adventurer in this aerial navigation. For this purpose M. Montgolfier constructed a new machine, of an oval shape, in a garden of the fauxbourg St. Antoine; its diameter being about 48 feet, and height 74 feet. To the aperture in the lower part was annexed a wicker gallery about three feet broad, with a ballustrade of three feet high. From the middle of the aperture an iron grate, or brazier, was suspended by chains, descending from the sides of the machine, in which a fire was lighted for inflating the machine; and towards the aperture port-holes were opened in the gallery, through which any person, who might venture to ascend, might feed the fire on the grate with fuel, and regulate at pleasure the dilatation of the air inclosed in the machine: the weight of the whole being upwards of 1600 pounds. On the 15th of October 1783, the fire being lighted, and the balloon inflated, M. P. de Rozier placed himself in the gallery, and, to the astonishment of a multitude of spectators, ascended as high as the length of the restraining cords would permit, which was about 84 feet from the ground, and there kept the machine afloat about four minutes and a half, by repeatedly throwing straw and wool upon the fire: the machine then descended gradually and gently, through a medium of increasing density, to the ground; and the intrepid adventurer assured the admiring spectators that he had not experienced the least inconvenience in this aerial excursion. This experiment was repeated on the 17th with nearly the same success; and again several times on the 19th, when M. P. de Rozier, by a partial ascent and descent, several times repeated, evinced to the multitude of observers that the machine may be made to ascend and descend at the pleasure of the aeronaut, by merely increasing or diminishing the fire in the grate. The balloon having been hauled down, by the ropes which always confined it, M. Gironde de Villette placed himself in the gallery  
opposite



opposite to M. de Rozier, and the machine being suffered to ascend, it hovered for about 9 minutes over Paris, in the sight of all its inhabitants, at the height of 330 feet. And on their descending, the marquis of Arlandes ascended with M. de Rozier much in the same manner.

In consequence of the report of these experiments, signed by the commissaries of the Academy of Sciences, it was ordered that the annual prize of 600 livres should be given to Messrs. Montgolfier for the year 1783.

In the experiments above-recited, the machine was always secured by long ropes, to prevent its entire escape: but they were soon succeeded by unconfined aerial navigation. For this purpose the same balloon of 74 feet in height was conveyed to La Muette, a royal palace in the Bois de Boulogne: and all things being got ready, on the 21st of November 1783, M. P. de Rozier and the marquis d'Arlandes took their post in opposite sides of the gallery, and at 54 minutes after one the machine was absolutely abandoned to the element, and it ascended calmly and majestically in the atmosphere. On reaching the height of about 280 feet the intrepid aeronauts waved their hats to the astonished multitude: but they soon after rose too high to be distinguished, and it is supposed they rose to more than 3000 feet in height. At first they were driven, by a north-west wind, horizontally over the river Seine and part of Paris, taking care to clear the steeples and high buildings by increasing the fire; and in rising they met with a current of air which carried them southward. Having thus passed the Boulevard, and finally desisting from supplying the fire with fuel, they descended very gently in a field beyond the new Boulevard, about 9000 yards, or a little more than 5 miles distant from the palace de La Muette, having been between 20 and 25 minutes in the air. The weight of the whole apparatus including that of the two travellers, was between 1600 and 1700 pounds.

Notwithstanding the rapid progress of aerostation in France, it is remarkable that we have no authentic account of any experiments of this kind being attempted in other countries. Even in our own island, where all arts and sciences find an indulgent nursery, and many their birth, no aerostatic machine was seen before the month of November 1783. Various speculations have been made on the reasons of this strange neglect of so novel and brilliant an experiment. But none seemed to carry any shew of probability except that it was said to be discouraged by the leader of a philosophical society, expressly instituted for the improvement of natural knowledge, for the reason, as it was said, that it was the discovery of a neighbouring nation. Be this however as it may, it is a fact that the first aerostatic experiment was exhibited in England by a foreigner unconnected and unsupported. This was a count Zambecari, an ingenious Italian, who happened to be in London about that time. He made a balloon of oiled-silk, 10 feet in diameter, weighing only 11 pounds: it was gilt, both for ornament, and to render it more impermeable to the inflammable air with which it was to be filled. The balloon, after being publicly shewn for several days in London, was

carried to the Artillery Ground, and there being filled about three-quarters with inflammable air, and having a direction inclosed in a tin box for any person by whom it should afterwards be found, it was launched about one o'clock on the 25th of November 1783. At half past three it was taken up near Petworth in Suffex, 48 miles distant from London; so that it travelled at the rate of near 20 miles an hour. Its descent was occasioned by a rent in the silk, which must have been the effect of the rarefaction of the inflammable air when the balloon ascended to a rarer part of the atmosphere.

The French philosophers having executed the first aerial voyage with a balloon inflated by heated air, resolved to attempt a similar voyage with a balloon filled with inflammable air, which seemed to be preferable to dilated air in every respect, the expence of preparing it only excepted. A subscription was opened however to defray that expence, which was estimated at about ten thousand livres; and the balloon was constructed by Messrs. Roberts, of gores of silk, varnished with a solution of elastic gum. Its form was spherical, and it measured  $27\frac{1}{2}$  feet in diameter. The upper hemisphere was covered by a net, which was fastened to a hoop encircling its middle, and called its equator. To this equator was suspended by ropes a car or boat, covered with painted linen, and beautifully ornamented, which swung a few feet below the balloon. To prevent the bursting of the machine by the expansion of the inflammable air in a rarer medium, or to cause the balloon to descend, it was furnished with a valve, which might be opened by means of a string descending from it, for discharging a part of the internal air, without admitting the external to enter: And the car was ballasted with bags of sand, for the purpose of lightening it occasionally, and causing it to ascend: so that by letting some of the air escape through the valve, they might descend; and by discharging some of their sand ballast, ascend. To this balloon was likewise annexed a long pipe by which it was filled. The apparatus for filling it consisted of several casks placed round a large tub of water, each having a long tin tube, that terminated under a vessel or funnel which was inverted into the water of the tub, and communicated with the long pipe annexed to the lower part of the balloon. Iron filings and diluted vitriolic acid being put into the casks, the inflammable air which was produced from these materials, passed through the tin tubes, thence through the water of the tubs to the inverted funnel, and so through the pipe into the balloon. When inflated, the weight of the common air which was equal in bulk to the balloon, was  $771\frac{1}{2}$  pounds; also the power of ascension, or weight just necessary to keep it from ascending, was 20 pounds, and the weight of the balloon, with its car, passengers, and all its appendages, was  $604\frac{1}{2}$  pounds, which two together make  $624\frac{1}{2}$  pounds: and this taken from  $771\frac{1}{2}$  pounds, the weight of common air displaced, leaves 147 pounds for the weight of the inflammable air contained in the balloon, and which is to  $771\frac{1}{2}$  pounds, the weight of the same bulk of common air, nearly as 1 to  $5\frac{1}{4}$ ; that is, the inflammable air used in this experiment was  $5\frac{1}{4}$  times lighter than common air.

The first of December was fixed on for the display of this grand experiment; and every preparation was made



made for conducting it with advantage. The garden of the Thuilleries at Paris was the scene of operation; which was soon crowded and encompassed with a prodigious multitude of observers: Signals were given, from time to time, by the firing of cannon, waving of flags, &c: and a small montgolfier was launched, for shewing the direction of the wind, and for the amusement of the people previous to the general display. At three quarters after one o'clock, M. Charles and one of the Roberts, having seated themselves in the boat attached to the balloon, and being furnished with proper instruments, cloathing, and provisions, left the ground, and ascended with a moderately accelerated velocity to the height of about 600 yards; the surrounding multitude standing silent with fear and amazement; while the aerial navigators at this height made signals of their safety. When they left the ground, the thermometer, according to Fahrenheit's scale, stood at 59 degrees; and the barometer, at 30.18 inches: and at the utmost height to which they ascended, the barometer fell to 27 inches; from which they deduced their height as above to be 600 yards, or one third part of a mile. During the rest of the voyage the quicksilver in the barometer was generally between 27 and 27.65 inches, rising and falling, as part of the ballast was thrown out, or some of the inflammable air escaped from the balloon. The thermometer generally stood between 53 and 57 degrees. Soon after their ascent, they remained stationary for some time: they then moved horizontally in the direction north-north-west: and having crossed the Seine, and passed over several towns and villages, to the great amazement of the inhabitants, they descended in a field, about 27 miles distant from Paris, at three-quarters past 3 o'clock; so that they had travelled at the rate of near 15 miles an hour, without feeling the least inconvenience.

The balloon still containing a considerable quantity of inflammable air, M. Charles re-ascended alone, and it was computed he went to the height of 3100 yards, or almost 2 miles, the barometer being then at 20 English inches: having amused himself in the air about 33 minutes, he pulled the string of the valve, and descended at 3 miles distance from the place of his ascent. All the inconvenience he experienced in his great elevation, was a dry sharp cold, with a pain in one of his ears and a part of his face, which he ascribed to the dilatation of the internal air: a circumstance that usually happens to persons who suddenly change the density of their atmosphere, either by ascending into a rarer, or descending into a denser one. The small balloon, launched at the beginning by M. Montgolfier, was found to have moved in a direction opposite to that of the aeronauts; from which it is inferred that there were two currents of air at different heights above the earth.

In the month of December this year, several experiments were made at Philadelphia in America with air balloons, by Messrs. Rittenhouse and Hopkins. They constructed and filled a great many small balloons, and connected them together; in which a man went up several times, and was drawn down again; and finally, the ropes being cut, he ascended to the height of 100 feet, and floated to a considerable distance; but,

being afraid, he cut open the balloons with a knife, and so descended.

About the close of this year small balloons were sent up in many places, and were become very common in some parts of France and England. And in the beginning of the year following, their number and magnitude increased considerably; and some of the more remarkable ones were as follow:—On the 19th of January M. Joseph Montgolfier, accompanied by six other persons, ascended from Lyons with a rarefied air balloon, to the height of 1000 yards. This was the largest machine that had been hitherto made, being 131 feet high, and 104 feet in diameter: it was formed of a double covering of linen, with three layers of paper between them; and it weighed, when it went up, 1600 pounds, including the gallery, passengers, &c. It was at first intended for six passengers; but before it went up, it was not judged safe to freight it with more than three: however no authority nor solicitations could prevail upon any of the six to quit their place, nor even to cast lots which three should resign their pretensions: so that the spectators saw them all ascend with terror and anxiety; and to add to their distress, when the ropes were cut, and the machine had ascended a foot or two from the ground, a seventh person suddenly leaped into the gallery, and the fire being increased, the whole ascended together. To add to the terror of the scene, after being in the air about 15 minutes, a large rent of about 50 feet in length was made by the balloon taking fire, in consequence of which it descended very rapidly to the ground, though fortunately without injury to any of the aeronauts.

On the 22d of February an inflammable air balloon was launched from Sandwich in Kent. It was but a small one, being only 5 feet in diameter; but it was rendered remarkable by being the first machine that crossed the sea from England to France. It was found in a field at Warneton, about 9 miles from Lille in French Flanders, two hours and a half after it left Sandwich, the distance being about 74 miles; so that it floated at the rate of about 30 miles an hour.

The chevalier Paul Andreani, of Milan, was the first aerial traveller in Italy. The chevalier was at the sole expence of this machine, but was assisted in the construction by two brothers of the name of Gerli. They all three ascended together near Milan on the 25th of February, and remained in the atmosphere about 20 minutes, when they descended, all their fuel being exhausted. This machine was a montgolfier, of a spherical shape, and about 68 feet in diameter. From calculations made on the power of this, and other machines of the same sort, it appears that the included air is rarefied commonly but about one-third, or that the included warm air weighs about two-thirds of the same bulk of the external or common air.

The next aerial voyage was performed on the 2d of March 1784, by M. Jean Pierre Blanchard, a man who has since that time made more voyages than any other person, and who has rendered himself famous by being the first who has floated in the air over the channel from England to France. M. Blanchard it seems had for many years been in pursuit of mechanical means for flying through the air; but on hearing of the late invented air balloons, he dropped his



his former pursuits, and turned his attention to them. He accordingly constructed one of 27 feet diameter, to which a boat was suspended with two wings, and a rudder to steer it by, as also a large parachute spread horizontally between the boat and the balloon, designed to check the fall in case the balloon should burst. The machine being filled with inflammable air, he ascended, from the Champs de Mars at Paris, to the height of near ten thousand feet, or almost 2 miles; and after floating in the air for an hour and a quarter, he descended at Billancourt near Seve, having experienced by turns heat, cold, hunger, and an excessive drowsiness. It appears from his own account, and as might have been expected, that the wings and rudder of his boat had little or no power in turning the balloon from the direction of the wind.

In the course of this year, 1784, aerostatic experiments and aerial voyages became so frequent, that the limits of this article will not allow of any thing farther than mentioning those which were attended with any remarkable circumstances. On the 25th of April, Messrs. de Morveau and Bertrand ascended from Dijon, with an inflammable-air balloon, to the height of thirteen thousand feet, or near 2 miles and a half, where the thermometer marked 25 degrees. They were in the air 1 hour and 25 minutes, in which time they floated 18 miles.

On the 20th of May, four ladies and a gentleman ascended from Paris, in a large montgolfier, above the highest buildings, and remained suspended there a considerable time, the balloon being confined by ropes from flying away.

On the 23d of May, M. Blanchard, with the same balloon as before, ascended from Rouen, to such height that the mercury in the barometer stood at 20.57 inches, which on the earth had been at 30.16. It was observed that in this voyage M. Blanchard's wings or oars could not turn him aside from the direction of the wind.

On the 4th of June M. Fleurant and Madame Thible, the first lady who made an aerial voyage, ascended at Lyons in a machine of 70 feet diameter. They went to the height of 8500 feet, and floated about 2 miles in 45 minutes.

On the 14th of June, M. Coustard de Massi and M. Mouchet ascended at Nantes to a great height, with a balloon of  $32\frac{1}{2}$  feet diameter, filled with inflammable air extracted from zinc; and they floated to the distance of 27 miles in 58 minutes.

On the 23d of June, the first aerial traveller M. Pilatre de Rozier, accompanied with M. Prouts, ascended at Versailles, in the presence of the royal family and the king of Sweden, with a large montgolfier, whose diameter was 79 feet, and its height 91 feet and a half. They floated to the distance of 36 miles in three-quarters of an hour, when they descended, which is at the rate of 48 miles an hour. In consequence of this experiment the king granted to M. de Rozier a pension of 2000 livres.

On the 15th of July the duke of Chartres, the two brothers Roberts, and a fourth person ascended from the park of St. Cloud, with an inflammable-air machine, of an oblong form, its diameter being 34 feet, and its length, which went in a direction parallel to the horizon, was  $55\frac{1}{2}$  feet; and they remained in the atmosphere

for 45 minutes in the greatest fear and danger. The machine contained an interior small balloon, filled with common air, by means of which it was proposed to cause the machine to ascend or descend without the loss of any inflammable air or ballast; and the boat was furnished with a helm and oars, which were intended to guide it. Three minutes after ascending, the machine was lost in the clouds, and involved in a dense vapour. A violent agitation of the air, resembling a whirlwind, greatly alarmed the aeronauts, turned the machine three times round in a moment, and gave it such shocks as prevented them from using any of their instruments for managing the machine. After many struggles, with great difficulty they tore about 7 or 8 feet of the lower part of the covering, by which the inflammable air escaped, and they descended to the ground with great rapidity, though without any hurt. At the place of departure the barometer stood at 30.12 inches, and at their greatest elevation it stood at 24.36 inches; so that their ascent was about 5100 feet, or near one mile.

On the 18th of July, M. Blanchard, with a Mr. Boley, made his third voyage with the same balloon as he had before, and rose so high as to sink the barometer from 30.1 to 25.34 inches, answering to a height of about 4600 feet. In 2 hours and a quarter they floated 45 miles, which is at the rate of 20 miles an hour. In this voyage M. Blanchard pretended that he was able to turn the machine with his wings, and to make it ascend and descend at pleasure. After descending, it is said the balloon remained all the night at anchor full of air; and that the next day several ladies amused themselves by ascending successively to the height of 80 feet, the length of the ropes by which it was anchored.

In the course of this summer two persons had nearly lost their lives by ascending with machines of warmed air. The one in Spain, on the 5th of June, by the machine taking fire, was much burnt, and so hurt by the fall that his life was long despaired of. The latter having ascended a few feet, the machinery entangled under the eaves of a house, which broke the ropes, and the man fell about twenty feet: the machine presently took fire, and was consumed. Other montgolfiers were also burned about London, by taking fire, through the defects of their construction.

The first aerial voyage performed in England was by one Vincent Lunardi, a native of Italy, who ascended from the Artillery Ground, London, with an inflammable-air balloon on the 15th of September. His machine was made of oiled silk, painted in alternate stripes of blue and red; and its diameter was 33 feet. From a net, which covered about two-thirds of the balloon, 45 cords descended to a hoop hanging below the balloon, to which the gallery was attached. The machine had no valve; and its neck, which terminated in the form of a pear, was the aperture through which the inflammable air was introduced, and through which it might be let out. The balloon was filled with air produced from zinc by means of diluted vitriolic acid. And when the aeronaut departed, at 2 o'clock, he took up with him a dog, a cat, and a pigeon. After throwing out some sand to clear the houses, he ascended to a considerable height; and the direction of his motion at first was north-west by west; but



but as the balloon rose higher, it came into another current of air, which carried it nearly north. In the course of his voyage the thermometer was as low as 29 degrees, and the drops of water which had collected round the balloon were frozen. About half after three he descended very near the ground, and landed the cat, which was almost dead with cold: then rising, he prosecuted his voyage, till at 10 minutes past 4 o'clock he landed near Ware in Hertfordshire. He pretends that he descended by means of his oars or wings; but other circumstances related by him, strongly contradict the fact.

The longest and most interesting voyage performed about this time, was that of Messrs. Roberts and M. Colin Hullin, who ascended at Paris, at noon on the 19th of September, with an aerostat, filled with inflammable air, which was 27<sup>3</sup> feet in diameter, and 46<sup>3</sup> feet long, the machine being made to float with its longest part parallel to the horizon, and having a boat of near 17 feet long attached to it. The boat was fitted up with several wings or oars, shaped like an umbrella, and they ascended at 12 o'clock with 450 pounds of sand ballast, and after various manœuvres finally descended, at 40 minutes past 6 o'clock, near Arras in Artois, 150 miles from Paris, having still 200 pounds of ballast remaining in the boat. In one part they found the current of air uniform from 600 to 4200 feet high, which it seems was their greatest height, and the fall of the barometer had been near 5.6 degrees. They found that by means of their oars they could accelerate their course a little in the direction of the wind, when it moved slowly, which may be true; but there is great reason to doubt of the accuracy of their experiments by which they pretended to cause their path to deviate about 22 degrees from the wind, going with a considerable velocity.

The second aerial voyage in England, was performed by Mr. Blanchard, and Mr. Sheldon professor of anatomy to the Royal Academy, being the first Englishman who ascended with an aerostatic machine. They ascended at Chelsea the 16th of October, at 9 minutes past 12 o'clock. Mr. Blanchard having landed Mr. Sheldon at about 14 miles from Chelsea, re-ascended alone, and finally landed near Rumsey in Hampshire, about 75 miles distant from London, having gone nearly at the rate of 20 miles an hour. The wings used on this occasion it seems produced no deviation from the direction of the wind. Mr. Blanchard said that he ascended so high as to feel a great difficulty of breathing: and that a pigeon, which flew away from the boat, laboured for some time to sustain itself with its wings in the rarefied air, but after wandering a good while, returned, and rested on the side of the boat.

On the 4th of October, Mr. Sadler, an ingenious tradesman at Oxford, ascended at that place with an inflammable-air balloon of his own construction and filing. And again on the 12th of the same month he ascended at Oxford, and floated to the distance of 14 miles in 17 minutes, which is at the rate of near 50 miles an hour.

The 30th of November this year Mr. Blanchard's fifth aerial voyage, still with his old machine, was performed in company with Dr. J. Jeffries, a native of America. Their voyage was about 21 miles; and it

does not appear that the greatest action of their oars produced any effect in directing the course of the balloon.

On the 4th of January, 1785, a Mr. Harper ascended at Birmingham with an inflammable-air balloon, and went to the distance of 50 miles in an hour and a quarter, and suffered no other inconvenience than a temporary deafness, and what might be expected from the changes of wet and cold. The thermometer descended from 40 to 28 degrees.

On the 7th of January, Mr. Blanchard, accompanied with Dr. Jeffries, performed his sixth aerial voyage, by actually crossing the British channel from Dover to Calais, with the same balloon which had five times before carried him successfully through the air. They ascended with only 30 pounds of sand ballast, besides their provisions, some books, instruments, and other necessaries. The machine parted with the gas very rapidly, and their ballast was soon all exhausted; after which, from time to time they threw out every thing else in the boat, to prevent themselves from dropping into the sea. In this way they disposed of all their provisions, their books and instruments, and finally the most part of their very clothes themselves. This however bringing them near the French coast, they gradually ascended, cleared the cliffs and houses, and landed in the forest of Guiennes. It is remarkable that a bottle, being thrown out when they were in danger of falling into the sea, struck the water with such force, that they heard and felt the shock very sensibly on the car and balloon. In consequence of this voyage the king of France presented M. Blanchard with a gift of 12000 livres, and granted him a pension of 1200 livres a year.

On the 19th of January, Mr. Crosbie ascended at Dublin in Ireland, with an inflammable-air balloon to a great height. He rose so rapidly that he was out of sight in 3 minutes and a half. By suddenly opening the valve he descended just at the edge of the sea, as he was driving towards the channel, being unprovided for properly passing over to England.

On the 23d of March, Count Zambeccari and Admiral Sir Edward Vernon ascended at London, and sailed to Horsham in Sussex, at the distance of 35 miles in less than an hour. The voyage proved very dangerous, owing to some of the machinery about the valve being damaged, which obliged them to cut open some part of the balloon when they were about two miles perpendicular height above the earth, the barometer having fallen from 30.4 to 20.8 inches. In descending they passed through a dense cloud, which felt very cold, and covered them with snow. The observations they made were, that the balloon kept perpetually turning round its vertical axis, sometimes so rapidly as to make each revolution in 4 or 5 seconds; that a peculiar noise, like rustling, was heard among the clouds, and that the balloon was greatly agitated in the descent.

On May the 5th, Mr. Sadler, and William Windham, esq. member of parliament for Norwich, ascended at Moulsey-hurst. The machine took a south-east course, and the current of air was so strong that they were in great danger of being driven to sea. They had the good fortune however to descend near the con-



flux of the Thames and Medway, not a mile from the water's edge. By an accident they lost their balloon: for while the aeronauts were busied in securing their instruments, the country people, whom they had employed in holding down the machine, suddenly let go the cords, when the balloon instantly ascended, and was driven many miles out to sea, where it fell, and was taken up by a trading vessel. It was afterwards restored again, and another voyage made with it from Manchester to Pontefract, in which Mr. Sadler was still more unfortunate; for no person being near when it descended, and not being able to confine it by his own strength, he was dragged by it over trees and hedges; and at last was forced to quit it at the utmost peril of his life; after which it rose, and was out of sight in a few minutes. It was afterwards found near Gainsborough.

On the 12th of May Mr. Crosbie ascended, at Dublin, as high as the tops of the houses; but soon descended again with a velocity that alarmed all the spectators for his safety. On his stepping out of the car, in an instant Mr. McGuire, a college youth, sprung into it, and the balloon ascended with him to the astonishment of the beholders, and presently he was carried with great velocity towards the channel in the direction of Holyhead. This being observed, a crowd of horsemen pursued full speed the course he seemed to take, and could plainly perceive the balloon descending into the sea. Lord H. Fitzgerald, who was among the foremost, instantly dispatched a swift-sailing vessel mounted with oars, and all the boats that could be got, to the relief of the gallant youth; whom they found almost spent with swimming, just time enough to save his life.

The fate of M. Pilatre de Rozier, the first aerial traveller, and his companion M. Romain, has been much lamented. They ascended at Boulogne the 15th of June, with intent to cross the channel to England: for the first 20 minutes they seemed to take the proper direction; when presently the whole apparatus was seen in flames, and the unfortunate adventurers fell to the ground from the height of more than a thousand yards, and were killed on the spot, their bones being broken, and their bodies crushed in a shocking manner. The machine in which they ascended, consisted of a spherical balloon, 37 feet in diameter, filled with inflammable air; and under this balloon was suspended a small montgolfier, or fire balloon, of 10 feet diameter; the gallery which suspended the aeronauts, was attached to the net of the upper balloon by cords, which were fastened to a hoop rather larger than the montgolfier, and descended perpendicularly to the gallery. The montgolfier was intended to promote and prolong the ascension, by rarefying the atmospheric air, and by that means gaining levity. It is not certainly known whether the balloon was actually set on fire by the montgolfier, or, being over-rarefied by the heat beneath, burst, and by that means the inflammable air was set in a blaze.

On the 19th of July, at 20 minutes past 2 o'clock, Mr. Crosbie ascended at Dublin, with intent to cross the channel to Holyhead in England. The usual form of the boat had been changed, for a capacious wicker basket, of a circular form, round the upper

edges of which were fastened a great many bladders, which were intended to render his gallery buoyant, in case of a disaster at sea. About 300 pounds of ballast were put into the basket, but the aeronaut discharged half a hundred on his first rise. At first the current of air carried him due west; but it soon changed his course to nearly north-east, pointing nearly towards Whitehaven. At upwards of 40 miles from the Irish shore, he found himself within clear sight of both lands, and he said it was impossible to give any adequate idea of the unspeakable beauties which the scenery of the sea, bounded by both lands, presented. He rose at one time so high, that by the intense cold his ink was frozen, and the mercury sunk quite into the ball of the thermometer. He was sick, and felt a strong repulsion on the tympanum of the ears. At his utmost height he thought himself stationary; but on liberating some gas, he descended to a current of air blowing north, and extremely rough. He now entered a thick cloud, and encountered strong blasts of wind, with thunder and lightning, which brought him rapidly towards the surface of the water. Here the balloon made a circuit, but falling lower, the water entered his car, and he lost his notes of observation. All his endeavours to throw out ballast were of no avail; the force of the wind plunged him into the ocean; and with much difficulty he put on his cork jacket. The propriety of his idea was now very manifest in the construction of his boat; as by the admission of the water into the lower part of it, and the suspension of his bladders, which were arranged at the top, the water, added to his own weight, became proper ballast; and the balloon maintaining its poise, it became a powerful sail, by means of which, and a snatch block to his car, he went before the wind as regularly as a sailing vessel. In this situation he found himself inclined to eat, and he took a little fowl. At the distance of a league he discovered some vessels crowding after him; but as his progress outstripped all their endeavours, he lengthened the space of the balloon from the car, which gave a check to the rapidity of his sailing, and he was at length overtaken and saved by the Dunleary barge, which took him on board, and steered to Dunleary, towing the balloon after them.

A similar accident happened to Major Money, who ascended at Norwich, on the 22d of July, at 20 minutes past 4 in the afternoon; when meeting with an improper current, and not being able to let himself down, on account of the smallness of the valve, he was driven out to sea, where, after blowing about for near two hours, he dropped into the water. Here the struggles were astonishing which he made to keep the balloon up, which was torn, and hung only like an umbrella over his head. A ship was once within a mile, but, he adds, whether from want of humanity, or by mistaking the balloon for a sea monster, they sheered off, and left him to his fate: but a boat chased him for two hours, till just dark, and then bore away. He now gave up all hopes, and began to wish that providence had given him the fate of Pilatre de Rozier, rather than such a lingering death. Exerting himself however to preserve life as long as possible, by keeping the balloon floating over his head, to keep himself out of the water, into which nevertheless he sunk gradually inch



by inch, as it lost its power, till he was at length breast deep in water, when he was providentially taken up by a revenue cutter, at half past eleven at night, but so weak that he was obliged to be lifted out of the car into the ship.

About the latter end of August, the longest aerial voyage hitherto made, was performed by Mr. Blanchard, who ascended at Lille, accompanied by the Chevalier de L'Epinaud, and travelled 300 miles in their balloon before it descended. On this occasion, as on some former ones, Mr. Blanchard made trial of a parachute, like a large umbrella, invented to break the fall in case of an accident happening the balloon: with this machine he dropped a dog from the car soon after his ascension, which descended gently and unhurt.

On September the 8th, Thomas Baldwin, Esq. ascended from the city of Chester, at 40 minutes past one o'clock, and descended at Rixton-Moss, at 25 miles distance, after a voyage of 2 hours and a quarter. The greatest perpendicular altitude ascended was about a mile and a half, and the aeronaut computed that in some parts of the voyage he moved at the rate of 30 miles an hour. Mr. Baldwin published a very circumstantial account of his voyage, with many ingenious philosophical remarks relating to aerostation, of which subject his book may be considered as one of the best treatises yet given to the public.

October the 5th, Mr. Lunardi made the first aerial voyage in Scotland. He ascended at Edinburgh, and after various turnings, landed near Cupar in Fife, having described a track of 40 miles over the sea, and 10 over the land, in an hour and a half. He said the mercury in the barometer sunk as low as 18.3 inches at his greatest elevation.

November the 19th the celebrated Blanchard ascended at Ghent to a great height, and after many dangers descended at Delft without his car, which he cut away to lighten the machine when he was descending too rapidly, and flung himself by the cords to the balloon, which served him then in the nature of a parachute. On his first ascent, when he was almost out of sight, he let down a dog, by means of a parachute, which came easily to the ground.

November the 25th Mr. Lunardi ascended at Glasgow, and in two hours it is said he described a track of 125 miles. It is further remarkable that, being overcome with drowsiness, he says he slept for about 20 minutes in the bottom of the car, during this voyage.

Many other voyages were made in different countries, and with various success. But since the year 1785, the rage for balloons has considerably abated, and we have gradually had less and less of these aerial excursions, so that it is now become rather an uncommon thing to hear of one of them performed in any country whatever: which speedy decline in this new art is perhaps to be ascribed chiefly to the following causes; namely, a less degree of eagerness in people to pursue such experiments, from their curiosity having been satisfied; secondly, the trouble, danger, and great expence, attending them; and lastly, the want of the means of conducting them, and the small degree of utility to which they have hitherto been applied. The failure in the many attempts that have been made to direct balloons at pleasure through the air, cannot but

be felt as a very discouraging circumstance: and it is to be feared that it will ever be felt as such, notwithstanding the pretensions of some persons on this head; for they never have caused, nor is it to be expected they ever can cause the machine to deviate sensibly from the course of the wind, except only in the case when this moves with a very small celerity. For when the current blows only at the rate of 10 miles an hour, which is but a very gentle wind, it may be shewn that a balloon of 50 feet in diameter will require a force equal to the pressure of 72 pounds weight, to cause it to deviate 30 degrees from the course of the wind; and a force equal to double or triple that weight, when the wind blows with a double or triple velocity, that is, at the rate of 20 or 30 miles an hour; and so in proportion. To obviate the danger of a fall, arising from any accident happening to the balloon, some experiments have been made with a parachute, chiefly by Mr. Blanchard, whose endeavours and perseverance it seems have continued longer than in any other person: we still hear of his excursions in different parts of Europe, and improvements of the parachute, wings, &c; and have just read accounts of two voyages lately performed by him; with which, being very curious, we shall conclude our narration of these aerial excursions. They will be best related in Mr. Blanchard's own words, taken from his letter, dated Leipzick, October the 9th, 1787, to the editors of the Paris Journal. "I did not mention," says he, "in your interesting paper, my ascension at Strasburg on the 26th of last August: the weather was so horrible that I mounted only for the sake of contenting the astonishing crowd of strangers assembled there from all parts of the country. Every body seemed satisfied at the attempt, but I assure you, gentlemen, that I was far from being pleased with so common an experiment. The only remarkable thing that occurred at that time, was the following circumstance: At the height of about 2000 yards, or a mile and half a quarter, I let down a dog tied to the parachute, who, instead of descending gently, was forcibly carried, by a whirlwind, above the clouds. I met him soon after, bending his course directly downwards, and, as on recollecting his master, he began to bark a little, I was going to take hold of the parachute, when another whirlwind lifted him again to a great height. I lost him for the space of six minutes, and perceived him afterwards, with my telescope, as if sleeping in the cradle or basket belonging to the machine. Continually agitated, and impetuously tossed through every point of the compass, by the violence of the different currents of air, I determined to end my voyage on the other side of the Rhine, after having passed vertically over Zell. I descended at a small village, with an intention to be assisted a little, and about thirty men soon came within reach of the balloon very a-propos, and fixed me to the ground. The wind was so violent that anchors or ropes would have been of no service. I had however added to the large aerostatic globe a smaller one, of 60 pounds ascensional force, which would have contributed to fix me, when once I let it loose; but notwithstanding this precaution, the men's assistance was very necessary to me. The parachute was still wavering in the air, and did not come down till 12 minutes after."

"I per-



" I performed my 27th ascension at Leipfick the 29th of September, in the midst of an incredible number of spectators, forming one of the most brilliant assemblies I ever beheld. The sky was as clear and serene as possible, and the air so calm that many of my friends, and multitudes of others, could follow me on horseback, and even on foot. I was sometimes so near them that they thought they could reach me, but I could soon find the means of rising; and once, when they had actually taken hold of the cords, to see me float with the strings in their hands, I suddenly cut them, and mounted again in the air. All these amusing evolutions were in sight of the town and its environs. At length I yielded to the earnest solicitations of the company, and entered the town triumphantly in my car, followed by a concourse of people transported with joy, and amidst the acclamations of thousands. The next day I emptied the inflammable air into another globe, with which I intended to try some experiments; and I let it off with a cradle, in which a dog was fixed. The balloon, having reached a considerable height, made an explosion in its under part, as I had imagined it would, having previously disposed it in a proper manner for that purpose; by which means the little animal fell gently to the ground."

" The day before yesterday having repeated this experiment, at the town's request, I prepared the globe in such a manner as to cause an explosion in its upper part, and added a parachute with two small dogs fixed to it. They went so high that, notwithstanding the serenity of the sky, the balloon was lost in its immense expanse. Telescopes of the best sort became useless, and I began to be apprehensive for the death of the little animals, on account of the severity of the cold. They descended however about two hours after, quite safe and well, in the town of Delitzsch, three miles from Leipfick. I went yesterday to claim them, and found them again over the town in the air with the parachute. Such experiments had been already tried many times in the course of the day, and some officers had thrown them from the top of a steeple, in the sight of all the inhabitants of Delitzsch, from whence they descended safe."

We have lately heard of Mr. Blanchard's 32d ascension at Brunswick in the month of August 1788, in which he much assisted his ascent by means of his wings.

For several figures of balloons, see plate 1.

*Practice of AEROSTATION.* The first consideration in the practice of aerostation, is the form and the size of the machine. Various shapes have been tried and proposed, but the globular, or the egg-like figure, is the most proper and convenient, for all purposes; and this form also will require less cloth or silk than any other shape of the same capacity; so that it will both come cheaper, and have a greater power of ascension. The bag or cover of an inflammable-air balloon, is best made of the silk stuff called lustring, varnished over. But for a montgolfier, or heated-air balloon, on account of its great size, linen cloth has been used, lined within or without with paper, and varnished. Small balloons are made either of varnished paper, or simply of paper unvarnished, or of gold beater's skin, or such-like light substances.

With respect to the form of a balloon, it will be necessary that the operator remember the common proportions between the diameters, circumferences, surfaces, and solidities of spheres; for instance, that of different spheres, the circumferences are as the diameters; that the surfaces are as the squares of the diameters; and the solidities as the cubes of the same diameters: that any diameter is to its circumference as 7 to 22, or as 1 to  $3\frac{1}{7}$ ; and therefore 3 times and  $\frac{1}{7}$  of any diameter will be its circumference; so that if the diameter of a balloon be 35 feet, its circumference will be 110 feet. And if the diameter be multiplied by the circumference, the product will be the surface of the sphere; thus 35 multiplied by 110 gives 3850, which is the surface of the same sphere in square feet: and if this surface be divided by the breadth of the stuff, in feet, which the balloon is to be made of, the quotient will be the number of feet in length necessary to construct the balloon; so if the stuff be 3 feet wide, then 3850 divided by 3, gives  $1283\frac{1}{3}$  feet, or 428 yards nearly, the requisite quantity of stuff of 3 feet or one yard wide, to form the balloon of 35 feet diameter. Hence also, by knowing the weight of a given piece of the stuff, as of a square foot, or square yard, it is easy to find the weight of the whole bag, namely by multiplying the surface, in square feet or yards, by the weight of a square foot or yard: so if each square yard weigh 16 ounces or 1 pound, then the whole bag will weigh 428 pounds. Again, the capacity, or solid contents, of the sphere, will be found by multiplying  $\frac{2}{3}$  of the surface by the diameter, or by taking  $\frac{1}{6}$  of the cube of the diameter; which gives 22458 cubic feet for the capacity of the said balloon, that is, it will contain, or displace, 22458 cubic feet of air. From the content and surface of the balloon, so found, is to be derived its power or levity, thus: on an average, a cubic foot of common air weighs  $1\frac{1}{2}$  ounce, and therefore to the number 22458, which is the content of our balloon, adding its  $\frac{1}{2}$ th part, we have 26950 ounces, or 1684 pounds, for the weight of the common air displaced or occupied by the balloon. From this weight must be deducted the weight of the bag, namely 428 pounds, and then there remains 1256 pounds levity of the balloon, without however considering the contained air, whether it be heated air, or of the inflammable kind. If inflammable air be used, as it is of different weights, from  $\frac{1}{4}$  to  $\frac{1}{8}$  or  $\frac{1}{12}$  the weight of common air, according to the modes of preparing it, let us suppose for instance that it is  $\frac{1}{8}$  of the weight of common air; then  $\frac{1}{8}$  of 1684 is 210 pounds, which is the weight of the bag full of that air; which being taken from 1256, leaves 995 pounds for the levity of the balloon when so filled with that inflammable air, or the weight which it will carry up, consisting of the car, the ropes, the passengers, the necessaries, and ballast. But if heated air be used; then as it is known from experiment that, by heating, the contained air is diminished in density about one-third only, therefore from 1684, take  $\frac{1}{3}$  of itself, and there remains 1123 for the weight of the contained warm air; and this being subtracted from 1256, leaves only 133 pounds for the levity of the balloon in this case; which being too small to carry up the car, passengers, &c, it shews that for those purposes a larger balloon is necessary, on



Montgolfier's principles. But if now, from the preceding computation, it be required to find how much the size of the balloon must be increased, that its levity, or power of ascension, may be equal to any given weight, as suppose 1000 pounds; then because the levities are nearly as the cubes of the diameters, therefore the diameters will be nearly as the cube roots of the levities; but the levities 133 and 1000 are nearly as 1 to 8, the cube roots of which are as 1 to 2, and consequently  $1 : 2 :: 35 : 70$  feet, the diameter of a montgolfier, made of the same thickness of stuff as the former, capable of lifting 1000 pounds.


On the same principles we can easily find the size of a balloon that shall just float in air when made of stuff of a given thickness or weight, and filled with air of a given density; the rule for which is this: from the weight of a cubic foot of common air, subtract that of a cubic foot of the lighter or contained air; then divide 6 times the weight of a square foot of the stuff, by the remainder; and the quotient will be the diameter, in feet, of the balloon that will just float at the surface of the earth. Suppose, for instance, that the materials are as before, namely, the stuff 1 pound to the square yard, or  $\frac{1}{9}$  ounces to the square foot, which taken 6 times is  $\frac{2}{3}$ ; then the cubic foot of common air weighing  $1\frac{1}{5}$  ounce, and of heated air  $\frac{2}{3}$  of the same, whose difference is  $\frac{2}{5}$ ; therefore  $\frac{2}{3}$  divided by  $\frac{2}{5}$ , gives  $26\frac{2}{3}$  feet, which is the diameter of a montgolfier that will just float: but if inflammable air be used of  $\frac{2}{3}$  the weight of common air, the difference between  $1\frac{1}{5}$  and  $\frac{2}{3}$  of it, is 1; by which dividing  $\frac{2}{3}$  or  $10\frac{2}{3}$ , the quotient is the same  $10\frac{2}{3}$  feet, which therefore is the diameter of an inflammable-air balloon that will just float. And if the diameter be more than these dimensions, the balloons will rise up into the atmosphere.

The height nearly to which a given balloon will rise in the atmosphere, may be thus found, having given only the diameter of the balloon, and the weight which just balances it, or that is just necessary to keep it from rising: compute the capacity or content of the globe in cubic feet, and divide its restraining weight in ounces by that content, and the quotient will be the difference between the density or specific gravity of the atmosphere at the earth's surface and that at the height to which the balloon will rise; therefore subtract that difference or quotient from  $1\frac{1}{5}$  or 1.2, the density at the earth, and the remainder will be the density at that height: then the height answering to that density will be found sufficiently near in the annexed table. Thus, in the foregoing examples, in which the diameter of the balloon is 35 feet, its capacity 22458, and the levity of the first one 995 pounds, or 15920 ounces, the quotient of the latter number divided by the former, is .709, which is the density at the utmost height, and to which in the table answers a little more than  $2\frac{1}{2}$  miles, or  $2\frac{5}{8}$  miles nearly, which therefore is the height to which the balloon will ascend. And when the same balloon was filled with heated air, its levity was found equal to only 133 pounds, or 2128

Height in miles.	Density.
0	1.200
$\frac{1}{4}$	1.141
$\frac{1}{2}$	1.085
$\frac{3}{4}$	1.031
1	0.980
$1\frac{1}{4}$	0.932
$1\frac{1}{2}$	0.886
$1\frac{3}{4}$	0.842
2	0.800
$2\frac{1}{4}$	0.761
$2\frac{1}{2}$	0.723
$2\frac{3}{4}$	0.687
3	0.653

ounces, then dividing this by 22458 the capacity, the quotient .095 taken from 1.200, leaves 1.105 for the density; to which in the table corresponds almost half a mile, or nearer  $\frac{3}{8}$  of a mile. And so high nearly would these balloons ascend, if they keep the same figure, and lose none of the contained air: or rather, those are the heights they would settle at; for their acquired velocity would first carry them above that height, so far as till all their motion should be destroyed; then they would descend and pass below that height, but not so much as they had gone above; after which they would re-ascend, and pass that height again, but not so far as they had gone below it; and so on for many times, vibrating alternately above and below that point, but always less and less every time. The foregoing rule, for finding the height to which the balloon will ascend, is independent of the different states of the thermometer at that highest point, and at the surface of the earth; but for greater accuracy, including the allowances depending on the different states of the thermometer, see under the word ATMOSPHERE, where the more accurate rules are given at large.

The best way to make up the whole coating of the balloon, is by different pieces or slips joined lengthways from end to end, like the pieces composing the surface of a geographical globe, and contained between one meridian and another, or like the slices into which a melon is usually cut, and supposed to be spread flat out. Now the edges of such pieces cannot be exactly described by a pair of compasses, not being circular, but flatter or less round than circular arches; but if the slips are sufficiently narrow, or numerous, they will differ the less from circles, and may be described as such. But more accurately, the breadths of the slip, at the several distances from the point to the middle, where it is broadest, are directly as the sines of those distances, radius being the sine of the half length of the slip, or of the distance of either point from the middle of the slip: that is, If ACBD be one of the slips, AB being half the circumference, or AE a quadrant conceived to be equal to AC or AD; then will CD be to  $ab$ , as radius or the sine of AC, to the sine of Aa. So that if the quadrant AE or AC be divided into any number of equal parts, as here suppose 9, then divide the quadrant or 90 degrees by the number of parts 9, and the quotient 10 is the number of degrees in each part; and hence the arcs AC, Aa, Ac, &c, will be respectively  $90^\circ$ ,  $80^\circ$ ,  $70^\circ$ , &c; and CD being radius, the several breadths  $ab$ ,  $cd$ ,  $ef$ , &c, will be respectively the sines of  $80^\circ$ ,  $70^\circ$ ,  $60^\circ$ , &c, which are here placed opposite them, the radius being 1. Therefore when it is proposed to cut out slips for a globe of a given diameter; compute the circumference, and make AE or AC a quarter of that circumference, and CD of any breadth, as 3 feet, or 2 feet, or any other quantity;





quantity; then multiply each of the decimal numbers, set opposite the figure, by that quantity, or breadth of CD, so shall the products be the several breadths *ab*, *cd*, *ef*, &c.

Various schemes have been devised for conducting balloons in any direction, whether vertical or sideways. As to the vertical directions, namely upwards or downwards, the means are obvious, viz. in order to ascend, the aeronaut throws out some ballast; and that he may descend, he opens a valve in the top of his machine by means of a string, to let some of the gas escape; or if it be a montgolfier, he increases or diminishes the fire, as he would ascend or descend. But to direct the machine in a side or horizontal course, is a very difficult operation, and what has hitherto not been accomplished, except in a small degree, and when the current of air is very gentle indeed. The difficulty of moving the balloon sideways, arises from the want of wind blowing upon it; for as it floats along with the current of air, it is relatively in a calm, and the aeronaut feels no more wind than if the machine were at rest in a perfect calm. For this reason, any thing in the nature of sails, can be of no use; and all that can be hoped for, is to be attempted by means of oars; and how small the effect of these must be, may easily be conceived from the rarity of the medium against which they must act, and the great magnitude of the machine to be forced through it. We can easily assign what force is necessary to move a given machine in the air with any proposed velocity. From very accurate experiments I have determined, that a globe of  $6\frac{1}{2}$  inches in diameter, and moving with a velocity of 20 feet per second of time, suffers a resistance from the air which is just equal to the weight or pressure of one ounce Averdupois; and farther that with different surfaces, and the same velocity, the resistances are directly proportional to the surface nearly, a double surface having a double resistance, a triple surface a triple resistance, and so on; and also that with different velocities, the resistances are proportional to the squares of the velocities nearly, so that a double velocity produced a quadruple resistance, and three times the velocity nine times the resistance, and so on. And hence we can assign the resistance to move a given balloon, with any velocity. Thus, take the balloon as before of 35 feet diameter; then by comparison as above it is found that this globe, if moved with the velocity of 20 feet per second, or almost 14 miles per hour, will suffer a resistance equal to 271 pounds; to move it at the rate of 7 miles an hour, the resistance will be 68 pounds; and to move it  $3\frac{1}{2}$  miles an hour, the resistance will be 17 pounds; and so on: and with such force must the aeronauts act on the air in a contrary direction, to communicate such a motion to the machine. And if the balloon move through a rarer part of the atmosphere, than that at the surface of the earth, as  $\frac{1}{2}$ , or  $\frac{1}{4}$ , &c, rarer, and consequently the resistance be less in the same proportion; yet the force of the oars will be diminished as much; and therefore the same difficulty still remains. In general, the aeronaut must strike the air, by means of his oars, with a force just equal to the resistance of the air on the balloon, and therefore he must strike that air with a velocity which must be greater as the surface of the oar is less

than the resisted surface of the globe, but not in the same proportion, because the force is as the square of the velocity.

Now suppose the aeronaut act with an oar equal to 100 square feet of surface, to move the balloon above mentioned at the rate of 20 feet per second, or 14 miles an hour; then must he move this oar with the great velocity of 62 feet per second, or near 43 miles per hour: and so in proportion for other velocities of the balloon. From whence it is highly probable, that it will never be in the power of man to guide such machine with any tolerable degree of success, especially when any considerable wind blows, which is almost always the case.

As some aeronauts have thought of using parachutes, made something like umbrellas, to break their fall, in case of any accident happening to the balloon, we shall here consider the principles and power of such a machine. Let us suppose a person wants to know what the size of a parachute must be, that he may descend with it at the uniform rate of 10 feet in a second, which is nearly equal to the velocity he acquires by falling or leaping from the height only of 17 inches, and which it is presumed he may do with safety. Now in order to descend with any uniform velocity, the resistance of the air must be equal to the whole weight that descends: then suppose the weight of the aeronaut to be 150 pounds, and that the parachute is flat, and circular, and made of such materials as that every square foot of its surface weighs 2 ounces, and farther that the weight increases in the same proportion as the surface; then the diameter of the parachute necessary to descend with the moderate velocity of 10 feet per second, must be upwards of 78 feet in diameter: but if the parachute be not a flat surface, but concave on the lower side, its power will be rather the greater, and the diameter may be somewhat less. If it be required to know the power of a flat circular parachute, or what resistance it meets with from air of a mean density, when descending with a given velocity; say as the number 800 is to the square of the velocity in feet, so is the square of the diameter in feet, to a fourth number, which will be the resistance in pounds. And hence, if it be required to know with what velocity a parachute will descend with a given weight; say as the given diameter is to the square root of the weight, so is the number  $28\frac{1}{2}$  to a fourth term, which will be the velocity when the descent is in air of a mean density. So if the diameter of a balloon be 50, and its weight together with that of a man be 530 pounds, the square root of which is 23 very nearly; then as  $50 : 23 :: 28\frac{1}{2} : 13$ , so that the man and parachute will descend with the velocity of 13 feet per second; which it is presumed he may safely do, as he would meet with a shock only equal to that of leaping freely from a height only of 2 feet 2 inches.

The methods of extracting inflammable air from various substances, for filling balloons, and for other purposes, may be seen under the words Air and Gas. And as to the methods of filling and constructing balloons, being matters merely mechanical, they are omitted in this place.

Ample information however on these, and many other particulars, may be met with in several books expressly written on the subject; as in Cavallo's History and



and Practice of Aeroſtation, 8vo, 1785; in Baldwin's Airopaidia, 8vo, 1786; &c.

It has often been diſcuſſed, ſays the former of theſe gentlemen, whether the preference ſhould be given to machines raiſed by inflammable air, or to thoſe raiſed by heated air. Each of them has its peculiar advantages and diſadvantages; a juſt conſideration of which ſeems to decide in favour of thoſe made with inflammable air. The principal comparative advantages of the other fort are, that they do not require to be made of ſo expenſive materials; that they are filled with little or no expence; and that the combuſtibles neceſſary to fill them are found almoſt every where; ſo that when the ſtock of fuel is exhausted, the aeronaut may deſcend and recruit it again, in order to proceed on his voyage. But then this fort of machines muſt be made larger than the other, to take up the ſame weight; and the preſence of a fire is a continual trouble, and a continual danger: in fact, among the many aerial voyages that have been made and attempted with ſuch machines, very few have ſucceeded without an inconvenience, or an accident; and ſome indeed have been attended with dangerous and even fatal conſequences; from which the other fort is in a great meaſure exempt. But, on the other hand, the inflammable air balloon muſt be made of a ſubſtance impermeable to the ſubtle gas: the gas itſelf cannot be produced without a conſiderable expence; and it is not eaſy to find the materials and apparatus neceſſary for the production of it in every place. However, it has been found that an inflammable-air balloon, of 30 feet in diameter, may be made ſo cloſe as to ſuſtain two perſons, and a conſiderable quantity of ballaſt, in the air for more than 24 hours, when properly managed; and poſſibly one man might be ſupported by the ſame machine for three days: and it is probable that the ſtuff for theſe balloons may be ſo far improved, as to be quite impermeable to the gas, or very nearly ſo; in which caſe, the machine, once filled, would continue to float for a long ſpace of time. At Paris they have already attained to a great degree of perfection in this point; and ſmall balloons have been kept floating in a room for many weeks, without loſing any conſiderable quantity of their levity: but the difficulty lies in the large machines: for in theſe, the weight of the ſtuff itſelf, with the weight and ſtreſs of the ropes and boat, and the folding them up, may eaſily crack and rub off the varniſh, and make them leaky.

In regard to philoſophical obſervations, derived from the new ſubject of aeroſtation, there have been very few made; the novelty of the diſcovery, and of the proſpect enjoyed from the car of an aeroſtatic machine, have commonly diſtracted the attention of the aeronauts; not to remark that many of the adventurers were inadequate to the purpoſe of making improvements in philoſophy, being moſtly influenced either by pecuniary motives, or the vanity of adding their names to the liſt of aerial travellers.—The agreeable ſtillneſs and tranquillity experienced aloft in the atmosphere, have been matter of general obſervation. Some machines have aſcended to a great height, as far, it has been ſaid, as two miles; and they have commonly paſſed through fogs and clouds, above which they have enjoyed the clear light and heat of the ſun, whiſt the earth beneath

was actually covered by denſe clouds, which poured down abundance of rain. In aſcending very high, the aeronauts have often experienced a pain in their ears, ariſing, it is ſuppoſed, from the internal air being not of the ſame denſity as the air without; but the pain uſually went off in a ſhort time: and it ſeems that this effect is ſimilar to what is experienced by perſons who deſcend by a diving-bell to conſiderable depths in the ſea: I remember often to have heard the late unfortunate Mr. Spalding, the celebrated diver, ſpeak of this effect, with a marked and philoſophical accuracy: after deſcending two or three fathoms below the ſurface, he began to feel a pain in his ears, which gradually increaſed to a very great degree if the deſcent was too quick; his method was therefore to deſcend ſlowly, and to make a ſtop for ſome minutes at the depth of 5 fathom, which is equal nearly to the preſſure of the atmosphere, and where conſequently the air in his bell was of double the denſity of common air at the ſurface; after reſting here awhile, his ears, as he expreſſed it, gave a crack, and he was ſuddenly relieved of the pain. He then deſcended 5 fathoms more, with the ſame ſymptoms, and the ſame effect: and ſo on continually, from one five fathoms to another, deſcending leiſurely, and ſtopping a little at each ſtage, to give time for his conſtitution to adapt itſelf to the degree of condenſation of the air; after which he felt no more inconvenience, till he came to aſcend again, which was performed with the ſame caution and circumſtances. One experiment is recorded, in which the air of a high region, being brought down, and examined by means of nitrous air, was found to be purer than the air below. The temperature of the upper regions too, it has been found, is much colder than that of the air near the earth; the thermometer, in ſome aeroſtatic machines, having deſcended many degrees below the freezing point of water, while it was conſiderably higher than that degree at the earth's ſurface.

ÆSTIVAL, ſee ESTIVAL.

ÆSTUARY, or ESTUARY, in *Geography*, an arm of the ſea, running up a good way into the land. Such as Briſtol channel, many of the friths in Scotland, and ſuch like.

ÆTHER, ſee ETHER.

AFFECTED, or ADFFECTED, EQUATION, in *Algebra*, is an equation in which the unknown quantity riſes to two or more ſeveral powers or degrees. Such, for example, is the equation  $x^3 - px^2 + qx = r$ , in which there are three different powers of  $x$ , namely,  $x^3$ ,  $x^2$ , and  $x$ .

The term, affected, is alſo uſed ſometimes in algebra, when ſpeaking of quantities that have co-efficients.

Thus in the quantity  $2a$ ,  $a$  is ſaid to be affected with the co-efficient 2.

It is alſo ſaid, that an algebraic quantity is affected with the ſign  $+$  or  $-$ , or with a radical ſign; meaning no more than that it has the ſign  $+$  or  $-$ , or that it includes a radical ſign.

The term adaffected, or affected, I think, was introduced by Vieta.

AFFECTION, in *Geometry*, a term uſed by ſome ancient writers, ſignifying the ſame as property.

AFFECTION. *Phyſ.* The affections of a body are certain modifications occaſioned or induced by motion.



motion; in virtue of which the body is disposed after such, or such a manner.

The affections of bodies, are sometimes divided into *primary* and *secondary*.

*Primary Affections*, are those which arise either out of the idea of matter, as magnitude, quantity, and figure; or out of the idea of form, as quality and power; or out of both, as motion, place, and time.

*Secondary, or derivative Affections*, are such as arise out of primary ones, as divisibility, continuity, contiguity, &c, which arise out of quantity; regularity, irregularity, &c, which arise out of figure, &c.

**AFFIRMATIVE QUANTITY, or POSITIVE QUANTITY**, one which is to be added, or taken effectually; in contradistinction to one that is to be subtracted, or taken defectively.—The term affirmative was introduced by Vieta.

**AFFIRMATIVE SIGN, or POSITIVE SIGN**, in *Algebra*, the sign of addition, thus marked  $+$ , and is called *plus*, or *more*, or *added to*. When set before any single quantity, it serves to denote that it is an affirmative or a positive quantity; when set between two or more quantities, it denotes their sum, shewing that the latter are to be added to the former. So  $+6$ , and  $+a$ , and  $+AB$ , are affirmative quantities; also  $+6 + 8 + 10$  denote the sum of 6, 8, and 10, which is 24, and are read thus, 6 plus 8 plus 10. Also  $a + b + c$  denote the sum of the quantities represented by  $a$ ,  $b$  and  $c$ , when added together. It seems now not easy to ascertain with certainty, when, or by whom, this sign was first introduced; but it was probably by the Germans, as I find it first used by Stifelius in his *Arithmetic*, printed in 1544.

The early writers on Algebra used the word *plus* in Latin, or *piu* in Italian, for addition, and afterwards the initial  $p$  only, as a contraction; like as they used *minus*, or *meno*, or the initial  $m$  only, for subtraction: and thus these operations were denoted in Italy by Lucas de Burgo, Tartalea, and Cardan, while the signs  $+$  and  $-$  were employed much about the same time in Germany by Stifelius, Scheubelius, and others, for the same operations.

**AGE**, in *Chronology*, is used for a century, being a system or a period of a hundred years.

Chronologists also divide the time since the creation of the world into three ages: The first, from Adam till Moses, which they call the age of nature; the second from Moses to Jesus Christ, called the age of the law; and the third, or age of grace, from Jesus Christ till the end of the world.

**AGE of the Moon**, in *Astronomy*, is the number of days elapsed since the last new moon. To find the moon's age, for any time nearly, for ordinary uses; add together the epact, the day of the current month, and the number of months from March to the present month inclusive; the sum is the moon's age: but if the sum exceed 29, deduct 29 from it in months that have 30 days, or 30 in those that have 31; and the remainder will be the age.—At the end of 19 years the moon's age returns upon the same day of the month, but falls a little short of the same hour of the day.

**AGENT, Agens**, in *Physics*, that by which a thing is done or effected; or any thing having a power by

which it acts on another, called the patient, or by its action induces some change in it.

**AGGREGATE**, the sum or result of several things added together. See *Sum*.

**AGITATION**, in *Physics*, a brisk intestine motion, excited among the particles of a body. Thus fire agitates the subtlest particles of bodies. Fermentations, and effervescences, are produced by a brisk agitation of the particles of the fermenting body.

**AGUILON (FRANCIS), or AQUILON**, was a jesuit of Brussels, and professor of philosophy at Doway, and of theology at Antwerp. He was one of the first that introduced mathematical studies into Flanders. He wrote a large work on *Optics*, in 6 books, which was published in folio, at Antwerp, in 1613; and a treatise of *Projections of the Sphere*. He promised also to treat upon *Catoptrics* and *Dioptrics*, but this was prevented by his death, which happened at Seville, in the year 1617.

**AIR**, in *Physics*, a thin, fluid, transparent, elastic, compressible, and dilatable body, which surrounds this terraqueous globe, and covers it to a considerable height.

Some of the ancients considered air as an element, namely, one of the four elements, air, earth, water, and fire, of which they conceived all bodies to be composed; and though it be certain that air, taken in the common acceptation, be far from the simplicity of an elementary substance, yet some of its parts may properly be so called. So that air may be distinguished into proper or elementary, and vulgar or heterogeneous.

**Elementary AIR**, or *Air* properly so called, is a subtle, homogeneous, elastic fluid; being the basis, or fundamental ingredient, of the whole air of the atmosphere, from which it takes its name. And in this sense Dr. Hales, and other modern philosophers, consider it as entering into the composition of most, or perhaps all bodies; existing in them in a solid state, devoid of its elasticity, and most of its distinguishing properties, and serving as their cement; but, by certain processes, capable of being disengaged from them, recovering its elasticity, and resembling the air of our atmosphere.

The particular nature of this aerial matter we know but little about: what authors have said concerning it being chiefly conjectural. There is no way of examining air pure and defecated from the several matters with which it is mixed; and consequently we cannot pronounce what are its peculiar properties, abstractedly from other bodies.

Dr. Hook, and some others, maintain that it is the same with the *ether*, or that imaginary fine, fluid, active matter, conceived to be diffused through the whole expanse of the celestial regions: which comes to much the same thing as Newton's subtle medium, or spirit. In this sense it is supposed to be a body *sui generis*, incorruptible, immutable, incapable of being generated, but present in all places, and in all bodies.

Other philosophers place its essence in elasticity, making that its distinctive character. These suppose that it may be generated, and that it is nothing else but the matter of other bodies, rendered by the changes



changes it has undergone, susceptible of a permanent elasticity. Mr. Boyle produces a number of experiments, which he made on the production of air, that is, according to him, the extraction of a sensible quantity of air from a body in which there appeared to be little or none at all, by whatever means this may be effected. He observes that among the different methods for this purpose, the chief are fermentation, corrosion, dissolution, decomposition, ebullition of water and other fluids, the reciprocal action of bodies, especially saline ones, upon one another; he adds, that different solid and mineral bodies, in the parts of which no elasticity could be conceived to exist, being plunged into corrosive mediums, which also are quite unelastic, will, by the attenuation of their parts from their mutual collision, produce a considerable quantity of elastic air.

Sir Isaac Newton is of the same opinion, according to whom the particles of a dense compact fixed substance, adhering to each other by a powerful attractive force, cannot be separated but by a violent heat, and perhaps never without fermentation; and these bodies, rarefied by heat and fermentation, are finally transformed into a truly permanent elastic air. On these principles, he adds, gunpowder produces air on explosion. Optics, Qu. 31, &c.

*Common, or heterogeneous AIR*, is an assemblage of corpuscles of various kinds, which together constitute one fluid mass, in which we live and move, and which we constantly breathe; which compound mass altogether, is called the atmosphere.

In this popular and extensive meaning of the term, Mr. Boyle acknowledges that air is the most heterogeneous body in the universe; and Boerhaave proves that it is an universal chaos, a mere jumble of all species of created things. Besides the matter of light or fire, which continually flows into it from the celestial bodies, and perhaps the magnetic effluvia of the earth, whatever fire can volatilize must be found in the air.

Hence, for instance, 1. All sorts of vegetable matter must be contained in the air; being either exhaled from plants growing all over the face of the earth, or rendered volatile by putrefaction, not excepting even the more solid and vascular parts of them.

2. It is no less certain that the air must contain particles of every substance belonging to the animal kingdom. For the copious emanations which are perpetually issuing from the bodies of animals, in the perspiration constantly kept up by the vital heat, are absorbed by the air; and in such quantities too, during the course of an animal life, that, could they be collected, they would be sufficient to compose a good round number of the like animals. And besides, when a dead animal continues exposed to the air, all its particles evaporate, and are quickly dissipated; so that the substance which composed the animal, is almost wholly incorporated with the air.

3. The whole fossil kingdom must necessarily be found in the atmosphere; for all of that kind, as salts, sulphurs, stones, metals, &c, are convertible into fume, and must consequently take place among aerial substances. Gold itself, the most fixed of all natural bodies, is found among ores, closely adhering to sulphurs in mines, and so is raised along with the mineral.

Of all the emanations which float in the vast ocean of the atmosphere, perhaps the principal are such as consist of saline particles. Many writers suppose that they are of a nitrous kind; but it is probable that they are of all sorts, as vitriol, alum, marine salt, and many others. And Mr. Boyle thinks that there may be great quantities of compound salts, not to be met with on or in the bowels of the earth, formed by the fortuitous concurrence and mixture of different saline spirits.

We often find the window-glass of old buildings corroded, as if eaten by worms; though we know of no particular salt that is capable of producing such an effect.

Sulphurs too must make a considerable portion of this compound mass, on account of the many volcanos, grotts, caverns, and mines, dispersed over the face of the globe.

Finally, the various attritions, separations, dissolutions, and other mutual operations of matter of different sorts upon one another, may be regarded as the sources of many other neutral, or anonymous bodies, unknown to us, which rise and float in the air.

Air, taken in this extensive sense, is one of the most general and considerable agents in nature; being concerned in the preservation of animal and vegetable life, and in the production of most of the phenomena that take place in the material world.

Its properties and effects, having been the principal objects of the researches and discoveries of modern philosophers, have been reduced to precise laws and demonstrations, forming no inconsiderable branch of mixed mathematics, under the titles of Pneumatics, Aerometry, &c.

*Mechanical Properties and Effects of AIR.* Of these the most considerable are its fluidity, its weight, and its elasticity.

1. *Its Fluidity.*—The great fluidity of the air is manifest from the great facility with which bodies traverse it; as in the propagation of, and easy conveyance it affords to, sounds, odours and other effluvia and emanations that escape from bodies: for these effects prove that it is a body whose parts give way to any force, and in yielding are easily moved amongst themselves; which is the definition of a fluid. That the air is a fluid is also proved from this circumstance, that it is found to exert an equal pressure in all directions; an effect which could not take place otherwise than from its extreme fluidity. Neither has it been found that the air can be deprived of this property, whether it be kept for many years together confined in glass vessels, or be exposed to the greatest natural or artificial cold, or condensed by the most powerful pressure; for in none of these circumstances has it ever been reduced to a solid state. It is true indeed that real permanent air may be extracted from solid bodies, and may also be absorbed by them; and we also know that in this case it must be exceedingly condensed, and reduced to a bulk many hundred times less than in its natural state: but in what form it exists in those bodies, or how their particles are combined together, is a mystery which remains hitherto inexplicable.

Those philosophers who, with the Cartesians, make fluidity to consist in a perpetual intestine motion of the parts, think they can prove that this character belongs to air: thus, in a darkened room, where the representations



tations of external objects are introduced by a single ray, the corpuscles with which the air is replet, are seen to be in a continual fluctuation. Some moderns attribute the fluidity of the air, to the fire which is intermixed with it; without which, say they, the whole atmosphere would harden into a solid impenetrable mass: and indeed it must be allowed that the more fire it contains, the greater will its fluidity, mobility, and permeability be; and according as the different positions of the sun augment or diminish the degree of fire, the air always receives a proportional temperature, and is kept in a continual reciprocation.

2. *Its Weight or Gravity.*—The weight or gravity of the air, is a property belonging to it as a body; for gravity is a property essential to matter, or at least a property found in all bodies. But independent of this, we have many direct proofs of its gravity from sense and experiment: thus, the hand laid close upon the end of a vessel, out of which the air is drawn at the other end, soon feels the load of the incumbent atmosphere: thus also, thin glass vessels, exhausted of their air, are easily crushed to pieces by the weight of the external air: and so two hollow segments of a sphere, 4 inches in diameter, exactly fitting each other, being emptied of air, are, by the weight of the ambient air, pressed together with a force which requires the weight of 188 pounds to separate them; and that they are thus forcibly held together by the pressure of the air, is made evident by suspending them in an exhausted receiver, for then they quickly separate of themselves, and fall asunder. Again, if a tube, close at one end, be filled with quicksilver, and the open end be immersed in a basin of the same fluid, and so held upright, the quicksilver in the tube will be kept raised up in it to the height of about 30 inches above the surface of that in the basin, being supported and balanced by the pressure of the external air upon that surface: and that this is the cause of the suspension of the quicksilver in the tube, is made evident by placing the whole apparatus under the receiver of an air-pump; for then the fluid will descend in the tube in proportion as the receiver is exhausted of its air; and then on gradually letting in the air again, the quicksilver reascends to its former height in the tube: and this is what is called, from its inventor, the Torricellian experiment. Nay farther, air can actually be weighed like any other body: for a rigid vessel, full even of common air, by a nice balance is found to weigh more than when the air is exhausted from it; and the effect is proportionally more sensible, if the vessel be weighed full of condensed air, and more still if it be weighed in a receiver void of air.

But although we have innumerable proofs of the gravitating property of the air, yet the full discovery of the laws and circumstances of it are certainly due to the moderns. It cannot indeed be denied, that several of the ancients had some confused notions about this property: thus Aristotle says that all the elements have gravity, and even air itself; and as a proof of it, says that a bladder inflated with air, weighs more than the same when empty; and Plutarch and Stobæus quote him as teaching that the air in its weight is between that of fire and of earth; and farther, he himself, treating of respiration, reports it as the opinion of

Empedocles, that he ascribes the cause of it to the weight of the air, which by its pressure forces itself into the lungs; and much in the same way are the sentiments of Asclepiades expressed by Plutarch, who represents him as saying, among other things, that the external air, by its weight, forcibly opened its way into the breast. But nevertheless it is certain, however unreasonable it may seem, that Aristotle's followers departed in this instance from their master, by asserting the contrary for many ages together. Indeed many of the phenomena arising from this property, have been remarked from the highest antiquity. Many centuries since, it was known that by sucking the air from an open pipe, having its extremity immersed in water, this fluid rises above its level, and occupies the place of the air. In consequence of such observations, sucking pumps were contrived, and various other hydraulic machines; as Heron's syphons, described in his *Spiritualia* or *Pneumatics*, and the watering-pots known in Aristotle's time under the name of *clepsydra*, which alternately stop or run as the finger closes or opens their upper orifice. Indeed the reason assigned, by philosophers many ages after, for this phenomenon, was a pretended horror that nature conceives for a vacuum, which, rather than endure it, makes a body ascend contrary to the powerful solicitation of its gravity. Even Galileo, with all his sagacity, could not for some time hit upon any thing more satisfactory; for he only assigned a limit to this dread of vacuity: having observed that sucking pumps would not raise water higher than 16 brasses, or 34 English feet, he limited this abhorring force of nature, to one that was equivalent to the weight of a column of water 34 feet high, on the same base as the void space. Consequently he pointed out a way of making a vacuum, by means of a hollow cylinder, whose piston is charged with a weight sufficient to detach it from the close bottom turned upwards: this effort he called the measure of the force of vacuity, and made use of it for explaining the cohesion of the parts of bodies.

Galileo however was well apprised of the weight of the air as a body: in his *Dialogues* he shews two ways of demonstrating it, by weighing it in bottles: the transition was easy from one discovery to another: yet still Galileo's knowledge of the matter was imperfect, that is, as to the particular instance of the suspension of a fluid above its level, by the pressure of the external air.

At length Torricelli fell upon the lucky guess, that the counterpoise which keeps fluids above their level, when nothing presses upon their internal surface, is the mass of air resting upon the external one. He discovered it in the following manner: In the year 1643 this disciple of Galileo, on occasion of executing an experiment on the vacuum formed in pumps, above the column of water, when it exceeds 34 feet, thought of using some heavier fluid, such as quicksilver. He conceived that whatever might be the cause by which a column of water of 34 feet high is sustained above its level, the same force would sustain a column of any other fluid, which weighed as much as that column of water, on the same base; whence he concluded that quicksilver, being about 14 times as heavy as water, would not be sustained higher than 29 or 30 inches.



He therefore took a glass tube of several feet in length, sealed it hermetically at one end, and filled it with quicksilver; then inverting it, and holding it upright, by pressing his finger against the lower or open orifice, he immersed that end in a vessel of quicksilver; then removing his finger, and suffering the fluid to run out, the event verified his conjecture; the quicksilver, faithful to the laws of hydrostatics, descended till the column of it was about 30 inches high above the surface of that in the vessel below. And hence Torricelli concluded that it was no other than the weight of the air incumbent on the surface of the external quicksilver, which counterbalanced the fluid contained in the tube.

By this experiment Torricelli not only proved, what Galileo had done before, that the air had weight, but also that it was its weight which kept water and quicksilver raised in pumps and tubes, and that the weight of the whole column of it was equal to that of a like column of quicksilver of 30 inches high, or of water 34 or 35 feet high; but he did not ascertain the weight of any particular quantity of it, as a gallon, or a cubic foot of it, nor its specific gravity to water, which had been done by Galileo, though to be sure with no great accuracy, for he only proved that water was more than 400 times heavier than air.

Torricelli's experiment became famous in a short time. Father Merfenne, who kept up a correspondence with most of the literati in Italy, was informed of it in 1644, and communicated it to those of France, who presently repeated the experiment: Messrs. Pascal and Petit made it first, and varied it several ways; which gave occasion to the ingenious treatise which Pascal published at 23 years of age, intitled *Experiences Nouvelles touchant la Vuide*. In this treatise indeed he makes use of the old principle of *fuga vacui*; but afterwards getting some notion of the weight of the air, he soon adopted Torricelli's idea, and devised several experiments to confirm it. One of these was to procure a vacuum above the reservoir of quicksilver; in which case he found the column sink down to the common level: but this appearing to him not sufficiently powerful to dissipate the prejudices of the ancient philosophy, he prevailed on M. Perier, his brother-in-law, to execute the famous experiment of Puy-de-Domme, who found that the height of the quicksilver half-way up the mountain was less, by some inches, than at the foot of it, and still less at the top: so that it was now put out of doubt that it was the weight of the atmosphere which counterpoised the quicksilver.

Des Cartes too had a right notion of this effect of the air, to sustain fluids above their level, as appears by some of his letters about this time, and some years before; and in one of those he lays claim to the idea of the Puy-de-Domme experiment: After having desired M. de Carcavi to inform him of the success of that experiment, which public rumour had advertised him had been made by M. Pascal himself, he adds, "I had reason to expect this from him, rather than from you, because I first proposed it to him two years since, assuring him at the same time, that although I had not tried it, yet I could not doubt of the consequence; but as he is a friend of M. Roberval, who professes himself no friend to me, I suppose he is guided

by that gentleman's passions." See more of this history under BAROMETER.

As to the actual weight of any given portion of common air, it seems that Galileo was the first who determined it experimentally; and he gives two different methods, in his Dialogues, for weighing it in bottles: he did not however perform the experiment very accurately, as he stated from the result that the gravity of water was to that of air rather above 400 to 1.

A quantity of air was next weighed by Merfenne in a very ingenious manner. His idea was to weigh a vessel both when full of air, and when emptied of it: to make the vacuum for this purpose, he knew no better way than by expelling the air out of an eolipile by heating it red hot: by weighing it both when cold and hot, he found a certain difference; which however was not the exact weight of that capacity of air, because the vacuum was not perfect. But by plunging the eolipile, when red hot, into water, just so much water entered as was equal in bulk to the air that had been expelled; then he took it out and weighed it with the water, which gave the weight of the same bulk of water; and on comparing this with the former difference, or weight of air expelled, he found their proportion to be as 1300 to 1. Which is as wide of the truth as Galileo's proportion, namely 400 to 1, but the contrary way. And it is remarkable that the mean between the two, namely 850 to 1, is very near the true proportion as settled by other more accurate experiments.

Mr. Boyle, by a more accurate experiment, found the proportion to be that of 938 to 1. And Mr. Hawksbee found it as 850 to 1, proceeding on the same principles as Merfenne, with a three-gallon glass bottle, but extracting the air out of it with the air pump, instead of expelling it by fire; the height of the barometer being at that time 29.7 inches. Also by other accurate experiments made before the Royal Society by Mr. Hawksbee, Dr. Halley, Mr. Cotes, and others, the proportion was always between 800 and 900 to 1, but rather nearer the latter, namely, being first found as 840 to 1, then as 852 to 1, and a third time as 860 to 1; the barometer then standing at  $29\frac{3}{4}$  inches, and the weather warm. Mr. Cavendish determines the ratio 800 to 1, the barometer being  $29\frac{3}{4}$ , and the thermometer at  $50^{\circ}$ ; and Sir George Shuckburgh, by a very accurate experiment, finds it 836 to 1, the barometer being at that time at 29.27, and the thermometer at  $51^{\circ}$ . And the medium of all these is about 832 or 833 to 1, when reduced to the pressure of 30 inches of the barometer, and the mean temperature  $55^{\circ}$  of the thermometer. Upon the whole therefore it may be safely concluded that, when the barometer is at 30 inches, and the thermometer at the mean temperature  $55^{\circ}$ , the density or gravity of water is to that of air, as  $833\frac{1}{3}$  to 1, that is as  $\frac{2500}{3}$  to 1, or as 2500 to 3; and that for any changes in the height of the barometer, the ratio varies proportionally; and also that the density of the air is altered by the  $\frac{1}{40}$ th part for every degree of the thermometer above or below temperate.

This number, which is a very good medium among them all, I have chosen with the fraction  $\frac{1}{3}$ , because it gives exactly  $1\frac{1}{2}$  ounce for the mean weight of a cubic foot



foot of air, the weight of the cubic foot of water being just 1000 ounces averdupois, and that of quicksilver equal to 13600 ounces.

*Air*, then, having been shewn to be a heavy fluid substance, the laws of its gravitation and pressure must be the same as those of water and other fluids; and consequently its pressure must be proportional to its perpendicular altitude. Which is exactly conformable to experiment; for on removing the Torricellian tube to different heights, where the column of air is shorter, the column of quicksilver which it sustains is shorter also, and that nearly at the rate of 100 feet for  $\frac{1}{25}$  of an inch of quicksilver. And on these principles depend the structure and use of the barometer.

And from the same principle it likewise follows that air, like other fluids, presses equally in all directions. And hence it happens that soft bodies endure this pressure without change of figure, and hard or brittle bodies without breaking; being equally pressed on all parts; but if the pressure be taken off, or diminished, on one side, the effect of it is immediately perceived on the other. See *ATMOSPHERE*, for the total quantity of effects and pressure, and the laws of different altitudes, &c.

From the weight and fluidity of the air, jointly considered, many effects and uses of it may easily be deduced. By the combination of these two qualities, it closely invests the earth, with all the bodies upon it, constringing and binding them down with a great force, namely a pressure equal to about 15 pounds upon every square inch. Hence, for example, it prevents the arterial vessels of plants and animals from being too much distended by the impetus of the circulating juices, or by the elastic force of the air so copiously abounding in them. For hence it happens, that on a diminution of the pressure of the air, in the operation of cupping, we see the parts of the body grow tumid, which causes an alteration in the circulation of the fluids in the capillary vessels. And the same cause hinders the fluids from transpiring through the pores of their containing vessels, which would otherwise cause the greatest debility, and often destroy the animal. To the same two qualities of the air, weight and fluidity, is owing the mixture of bodies contiguous to one another, especially fluids; for several liquids, as oils and salts, which readily mix of themselves in air, will not mix at all in vacuo. With many other natural phenomena.

3. *Elasticity*. Another quality of the air, from whence arise a multitude of effects, is its elasticity; a quality by which it yields to the pressure of any other bodies, by contracting its volume; and dilates and expands itself again on the removal or diminution of the pressure. This quality is the chief distinctive property of air, the other two being common to other fluids also.

Of this property we have innumerable instances. Thus, for example, a blown bladder being squeezed in the hand, we find a sensible resistance from the included air; and upon taking off the pressure, the compressed parts immediately restore themselves to their former round figure. And on this property of elasticity depend the structure and uses of the air-pump.

Every particle of air makes a continual effort to dilate itself, and so it acts forcibly against all the neigh-

bouring particles, which also exert the like force in return; but if their resistance happen to cease, or be weakened, the particle immediately expands to an immense extent. Hence it is that thin glass bubbles, or bladders, filled with air, and placed under the receiver of an air-pump, do, upon pumping out the air, burst asunder by the force of the air which they contain. So likewise a close flaccid bladder, containing only a small quantity of air, being put under the receiver, swells as the receiver is exhausted, and at length appears quite full. And the same thing happens by carrying the flaccid bladder to the top of a very high mountain.

The same experiment shews that this elastic property of the air is very different from the elasticity of solid bodies, and that these are dilated after a different manner from the air. For when air ceases to be compressed, it not only dilates, but then occupies a far greater space, and exists under a volume immensely greater than before; whereas solid elastic bodies only resume the figure they had before they were compressed.

It is plain that the weight or pressure of the air does not at all depend on its elasticity, and that it is neither more nor less heavy than if it were not at all elastic. But from its being elastic, it follows that it is susceptible of a pressure, which reduces it to such a space, that the force of its elasticity, which re-acts against the pressing weight, is exactly equal to that weight. Now the law of the elasticity is such, that it increases in proportion to the density of the air, and that its density increases in proportion to the forces or weights which compress it. But there is a necessary equality between action and re-action; that is, the gravity of the air, which effects its compression, and the elasticity of it, which gives it its tendency to expansion, are equal.

So that, the elasticity increasing or diminishing, in the same proportion as the density increases or diminishes, that is, as the distance between its particles decrease or increase; it is no matter whether the air be compressed, and retained in any space, by the weight of the atmosphere, or by any other cause; as in either case it must endeavour to expand with the same force. And therefore, if such air as is near the earth be inclosed in a vessel, so as to have no communication with the external air, the pressure of such inclosed air will be exactly equal to that of the whole external atmosphere. And accordingly we find that quicksilver is sustained to the same height, by the elastic force of air inclosed in a glass vessel, as by the whole pressure of the atmosphere.—And on this principle of the condensation and elasticity of the air, depends the structure and use of the air-gun.

That the density of the air is always directly proportional to the force or weight which compresses it, was proved by Boyle and Mariotte, at least as far as their experiments go on this head: and Mr. Mariotte has shewn that the same rule takes place in condensed air. However, this rule is not to be admitted as scrupulously exact; for when air is very forcibly compressed, so as to be reduced to  $\frac{1}{4}$ th of its ordinary bulk, the effect does not answer precisely to the rule; for in this case the air begins to make a greater resistance, and requires a stronger compression, than according to the rule. And hence it would seem, that the particles of air cannot, by means of any possible weight or pressure,



how great soever, be brought into perfect contact, or that it cannot thus be reduced to a solid mass; and consequently that there must be a limit to which this condensation of the air can never arrive. And the same remark is true with regard to the rarefaction of air, namely, that in very high degrees of rarefaction, the elasticity is decreased rather more than in proportion to the weight or density of the air: and hence there must also be a limit to the rarefaction and expansion of the air, by which it is prevented from expanding to infinity.

We know not however how to assign those limits to the elasticity of the air, nor to destroy or alter it, without changing the very nature of air, which is effected by chemical processes. To what degree air is susceptible of condensation, by compression, is not certainly known. Mr. Boyle condensed it 13 times more than in its natural state, by this means: others have compressed it into  $\frac{1}{80}$ th part of its ordinary volume; Dr. Hales made it 38 times more dense, by means of a press; but by freezing water in a hollow cast-iron ball or shell, he reduced it to 1838 times less space than it naturally occupies; in which state it must have been of more than twice the density or specific gravity of water: And as water is not compressible, except in a very small degree, it follows from this experiment, that the particles of air must be of a nature very different from those of water; since it would otherwise be impossible to reduce air to a volume above 800 times less than in its common state; an inference however which militates directly against an assertion made by Dr. Halley, from some experiments performed in London, and others at Florence by the Academy del Cimento, namely, that it may be safely concluded that no force whatever is capable to reduce air into a space 800 times less than that which it naturally occupies near the surface of the earth.

The elasticity of the air exerts its force equally in all directions; and when it is at liberty, and freed from the cause which compressed it, it expands equally in all directions, and in consequence always assumes a spherical figure in the interstices of the fluids in which it is lodged. This is evident in liquors placed in the receiver of an air pump, by exhausting the air; at first there appears a multitude of exceeding small bubbles, like grains of fine sand, dispersed through the fluid mass, and rising upwards; and as more air is pumped out, they enlarge in size; but still they continue round. Also if a plate of metal be immersed in the liquor, on pumping, its surface will be seen covered over with small round bubbles, composed of the air which adhered to it, now expanding itself. And for the same reason it is that large glass globes are always blown up of a spherical shape, by blowing air through an iron tube into a piece of melted glass at the end of the pipe.

The expansion of the air, by virtue of its elastic property, when only the compressing force is taken off, or diminished, is found to be surprisingly great; and yet we are far from knowing the utmost dilatation of which it is capable. In several experiments made by Mr. Boyle, it expanded first into 9 times its former space; then into 31 times; then into 60, and then into 150 times. Afterwards, it was brought to dilate into

8000 times its first space; then into 10000, and at last even into 13679 times its space; and this solely by its own natural expansive force, by only removing the pressure, but without the help of fire. And on this principle depends the construction and use of the MANOMETER.

The elasticity of the air, under one and the same pressure, is still farther increased by heat, and diminished by cold, and that, by some late accurate experiments made by Sir George Shuckburgh, at the rate of the 440th part of its volume nearly, for each degree of the variation of heat, from that of temperate, in Fahrenheit's thermometer.

Mr. Hawksbee observed that a portion of air inclosed in a glass tube, when the temperature was at the freezing point, formed a volume which was to that of the same quantity of air in the greatest heat of summer here in England, as 6 to 7. And it has been found by several experiments, that air is expanded  $\frac{1}{3}$  of its natural bulk by applying the heat of boiling water to it.

Dr. Hales found that the air in a retort, when the bottom of the vessel just became red hot, was dilated into twice its former space; and that in a white, or almost melting heat, it filled thrice its former space: but Mr. Robins found that air was expanded, by means of the white or fusing heat of iron, to 4 times its former bulk.

See several ingenious experiments on the elasticity of the air, in the *Philos. Transf.* for the year 1777, by Sir George Shuckburgh and Colonel Roy.

This properly explains the common effect observed on bringing a close flaccid bladder near the fire to warm it; when it is presently found to swell as if more air were blown into it. And upon this principle depends the structure and office of the thermometer; as also the air balloons, lately invented by Mr. Montgolfier, for floating in the atmosphere.

M. Amontons first discovered that, with the same degree of heat, air will expand in a degree proportioned to its density. And on this foundation the ingenious author has formed a discourse, to prove "that the spring and weight of the air, with a moderate degree of warmth, may enable it to produce even earthquakes, and others of the most vehement commotions of nature." He computes that at the depth of the 74th part of the earth's radius below the surface, the natural pressure of the air would reduce to the density of gold; and thence infers that all matter below that depth, is probably heavier than the heaviest metal that we know of. And hence again, as it is proved that the more the air is compressed, the more does the same degree of fire increase the force of its elasticity; we may infer that a degree of heat, which in our orb can produce only a moderate effect, may have a very violent one in such lower orb; and that, as there are many degrees of heat in nature, beyond that of boiling water, it is probable there may be some whose violence, thus assisted by the weight of the air, may be sufficiently powerful to tear asunder the solid globe. *Mem. de l'Acad.* 1703.

Many philosophers have supposed that the elastic property of the air depends on the figure of its corpuscles, which they take to be ramous: some maintain that



that they are so many minute *floculi*, resembling fleeces of wool: others conceive them rolled up like hoops, and curled like wires, or shavings of wood, or coiled like the springs of watches, and endeavouring to expand themselves by virtue of their texture.

But Sir Isaac Newton (*Optics*, Qu. 31, &c.) explains the matter in a different way; such a contexture of parts he thinks by no means sufficient to account for that amazing power of elasticity observed in air, which is capable of dilating itself into above a million of times more space than it occupied before: but, he observes, as it is known that all bodies have an attractive and a repelling power; and as both these are stronger in bodies, the denser, more compact, and solid they are; hence it follows that when, by heat, or any other powerful agent, the attractive force is overcome, and the particles of the body separated so far as to be out of the sphere of attraction; the repelling power, then commencing, makes them recede from each other with a strong force, proportionable to that with which they before cohered; and thus they become permanent air.

And hence, he says, it is, that as the particles of air are grosser, and rise from denser bodies, than those of transient air, or vapour, true air is more ponderous than vapour, and a moist atmosphere lighter than a dry one.

And M. Amontons makes the elasticity of air to arise from the fire it contains; so that by augmenting the degree of heat, the rarefaction will be increased to a far greater degree than by a mere spontaneous dilatation.

The elastic power of the air becomes the second great source of the remarkable effects of this important fluid. By this property it insinuates itself into the pores of bodies, where, by means of this virtue of expanding, which is so easily excited, it must put the particles of those bodies into perpetual vibrations, and maintain a continual motion of dilatation and contraction in all bodies, by the incessant changes in its gravity and density, and consequently its elasticity and expansion.

This reciprocation is observable in several instances, particularly in plants, in which the tracheæ or air-vessels perform the office of lungs; for as the heat increases or diminishes, the air alternately dilates and contracts, and so by turns compresses the vessels, and eases them again; thus promoting a circulation of their juices. And hence it is found that no vegetation or germination is carried on in vacuo.

It is from the same cause too, that ice is burst by the continual action of the air contained in its bubbles. Thus, too, glasses and other vessels are frequently cracked, when their contained liquors are frozen; and thus also large blocks of stone, and entire columns of marble, sometimes split in the winter season, from some little bubble of included air acquiring an increased elasticity: and for the same reason it is that so few stones will bear to be heated by a fire, without cracking into many pieces, by the increased expansive force of some air confined within their pores. From the same source arise also all putrefaction and fermentation; neither of which can be carried on in vacuo, even in the best disposed subjects. And even respiration, and animal life itself, are supposed, by many authors, to be

conducted, in a great measure, by the same principle of the air. And as we find such great quantities of air generated by the solution of animal and vegetable substances, a good deal must constantly be raised from the dissolution of these elements in the stomach and bowels.

In fact, all natural corruption and alteration seem to depend on air; and even metals, particularly gold, only seem to be durable and incorruptible, in so far as they are impervious to air.

As to the different kinds of air, with its generation, and the effects of different ingredients of it, &c, they are omitted here, as properly belonging to a Chemical Dictionary, or to a General Dictionary of Arts, &c.

For the resistance of the air, see RESISTANCE.

AIR-GUN, in PNEUMATICS, is a machine for propelling bullets with great violence, by the sole means of condensed air.

The first account we meet with of an air-gun, is in the *Elemens d'Artillerie* of David Rivaut, who was preceptor to Louis XIII. of France. He ascribes the invention to one Marin, a burgher of Lisieux, who presented one to Henry IV.

To construct a machine of this kind, it is only necessary to take a strong vessel of any sort, into which the air is to be thrown or condensed by means of a syringe, or otherwise, the more the better; then a valve is suddenly opened, which lets the air escape by a small tube in which a bullet is placed, and which is thus violently forced out before the air.

It is evident then that the effect is produced by virtue of the elastic property of the air; the force of which, as has been shewn in the last article, is directly proportional to its condensation; and therefore the greater quantity that can be forced into the engine, the greater will be the effect. Now this effect will be exactly similar to that of a gun charged with powder, and therefore we can easily form a comparison between them: for inflamed gun-powder is nothing more than very condensed elastic air; so that the two forces are exactly similar. Now it is shewn by Mr. Robins, in his *New Principles of Gunnery*, that the fluid of inflamed gun-powder, has, at the first moment, a force of elasticity equal to about a 1000 times that of common air; and therefore it is necessary that air should be condensed a 1000 times more than in its natural state, to produce the same effect as gun-powder. But then it is to be considered, that the velocities with which equal balls are impelled, are directly proportional to the square roots of the forces; so that if the air in an air-gun be condensed only 10 times, then the velocity it will project a ball with, will be, by that rule,  $\frac{1}{5}$ th of that arising from gun-powder; and if the air were condensed 20 times, it would communicate a velocity of  $\frac{1}{7}$  of that of gun-powder. But in reality the air-gun shoots its ball with a much greater proportion of velocity than as above, and for this reason, namely, that as the reservoir, or magazine of condensed air, is commonly very large in proportion to the tube which contains the ball, its density is very little altered by expanding through that narrow tube, and consequently the ball is urged all the way by nearly the same uniform force as at the first instant; whereas the elastic fluid arising from inflamed gun-powder is but very small in proportion



proportion to the tube or barrel of the gun, occupying at first indeed but a very small portion of it next the but-end : and therefore by dilating into a comparatively large space, as it urges the ball along the barrel, its elastic force is proportionally weakened, and it acts always less and less on the ball in the tube. From which cause it happens, that air condensed into a good large machine only 10 times, will shoot its ball with a velocity but little inferior to that given by the gun-powder. And if the valve of communication be suddenly shut again by a spring, after opening it to let some air escape, then the same collection of it may serve to impel many balls, one after another.

In all cases in which a considerable force is required, and consequently a great condensation of air, it will be requisite to have the condensing syringe of a small bore, perhaps not more than half an inch in diameter : otherwise the force to produce the compression will become so great, that the operator cannot work the machine : for, as the pressure against every square inch is about 15 pounds, and against every circular inch about 12 pounds, if the syringe be one inch in diameter, when one atmosphere is injected, there will be a resistance of 12 pounds against the piston ; when 2, of 24 pounds ; and when 10 are injected, there will be a force of 120 pounds to overcome ; whereas 10 atmospheres act against the half-inch piston, whose area is but  $\frac{1}{4}$  of the former, with  $\frac{1}{4}$  of the force only, namely, 30 pounds ; and 40 atmospheres may be injected with such a syringe, as well as ten with the larger.

There are air-guns of various constructions ; an easy and portable one is represented in Plate II, fig. 1. which is a section lengthways through the axis, to shew the inside. It is made of brass, and has two barrels ; the inner barrel D A of a small bore, from which the bullets are shot ; and a larger barrel E S C D R, on the outside of it. In the stock of the gun there is a syringe M N P S, whose rod M draws out to take in air ; and by pushing it in again, the piston S N drives the air before it, through the valve P E into the cavity between the two barrels. The ball K is put down into its place in the small barrel, with the rammer, as in another gun. There is another valve at S L, which, being opened by the trigger O, permits the air to come behind the ball, so as to drive it out with great force. If this valve be opened and shut suddenly, one charge of condensed air may make several discharges of bullets ; because only part of the injected air will then go out at a time, and another bullet may be put into the place K : but if the whole air be discharged on a single bullet, it will impel it more forcibly. This discharge is effected by means of a lock (fig. 2) when fixed to its place as usual in other guns ; for the trigger being pulled, the cock will go down and drive a lever which opens the valve.

Dr. Macbride (Exper. Eff. p. 81) mentions an improvement of the air-gun, made by Dr. Ellis ; in which the chamber for containing the condensed air is not in the stock, which renders the machine heavy and unweildy, but has five or six hollow spheres belonging to it, of about 3 inches diameter, fitted to a screw on the lock of the gun. These spheres are contrived with valves, to confine the air which is forced into their cavities, so that a servant may carry them ready charged with condensed air : and thus the gun of this

construction is rendered as light and portable as one of the smallest fowling-pieces.

Fig. 3 represents one made by the late Mr. B. Martin of London, and now by several of the mathematical instrument and gun-makers of the metropolis ; which, for simplicity and perfection, perhaps exceeds any other that has been contrived. A is the gun-barrel, of the size and weight of a common fowling-piece, with the lock, stock, and ramrod. Under the lock, at b, is a round steel tube, having a small moveable pin in the inside, which is pushed out when the trigger a is pulled, by the springwork within the lock ; to this tube b is screwed a hollow copper ball, perfectly airtight. This copper ball is fully charged with condensed air by means of a syringe, previous to its being applied to the tube b. Hence, if a bullet be rammed down in the barrel, the copper ball screwed fast at b, and the trigger a be pulled ; then the pin in b will forcibly push open a valve within the copper ball, and let out a portion of the condensed air ; which air will rush up through the aperture of the lock, and forcibly act against the bullet, driving it to the distance of 60 or 70 yards, or farther. If the air be strongly condensed at every discharge, only a portion of the air escapes from the ball ; therefore, by re-cocking the piece, another discharge may be made ; and this repeated 15 or 16 times. An additional barrel is sometimes made, and applied for the discharge of shot, instead of the ball above described.

Sometimes the syringe is applied to the end of the barrel C (fig. 4) ; the lock and trigger shut up in a brass case d ; and the trigger pulled, or the discharge made, by pulling the chain b. In this contrivance there is a round chamber for the condensed air at the end of the spring at e, and it has a valve acting in a similar manner to that of the copper ball. When this instrument is not in use, the brass case d is made to slide off, and the instrument then becomes a walking stick : from which circumstance, and the barrel being made of cane, or brass, &c, it has been called the *Air-cane*. The head of the cane unscrews and takes off at a, where the extremity of the piston-rod in the barrel is shewn. An iron rod is placed in a ring at the end of this, and the air is condensed in the barrel in a manner similar to that of the gun as above ; but its force and action is not near so strong as in the gun.

**MAGAZINE AIR-Gun.** This is an improvement of the common air-gun, made by an ingenious artist, called L. Colbe. By his contrivance, ten bullets are so lodged in a cavity, near the place of discharge, that they may be successively drawn into the barrel, and shot so quickly as to be nearly of the same use as so many different guns ; the only motion required, after the air has been injected, being that of shutting and opening the hammer, and cocking and pulling the trigger. Fig. 3 is a longitudinal section of this gun, as large in every part as the gun itself ; and as much of its length is shewn as is peculiar to this construction ; the rest of it being like the ordinary air-gun. E E is part of the stock ; G is the end of the injecting syringe, with its valve H, opening into the cavity F F F F between the barrels. K K is the small or shooting barrel, which receives the bullets, one at a time, from the magazine D E, being a serpentine cavity, in which the bullets b, b, b, &c, are lodged,



lodged, and closed at the end D; from whence, by one motion of the hammer, they are brought into the barrel at I, and thence are shot out by the opening of the valve V, which lets in the condensed air from the cavity F F F into the channel V K I, and so along the inner barrel K K K, whence the bullet is discharged. *s I s i* M k is the key of a cock, having a hole through it; which hole, in the present situation, makes part of the barrel K K, being just of the same bore: so that the air, which is let in at every opening of the valve V, comes behind this cock, and taking the ball out of it, carries it forward, and so out of the mouth of the piece.

To bring in another bullet to succeed I, which is done in an instant, bring the cylindrical cavity of the key of the cock, which made part of the barrel K K K, into the situation *i k*, so that the part l may be at K; then turning the gun upside-down, one bullet next the cock will fall into it out of the magazine, but will go no farther into this cylindrical cavity, than the two little pieces *s s* will permit it; by which means only one bullet at a time will be taken in to the place I, to be discharged again as before.

A more particular description of the several parts may be seen in Desaguliers' Exper. Philos. vol. ii. pa. 399 et seq.

AIR-PUMP, in *Pneumatics*, is a machine for exhausting the air out of a proper vessel, and so to make what is commonly called a vacuum; though in reality the air in the receiver is only rarefied to a great degree, so as to take off the ordinary effects of the atmosphere. So that by this machine we learn, in some measure, what our earth would be without air; and how much all vital, generative, nutritive, and alterative powers depend upon it.

The principle on which the air-pump is constructed, is the spring or elasticity of the air; as that on which the common, or water-pump is formed, is the gravity of the same air: the one gradually exhausting the air from a vessel by means of a piston, with a proper valve, working in a cylindrical barrel or tube; and the other exhausting water in a similar manner.

The air-pump has proved one of the principal means of performing philosophical discoveries, that has been invented by the moderns. The idea of such a machine occurred to several persons, nearly about the same time. But the first it seems was completed by Otto Guericke, the celebrated consul of Magdeburg, who exhibited his first public experiments with it, before the emperor and the states of Germany, at the breaking up of the imperial diet at Ratisbon, in the year 1654. But it was not till the year 1672 that Guericke published a description of the instrument, with an account of his experiments, in his *Experimenta Nova Magdeburgica de Vacuo Spacio*: though an account of them had been published by Schottus in 1657, in his *Mechanica Hydraulico Pneumatica*.

Dr. Hook and M. Duhamel ascribe the invention of the air-pump to Mr. Boyle. But that great man frankly confesses that Guericke was beforehand with him in the execution. Some attempts, he assures us, he had indeed made upon the same foundation, before he knew any thing of what had been done abroad: but the information he afterwards received from the account given

by Schottus, enabled him, with the assistance of Dr. Hook, after two or three unsuccessful trials, to bring his design to maturity. The product of their labours was a new air-pump, much more easy, convenient, and manageable, than the German one. And hence, or rather from the great variety of experiments to which this illustrious author applied the machine, it was afterwards called *Machina Boyleana*, and the vacuum produced by it, *Vacuum Boyleanum*.

*Structure of the Air-Pump.* Most of the air-pumps that were first made, consisted of only one barrel, or hollow cylinder of brass, with a valve at the bottom, opening inwards; and a moveable embolus or piston, having likewise a valve opening upwards, and so exactly fitted to the barrel, that when it is drawn up from the bottom, by means of an indented iron rod or rack, and a handle turning a small indented wheel, playing in the teeth of that rod, all the air will be drawn up from the cavity of the barrel: there is also a small pipe opening into the bottom of the barrel, by means of which it communicates with any proper vessel to be exhausted of air, which is called a receiver, from its office in receiving the subjects upon which experiments are to be made in vacuo: the whole being fixed in a convenient frame of wood-work, where the end of the pipe turns up into a horizontal plate, upon which the receiver is placed, just over that end of the pipe.

The other parts of the machine, being only accidental circumstances, chiefly respecting conveniency, have been diversified and improved from time to time, according to the address and several views of the makers. That of Otto Guericke was very rude and inconvenient, requiring the labour of two strong men, for more than two hours, to extract the air from a glass, which was also placed under water; and yet allowed of no change of subjects for experiments.

Mr. Boyle, from time to time, removed several of these inconveniences, and lessened others: but still the working of his pump, which had but one barrel, was laborious, by reason of the pressure of the atmosphere, a great part of which was to be removed at every lift of the piston, when the exhaustion was nearly completed. Various improvements were successively made in the machine by the philosophers about that time, and soon after, who cultivated this new and important branch of pneumatics; as Papin, Merfenne, Mariotte, and others; but still they laboured under a difficulty of working them, from the circumstance of the single barrel, till Papin, in his farther improvements of the air-pump, removed that inconvenience, by the use of a second barrel and piston, contrived to rise, as the other fell, and to fall as that rose; by which, and the great improvements made by Mr. Hawksbee, the pressure of the atmosphere on the descending piston, always nearly balanced that of the ascending one; so that the winch, which worked them up and down, was easily moved by a very gentle force with one hand: and besides, the exhaustion was hereby made in less than half the time.

Some of the Germans, and others likewise, made improvements in the air-pump, and contrived it to perform the counter office of a condenser, in order to examine the properties of the air depending on its condensation.

Mr.



Mr. Boyle contrived a mercurial gauge or index to the air-pump, which is described in his first and second *Physico-Mechanical Continuations*, for measuring the degrees of the air's rarefaction in the receiver. This gauge is similar to the barometer, being a long glass tube, having its lower end immersed in an open basin of quicksilver, but its other end, which was open also, communicating with the receiver: which being exhausted, this tube is equally exhausted of air at the same time, and the external air presses the quicksilver up into the tube, to a height proportioned to the degree of exhaustion.

Mr. Vream, an ingenious pneumatic operator, made an improvement in Hawksbee's air-pump, by reducing the alternate up-and-down motion of the hand and winch to a circular one. In his method, the winch is turned quite round, and yet the pistons are alternately raised and depressed: by which the trouble of shifting the hand backwards and forwards, as well as the loss of time, and the shaking of the pump, are prevented.

The air-pump, thus improved, is represented in plate III. fig. 1; where *o o* is the receiver to be exhausted, ground truly level at the bottom, set over a hole in the plate, from which descends the bent pipe *b b* to the cistern *d d*, with which the two barrels *a a* communicate, in which the pistons are worked by a toothed wheel, by turning the handle *b b*; by which the racks *c c*, with the pistons, are worked alternately up and down. *l l* is the gauge tube, immersed in a basin of quicksilver *m* at bottom, and communicating with the receiver at top; from which however it may be occasionally disengaged, by turning a cock. And *n* is another cock, by turning of which, the air is again let in to the exhausted receiver; into which it is heard to rush with a considerable hissing noise.

Notwithstanding the great excellency of Mr. Hawksbee's air-pump, it was still subject to inconveniences, from which it was in a great measure relieved by some contrivances of Mr. Smeaton, which are described at large in the *Philos. Trans.* for the year 1752. The principal improvements suggested by Mr. Smeaton, relate to the gauge, the valves of the piston, and the piston going closer down to the bottom of the barrel; for his pump has only one. By the last of these, the air was extracted more perfectly at each stroke. By the second, he remedied an inconvenience arising from the valve hole of the piston being too wide properly to support the bladder valve which covered it: instead of the usual circular orifice, Mr. Smeaton perforated the piston with seven small and equal hexagonal holes, one in the centre, and the other six around, forming together the appearance of a transverse section of a honeycomb; the bars or divisions between which, served to support the pressure of the air on the valve. His gauge consists of a bulb of glass, of a pear-like shape, and capable of holding about half a pound of quicksilver: it is open at the lower end, the other terminating in a tube hermetically sealed; and it has annexed to it a scale, divided into parts of about  $\frac{1}{16}$  of an inch, and answering to the 1000th part of the whole capacity. During the exhaustion of the receiver, the gauge is suspended in it by a wire; but when the pump has been worked as much as necessary, the gauge is pushed down, till the open end be immersed in a basin of quicksilver

placed underneath. The air is then let into the receiver again, and the quicksilver driven by it from the basin, up into the gauge, till the air remaining in it become of the same density as the air without; and as the air always takes the highest place, the tube being uppermost, the expansion will be determined by the number of divisions occupied by the air at the top. This air-pump is made to act also as a condensing engine, as some German machines had done before, by the very simple apparatus of turning a cock.

By means of this gauge, Mr. Smeaton judged that his machine was incomparably better than any former ones, as it seemed to rarefy the air in the receiver 1000, or even 2000 times, while the best of the former construction only rarefied about 140 times: and so the case has since been always understood, an implicit confidence being placed in Mr. Smeaton's accuracy, till the fallacy was accidentally detected in the manner related at large by Mr. Nairne in the *Philos. Trans.* for the year 1777. This accurate and ingenious artist wanting to make trial of Mr. Smeaton's pear-gauge, executed an air-pump of his improved construction, in the best manner possible; which, in various experiments made with it, appeared, by the pear-gauge, to rarefy the air to an amazing degree indeed, being at times from 4000 to 10000, or 50000, or even 100000 times rarefied. But upon measuring the same expansion by the usual long and short tube gauges, which both accurately agreed together, he found that these never shewed a rarefaction of more than 600 times: widely different from the same as measured by the pear or internal gauge, by experiments often repeated. 'Finding, says Mr. Nairne, still this disagreement between the pear-gauge and the other gauges, I tried a variety of experiments; but none of them appeared to me satisfactory, till one day in April 1776, shewing an experiment with one of these pumps to the honourable Henry Cavendish, Mr. Smeaton, and several other gentlemen of the Royal Society, when the two gauges differed some thousand times from one another, Mr. Cavendish accounted for it in the following manner. "It appeared, he said, from some experiments of his father's, Lord Cavendish, that water, whenever the pressure of the atmosphere on it is diminished to a certain degree, is immediately turned into vapour, and is as immediately turned back again into water on restoring the pressure. This degree of pressure is different according to the heat of the water: when the heat is 72° of Fahrenheit's scale, it turns into vapour as soon as the pressure is no greater than that of three quarters of an inch of quicksilver, or about 1.40th of the usual pressure of the atmosphere; but when the heat is only 41°, the pressure must be reduced to that of a quarter of an inch of quicksilver before the water turns into vapour. It is true, that water exposed to the open air, will evaporate at any heat, and with any pressure of the atmosphere; but that evaporation is intirely owing to the action of the air upon it; whereas the evaporation here spoken of, is performed without any assistance from the air. Hence it follows, that, when the receiver is exhausted to the above-mentioned degree, the moisture adhering to the different parts of the machine will turn into vapour, and supply the place of the air, which is continually drawn away by the working of the



the pump; so that the fluid in the pear-gauge, as well as that in the receiver, will consist in a good measure of vapour. Now letting the air into the receiver, all the vapour within the pear-gauge will be reduced to water, and only the real air will remain uncondensed; consequently the pear-gauge shews only how much real air is left in the receiver, and not how much the pressure or spring of the included fluid is diminished; whereas the common gauges shew how much the pressure of the included fluid is diminished, and that equally, whether it consist of air or of vapour." Mr. Cavendish having explained so satisfactorily the cause of the disagreement between the two gauges, Mr. Nairne considered that, if he were to avoid moisture as much as possible, the two gauges should nearly agree. And in fact they were found so to do, each shewing a rarefaction of about 600, when all moisture was perfectly cleared away from the pump, and the plate and the edges of the receiver were secured by a cement instead of setting it upon a soaked leather, as in the usual way. But by future experiments, Mr. Nairne found that the same excellent machine would not exhaust more than 50 or 60 times, when the receiver was set upon leather soaked in water, the heat of the room being about  $57^{\circ}$ . And from the whole, Mr. Nairne concludes that the air-pump of Otto Guericke, and those contrived by Mr. Gratorix, and Dr. Hook, and the improved one by Mr. Papin, both used by Mr. Boyle, as also Hawksbee's, s'Gravesande's, Muschenbroeck's, and those of all who have used water in the barrels of their pumps, could never have exhausted to more than between 40 and 50, if the heat of the place was about  $57^{\circ}$ ; and although Mr. Smeaton, with his pump, where no water was in the barrel, but where leather soaked in a mixture of water and spirit of wine was used on the pump-plate, to set the receiver upon, may have exhausted all but a thousandth, or even a tenthousandth part of the common air, according to the testimony of his pear-gauge; yet so much vapour must have arisen from the wet leather, that the contents of the receiver could never be less than a 70th or 80th part of the density of the atmosphere. But when nothing of moisture is used about this machine, it will, when in its greatest perfection, rarefy its contents of air about 600 times.

It is evident that by means of these two gauges we can ascertain the several quantities of vapour and permanent air which make up the contents of the receiver, after the exhaustion is made as perfect as can be; for the usual external gauge determines the whole contents, made up of the vapour and air, whilst the pear-gauge shews the quantity of real permanent air; consequently the difference is the quantity of vapour.

The principal cause which prevents this pump from exhausting beyond the limit above-mentioned, is the weakened elasticity of the air within the receiver, which, decreasing in proportion as the quantity of the air within is diminished, becomes at last incapable of lifting up the valve of communication between the receiver and the barrel; and consequently no more air can then pass from the former to the latter.

Several ingenious persons have used their endeavours to remove this imperfection in the best air-pumps. Amongst these it seems that one Mr. Haas has succeeded tolerably well; having, by means of a contri-

vance to open the communication valve in the bottom of the barrel, made his machine so perfect, that when every thing is in the greatest perfection, it rarefies the contents of the receiver as far as 1000 times, even when measured by the exterior gauge. The description of this machine, and an account of some experiments performed with it, are given by Mr. Cavallo in the *Philos. Trans.* for the year 1783.

But the imperfections it seems have more recently been removed by an ingenious contrivance of Mr. Cuthbertson, a mathematical instrument maker at Amsterdam, now of London, whose air-pump has neither cocks nor valves, and is so constructed, that what supplies their place has the advantages of both, without the inconveniences of either. He has also made improvements in the gauges, by means of which he determines the height of the mercury in the tube, by which the degree of exhaustion is indicated, to the hundredth part of an inch. And to obviate the inconvenience of the elastic vapour arising from the wet leather, upon which the receiver is placed, for common experiments, he recommends the use of leather dressed with allum, and soaked in hog's lard, which he found to yield very little of this vapour; but when the utmost degree of exhaustion is required, his advice is, to dry the receiver well, and set it upon the plate without any leather, only smearing its outer edges with hog's lard, or with a mixture of three parts of hog's lard and one of oil. But the use of the leather has long been laid aside by our English instrument-makers, a circumstance which probably had not come to Mr. Cuthbertson's knowledge. An account of this instrument, and of some experiments performed with it, was published at Amsterdam in the year 1787; from which experiments it appears that, by a coincidence of the several gauges, a rarefaction of 1200 times was shewn; but when the atmosphere was very dry, the exhaustion has been so complete, that the gauges have shewn the air in the receiver to be rarefied above 2400 times.

There are made also by different persons, portable, or small air-pumps, of various constructions, to set upon a table, to perform experiments with. In these, the gauge is varied according to the fancy of the maker, but commonly it consists of a bent glass tube, like a syphon, open only at one end. The gauge is placed under a small receiver communicating, by a pipe, with the principal pipe leading from the general receiver to the barrels. The close end of the gauge, of 3 or 4 inches long, before the exhaustion, has the quicksilver forced close up to the top by the pressure of the air on the open end; but when the exhaustion is considerably advanced, it begins to descend, and then the difference of the heights of the quicksilver in the two legs, compared with the height in the barometrical tube, determines the degree of exhaustion: so if the difference between the two be one inch, when the barometer stands at 30, the air is rarefied 30 times; but if the difference be only half an inch, the rarefaction is 60 times, and so on. See Plate III. fig. 2.

*The Use of the Air-Pump.* In whatever manner or form this machine be made, the use and operation of it are always the same. The handle, which works the piston, is moved up and down in the barrel, by which



means a barrel of the contained air is drawn out at every stroke of the piston, in the following manner: by pushing the piston down to the bottom of the barrel, where the air is prevented from escaping downwards, by its elasticity it opens the valve of the piston, and escapes upwards above it into the open air; then raising the piston up, the external atmosphere shuts down its valve, and a vacuum would be made below it, but for the air in the receiver, pipe, &c, which now raises the valve in the bottom of the barrel, and rushes in and fills it again, till the whole air in the receiver and barrel be of one uniform density, but less than it was before the stroke, in proportion as the sum of all the capacities of the receiver, pipe, and barrel together, is to the same sum wanting the barrel. And thus is the air in the receiver diminished at each stroke of the piston, by the quantity of the barrel or cylinder full, and therefore always in the same proportion: so that by thus repeating the operation again and again, the air is rarefied to any proposed degree, or till it has not elasticity enough to open the valve of the piston or of the barrel, after which the exhaustion cannot be any farther carried on: the gauge, in comparison with the barometer, shewing at any time what the degree of exhaustion is, according to the particular nature and construction of it.

But, supposing no vapour from moisture, &c, to rise in the receiver, the degree of exhaustion, after any number of strokes of the piston, may be determined by knowing the respective capacities of the barrel and the receiver, including the pipe, &c. For as we have seen above that every stroke diminishes the density in a constant proportion, namely as much as the whole content exceeds that of the cylinder or barrel; and consequently the sum of as many diminutions as there are strokes of the piston, will shew the whole diminution by all the strokes. So, if the capacity of the barrel be equal to that of the receiver, in which the communication pipe is always to be included; then, the barrel being half the sum of the whole contents, half the air will be drawn out at one stroke; and consequently the remaining half, being dilated through the whole or first capacity, will be of only half the density of the first: in like manner, after the second stroke, the density of the remaining contents will be only half of that after the first stroke, that is only  $\frac{1}{4}$  of the original density: continuing this operation, it follows that the density of the remaining air will be  $\frac{1}{8}$  after 3 strokes of the piston,  $\frac{1}{16}$  after 4 strokes,  $\frac{1}{32}$  after 5 strokes, and so on, according to the powers of the ratio  $\frac{1}{2}$ ; that is, such power of the ratio as is denoted by the number of the strokes. In like manner, if the barrel be  $\frac{1}{3}$  of the whole contents, that is, the receiver double of the barrel, or  $\frac{2}{3}$  of the whole contents; then the ratio of diminution of density being  $\frac{2}{3}$ , the density of the contents, after any number of strokes of the piston, will be denoted by such power of  $\frac{2}{3}$  whose exponent is that number; namely, the density will be  $\frac{2}{3}$  after one stroke,  $(\frac{2}{3})^2$  or  $\frac{4}{9}$  after two strokes,  $(\frac{2}{3})^3$  or  $\frac{8}{27}$  after 3 strokes, and in general it will be  $(\frac{2}{3})^n$  after  $n$  strokes: the original density of the air being 1. Hence then, universally, if  $s$  denote the sum of the contents of the receiver and barrel, and  $r$  that of the receiver only without the barrel, and  $n$  any number of strokes of the piston; then, the original density of the air being 1,

the density after  $n$  strokes will be  $(\frac{r}{s})^n$  or  $\frac{r^n}{s^n}$ , namely

the  $n$  power of the ratio  $\frac{r}{s}$ . So, for example, if the capacity of the receiver be equal to 4 times that of the barrel; then their sum  $s$  is 5, and  $r$  is 4; and the density of the contents after 30 strokes, will be  $(\frac{4}{5})^{30}$ , or the 30th power of  $\frac{4}{5}$ , which is  $\frac{1}{808}$  nearly; so that the air in the receiver is rarefied 808 times.

See also the Memoires de l'Acad. Royale des Sciences for the years 1693 and 1705.

From the same formula, namely  $(\frac{r}{s})^n = d$  the density, we easily derive a rule for finding the number of strokes of the piston, necessary to rarefy the air any number of times, or to reduce it to a given density  $d$ , that of the natural air being 1. For since  $(\frac{r}{s})^n = d$ , by taking the logarithm of this equation, it is  $n \times \log. \frac{r}{s} = \log. d$ ; and hence  $n = \frac{\log. d}{\log. \frac{r}{s}} = \frac{\log. d}{\log. r - \log. s}$ .

that is, divide the log. of the proposed density by the log. of the ratio of the receiver to the sum of the receiver and barrel together, and the quotient will shew the number of strokes of the piston requisite to produce the degree of exhaustion required. So, for example, if the receiver be equal to 5 times the barrel, and it be proposed to find how many strokes of the piston will rarefy the air 100 times; then  $r = 5$ ,  $s = 6$ ,  $d = \frac{1}{100}$ , whose log. is  $-2$ , and  $\frac{r}{s} = \frac{5}{6}$ , whose log. is  $-.07918$ ;

therefore  $\frac{2}{.07918} = 25\frac{1}{4}$  nearly, which is the number of strokes required.

And, farther, the same formula reduced, would give us the proportion between the receiver and barrel, when the air is rarefied to any degree by an assigned number of strokes of the piston. For since  $(\frac{r}{s})^n = d$  the density, therefore, extracting the  $n$  root of both sides, it is  $\frac{r}{s} = \sqrt[n]{d}$ : that is, the  $n$  root of the density is equal to the ratio of the receiver to the sum of the receiver and barrel. So, if the density  $d$  be  $\frac{1}{128}$ , and the number of strokes  $n = 7$ ; then the 7th root of  $\frac{1}{128}$  is  $\frac{1}{2}$ ; which shews that the receiver is equal to half the receiver and barrel together, or that the capacity of the barrel is just equal to that of the receiver.

Some of the principal effects and phenomena of the air-pump, are the following: That, in the exhausted receiver, heavy and light bodies fall equally swift; so, a guinea and feather fall from the top of a tall receiver to the bottom exactly together. That most animals die in a minute or two; but however, That vipers and frogs, though they swell much, live an hour or two; and after being seemingly quite dead, come to life again in the open air: That snails survive about ten hours; efts, or slow-worms, two or three days; and leeches five or six. That oysters live for 24 hours. That the heart of an eel taken out of the body, continues to beat



beat for good part of an hour, and that more briskly than in the air. That warm blood, milk, gall, &c, undergo a considerable intumescence and ebullition. That a mouse or other animal may be brought, by degrees, to survive longer in a rarefied air, than naturally it does. That air may retain its usual pressure, after it is become unfit for respiration. That the eggs of silk-worms hatch in vacuo. That vegetation stops. That fire extinguishes; the flame of a candle usually going out in one minute; and a charcoal in about five minutes. That red-hot iron, however, seems not to be affected; and yet sulphur or gun-powder are not lighted by it, but only fused. That a match, after lying seemingly extinct a long time, revives again on re-admitting the air. That a flint and steel strike sparks of fire as copiously, and in all directions, as in air. That magnets, and magnetic needles, act the same as in air. That the smoke of an extinguished luminary gradually settles to the bottom in a darkish body, leaving the upper part of the receiver clear and transparent; and that on inclining the vessel sometimes to one side, and sometimes to another, the fume preserves its surface horizontal, after the nature of other fluids. That heat may be produced by attrition. That camphire will not take fire; and that gun-powder, though some of the grains of a heap of it be kindled by a burning glass, will not give fire to the contiguous grains. That glow-worms lose their light in proportion as the air is exhausted, and at length become totally obscure; but on re-admitting the air, they presently recover it all. That a bell, on being struck, is not heard to ring, or very faintly. That water freezes. But that a syphon will not run. That electricity appears like the aurora borealis. With multitudes of other curious and important particulars, to be met with in the numerous writings on this machine, namely, besides the *Philos. Transactions* of most academies and societies, in the writings of Torricelli, Pascal, Merfenne, Guericke, Schottus, Boyle, Hook, Duhamel, Mariotte, Hawksbee, Hales, Muschenbroeck, Gravesande, Desaguliers, Franklin, Cotes, Helsham, and a great many other authors.

**AIR-VESSEL**, in *Hydraulics*, is a vessel of air within some water engines, which being compressed, by forcing in a considerable quantity of water, by its uniform spring, forces it out at the pipe in a constant uninterrupted stream, to a great height.

**Air-vessel** too, in the improved fire engines, is a metallic cylinder, placed between the two forcing pumps, by the action of whose pistons the water is forced into this vessel, through two pipes, with valves; then the air, previously contained in it, is compressed by the water, in proportion to the quantity admitted, and this air, by its spring, forces the water through a pipe by a constant and equal stream; whereas in the common squirting engine, the stream is discontinued between the several strokes.

**AIRY TRIPPLICITY**, in *Astrology*, the signs of Gemini, Libra, and Aquarius.

**AJUTAGE**, or **ANJUTAGE**, in *Hydraulics*, part of the apparatus of a *jet d'eau*, or artificial fountain; being a kind of tube fitted to the aperture or mouth of the cistern, or the pipe; through which the water is to be played in any direction, and in any shape or figure.

It is chiefly the diversity in the ajutage, that makes the different kinds of fountains. So that, by having several ajutages, to be applied occasionally, one fountain is made to have the effect of many.

Mariotte, Gravesande, and Desaguliers have written pretty fully on the nature of ajutages, or spouts for jets d'eau, and especially the former. He affirms, from experiment, that an even polished round hole, made in the thin end of a pipe, gives a higher jet than either a cylindrical or a conical ajutage; but that, of these two latter however, the conical is better than the cylindrical figure. See his *Traite du Mouvement des Eaux*, part 4.

The quantity of water discharged by ajutages of equal area, but of different figures, is the same. And for like figures, but of different sizes, the quantity discharged, is directly proportional to the area of the ajutage, or to the square of its diameter, or of any side or other linear dimension: so, an ajutage of a double diameter, or side, will discharge 4 times the quantity of water; of a triple diameter, 9 times the quantity; and so on; supposing them at an equal depth below the surface or head of water. But if the ajutage be at different depths below the head, then the celerity with which the water issues, and consequently the quantity of it run out in any given time, is directly proportional to the square-root of the altitude of the head, or depth of the hole: so at 4 times the depth, the celerity and quantity is double; at 9 times the depth, triple; and so on.

It has been found that jets do not rise quite so high as the head of water; owing chiefly to the resistance of the air against it, and the pressure of the upper parts of the jet upon the lower: and for this reason it is, that if the direction of the ajutage be turned a very little from the perpendicular, it is found to spout rather higher than when the jet is exactly upright.

It is found by experiment too, that the jet is higher or lower, according to the size of the ajutage: that a circular hole of about an inch and a quarter in diameter, jets highest; and that the farther from that size, the worse. Experience also shews that the pipe leading to the ajutage, should be much larger than it; and if the pipe be a long one, that it should be wider the farther it is from the ajutage.

For the other circumstances relating to jets and the issuing of water under various circumstances, see **EXHAUSTION**, **FLUX**, **FOUNTAIN**, **JET D'EAU**, &c, to which they more properly belong.

**ALBATEGNI**, an Arabic prince of Batan in Mesopotamia, who was a celebrated astronomer, about the year of Christ 880, as appears by his observations. He is also called *Muhammed ben Geber Albatani*, *Mahomet the son of Geber*, and *Muhamedes Araçensis*. He made astronomical observations at Antioch, and at Racah or Araçta, a town of Chaldea, which some authors call a town of Syria or of Mesopotamia. He is highly spoken of by Dr. Halley, as a *vir admirandi acuminis, ac in administrandis observationibus exercitissimus*.

Finding that the tables of Ptolomy were imperfect, he computed new ones, which were long used as the best among the Arabs: these were adapted to the meridian of Araçta or Racah. Albategni composed in



Arabic a work under the title of *The Science of the Stars*, comprising all parts of astronomy, according to his own observations and those of Ptolomy. This work, translated into Latin by Plato of Tibur, was published at Nuremberg in 1537, with some additions and demonstrations of Regiomontanus; and the same was reprinted at Bologna in 1645, with this author's notes. Dr. Halley detected many faults in these editions: *Philos. Trans.* for 1693, N<sup>o</sup> 204.

In this work, Albategni gives the motion of the sun's apogee since Ptolomy's time, as well as the motion of the stars, which he makes 1 degree in 70 years. He made the longitude of the first star of Aries to be  $18^{\circ} 2'$ ; and the obliquity of the ecliptic  $23^{\circ} 35'$ . And upon Albategni's observations were founded the Alphonsine tables of the moon's motions; as is observed by Nic. Muler, in the *Tab. Frificæ*, pa. 248.

ALBERTUS MAGNUS, a very learned man in the 13th century, who, among a multitude of books, wrote several upon the various mathematical sciences, as Arithmetic, Geometry, Perspective or Optics, Music, Astrology and Astronomy, particularly under the titles, *de sphaera, de astris, de astronomia, item speculum astronomicum*.

Albertus Magnus was born at Lawingen on the Danube, in Suabia, in 1205, or according to some in 1193; and he died at a great age, at Cologn, November 15, 1280. Vossius and other authors speak of him as a great genius, and deeply skilled in all the learning of the age. His writings were so numerous, that they make 21 volumes in folio, in the Lyons edition of 1615. He has passed also for the author of some writings relating to midwifery, &c, under the title of *De natura rerum*, and *De secretis mulierum*, in which there are many phrases and expressions unavoidable on such a subject, which gave great offence, and raised a clamour against him as the supposed author, and inconsistent with his character, being a Dominican friar, and sometime bishop of Ratisbon; which dignity however he soon resigned, through his love for solitude, to enter again into the monastic life. But the advocates of Albert assert, that he was not the author of either of these two works. It must be acknowledged however, that there are, in his *Comment upon the Master of Sentences*, some questions concerning the practice of conjugal duty, in which he has used some words rather too gross for chaste and delicate ears: but they allege what he himself used to say in his own vindication, that he came to the knowledge of so many monstrous things at confession, that it was impossible to avoid touching upon such questions. Albert was certainly a man of a most curious and inquisitive turn of mind, which gave rise to other accusations against him; such as, that he laboured to find out the philosopher's stone; that he was a magician; and that he made a machine in the shape of a man, which was an oracle to him, and explained all the difficulties he proposed: the common cant accusations of those times of ignorance and superstition. But having great knowledge in the mathematics and mechanics, by his skill in these sciences he probably formed a head, with springs capable of articulate sounds; like the machines of Boetius and others.

John Matthæus de Luna, in his treatise *De Rerum Inventoribus*, has attributed the invention of fire-arms

to Albert; but in this he is refuted by Naude, in his *Apologie des grands hommes*.

ALBUMAZAR, otherwise called ABUASSAR, and JAPHAR, was a celebrated Arabian philosopher and astrologer, of the 9th or 10th century, or according to some authors much earlier. Blancanus, Vossius, &c, speak of him as one of the most learned astronomers of his time, or astrologers, which was then the same thing. He wrote a work *De Magnis Conjunctionibus Annorum Revolutionibus, ac eorum Perfectionibus*, printed at Venice in 1515, at the expence of Melchior Seffa, a work chiefly astrological.

He wrote also *Introductio in Astronomiam*, printed in the year 1489. And it is reported that he observed a comet in his time, above the orb of Venus.

ALCOHOL, in the Arabian Astrology, is when a heavy slow-moving planet receives another lighter one within its orb, so as to come in conjunction with it.

ALDEBARAN, an Arabian name of a fixed star, of the first magnitude, just in the eye of the sign or constellation Taurus, or the bull, and hence it is popularly called the bull's eye. For the beginning of the year 1800, its

Right Ascension is	—	66° 6' 51".10
Annual variation in AR	0 0	51".31
Declination	—	16 5 52".00 N.
And Annual variat. in Decl.	0 0	8".30

ALDERAIMIN, a star of the third magnitude in the right shoulder of the constellation Cepheus.

ALDHAFERA, or Aldhaphra, in the Arabian Astronomy, denotes a fixed star of the third magnitude, in the mane of the sign or constellation Leo, the lion.

ALEMBERT (JOHN LE ROND D'), an eminent French mathematician and philosopher, and one of the brightest ornaments of the 18th century. He was perpetual secretary to the French Academy of Sciences, and a member of most of the philosophical academies and societies of Europe.

D'Alembert was born at Paris, the 16th of November 1717. He derived the name of John le Rond from that of the church near which, after his birth, he was exposed as a foundling. But his father, informed of this circumstance, listening to the voice of nature and duty, took measures for the proper education of his child, and for his future subsistence in a state of ease and independence. His mother, it is said, was a lady of of rank, the celebrated Mademoiselle Tencin, sister to cardinal Tencin, archbishop of Lyons.

He received his first education among the Jansenists, in the College of the Four Nations, where he gave early signs of genius and capacity. In the first year of his philosophical studies, he composed a Commentary on the Epistle of St. Paul to the Romans. The Jansenists considered this production as an omen, that portended to the party of Port-Royal a restoration to some part of their former splendor, and hoped to find one day in d'Alembert a second Pascal. To render this resemblance more complete, they engaged their pupil in the study of the mathematics; but they soon perceived that his growing attachment to this science was likely to disappoint the hopes they had formed with respect to his future destination: they therefore endeavoured to divert him from this line; but their endeavours were fruitless.

On



On his quitting the college, finding himself alone, and unconnected in the world, he sought an asylum in the house of his nurse. He hoped that his fortune, though not ample, would enlarge the subsistence, and better the condition of her family, which was the only one that he could consider as his own. It was here therefore that he fixed his residence, resolving to apply himself entirely to the study of geometry.—And here he lived, during the space of 40 years, with the greatest simplicity, discovering the augmentation of his means only by increasing displays of his beneficence, concealing his growing reputation and celebrity from these honest people, and making their plain and uncouth manners the subject of good-natured pleasantry and philosophical observation. His good nurse perceived his ardent activity; heard him mentioned as the writer of many books; but never took it into her head that he was a great man, and rather beheld him with a kind of compassion. “You will never, said she to him one day, be any thing but a philosopher—and what is a philosopher?—a fool, who toils and plagues himself all his life, that people may talk of him when he is dead.”

As d’Alembert’s fortune did not far exceed the demands of necessity, his friends advised him to think of some profession that might enable him to increase it. He accordingly turned his views to the law, and took his degrees in that faculty: but soon after, abandoning this line, he applied himself to the study of medicine. Geometry however was always drawing him back to his former pursuits; so that after many ineffectual struggles to resist its attractions, he renounced all views of a lucrative profession, and gave himself up entirely to mathematics and poverty.

In the year 1741 he was admitted a member of the Academy of Sciences; for which distinguished literary promotion, at so early an age (24), he had prepared the way by correcting the errors of a celebrated work (The *Analyse Démontrée* of Reyneau), which was esteemed classical in France in the line of analytics. He afterwards set himself to examine, with close attention and assiduity, what must be the motion and path of a body, which passes from one fluid into another denser fluid, in a direction oblique to the surface between the two fluids. Every one knows the phenomenon which happens in this case, and amuses children, under the denomination of *Ducks and Drakes*; but it was d’Alembert who first explained it in a satisfactory and philosophical manner.

Two years after his election to a place in the academy, he published his *Treatise on Dynamics*. The new principle developed in this treatise, consisted in establishing an equality, at each instant, between the changes that the motion of a body has undergone, and the forces or powers which have been employed to produce them: or, to express the same thing otherwise, in separating into *two parts* the action of the moving powers, and considering the *one* as producing alone the motion of the body, in the second instant, and the *other* as employed to destroy that which it had in the first.

So early as the year 1744, d’Alembert had applied this principle to the theory of the equilibrium, and the motion of fluids: and all the problems before resolved

in physics, became in some measure its corollaries. The discovery of this new principle was followed by that of a new calculus, the first essays of which were published in a *Discourse on the General Theory of the Winds*, to which the prize-medal was adjudged by the Academy of Berlin in the year 1746, which proved a new and brilliant addition to the fame of d’Alembert. This new calculus of *Partial Differences* he applied, the year following, to the problem of vibrating chords, the resolution of which, as well as the theory of the oscillations of the air and the propagation of sound, had been but imperfectly given by the mathematicians who preceded him; and these were his masters or his rivals.

In the year 1749 he furnished a method of applying his principle to the motion of any body of a given figure. He also resolved the problem of the precession of the equinoxes; determining its quantity, and explaining the phenomenon of the nutation of the terrestrial axis discovered by Dr. Bradley.

In 1752, d’Alembert published a treatise on the *Resistance of Fluids*, to which he gave the modest title of an *Essay*; though it contains a multitude of original ideas and new observations. About the same time he published, in the *Memoirs of the Academy of Berlin*, *Researches concerning the Integral Calculus*, which is greatly indebted to him for the rapid progress it has made in the present century.

While the studies of d’Alembert were confined to mere mathematics, he was little known or celebrated in his native country. His connections were limited to a small society of select friends. But his cheerful conversation, his smart and lively sallies, a happy knack at telling a story, a singular mixture of malice of speech with goodness of heart, and of delicacy of wit with simplicity of manners, rendering him a pleasing and interesting companion, his company began to be much sought after in the fashionable circles. His reputation at length made its way to the throne, and rendered him the object of royal attention and beneficence. The consequence was a pension from government, which he owed to the friendship of count d’Argenson.

But the tranquillity of d’Alembert was abated when his fame grew more extensive, and when it was known beyond the circle of his friends, that a fine and enlightened taste for literature and philosophy accompanied his mathematical genius. Our author’s enlogist ascribes to envy, detraction, &c, all the opposition and censure that d’Alembert met with on account of the famous *Encyclopédie*, or Dictionary of Arts and Sciences, in conjunction with Diderot. None surely will refuse the well-deserved tribute of applause to the eminent displays of genius, judgment, and true literary taste, with which d’Alembert has enriched that great work. Among others, the Preliminary Discourse he has prefixed to it, concerning the rise, progress, connections, and affinities of all the branches of human knowledge, is perhaps one of the most capital productions the philosophy of the age can boast of.

Some time after this, d’Alembert published his *Philosophical, Historical, and Philological Miscellanies*. These were followed by the *Memoirs of Christina queen of Sweden*; in which d’Alembert shewed that he was acquainted with the natural rights of mankind, and



and was bold enough to assert them. His *Essay on the Intercourse of Men of Letters with Persons high in Rank and Office*, wounded the former to the quick, as it exposed to the eyes of the public the ignominy of those servile chains, which they feared to shake off, or were proud to wear. A lady of the court hearing one day the author accused of having exaggerated the despotism of the great, and the submission they require, answered slyly, "If he had consulted me, I would have told him still more of the matter."

D'Alembert gave elegant specimens of his literary abilities in his translations of some select pieces of Tacitus. But these occupations did not divert him from his mathematical studies: for about the same time he enriched the *Encyclopédie* with a multitude of excellent articles in that line, and composed his *Researches on several Important Points of the System of the World*, in which he carried to a higher degree of perfection the solution of the problem concerning the perturbations of the planets, that had several years before been presented to the Academy.

In 1759 he published his *Elements of Philosophy*: a work much extolled as remarkable for its precision and perspicuity.

The resentment that was kindled (and the disputes that followed it) by the article *Geneva*, inserted in the *Encyclopédie*, are well known. D'Alembert did not leave this field of controversy with flying colours. Voltaire was an auxiliary in the contest: but as he had no reputation to lose, in point of candour and decency; and as he weakened the blows of his enemies, by throwing both them and the spectators into fits of laughter, the issue of the war gave him little uneasiness. It fell more heavily on d'Alembert; and exposed him, even at home, to much contradiction and opposition.

It was on this occasion that the late king of Prussia offered him an honourable asylum at his court, and the office of president of his academy: and the king was not offended at d'Alembert's refusal of these distinctions, but cultivated an intimate friendship with him during the rest of his life. He had refused, some time before this, a proposal made by the empress of Russia to entrust him with the education of the Grand Duke;—a proposal accompanied with all the flattering offers that could tempt a man, ambitious of titles, or desirous of making an ample fortune: but the objects of his ambition were tranquillity and study.

In the year 1765, he published his *Dissertation on the Destruction of the Jesuits*. This piece drew upon him a swarm of adversaries, who only confirmed the merit and credit of his work by their manner of attacking it.

Beside the works already mentioned, he published nine volumes of memoirs and treatises, under the title of *Opuscules*; in which he has resolved a multitude of problems relating to astronomy, mathematics, and natural philosophy; of which his panegyrist, Condorcet, gives a particular account, more especially of those which exhibit new subjects, or new methods of investigation.

He published also *Elements of Music*; and rendered, at length, the system of Rameau intelligible: but he did not think the mathematical theory of the sonorous body sufficient to account for the rules of that art.

In the year 1772 he was chosen secretary to the French Academy of Sciences. He formed, soon after this preferment, the design of writing the lives of all the deceased academicians, from 1700 to 1772; and in the space of three years he executed this design, by composing 70 eulogies.

D'Alembert died on the 29th of October 1783, being nearly 66 years of age. In his moral character there were many amiable lines of candour, modesty, disinterestedness, and beneficence; which are described, with a diffusive detail, in his eulogium, by Condorcet, in the *Hist. de l'Acad. Royale des Sciences*, 1783.

As it may be curious and useful to have in one view an entire list of d'Alembert's writings, I shall here insert a catalogue of them, from Rozier's *Nouvelle Table des Articles contenus dans les volumes de l'Académie Royale des Sciences de Paris*, &c, as follows:

*Traité de Dynamique*, in 4to, Paris, 1743. The 2d ed. in 1758.

*Traité de l'Equilibre et du Mouvement des Fluides*. Paris, 1744; and the 2d edition in 1770.

*Reflexions sur la Cause Générale des Vents*; which gained the prize at Berlin in 1746; and was printed at Paris in 1747, in 4to.

*Recherches sur la Précession des Équinoxes, & sur la Nutation de l'Axe de la Terre dans le Système Newtonien*. Paris, 1749, in 4to.

*Essais d'une Nouvelle Théorie du Mouvement des Fluides*. Paris, 1752, in 4to.

*Recherches sur différens Points importans du Système du Monde*. Paris, 1754 and 1756, 3 vol. in 4to.

*Elemens de Philosophie*, 1759.

*Opuscules Mathématiques*, ou Memoires sur différens Sujets de Géométrie, de Mécaniques, d'Optiques, d'Astronomie. Paris, 9 vol. in 4to; 1761 to 1773.

*Elémens de Musique*, théorique & pratique, suivant les Principes de M. Rameau, éclairés, développés, & simplifiés. 1 vol. in 8vo. à Lyon.

*De la Destruction des Jésuites*, 1765.

In the Memoirs of the Academy of Paris are the following pieces, by d'Alembert: viz,

*Précis de Dynamique*, 1743, Hist. 164.

*Précis de l'Equilibre & de Mouvement des Fluides*, 1744, Hist. 55.

*Méthode générale pour déterminer les Orbites & les Mouvements de toutes les Planètes, en ayant égard à leur action mutuelle*, 1745, p. 365.

*Précis des Reflexions sur la Cause Générale des Vents*, 1750, Hist. 41.

*Précis des Recherches sur la Précession des Équinoxes, et sur la Nutation de l'Axe de la Terre dans le Système Newtonien*, 1750, Hist. 134.

*Essai d'une Nouvelle Théorie sur la Résistance des Fluides*, 1752, Hist. 116.

*Précis des Essais d'une Nouvelle Théorie de la Résistance des Fluides*, 1753, Hist. 289.

*Précis des Recherches sur les différens Points importans du Système du Monde*, 1754, Hist. 125.

*Recherches sur la Précession des Équinoxes, & sur la Nutation de l'Axe de la Terre, dans l'Hypothèse de la Dissimilitude des Méridiens*, 1754, p. 413, Hist. 116.

*Reponse à un Article du Mémoire de M. l'Abbé de la*



la Caille, sur la Théorie du Soleil, 1757, p. 145, Hist. 118.

*Addition à ce Mémoire*, 1757, p. 567, Hist. 118.

*Précis des Opuscules Mathématiques*, 1761, Hist. 86.

*Précis du troisième volume des Opuscules Mathématiques*, 1764, Hist. 92.

*Nouvelles Recherches sur les Verres Optiques*, pour servir de suite à la théorie qui en a été donnée dans le volume 3<sup>e</sup> des Opuscules Mathématiques. Premier Mémoire, 1764, p. 75, Hist. 175.

*Nouvelles Recherches sur les Verres Optiques*, pour servir de suite à la théorie qui en a été donnée dans le troisième volume des Opuscules Mathématiques. Second Mémoire, 1765, p. 53.

*Observations sur les Lunettes Achromatiques*, 1765, p. 53, Hist. 119.

*Suite des Recherches sur les Verres Optiques*. Troisième Mémoire, 1767, p. 43, Hist. 153.

*Recherches sur le Calcul Intégral*, 1767, p. 573.

*Accident arrivé par l'Explosion d'une Meule d'Emouleur*, 1768, Hist. 31.

*Précis des Opuscules de Mathématiques*, 4<sup>e</sup> & 5<sup>e</sup> volumes. Leur Analyse, 1768, Hist. 83.

*Recherches sur les Mouvements de l'Axe d'une Planète quelconque dans l'hypothèse de la Dissimilitude des Méridiennes*, 1768 p. 1, Hist. 95.

*Suite des Recherches sur les Mouvements, &c.*, 1768 p. 332, Hist. 95.

*Recherches sur le Calcul Intégral*, 1769, p. 73.

*Mémoire sur les Principes de la Mech.* 1769, p. 278.

And in the Memoirs of the Academy of Berlin, are the following pieces, by our author: viz,

*Recherches sur le Calcul Integral*, première partie, 1746.

*Solution de quelques problèmes d'astronomie*, 1747.

*Recherches sur la courbe que forme une Corde Tendue, mise en Vibration*, 1747.

*Suite des recherches sur le Calcul Intégral*, 1748.

*Lettre à M. de Maupertuis*, 1749.

*Addition aux recherches sur la courbe que forme une Corde Tendue mise en Vibration*, 1750.

*Addition aux recherches sur le Calcul Intégral*, 1750.

*Lettre à M. le professeur Formey*, 1755.

*Extr. de differ. lettres à M. de la Grange*, 1763.

*Sur les Tautochrones*, 1765.

*Extr. de differ. lettres à M. de la Grange*, 1769.

Also in the Memoirs of Turin are,

*Differentes Lettres à M. de la Grange*, en 1764 & 1765, tom. 3 of these Memoirs.

*Recherches sur differens sujets de Math.* t. 4.

ALFECCA, or Alfeta, a name given to the star commonly called Lucida Coronæ.

ALFRAGAN, ALFERGANI, or FARGANI, a celebrated Arabic astronomer, who flourished about the year 800. He was so called from the place of his nativity, Fergan, in Sogdiana, now called Maracanda, or Samarcand, anciently a part of Bactria. He is also called Ahmed (or Muhammed) ben-Cothair, or Katir. He wrote the Elements of Astronomy, in 30 chapters or sections. In this work the author chiefly follows Ptolomy, using the same hypotheses, and the same terms, and frequently citing him.

There are three Latin translations of Alfragan's

work. The first was made in the 12th century, by Joannes Hispalensis; and was published at Ferrara in 1493, and at Nuremberg in 1537, with a preface by Melancthon. The second was by James Christman, from the Hebrew version of James Antoli, and appeared at Frankfort in 1590. Christman added to the first chapter of the work an ample commentary, in which he compares together the calendars of the Romans, the Egyptians, the Arabians, the Persians, the Syrians, and the Hebrews, and shews the correspondence of their years.

The third and best translation was made by Golius, professor of mathematics and Oriental languages at Leyden: this work, which came out in 1669, after the death of Golius, is accompanied with the Arabic text, and many learned notes upon the first nine chapters; for this author was not spared to carry them farther.

ALGAROTI, commonly called *Count Algaroti*, a celebrated Italian of the present century, well skilled in Architecture and the Newtonian philosophy, &c. Algaroti was born at Padua, but in what year has not been mentioned. Led by curiosity, as well as a desire of improvement, he travelled early into foreign countries; and was very young when he arrived in France in 1736. It was here that he composed his *Newtonian Philosophy for the Ladies*, as Fontenelle had done his Cartesian Astronomy, in the work intitled *The Plurality of Worlds*. He was much noticed by the king of Prussia, who conferred on him many marks of his esteem. He died at Pisa the 23d of May, 1764, and gave orders for his own mausoleum, with this inscription upon it; *Hic jacet Algarotus, sed non omnis*. He was esteemed to be well skilled in painting, sculpture, and architecture. His works, which are numerous, and upon a variety of subjects, abound with vivacity, elegance, and wit: a collection of them has lately been made, and printed at Leghorn; but that for which he is chiefly intitled to a place in this work is his *Newtonian Philosophy for the Ladies*, a sprightly, ingenious, and popular work.

ALGEBRA, a general method of resolving arithmetical problems by means of equations. Or, it is a method of performing the calculations of all sorts of quantities by means of general signs or characters. At first, numbers and things were expressed by their names at full length; but afterwards these were abridged, and the initials of the words used instead of them; and, as the art advanced farther, the letters of the alphabet came to be employed as general representations of all sorts of quantities; and other marks were gradually introduced, to express all sorts of operations and combinations; so as to entitle it to different appellations—universal arithmetic, and literal arithmetic, and the arithmetic of signs.

The etymology of the name, *Algebra*, is given in various ways. It is pretty certain, however, that the word is Arabian, and that from those people we had the name, as well as the art itself, as is testified by Lucas le Burgo, the first European author whose treatise was printed on this art, and who also refers to former authors and masters, from whose writings he had learned it. The Arabic name he gives it, is *Algebra* & *Almucabala*, which is explained to signify the art of  
restitution



*restitution and comparison, or opposition and comparison, or resolution and equation*, all which agree well enough with the nature of this art. Some however derive it from various other arabic words; as from Geber, a celebrated philosopher, chemist, and mathematician, to whom also they ascribe the invention of this science: some likewise derive it from the word Geber, which with the particle *al*, makes Algeber, which is purely Arabic, and signifies the reduction of broken numbers or fractions to integers.

But Peter Ramus, in the beginning of his Algebra, says “the name Algebra is Syriac, signifying the art and doctrine of an excellent man. For *Geber*, in Syriac, is a name applied to men, and is sometimes a term of honour, as master or doctor among us. That there was a certain learned mathematician, who sent his Algebra, written in the Syriac language, to Alexander the Great, and he named it *Almucabala*, that is, the book of dark or mysterious things, which others would rather call the doctrine of Algebra. And to this day the same book is in great estimation among the learned in the oriental nations, and by the Indians who cultivate this art it is called *Aljabra*, and *Alboret*; though the name of the author himself is not known.” But Ramus gives no authority for this singular paragraph. It has however on various occasions been distinguished by other names. Lucas Pacioli, or de Burgo, in Italy, called it *l'Arte Maggiore: ditta dal vulgo la Regola de la Cosa over Algebra e Almucabala*; calling it *l'Arte Maggiore*, or the greater art, to distinguish it from common arithmetic, which is called *l'Arte Minore*, or the lesser art. It seems too that it had been long and commonly known in his country by the name *Regola de la Cosa*, or *Rule of the Thing*; from whence came our rule of coefs, coefs numbers, and such like terms. Some of his countrymen followed his denomination of the art; but other Italian and Latin writers called it *Regula rei & census*, the rule of the thing and the product, or the root and the square, as the unknown quantity in their equations commonly ascended no higher than the square or second power. From this Italian word *census*, pronounced *chenfus*, came the barbarous word *zenzus*, used by the Germans and others, for quadratics; with the several *zenzic* or square roots. And hence  $\mathcal{Z}$ ,  $\mathcal{S}$ ,  $\mathcal{C}$ , which are derived from the letters *r*, *z*, *c*, the initials of *res*, *zenzus*, *cubus*, or root, square, cube, came to be the signs or characters of these words: like as  $\mathcal{R}$  and  $\sqrt{\phantom{x}}$ , derived from the letters *R*, *r*, became the signs of radicality.

Later authors, and other nations, used some the one of those names, and some another. It was also called *Specious Arithmetic* by Vieta, on account of the species, or letters of the alphabet, which he brought into general use; and by Newton it was called *Universal Arithmetic*, from the manner in which it performs all arithmetical operations by general symbols, or indeterminate quantities.

Some authors define algebra to be *the art of resolving mathematical problems*: but this is the idea of analysis, or the analytic art in general, rather than of algebra, which is only one particular species of it.

Indeed algebra properly consists of two parts: first, the method of calculating magnitudes or quantities, as represented by letters or other characters: and secondly

the manner of applying these calculations in the solution of problems.

In algebra, as applied to the resolution of problems, the first business is to translate the problem out of the common into the algebraic language, by expressing all the conditions and quantities, both known and unknown, by their proper characters, arranged in an equation, or several equations if necessary, and treating the unknown quantity, whether it be number, or line, or any other thing, in the same way as if it were a known one: this forms the composition. Then the resolution, or analytic part, is the disentangling the unknown quantity from the several others with which it is connected, so as to retain it alone on one side of the equation, while all the other, or known, quantities, are collected on the other side, and so giving the value of the unknown one. And as this disentangling of the quantity sought, is performed by the converse of the operations by which it is connected with the others, taking them always backwards in the contrary order, it hence becomes a species of the analytic art, and is called the modern analysis, in contradistinction to the ancient analysis, which chiefly respected geometry, and its applications.

There have arisen great controversies and sharp disputes among authors, concerning the history of the progress and improvements of Algebra; arising partly from the partiality and prejudices which are natural to all nations, and partly from the want of a closer examination of the works of the older authors on this subject. From these causes it has happened, that the improvements made by the writers of one nation, have been ascribed to those of another; and the discoveries of an earlier author, to some one of much later date. Add to this also, that the peculiar methods of many authors have been described so little in detail, that our information derived from such histories, is but very imperfect, and amounting only to some general and vague ideas of the true state of the arts. To remedy this inconvenience therefore, and to reform this article, I have taken the pains carefully to read over in succession all the older authors on this subject, which I have been able to meet with, and to write down distinctly a particular account and description of their several compositions, as to their contents, notation, improvements, and peculiarities; from the comparison of all which, I have acquired an idea more precise and accurate than it was possible to obtain from other histories, and in a great many instances very different from them. The full detail of these descriptions would employ a volume of itself, and would be far too extensive for this place: I must therefore limit this article to a very brief abridgment of my notes, remarking only the most material circumstances in each author; from which a general idea of the chain of improvements may be perceived, from the first rude beginnings, down to the more perfect state; from which it will appear that the discoveries and improvements made by any one single author, are scarcely ever either very great or numerous; but that, on the contrary, the improvements are almost always very slow and gradual, from former writers, successively made, not by great leaps, and after long intervals of time, but by gradations which, viewed in succession, become almost imperceptible.



As to the origin of the analytic art, of which Algebra is a species, it is doubtless as old as any science in the world, being the natural method by which the mind investigates truths, causes, and theories, from their observed effects and properties. Accordingly, traces of it are observable in the works of the earliest philosophers and mathematicians, the subject of whose enquiries most of any require the aid of such an art. And this process constituted their Analytics. Of that part of analytics however which is properly called Algebra, the oldest treatise which has come down to us, is that of Diophantus of Alexandria, who flourished about the year 350 after Christ, and who wrote, in the Greek language, 13 books of Algebra or Arithmetic, as mentioned by himself at the end of his address to Dionysius, though only 6 of them have hitherto been printed; and an imperfect book on multangular numbers, namely in a Latin translation only, by Xilander, in the year 1575, and afterwards in 1621 and 1670 in Greek and Latin by Gaspar Bachet. These books however do not contain a treatise on the elementary parts of Algebra, but only collections of difficult questions relating to square and cube numbers, and other curious properties of numbers, with their solutions. And Diophantus only prefaces the books by an address to one Dionysius, for whose use it was probably written, in which he just mentions certain *precognita*, as it were to prepare him for the problems themselves. In these remarks he shews the names and generation of the powers, the square, cube, 4th, 5th, 6th, &c, which he calls *dynamis*, *cubus*, *dynamodinamis*, *dynamocubus*, *cubocubus*, according to the sum of the indices of the powers; and he marks these powers with the initials thus  $\delta^{\bar{v}}$ ,  $\kappa^{\bar{v}}$ ,  $\delta\delta^{\bar{v}}$ ,  $\delta\kappa^{\bar{v}}$ ,  $\kappa\kappa^{\bar{v}}$ , &c: the unknown quantity he calls simply  $\alpha\rho\theta\mu\omicron\varsigma$ , *numerus*, the number; and in the solutions he commonly marks it by the final thus  $\zeta$ ; also he denotes the monades, or indefinite unit, by  $\mu^{\bar{o}}$ . Diophantus there remarks on the multiplication and division of simple species together, shewing what powers or species they produce; declares that minus ( $\lambda\epsilon\iota\psi\iota\varsigma$ ) multiplied by minus produces plus ( $\nu\pi\alpha\rho\epsilon\iota\mu$ ); but that minus multiplied by plus, produces minus; and that the mark used for minus is  $\pi$ , namely the  $\psi$  inverted and curtailed, but he uses no mark for plus, but a word or conjunction copulative. As to the operations, viz. of addition, subtraction, multiplication, and division of compound species, or those connected by plus and minus, Diophantus does not teach, but supposes his reader to know them. He then remarks on the preparation or simplifying of the equations that are derived from the questions, which we call reduction of equations, by collecting like quantities together, adding quantities that are minus, and subtracting such as are plus, called by the moderns Transposition, so as to bring the equation to simple terms, and then depressing it to a lower degree by equal division when the powers of the unknown quantity are in every term: which preparation, or reduction of the complex equation, being now made, or reduced to what we call a final equation, Diophantus goes no farther, but barely says what the root or *res ignota* is, without giving any rules for finding it, or for the resolution of equations; thereby intimating that such rules were to be found in some other work, done either by himself or others. Of the body of the

work, *Lib. 1* contains 43 questions, concerning one, two, three, or four unknown numbers, having certain relations to each other, viz. concerning their sums, differences, ratios, products, squares, sums and differences of squares, &c, &c; but none of them concerning either square or cubic numbers. *Lib. 2* contains 36 questions. The first five questions are concerning two numbers, though only one condition is given in each question; but he supplies another by assuming the numbers in a given ratio, viz. as 2 to 1. The 6th and 7th contain each two conditions: then in the 8th question he first comes to treat of square numbers, which is this, to divide a given square number into two other squares; and the 9th is the same, but performed in a different way: the rest, to the end, are, almost all, about one, two, or three squares. *Lib. 3* contains 24 questions concerning squares, chiefly including three or four numbers. *Lib. 4* begins with cubes; the first of which is this, to divide a given number into two cubes whose sides shall have a given sum: here he has occasion to cube the two binomials  $5+n$  and  $5-n$ ; the manner of doing which shews that he knew the composition of the cube of a binomial; and many other places manifest the same thing. Only part of the questions in this book are concerning cubes; the rest are relating to squares. Two or three questions in this book have general solutions, and the theorems deduced are general, and for any numbers indefinitely; but all the other questions, in all the four books, find only particular numbers. *Lib. 5* is also concerning square and cube numbers, but of a more difficult kind, beginning with some that relate to numbers in geometrical progression. *Lib. 6* contains 26 propositions, concerning right-angled triangles; such as to make their sides, areas, perimeters, &c, &c, squares or cubes, or rational, &c. In some parts of this book it appears, that he was acquainted with the composition of the 4th power of the binomial root, as he sets down all the terms of it; and, from his great skill in such matters, it seems probable that he was acquainted with the composition of other higher powers, and with other parts of Algebra, besides what are here treated of. At the end is part of a book, in 10 propositions, concerning arithmetical progressions, and multangular or polygonal numbers. Diophantus once mentions a compound quadratic equation; but the resolution of his questions is by simple equations, and by means of only one unknown letter or character, which he chooses so ingeniously, that all the other unknown quantities in the question are easily expressed by it, and the final equation reduced to the simplest form which it seems the question can admit of. Sometimes he substitutes for a number sought immediately, and then expresses the other numbers or conditions by it: at other times he substitutes for the sum or difference, &c, and thence derives the rest, so as always to obtain the expressions in the simplest form. Thus, if the sum of two numbers be given, he substitutes for their difference; and if the difference be given, he substitutes for their sum: and in both cases he has the two numbers easily expressed by adding and subtracting the half sum and half difference; and so in other cases he uses other similar ingenious notations. In short, the chief excellence in this collection of questions, which seems to be only a set of exercises to some rules which had been given elsewhere, is the neat mode of substitution or notation; which being once made, the reduc-



tion to the final equation is easy and evident : and there he leaves the solution, only mentioning that the root or *επιθυμιος* is so much. Upon the whole, this work is treated in a very able and masterly manner, manifesting the utmost address and knowledge in the solutions, and forcing a persuasion that the author was deeply skilled in the science of Algebra, to some of the most abstruse parts of which these questions or exercises relate. However, as he contrives his assumptions and notations so as to reduce all his conditions to a simple equation, or at least a simple quadratic, it does not appear what his knowledge was in the resolution of compound or affected equations.

But although Diophantus was the first author on Algebra that we now know of, it was not from him, but from the Moors or Arabians that we received the knowledge of Algebra in Europe, as well as that of most other sciences. And it is matter of dispute who were the first inventors of it ; some ascribing the invention to the Greeks, while others say that the Arabians had it from the Persians, and these from the Indians, as well as the arithmetical method of computing by ten characters, or digits ; but the Arabians themselves say it was invented amongst them by one *Mahomet ben Musa*, or son of Moses, who it seems flourished about the 8th or 9th century. It is more probable, however, that Mahomet was not the inventor, but only a person well skilled in the art ; and it is farther probable, that the Arabians drew their first knowledge of it from Diophantus or other Greek writers, as they did that of Geometry and other sciences, which they improved and translated into their own language ; and from them it was that we received these sciences, before the Greek authors were known to us, after the Moors settled in Spain, and after the Europeans began to hold communications with them, and that our countrymen began to travel amongst them to learn the sciences. And according to the testimony of Abulpharagius, the Arithmetic of Diophantus was translated in Arabic by Mahomet ben-yahya Baziani. But whoever were the inventors and first cultivators of Algebra, it is certain that the Europeans first received the knowledge, as well as the name, from the Arabians or Moors, in consequence of the close intercourse which subsisted between them for several centuries. And it appears that the art was pretty generally known, and much cultivated, at least in Italy, if not in other parts of Europe also, long before the invention of printing, as many writers upon the art are still extant in the libraries of manuscripts ; and the first authors, presently after the invention of printing, speak of many former writers on this subject, from whom they learned the art.

It was chiefly among the Italians that this art was first cultivated in Europe. And the first author whose works we have in print, was Lucas Pacioli, or Lucas de Burgo, a Cordelier, or Minorite Friar. He wrote several treatises of Arithmetic, Algebra, and Geometry, which were printed in the years 1470, 1476, 1481, 1487, and in 1494 his principal work, intitled *Summa de Arithmetica, Geometria, Proportioni, et Proportionalita*, is a very masterly and complete treatise on those sciences, as they then stood. In this work he mentions various former writers, as Euclid, St. Augustine, Sacrobosco or Halifax, Boetius, Prodocimo, Giordano,

Biagio da Parma, and Leonardus Pisanus, from whom he learned those sciences. The order of the work is, 1st Arithmetic, 2d Algebra, and 3d Geometry. Of the Arithmetic the contents, and the order of them, are nearly as follow. First, of numbers figurate, odd and even, perfect, prime and composite, and many others. Then of Common Arithmetic in 7 parts, namely numeration or notation, addition, subtraction, multiplication, division, progression, and extraction of roots. Before him, he says, duplation and mediation, or doubling and halving, were accounted two rules in Arithmetic ; but that he omits them, as being included in multiplication and division. He ascribes the present notation and method of Arithmetic to the Arabs ; and says that according to some the word *Abaco* is a corruption of *Modo Arabico*, but that according to others it was from a Greek word. All those primary operations he both performs and demonstrates in various ways, many of which are not in use at present, proving them not only by what is called casting out the nines, but also by casting out the sevens, and otherwise. In the extraction of roots he uses the initial *R* for a root ; and when the roots can be extracted, he calls them discrete or rational ; otherwise surd, or indiscrete, or irrational. The square root is extracted much the same way as at present, namely, dividing always the last remainder by double the root found ; and so he continues the surd roots continually nearer and nearer in vulgar fractions. Thus, for the root of 6, he first finds the nearest whole number 2, and the remainder 2 also ; then  $\frac{2}{4}$  or  $\frac{1}{2}$  is the first correction, and  $2\frac{1}{2}$  the second root : its square is  $6\frac{1}{4}$ , therefore  $\frac{1}{4}$  divided by 5, or  $\frac{1}{25}$  is the next correction, and  $2\frac{1}{2}$  minus  $\frac{1}{25}$ , or  $2\frac{24}{25}$  is the 3d root : its square is  $6\frac{1}{5}$ , therefore  $\frac{1}{5}$  divided by  $4\frac{24}{25}$ , or  $\frac{1}{98}$  is the 3d correction, which gives  $2\frac{48}{98}$  for the 4th root, whose square exceeds 6 by only  $\frac{1}{384}$  : and so on continually : and this process he calls approximation. He observes that fractions, which he sets down the same way as we do at present, are extracted, by taking the root of the denominator, and of the denominated, for so he calls the numerator : and when mixed numbers occur, he directs to reduce the whole to a fraction, and then extract the roots of its two terms as above : as if it be  $12\frac{1}{4}$  ; this he reduces to  $\frac{49}{4}$ , and then the roots give  $\frac{7}{2}$  or  $3\frac{1}{2}$  : in like manner he finds that  $4\frac{1}{2}$  is the root of  $20\frac{1}{4}$  ;  $5\frac{1}{2}$  the root of  $30\frac{1}{4}$  ; “ and so on (he adds) in infinitum ;” which shews that he knew how to form the series of squares by addition. He then extracts the cube root, by a rule much the same as that which is used at present ; from which it appears that he was well acquainted with the co-efficients of the binomial cubed, namely 1, 3, 3, 1 ; and he directs how the operation may be continued “ in infinitum ” in fractions, like as in the square root. After this, he describes geometrical methods for extracting the square and cube roots instrumentally : he then treats professedly of vulgar fractions, their reductions, addition, subtraction, and other operations, much the same as at present : then of the rule-of-three, gain-and-lost, and other rules used by merchants.

Pacioli next enters on the algebraical part of this work, which he calls “ *L'Arte Magiore ; ditta dal vulgo la Regola de la Cosa, over Alghebra e Almucabala* :” which last name he explains by *restauratio & oppositio*, and assigns



assigns as a reason for the first name, because it treats of things above the common affairs in business, which make the *Arte Minore*. Here he ascribes the invention of Algebra to the Arabians, and denominates the series of powers, with their marks or abbreviations, as *n<sup>o</sup>*. or *numero*, the absolute or known number; *co.* or *cosa*, the thing or 1st power of the unknown quantity; *ce.* or *censo*, the product or square; *cu.* or *cubo*, the cube, or 3d power; *ce. ce.* or *censo de censo*, the square-squared, or 4th power; *p<sup>o</sup>.* *r<sup>o</sup>.* or *primo relato*, or 5th power; *ce. cu.* or *censo de cubo*, the square of the cube, or 6th power; and so on, compounding the names or indices according to the multiplication of the numbers 2, 3, &c, and not according to their sum or addition, as used by Diophantus. He describes also the other characters made use of in this part, which are for the most part no more than the initials or other abbreviations of the words themselves; as *R.* for *radici*, the root; *R. R.* *radici de radici*, the root of the root; *R. u.* *radici universale*, or *radici legata*, or *radici unita*; *R. cu.* *radici cuba*; and  $\sqrt[3]{\phantom{x}}$  *quantita*, quantity; *p* for *piu* or plus, and *m* for *meno* or minus; and he remarks that the necessity and use of these two last characters are for connecting, by addition or subtraction, different powers together; as 3 *co.* *p.* 4 *ce.* *m.* 5 *cu.* *p.* 2 *ce. ce.* *m.* 6 *n<sup>o</sup>.* that is, 3 *cosa* *piu* 4 *censo* *meno* 5 *cubo* *piu* 2 *censo-censo* *meno* 6 *numeri*, or, as we now write the same thing,  $3x + 4x^2 - 5x^3 + 2x^4 - 6$ . He first treats very fully of proportions and proportionalities, both arithmetical and geometrical, accompanied with a large collection of questions concerning numbers in continued proportion, resolved by a kind of Algebra. He then treats of *el Cataym*, which he says, according to some, is an Arabic or Phenician word, and signifies the Double Rule of False Position: but he here treats of both single and double position, as we do at present, dividing the *el Cataym* into single and double. He gives also a geometrical demonstration of both the cases of the errors in the double rule, namely when the errors are both plus or both minus, and when the one error is plus and the other minus; and adds a large collection of questions, as usual. He then goes through the common operations of Algebra, with all the variety of signs, as to plus and minus; proving that, in multiplication and division, like signs give plus, and unlike signs give minus. He next treats of different roots in *infinitum*, and the extraction of roots; giving also a copious treatise on radicals or surds, as to their addition, subtraction, multiplication and division, and that both in square roots and cube roots, and in the two together, much the same as at present. He makes here a digression concerning the 15 lines in the 10th book of Euclid, treating them as surd numbers, and teaching the extraction of the roots of the same, or of compound surds or binomials, such as of  $23 p R 448$ , or of  $R 18 p R 10$ ; and gives this rule, among several others, namely: Divide the first term of the binomial into two such parts that their product may be  $\frac{1}{4}$  of the number in the second term; then the roots of those two parts, connected by their proper sign *p* or *m*, is the root of the binomial; as in this  $23 p R 448$ , the two parts of 23 are 7 and 16, whose product, 112, is  $\frac{1}{4}$  of 448, therefore their roots give  $4 p R 7$  for the root  $R u. 23 p R 448$ . He next treats of equations both simple and quadratic, or simple and compound, as he

calls it; and this latter he performs by completing the square, and then extracting the root, just as we do at present. He also resolves equations of the simple 4th power, and of the 4th combined with the 2d power, which he treats the same way as quadratics; expressing his rules in a kind of bad verse, and giving geometrical demonstrations of all the cases. He uses both the roots or values of the unknown quantity, in that case of the quadratics which has two positive roots; but he takes no notice of the negative roots in the other two cases. But as to any other compound equations, such as the cube and any other power, or the 4th and 1st, or 4th and 3d, &c, he gives them up as impossible, or at least says that no general rule has yet been found for them, any more, he adds, than for the quadrature of the circle. —The remainder of this part is employed on rules in trade and merchandize, such as Fellowship, Barter, Exchange, Interest, Composition or Alligation, with various other cases in trade. And in the third part of the work, he treats of Geometry, both theoretical and practical.

From this account of Lucas de Burgo's book, we may perceive what was the state of Algebra about the year 1500, in Europe; and probably it was much the same in Africa and Asia, from whence the Europeans had it. It appears that their knowledge extended only to quadratic equations, of which they used only the positive roots; that they used only one unknown quantity; that they had no marks or signs for either quantities or operations, excepting only some few abbreviations of the words or names themselves; and that the art was only employed in resolving certain numeral problems. So that either the Africans had not carried Algebra beyond quadratic equations, or else the Europeans had not learned the whole of the art, as it was then known to the former. And indeed it is not improbable but this might be the case: for whether the art was brought to us by an European, who, travelling in Africa, there learned it; or whether it was brought to us by an African; in either case we might receive the art only in an imperfect state, and perhaps far short of the degree of perfection to which it had been carried by their best authors. And this suspicion is rendered rather probable by the circumstance of an Arabic manuscript, said to be on cubic equations, deposited in the Library of the university of Leyden by the celebrated Warner, bearing a title which in Latin signifies *Omar Ben Ibrahim al'Ghajamai Algebra cubicarum aequationum, sive de problematum solidorum resolutione*; and of which book I am in some hopes of procuring either a copy or a translation, by means of my worthy friend Dr. Damen, the learned Professor of Mathematics in that university, and by that means to throw some light on this doubtful subject.

Since this was written, death has prematurely put an end to the useful labours of this ingenious and worthy successor of Gravesande.

After the publication of the books of Lucas de Burgo, the science of Algebra became more generally known, and improved, especially by many persons in Italy; and about this time, or soon after, namely about the year 1505, the first rule was there found out by Scipio Ferreus, for resolving one case of a compound cubic equation. But this science, as well as other branches of



Mathematics, was most of all cultivated and improved there by Hieronymus Cardan of Bononia, a very learned man, whose arithmetical writings were the next that appeared in print, namely in the year 1539, in 9 books, in the Latin language, at Milan, where he practised physic, and read public lectures on Mathematics; and in the year 1545 came out a 10th book, containing the whole doctrine of cubic equations, which had been in part revealed to him about the time of the publication of his first 9 books. And as it is only this 10th book which contains the new discoveries in Algebra, I shall here confine myself to it alone, as it will also afford sufficient occasion to speak of his manner of treating Algebra in general. This book is divided into 40 chapters, in which the whole science of cubic equations is most amply and ably treated. Chap. 1 treats of the nature, number and properties of the roots of equations, and particularly of single equations that have double roots. He begins with a few remarks on the invention and name of the art: calls it *Ars Magna*, or *Cosa*, or *Rules of Algebra*, after Lucas de Burgo and others: says it was invented by Mahomet, the son of one Moses an Arabian, as is testified by Leonardus Pisanus; and that he left four rules or cases, which perhaps only included quadratic equations: that afterwards three derivatives were added by an unknown author, though some think by Lucas Pacioli; and after that again three other derivatives, for the cube and 6th power, by another unknown author; all which were resolved like quadratics: that then Scipio Ferreus, Professor of Mathematics at Bononia, about 1505, found out the rule for the case *cubum & rerum numero equalium*, or, as we now write it,  $x^3 + bx = c$ , which he speaks of as a thing admirable: that the same thing was next afterwards found out, in 1535, by Tartalea, who revealed it to him, Cardan, after the most earnest intreaties: that, finally, by himself and his quondam pupil Lewis Ferrari, the cases are greatly augmented and extended, namely, by all that is not here expressly ascribed to others; and that all the demonstrations of the rules are his own, except only three adopted from Mahomet for the quadratics, and two of Ferrari for cubics.

He then delivers some remarks, shewing that all square numbers have two roots, the one positive, and the other negative, or, as he calls them, *vera & falsa*, true and fictitious or false; so the *æstimatio rei*, or root, of 9, is either 3 or  $-3$ ; of 16 it is 4 or  $-4$ ; the 4th root of 81 is 3 or  $-3$ ; and so on for all even *denominations* or powers. And the same is remarked on compound cases of even powers that are added together; as if  $x^4 + 3x^2 = 28$ , then the *æstimatio x* is  $= 2$  or  $-2$ ; but that the form  $x^4 + 12 = 7x^2$  has four answers or roots, in real numbers, two plus and two minus, viz. 2 or  $-2$ , and  $\sqrt{3}$  or  $-\sqrt{3}$ ; while the case  $x^4 + 12 = 6x^2$  has no real roots; and the case  $x^4 = 2x^2 + 8$  has two, namely 2 and  $-2$ : and in like manner for other even powers. So that he includes both the positive and negative roots; but rejects what we now call imaginary ones. I here express the cases in our modern notation, for brevity sake, as he commonly expresses the terms by words at full length, calling the absolute or known term the *numero*, the 1st power the *res*, the 2d the *quadratum*, the 3d the *cubum*, and so on, using no mark for the unknown

quantity, and only the initials *p* and *m* for plus and minus, and *R* for radix or root. The *res* he sometimes calls *positio*, and *quantitas ignota*; and in stating or setting down his equations, he, as well as Lucas de Burgo before him, sets down the terms on that side where they will be plus, and not minus.

On the other hand, he remarks that the odd denominations, or powers, have only one *æstimatio*, or root, and that true or positive, but none fictitious or negative, and for this reason, that no negative number raised to an odd power, will give a positive number; so of  $2x = 16$ , the root is 8 only; and if  $2x^3 = 16$ , the root is 2 only: and if there be ever so many odd denominations, added together, equal to a number, there will be only one *æstimatio* or root; as if  $x^3 + 6x = 20$ , the only root is 2. But that when the signs of some of the terms are different as to plus and minus, they may have more roots; and he shews certain relations of the co-efficients, when they have two or more roots: so the equation  $x^3 + 16 = 12x$  has two *æstimations*, the one true or 2, and the other fictitious or  $-4$ , which he observes is the same as the true *æstimatio* of the case  $x^3 = 12x + 16$ , having only the sign of the absolute number changed from the former, the 3d root 2 being the same as the first, which therefore he does not count. He next shews what are the relations of the co-efficients when a cubic equation has three roots, of which two are true, and the 3d fictitious, which is always equal to the sum of the other two, and also equal to the true root of the same equation with the sign of the absolute number changed: thus, in the equation  $x^3 + 9 = 12x$ , the two true roots are 3 and  $\sqrt{5\frac{1}{4}} - 1\frac{1}{2}$ , and the fictitious one is  $-\sqrt{5\frac{1}{4}} - 1\frac{1}{2}$ , which last is the same as the true root of  $x^3 = 12x + 9$ , viz.  $\sqrt{5\frac{1}{4}} + 1\frac{1}{2}$ ; and he here infers generally that the fictitious *æstimatio* of the case  $x^3 + c = bx$ , always answers to the true root of  $x^3 = bx + c$ . Cardan also shews what the relation of the co-efficients is, when the case has no true roots, but only one fictitious root, which is the same as the true root of the reciprocal case, formed by changing the sign of the absolute number. Thus, the case  $x^3 + 21 = 2x$  has no true root, and only one false root, viz.  $-3$ , which is the same as the true root of  $x^3 = 2x + 21$ : and he shews in general, that changing the sign of the absolute number in such cases as want the 2d term, or changing the signs of the even terms when it is not wanting, changes the signs of all the three roots, which he also illustrates by many examples; thus, the roots of  $x^3 + 11x^2 = 72$ , are  $+\sqrt{40} - 4$ , and  $-3$ , and  $-\sqrt{40} - 4$ ; and the roots of  $x^3 + 72 = 11x^2$ , are  $-\sqrt{40} + 4$ , and  $+3$ , and  $+\sqrt{40} + 4$ .

And he further observes, that the sum of the three roots, or the difference between the true and fictitious roots, is equal to 11, the co-efficient of the 2d term. He also shews how certain cubic cases have one, or two, or three roots, according to circumstances: that the case  $x^4 + d = bx^2$  has sometimes four roots, and sometimes none at all, that is, no real ones: that the case  $x^3 + bx = ax^2 + c$  may have three true *æquations*, or positive roots, but no fictitious or negative ones; and for this reason, that the odd powers of minus being minus, and the even powers plus, the two terms  $x^3 + bx$  would be negative, and equal to a positive sum  $ax^2 + c$ , which



which is absurd: and farther, that the case  $x^3 + ax^2 + bx = c$  has three roots, one plus and two minus, which are the same, with the signs changed, as the roots of the case  $x^3 + bx + c = ax^2$ . He also shews the relation of the co-efficients when the equation has only one real root, in a variety of cases: but that the case  $x^3 + c = ax^2 + bx$  has always one negative root, and either two positive roots, or none at all; the number of roots failing by pairs, or the impossible roots, as we now call them, being always in pairs. Of all these circumstances Cardan gives a great many particular examples in numeral co-efficients, and subjoins geometrical demonstrations of the properties here enumerated; such as, that the two corresponding or reciprocal cases have the same root or roots, but with different signs or affections; and how many true or positive roots each case has.

Upon the whole, it appears from this short chapter, that Cardan had discovered most of the principal properties of the roots of equations, and could point out the number and nature of the roots, partly from the signs of the terms, and partly from the magnitude and relations of the co-efficients. He shews in effect, that when the case has all its roots, or when none are impossible, the number of its positive roots is the same as the number of changes in the signs of the terms, when they are all brought to one side: that the co-efficient of the 2d term is equal to the sum of all the roots positive and negative collected together, and consequently that when the 2d term is wanting, the positive roots are equal to the negative ones: and that the signs of all the roots are changed, by changing only the signs of the even terms: with many other remarks concerning the nature of equations.

In chap. 2, Cardan enumerates all the cases of compound equations of the 2d and 3d order, namely, 3 quadratics, and 19 cubics; with 44 derivatives of these two, that is, of the same kind, with higher denominations.

In chap. 3 are treated the roots of simple cases, or simple equations, or at least that will reduce to such, having only two terms, the one equal to the other. He directs to depress the denominations equally, as much as they will, according to the height of the least; then divide by the number or co-efficient of the greatest; and lastly extract the root on both sides. So if  $20ax^3 = 180ax^5$ , then  $20 = 180x^2$ , and  $\frac{1}{9} = x^2$ , and  $x = \frac{1}{3}$ .

Chap. 4 treats of both general and particular roots, and contains various definitions and observations concerning them. It is here shewn that the several cases of quadratics and cubics have their roots of the following forms or kinds, namely that the case

$x^2 = ax + b$  has its root of this kind  $\sqrt{19 + 3}$ ,

$x^2 + ax = b$  has its root of this kind  $\sqrt{19 - 3}$ ,

$x^2 + b = ax$  has two roots like  $3 + \sqrt{2}$  and  $3 - \sqrt{2}$ ,

$x^3 = bx + c$  has its root of this kind  $\sqrt[3]{4} + \sqrt[3]{2}$ .

$x^3 + bx = c$  has its root of this kind  $\sqrt[3]{4} - \sqrt[3]{2}$ ,

$a^3 = ax^2 + c$  has this kind of root  $\sqrt[3]{16 + 2} + \sqrt[3]{4}$

$x^3 + ax^2 = c$  has this kind of root  $\sqrt[3]{16 - 2} + \sqrt[3]{4}$ ,

where the three parts  $\sqrt[3]{16}$ , 2,  $\sqrt[3]{4}$ , are in continual proportion.

Chap. 5 treats of the æstimation of the lowest degree of compound cases, that is, affected quadratic equations; giving the rule for each of the three cases, which con-

sists in completing the square, &c, as at present, and which it seems was the method given by the Arabians; and proving them by geometrical demonstrations from Eucl. I. 43, and II. 4 and 5, in which he makes some improvement of the demonstrations of Mahomet. And hence it appears that the work of this Arabian author was in being, and well known in Cardan's time.

Chap. 6, on the methods of finding new rules, contains some curious speculations concerning the squares and cubes of binomial and residual quantities, and the proportions of the terms of which they consist, shewn from geometrical demonstrations, with many curious remarks and properties, forming a foundation of principles for invelligating the rules for cubic equations.

Chap. 7 is on the transmutation of equations, shewing how to change them from one form to another, by taking away certain terms out of them; as  $x^3 + ax^2 = c$ , to  $y^3 = by + d$ , &c. The rules are demonstrated geometrically; and a table is added, of the forms into which any given cases will reduce; which transformations are extended to equations of the 4th and 5th order. And hence it appears that Cardan knew how to take away any term out of an equation.

Chap. 8 shews generally how to find the root of any such equation as this  $ax^m = x^n + b$ , where  $m$  and  $n$  are any exponents whatever, but  $n$  the greater; and the rule is, to separate or divide the co-efficient  $a$  into two such parts  $z$  and  $a - z$ , as that the absolute number  $b$

shall be equal to  $\frac{a - z}{a - z} \cdot z^{\frac{n}{n-m}}$ , the product of the one part  $a - z$ , and the  $\frac{m}{n-m}$  power of the other

part: then the root  $x$  is  $= z^{\frac{1}{n-m}}$ . The rule is general for quadratics, cubics, and all the higher powers; and could not have been formed without the knowledge of the composition of the terms from the roots of the equation.

Chap. 9 and 10 contain the resolution of various questions producing equations not higher than quadratics.

Chap. 11 is of the case or form  $x^3 + 3bx = 2c$ . Cardan now comes to the actual resolution of the first case of cubic equations. He begins with relating a short history of the invention of it, observing that it was first found out, about 30 years before, by Scipio Ferrens of Bononia, and by him taught to Antonio Maria Florido of Venice, who having a contest afterwards with Nicolas Tartalea of Brescia, it gave occasion to Tartalea to find it out himself, who after great entreaties taught it to Cardan, but suppressed the demonstration. By help of the rule alone, however, Cardan of himself discovered the source or geometrical investigation, which he gives here at large, from Eucl. II. 4. In this process he makes use of the Greek letters  $\alpha$ ,  $\epsilon$ ,  $\gamma$ ,  $\delta$ , &c, to denote certain indefinite numbers or quantities, to render the investigation general; which may be considered as the first instance of such literal notation in Algebra. He then gives the rule in words at length, which comes to this,

$$x = \sqrt[3]{\sqrt{c^2 + b^3} + c} - \sqrt[3]{\sqrt{c^2 + b^3} - c};$$

illustrating it in a variety of examples; in the resolution of



of which, he extracts the cubic roots of such of the binomials as will admit of it, by some rule which he had for that purpose; such as  $x^3 + 6x = 20$ , which  $x = \sqrt[3]{\sqrt{108} + 10} - \sqrt[3]{\sqrt{108} - 10} - \sqrt{3} - 1 = 2 = \sqrt{3} + 1$ .

Chap. 12, of the case  $x^3 = 3bx + 2c$ . This he treats exactly as the last, and finds the rule

$$x = \sqrt[3]{c + \sqrt{c^2 - b^3}} + \sqrt[3]{c - \sqrt{c^2 - b^3}};$$

which he illustrates by many examples, as usual. But when  $b^3$  exceeds  $c^2$ , which has since been called the irreducible case, he refers to another following book, called *Aliza*, for other rules of solution, to overcome this difficulty, about which he took infinite pains.

Chap. 13, of the case  $x^3 + 2c = 3bx$ . This case, by a geometrical process, he reduces to the case in the last chapter: thus, find the æstimation  $y$  of the case  $y^3 = 3by + 2c$ , having the same co-efficients as the given case  $x^3 + 2c = 3bx$ ; then is  $x = \frac{1}{2}y \pm \frac{1}{2}\sqrt{12b - 3y^2}$ , giving two roots. He shews also how to find the second root, when the first is known, independent of the foregoing case. From this relation of these two cases he deduces several corollaries, one of which is, that the æstimation or root of the case  $y^3 = 3by + 2c$ , is equal to the sum of the roots of the case  $x^3 + 2c = 3bx$ . As in the example  $y^3 = 16y + 21$ , whose æstimation is  $\sqrt[3]{9\frac{1}{4}} + 1\frac{1}{2}$ , which is equal to the sum of 3 and  $\sqrt[3]{9\frac{1}{4}} - 1\frac{1}{2}$ , the two roots of the case  $x^3 + 21 = 16x$ .

In chapters 14, 15, and 16, he treats of the three cases which contain the 2d and 3d powers, but wanting the first power, according to all the varieties of the signs; which he performs by exterminating the 2d term, or that which contains the 2d power of the unknown quantity  $x$ , by substituting  $y \pm \frac{1}{3}$  the co-efficient of that term for  $x$ , and so reducing these cases to one of the former. In these chapters Cardan sometimes also gives other rules; thus, for the case  $x^3 + 3ax^2 = 2c$ , find first the æstimation  $y$  of the case  $y^3 = 3ay + \sqrt{2c}$ , then is  $x = y^2 - 3a$ : also for the case  $x^3 + 2c = 3ax^2$ , first find the two roots of  $y^3 + 2c = 3ay\sqrt[3]{2c}$ , then is  $x = \frac{\sqrt[3]{4c^2}}{y}$  the two values of  $x$  according to the two values of  $y$ .

He here also gives another rule, by which a second æstimation or root is found, when the first is known, namely, if  $e$  be the first estimation or value of  $x$  in the case  $x^3 + 2c = 3ax^2$ , then is the other value of

$$x = \frac{1}{2}\sqrt{3a - e \cdot 3a + 3e} + \frac{1}{2} \cdot 3a - e.$$

In chapters 17, 18, 19, 20, 21, 22, 23, Cardan treats of the cases in which all the four terms of the equation are present; and this he always effects by taking away the 2d term out of the equation, and so reducing it to one of the foregoing cases which want that term, giving always geometrical investigations, and adding a great many examples of every case of the equations.

Chap. 24, of the 4d derivative cases; which are only higher powers of the forms of quadratics and cubics.

Chap. 25, of imperfect and special cases: containing many particular examples when the co-efficients have certain relations amongst them, with easy rules for

finding the roots; also 8 other rules for the irreducible case  $x^3 = bx + c$ .

Chap. 26, in like manner, contains easy rules for biquadratics, when the co-efficients have certain special relations.

Then the following chapters, from chap. 27 to chap. 38, contain a great number of questions and applications of various kinds, the titles of which are these: *De transitu capituli specialis in capitulum speciale*; *De operationibus radicum pronicarum seu mixtarum* & *Allellarum*; *De regula modi*; *De regula Aurca*; *De regula Magna*, or the method of finding out solutions to certain questions; *De regula equalis positionis*, being a method of substituting for the half sum and half difference of two quantities, instead of the quantities themselves; *De regula inæqualiter ponendi, seu proportionis*; *De regula medii*; *De regula aggregati*; *De regula liberæ positionis*; *De regula falsum ponendi*, in which some quantities come out negative; *Quomodo excident partes & denominationes multiplicando*. Among the foregoing collection of questions, which are chiefly about numbers, there are some geometrical ones, being the application of Algebra to Geometry, such as, In a right-angled triangle, given the sum of each leg and the adjacent segment of the hypotenuse, made by a perpendicular from the right angle, to determine the area &c; with other such geometrical questions, resolved algebraically.

Chap. 39, *De regula qua pluribus positionibus invenimus ignotam quantitatem*; which is employed on biquadratic equations. After some examples of his own, Cardan gives a rule of Lewis Ferrari's, for resolving all biquadratics, namely by means of a cubic equation, which Ferrari investigated at his request, and which Cardan here demonstrates, and applies in all its cases. The method is very general, and consists in forming three squares, thus: first, complete one side of the equation up to a square, by adding or subtracting some multiples or parts of some of its own terms on both sides, which it is always easy to do: 2d, supposing now the three terms of this square to be but one quantity, viz, the first term of another square to which this same side is to be completed, by annexing the square of a new and assumed indeterminate quantity, with double the product of the roots of both; which evidently forms the square of a binomial, consisting of the assumed indeterminate quantity and the root of the first square: 3d, the other side of the equation is then made to become the square of a binomial also, by supposing the product of its 1st and 3d terms to be equal to the square of half its 2d term; for it consists of only three terms, or three different denominations of the original unknown quantity: then this equality will determine the value of the assumed indeterminate quantity, by means of a cubic equation, and from it, that of the original ignota, by the equal roots of the 2d and 3d squares. Here we have a notable example of the use of assuming a new indeterminate quantity to introduce into an equation, long before Des Cartes was born, who made use of a like assumption for a similar purpose. And this method is very general, and is here applied to all forms of biquadratics, either having all their terms, or wanting some of them. To illustrate this rule I shall here set down the process of one of his examples, which is this,  $x^4 + 4x + 8 = 10x^2$ . Now first sub-



tract  $2x^2 + 4x + 7$  from both sides, then the first becomes a square, viz,  $x^4 - 2x^2 + 1$  or  $\overline{x^2 - 1}^2 = 8x^2 - 4x - 7$ . Next assume the indeterminate  $y$ , and subtract  $2y(x^2 - 1) - y^2$  from both sides, making the first side again a square, viz,  $\overline{x - 1}^2 = 2y \cdot \overline{x - 1} + y^2$  or  $\overline{x^2 - 1 - y^2} = \overline{8 - 2y \cdot x^2 - 4x + y^2 + 2y - 7}$ . Of this latter side, make the product of the 1st and 3d terms equal to the square of half the 2d term, that is,  $\overline{8 - 2y \cdot y^2 + 2y - 7} = 2^2$ , which reduces to  $y^3 + 30 = 2y^2 + 15y$ ; the positive roots of which are  $y = 2$  or  $\sqrt{15}$ ; and hence, using 2 for  $y$ , the equation of equal squares becomes  $\overline{x^2 - 1 - y^2}$  or  $\overline{x^2 - 3}^2 = 4x^2 - 4x + 1$ , the roots of which give  $x^2 - 3 = 2x \pm 1$ ; and hence  $x^2 = 2x + 2$ , or  $x^2 + 2x = 4$ ; the two positive roots of which are  $\sqrt{3} + 1$  and  $\sqrt{5} - 1$ , which are two of the values of  $x$  in the given equation  $x^4 + 4x + 8 = 10x^2$ . The other roots he leaves to be tried by the reader.

The 40th, or last, chap. is entitled, Of modes of general supposition relating to this art; with some rules of an unusual kind; and æstimation or roots of a nature different from the foregoing ones. Some of these are as follow: If  $x^3 = ax^2 + c$ , and  $x - a = y$ , and  $x : y :: c : d$ ; then is  $y^3 + ay^2 = d$ .

Secondly, if  $x^3 + ax^2 = c$ ,

and  $y^3 = ay^2 + c$ ,

then is  $x + a : y - a :: y^2 : x^2$ .

Thirdly, when  $x^3 + c = ax^2$ , the square will be taken away, by putting  $x = y + \frac{1}{3}a$ ; and then the equation becomes  $y^3 + c - 2(\frac{1}{3}a)^3 = \frac{1}{3}a^2y$ .

Cardan adds some other remarks concerning the solutions of certain cases and questions, all evincing the accuracy of his skill, and the extent of his practice; and then he concludes the book with a remark concerning a certain transformation of equations, which quite astonishes us to find that the same person who, through the whole work, has shewn such a profound and critical skill in the nature of equations, and the solution of problems, should yet be ignorant of one of the most obvious transmutations attending them, namely increasing or diminishing the roots in any proportion. Cardan having observed that the form  $x^3 = bx + c$  may

be changed into another similar one, viz,  $y^3 = \frac{b}{c}y + \sqrt{\frac{1}{c}}$ ,

of which the co-efficient of the term  $y$  is the quotient arising from the co-efficient of  $x$  divided by the absolute number of the first equation: and that the absolute number of the 2d equation is the root of the quotient of 1 divided by the said absolute number of the first; he then adds, that finding the æstimation or root of the one equation from that of the other is very difficult, *valde difficilis*.

It is matter of wonder that Cardan, among so many transmutations, should never think of substituting instead of  $x$  in such equations, another positio or root, greater or less than the former in any indefinite proportion, that is, multiplied or divided by a given number; for this would have led him immediately to the same transformation as he makes above, and that by a way which would have shewn the constant proportion be-

tween the two roots. Thus, instead of  $x$  in the given form - - - - -  $x^3 = bx + c$ , substitute  $dy$ , and it becomes - - -  $d^3y^3 = bdy + c$ ; and this divided

by  $d^3$  becomes - - -  $y^3 = \frac{b}{d^2}y + \frac{c}{d^3}$ ; and here if  $d$

be taken  $= \sqrt{c}$ , it becomes  $y^3 = \frac{b}{c}y + \sqrt{\frac{1}{c}}$ ; which is

the transformation in question, and in which it is evident

that  $x$  is  $= y\sqrt{c}$ , and  $y = \frac{x}{\sqrt{c}}$ . Instead of this, Car-

dan gives the following strange way of finding the one root  $x$  from the other  $y$ , when this latter is by any means known; viz, Multiply the first given equation

by  $y^2x + 1$ , then add  $\frac{x^2}{4y^2}$  to both sides, and lastly ex-

tract the roots of both, which can always be done, as they will always be both of them squares; and the roots will give the value of  $x$  by a quadratic equation.

Thus,  $x^3 = bx + c$  multiplied by  $y^2x + 1$  gives

$y^2x^4 + x^3 = by^2x^2 + \overline{cy^2 + b} \cdot x + c$ ; add  $\frac{x^2}{4y^2}$ , then

$y^2x^4 + x^3 + \frac{x^2}{4y^2} = by^2 + \frac{1}{4y^2} \cdot x^2 + \overline{b + cy^2} \cdot x + c$ ; and

the roots are  $yx^2 + \frac{x}{2y} = \sqrt{(by^2 + \frac{1}{4y^2})x^2 + (b + cy^2)x + c}$ ;

and this 2d side of the equation he says will always have a root also. It is indeed true that it will have an exact root; but the reason of it is not obvious, which is, be-

cause  $y$  is the root of the equation  $y^3 = \frac{b}{c}y + \sqrt{\frac{1}{c}}$ .

Cardan has not shewn the reason why this happens; but I apprehend he made it out in this manner, viz, similar to the way in which he forms the last square in the case of biquadratic equations, namely, by making the product of the 1st and 3d terms equal to the square of half the 2d term: thus, in the present case, it is

$4c(by^2 + \frac{1}{4y^2}) = (b + cy^2)^2$ , which reduces to  $y^3 = \frac{b}{c}y$

+  $\sqrt{\frac{1}{c}}$  the equation in question. Therefore taking  $y$

the root of the equation  $y^3 = \frac{b}{c}y + \sqrt{\frac{1}{c}}$ , and sub-

stituting its value in the quantity  $(by^2 + \frac{1}{4y^2})x^2 + (b + cy^2)x + c$ , this will become a complete square.

#### Of Cardan's *Libellus de Aliza Regula*.

Subjoined to the above Treatise on cubic equations, is this *Libellus de Aliza regula*, or the algebraic logistics, in which the author treats of some of the abstruser parts of Arithmetic and Algebra, especially cubic equations, with many more attempts on the irreducible case  $x^3 = bx + c$ . This book is divided into 60 chapters; but



but I shall only set down the titles of some few of them, whose contents require more particular notice.

Chap. 4. *De modo redigendi quantitates omnes, que dicuntur latera prima ex decimo Euclidis in compendium.* He treats here of all Euclid's irrational lines, as surd numbers, and performs various operations with them.

Chap. 5. *De consideratione binomiorum & reciforum, &c; ubi de estimatione capitulorum.* Contains various operations of multiplying compound numbers and surds.

Chap. 6. *De operationibus p: & m: (i. e. + and -) secundum communem usum.* Here it is shewn that, in multiplication and division, plus always gives the same signs, and minus gives the contrary signs. So also in addition, every quantity retains its own sign; but in subtraction they change the signs. That the  $\sqrt{+}$ , or the square root of plus, is +; but the  $\sqrt{-}$ , or the square root of minus, is nothing as to common use: (but of this below.) That  $\sqrt{-}$  is -; as  $\sqrt{-8}$  is -2. That a residual, composed of + and - may have a root also composed of + and -: So  $\sqrt{5} - \sqrt{24}$  is  $= \sqrt{3} - \sqrt{2}$ . The rules for the signs in multiplication and division are illustrated by this example; to divide 8 by  $2 + \sqrt{6}$  or  $\sqrt{6} + 2$ . Take the two corresponding residuals  $2 - \sqrt{6}$  and  $\sqrt{6} - 2$ , and by these multiply both the divisor and dividend; then the products are + and - respectively, and the quotients still both alike. Thus,

Divid.	Divis.	Divid.	Divis.
8	$\sqrt{6} + 2$	8	$2 + \sqrt{6}$
$\sqrt{6} - 2$	$\sqrt{6} - 2$	$2 - \sqrt{6}$	$2 - \sqrt{6}$
$\sqrt{384} - 16$	divide + 4	$16 - \sqrt{384}$	div. - 2
Quot. $\sqrt{96} - 8$ .		Quot. $\sqrt{96} - 8$ .	

And this method of performing division of compound surds, was fully taught before him, by Lucas de Burgo, namely, reducing the compound divisor to a simple quantity, by multiplying by the corresponding quantity, having the sign changed.

In chap. 11 and 18, and elsewhere, Cardan makes a general notation of  $a, b, c, d, e, f$ , for any indefinite quantities, and treats of them in a general way.

Cap. 2. *De contemplatione p: & m: (or + and -), & quod m: in m: facit p: & de causis horum juxta veritatem.* Cardan here demonstrates geometrically that, in multiplication and division, like signs give plus, and unlike signs give minus. And he illustrates this numerically, by squaring the quantity 8, or  $6 + 2$ , or  $10 - 2$ , which must all produce the same thing, namely 64.

Among many of the chapters which treat of the irreducible case  $x^3 = bx + c$ , there is a peculiar kind of way given in chap. 31, which is entitled *De estimatione generali  $x^3 = bx + c$  solida vocata, & operationibus ejus*; in which he shews how to approximate to the root of that case, in a manner similar to approximating the square root and cube root of a number. The rule he uses for this purpose, is the 3d in chap. 25 of the last book, and it is this: Divide  $b$  into two parts, such that the sum of the products of each, multiplied by the square of the other, may be equal to  $\frac{1}{4}c$ ; then the sum of the roots of these parts is the estimation or value of  $x$  required. So, of this equation  $x^3 = 10x + 24$ ; the two parts are 9 and 1, and their roots 3 and 1, and their sum  $4 = x$ , as in the margin.

$$\begin{array}{r} 9 \quad 1 \\ \times \\ 3 \quad 1 \\ \hline 3 + 9 \text{ or } 12 = \frac{1}{4}c. \\ x = 3 + 1 = 4 \end{array}$$

Again, take  $x^3 = 6x + 1$ . Here he invents a new notation to express the root or *radix*, which he calls *solida*, viz,  $x = \sqrt{\text{solida } 6 \text{ in } \frac{1}{2}}$ , that is, the roots of the two parts of 6, so that each part multiplied by the root of the other, the two products may be  $\frac{1}{2}$  or  $\frac{1}{2}c$ . Then to free this from fractions, and make the operation easier, multiply that root by some number as suppose 4, that is the square part 6 by the square of 4, and the solid part  $\frac{1}{2}$  by the cube of 4; then  $x = \frac{1}{4}\sqrt{\text{solida } 96 \text{ in } 32}$ . Now, by a few trials, it is found that the parts are nearly  $95\frac{8}{9}$  and  $\frac{1}{9}$ , which give too much,

or  $95\frac{1}{9}$  and  $\frac{1}{9}$ , which give too little, and thereof  $95\frac{1}{9}$  and  $\frac{1}{9}$  are still nearer. Divide both by  $4^2$  or 16, then  $5\frac{1}{16}$  and  $\frac{1}{16}$  are the quot. And the sum of their roots, or  $x = \sqrt{5\frac{1}{16}} + \sqrt{\frac{1}{16}}$  is nearly the value of the root  $x$ .

Cap. 42. *De duplici equatione comparanda in capitulo cubi & numeri equalium rebus.* Treats of the two positive roots of that case, neglecting the negative one; and shewing, not only that that case has two such roots, but that the same number may be the common root of innumerable equations.

Cap. 57. *De tractatione estimationis generalis capituli  $x^3 = bx + c$ .* Cardan here again resumes the consideration of the irreducible case, making ingenious observations upon it, but still without obtaining the root by a general rule. In this place also, as well as elsewhere, he shews how to form an equation in this case, that shall have a given binomial root, as suppose  $\sqrt{m} + n$ , where the equation will be  $x^3 = (m + 3n^2)x + 2n(m - n^2)$ , having  $\sqrt{m} + n$  for one root, namely the positive root. From which it appears that he was well acquainted with the composition of cubic equations from given roots.

Cap. 59. *De ordine & exemplis in binomiis secundo & quinto.* Contains a great many numeral forms of the same irreducible case  $x^3 = bx + c$ , with their roots; from which are derived these following cases, with many curious remarks. When

$$x^3 = (c + 1)x + c, \text{ then } x = \sqrt{c + \frac{1}{4}} + \frac{1}{2}.$$

$$x^3 = (\frac{1}{2}c + 4)x + c, \text{ then } x = \sqrt{\frac{1}{2}c + \frac{1}{4}} + \frac{1}{2}.$$

$$x^3 = (\frac{1}{3}c + 9)x + c, \text{ then } x = \sqrt{\frac{1}{3}c + \frac{9}{4}} + \frac{3}{2}.$$

$$x^3 = (\frac{1}{4}c + 16)x + c, \text{ then } x = \sqrt{\frac{1}{4}c + \frac{16}{4}} + \frac{4}{2}.$$

$$x^3 = (\frac{1}{n}c + n^2)x + c, \text{ then } x = \sqrt{\frac{1}{n}c + \frac{n^2}{4}} + \frac{n}{2}.$$

Cap. 60. *Demonstratio generalis capituli cubi equalis rebus & numero.* This demonstration of the irreducible case is geometrical, like all the rest. Some more ingenious remarks are again added, as if he reluctantly finished the book without perfectly overcoming the difficulty of the irreducible case. Cardan here also uses the letters  $a$  and  $b$  for any two indefinite numbers, in order to shew the form and manner of the arithmetical operations: thus  $\frac{a}{b}$  is the fraction for their quotient; also

$$\sqrt{\frac{a}{b}} \text{ or } \frac{\sqrt{a}}{\sqrt{b}} \text{ the square root of that quotient, and}$$

$$\sqrt[3]{\frac{a}{b}} \text{ or } \frac{\sqrt[3]{a}}{\sqrt[3]{b}} \text{ the cube root of it, \&c.}$$

Having



Having considered the chief contents of Cardan's algebra, it will now be proper to sum them up, and set down a list of the improvements made by him, as collected from his writings :

And 1st, Tartalea having only communicated to him the rules for resolving these three cases of cubic equations, viz,

$x^3 + bx = c$ , } he from thence raised a very large and  
 $x^3 = bx + c$ , } complete work, laying down rules for all  
 $x^3 + c = bx$ , } forms and varieties of cubic equations, having all their terms, or wanting any of them, and having all possible varieties of signs ; demonstrating all these rules geometrically ; and treating very fully of almost all sorts of transformations of equations, in a manner heretofore unknown.

2nd, It appears that he was well acquainted with all the roots of equations that are real, both positive and negative ; or, as he calls them, true and fictitious ; and that he made use of them both occasionally. He also shewed, that the even roots of positive quantities, are either positive or negative ; that the odd roots of negative quantities, are real and negative ; but that the even roots of them are impossible, or nothing as to common use. He was also acquainted with,

3d, The number and nature of the roots of an equation, and that partly from the signs of the terms, and partly from the magnitude and relation of the coefficients. He also knew,

4th, That the number of positive roots is equal to the number of changes of the signs of the terms.

5th, That the coefficient of the second term of the equation, is the difference between the positive and negative roots.

6th, That when the second term is wanting, the sum of the negative roots is equal to the sum of the positive roots.

7th, How to compose equations that shall have given roots.

8th, That, changing the signs of the even terms, changes the signs of all the roots.

9th, That the number of roots failed in pairs ; or what we now call impossible roots were always in pairs.

10th, To change the equation from one form to another, by taking away any term out of it.

11th, To increase or diminish the roots by a given quantity. It appears also,

12th, That he had a rule for extracting the cube root of such binomials as admit of extraction.

13th, That he often used the literal notation  $a, b, c, d$ , &c.

14th, That he gave a rule for biquadratic equations, suiting all their cases ; and that, in the investigation of that rule, he made use of an assumed indeterminate quantity, and afterwards found its value by the arbitrary assumption of a relation between the terms.

15th, That he applied Algebra to the resolution of geometrical problems. And

16th, That he was well acquainted with the difficulty of what is called the irreducible case, viz,  $x^3 = bx + c$ , upon which he spent a great deal of time, in attempting to overcome it. And though he did not fully succeed in this case, any more than other persons have done since, he nevertheless made many ingenious observations about it, laying down rules for many particular

forms of it, and shewing how to approximate very nearly to the root in all cases whatever.

#### OF TARTALEA.

Nicholas Tartalea, or Tartaglia, of Brescia, was contemporary with Cardan, and was probably older than he was, but I do not know of any book of Algebra published by him till the year 1546, the year after the date of Cardan's work on Cubic Equations, when he printed his *Questi & Inventioni diverse*, at Venice, where he resided as a public lecturer on mathematics. This work is dedicated to our king Henry the VIIIth of England, and consists of 9 books, containing answers to various questions which had been proposed to him at different times, concerning mechanics, statics, hydrostatics, &c. ; but it is only the 9th, or last, that we shall have occasion to take notice of in this place, as it contains all those questions which relate to arithmetic and algebra. These are all set down in chronological order, forming a pretty collection of questions and solutions on those subjects, with a short account of the occasion of each of them. Among these, the correspondence between him and Cardan forms a remarkable part, as we have here the history of the invention of the rules for cubic equations, which he communicated to Cardan under the promise, and indeed oath, to keep them secret, on the 25th of March 1539. But, notwithstanding his oath, finding that Cardan published them in 1545, as above related, it seems Tartalea published the correspondence between them in revenge for his breach of faith ; and it elsewhere appears, that many other sharp bickerings passed between them on the same account, which only ended with the death of Tartalea, in the year 1557. It seems it was a common practice among the mathematicians, and others, of that time, to send to each other nice and difficult questions, as trials of skill, and to this cause it is that we owe the principal questions and discoveries in this collection, as well as many of the best discoveries of other authors. The collection now before us contains questions and solutions, with their dates, in a regular order, from the year 1521, and ending in 1541, in 42 dialogues, the last of which is with an English gentleman, namely, Mr. Richard Wentworth, who it seems was no mean mathematician, and who learned some algebra, &c, of Tartalea, while he resided at Venice. The questions at first are mostly very easy ones in arithmetic, but gradually become more difficult, and exercising simple and quadratic equations, with complex calculations of radical quantities : all shewing that he was well skilled in the art of Algebra as it then stood, and that he was very ingenious in applying it to the solutions of questions. Tartalea made no alteration in the notation or forms of expression used by Lucas de Burgo, calling the first power of the unknown quantity, in his language, *cosa*, the second power *censu*, the third *cubo*, &c, and writing the names of all the operations in words at length, without using any contractions, except the initial *R* for root or radicality. So that the only thing remarkable in this collection, is the discovery of the rules for cubic equations, with the curious circumstances attending the same.



The first two of these were discovered by Tartalea in the year 1530, namely for the two cases  $x^3 + ax^2 = c$ , and  $x^3 = ax^2 + c$ , as appears by Quest. 14 and 25 of this collection, on occasion of a question then proposed to him by one Zuanni de Tonini da Coi or Colle, John Hill, who kept a school at Brescia. And from the 25th letter we learn, that he discovered the rules for the other two cases  $x^3 + bx = c$ , and  $x^3 = bx + c$ , on the 12th and 13th of February 1535, at Venice, where he had come to reside the year before. And the occasion of it was this: There was then at Venice one Antonio Maria Fiore or Florido, who, by his own account, had received from his preceptor Scipio Farreo, about thirty years before, a general rule for resolving the case  $x^3 + bx = c$ . Being a captious man, and presuming on this discovery, Florido used to brave his contemporaries, and by his insults provoked Tartalea to enter into a wager with him, that each should propose to the other thirty different questions; and that he who soonest resolved those of his adversary, should win from him as many treats for himself and friends. These questions were to be proposed on a certain day at some weeks distance; and Tartalea made such good use of his time, that eight days before the time appointed for delivering the propositions, he discovered the rules both for the case  $x^3 + bx = c$ , and the case  $x^3 = bx + c$ . He therefore proposed several of his questions so as to fall either on this latter case, or on the cases of the cube and square, expecting that his adversary would propose his in the former. And what he suspected fell out accordingly; the consequence of which was, that on the day of meeting Tartalea resolved all his adversary's questions in the space of two hours, without receiving one answer from Florido in return; to whom, however, Tartalea generously remitted the forfeit of the thirty treats won of him.

Question 31 first brings us acquainted with the correspondence between Tartalea and Cardan. This correspondence is very curious, and would well deserve to be given at full length in their own words, if it were not too long for this place. I may enlarge farther upon it under the article *Cubic Equations*; but must here be content with a brief abstract only. Cardan was then a respectable physician, and lecturer in mathematics at Milan; and having nearly finished the printing of a large work on Arithmetic, Algebra, and Geometry, and having heard of Tartalea's discoveries in cubic equations, he was very desirous of drawing those rules from him, that he might add them to his book before it was finished. For this purpose he first applied to Tartalea, by means of a third person, a bookseller, whom he sent to him, in the beginning of the year 1539, with many flattering compliments, and offers of his services and friendship, &c, accompanied with some critical questions for him to resolve, according to the custom of the times. Tartalea however refused to disclose his rules to any one, as the knowledge of them gained him great reputation among all people, and gave him a great advantage over his competitors for fame, who were commonly afraid of him on account of those very rules. He only sent Cardan therefore, at his request, a copy of the thirty questions which had been proposed to him in the contest with Florido. Not to be rebuffed so easily, Cardan next applied, in the most urgent manner,

by letter to Tartalea; which however procured from him only the solution of some other questions proposed by Cardan, with a few of the questions that had been proposed to Florido, but none of their solutions. Finding he could not thus prevail, with all his fair promises, Cardan then fell upon another scheme. There was at Milan a certain Marquis dal Vasto, a great patron of Cardan, and, it was said, of learned men in general. Cardan conceived the idea of making use of the influence of this nobleman to draw Tartalea to Milan, hoping that then, by personal intreaties, he should succeed in drawing the long-concealed rules from him. Accordingly he wrote a second letter to Tartalea, much in the same strain with the former, strongly inviting him to come and spend a few days in his house at Milan, and representing that, having often commended him in the highest terms to the marquis, this nobleman desired much to see him; for which reason Cardan advised him, as a friend, to come to visit them at Milan, as it might be greatly to his interest, the marquis being very liberal and bountiful; and he besides gave Tartalea to understand, that it might be dangerous to offend such a man by refusing to come, who might, in that case, take offence, and do him some injury. This manœuvre had the desired effect: Tartalea on this occasion laments to himself in these words, "By this I am reduced to a great dilemma; for if I go not to Milan, the marquis may take it amiss, and some evil may befall me on that account; I shall therefore go, although very unwillingly." When he arrived at Milan however, the marquis was gone to Vigevano, and Tartalea was prevailed on to stay three days with Cardan, in expectation of the marquis returning, at the end of which he set out from Milan, with a letter from Cardan, to go to Vigevano to that nobleman. While Tartalea was at Milan the three days, Cardan plied him by all possible means to draw from him the rules for the cubic equations; and at length, just as Tartalea was about to depart from Milan, on the 25th of March 1539, he was overcome by the most solemn protestations of secrecy that could be made. Cardan says, "I shall swear to you on the holy evangelists, and by the honour of a gentleman, not only never to publish your inventions, if you reveal them to me; but I also promise to you, and pledge my faith as a true christian, to note them down in cyphers, so that after my death no other person may be able to understand them." To this Tartalea replies, "If I refuse to give credit to these assurances, I should deservedly be accounted utterly void of belief. But as I intend to ride to Vigevano, to see his excellency the marquis, as I have been here now these three days, and am weary of waiting so long; whenever I return therefore, I promise to shew you the whole." Cardan answers, "Since you determine at any rate to go to Vigevano, to the marquis, I shall give you a letter for his excellency, that he may know who you are. But now before you depart, I intreat you to shew me the rule for the equations, as you have promised." "I am content," says Tartalea: "But you must know, that to be able on all occasions to remember such operations, I have brought the rule into rhyme; for if I had not used that precaution, I should often have forgot it; and although my rhymes are not very good, I do not value that, as it is sufficient



sufficient that they serve to bring the rule to mind as often as I repeat them. I shall here write the rule with my own hand, that you may be sure I give you the discovery exactly." These rude verses contain, in rather dark and enigmatical language, the rule for these three cases, viz.

$x^3 + bx = c$ ,  $x^3 = bx + c$ ,  $x^3 + c = bx$ , } which differ however only in the sign of one quantity, and the rule amounts to this: Find two numbers,  $z$  and  $y$ , such that their difference in the first case, and their sums in the 2d and 3d, may be equal to  $c$  the absolute number, and their product equal to the cube of  $\frac{1}{3}$  of  $b$  the coefficient of the less power; then the difference of their cube roots will be equal to  $x$  in the first case, and the sum of their cube roots equal to  $x$  in the 2d and 3d cases: that is, taking  $z - y = c$  in the 1st case, or  $z + y = c$  in the 2d and 3d, and  $zy = (\frac{1}{3}b)^3$ ; then  $x = \sqrt[3]{z} - \sqrt[3]{y}$  in the first case, and  $x = \sqrt[3]{z} + \sqrt[3]{y}$  in the other two. At parting, T. fails not again to remind C. of his obligation:

"Now your excellency will remember not to break your promised faith, for if unhappily you should insert these rules either in the work you are now printing, or in any other, although you should even give them under my name, and as of my invention, I promise and swear that I shall immediately print another work that will not be very pleasing to you." "Doubt not, says C. but that I shall observe what I have promised: Go, and rest secure as to that point: and give this letter of mine to the marquis." It should seem however that T. was much displeased at having suffered himself to be worried as it were out of his rules, for as soon as he quitted Milan, instead of going to wait upon the marquis, he turned his horse's head, and rode straight home to Venice, saying to himself, "By my faith I shall not go to Vigeveno, but shall return to Venice, come of it what will."

After T's departure it seems C. applied himself immediately to resolving some examples in the cubic equations by the new rules, but not succeeding in them, for indeed he had mistaken the words, as it was very easy to do in such bad verses, having mistaken  $(\frac{1}{3}b)^3$  for  $\frac{1}{3}b^3$ , or the cube of  $\frac{1}{3}$  of the coefficient, for  $\frac{1}{3}$  of the cube of the coefficient; accordingly we find him writing to T. in fourteen days after the above, blaming him much for his abrupt departure without seeing the marquis, who was so liberal a prince he said, and requesting T. to resolve him the example  $x^3 + 3x = 10$ . This T. did to his satisfaction, rightly guessing at the nature of his mistake; and concludes his answer with these emphatical words, "Remember your promise." On the 12th of May following C. returns him a letter of thanks, together with a copy of his book, saying, "As to my work, just finished, to remove your suspicion, I send you a copy, but unbound, as it is yet too fresh to be beaten. But as to the doubt you express lest I may print your inventions, my faith which I gave you with an oath should satisfy you; for as to the finishing of my book, that could be no security, as I could always add to it whenever I please. But on account of the dignity of the thing, I excuse you for not relying on that which you ought to have done, namely on the faith of a gentleman, instead of the finishing of a book, which might at any time be enlarged by the addition of new chapters; and there are besides a thousand other ways. But the security consists in this, that there is no

greater treachery than to break one's faith, and to aggrrieve those who have given us pleasure. And when you shall try me, you will find whether I be your friend or not, and whether I shall make an ungrateful return for your friendship, and the satisfaction you have given me."

It was within less than two months after this, however, that T. received the alarming news of Cardan's shewing some symptoms of breaking the faith he had so lately pledged to him; this was in a letter from a quondam pupil of his, in which he writes, "A friend of mine at Milan has written to me, that Dr. Cardano is composing another algebraical work, concerning some lately-discovered rules; hence I imagine they may be those same rules which you told me you had taught him; so that I fear he will deceive you." To which T. replies, "I am heartily grieved at the news you inform me of, concerning Dr. Cardano of Milan; for if it be true, they can be no other rules but those I gave him; and therefore the proverb truly says, 'That which you wish not to be known, tell to nobody.' Pray endeavour to learn more of this matter, and inform me of it."

Tartalea, after this, kept on the reserve with Cardan, not answering several letters he sent him, till one written on the 4th of August the same year, 1539, complaining greatly of T's neglect of him, and farther requesting his assistance to clear up the difficulty of the irreducible case  $x^3 = bx + c$ , which C. had thus early been embarrassed with: he says that when  $(\frac{1}{3}b)^3$  exceeds  $(\frac{1}{2}c)^2$ , the rule cannot be applied to the equation in hand, because of the square root of the negative quantities. On this occasion T. turns the tables on C. and plays his own game back upon him; for being aware of the above difficulty, and unable to overcome it himself, he wanted to try if C. could be encouraged to accomplish it, by pretending that the case might be done, though in another way. He says thus to himself, "I have a good mind to give no answer to this letter, no more than to the other two. However I will answer it, if it be but to let him know what I have been told of him. And as I perceive that a suspicion has arisen concerning the difficulty or obstacle in the rule for the case  $x^3 = bx + c$ , I have a mind to try if he can alter the data in hand, so as to remove the said obstacle, and to change the rule into another form, although I believe indeed that it cannot be done; however there is no harm in trying."—"M. Hieronime, I have received your letter, in which you write that you understand the rule for the case  $x^3 = bx + c$ , but that when  $(\frac{1}{3}b)^3$  exceeds  $(\frac{1}{2}c)^2$ , you cannot resolve the equation by following the rule, and therefore you request me to give you the solution of this equation  $x^3 = 9x + 10$ . To which I reply, that you have not used a good method in that case, and that your whole process is intirely false. And as to resolving you the equation you have sent, I must say that I am very sorry that I have already given you so much as I have done, for I have been informed, by a credible person, that you are about to publish another Algebraical work, and that you have been boasting through Milan of having discovered some new rules in algebra. But, take notice, that if you break your faith with me, I shall certainly keep my word with you, nay, I even assure you to do more than I promised." In Cardan's answer to this he says, "You have been misinformed as to my intention to



publish more on Algebra. But I suppose you have heard something about my work *de mysteriis eternitatis*, which you take for some Algebra I intend to publish. As to your repenting of having given me your rules, I am not to be moved from the faith I promised you for any thing you say." To this, and many other things contained in the same letter, T. returned no answer, being still suspicious of Cardan's intentions, and declining any more correspondence with him. This however did not discourage C. for we find him writing again to T. on the 5th of January, 1540, to clear up another difficulty which had occurred in this business, namely to extract the cube root of the binomials, of which the two parts of the rule always consisted, and for which purpose it seems C. had not yet found out a rule. On this occasion he informs T. that his quondam competitor Zuanne Colle had come to Milan, where, in some contests between them, Colle gave Cardan to understand that he had found out the rules for the two cases  $x^3 + bx = c$ , and  $x^3 = bx + c$ , and farther that he had discovered a general rule for extracting the cube roots of all such binomials as can be extracted; and that, in particular, the cube root of  $\sqrt{108} + 10$  is  $\sqrt{3} + 1$ , and that of  $\sqrt{108} - 10$  is  $\sqrt{3} - 1$ , and consequently that  $\sqrt[3]{\sqrt{108} + 10} - \sqrt[3]{\sqrt{108} - 10}$  is  $= \sqrt{3} + 1 - \sqrt{3} - 1 = 2$ . He then earnestly entreats T. to try to find out the rule, and the solution of certain other questions which had been proposed to him by Colle. By this letter T. is still more confirmed in his resolution of silence; so that, without returning any answer, he only sets down among his own memorandums some curious remarks on the contents of the letter, and then concludes to himself, "Wherefore I do not choose to answer him again, as I have no more affection for him than for M. Zuanne, and therefore I shall leave the matter between them." Among those remarks he sets down a rule for extracting the cube root of such binomials as can be extracted, and that is done from either member of the binomial alone, thus: Take either term of the binomial, and divide it into two such parts that one of them may be a complete cube, and the other part exactly divisible by 3; then the cube root of the said cubic part will be one term of the required root, and the square root of the quotient arising from the division of  $\frac{1}{3}$  of the 2d part by the cube root of the first, will be the other member of the root sought. This rule will be better understood in characters thus: let  $m$  be one member of the given binomial, whose cube root is sought, and let it be divided into the two parts  $a^3$  and  $3b$ , so that  $a^3 + 3b$  be  $= m$ ; then is  $a + \sqrt{\frac{b}{a}}$  the cube root required, if it have one. Thus in the quantity  $\sqrt{108} + 10$ , taking the term 10 for  $m$ , then 10 divides into 1 and 9, where  $a^3 = 1$  or  $a = 1$ , and  $3b = 9$  or  $b = 3$ : therefore  $a + \sqrt{\frac{b}{a}}$  becomes  $1 + \sqrt{3}$  for the cube root of  $\sqrt{108} + 10$ . And taking the other member  $\sqrt{108}$ , this divides into the two equal parts  $\sqrt{27}$  and  $\sqrt{27}$ , making  $a^3 = \sqrt{27}$ , and  $3b = \sqrt{27}$ ; hence  $a = \sqrt{3}$ , and  $b = \sqrt{3}$  also; consequently  $a + \sqrt{\frac{b}{a}}$  is  $= \sqrt{3} + \sqrt{\frac{\sqrt{3}}{\sqrt{3}}}$  or  $\sqrt{3} + 1$  for the cube root of the binomial sought, the same as before. "And thus, he adds, we may know whether any proposed binomial

or residual be a cube or a noncube; for if it be a cube, the same two terms for the root must arise from both the given terms separately; and if the two terms of the root cannot thus be brought to agree both ways, such binomial or residual will not be a cube." And thus ends the correspondence between them, at least for this time. But it seems they had still more violent disputes when C. in violation of his faith so often pledged to the contrary, published his work on cubic equations 4 years afterwards, viz, in the year 1545, of which we have before given an account, which disputes, it is said, continued till the death of Tartalea in the year 1557.

The last article in the volume contains a dialogue on some other forms of the cubic equations, in the year 1541, between T. and a Mr. Richard Wentworth, an English gentleman, who it seems had resided some time at Venice, on some public service from England, as T. in the dedication of the volume to Henry VIII. king of England, makes mention of him as "a gentleman of his sacred majesty." Mr. Wentworth had learned some mathematics of T. and being about to depart for England, requests T. to shew him his newly discovered rules for cubic equations, as a farewell-lesson; and it is worth while to note a few particulars in this conference, as they shew pretty nicely the limited knowledge of T. at that time, as to the nature and roots of such equations. T. had before, it seems, shewed Mr. W. the rules for the cases of the 3d and 1st powers, and now the latter desires him to do the same as to the three cases in which the 3d and 2d powers only are concerned. On this T. professes great gratitude to Mr. W. for many obligations, but desires to be excused from giving him the rules for these, because he says he intends soon to compose a new work on Arithmetic, Geometry, and Algebra, which he intends to dedicate to him, and in which he means to insert all his new discoveries. On Mr. W. urging him further, T. gives him the roots of some equations of that kind, as for instance:

If  $x^3 + 6x^2 = 100$ , then

$$x = \sqrt[3]{42} + \sqrt{17000} + \sqrt[3]{42} - \sqrt{17000} - 2,$$

$$\text{If } x^3 + 9x^2 = 100, \text{ then } x = \sqrt{24} - 2,$$

$$\text{If } x^3 + 3x^2 = 2, \text{ then } x = \sqrt{3} - 1,$$

$$\text{If } x^3 + 4 = 5x^2, \text{ then } x = \sqrt{8} + 2,$$

$$\text{If } x^3 + 6 = 7x^2, \text{ then } x = \sqrt{15} + 3;$$

but not the rules for finding them.

In the course of the conversation T. tells him that "all such equations admit of two different answers, and perhaps more; and hence it follows that they have, or admit of, two different rules, and perhaps more, the one more difficult than the other." And on Mr. W. expressing his wonder at this circumstance of a plurality of roots, T. replies, "It is however very true, though hardly to be believed, and indeed if experience had not confirmed it, I should scarcely have believed it myself." He then commits a strange blunder in an example which he takes to illustrate this by, namely the equation  $x^3 + 3x = 14$ , which, he says, it is evident has the number 2 for one of its roots; and yet, he adds, "whoever shall resolve the same equation by my rule, will find the value of  $x$  to be  $\sqrt[3]{7} + \sqrt{50} + \sqrt[3]{7} - \sqrt{50}$ , which is proved to be a true root by substituting



stituting it in the equation for  $x$ . And therefore, continues he, it is manifest that the case  $x^3 + bx = c$  admits of two rules, namely, one (as in the above example) which ought to give the value of  $x$  rational, viz 2, and the other is my rule, which gives the value of  $x$  irrational, as appears above; and there is reason to think that there may be such a rule as will give the value of  $x = 2$ , although our ancestors may not have found it out."—"And these two different answers will be found not only in every equation of this form  $x^3 + bx = c$ , when the value of  $x$  happens to be rational, as in the example  $x^3 + 3x = 14$  above, but the same will also happen in all the other five forms of cubic equations: and therefore there is reason to think that they also admit of two different rules; and by certain circumstances attending some of them, I am almost certain that they admit of more than two rules, as, God willing, I shall soon demonstrate." Now all this discourse shews a strange mixture of knowledge and ignorance: it is very probable that he had met with some equations which admit of a plurality of roots; indeed it was hardly possible for him to avoid it; but it seems he had no suspicion what the number of roots might be, nor that his reasoning in this instance was founded on an error of his own, mistaking the root  $x = \sqrt[3]{7 + \sqrt{50}}$

+  $\sqrt[3]{7 - \sqrt{50}}$ , of the equation  $x^3 + 3x = 14$ , for a different root from the number or root 2, when in reality it is the very same, as he might easily have found, if he had extracted the cube roots of the binomials by the rule which he himself had just given above for that purpose: for by that rule he would have found  $\sqrt[3]{7 + \sqrt{50}} = 1 + \sqrt{2}$ , and  $\sqrt[3]{7 - \sqrt{50}} = 1 - \sqrt{2}$ , and therefore their sum is  $2 = x$ , the same root as the other, which T. thought had been different. And besides this root 2, the equation in hand,  $x^3 + 3x = 14$ , admits of no other real roots. Nor does any equation of the same form,  $x^3 + bx = c$ , admit of more than one real root.

It seems also they had not yet discovered that all cases belong to the rules and forms for quadratic equations, which have only two powers in them, in which the exponent of the one is just double of the exponent of the other, as  $x^{2n} + bx^n = c$ ; but some particular cases only of this sort they had as yet ventured to refer to quadratics, as the case  $x^4 + bx^2 = c$ . But in the conclusion of this dialogue T. informs W. of another case of this sort which he had accomplished, as a notable discovery, in these words: "I well remember, says he, that in the year 1536, on the night of St. Martin, which was on a Saturday, meditating in bed when I could not sleep, I discovered the general rule for the case  $x^6 + bx^3 = c$ , and also for the other two, its accompanying cases, in the same night." And then he directs that they are to be resolved like quadratics, by completing the square, &c. And in these resolutions it is remarkable that he uses only the positive roots, without taking any notice of the negative ones.

Tartalea also published at Venice, in 1556, &c, a very large work, in folio, on Arithmetic, Geometry, and Algebra. This is a very complete and curious work upon the first two branches; but that of Algebra is carried no farther than quadratic equations,

called *book the first*, with which the work terminates. It is evidently incomplete, owing to the death of the author, which happened before this latter part of the work was printed, as appears by the dates, and by the prefaces. It appears also, from several parts of this work, that the author had many severe conflicts with Cardan and his friend Lewis Ferrari: and particularly, there was a public trial of skill between them, in the year 1547; in which it would seem that Tartalea had greatly the advantage, his questions mostly remaining unanswered by his antagonists.

OF MICHAEL STIFELIUS.

After the foregoing analysis of the works of the first algebraic writers in Italy, it will now be proper to consider those of their contemporaries in Germany; where, excepting for the discoveries in cubic equations, the art was in a more advanced state, and of a form approaching nearer to that of our modern Algebra; the state and circumstances indeed being so different, that one would almost be led to suppose they had derived their knowledge of it from a different origin.

Here Stifelius and Scheubelius were writers of the same time with Cardan and Tartalea, and even before their discoveries, or publication, concerning the rules for cubic equations, Stifelius's *Arithmetica Integra* was published at Norimberg in 1544, being the year before Cardan's work on cubic equations, and is an excellent treatise, both on Arithmetic and Algebra. The work is divided into three books, and is prefaced with an Introduction by the famous Melanchthon. The first book contains a complete and ample Treatise on Arithmetic, the second an Exposition of the 10th book of Euclid's Elements, and the third a Treatise of Algebra, and it is therefore properly the part with which we are at present concerned. In the dedication of this part, he ascribes the invention of Algebra to Geber, an Arabic Astronomer; and mentions besides, the authors Campanus, Christ. Rudolph, and Adam Ris, Risen, or Gigas, whose rules and examples he has chiefly given. In other parts of the book he speaks, and makes use also, of the works of Bretius, Campanus, Cardan (i. e. his Arithmetic published in 1539, before the work on cubic equations appeared), de Cusa, Euclid, Jordan, Milichius, Schonerus, and Stapulensis.

*Chap. 1. Of the Rule of Algebra, and its parts.* Stifelius here describes the notation and marks of powers, or denominations as he calls them, which marks for the several powers are thus:

1st, 2d, 3d, 4th, 5th, 6th, &c.  
 $\mathcal{Z}$ ,  $\mathcal{S}$ ,  $\mathcal{C}$ ,  $\mathcal{Q}$ ,  $\mathcal{F}$ ,  $\mathcal{H}$ , &c.

which are formed from the initials of the barbarous way in which the Germans pronounced and wrote the Latin and Italic names of the powers, namely, *res* or *cosa*, *zensus*, *cubo*, *zensi-zensus*, *sur-solid*, *zensi-cubo*, &c. And the *cos* or first power  $\mathcal{Z}$ , he calls the *radix* or root, which is the first time that we meet with this word in the printed authors. He also here uses the signs or characters,  $+$  and  $-$ , for addition and subtraction, and the first of any that I know of: for in Italy they used none of these characters for a long time after. He has no mark however for equality, but makes use of the word itself.

*Chap.*



*Chap. 2.* Of the Part of the Rule of Geber or Algebra: teaching the various reductions by addition, subtraction, multiplication, division, involution, and evolution, &c.

*Chap. 3.* Of the Algorithm of Cossic Numbers: teaching the usual operations of addition, subtraction, multiplication, division, involution, and extraction of roots, much the same as they are at present. Single terms, or powers, he calls simple quantities; but such as  $13 + 12$  a composite or compound, and  $22 - 8$  a defective one. In multiplication and division, he proves that like signs give  $+$ , and unlike signs  $-$ . He shews that the powers  $1, 2, 3, 4, \&c.$  form a geometrical progression from unity; and that the natural series of numbers  $0, 1, 2, 3, \&c.$  from  $0$ , are the exponents of the cossic powers; and he, for the first time, expressly calls them exponents: thus,

Exponents,  $0, 1, 2, 3, 4, 5, 6, \&c.$

Powers,  $1, 2, 3, 4, 5, 6, 7, 8, \&c.$

And he shews the use of the exponents, in multiplication, division, powers, and roots, as we do at present; viz, adding the exponents in multiplication, and subtracting them in division, &c. And these operations he demonstrates from the nature of arithmetical and geometrical progressions. It is remarkable that these compound denominations of the powers are formed from the simple ones according to the *products* of the exponents, while those of Diophantus are formed according to the *sums* of them; thus the 6th power here is  $3^4$  or quadrato-cubi, but with Diophantus it is cubo-cubi; and so of others. Which is presumptive evidence that the Europeans had not taken their Algebra immediately from him, independent of other proofs.

*Chap. 4.* Of the extraction of the roots of cossic numbers. He here treats of quadratic equations, which he resolves by completing the square, from Euclid II. 4 &c. Also quadratics of the higher orders, shewing how to resolve them in all cases, whatever the height may be, provided the exponents be but in arithmetical progression, as

$2, 1, 0$  } &c; where it is plain that he always  
 $4, 2, 0$  } counts  $0$  for the exponent of the un-  
 $6, 3, 0$  } known quantity in the absolute term.  
 $8, 4, 0$  }

*Chap. 5.* Of irrational cossic numbers, and of furd or negative numbers. In this treatise of radicals, or irrationals, he first uses the character  $\sqrt{\quad}$  to denote a root, and sets after it the mark of the power whose root is intended; as  $\sqrt{320}$  for the square root of  $20$ , and  $\sqrt[3]{20}$  for the cube root of the same, and so on. He treats here also of negative numbers, or what he calls furd or fictitious, or numbers less than  $0$ . On which he takes occasion to observe, that when a geometrical progression is continued downwards below  $1$ , then the exponents of the terms, or the arithmetical progression, will go below  $0$  into negative numbers, and will yet be the true exponents of the former; as in these,

Expon.	$-3$	$-2$	$-1$	$0$	$1$	$2$	$3$
Pow.	$\frac{1}{8}$	$\frac{1}{4}$	$\frac{1}{2}$	$1$	$2$	$4$	$8$

And he gives examples to shew that these negative exponents perform their office the same as the positive ones, in all the operations.

*Chap. 6.* Of the perfection of the Rule of Algebra, and of Secondary Roots. In the reduction of equations he uses a more general rule than those who had preceded him, who detailed the rule in a multitude of cases; instead of which, he directs to multiply or divide the two sides equally, to transpose the terms with  $+$  or  $-$ , and lastly to extract such root as may be denoted by the exponent of the highest power.

As to secondary roots, Cardan treated of a 2d *ignota* or unknown, which he called *quantitas*, and denoted it by the initial  $q$ , to distinguish it from the first. But here Stifelius, for distinction sake, and to prevent one root from being mistaken for others, assigns literal marks to all of them, as  $A, B, C, D, \&c.$  and then performs all the usual operations with them, joining them together as we do now, except that he subjoins the initial of the power, instead of its numeral exponent:

thus,  $3A$  into  $9B$  makes  $27AB$ ,

$33$  into  $4B$  makes  $123B$ ,

$24$  into  $4A3$  makes  $84A3$ ,

$1A$  squared makes  $1A3$ ,

$6$  into  $3C$  makes  $18C$ ,

$2A3$  into  $5A4$  makes  $10A5$ , &c, &c.

$84A3$  divided by  $44$  makes  $2A3$ , &c.

The square root of  $25A3$  is  $5A$ , &c.

Also  $2A$  added to  $22$  makes  $22 + 2A$ ,

and  $2A$  subtr. from  $22$  makes  $22 - 2A$ .

And he shews how to use the same, in questions concerning several unknown numbers; where he puts a different character for each of them, as  $2, A, B, C, \&c.$ ; he then makes out, from the conditions of the question, as many equations as there are characters; from these he finds the value of each letter, in terms of some one of the rest; and so, expelling them all but that one, reduces the whole to a final equation, as we do at present.

The remainder of the book is employed with the solutions of a great number of questions to exercise all the rules and methods; some of which are geometrical ones.

From this account of the state of Algebra in Stifelius, it appears that the improvements made by himself, or other Germans, beyond those of the Italians, as contained in Cardan's book of 1539, were as follow:

1st. He introduced the characters  $+$ ,  $-$ ,  $\sqrt{\quad}$ , for plus, minus, and root, or *radix*, as he calls it.

2d. The initials  $2, 3, 4, \&c.$  for the powers.

3d. He treated all the higher orders of quadratics by the same general rule.

4th. He introduced the numeral exponents of the powers,  $-3, -2, -1, 0, 1, 2, 3, \&c.$  both positive and negative, so far as integral numbers, but not fractional ones; calling them by the name *exponens*, exponent; and he taught the general uses of the exponents, in the several operations of powers, as we now use them, or the logarithms.

5th. And lastly, he used the general literal notation  $A, B, C, D, \&c.$  for so many different unknown or general quantities.

OF SCHEUBELIUS.

John Scheubelius published several books upon Arithmetic and Algebra. The one now before me, is intitled *Algebrae Compendiosa Facilisque Descriptio, quae depro-*  
*muntur*



*muntur magna Arithmetices miracula. Authore Johanne Scheubelio Mathematicarum Professore in Academia Tubingensi. Parisiis 1552.* But at the end of the book it is dated 1551. The work is most beautifully printed, and is a very clear and succinct treatise; and both in the form and matter much resembles a modern printed book. He says that the writers ascribe this art to Diophantus, which is the first time that I find this Greek author mentioned by the modern algebraists: he farther observes, that the Latins call it *Regula Rei & Censur*, the rule of the thing and the square (or of the 1st and 2d power); and the Arabs, Algebra. His characters and operations are much the same as those of Stifelius, using the signs and characters +, −, √, and the powers 2, 3, 4, &c, where the character 2 is used for 1 or unity, or a number, or the 0 power; prefixing also the numeral coefficients; thus  $44/3 + 113 + 312 - 534$ . He uses also the exponents 0, 1, 2, 3, &c, of the powers, the same way as Stifelius, before him. He performs the algebraical calculations, first in integers, and then in fractions, much the same as we do at present. Then of equations, which he says may be of infinite degrees, though he treats only of two, namely the first and second orders, or what we call simple and quadratic equations, in the usual way, taking however only the positive roots of these; and adverting to all the higher orders of quadratics, namely,  $x^4, ax^2, b$ ;

$$x^6, ax^3, b;$$

$$x^8, ax^4, b; \&c.$$

Next follows a tract on surds, both simple and compound, quadratic, cubic, binomial, and residual. Here he first marks the notation, observing that the root is either denoted by the initial of the word, or, after some authors, by the mark √: viz. the sq. root √:, the cube root w√:, and the 4th root, or root of the root thus v√:, which latter method he mostly uses. He then gives the Arithmetic of surds, in multiplication, division, addition, and subtraction. In these last two rules he squares the sum or difference of the surds, and then sets the root to the whole compound, which he calls *radix collecti*, what Cardan calls *radix universalis*. Thus  $\sqrt{12} \pm \sqrt{20}$  is ra. col.  $32 \pm \sqrt{960}$ . But when the terms will reduce to a common surd, he then unites them into one number; as  $\sqrt{27} + \sqrt{12}$  is equal  $\sqrt{75}$ . Also of cubic surds, and 4th roots. In binomial and residual surds, he remarks the different kinds of them which answer to the several irrational lines in the 10th book of Euclid's elements; and then gives this general rule for extracting the root of any binomial or residual  $a \pm b$ , where one or both parts are surds, and  $a$  the greater quantity, namely, that the square root of it is

$$\sqrt{\frac{a + \sqrt{a^2 - b^2}}{2}} \pm \sqrt{\frac{a - \sqrt{a^2 - b^2}}{2}}; \text{ which he illustrates by many examples. This rule will only succeed however, so as to come out in simple terms, in certain cases, namely, either when } a^2 - b^2 \text{ is a square, or when } a \text{ and } \sqrt{a^2 - b^2} \text{ will reduce to a common surd, and unite: in all other cases the root is in two compound surds, instead of one. He gives also another rule, which comes however to the same thing as the former, though by the words of them they seem to be different.}$$

Scheubelius wrote much about the time of Cardan and Stifelius. And as he takes no notice of cubic equations, it is probable he had neither seen nor heard any thing about them; which might very well happen, the one living in Italy, and the other in Germany. And, besides, I know not if this be the first edition of Scheubel's book: it is rather likely it is not, as it is printed at Paris, and he himself was professor of mathematics at Tubingen in Germany.

ROBERT RECORDE.

The first part of his Arithmetic was published in 1552; and the second part in 1557, under the title of, "The Whetstone of Witte, which is the seconde parte of Arithmetike: containing the Extraction of Rootes: The Cossike Practise, with the Rule of Equation: and the Workes of Surde Nombers." The work is in dialogue between the master and scholar; and is nearly after the manner of the Germans, Stifelius and Scheubelius, but especially the latter, whom he often quotes, and takes examples from. The chief parts of the work are, 1st. The properties of abstract and figurate numbers. 2nd. The extraction of the square and cube roots, much the same as at present. Here, when the number is not an exact power, but having some remainder over, he either continues the root into decimals as far as he pleases, by adding to the remainders always periods of cyphers; or else makes a vulgar fraction for the remaining part of the root, by taking the remainder for the numerator, and double the root for the denominator, in the square root; but in the cube root he takes for the nominator either the triple square of the root, which is Cardan's rule, or the triple square and triple root, with one more, which is Scheubel's rule. 3d. Of Algebra, or "Cossike Nombers." He uses the notation of powers with their exponents the same as Stifel, with all the operations in simple and compound quantities, or integers and fractions. And he gives also many examples of extracting the roots of compound algebraic quantities, even when the roots are from two to six terms, in imitation of the same process in numbers, just as we do at present; which is the first instance of this kind that I have observed. As of this quantity:

$$253 \text{ 2} + 80 \text{ 3} - 2633 - 633 (5 \text{ 4} + 83 - 94).$$

4th. "The Rule of Equation, commonly called Alber's Rule." He here, first of any, introduces the character =, for brevity sake. His words are, "And to avoide the tedious repetition of these woordes: is equalle to: I will sette as I doe often in woorke use. a paire of paralleles, or gemowe lines of one lengthe, thus: =, bicause noe 2 thynges can be moare equalle." He gives the rules for simple and quadratic equations, with many examples. He gives also some examples in higher compound equations, with a root for each of them, but gives no rule how to find it. 5th. "Of Surde Nombers." This is a very ample treatise on surds, both simple and compound, and surds of various degrees, as square, cubic, and biquadratic, marking the roots in Scheubel's manner, thus: √, w√, v√. He here uses the names binomial, binomial, and residual; but says they have been used by others before him, though this is the first place where I have observed the two latter.—Hence it appears that the things which chiefly are new in this author, are these three, viz.



1. The extraction of the roots of compound algebraic quantities.
2. The use of the terms binomial and residual.
3. The use of the sign of equality, or =.

OF PELETARIUS.

The first edition of this author's algebra was printed in 4to at Paris, in 1558, under this title, *Jacobi Peletarii Cenomani, de occulta parte Numerorum, quam Algebram vocant. Lib. duo.*

In the preface he speaks of the supposed authors of Algebra, namely Geber, Mahomet the son of Moses, an Arabian, and Diophantus. But he thinks the art older, and mentions some of his contemporary writers, or a very little before him, as Cardan, Stifel, Scheubel, Chr. Januarius; and a little earlier again, Lucas Paci-olus of Florence, and Stephen Villafrancus a Gaul.

Of the two books, into which the work is divided, the first is on rational, and the second on irrational or furd quantities; each being divided into many chapters. It will be sufficient to mention only the principal articles.

He calls the series of powers *numeri creati*, or derived numbers, or also radicals, because they are all raised from one root or *radix*. He names them thus, radix, quadratus cubus, quadrato-quadratus, or biquadratus, supersolidus, quadrato-cubus, &c; and marks them thus R, q, q<sup>2</sup>, q<sup>3</sup>, q<sup>4</sup>, q<sup>5</sup>, q<sup>6</sup>, q<sup>7</sup>, q<sup>8</sup>, &c. Of these he gives the following series in numbers, having the common ratio 2, with their marks set over them, and the exponents set over these again, in an arithmetical series, beginning at 0, thus:

0	1	2	3	4	5	6	7	8
1	R	q	q <sup>2</sup>	q <sup>3</sup>	q <sup>4</sup>	q <sup>5</sup>	q <sup>6</sup>	q <sup>7</sup>
1	2	4	8	16	32	64	128	256 &c.

And he shews the use of the exponents, the same as Stifelius and Scheubelius; like whom also he prefixes coefficients to quantities of all kinds, as also the radical  $\sqrt{\phantom{x}}$ . But he does not follow them in the use of the signs + and —, but employs the initials *p* and *m* for the same purpose. After the operations of addition, &c, he performs involution and evolution also much the same way as at present: thus, in powers, raise the coefficient to the power required, and multiply the exponent, or sign, as he calls it, by 2, or 3, or 4, &c, for the 2nd, 3d, 4th, &c, power; and the reverse for extraction: and hence he observes, if the number or coefficient will not exactly extract, or the sign do not exactly divide, the quantity is a furd.

After the operations of compound quantities, and fractions, and reduction of equations, namely, simple and quadratic equations, as usual, in chap. 16, *De Inveniendis generatim Radicibus Denominatorum*, he gives a method of finding the roots of equations among the divisors of the absolute number, when the root is rational, whether it be integral or fractional; for then, he observes, the root always lies hid in that number, and is some one of its divisors. This is exemplified in several instances, both of quadratic and cubic equations, and both for integral and fractional roots. And he here observes, that he knows not of any person who has yet given general rules for the solution of cubic equations; which shews that when he wrote this book, either Cardan's last book was not published, or else it had not yet come to his knowledge.

Chap. 17 contains, in a few words, directions for

bringing questions to equations, and for reducing these. He here observes, that some authors call the unknown number *res*, and others the *positio*; but that he calls it *radix*, or root, and marks it thus R: hence the term, root of an equation. But it was before called radix by Stifelius.

Chap. 21 & seq. treat of secondary roots, or a plurality of roots, denoted by A, B, C, &c, after Stifelius.

The 2d book contains the like operations in furd, or irrational numbers, and is a very complete work on this subject indeed. He treats first of simple or single furd, then of binomial furd, and lastly of trinomial furd. He gives here the same rule for extracting the root of a binomial and residual as Scheubelius, viz,  $\sqrt{a \pm b} =$

$$\sqrt{\frac{a + \sqrt{a^2 - b^2}}{2}} \pm \sqrt{\frac{a - \sqrt{a^2 - b^2}}{2}}. \text{ Individing by a}$$

binomial or residual, he proceeds as all others before him had done, namely, reducing the divisor to a simple quantity, by multiplying it by the same two terms with the sign of one of them changed, that is by the binomial if it be a residual, or by the same residual if it be a binomial; and multiplying the dividend by the same thing:

$$\text{thus } \frac{3}{\sqrt{5} - 2} = \frac{3}{\sqrt{5} - 2} \times \frac{\sqrt{5} + 2}{\sqrt{5} + 2} = \frac{3\sqrt{5} + 6}{5 - 4} = 3\sqrt{5} + 6.$$

And, in imitation of this method, in division by trinomial furd, he directs to reduce the trinomial divisor first to a binomial or residual, by multiplying it by the same trinomial with the sign of one term changed, and then to reduce this binomial or residual to a simple nomial as above; observing to multiply the dividend by the same quantities as the divisor. Thus, if the divisor be  $4 + \sqrt{2} - \sqrt{3}$ ; multiplying this by  $4 + \sqrt{2} + \sqrt{3}$ , the product is  $15 + 8\sqrt{2}$ ; then this binomial multiplied by the residual  $15 - 8\sqrt{2}$ , gives  $225 - 128$  or 97 for the simple divisor: and the dividend, whatever it is, must also be multiplied by the two  $4 + \sqrt{2} + \sqrt{3}$  and  $15 - 8\sqrt{2}$ . Or in general, if the divisor be  $a + \sqrt{b} - \sqrt{c}$ ; multiply it by  $a + \sqrt{b} + \sqrt{c}$ , which

$$\text{gives } (a + \sqrt{b})^2 - c = a^2 + b - c + 2a\sqrt{b}; \text{ then multiply this by } a^2 + b - c - 2a\sqrt{b},$$

and it gives  $(a^2 + b - c)^2 - 4a^2b$ , which will be rational, and will all collect into one single term. But Tartalea must have been in possession of some such rule as this, as one of the questions he proposed to Florido was of this nature, namely to find such a quantity as multiplied by a given trinomial furd, shall make it rational: and it appears, from what is done above,

that, the given trinomial being  $a + \sqrt{b} - \sqrt{c}$ , the answer will be  $a + \sqrt{b} + \sqrt{c} \times a^2 + b - c + 2a\sqrt{b}$ .

Chap. 24 shews the composition of the cube of a binomial or residual, and thence remarks on the root of the case or equation  $1 \text{ of } p \text{ } 3R$  equal to 10, which he seems to know something about, though he had not Cardan's rules.

Chap. 30, which is the last, treats of certain precepts relating to square and cubic numbers, with a table of such squares and cubes for all numbers to 140; also shewing how to compute them both, by adding always their differences.

He



He then concludes with remarking that there are many curious properties of these numbers, one of which is this, that the sum of any number of the cubes, taken from the beginning, always makes a square number, the root of which is the sum of the roots of the cubes; so that the series of squares so formed, have for their roots — 1, 3, 6, 10, 15, 21, &c. whose diff. are the natural nos 1, 2, 3, 4, 5, 6, &c. Namely,  $1^3 = 1^2$ ;  $1^3 + 2^3 = 3^2$ ;  $1^3 + 2^3 + 3^3 = 6^2$ , &c. Or in general,  $1^3 + 2^3 + 3^3 + \dots + n^3 = (1 + 2 + 3 + \dots + n)^2 = \frac{1}{2}n \cdot n + 1$ .

This work of Peletarius is a very ingenious and masterly composition, treating in an able manner of the several parts of the subject then known, excepting the cubic equations. But his real discoveries, or improvements, may be reduced to these three, viz.

1st. That the root of an equation, is one of the divisors of the absolute term.

2d. He taught how to reduce trinomials to simple terms, by multiplying them by compound factors.

3d. He taught curious precepts and properties concerning square and cube numbers, and the method of constructing a series of each by addition only, namely by adding successively their several orders of differences.

## RAMUS.

Peter Ramus wrote his arithmetic and algebra about the year 1560. His notation of the powers is thus,  $l$ ,  $q$ ,  $c$ ,  $bq$ , being the initials of *latus*, *quadratus*, *cubus*, *biquadratus*. He treats only of simple and quadratic equations. And the only thing remarkable in his work, is the first article, on the names and invention of Algebra, which we have noticed at the beginning of this history.

## BOMBELLI.

Raphael Bombelli's Algebra was published at Bologna in the year 1579, in the Italian language. It seems however it was written some time before, as the dedication is dated 1572. In a short, but neat, introduction, he first adverts, in a few words, to the great excellence and usefulness of arithmetic and algebra. He then laments that it had hitherto been treated in so imperfect and irregular a way; and declares it is his intention to remedy all defects, and to make the science and practice of it as easy and perfect as may be. And for this purpose he first resolved to procure and study all the former authors. He then mentions several of these, with a short history or character of them; as Mahomet the son of Moses, an Arabian; Leonard Pisano; Lucas de Burgo, the first printed author in Europe; Oroncius; Seribellus; Boglione Francesi; Stifelius in Germany; a certain Spaniard, doubtless meaning Nunez or Nonius; and lastly Cardan, Ferrari, and Tartalea; with some others since, whose names he omits. He then adds a curious paragraph concerning Diophantus: he says that some years since there had been found, in the Vatican library, a Greek work on this art, composed by a certain Diophantus, of Alexandria, a Greek author, who lived in the time of Antoninus Pius; which work having been shewn to him by Mr. Antonio Maria Pazzi Reggiano, public lecturer on mathematics at Rome; and finding it to be a good work, these two formed the re-

solution of giving it to the world, and he says that they had already translated five books, of the six which were then extant, being as yet hindered by other avocations from completing the work. He then adds the following strange circumstance, viz. that they had found that in the said work the Indian authors are often cited; by which they learned that this science was known among the Indians before the Arabians had it: a paragraph the more remarkable as I have never understood that any other person could ever find, in Diophantus, any reference to Indian writers: and I have examined his work with some attention, for that purpose.

Bombelli's work is divided into three books. In the first, are laid down the definitions and operations of powers and roots, with various sorts of radicals, simple and compound, binomial, residual, &c; mostly after the rules and manner of former writers, excepting in some few instances, which I shall here take notice of. And first of his rule for the cube root of binomials or residuals, which for the sake of brevity, may be expressed in modern notation as follows: let  $\sqrt[3]{b+a}$  be the binomial, the term  $\sqrt[3]{b}$  being greater than  $a$ ; then the rule for the cube root of  $\sqrt[3]{b+a}$  comes

to this,  $P - Q + \sqrt{P - Q^2} + \sqrt[3]{b - a^2}$  where  $P = \sqrt[3]{\frac{a^2}{64} + \frac{b - a^2}{64} + \frac{a}{8}}$ , and  $Q = \sqrt[3]{\frac{a^2}{64} + \frac{b - a^2}{64} - \frac{a}{8}}$ .

Which is a rule that can be of little or no use; for, in the first place,  $\sqrt{P - Q^2} + \sqrt[3]{b - a^2}$  is the same as

$\sqrt{P + Q^2}$ ; and  $P$  or  $\sqrt[3]{\frac{a^2}{64} + \frac{b - a^2}{64} + \frac{a}{8}}$  is =

$$\sqrt[3]{\sqrt[3]{\frac{b}{64} + \frac{a}{8}}} = \sqrt[3]{\frac{\sqrt[3]{b+a}}{8}} = \frac{1}{2} \sqrt[3]{\sqrt[3]{b+a}};$$

therefore the whole  $P - Q + \sqrt{P - Q^2} + \sqrt[3]{b - a^2}$  reduces to  $P - Q + P + Q = 2P = 2 \times \frac{1}{2} \sqrt[3]{\sqrt[3]{b+a}}$

=  $\sqrt[3]{\sqrt[3]{b+a}}$ , the original quantity first proposed. The next thing remarkable in this 1st book, is his method for the square roots of negative quantities, and his rule for the cube roots of such imaginary binomials as arise from the irreducible case in cubic equations. His words, translated, are these: "I have found another sort of cubic root, very different from the former, which arises from the case of the cube equal to the first power and a number, when the cube of the  $\frac{1}{3}$ d part of the (coef. of the) 1st power, is greater than the square of half the absolute number, which sort of square root hath in its algorism, names and operations different from the others; for in that case, the excess cannot be called either plus or minus; I therefore call it *plus of minus* when it is to be added, and *minus of minus* when it is to be subtracted." He then gives a set of rules for the signs when such roots are multiplied, and illustrates them by a great many examples. His rule for the cube roots of such binomials, viz. such

as  $a + \sqrt{-b}$ , is this: First find  $\sqrt[3]{a^2 + b}$ ; then, by trials search out a number  $c$ , and a sq. root  $\sqrt{d}$ , such, that the sum of their squares  $c^2 + d$  may be =  $\sqrt[3]{a^2 + b}$  and



and also  $c^3 - 3cd = a$ ; then shall  $c + \sqrt{-d}$  be =  $\sqrt[3]{a + \sqrt{-b}}$  sought. Thus, to extract the cube root of  $2 + \sqrt{-121}$ : here  $\sqrt[3]{a^2 + b} = \sqrt[3]{125} = 5$ ; then taking  $c = 2$ , and  $d = 1$ , it is  $c^2 + d = 5 = \sqrt[3]{a^2 + b}$ , and  $c^3 - 3cd = 8 - 6 = 2 = a$ , as it ought; and therefore  $2 + \sqrt{-1}$  is = the cube root of  $2 + \sqrt{-121}$ , as required.

The notation in this book, is the initial *R* for root, with *q* or *c* &c after it, for quadrate or cubic, &c root. Also *p* for *plus*, and *m* for *minus*.

In the 2d book, Bombelli treats of the algorism with unknown quantities, and the resolution of equations. He first gives the definitions and characters of the unknown quantity and its powers, in which he deviates from the former authors, but professes to imitate Diophantus. He calls the unknown quantity *tanto*, and marks it thus

Its square or 2d power *potenza*, <sup>2</sup>,  
Its cube - - - *cubo*, <sup>3</sup>,

and the higher names are compounded of these, and marked <sup>4</sup>, <sup>5</sup>, <sup>6</sup>, <sup>7</sup>, &c, so that he denotes all the powers by their exponents set over the common character  $\sqrt{\phantom{x}}$ . And all these powers he calls by the general name *dignita*, dignity. He then performs all the algorism of these powers, by means of their exponents, as we do at present, viz, adding them in multiplication, subtracting in division, multiplying them by the index in involution, and dividing by the same in evolution.

In equations he goes regularly through all the cases, and varieties of the signs and terms; first all the simple or single powers, and then all the compound cases; demonstrating the rules geometrically, and illustrating them by many examples.

In compound quadratics, he gives two rules: the first is by freeing the *potenza* or square from its coefficient by division, and then completing the square, &c, in the usual way: and the 2d rule, when the first term has its coefficient, may be thus expressed; if  $ax^2 + bx = c$ ,

then  $x = \frac{\sqrt{ac + \frac{1}{4}b^2} - \frac{1}{2}b}{a}$ . He takes only the positive

root or roots; and in the case  $ax^2 + c = bx$ , which has two, he observes that the nature of the problem must shew which of the two is the proper one.

In the cubic equations, he gives the rules and transformations, &c, after the manner of Cardan; remarking that some of the cases have only one root, but others two or three, of which some are true, and others false or negative. And in one place he says that by means of the case  $x^3 = bx + c$  he *trifects* or *divides an angle into three equal parts*.

When he arrives at biquadratic equations, and particularly to this case  $x^4 + ax = b$ , he says, "Since I have seen Diophantus's work, I have always been of opinion that his chief intention was to come to this equation, because I observe he labours at finding always square numbers, and such, that adding some number to them, may make squares; and I believe that the six books, which are lost, may treat of this equation, &c." — "But Lewis Ferrari," he adds, "of this city, also laboured in this way, and found out a rule for such cases, which was a very fine invention, and therefore I

shall here treat of it the best I can." This he accordingly does, in all the cases of biquadratics, both with respect to the number of terms in the equation, and the signs of the terms, except I think this most general case only  $rx - qx^2 + px^3 - x^4 = s$ ; fully applying Ferrari's method in all cases. Which concludes the 2d book.

The 3d book consists only of the resolution of near 300 practical questions, as exercises in all the rules and equations, some of which are taken from Diophantus and other authors.

Upon the whole it appears that this is a plain, explicit, and very orderly treatise on algebra, in which are very well explained the rules and methods of former writers. But Bombelli does not produce much of improvement or invention of his own, except his notation, which varies from others, and is by means of one general character, with the numeral indices of Stifelius. He also first remarks that angles are trifected by a cubic equation. But I know not how to account for his assertion, that Diophantus often cites the Indian authors; which I think must be a mistake in Bombelli.

#### CLAVIUS.

Christopher Clavius wrote his Algebra about the year 1580, though it was not published till 1608, at Orleans. He mostly follows Stifelius and Scheubelius in his notation and method, &c, having scarcely any variations from them; nor does he treat of cubic equations. He mentions the names given to the art, and the opinions about its origin, in which he inclines to ascribe it to Diophantus, from what Diophantus says in his preface to Dyonisius.

#### STEVINUS.

The Arithmetic of Simon Stevin of Bruges, was published in 1585, and his Algebra a little afterwards. They were also printed in an edition of his works at Leyden in 1634, with some notes and additions of Albert Girard, who it seems died the year before, this edition being published for the benefit of Girard's widow and children. The Algebra is an ingenious and original work. He denotes the *res*, or unknown quantity, in a way of his own, namely by a small circle  $\circ$ , within which he places the numeral exponent of the power, as  $\textcircled{0}$ ,  $\textcircled{1}$ ,  $\textcircled{2}$ ,  $\textcircled{3}$ , &c, which are the 0, 1, 2, 3, &c power of the quantity  $\circ$ ; where  $\textcircled{0}$ , or the 0 power, is the beginning of quantity, or arithmetical unit. He also extends this notation to roots or fractional exponents, and even to radical ones.

Thus  $\textcircled{\frac{1}{2}}$ ,  $\textcircled{\frac{1}{3}}$ ,  $\textcircled{\frac{1}{4}}$ , &c, are the sq. root, cube root, 4th root, &c;

and  $\textcircled{4}$  is the cube root of the square;

and  $\textcircled{\frac{2}{3}}$  is the sq. root of the cube. And so of others.

The first three powers,  $\textcircled{1}$ ,  $\textcircled{2}$ ,  $\textcircled{3}$ , he also calls *coste* (side), *quarre* (square), *cube* (cube); and the first of them,  $\textcircled{1}$ , the prime quantity, which he observes is also *metaphorically* called the *racine* or root, (the mark of which is also  $\sqrt{\phantom{x}}$ ), because it represents the root or origin from whence all other quantities spring or arise, called the *potences* or powers of it. He condemns the terms *furfolds*, and numbers absurd, irrational, irregular, inexplicable, or furd, and shews that all numbers are denoted the same way, and are all equally proper expressions



pressions of some length or magnitude, or some power of the same root. He also rejects all the compound expressions of square-squared, cube-squared, cube-cubed, &c, and shews that it is best to name them all by their exponents, as the 1st, 2d, 3d, 4th, 5th, 6th, &c power or quantity in the series. And on his extension of the new notation he justly observes that what was before obscure, laborious, and tiresome, will by these marks be clear, easy, and pleasant. He also makes the notation of algebraic quantities more general in their coefficients, including in them not only integers, as  $3\textcircled{1}$ , but also fractions and radicals, as  $\frac{3}{4}\textcircled{2}$ , and  $\sqrt{2}\textcircled{3}$ , &c. He has various other peculiarities in his notations; all shewing an original and inventive mind. A quantity of several terms, he calls a multinomial, and also binomial, trinomial, &c, according to the number of the terms. He uses the signs + and -, and sometimes : for equality; also  $\times$  for division of fractions, or to multiply crosswise thus,  $\frac{5}{7} \times \frac{2}{3} : \frac{10}{21}$ .

He teaches the generation of powers by means of the annexed table of numbers, which are the coefficients of all the terms except the first and last. And he makes use of the same numbers also for extracting all roots whatever: both which things had first been done by Stifelius. In extracting the roots of non-quadrate or non-cubic numbers, he has the same approximations as at present, viz, either to continue the extraction indefinitely in decimals, by adding periods of ciphers, or by making a fraction of the remainder

in this manner, viz,  $\sqrt{N} = n + \frac{N - n^2}{2n + 1}$  nearly, and

$\sqrt[3]{N} = n + \frac{N - n^3}{3n^2 + 3n + 1}$  nearly; where  $n$  is the

nearest exact root of  $N$ ; which is Peletarius's rule, and which differs from Tartalea's rule, as this wants the 1 in the denominator. And in like manner he goes on to the roots of higher powers.

He then treats of equations, and their inventors, which according to him are thus:

Mahomet, son of Moses, an Arabian, invented these

And some unknown author, the derivatives of this.

Some unknown author invented these

But afterwards he mentions Ferreus, Tartalea, Cardan, &c, as being also concerned in the invention of them.

Lewis Ferrari invented

He says also that Diophantus once resolves the case  $\textcircled{2}$  egale à  $\textcircled{1} \textcircled{0}$ . In his reduction of equations, which is full and masterly, he always puts the highest power on one side alone, equal to all the other terms, set in their order, on the other side, whether they be + or -. And he demonstrates all the rules both arithmetically and geometrically. In cubics, he gives up the irreducible case, as hopeless: but says that Bombelli resolves it by *plus of minus*, and *minus of minus*; thus, if  $1\textcircled{3} = 30\textcircled{1} + 36$ , then  $1\textcircled{1} = \sqrt[3]{18 + \text{of} - 26} + \sqrt[3]{18 - \text{of} - 26}$ , that is,  $1\textcircled{1} = \sqrt[3]{18 + 26\sqrt{-1}} + \sqrt[3]{18 - 26\sqrt{-1}}$ . He resolves biquadratics by

means of cubics and quadratics. In quadratics, he takes both the two roots, but looks for no more than two in cubics or biquadratics. He gives also a general method of approaching indefinitely near, in decimals, to the root of any equation whatever: but it is very laborious, being little more than trying all numbers, one after another, finding thus the 1st figure, then the 2d, then the 3d, &c, among these ten characters 0, 1, 2, 3, 4, 5, 6, 7, 8, 9. And finally he applies the rules in the resolution of a great many practical questions.

Although a general air of originality and improvement runs through the whole of Stevinus's work, yet his more remarkable or peculiar inventions, may be reduced to these few following: viz,

1st. He invented not only a new character for the unknown quantity, but greatly improved the notation of powers, by numeral indices, first given by Stifelius as to integral exponents; which Stevinus extended to fractional and all other sorts of exponents, thereby denoting all sorts of roots the same way as powers, by numeral exponents. A circumstance hitherto thought to be of much later invention.

2d. He improved and extended the use and notation of coefficients, including in them fractions and radicals, and all sorts of numbers in general.

3d. A quantity of several terms, he called generally a multinomial; and he denoted all nomials whatever by particular names expressing the number of their terms, binomial, trinomial, quadrinomial, &c.

4th. A numeral resolution of all equations whatever by one general method.

Besides which, he hints at some unknown author as the first inventor of the rules for cubic equations; by whom may probably be intended the author of the Arabic manuscript treatise on cubic equations, given to the library at Leyden by the celebrated Warner.

#### VIETA.

Most of Vieta's algebraical works were written about or a little before, the year 1600, but some of them were not published till after his death, which happened in the year 1603. And his whole mathematical works were collected together by Francis Schooten, and elegantly printed in a folio volume in 1646. Of these, the algebraical parts are as follow:

1. Isagoge in Artem Analyticam.
2. Ad Logisticen Speciosam Notæ priores.
3. Zeticorum libri quinque.
4. De Aequationum Recognitione, & Emendatione.
5. De Numerosa Potestatum ad Exegesein Resolutione.

Of all these I shall give a very minute account, especially in such parts as contain any discoveries, as we here meet with more improvements and inventions in the nature of equations, than in almost any former author. And first of the *Isagoge* or Introduction to the Analytic Art. In this short introduction Vieta lays down certain præcognita in this art, as definitions, axioms, notations, common precepts or operations of addition, subtraction, multiplication, and division, with rules for questions, &c. From which we find, 1st. That the names of his powers are *latus*, *quadratum*, *cubus*, *quadrato-quadratum*, *quadrato-cubus*, *cubocubus*, &c; in which he follows the method of Diophantus, and not that derived from the Arabians.



2d. That he calls powers pure or adfectèd, and first here uses the terms coefficient, affirmative, negative, specious logisticks or calculations, homogeneous comparisonis, or the absolute known term of an equation, homogeneous adfectionis, or the 2d or other term which makes the equation adfectèd, &c. 3d. That he uses the capital letters to denote the known as well as unknown quantities, to render his rules and calculations general, namely, the vowels A, E, I, O, U, Y for the unknown quantities, and the consonants B, C, D, &c, for the known ones. 4th. That he uses the sign + between two terms for addition; - for subtraction, placing the greater before the less; and when it is not known which term is the greater, he places = between them for the difference, as we now use  $\infty$ ; thus  $A = B$  is the same as  $A \infty B$ ; that he expresses division by placing the terms like a fraction, as at present; though he was not first in this. But that he uses no characters for multiplication or equality, but writes the words themselves, as well as the names of all the powers, as he uses no exponents, which causes much trouble and prolixity in the progress of his work; and the numeral coefficients set after the literal quantities, have a disagreeable effect.

II. *Ad Logisticen Speciosam, Notæ Priores.* These consist of various theorems concerning sums, differences, products, powers, proportionals, &c, with the genesis of powers from binomial and residual roots, and certain properties of rational right-angled triangles.

III. *Zeticorum libri quinque.* The zeticus or questions in these 5 books are chiefly from Diophantus, but resolved more generally by literal arithmetic. And in these questions are also investigated rules for the resolution of quadratic and cubic equations. In these also Vieta first uses a line drawn over compound quantities, as a vinculum.

IV. *De Aequationum Recognitione, & Emendatione.* These two books, which contain Vieta's chief improvements in Algebra, were not published till the year 1615, by Alexander Anderson, a learned and ingenious Scotchman, with various corrections and additions. The 1st of these two books consists of 20 chapters. In the first six chapters, rules are drawn from the zeticus for the resolution of quadratic and cubic equations. These rules are by means of certain quantities in continued proportion, but in the resolution they come to the same thing as Cardan's rules. In the cubics, Vieta sometimes changes the negative roots into affirmative, as Cardan had done, but he finds only the affirmative roots. And he here refers the irreducible case to angular sections for a solution, a method which had been mentioned by Bombelli.

*Chap. 7* treats of the general method of transforming equations, which is done either by changing the root in various ways, namely by substituting another instead of it which is either increased or diminished, or multiplied or divided by some known number, or raised or depressed in some known proportion; or by retaining the same root, and equally multiplying all the terms. Which sorts of transformation, it is evident, are intended to make the equation become simpler, or more convenient for solution. And all or most of these reductions and transformations were also practised by Cardan.

*Chap. 8* shews what purposes are answered by the

foregoing transformations; such as taking away some of the terms out of an equation, and particularly the 2d term, which is done by increasing or diminishing the root by the coefficient of the 2d term divided by the index of the first: by which means also the affected quadratic is reduced to a simple one. And various other effects are produced.

*Chap. 9* shews how to deduce compound quadratic equations from pure ones, which is done by increasing or diminishing the root by a given quantity, being one application of the foregoing reductions.

*Chap. 10*, the reduction of cubic equations affected with the 1st power, to such as are affected with the 2d power; by the same means.

In *chap. 11*, by the same means also, the 2d term is restored to such cubic equations as want it.

In *chap. 12*, quadratic and cubic equations are raised to higher degrees by substituting for the root, the square or cube of another root divided by a given quantity.

In *chap. 13* affected biquadratic equations are deduced from affected quadratics in this manner, when expressed in the modern notation:

If  $A^2 + BA = Z$ , then shall  $A^4 + B^2 + 2BZ \cdot A = Z^2 + B^2Z$ .

For since  $A^2 + BA = Z$ , therefore  $A^2 = Z - BA$ , and its square is

$A^4 = Z^2 - 2BAZ + B^2A^2$ : but  $B^2A^2 = B^2Z - B^3A$ , therefore

$A^4 = Z^2 - 2BAZ + B^2Z - B^3A$ , or  $A^4 + B^2 + 2BZ \cdot A = Z^2 + B^2Z$ .

And in like manner for the biquadratic affected with its other terms. And in a similar manner also, in *chap. 14*, affected cubic equations are deduced from the affected quadratics.

In *chap. 15* it is shewn that the quadratic  $BA - A^2 = Z$  has two values of the root A, or has ambiguous roots, as he calls them; and also that the cubics, biquadratics, &c, which are raised or deduced from that quadratic, have also double roots.

Having, in the foregoing chapters, shewn how the coefficients of equations of the 3d and 4th degree are formed from those of the 2d degree, of the same root; and that certain quadratics, and others raised from them, have double roots; then in the 16th *chap.* Vieta shews what relation those two roots bear to the coefficients of the two lowest terms of an equation consisting of only three terms. Thus,

If  $BA - A^2 = Z$  } then  $B = \frac{A^2 - E^2}{A - E} = A + E$ ,  
and  $BE - E^2 = Z$  } and  $Z = \frac{A^2E - AE^2}{A - E} = AE$ .

If  $BA - A^3 = Z$  } then  $B = \frac{A^3 - E^3}{A - E} = A^2 + AE + E^2$ ,  
and  $BE - E^3 = Z$ , } and  $Z = \frac{A^3E - AE^3}{A - E} = A^2E + AE^2$ .

If  $A^m + BA^n = Z$ , } then  $B = \frac{E^m - A^m}{E^n + A^n}$ , &  $Z = \frac{A^mE^n + A^nE^m}{E^n + A^n}$ ,  
and  $E^m - BE^n = Z$ , }

And so on for the same terms with their signs variously changed.



*Chap. 17* contains several theorems concerning quantities in continued geometrical progression. Which are preparatory to what follows, concerning the double roots of equations, the nature of which he expounds by means of such properties of proportional quantities.

*Chap. 18, Æquationum ancipitum constitutiva*; treating of the nature of the double roots of equations. Thus, if  $a, b, c, d$ , &c, be quantities in continual progression; then, 1st, of equations affected with the first power, If  $BA - A^2 = Z$ ; then  $B = a + b, Z = ab$ , &  $A = a$  or  $b$ . If  $BA - A^3 = Z$ ; then  $B = a^2 + b^2 + c^2, Z = a(b^2 + c^2)$ , &  $A = a$  or  $c$ .

And in general, if  $BA - A^{n+1} = Z$ ; then  $B = a^n + b^n + c^n + \dots + k^n, Z = a(b^n + c^n + d^n + \dots + k^n)$ , and  $A = a$  or  $k$  the first or last term. Where the number of terms  $a^n, b^n$ , &c, in  $B$  is  $n + 1$ , and the number of terms in  $Z$  is  $n$ .

2d, For equations containing only the highest two powers.

If  $BA^2 - A^3 = Z$ ; then  $B = a + b + c, Z = a(b + c)^2$ , or  $= c(a + b)^2$ , and  $A = a + b$ , or  $b + c$ .

If  $BA^3 - A^4 = Z$ ; then  $B = a + b + c + d, Z = a(b + c + d)^3$  or  $= d(a + b + c)^3$ , and  $A = a + b + c$  or  $b + c + d$ .

And, in general, if  $BA^n - A^{n+1} = Z$ ;

then  $B = a + b + c + \dots + i + k$ ,

$Z = a(b + c + \dots + k)$  or  $= k(a + b + c + \dots + i)$ ,

and  $A = a + b + c + \dots + i$  or  $= b + c + d + \dots + k$ , the sum of all except the last, or sum of all except the first; where the number of terms in  $B$  is  $n + 1$ , and the number of terms in  $Z$  is  $n$ .

3d. Of equations affected by the intermediate powers.

If  $BA^2 - A^4 = Z$ ; then  $B = a^2 + b^2, Z = a^2 b^2$ , &  $A^2 = a^2$  or  $b^2$ .

If  $BA^3 - A^6 = Z$ ; then  $B = a^3 + b^3, Z = a^3 b^3$ , &  $A^3 = a^3$  or  $b^3$ .

If  $BA^4 - A^6 = Z$ ; then  $B = a^2 + b^2 + c^2, Z = a(b^2 + c^2)$ , and  $A^2 = a + b$  or  $b + c$ .

4th. Of the remaining cases.

If  $BA^2 - A^5 = Z$ ;

then  $B = (a + b)^3 + (b + c)^3 + a(b + c)^2$  or  $+ c(a + b)^2$ ,

and  $Z = \overline{B - (a + b)^3} \cdot (a + b)^2$

or  $= \overline{(b + c)^3 + c(a + b)^2} \cdot (a + b)^2$ ,

and  $A = a + b$  or  $b + c$ .

If  $BA^3 - A^5 = Z$ ;

then  $B = (a + b + c)^2 + (b + c + d)^2 - c(a + b + c)$

or  $= b(b + c + d)$ ,

and  $Z = \overline{B - (b + c + d)^2} \cdot (b + c + d)^2$

or  $= \overline{B - (a + b + c)^2} \cdot (a + b + c)^2$ ;

and  $A = a + b + c$  or  $b + c + d$ .

*Chap. 19. Æqualitatum contradicentium constitutiva.*

Of the relation of equations of like terms, but the sign of one term different; containing these 5 theorems, viz,

1. If  $A^2 + BA = Z$ , } then  $B = b - a, Z = ab$ ;  
and  $E^2 - BE = Z$ ; } and  $A = a, E = b$ .

2. If  $A^4 + BA = Z$ , } then  $B = (b + d)^3 - (a + c)^3$ ,  
and  $E^4 - BE = Z$ ; }  $Z = a(d^3 - b^3)$  or  $= d(c^3 - a^3)$ ;  
and  $A = a, E = d$ .

3. If  $A^5 + BA = Z$ , } then  $B = b^5 + d^5 + f^5 - a^5 - c^5 - e^5$ ,  
and  $E^6 - BE = Z$ ; }  $Z = a(b^5 + d^5 + f^5 - c^5 - e^5)$   
or  $= f(a^5 + c^5 + e^5 - b^5 - d^5)$ ;  
and  $A = a, E = f$ .

4. If  $A^4 + BA^3 = Z$ , } then  $B = b + d - a - c, Z =$   
and  $E^4 - BE^3 = Z$ ; }  $a(d - b)^3$  or  $= d(c - a)^3$ ;  
and  $A = c - a, E = d - b$ .

5. If  $A^6 + BA^5 = Z$ , } then  $B = b + d + f - a - c - e$ ,  
and  $E^6 - BE^5 = Z$ ; }  $Z = a(b + d + f - c - e)^5$   
or  $= f(a + c + e - b - d)^5$ ;  
and  $A = a + c + e - b - d$ ,  
 $E = b + d + f - c - e$ .

*Chap. 20. Æqualitatum inversarum constitutiva.* Containing these six theorems, viz,

1. If  $BA - A^3 = Z$ , } then  $B = c^2 - a^2, Z = a(c^2 - b^2)$   
and  $E^3 - BE = Z$ ; } or  $= c(b^2 - a^2)$ ;  
and  $A = a, E = c$ .

2. If  $BA - A^5 = Z$ , } then  $B = a^4 + c^4 + e^4 - b^4 - d^4$ ,  
and  $E^5 - BE = Z$ ; }  $Z = a(c^4 + e^4 - b^4 - d^4)$   
or  $= e(b^4 + d^4 - a^4 - c^4)$ ;  
and  $A = a, E = e$ .

3. If  $BA^2 + A^3 = Z$ , } then  $B = c - a, Z = a(c - b)^2$   
and  $BE^2 - E^3 = Z$ ; } or  $= c(b - a)^2$ ;  
and  $A = b - a, E = c - b$ .

4. If  $BA^4 + A^5 = Z$ , } then  $B = a + c + e - b - d$ ,  
and  $BE^4 - E^5 = Z$ ; }  $Z = a(c + e - b - d)^4$   
or  $= e(b + d - a - c)^4$ ;  
and  $A = b + d - a - c$ ,  
 $E = c + e - b - d$ .

5. If  $BA^3 + A^5 = Z$ , } then  $B = (a + f) \cdot (d - a)$ ,  
and  $BE^3 - E^5 = Z$ ; }  $Z = (d - a)^3 B + (d - a)^5$   
or  $= (e - b)^3 B - (e - b)^5$ ;  
and  $A = d - a, E = e - b$ .

6. If  $BA^2 + A^5 = Z$ , } then  $B = (a + f) \cdot (c - a)^2$ ,  
and  $BE^2 - E^5 = Z$ ; }  $Z = \overline{B + (c - a)^3} \cdot (c - a)^2$   
or  $= \overline{B - (d - b)^3} \cdot (d - b)^2$   
and  $A = c - a, E = d - b$ .

*Chap. 21. Alia rursus æqualitatum inversarum constitutiva.* In these two theorems:

1. If  $B^2 A - A^3 = Z$ , } then  $B = a^2 + b^2 + c^2$ ,  
and  $E^3 - B^2 A = Z$ ; }  $Z = (a + c)b^2$  or  $= a(b^2 + c^2)$   
or  $= c(a^2 + b^2)$ ;  
and  $A = a$  or  $c, E = a + c$ .

2. If  $BA^2 - A^3 = Z$ , } then  $B = a + b + c$ ,  
and  $BE^2 + E^3 = Z$ ; }  $Z = (a + b + c)b^2$  or  $=$   
 $a(b + c)^2$  or  $= c(a + b)^2$ ;  
and  $A = a + b$  or  $b + c, E = b$ .

Next follows the 2d of the pieces published by Alexander Anderson, namely,

*De Emendatione Æquationum*, in 14 chapters.

*Chap. 1.* Of preparing equations for their resolution in numbers, by taking away the 2d term; by which affected quadratics are reduced to pure ones, and cubic equations affected with the 2d term are reduced to such as are affected with the 3d only. Several examples of both sorts of equations are given. He here too remarks upon the method of taking away any other term out of an equation, when the highest power is combined with that other term only; and this Vieta effects by means of the coefficients, or, as he calls them, the unciæ of the power of a binomial. All which was also performed by Cardan for the same purpose.

*Chap. 2. De transmutatione Περὶ τὸν ἑκαστον, quæ remedium est adversus vitium negationis.* Concerning the transformations by changing the given root  $A$  for another root  $E$ , which is equal to the homogeneous compara-



comparationis divided by the first root  $A$ ; by which means negative terms are changed to affirmative, and radicals are taken out of the equation when they are contained in the homogeneous comparisonis.

*Chap. 3, De Anastrophe*, shewing the relation between the roots of correlate equations; from whence, having given the root of the one equation, that of the other becomes known; and it consists of these following 8 theorems, mostly deduced from the last 4 chapters of the foregoing *recognitio æquationum*.

1. If  $BA - A^3 = Z$ ,  
and  $E^3 - BE = Z$ ; } then  $EA - A^2 = E^2 - B$ .
2. If  $BA^2 - A^3 = Z$ ,  
and  $BE^2 + E^3 = Z$ ; } then  $(E + B)A - A^2 = BD + D^2$ .
3. If  $BA - A^5 = Z$ ,  
and  $E^3 - BE = Z$ ; } then  $E^3A - E^2A^2 + EA^3 - A^4 = E^4 - B$ .
4. If  $BA^4 - A^5 = Z$ ,  
and  $BE^4 + E^5 = Z$ ; } then  $(BE^2 + E^3)A - (BE + E^2)A^2 + (B + E)A^3 - A^4 = BE^3 + E^4$ .
5. If  $A^3 - BA = Z$ ,  
and  $BE - E^3 = Z$ ; } then  $A^2 - EA = B - E^2$ .
6. If  $BA^2 + A^3 = Z$ ,  
and  $BE^2 - E^3 = Z$ ; } then  $A^2 + (B - E)A = BE - E^2$ .
7. If  $BA - A^3 = Z$ ,  
and  $BE - E^3 = Z$ ; } then  $A^2 + EA = B - E^2$ .
8. If  $BA^2 - A^3 = Z$ ,  
and  $BE^2 - E^3 = Z$ ; } then  $A^2 + (E - B)A = BE - E^2$ .

*Chap. 4, De Isomæria, adversus vitium fractionis.* To take away fractions out of an equation. Thus,

if  $A^3 + \frac{B}{D}A = Z$ . Put  $A = \frac{E}{D}$ ; then  $E^3 + BDE = ZD^3$ .

*Chap. 5, De Symmetrica Climacismo adversus vitium asymmetriæ.* To take away radicals or surds out of equations, by squaring &c the other side of the equation.

*Chap. 6.* To reduce biquadratic equations by means of cubics and quadratics, by methods which are small variations from those of Ferrari and Cardan.

*Chap. 7.* The resolution of cubic equations by rules which are the same with Cardan's.

*Chap. 8. De Canonica æquationum transmutatione, ut coefficientes subgraduales sint quæ præscribuntur.* To transmute the equation so that the coefficient of the lower term, or power, may be any given number, he changes the root in the given proportion, thus: Let  $A$  be the root of the equation given,  $E$  that of the transmuted equation,  $B$  the given coefficient, and  $X$

the required one; then take  $A = \frac{BE}{X}$ , which substitute in the given equation, and it is done.—He commonly changes it so, that  $X$  may be 1; which he does, that the numeral root of the equation may be the easier found; and this he here performs by trials, by taking the nearest root of the highest power alone; and if that does not turn out to be the root of the whole equation, he concludes that it has no rational root.

*Chap. 9.* To reduce certain peculiar forms of cubics to quadratics, or to simpler forms, much the same as Cardan had done. Thus,

1. If  $A^3 - 2B^2A = B^3$ ; then is  $A^2 - BA = B^2$ .
2. If  $2B^2A - A^3 = B^3$ ; then is  $A^2 + BA = B^2$ .
3. If  $A^3 - 3B^2A = 2B^3$ ; then is  $A = 2B$ .
4. If  $3B^2A - A^3 = 2B^3$ ; then is  $A = B$ .
5. If  $A^3 - BA^2 + DA = BD$ ; then is  $A = B$ .
6. If  $A^3 + BA^2 - D^2A = BD^2$ ; then is  $A = D$ .
7. If  $BA^2 + D^2A - A^3 = BD^2$ ; then is  $A = B$  or  $A = D$ .
8. If  $D^2A + BDA - A^3 = B^3$ ; then is  $(D + B)A - A^2 = B^2$ .
9. If  $A^3 - DA^2 + BDA = B^3$ ; then is  $(D - B)A - A^2 = B^2$ .
10. If  $A^3 - 3BA = B\sqrt{2}B$ ; then is  $A = \sqrt{\frac{3}{2}}B - \sqrt{\frac{1}{2}}B$ .
11. If  $3BA - A^3 = B\sqrt{2}B$ ; then is  $A = \sqrt{\frac{3}{2}}B - \sqrt{\frac{1}{2}}B$ .

*Chap. 10. Similium reductionum continuatio.* Being some more similar theorems, when the equation is affected with all the powers of the unknown quantity  $A$ .

*Chap. 11, 12, 13* relate also to certain peculiar forms of equations, in which the root is one of the terms of a certain series of continued proportionals.

*Chap. 14*, which is the last in this tract, contains, in four theorems, the general relation between the roots of an equation and the coefficients of its terms, when all its roots are positive. Namely,

1. If  $\overline{B + D} \cdot A - A^2 = BD$ ; then is  $A = B$  or  $A = D$ .
2. If  $\overline{A^3 - B - D - G} \cdot \overline{A^2 + BD + BG + DG} \cdot A = BDG$ ; then is  $A = B$  or  $A = D$  or  $A = G$ .
3. If  $\overline{BDG + BDH + BGH + DGH} \cdot \overline{A - BD - BG - BH - DG - DH - GH} \cdot \overline{A^2 + B + D + G + H} \cdot \overline{A^3 - A^4} = BDGH$ ; then is  $A = B$  or  $A = D$  or  $A = G$  or  $A = H$ .
4. If  $\overline{A^3 - B - D - G - H - K} \cdot \overline{A^4 + BD + BG + BH + BK + DG + DH + DK + GH + GK + HK} \cdot \overline{A^3 - BDG - BDH - BDK - BGH - BGK - BHK - DGH - DGK - DHK - GHK} \cdot \overline{A^2 + BDGH + BDGK + BDHK + BGHK + DGHK} \cdot A = BDGHK$ ; then is  $A = B$  or  $A = D$  or  $A = G$  or  $A = H$  or  $A = K$ .

And from these last 4 theorems it appears that Vieta was acquainted with the composition of these equations, that is, when all their roots are positive, for he never adverts to negative roots; and from other parts of the work it appears that he was not aware that the same properties will obtain in all sorts of roots whatever. But it is not certain in what manner he obtained these theorems, as he has not given any account of the investigations, though that was usually his way on other occasions; but he here contents himself with barely announcing the theorems as above, and for this strange reason, that he might at length bring his work to a conclusion.

To this piece is added, by Alexander Anderson, an Appendix, containing the construction of the cubic equations by the trisection of an angle, and a demonstration of the property referred to by Vieta for this purpose.

*De Numerosa Potestatum Purarum Resolutione.* Vieta here gives some examples of extracting the roots of pure powers, in the way that had been long before practised, by pointing the number into periods of figures according to the index of the root to be extracted, and then proceeding from one period to another, in the usual way.



*De Numerosa Potestatum adfectarum Resolutione.* And here, in close imitation of the above method for the roots of pure powers, Vieta extracts those of adaffected ones; or finding the roots of affected equations, placing always the homogeneous comparisonis, or absolute term, on one side, and all the terms affected with the unknown quantity, and their proper signs, on the other side. The method is very laborious, and is but little more than what was before done by Stevinus on this subject, depending not a little upon trials. The examples he uses are such as have either one or two roots, and indeed such as are affected commonly with only two powers of the unknown quantity, and which therefore admit only of those two varieties as to the number of roots, namely according as the higher of the two powers is affirmative or negative, the homogeneous comparisonis, on the other side of the equation, being always affirmative; and he remarks this general rule, if the higher power be negative, the equation has two roots; otherwise, only one; that is, affirmative roots; for as to negative and imaginary ones, Vieta knew nothing about them, or at least he takes no notice of them. By the foregoing extraction, Vieta finds both the greater and less root of the two that are contained in the equation, and either of them that he pleases; having first, for this purpose, laid down some observations concerning the limits within which the two roots are contained. Also, having found one of the roots, he shews how the other root may be found by means of another equation, which is a degree lower than the given one; though not by depressing the given equation, by dividing it as is now done; but from the nature of proportionals, and the theorems relating to equations, as given in the former tracts, he finds the terms of another equation, different from that last mentioned, from the root &c of which, the 2d root of the original equation may be obtained.

In the course of this work, Vieta makes also some observations on equations that are ambiguous, or have three roots; namely, that the equation  $1C - 6Q + 11N = 6$ , or as we write it  $x^3 - 6x^2 + 11x = 6$  is ambiguous, when the 2d term is negative, and the 3d term affirmative, and when  $\frac{1}{3}$  of the square of 6 the coefficient of the 2d term, exceeds 11, the coefficient of the 3d term, and has then three roots. Or in general, if  $x^3 - ax^2 + bx = c$ , and  $\frac{1}{3}a^2 > b$ , the equation is ambiguous, and has three roots. He shews also, from the relation of the coefficients, how to find whether the roots are in arithmetical progression or not, and how far the middle root differs from the extremes, by means of a cubic equation of this form  $x^3 - ba = c$ . In all or most of which remarks he was preceded by Cardan.—Vieta also remarks that the case  $x^3 - 9x^2 + 24x = 20$ , has three roots by the same rule, viz, 2, 2, 5, but that two of them are equal. And farther, that when  $\frac{1}{3}a^2 = b$ , then all the three roots are equal, as in the case  $x^3 - 6x^2 + 12x = 8$ , the three roots of which are 2, 2, 2. But when  $\frac{1}{3}a^2$  is less than  $b$ , the case is not ambiguous, having but one root. And when  $ab = c$ , then  $a = x$  is one root itself.

Many curious notes are added at the end, with remarks on the method of finding the approximate roots, when they are not rational, which is done in two ways, in imitation of the same thing in the extraction

of pure powers, viz, the one by forming a fraction of the remainder after all the figures of the homogeneous comparisonis are exhausted; the other by increasing the root of the equation in a 10 fold, or 100 fold, &c, proportion, and then dividing the root which results by 10, or 100, &c: and this is a decimal approximation. And Vieta observes that the roots will be increased 10 or 100 fold, &c, by adding the corresponding number of ciphers to the coefficient of the 2d term, double that number to the 3d, triple the same number to the 4th, and so on. So if the equation were

$$1C + 4Q + 6N = 8,$$

then  $1C + 40Q + 600N = 8000$  will have its root 10 fold, and  $1C + 400Q + 60000N = 8000000$  will have it 100 fold.

Besides the foregoing algebraical works, Vieta gave various constructions of equations by means of circles and right lines, and angular sections, which may be considered as an algebraical tract, or a method of exhibiting the roots of certain equations having all their roots affirmative, and by means of which he resolved the celebrated equation of 45 powers, proposed to all the world by Adrianus Romanus.

Having now delivered a particular analysis of Vieta's algebraical writings, it will be proper, as with other authors, to collect into one view the particulars of his more remarkable peculiarities, inventions, and improvements.

And first it may be observed, that his writings shew great originality of genius and invention, and that he made alterations and improvements in most parts of algebra; though in other parts and respects his method is inferior to some of his predecessors; as, for instance, where he neglects to avail himself of the negative roots of Cardan; the numeral exponents of Stifelius, instead of which he uses the names of the powers themselves; or the fractional exponents of Stevinus; or the commodious way of prefixing the coefficient before the quantity or factor; and such like circumstances; the want of which gives his Algebra the appearance of an age much earlier than its own. But his real inventions of things before not known, may be reduced to the following particulars.

1st. Vieta introduced the general use of the letters of the alphabet to denote indefinite given quantities; which had only been done on some particular occasions before his time. But the general use of letters for the unknown quantities was before pretty common with Stifelius and his successors. Vieta uses the vowels A, E, I, O, U, Y for the unknown quantities, and the consonants B, C, D, &c, for known ones.

2d. He invented, and introduced many expressions or terms, several of which are in use to this day: such as coefficient, affirmative and negative, pure and adaffected or affected, uncix, homogeneous adfectionis, homogeneous comparisonis, the line or vinculum over

compound quantities thus  $\overline{A + B}$ . And his method of setting down his equations, is to place the homogeneous comparisonis, or absolute known term, on the right-hand side alone, and on the other side all the terms which contain the unknown quantity, with their proper signs.

3d. In most of the rules and reductions for cubic and other



other equations, he made some improvements, and variations in the modes.

4th. He shewed how to change the root of an equation in a given proportion.

5. He derived or raised the cubic and biquadratic, &c equations, from quadratics; but not by composition in Harriot's way, but by squaring and otherwise multiplying certain parts of the quadratic. And as some quadratic equations have two roots, therefore the cubics and others raised from them, have also the same two roots, and no more. And hence he comes to know what relation these two roots bear to the coefficients of the two lowest terms of cubic and other equations, when they have only 3 terms, namely, by comparing them with similar equations so raised from quadratics. And, on the contrary, what the roots are, in terms of such coefficients.

6. He made some observations on the limits of the two roots of certain equations.

7. He stated the general relation between the roots of certain equations and the coefficients of its terms, when the terms are alternately plus and minus, and none of them are wanting, or the roots all positive.

8. He extracted the roots of affected equations, by a method of approximation similar to that for pure powers.

9. He gave the construction of certain equations, and exhibited their roots by means of angular sections; before adverted to by Bombelli.

#### OF ALBERT GIRARD.

Albert Girard was an ingenious Dutch or Flemish mathematician, who died about the year 1633. He published an edition of Stevinus's Arithmetic in 1625, augmented with many notes; and the year after his death was published by his widow, an edition of the whole works of Stevinus, in the same manner, which Girard had left ready for the press. But the work which entitles him to a particular notice in this history, is his "*Invention Nouvelle en l'Algebre, tant pour la solution des equations, que pour recognoistre le nombre des solutions qu'elles reçoivent, avec plusieurs choses qui sont nécessaires a la perfection de ceste divine science*;" which was printed at Amsterdam 1629, in small quarto in 63 pages, viz, 49 pages on Arithmetic and Algebra, and the rest on the measure of the superficies of spherical triangles and polygons, by him then lately discovered.

In this work Girard first premises a short tract on Arithmetic; in the notation of which he has something peculiar, viz, dividing the numbers into the ranks of millions, billions, trillions, &c.

He next delivers the common rules of Algebra; both in integers, fractions and radicals; with the notation of the quantities and signs. In this part he uses sometimes the letters A, B, C, &c, after the manner of Vieta, but more commonly the characters of Stevinus, viz,  $\odot$ ,  $\textcircled{1}$ ,  $\textcircled{2}$ ,  $\textcircled{3}$ , &c, for the powers of the unknown quantity, with their roots  $\sqrt[5]{\textcircled{2}}$ ,  $\sqrt[1]{\textcircled{2}}$ ,  $\sqrt[2]{\textcircled{2}}$ ,  $\sqrt[3]{\textcircled{2}}$ ,  $\sqrt[4]{\textcircled{2}}$ , &c, used by Stevinus; and sometimes the more usual marks of the roots as,  $\sqrt{\textcircled{2}}$  or  $\sqrt[2]{\textcircled{2}}$ ,  $\sqrt[3]{\textcircled{2}}$ ,  $\sqrt[4]{\textcircled{2}}$ , &c; prefixing the coefficients, as  $6\textcircled{2}$ , or  $3\sqrt[5]{32}$ , or  $2\sqrt[4]{\textcircled{2}}$ . In the signs he follows his predecessors so far as to have + for plus, - or  $\div$  for minus, = for general or indefinite difference,  $A + B$  for the sum,  $A - B$  or  $A = B$  for

the difference,  $AB$  the product, and  $\frac{A}{B}$  for the quotient

of A and B. He uses the parentheses ( ) for the vinculum or bond of compound quantities, as is now commonly practised on the continent; as  $A(AB + Bq)$ , or  $\sqrt[3]{(A \text{ cub.} - 3AqB)}$ ; and he introduces the new characters ff for *greater than*, and § for *less than*; but he uses no character for equality, only the word itself.

Girard gives a new rule for extracting the cube root of binomials, which however is in a good measure tentative, and which he explains thus: To extract the cube root of  $72 + \sqrt{5120}$ .

$$\text{The squares of the terms } \begin{cases} 5184 \\ 5120 \end{cases}$$

their difference 64, and its cube root 4. Which shews that the difference between the squares of the terms required is 4; and the rational part 72 being the greater, the greater term of the root will be rational also; and farther, that the greater terms of the power and root are commensurable, as also the two less terms. Then having made a table as in the margin, where the square of the rational

$2 + \sqrt{0}$	term always exceeds that of the other, by the
$3 + \sqrt{5}$	number 4 above mentioned, one of these bi-
$4 + \sqrt{12}$	nomials must be the cubic root sought, if the
$5 + \sqrt{21}$	given quantity have such a root, and it must

be one of these four forms, for it is known to be carried far enough by observing that the cube root of 72 is less than 5, and the cube root of 5120 less than 21; indeed, this being the case, the last binomial is excluded, as evidently too great; and the first is excluded because one of its terms is 0; therefore the root must be either  $3 + \sqrt{5}$  or  $4 + \sqrt{12}$ . And to know whether of these two it must be, try which of them has its two terms exact divisors of the corresponding terms of the given quantity; then it is found that 3 and 4 are both divisors of 72, but that only 5, and not 12, is a divisor of 5120; therefore  $3 + \sqrt{5}$  is the root sought, which upon trial is found to answer. It is remarkable here that Girard uses  $4 + \sqrt{20}$  instead of  $4 + \sqrt{12}$ , and  $5 + \sqrt{29}$  instead of  $5 + \sqrt{20}$ , contrary to his own rule.

Girard then gives distinct and plain rules for bringing questions to equations, and for the reduction of those equations to their simplest form, for solution, by the usual modes, and also by the way called by Vieta *Isomeria*, multiplying the terms of the equation by the terms of a geometrical progression, by which means the roots are altered in the proportion of 1 to the ratio of the progression. He then treats of the methods of finding the roots of the several sorts of equations, quadratic, cubic, &c; and adds remarks on the proper number of conditions or equations for limiting questions. The quadratics are resolved by completing the square, and both the positive and negative roots are taken; and he observes that sometimes the equation is impossible, as  $2)$  equ.  $6\textcircled{1} - 25$ , whose roots, he adds, are  $3 + \sqrt{-16}$  and  $3 - \sqrt{-16}$ .

The cubic equations he resolves by Cardan's rule, except the irreducible case, which he the first of any resolves by a table of sines; the other cases also he resolves by tables of sines and tangents; and adds geometrical constructions by means of the hyperbola or the trisection



trisection of angles. He next adds a particular mode of resolving all sorts of equations, that have rational roots, upon the principle of the roots being divisors of the last or absolute term, as before mentioned by Peletarius; and then gives the method of approximating to other roots that are not rational, much the same way as Stevinus.

Having found one root of an equation, by any of the former methods, by means of it he depresses the equation one degree lower, then finds another root, and so on till they are all found; for he shews that every algebraic equation admits of as many solutions or roots, as there are units in the index of the highest power, which roots may be either positive or negative, or imaginary, or, as he calls them, greater than nothing, or less than nothing, or involved; so the roots of the equation 1 ③ equ. 7 ① - 6, are 2, 1, and - 3; and the roots of the equation 1 ④ equ. 4 ① - 3 are

$$\begin{aligned} &1, \\ &1, \\ &-1 + \sqrt{-2}, \\ &-1 - \sqrt{-2}. \end{aligned}$$

In depressing an equation to lower degrees, he does not use the method of resolution of Harriot, but that which is derived from the general relation of the roots and coefficients of the terms, which he here fully and universally states, viz, that the coefficient of the 2d term is equal to the sum of all the roots; that of the 3d term equal to the sum of all the products of the roots, taken two by two; that of the 4th term, the sum of the products, taken three by three; and so on, to the last or absolute term, which is the continual product of all the roots; a property which was before stated by Vieta, as to the equations that have all their roots positive; and here extended by Girard to all sorts of roots whatever: but how either Vieta or he came by this property, no where appears that I know of. From this general property, among other deductions, Girard shews how to find the sums of the powers of the roots of an equation; thus, let A, B, C, D, &c, be the 1st, 2d, 3d, 4th, &c, coefficient, after the first term, or the sums of the products taken one by one, two by two, three by three, &c; then, in all sorts of equations,

$$\left. \begin{aligned} &A \\ &Aq - 2B \\ &A \text{ cub.} - 3AB + 3C \\ &Aqq - 4AqB + 4AC + 2Bq - 4D \end{aligned} \right\} \begin{array}{l} \text{will be the} \\ \text{sum of the} \end{array} \left\{ \begin{array}{l} \text{roots,} \\ \text{squares,} \\ \text{cubes,} \\ \text{biquadrates.} \end{array} \right.$$

Girard next explains the use of negative roots in Geometry, shewing that they represent lines only drawn in a direction contrary to those representing the positive roots; and he remarks that this is a thing hitherto unknown. He then terminates the Algebra by some questions having two or more unknown quantities; and subjoins to the whole a tract on the mensuration of the surfaces of spherical triangles and polygons, by him lately discovered.

From the foregoing account it appears that,

1st, He was the first person who understood the general doctrine of the formation of the coefficients of the powers, from the sums of their roots, and their products, &c.

2d, He was the first who understood the use of negative roots in the solution of geometrical problems.

3d, He was the first who spoke of the imaginary roots, and understood that every equation might have as many roots real and imaginary, and no more, as there are

VOL. I.

units in the index of the highest power. And he was the first who gave the whimsical name of *quantities less than nothing* to the negative. And,

4th, He was the first who discovered the rules for summing the powers of the roots of any equation.

OF HARRIOT.

Thomas Harriot, a celebrated astronomer, philosopher, and mathematician, flourished about the year 1610, about which time it is probable he wrote his Algebra, as he was then, and had been for many years before, celebrated for his mathematical and astronomical labours. In that year he made observations on the spots in the sun, and on Jupiter's satellites, the same year also in which Galileo first observed them: he left many other curious astronomical observations, and amongst them, some on the remarkable comets of the years 1607 and 1618. His Algebra was left behind him unpublished, as well as those other papers, at his death, which happened in the year 1621, being then 60 years of age, and but six years after the first publication of the principal parts of Vieta's Algebra by Alexander Anderson; so that it is probable that Harriot's Algebra was written before this time, and indeed that he had never seen these pieces. Harriot's Algebra was published by his friend Walter Warner, in the year 1631: and it would doubtless be highly grateful to the learned in these sciences, if his other curious algebraical and astronomical works were published from his original papers in the possession of the Earl of Egremont, to whom they have descended from Henry Percy, the Earl of Northumberland, that noble Mæcenas of his day. The book is in folio, and intitled *Artis Analytica Praxis, ad Aequationes Algebraicas nova, expedita, & generali methodo, resolvendas*; a work in all parts of it shewing marks of great genius and originality, and is the first instance of the modern form of Algebra in which it has ever since appeared. It is prefaced by 18 definitions, which are these: 1st, Logistica Speciosa; 2d, Equation; 3d, Synthesis; 4, Analysis; 5, Composition and Resolution; 6, Forming an Equation; 7, Reduction of an Equation; 8, Verification; 9, Numerosa & Speciosa; 10, Excogitata; 11, Resolution; 12, Roots; 13 and 14, The kinds and generation of equations by multiplication, from binomial roots or factors, called original equations,

$$\begin{aligned} \text{as } \left. \begin{array}{l} a + b \\ a - c \end{array} \right| &= \begin{array}{l} aa + ba \\ -ca - bc, \end{array} \\ \text{or } \left. \begin{array}{l} a + b \\ a + c \\ a - d \end{array} \right| &= \begin{array}{l} aa + baa + bca \\ +caa - bda \\ -daa - cda - bcd, \end{array} \end{aligned}$$

where he puts *a* for the unknown quantity, and the small consonants, *b, c, d, &c*, for its literal values or roots; 15, The first form of canonical equations, which are derived from the above originals, by transposing the homogeneous, or absolute term,

$$\begin{aligned} \text{thus } aa + ba \\ -ca &= +bc, \text{ \&c; } \end{aligned}$$

16, The secondary canonicals, formed from the primary by expelling the 2d term,

$$\begin{aligned} \text{thus } aa &= +bb, \\ \text{or } aaa - bba \\ -bca \\ -cca &= +bbc \\ &+bcc; \end{aligned}$$



17, That these are called canonicals, because they are adapted to canons or rules for finding the numeral roots, &c. 18, Reciprocal equations, in which the homogeneous is the product of the coefficients of the other terms, and the first term, or highest power of the root, is equal to the product of the powers in the other terms, as  $aaa - caa + bba = +bbc$ .

After these definitions, the work is divided into two principal parts; 1st, of various generations, reductions, and preparations of equations for their resolution in the 2d part. The former is divided into 6 sections as follows.

*Sect. 1. Logistices Speciosæ*, exemplified in the 4 operations of addition, subtraction, multiplication, and division; as also the reduction of algebraic fractions, and the ordinary reduction of irregular equations to the form proper for the resolution of them, namely, so that all the unknown terms be on one side of the equation, and the known term on the other, the powers in the terms ranged in order, the greatest first, and the first or highest power made positive, and freed from its coefficient; as  $aa + ba = cd$ ,

$$\text{or } aaa + baa - cda = -c^2d.$$

In this part he explains some unusual characters which he introduces, namely

= for equality, as  $a = b$ .

> for majority, as  $a > b$ ,

< for minority, as  $a < b$ ;

but the first had been before introduced by Robert Recorde.

*Sect. 2.* The generation of original equations from binomial factors or roots, and the deducing of canonicals from the originals. He supposes that every equation has as many roots as dimensions in its highest power; then supposing the values of the unknown letter  $a$  in any equation to be  $b, c, d, f$ , &c, that is  $a = b$ , and  $a = c$ , and  $a = d$ , &c; by transposition, or equal subtraction, these become  $a - b = 0$ , and  $a - c = 0$ , and  $a - d = 0$ , &c, or the same letters with contrary signs, for negative values or roots; then two of these binomial factors multiplied together, gives a quadratic equation, three of them a cubic, four of them a biquadratic, and so on, with all the terms on one side of the equation, and 0 on the other side, since, every binomial factor being  $= 0$ , the continual product of all of them must also be  $= 0$ . Thus,

$$\begin{array}{l|l} a + b & = aaa + baa + bca \\ a + c & + caa - bda \\ a - d & - daa - cda - bcd = 0 \end{array}$$

an original equation,

and

$$\begin{array}{l} aaa + baa + bca \\ + caa - bda \\ - daa - cda = +bcd \end{array}$$

its canonical, deduced from it. And these operations are carried through all the cases of the 2d, 3d and 4th powers, as to the varieties of the signs  $+$  and  $-$ , and the proportions of the roots as to equal and unequal, with the reciprocals, &c. From which are made evident, at one glance of the eye, all the relations and properties between the roots of equations, and the coefficients of the terms.

*Sect. 3. Aequationum canonicarum secundariorum a primariis reductio per gradus alicujus pædici sublationem*

*radice supposititia invariata manente.* Containing a great many examples of preparing equations by taking away the 2d, 3d, or any other of the intermediate terms, which is done by making the positive coefficients in that term, equal to the negative ones, by which means the whole term vanishes, or becomes equal to nothing.

They are extended as far as equations of the 5th degree; and at the end are collected, and placed in regular order, all the secondary canonicals, so reduced, so that by the uniform law which is visible through them all, the series may be continued to the higher degrees as far as we please.

*Sect. 4. Aequationum canonicarum tam primariorum, quam secundariorum, radicum designatio.* A great many literal equations are here set down, and their roots assigned from the form of the equation, that is all their positive roots; for their negative roots are not noticed here; and it is every where proved that they cannot have any more positive roots than these, and consequently the rest are negative. That those are roots, he proves by substituting them instead of the unknown letter  $a$  in the equation, when they make all the terms on one side come to the same thing as the homogeneous on the other side.

*Sect. 5. In qua aequationum communium per canonicarum equipollentiam, radicum numerus determinatur.* On the number of the roots of common equations, that is the positive roots. This Harriot determines by comparing them with the like cases found among his canonical forms, which two equations, having the same number of terms with the same signs, and the relations of the coefficients and homogeneous correspondent, he calls equipollents. And whatever was the number of positive roots used in the composition of the canonical, the same, he infers, is the number in the proposed common equation. It is remarkable that in all the examples here used, the number of positive roots is just equal to the number of the changes in the signs from  $+$  to  $-$  and from  $-$  to  $+$ , which is a circumstance, though not here expressly mentioned, that could not escape the observation, or the eye, of any one, much less of so clear and comprehensive a sight as that of Harriot. In this section are contained many ingenious disquisitions concerning the limits and magnitudes of quantities, with several curious lemmas laid down to demonstrate the propositions by, which lemmas are themselves demonstrated in a pure mathematical way, from the magnitudes themselves, independent of geometrical figures; such as, 1, If a quantity be divided into any two unequal parts, the square of half the line will be greater than the product of the two unequal parts. 2, In three continued proportionals, the sum of the extremes is greater than double the mean. 3, In four continued proportionals, the sum of the extremes is greater than the sum of the two means. 4, In any two quantities, one-fourth the square of the sum of the cubes, is greater than the cube of the product of the two quantities. 5, Of any two quantities  $q$  and  $r$ , then  $\frac{1}{2}(qq + qr + rr)^3 > \frac{1}{4}(qqr + qrr)^2$ . 6, If any quantity be divided into three unequal parts, the square of  $\frac{1}{3}$  of the whole quantity is greater than  $\frac{1}{3}$  of the sum of the three products made of the three unequal parts. 7, Also the cube of the  $\frac{1}{3}$  part of the whole, is greater than the solid or continual product of the three unequal parts.



*Scct. 6. Aequationum communium reductio per gradus alicujus parodici exclusionem & radice supposititiae mutationem.* Here are a great many examples of reducing and transforming equations of the 2d, 3d, and 4th degrees; chiefly either by multiplying the roots of equations in any proportion, as was done by Vieta, or increasing or diminishing the root by a given quantity, after the manner of Cardan. The former of these reductions is performed by multiplying the terms of the equation by the corresponding terms of a geometrical progression, the 1st term being 1, and the 2d term the quantity by which the root is to be multiplied. And the other reduction, or transforming to another root, which may be greater or less than the given root by a given quantity, is performed commonly by substituting  $e +$  or  $- b$  for the given root  $a$ , by which the equation is reduced to a simpler form. Other modes of substitution are also used; one of which is this, viz, substituting  $\frac{ee \pm bb}{e}$  or  $e \pm \frac{bb}{e}$  for the root  $a$  in the given equation

$aaa \mp 3.bba = 2.ccc$ , by which it reduces to this quadratic form  $e^6 \mp 2c^3 e^3 = -b^6$ , from whence Cardan's forms are immediately deduced; namely  $e = \sqrt[3]{c^3} \pm \sqrt[3]{c^6 \pm b^6}$ , and therefore  $a$  or  $e \pm \frac{bb}{e} = \sqrt[3]{c^3} \pm \sqrt[3]{c^6 \pm b^6} \pm \sqrt[3]{c^3} = \sqrt[3]{c^3} \pm \sqrt[3]{c^6 \pm b^6}$ ; where he denotes the cube or 3d root thus  $\sqrt[3]{\phantom{x}}$ , but without any vinculum over the compound quantities.

In this section, Harriot makes various remarks as they occur: thus he remarks, and demonstrates, that  $eee - 3.bbe = -ccc$

$-2.bbb$  is an impossible equation, or has no affirmative root. He remarks also that the three cases of the equation  $aaa - 3.bba = + 2.ccc$  are similar to the three conic sections; namely to the hyperbola when  $c > b$ , to the parabola when  $c = b$ , or to the ellipsis when  $c < b$ , and for which reason this case is not generally resolvable in species.

Having thus shewn how to simplify equations, and prepare them for solution, Harriot enters next upon the second part of his work, being the

*Exegetice Numerosa,*

or the numeral resolution of all sorts of equations by a general method, which is exemplified in a great number of equations, both simple and affected as far as the 5th power inclusive; and they are commonly prepared, by the foregoing parts, by freeing them from their 2d term, &c. These extractions are explained and performed in a way different from that of Vieta; and the examples are first in perfect or terminate roots, and afterwards for irrational or interminate ones, to which Harriot approximates by adding always periods of ciphers to the given number or resolvend, as far as necessary in decimals, which are continued and set down as such, but with their proper denominator 10, or 100, or 1000, &c.

He then concludes the work with

*Canones Directorii,*

which form a collection of the cases or theorems for making the foregoing numeral extractions, ready arranged for use, under the various forms of equations, with the factors necessary to form the several resolvends and subtrahends.

And from a review of the whole work, it appears that Harriot's inventions, peculiarities, and improvements in algebra, may be comprehended in the following particulars.

1st. He introduced the uniform use of the small letters  $a, b, c, d$ , &c, viz, the vowels  $a, e$ , &c for unknown quantities, and the consonants  $b, c, d, f$ , &c for the known ones; which he joins together like the letters of a word, to represent the multiplication or product of any number of these literal quantities, and prefixing the numeral coefficient as we do at present, except only separated by a point, thus  $5.bbc$ . For a root he set the index of the root after the mark  $\sqrt{\phantom{x}}$ ; as  $\sqrt[3]{\phantom{x}}$  for the cube root. He also introduced the characters  $>$  and  $<$  for greater and less; and in the reduction of equations, he arranged the operations in separate steps or lines, setting the explanations in the margin on the left hand, for each line. By which, and other means, he may be considered as the introducer of the modern state of Algebra, which quite changed its form under his hands.

2d. He shewed the universal generation of all the compound or affected equations, by the continual multiplication of so many simple ones, or binomial roots; thereby plainly exhibiting to the eye the whole circumstances of the nature, mystery and number of the roots of equations; with the composition and relations of the coefficients of the terms; and from which many of the most important properties have since been deduced.

3d. He greatly improved the numeral exegetis, or extraction of the roots of all equations, by clear and explicit rules and methods, drawn from the foregoing generation or composition of affected equations of all degrees.

OF OUGHTRED'S CLAVIS.

Oughtred was contemporary with Harriot, but lived a long time after him. His Clavis was first published in 1631, the same year in which Harriot's Algebra was published by his friend Warner. In this work, Oughtred chiefly follows Vieta, in the notation by the capitals  $A, B, C, D$ , &c, in the designation of products, powers, and roots, though with some few variations. His work may be comprehended under the following particulars.

1. *Notation.* This extends to both Algebra and Arithmetic, vulgar and decimal. The Algebra chiefly after the manner of Vieta, as abovesaid. And he separates the decimals from the integers thus,  $21\overline{56}$ , which is the first time I have observed such a separation, and the decimals set down without their denominator.

2. The common rules or operations of Arithmetic and Algebra. In algebraic multiplication, he either joins the letters together like a word, or connects them by the mark  $\times$ , which is the first introduction of this character of multiplication: thus  $A \times A$  or  $AA$  or  $A^2$ . But omitting the vinculum over compound factors, used by Vieta. He introduces here many neat and useful contractions in multiplication and division of decimals: as that common one of inverting the multiplier, to have fewer decimals, and abridge the work; that of omitting always one figure at a time, of the divisor, for the same purpose; dividing by the component factors of a number instead of the number itself; as 4 and 6 for 24; and many other neat contractions. He



states his proportions thus  $7.9 :: 28.36$ , and denotes continued proportion thus  $\div$ ; which is the first time I have observed these characters.

3. Invents and describes various symbolical marks or abbreviations, which are not now used.

4. *The genesis and analysis of powers.* Denotes powers like Vieta, and also roots, thus  $\sqrt{q6}$ ,  $\sqrt{c20}$ ,  $\sqrt{qq24}$ , &c; and much in his manner too performs the numeral extraction of roots. He here gives a table of the powers of the binomial  $A + E$  as far as the 10th power, with all their terms and coefficients, or uncixæ as he calls them, after Vieta.

5. *Equations.* He here gives express and particular directions for the several sorts of reductions, according as the form of the equation may require. And he uses the letter  $u$  after  $\sqrt{\phantom{x}}$ , for universal, instead of the vinculum of Vieta. And observes that the signs of all the terms of the powers of  $A + E$  are positive, but those of  $A - E$  are alternately positive and negative.

6. Next follow many properties of triangles and other geometrical figures; and the first instance of applying Algebra to Geometry, so as to investigate new geometrical properties; and after the algebraical resolution of each problem, he commonly deduces and gives a geometrical construction adapted to it. He gives also a good tract on angular sections.

7. The work concludes with the numeral resolution of affected equations, in which he follows the manner of Vieta, but he is more explicit.

#### OF DESCARTES.

Descartes's Geometry was first published in 1637, being six years after the publication of Harriot's Algebra. That work was rather an application of Algebra to Geometry, than the science either of Algebra or Geometry itself, purely and properly so called. And yet he made improvements in both. We must observe however, that all the properties of equations, &c, which he sets down, are not to be considered as even meant by himself for new inventions or discoveries; but as statements and enumerations of properties, before known and taught by other authors, which he is about to make some use or application of, and for which reason it is that he mentions those properties.

Descartes's Geometry consists of three books. The first of these is, *De Problematibus, quæ construi possunt, adhibendo tantum rectas lineas & circulos.* He here accommodates, or performs arithmetical operations by Geometry, supposing some line to represent unity, and then, by means of proportionals, shewing how to multiply, divide, and extract roots by lines. He next describes the notation he uses, but not because it is a new one, for it is the same as had been used by former authors, viz,  $a + b$  for the addition of  $a$  and  $b$ , also  $a - b$  for their subtraction,  $ab$  multiplication,  $\frac{a}{b}$  division,  $aa$

or  $a^2$  the square of  $a$ ,  $a^3$  its cube, &c: also  $\sqrt{a^2 + b^2}$  for the square root of  $a^2 + b^2$ , and  $\sqrt[3]{C.a^3 - b^3 + abb}$  for the cube root, &c. He then observes, after Stifelius, that there must be as many equations as there are unknown lines or quantities; and that they must be reduced all to one final equation, by exterminating all the unknown letters except one; when the final equation will appear like these,

$$z \propto b, \text{ or}$$

$$z^2 \propto -az + b^2, \text{ or}$$

$$z^3 \propto +az^2 + b^2z - c^3, \text{ or}$$

$$z^4 \propto +az^3 + b^2z^2 - c^3z + d^4, \text{ \&c.}$$

Where he uses  $\propto$  for  $=$  or equality, setting the highest term or power alone on one side of the equation, and all the other terms on the other side, with their proper signs.

Descartes next defines plane problems, namely, such as can be resolved by right lines and circles, described on a plane superficies; and then the final equation rises only to the 2d power of the unknown letter. He then constructs such equations, viz. quadratics, by the circle, thus finding geometrically the root or roots, that is, the positive ones. But when the lines, by which the roots are determined, neither cut nor touch, he observes that the equation has then no possible root, or that the problem is impossible. He then concludes this book with the algebraical solution of the celebrated problem, before treated of by the antients, namely, to find a point, or the locus of all the points, from whence a line being drawn to meet any number of given lines in given angles, the product of the segments of some of them shall have a given ratio to that of the rest.

*Lib. 2. De Natura Linearum Curvarum.* This is a good algebraical treatise on curve lines in general, and the first of the kind that has been produced by the moderns. Here the nature of the curve is expressed by an equation containing two unknown or variable lines, and others that are known or constant, as  $y^2 \propto cy - \frac{cxy}{b}$

+  $ay - ac$ . But, not relating to pure Algebra, the particulars will be most properly placed under the article of curve lines, and other terms relating to them. Only one discovery, among many ingenious applications of Algebra to Geometry, may here be particularly noticed, as it may be considered as the first step towards the arithmetic of infinites; and that is the method of tangents, here given, or, which comes to the same thing, of drawing a line perpendicular to a curve at any point, which is an ingenious application of the general form of an equation, generated in Harriot's way, that has two equal roots, to the equation of the curve. Of which a particular account will be given at the article TANGENTS.

*Lib. 3. De Construtione Problematum Solidorum, et Solida excedentium.* Descartes begins this book with remarks on the nature and roots of equations, observing that they have as many roots as dimensions, which he shews, after Harriot, by multiplying a certain number of simple binomial equations together, as  $x - 2 \propto 0$ , and  $x - 3 \propto 0$ , and  $x - 4 \propto 0$ , producing  $x^3 - 9xx + 26x - 24 \propto 0$ . He here remarks that equations may sometimes have their roots false, or what we call negative, which he opposes to those that are positive, or as he calls them true, as Cardan had done before. As a natural deduction from the generation or composition of equations, by multiplication, he infers their resolution, or depression, or decomposition, namely, dividing them by the binomial factors which were multiplied to produce the equation: and he observes that by this operation it is known that this divisor is one of the binomial roots, and that there can be no more roots than dimensions, or than those which form with the unknown letter  $x$ , bino-



binomials that will exactly divide the equation, as Harriot had shewn before. Descartes adverts to several other properties, mostly known before, which he has occasion to make use of in the progress of his work; such as, that equations may have as many true roots as the terms have changes of the signs + and -, and as many false ones as successions of the same signs: which number and nature of the roots had before been partly shewn by Cardan and Vieta, from the relation of the coefficients, and their signs, and more fully by Harriot in his 5th section. And hence Descartes infers the method of changing the true roots to false, and the false to true, namely by changing the signs of the even terms only, as Cardan had taught before. Descartes then adverts to other reductions and transmutations which had been taught by Cardan, Vieta, and Harriot, such as, To increase or diminish the roots by any quantity; To take away the 2d term: To alter the roots in any proportion, and thence to free the equation from fractions and radicals.

Descartes next remarks that the roots of equations, whether true or false, may be either real or imaginary; as in the equation  $x^3 - 6xx + 13x - 10 = 0$ , which has only one real root, namely 2. The imaginary roots were first noticed by Albert Girard, as before mentioned. He then treats of the depression of a cubic equation to a quadratic, or plane problem, that it may be constructed by the circle, by dividing it by some one of the binomial factors, which, in Harriot's way, compose the equation. Peletarius having shewn that the simple root is one of the divisors of the known term of the equation, and Harriot that that term is the continual product of all the roots, Descartes therefore tries all the simple divisors of that term, till he finds one of them which, connected with the unknown letter  $x$ , by + or -, will exactly divide the equation. And the process is the same for higher powers than the cube. But when a divisor cannot be thus found, for depressing a biquadratic equation to a cubic, he gives another rule, which is a new one, for dissolving it into two quadratics, by means of a cubic equation, in this manner:

Let the given biqu. be  $+ x^4 + pxx + qx + r = 0$ ;  
Which suppose  
equal to the  $+ xx - yx + \frac{1}{2}yy + \frac{1}{2}p \cdot \frac{q}{2y} = 0$ ,  
product of  $+ xx + yx + \frac{1}{2}yy + \frac{1}{2}p \cdot \frac{q}{2y} = 0$ ;  
these two

where the sign of  $\frac{1}{2}p$  in the two quadratics must be the same as the sign of  $p$  in the given equation, and in the 1st quadratic the sign of  $\frac{q}{2y}$  must be the same as the sign of  $q$ , but in the 2d quadratic the contrary. Then if there be found the root  $yy$  of this cubic equation

$$y^6 + 2py^4 + \frac{1}{4}ppyy - qq = 0,$$

where the sign of  $2p$  is the same as of  $p$  in the given biquadratic, but the sign of  $4r$  contrary to that of  $r$  in the same: Then the value of  $y$ , hence deduced, being substituted for it in the two quadratic equations, and their two pairs of roots taken, they will be the four roots of the proposed biquadratic. And thus also, he hints, may equations of the 6th power be reduced to those of the 5th, and those of the 8th power to those of the 7th,

and so on. Descartes does not give the investigation of this rule; but it has evidently been done, by assuming indeterminate quantities, after the manner of Ferrari and Cardan, as coefficients of the terms of the two quadratic equations, and, after multiplying the two together, determining their values by comparing the resulting terms with those of the proposed biquadratic equation.

After these reductions, which are only mentioned for the sake of the geometrical constructions which follow, by simplifying and depressing the equations as much as they will admit, Descartes then gives the construction of solid and other higher problems, or of cubic and higher equations, by means of parabolas and circles; where he observes that the false roots are denoted by the ordinates to the parabola lying on the contrary side of the axis to the true roots. Finally, these constructions are illustrated by various problems concerning the trisection of an angle, and the finding of two or four mean proportionals; which concludes this ingenious work.

From the foregoing analysis may easily be collected the real inventions and improvements made in algebra by Descartes. His work, as has been observed before, is not algebra itself, but the application of algebra to geometry, and the algebraical doctrine of curve lines, expressing and explaining their nature by algebraical equations, and on the contrary, constructing and explaining equations by means of the curve lines. What respects the geometrical parts of this tract we shall have occasion to advert to elsewhere; and therefore shall here only enumerate the circumstances which belong more peculiarly to the science of Algebra, which I shall distinguish into the two heads of improvements and inventions. And

1st. Of his improvements. That he might fit equations the better for their application in the construction of problems, Descartes mentions, as it were by-the-by, many things concerning the nature and reduction of equations, without troubling himself about the first inventors of them, stating them in his own terms and manner, which is commonly more clear and explicit, and often with improvements of his own. And under this head we find that he chiefly followed Cardan, Vieta, and Harriot, but especially the last, and explains some of their rules and discoveries more distinctly, and varies but a little in the notation, putting the first letters of the alphabet for the known, and the latter letters for the unknown quantities; also  $x^3$  for  $aaa$ , &c; and  $\infty$  for  $=$ . But Herigone used the numeral exponents in the same manner two years before. Descartes explained or improved most parts of the reductions of equations, in their various transmutations, the number and nature of their roots, true and false, real and what he calls imaginary, called involved by Girard; and the depression of equations to lower degrees.

2d. As to his inventions and discoveries in algebra, they may be comprehended in these particulars, namely, the application of algebra to the geometry of curve lines, the constructing equations of the higher orders, and a rule for resolving biquadratic equations by means of a cubic and two quadratics.

Having now traced the science of Algebra from its origin and rude state, down to its modern and more polished form, in which it has ever since continued, with very little variation; having analysed all or most



of the principal authors, in a chronological order, and deduced the inventions and improvements made by each of them; from this time the authors both become too numerous, and their improvements too inconsiderable, to merit a detail in the same minute and circumstantial way: and besides, these will be better explained in a particular manner under the word or article to which each of them severally belongs. It may therefore now suffice to enumerate, or announce only in a cursory manner, the chief improvements and authors on algebra down to the present time.

After the publication of the Geometry of Descartes, a great many other ingenious men followed the same course, applying themselves to algebra and the new geometry, to the mutual improvement of them both; which was done chiefly by reasoning on the nature and forms of equations, as generated and composed by Harriot. Before proceeding upon these however, it is but proper to take notice here of Fermat, a learned and ingenious mathematician, who was contemporary and a competitor of Descartes for his brightest discoveries, which he was in possession of before the geometry of Descartes appeared. Namely, the application of algebra to curve lines, which he expressed by an algebraical equation, and by them constructing equations of the 3d and 4th orders; also a method of tangents, and a method de maximis et minimis, which approach very near to the method of Fluxions or Increments, which they strikingly resemble both in the manner of treating the problems, and in the algebraic notation and process. The particulars of which, see under their proper heads. Besides these, Fermat was deeply learned in the Diophantine problems, and the best edition of Diophantus's Arithmetic, is that which contains the notes of Fermat on that ingenious work.

But to return to the successors of Descartes. His geometry having been published in Holland, several learned and ingenious mathematicians of that country, presently applied themselves to cultivate and improve it; as Schooten, Hudde, Van-Heuraet, De Witte, Slusius, Huygens, &c; besides M. de Beaune, and perhaps some others in France.

Francis Schooten, professor of mathematics in the university of Leyden, was one of the first cultivators of the new geometry. He translated Descartes's Geometry out of French into Latin, and published it in 1649, with his commentary upon it, as also Brief Notes of M. de Beaune; both of them containing many ingenious and useful things. And in 1659 he gave a new edition of the same in two volumes, with the addition of several other ingenious pieces: as two posthumous tracts of de Beaune, the one on the nature and constitution, the other on the limits of equations, shewing how to assign the limits between which are contained the greatest and least roots of equations, extended and completed by Erasmus Bartholine: two letters of M. Hudde on the reduction of equations, and on the maxima and minima of quantities, containing many ingenious rules; among which are some concerning the drawing of tangents, and on the equal roots of equations, which he determines by multiplying the terms of the equation by the terms of any arithmetical progression, 0 being one of the terms, the equation is commonly depressed one degree lower: also a tract of Van Heuraet on the rectifi-

cation of curve lines; the elements of curves by De Witte; Schooten's principles of universal mathematics, or introduction to Descartes's geometry, which had before been published by itself in 1651; and to the end of the work is added a posthumous piece of Schooten's (for he died while the 2d vol. was printing) intitled *Tractatus de concinnandis demonstrationibus geometricis ex calculo algebraico*. Schooten also published, in 1657, *Exercitationes Mathematicæ*, in which are contained many curious algebraical and analytical pieces, amongst others of a geometrical nature.

An elaborate commentary on Descartes's Geometry was also published by F. Rabuel, a Jesuit; and James Bernoulli, enriched with notes, an edition of the same, printed at Basil in 169—.

The celebrated Huygens also, among his great discoveries, very much cultivated the algebraical analysis: and he is often cited by Schooten, who relates divers inventions of his, while he was his pupil.

Slusius, a canon of Liege, published in 1659, *Mesolabum, seu duæ mediæ propor. per circulum & ellips. vel hyperb. infinitis modis exhibitæ*; by which, any solid problem may be constructed by infinite different ways. And in 1668 he gave a second edition of the same, with the addition of the analysis, and a miscellaneous collection of curious and important problems, relating to spirals, centres of gravity, maxima and minima, points of inflexion, and some Diophantine problems; all shewing him deeply skilled in Algebra and Geometry.

There have been a great number of other writers and improvers of Algebra, of which it may suffice slightly to mention the chief part, as in the following catalogue.

Peter Nonius, or Nunez, a Spaniard, wrote about the time of Cardan, or soon after.

In 1619 several pieces of Van Collen, or Ceulen, were translated out of Dutch into Latin, and published at Leyden by W. Snell; among which are contained a particular treatise on surds, and his proportion of the circumference of a circle, to its diameter.

In 1621 Bachet published, in Greek and Latin, an edition of Diophantus, with many notes. And another edition of the same was published in 1670, with additions by Fermat.

In 1624 Bachet's *Problemes Plaisans et Delectables*, being curious problems in mathematical recreations.

In 1634 Herigone published, at Paris, the first course of mathematics, in 5 vols. 8vo; in the 2d of which is contained a good treatise on Algebra; in which he uses the notation by small letters, introduced by the Algebra of Harriot, which was published three years before, though the rest of it does not resemble that work, and one would suspect that Herigone had not seen it. The whole of this piece bears evident marks of originality and ingenuity. Besides + for plus, he uses ∞ for minus, and | for equality, with several other useful abbreviations and marks of his own. In the notation of powers, he does not repeat the letters like Harriot, but subjoins the numeral exponents, to the letter, as Descartes did two years afterwards. And Herigone uses the same numeral exponents for roots, as  $\sqrt[3]{}$  for the cube root.

In 1635 Cavalerius published his *Indivisibles*; which proved a new æra in analytics, and gave rise to other new modes of computation in analytics.

About



About 1640, et seq. Roberval made several notable improvements in analytics, which are published in the early volumes of the Memoirs of the Academy of Sciences; as, 1. A tract on the composition of motion, and a method of tangents. 2, *De recognitione æquationum*. 3, *De geometrica planarum & cubicarum æquationum resolutione*. 4, A treatise on indivisibles, &c.

In 1643 De Billy published *Nova Geometriae Clavis Algebra*. And in 1670 *Diophantus Redivivus*. He was an author particularly well skilled in Diophantine problems.

In 1644 Renaldine published, in 4to, *Opus Algebraicum*, both ancient and modern, with mathematical resolution and composition. And in 1665, in folio, the same, greatly enlarged, or rather a new work, which is very heavy and tedious. In this work Renaldine uses the parentheses ( $a^2 + b^2$ ) as a vinculum, instead of the line over, as  $a^2 + b^2$ .

In 1655 was published Wallis's *Arithmetica Infinitorum*, being a new method of reasoning on quantities, or a great improvement on the Indivisibles of Cavalierius, and which in a great measure led the way to infinite series, the binomial theorem, and the method of fluxions. Wallis here treats ingeniously of quadratures and many other problems, and gives the first expression for the quadrature of the circle by an infinite series. Another series is here added for the same purpose, by the Lord Brouncker.

In 1659 was published *Algebra Rhonii Germanice*; which was in 1668, translated into English by Mr. Thomas Brancker, with additions and alterations by Dr. John Pell.

In 1661 was published in Dutch, a neat piece of Algebra by Mr. Kinckhuysen; which Sir I. Newton, while he was professor of mathematics at Cambridge, made use of and improved, and he meant to republish it, with the addition of his method of fluxions and infinite series; but he was prevented by the accidental burning of some of his papers.

In 1665 or 1666 Sir Isaac Newton made several of his brightest discoveries, though they were not published till afterwards: such as the binomial theorem; the method of fluxions and infinite series; the quadrature, rectification, &c of curves; to find the roots of all sorts of equations, both numeral and literal, in infinite converging series; the reversion of series, &c. Of each of which a particular account may be seen in their proper articles.

In 1666 M. Frenicle gave several curious tracts concerning combinations, magic squares, triangular numbers, &c; which were printed in the early volumes of Memoirs of the Academy of Sciences.

In 1668 Thomas Brancker published a translation of Rhonius's Algebra, with many additions by Dr. John Pell, who used a peculiar method of registering the steps in any algebraical process, by means of marks and abbreviations in a small column drawn down the margin, by which each line, or step, is clearly explained, as was before done by Harriot in words at length.

In 1668 Mercator published his *Logarithmotechnia*, or method of constructing logarithms; in which he gives the quadrature of the hyperbola, by means of an infinite series of algebraical terms, found by dividing a simple algebraic quantity by a compound one, and for the first time that this operation was given to the

public, though Newton had before that expanded all sorts of compound algebraical quantities into infinite series.

In the same year was published James Gregory's *Excercitationes Geometricæ*, containing, among other things, a demonstration of Mercator's quadrature of the hyperbola, by the same series.

And in the same year was published, in the Philosophical Transactions, Lord Brouncker's quadrature of the hyperbola by another infinite series of simple rational terms, which he had been in possession of since the year 1657, when it was announced to the public by Dr. Wallis. Lord Brouncker's series for the quadrature of the circle, had been published by Wallis in his Arithmetic of Infinites.

In 1669 Dr. Isaac Barrow published his Optical and Geometrical Lectures, abounding with profound researches on the dimensions and properties of curve lines; but particularly to be noticed here for his method of tangents, by a mode of calculation similar to that of Fluxions, or Increments, from which these differ but little, except in the notation.

In 1673 was published, in 2 vols. folio, *Elements of Algebra*, by John Kersey; a very ample and complete work, in which Diophantus's problems are fully explained.

In 1675 were published *Nouveaux Elemens des Mathematiques*, par J. Prestet, prêtre: a prolix and tedious work, which he presumptuously dedicated to God Almighty.

About 1677 Leibnitz discovered his *Methodus Differentialis*, or else made a variation in Newton's Fluxions, or an extension of Barrow's method, for it is not certain which. He gave the first instance of it in the Leipzig Acts for the year 1684. He also improved infinite series, and gave a simple one for the quadrature of the circle, in the same acts for 1682.

In 1682 Ismael Bulliald published, in folio, his *Opus Novum ad Arithmetica Infinitorum*, being a large amplification of Wallis's Arithmetic of Infinites.

In 1683 Tschirnhausen gave a memoir, in the Leipzig Acts, concerning the extraction of the roots of all equations in a general way; in which he promised too much, as the method did not succeed.

In 1684 came out, in English and Latin, 4to, Thomas Baker's *Geometrical Key, or Gate of Equations Unlocked*; being an improvement of Descartes's construction of all equations under the 5th degree, by means of a circle and only one and the same parabola for all equations, using any diameter instead of the axis of the parabola.

In 1685 was published, in folio, Wallis's *Treatise of Algebra, both Historical and Practical*, with the addition of several other pieces; shewing the origin, progress, and advancement of that science, from time to time. It cannot be denied that, in this work, Wallis has shewn too much partiality to the Algebra of Harriot. Yet, on the other hand, it is as true, that M. de Gua, in his account of it, in the Memoirs of the Academy of Sciences for 1741, has run at least as far into the same extreme on the contrary side, with respect to the discoveries of Vieta; and both these I believe from the same cause, namely, the want of examining the works of all former writers on Algebra, and specifying their several discoveries; as has been done in the course of this article,



In 1687 Dr. Halley gave, in the *Philos. Transf.* the construction of cubic and biquadratic equations, by a parabola and circle; with improvements on what had been done by Descartes, Baker, &c. Also, in the same Transactions, a memoir on the number of the roots of equations, with their limits and signs.

In 1690 was published, in 4to, by M. Rolle, *Traité d'Algèbre*; in 1699 *Une Méthode pour la Résolution des Problèmes indéterminés*; and in 1704 *Mémoires sur l'inverse des tangents*; and other pieces.

In 1690 Joseph Raphson published *Analysis Aequationum Universalis*; being a general method of approximating to the roots of equations in numbers. And in 1715 he published the *History of Fluxions*, both in English and Latin.

In 1690 was also published, in 4 vols 4to, Dechale's *Cursus seu mundus mathematicus*; in which is a piece of algebra, of a very old-fashioned sort, considering the time when it was written.

About 1692, and at different times afterwards, De Lagny published many pieces on the resolution of equations in numbers, with many theorems and rules for that purpose.

In 1693 was published, in a neat little volume, *Synopsis Algebraica, opus posthumum Johannis Alexandri*.

In 1694, Dr. Halley gave, in the *Philos. Transf.* an ingenious tract on the numeral extraction of all roots, without any previous reduction. And this tract is also added to some editions of Newton's *Universal Arithmetic*.

In 1695 Mr. John Ward, of Chester, published, in 8vo, *A Compendium of Algebra*, containing plain, easy, and concise rules, with examples in an easy and clear way. And in 1706 he published the first edition of his *Young Mathematician's Guide*, or a plain and easy introduction to the mathematics: a book which is still in great request, especially with beginners, and which has been ever since the ordinary introduction of the greatest part of the mathematicians of this country.

In 1696 the Marquis de l'Hôpital published his *Analyse des infiniment petits*. And gave several papers to the *Leipfic Acts* and the *Mémoires* of the Academy of Sciences. He left behind him also an ingenious treatise, which was published in 1707, intitled *Traité analytique des Sections Coniques, et de la construction des lieux géométriques*.

In 1697, and several other years, Mr. Ab. Demoivre gave various papers, in the *Philos. Transf.* containing improvements in Algebra: viz. in 1697, A method of raising an infinite multinomial to any power, or extracting any root of the same. In 1698, The extraction of the root of an infinite equation. In 1707, Analytical solution of certain equations of the 3d, 5th, 7th, &c degree. In 1722, Of algebraic fractions and recurring series. In 1738, The reduction of radicals into more simple forms. Also in 1730, he published *Miscellanea analytica de seriebus & quadraturis*, containing great improvements in series, &c.

In 169, Mr. Richard Sault published, in 4to, *A New Treatise of Algebra, apply'd to numeral questions and geometry. With a converging series for all manner of Adfected Equations*. The series here alluded to, is Mr. Raphson's method of approximation, which had been lately published.

In 1699 Hyac. Christopher published at Naples, in 4to, *De constructione equationum*.

In 1702 was published Ozanam's Algebra; which is chiefly remarkable for the Diophantine analysis. He had published his mathematical dictionary in 1691, and in 1693 his course of mathematics, in 5 vols 8vo, containing also a piece of algebra.

In 1704, Dr. John Harris published his *Lexicon Technicum*, the first dictionary of arts and sciences: a very plain, and useful book, especially in the mathematical articles. And in 1705 a neat little piece on algebra and fluxions.

In 1705 M. Guisnée published, in 4to, his *Application de l'algebre a la geometrie*: a useful book.

In 1706 Mr. William Jones published his *Synopsis Palmariorum Matheseos*, or a new introduction to the mathematics: a very useful compendium in the mathematical sciences. And in 1711 he published, in 4to, a collection of Sir Isaac Newton's papers, intitled *Analysis per quantitatum series, fluxiones, ac differentias: cum enumeratione linearum tertii ordinis*.

In 1707 was published by Mr. Whiston, the first edition of Sir Isaac Newton's *Arithmetica Universalis: sive de compositione et resolutione arithmetica liber*: and many editions have been published since. This work was the text book used by our great author in his lectures, while he was professor of mathematics in the university of Cambridge. And although it was never intended for publication, it contains many and great improvements in analytics; particularly in the nature and transmutation of equations; the limits of the roots of equations; the number of impossible roots; the invention of divisors, both surd and rational; the resolution of problems, arithmetical and geometrical; the linear construction of equations; approximating to the roots of all equations, &c. To the later editions of the book is commonly subjoined Dr. Halley's method of finding the roots of equations. As the principal parts of this work are not adapted to the circumstance of beginners, there have been published commentaries upon it by several persons, as s'Gravande, Castillon, Wilder, &c.

In 1708 M. Reyneau published his *Analyse Démontrée*, in 2 vols 4to. And in 1714 *La Science du Calcul, &c.*

In 1709 was published an English translation of Alexander's algebra. With an ingenious appendix by Humphry Ditton.

In 1715 Dr. Brooke Taylor published his *Methodus Incrementorum*: an ingenious and learned work. And in the *Philos. Transf.* for 1718, An improvement of the method of approximating to the roots of equations in numbers.

In 1717 M. Nicole gave, in the *memoirs* of the academy of sciences, a tract on the calculation of finite differences. And in several following years, he gave various other tracts on the same subject, and on the resolution of equations of the 3d degree, and particularly on the irreducible case in cubic equations.

Also in 1717 was published a treatise on Algebra by Philip Ronayne.

Also in 1717 Mr. James Sterling published *Linea tertii Ordinis*; an ingenious work, containing good improvements in analytics. Also in 1730 *Methodus Differentialis: sive tractatus de summatione et interpolatione serierum infinitarum*: with great improvements on infinite series.



In 1726 and 1729 Maclaurin gave, in the *Philos. Trans.* tracts on the imaginary roots of equations. And afterwards was published, from his posthumous papers, his treatise on Algebra, with its application to curve lines.

In 1727 came out s'Gravesande's Algebra, with a specimen of a commentary on Newton's universal arithmetic.

In 1728 Mr. Campbell gave, in the *Philos. Trans.* an ingenious paper on the number of impossible roots of equations.

In 1732 was published Wolfius's Algebra, in his course of mathematics, in 5 vols. 4to.

In 1735 Mr. John Kirkby published his arithmetic and algebra. And in 1748 his doctrine of ultimators.

In 1740 were published Mr. Thomas Simpson's Essays; in 1743 his Dissertations, and in 1757 his Tracts; in all which are contained several improvements in series and other parts of Algebra. As also in his algebra, first printed in 1745, and in his Select Exercises, in 1752.

Also in 1740 was published professor Saunderson's Elements of Algebra, in 2 vols. 4to.

In 1741 M. de la Caille published *leçons de mathématiques; ou élémens d'algebre & de geometrie*.

Also in 1741, in the memoirs of the academy of sciences, were given two articles by M. de Gua, on the number of positive, negative, and imaginary roots of equations. With an historical account of the improvements in Algebra; in which he severely censures Wallis for his partiality; a circumstance in which he himself is not less faulty.

In 1746 M. Clairaut published his *Elémens d'algebre*, in which are contained several improvements, especially on the irreducible case in cubic equations. He has also several good papers on different parts of analytics, in the memoirs of the academy of sciences.

In 1747 M. Fontaine gave, in the memoirs of the academy of sciences, a paper on the resolution of equations. Besides some analytical papers in the memoirs of other years.

In 1761 M. Castillon published, in 2 vols 4to, Newton's universal arithmetic, with a large commentary.

In 1763 Mr. Emerson published his *Increments*. In 1764 his Algebra, &c.

In 1764 Mr. Landen published his *Residual Analysis*. In 1765 his *Mathematical Lucubrations*. And in 1780 his *Mathematical Memoirs*. All containing good improvements in infinite series, &c.

In 1770 was published, in the German language, *Elements of Algebra* by M. Euler. And in 1774 a French translation of the same. The memoirs of the Berlin and Petersburg academies also abound with various improvements on series and other branches of analysis by this great man.

In 1775 was published at Bologna, in 2 vols 4to, *Compendio d'Analisi di Girolamo Saladini*.

Besides the foregoing, there have been many other authors who have given treatises on Algebra, or who have made improvements on series and other parts of Algebra; as Schonerus, Coignet, Salignac, Laloubere, Hemischi, Degraave, Mescher, Henischi, Roberval, the Bernoullis, Malbranche, Agnesi, Wells, Dodson, Manfredi, Regnault, Rowning, Maseres, Waring, Lorg-

VOL. I.

na, de la Grange, de la Place, Bertrand, Kuhnus, Hales, and many others.

ALGEBRA, *numeral*, is that which is chiefly concerned in the solution of numeral problems, and in which all the given quantities are expressed by numbers only. As used by the more early authors, Diophantus, Pacioli, Stifelius, &c.

ALGEBRA, *specious*, or *literal*, is that commonly used by the moderns, in which all the quantities, both known and unknown, are represented or expressed by species or general characters, as the letters of the alphabet, &c; in consequence of which general designation, all the conclusions become universal theorems for performing every operation of the like kind. There are specimens of this method from Cardan and others about his time, but it was more generally employed and introduced by Vieta.

ALGEBRAICAL, something relating to algebra.

Thus we say *algebraical* solutions, curves, characters or symbols, &c.

ALGEBRAICAL *Curve*, is a curve in which the general relation between the abscisses and ordinates can be expressed by a common algebraical equation.

These are also called *geometrical lines* or *curves*, in contradistinction to *mechanical* or *transcendental* ones.

ALGEBRAIST, a person skilled in algebra.

ALGENEB, or ALGENIB, a fixed star of the second magnitude, on the right side of Perseus.

ALGOL, or *Medusa's Head*, a fixed star of the third magnitude, in the constellation Perseus.

ALGORAB, a fixed star of the third magnitude, in the right wing of the constellation Corvus.

ALGORISM, or ALGORITHM, is similar to logistics, signifying the art of computing in any particular way, or about some particular subject; or the common rules of computing in any art. As the *algorithm* of numbers, of algebra, of integers, of fractions, of surds, &c; meaning the common rules for performing the operations of arithmetic, or algebra, or fractions, &c.

ALHAZEN, ALLACEN, or ABDILAZUM, was a learned Arabian, who lived in Spain about the year 1100, according to his editor Risner, and Weidler. He wrote upon Astrology; and his work upon Optics was printed, in Latin, at Basil, in 1572, under the title of *Optica Thesaurus*, by Risner. Alhazen was the first who shewed the importance of refractions in astronomy, so little known to the ancients. He is also the first author who has treated on the twilight, upon which he wrote a work, in which he also speaks of the height of the clouds.

ALIDADE, an Arabic name for the label, index, or ruler, which is moveable about the centre of an astrolabe, quadrant, &c, and carrying the sights or telescope, and by which are shewn the degrees cut off the limb or arch of the instrument.

ALIQUANT *part*, is that part which will not exactly measure or divide the whole, without leaving some remainder. Or the *aliquant* part is such, as being taken or repeated any number of times, does not make up the whole exactly, but is either greater or less than it. Thus 4 is an aliquant part of 10; for 4 twice taken makes 8 which is less than 10, and three times taken makes 12 which is greater than 10.

O

ALIQUOT



**ALIQUOT** *part*, is such a part of any whole, as will exactly measure it without any remainder. Or the *aliquot* part is such, as being taken or repeated a certain number of times, exactly makes up, or is equal to the whole. So 1 is an aliquot part of 6, or of any other whole number; 2 is also an aliquot part of 6, being contained just 3 times in 6; and 3 is also an aliquot part of 6, being contained just 2 times: so that all the aliquot parts of 6 are 1, 2, 3.

All the *aliquot* parts of any number may be thus found: Divide the given number by its least divisor; then divide the quotient also by its least divisor; and so on always dividing the last quotient by its least divisor, till the quotient 1 is obtained; and all the divisors, thus taken, are the prime aliquot parts of the given number. Then multiply continually together these prime divisors, viz. every two of them, every three of them, every four of them, &c; and the products will be the other or compound *aliquot* parts of the given number. So if the *aliquot* parts of 60 be required; first divide it by 2, and the quotient is 30: then 30 divided by 2 also, gives 15, and 15 divided by 3 gives 5, and 5 divided by 5 gives 1: so that all the prime divisors or aliquot parts are 1, 2, 2, 3, 5. Then the compound ones, by multiplying every two, are 4, 6, 10, 15; and every three 10, 20, 30. So that all the *aliquot* parts of the given number 60, are 1, 2, 3, 4, 5, 6, 10, 12, 15, 20, 30.—In like manner it will be found that all the *aliquot* parts of 360 are 1, 2, 3, 4, 5, 6, 8, 9, 10, 12, 15, 18, 20, 24, 30, 36, 40, 45, 60, 72, 180.

**ALLEN** (THOMAS) a celebrated mathematician of the 16th century. He was born at Uttoxeter in Staffordshire, in 1542; was admitted a scholar of Trinity college, Oxford, in 1561; where he took his degree of master of arts in 1567. In 1570 he quitted his college and fellowship, and retired to Gloucester hall, where he studied very closely, and became famous for his knowledge in antiquity, philosophy and mathematics. He received an invitation from Henry earl of Northumberland, a great friend and patron of the mathematicians, and he spent some time at the earl's house; where he became acquainted with those celebrated mathematicians Thomas Harriot, John Dee, Walter Warner, and Nathaniel Torporley. Robert earl of Leicester, too, had a great esteem for Allen, and would have conferred a bishopric upon him; but his love for solitude and retirement made him decline the offer. His great skill in the mathematics gave occasion to the ignorant and vulgar to look upon him as a magician or conjurer. Allen was very curious and indefatigable in collecting scattered manuscripts relating to history, antiquity, astronomy, philosophy, and mathematics: which collections have been quoted by several learned authors, and mentioned as in the Bibliotheca Alleniana. He published in Latin the second and third books of Ptolemy, *Concerning the Judgment of the Stars*, or, as it is usually called, of the *quadripartite construction*, with an exposition. He wrote also notes on many of Lilly's books, and some on John Bale's work, *De scriptoribus Maj. Britannia*. He died at Gloucester hall in 1632, being 90 years of age.

Mr. Burton, the author of his funeral oration, calls him the very soul and sun of all the mathematicians of his age. And Selden mentions him as a person of the most extensive learning and consummate judgment, the bright-

est ornament of the university of Oxford. Also Camden says he was skilled in most of the best arts and sciences. A. Wood has also transcribed part of his character from a manuscript in the library of Trinity college, in these words: "He studied polite literature with great application; he was strictly tenacious of academic discipline, always highly esteemed both by foreigners and those of the university, and by all of the highest stations in the church of England, and the university of Oxford. He was a sagacious observer, an agreeable companion, &c."

**ALLIGATION**, one of the rules in arithmetic, by which are resolved questions which relate to the compounding or mixing together of divers simples or ingredients, being so called from *alligare*, to tie or connect together, probably from certain vincula, or crooked ligatures, commonly used to connect or bind the numbers together.

It is probable that this rule came to us from the Moorish or Arabic writers, as we find it, with all the other rules of arithmetic, in *Lucas de Burgo*, and the other early authors in Europe.

*Alligation* is of two kinds, *medial* and *alternate*.

**ALLIGATION** *medial* is the method of finding the rate or quality of the composition, from having given the rates and quantities of the simples or ingredients.

The rule of operation is this: multiply each quantity by its rate, and add all the products together; then divide the sum of the products by the sum of the quantities, or whole compound, and the quotient will be the rate sought.

*For example*, Suppose it were required to mix together 6 gallons of wine, worth 5s. a gallon; 8 gallons, worth 6s. the gallon; and 4 gallons, worth 8s. the gallon; and to find the worth or value, per gallon, of the whole mixture.

	Gal.		s.	products.
Here	6	mult. by 5	gives	30
	8	- by 6	-	48
	4	- by 8	-	32
<hr/>				
whole comp.	18			110 sum of prod.
Then 18 ) 110 ( 6 $\frac{2}{3}$ or 6 $\frac{1}{3}$ s. is the rate sought.				
				108
				<hr/>
				2

**ALLIGATION** *alternate* is the method of finding the quantities of ingredients or simples, necessary to form a compound of a given rate.

The *rule* of operation is this: 1st, Place the given rates of the simples in a column, under each other; noting which rates are less, and which are greater than the proposed compound. 2d, Connect or link with a crooked line, each rate which is less than the proposed compound rate, with one or any number of those which are greater than the same; and every greater rate with one or any number of the less ones. 3d, Take the difference between the given compound rate and that of each simple rate, and set this difference opposite every rate with which that one is linked. 4th, Then if only one difference stand opposite any rate, it will be the quantity belonging to that rate; but when there are several differences to any one, take their sum for its quantity.

*For example*, Suppose it were required to mix together gold of various degrees of fineness, viz. of 19, of



of 21, and of 23 caracts fine, so that the mixture shall be of 20 caracts fine. Hence,

	Rates	Diff.	Sum of Diff.
Comp. rate 20	21	- - - 1	{ 1 of 21 caracts fine, 4 of 19 caracts fine, 1 of 23 caracts fine.
	19	- - - 1 + 3	
	23	- - - 1	

That is, there must be an equal quantity of 21 and 23 caracts fine, and 4 times as much of 19 caracts fine.

Various limitations, both of the compound and the ingredients, may be conceived; and in such cases, the differences are to be altered proportionally.

Questions of this sort are however commonly best and easiest resolved by common Algebra, of which they form a species of indeterminate problems, as they admit of many, or an indefinite number of answers.

There is recorded a remarkable instance of a discovery made by Archimedes, both by *alligation* and specific gravity at the same time, namely, concerning the crown of Hiero, king of Syracuse. The king had ordered a crown to be made of pure gold, but when brought to him, a suspicion arose that it was mixed with alloy of either silver or copper, and the king recommended it to Archimedes to discover the cheat without defacing the crown. Archimedes, after long thinking on the matter, without lighting on the means of doing it, being one day in the bath, and observing how his body raised the water higher, conceived the idea that different metals of the same weight would occupy different spaces, and so raise or expel different quantities of water. Upon which he procured two other masses, each of the same weight with the crown, the one of pure gold, and the other of alloy; then immersing them all three, separately, in water, and observing the space each occupied, by the quantity it raised the water, he from thence computed the quantities of gold and alloy contained in the crown.

ALLIOTH, a star in the tail of the great bear. The word in Arabic denotes a horse; and they gave this name to each of the three stars, in the tail of the great bear, as they are placed like three horses, thus arranged for the purpose of drawing the waggon commonly called Charles's wain, represented by the four stars on the body of the same constellation.

ALMACANTAR. See ALMUCANTAR.

ALMAGEST, the name of a celebrated book composed by Ptolemy; being a collection of a great number of the observations and problems of the ancients, relating to geometry and astronomy, but especially the latter. And being the first work of this kind which has come down to us, and containing a catalogue of the fixed stars, with their places, beside numerous records and observations of eclipses, the motions of the planets, &c, this work will ever be held dear and valuable to the cultivators of astronomy.

In the original Greek it is called *συντάξις μεγίστη*, the *great composition* or *collection*. And to the word *μεγίστη*, *megiste*, the Arabians joined the particle *al*, and thence called it *Almagesti*, or, as we call it, from them, the *Almagest*.

Ptolemy was born about the year of Christ 69, and died in 147, and wrote this work, consisting of 13 books, at Alexandria in Egypt, where the Arabians found it on the capture of that kingdom. It was by them

translated out of Greek, into Arabic, by order of the caliph Alnaimon, about the year 827; and first into Latin about 1230, by favour of the emperor Frederic II. The Greek text however was not known in Europe till about the beginning of the 15th century, when it was brought from Constantinople, then taken by the Turks, by *George*, a monk of Trabezond, who translated it into Latin, which translation has several times been published.

Riccioli, an Italian jesuit, also published, in 1651, a body of Astronomy, which, in imitation of Ptolemy, he called *Almagestum Novum*, the *New Almagest*; being a large collection of ancient and modern observations and discoveries, in the science of Astronomy.

ALMAMON, caliph of Bagdat, a philosopher and astronomer in the beginning of the 9th century, he having ascended the throne in the year 814. He was son of Harun Al-Rashid, and grand-son of Almanfor. His name is otherwise written *Mamon*, *Almaon*, *Almamun*, *Alamoun*, or *Al-Maimon*. Having been educated with great care and with a love for the liberal sciences, he applied himself to cultivate and encourage them in his own country. For this purpose he requested the Greek emperors to supply him with such books on philosophy as they had among them; and he collected skilful interpreters to translate them into the Arabic language. He also encouraged his subjects to study them; frequenting the meetings of the learned, and assisting at their exercises and deliberations. He caused Ptolemy's *Almagest* to be translated in 827, by Isaac Ben-honain, and Thabet Ben-korah, according to Herbelot, but according to others by Sergius, and Alhazen, the son of Joseph. In his reign, and doubtless by his encouragement, an astronomer of Bagdat, named Habash, composed three sets of astronomical tables.

Almamon himself made many astronomical observations, and determined the obliquity of the ecliptic to be then  $23^{\circ} 35'$  (or  $23^{\circ} 33'$  in some manuscripts), but Vossius says  $23^{\circ} 51'$  or  $23^{\circ} 34'$ . He also caused skilful observers to procure proper instruments to be made, and to exercise themselves in astronomical observations; which they did accordingly at Shemasi in the province of Bagdat, and upon Mount Casius near Damas.

Under the auspices of Mamon also a degree of the meridian was measured on the plains of Sinjar or Sindgiar (or according to some Fingar), upon the borders of the Red Sea; by which the degree was found to contain  $56\frac{2}{3}$  miles, of 4000 coudees each, the coudee being a foot and a half: but it is not known what foot is here meant, whether the Roman, the Alexandrian, or some other. Riccioli makes this measure of the degree amount to 81 ancient Roman miles, which value answers to 62046 French toises; a quantity more than the true value of the degree by almost one-third.

Finally, Mamon revived the sciences in the East to such a degree, that many learned men were found, not only in his own time, but after him, in a country where the study of the sciences had been long forgotten. This learned king died near Tarsus in Cilicia, by having eaten too freely of some dates, on his return from a military expedition, in the year 833.

ALMANAC, a calendar or table, in which are set down and marked the days and feasts of the year, the common ecclesiastical notes, the course and phases of



the moon, &c, for each month: and answers to the *fasti* of the ancient Romans.

The etymology of the word is much controverted among grammarians.—Some derive it from the Arabic, viz, from the particle *al*, and *manab*, to count. While Scaliger, and others, derive it from the same *al*, and the Greek *μηναιος*, the course of the months. But Golius controverts these opinions, and ascribes the word to another origin, though he still makes it of Arabic extract, which it more evidently is. He says that, in the East it is the custom for the people, at the beginning of the year, to make presents to their princes; and that, among the rest, the astrologers present them with their almanacs, or ephemerides, for the year ensuing; whence these came to be called *almanba*, that is, new-year's gifts. But this derivation seems rather strained and improbable; for, by the same rule, the gifts or productions of other artists, or classes of men, might also be called almanacs. There are other guesses at the etymology; and Verstegan writes the word *almonac*, and makes it of German original. Our ancestors, he observes, used to carve the courses of the moon, for the whole year, upon a square piece of wood, which they called *al-monaght*, which is as much as to say, in old English or Saxon, *all-moon-beed*.

Almanacs are of various kinds and composition, some books, others sheets, &c, some annual, others perpetual. The essential part is the calendar of months, weeks, and days; the motions, changes, and phases of the moon; with the rising and setting of the sun and moon. To these are commonly added various matters, astronomical, astrological, chronological, meteorological, and even political, rural, medical, &c; as also eclipses, solar ingresses, aspects, and configurations of the heavenly bodies, lunations, heliocentric and geocentric motions of the planets, prognostications of the weather, and predictions of other events, the tides, twilight, equation of time, &c.

Till about the 4th century, almanacs bore the marks of heathenism only; from thence to the 7th century, they were a mixture of heathenism and christianity; and ever since they have been altogether christian: but at all times, astrological and other predictions have been considered as an essential part, and still are so to this day with several of them, notwithstanding that most people affect to disbelieve in such predictions.

*Nautical ALMANAC*, and *Astronomical Ephemeris*, is a kind of national almanac, chiefly for nautical purposes, which was begun in the year 1767 under the direction of the Board of Longitude, on the recommendation of the present worthy Astronomer Royal, who has the immediate conducting of it. It is still published by anticipation for several years before hand, for the convenience of ships going out upon long voyages, for which it is highly useful, and was found eminently so in the course of the late voyages round the world for making discoveries. Besides most things essential to general use, that are to be found in other almanacs, it contains many new and important particulars; more especially, the distances of the moon from the sun and fixed stars, which are computed for the meridian of the Royal Observatory of Greenwich, and set down to every three hours of time, expressly designed for computing the longitude at sea, by comparing these with the like distances observed there.

*ALMANAR*, in the Arabian astrology, denotes the pre-eminence or prevalence of one planet over another.

*ALMUCANTARS*, *ALMACANTARS*, or *ALMICANTARS*, from the Arabic *almocantharat*, are circles parallel to the horizon, conceived to pass through every degree of the meridian; serving to shew the height of the sun, moon, or stars, &c; and are the same as the parallels of altitude.

*ALMUCANTAR-Staff*, was an instrument formerly used at sea to observe the sun's amplitude at rising or setting, and thence to determine the variation of the compass, &c. The instrument had an arch of 15 degrees, made of some smooth wood.

*ALPHONSINE Tables*, are astronomical tables compiled by order of Alphonfus, king of Castile. In the compiling of these it is thought that prince himself assisted. See *Astronomical tables*.

*ALPHONSUS* the 10th, king of Leon and Castile, who has been surnamed The Wise, on account of his attachment to literature, and is now more celebrated for having been an astronomer than a king. He was born in 1203; succeeded his father Ferdinand the 3d, in 1252; and died in 1284, consequently at the age of 81.

The affairs of the reign of Alphonfus were very extraordinary and unfortunate for him. But we shall here only consider him in that part of his character, on account of which he has a place in this work, namely, as an astronomer and man of letters. He understood astronomy, philosophy, and history, as if he had been only a man of letters; and composed books upon the motions of the heavens, and on the history of Spain, which are highly commended. "What can be more surprising," says Mariana, "than that a prince, educated in a camp, and handling arms from his childhood, should have such a knowledge of the stars, of philosophy, and the transactions of the world, as men of leisure can scarcely acquire in their retirements? There are extant some books of Alphonfus on the motions of the stars, and the history of Spain, written with great skill and incredible care." In his astronomical pursuits he discovered that the tables of Ptolemy were full of errors; and thence he conceived the first of any the resolution of correcting them. For this purpose, about the year 1240, and during the life of his father, he assembled at Toledo the most skilful astronomers of his time, Christians, Moors, or Jews, when a plan was formed for constructing new tables. This task was accomplished about 1252, the first year of his reign; the tables being drawn up chiefly by the skill and pains of Rabbi Isaac Hazan a learned Jew, and the work called the *Alphonsine Tables*, in honour of the prince, who was at vast expences concerning them. He fixed the epoch of the tables to the 30th of May 1252, being the day of his accession to the throne. They were printed for the first time in 1483, at Venice, by Radtolt, who excelled in printing at that time; an edition extremely rare: there are others of 1492, 1521, 1545, &c. (Weidler, p. 280).


We must not omit a memorable saying of Alphonfus, which has been recorded for its boldness and pretended impiety; namely, "that if he had been of God's privy council when he made the world, he could have advised him better." Mariana however says only in general, that Alphonfus was so bold as to blame the works of Providence,



Providence, and the construction of our bodies; and he says that this story concerning him rested only upon a vulgar tradition. The Jesuit's words are curious: "Emanuel, the uncle of Sanchez (the son of Alphonsus), in his own name, and in the name of other nobles, deprived Alphonsus of his kingdom by a public sentence: which that prince merited, for daring severely and boldly to censure the works of divine Providence, and the construction of the human body, as tradition says he did. Heaven most justly punished the folly of his tongue." Though the silence of such an historian as Mariana, in regard to Ptolemy's system, ought to be of some weight, yet we cannot think it improbable, that if Alphonsus did pass so bold a censure on any part of the universe, it was on the celestial sphere, and meant to glance upon the contrivers and supporters of that system. For, besides that he studied nothing more, it is certain that at that time astronomers explained the motions of the heavens by intricate and confused hypotheses, which did no honour to God, nor anywise answered the idea of an able workman. So that, from considering the multitude of spheres composing the system of Ptolemy, and those numerous eccentric cycles and epicycles with which it is embarrassed, if we suppose Alphonsus to have said, "That if God had asked his advice when he made the world, he would have given him better council," the boldness and impiety of the censure will be greatly diminished.

ALSTED (JOHN-HENRY), a German protestant divine, and one of the most indefatigable writers of the 17th century. He was some time professor of philosophy and divinity at Herborn in the county of Nassau: from thence he went into Transilvania, to be professor at Alba Julia; where he continued till his death, which happened in 1638, being then 50 years of age. He applied himself chiefly to compose methods, and to reduce the several branches of arts and sciences into systems. His *Encyclopædia* has been much esteemed even by Roman Catholics; it was printed at Lyons, and sold very well throughout all France. Vossius mentions the *Encyclopædia* in general, but speaks of his treatise of *Arithmetic* more particularly, and allows the author to have been a man of great reading and universal erudition. His *Theſaurus Chronologicus* is by some esteemed one of his best works, and has gone through several editions, though others speak of it with contempt. In his *Triumphus Biblicus* Alsted endeavours to prove that the materials and principles of all the arts and sciences may be found in the scriptures; but he gained very few to his opinion. John Himmeliſius wrote a piece against his *Theologia Polemica*, which was one of Alsted's best performances. It seems he was a millenarian, having published, in 1672, a treatise *De Mille Annis*, in which he asserts that the faithful shall reign with Jesus Christ upon earth a thousand years; after which will be the general resurrection, and the last judgment; and he pretended that this reign would commence in the year 1694.

ALTERNATE angles, are the internal angles, A and B, or  $a$  and  $b$ , made by a line cutting two parallel lines, and lying on opposite sides of the cutting line. It is the property of these angles to be always equal to each other, namely the angle  $A =$  the



angle B, and the angle  $a =$  the angle  $b$ . And the exterior alternate angles are also equal.

ALTERNATE Ratio or Proportion, is the ratio of the one antecedent to the other, or of one consequent to the other, in any proportion, in which the quantities are of the same kind. So if  $A : B :: C : D$ , then alternately, or by alternation  $A : C :: B : D$ .

ALTERNATION, or Permutation, of quantities or things, is the varying or changing the order or position of them.

As suppose two things  $a$  and  $b$ ; these may be placed either thus  $ab$  or  $ba$  that is two ways, or  $1 \times 2$ . If there be three things,  $a, b, c$ , then the 3d thing  $c$ , may be placed three different ways with respect to each of the two positions  $ab$  and  $ba$  of the other two things, it may stand either before them, or between them, or after them both, that is, it may stand either 1st, 2d, or 3d; and therefore with three things there will be three times as many changes as with two, that is  $1 \times 2 \times 3$  or six changes with three things. Again, if there be four things  $a, b, c, d$ ; then the fourth thing  $d$  may be placed in four different ways with respect to each of the six positions of the other three; for it may be set either 1st or 2d or 3d or 4th in the order of each position; consequently from four things there will be four times as many alternations as there are from three things; and therefore  $1 \times 2 \times 3 \times 4 = 24$  is the number of changes with four things. And so on, always multiplying the last found number of alternations by the next number of things; or to find the number of changes for any number of things, as  $n$ , multiply the series of natural numbers 1, 2, 3, 4, 5, &c, to  $n$ , continually together, and the last product will be the number of alternations sought; so  $1 \times 2 \times 3 \times 4 \times 5 \dots n$  is the number of changes in  $n$  things.

So if, for example, it were required to find how many changes may be rung on 12 bells; it would be  $1 \times 2 \times 3 \times 4 \times 5 \times 6 \times 7 \times 8 \times 9 \times 10 \times 11 \times 12 = 479001600$ , the number of changes. Now supposing there might be rung 10 changes in one minute, that is  $10 \times 12$  or 120 strokes in a minute, or 2 strokes in each second of time; then, according to this rate, it would take upwards of 91 years to ring over all these changes on the 12 bells only. Also, if but two more bells were added, making 14 bells; then, at the same rate of ringing, it would require about 16575 years to ring all the changes on 14 bells but once over. And if the number of bells were 24, it would require more than 11700000000000000 years to ring all the different changes upon them!

ALTIMETRY, ALTIMETRIA, the art of taking or measuring altitudes or heights, whether accessible or inaccessible. Or

Altimetria is the part of practical Geometry which respects the theory and practice of measuring both heights and depths, and both in respect of perpendicular and oblique lines.

ALTING (JAMES), was born at Heidelberg in 1618. He travelled into England in 1640, where he was ordained by the learned Dr. Prideaux, bishop of Worcester. He afterwards succeeded Gomarus in the professorship of Groningen. He died in 1697; and recommended the edition of his works to Menſio Alting (author of *Notitia German. Infer. Antiquæ*); but they were published in 5 vols folio, with his life, by Bekker



Bekker of Amsterdam. They contain various analytical, exegetical, practical, problematical, and philosophical tracts, which shew his great industry and knowledge.

**ALTITUDE**, in Geometry is the third dimension of body, considered with respect to its elevation above the ground: and is otherwise called its height when measured from bottom to top, or its depth when measured from top to bottom.

**ALTITUDE** of a figure, is the distance of its vertex from the base, or the length of a perpendicular let fall from its vertex to the base. The altitudes of figures are useful in computing their areas or solidities.

**ALTITUDE**, or *Height* of any point of a terrestrial object, is the perpendicular let fall from that point to the plane of the horizon. *Altitudes* are distinguished into *accessible* and *inaccessible*.

*Accessible ALTITUDE* of an object, is that whose base there is access to, to measure the nearest distance to it on the ground, from any place.


*Inaccessible ALTITUDE*, of an object, is that whose base there is not free access to, by which a distance may be measured to it, by reason of some impediment, such as water, wood, or the like.

*To measure or take ALTITUDES.* If an altitude cannot be measured by stretching a string from top to bottom, which is the direct and most accurate way, then some indirect way is used, by actually measuring some other line or distance which may serve as a basis, in conjunction with some angles, or other proportional lines, either to compute, or geometrically determine, the altitude of the object sought.

There are various ways of measuring altitudes, or depths, by means of different instruments, and by shadows or reflected images, on optical principles. There are also various ways of computing the altitude in numbers, from the measurements taken as above, either by geometrical construction, or trigonometrical calculation, or by simple numeral computation from the property of parallel lines, &c.


The instruments mostly used in measuring altitudes, are the quadrant, theodolite, geometrical square, line of shadows, &c; the descriptions of each of which may be seen under their respective names.

*To measure an Accessible Altitude Geometrically.* Thus, suppose the height of the accessible tower AB be required. First, by means of two rods, the one longer than the other: plant the longer upright at C; then move the shorter back from it, till by trials you find such a place, D, that the eye placed at the top of it at E, may see the top of the other, F, and the top of the object B straight in a line: next measure the distances DA or EG and DC or EH, also HF the difference between the heights of the rods: then, by similar triangles, as EH: EG:: HF: the 4th proportional GB; to which add AG or DE, and the sum will be the whole altitude AB sought.




Or, with one rod CF only: plant it at such a place C, that the eye at the ground, or near it, at I, may see the tops F and B in a right line: then, having measured IC, IA, CF, the 4th proportional to these will be the altitude AB sought.


Or thus, by means of Shadows. Plant a rod *ab* at *a*, and measure its shadow *ac*, as also the shadow AC of the object AB; then the 4th proportional to *ac*, *ab*, AC will be the altitude AB sought.



Or thus, by means of Optical Reflection. Place a vessel of water, or a mirror or other reflecting surface, horizontal at C; and move off from it to such a distance, D, that the eye E may see the image of the top of the object in the mirror at C: then, by similar figures, CD: DE:: CA: AB the altitude sought.



Or thus, by the Geometrical Square. At any place, C, fix the stand, and turn the square about the centre of motion, D, till the eye there see the top of the object through the sights or telescope on the side DE of the quadrant, and note the number of divisions cut off the other side by the plumb line EG: then as EF: FG:: DH: HB; to which add AH or CD, for the whole height AB.




*To measure an Accessible Altitude Trigonometrically.* At any convenient station, C, with a quadrant, theodolite, or other graduated instrument, observe the angle of elevation ACB above the horizontal line AC; and measure the distance AC. Then, A being a right angle, it will be, as radius is to the tangent of the angle A, so is AC to AB sought.

If AC be not horizontal, but an inclined plane; then the angle above it must be observed at two stations C and D in a right line, and the distances AC, CD both measured. Then, from the angle C take the angle D, and there remains the angle CBD; hence in the triangle BCD, are given the angles and the side DC, to find the side CB; and then in the triangle ABC, are given the

two




two sides CA and CB, with the included angle C, to find the third side AB.




Or thus, measure only the distance AC, and the angles A and C: then, in the triangle ABC, are given all the angles and the side AC, to find the side AB.

To measure an Inaccessible Altitude, as a hill, cloud, or other object. This is commonly done, by observing the angle of its altitude at two stations, and measuring the distance between them. Thus, for the height AB of a hill, measure the distance CD at the foot of it, and observe the quantity of the two angles C and D. Then, from the angle C taking the angle D, leaves the angle CBD; hence

As sine  $\angle CBD$  : sine  $\angle D$  :: CD : CB; and  
As rad. : sine  $\angle ACB$  :: CB : AB the altitude.



And for a balloon, or cloud, or other moveable object C, let two observers at A and B, in a plane with C, take at the same time the angles A and B, and measure the distance between them AB; then calculate the altitude CD exactly as in the last example.



To find the height of an object, by knowing the utmost distance at which its top can be just seen in the horizon. As suppose the top H of a tower FH can be just seen from E when the distance EF is 25 miles, supposing the circumference of the earth to be 25000 miles, or the radius 3979 miles or 21009120 feet. First, as  $25000 : 25 :: 360^\circ : 21' 36''$  equal to the angle G; then as radius : sec.  $\angle G$  :: EG : GH, which will be found to be 21009536 feet; from which take EG or GF, and there remains 416 feet, for FH the height of the tower sought.— Or rather thus, as 10000000 radius : 198 = sec.  $\angle G$  — radius :: 21009120 = EG : 416 = FH, as before.

Or the same may be found easier thus: The horizon dips nearly 8 inches or  $\frac{2}{3}$  of a foot at the distance of 1 mile, and according to the square of the distance for

other distances; therefore as  $1^2$  or  $1 : 25^2$  or  $625 :: \frac{2}{3} : \frac{2}{3}$  of 625 or 416 feet, the same as before.

There is a very easy method of taking great terrestrial altitudes, such as mountains &c, by means of the difference between the heights of the barometer observed at the bottom and top of the same. Which see under the article BAROMETER.

ALTITUDE of the Eye, in Perspective, is a right line let fall from the eye, perpendicular to the geometrical plane.



ALTITUDE, in Astronomy, is the arch of a vertical circle, measuring the height of the sun, moon, star, or other celestial object, above the horizon.

This altitude may be either true or apparent. The apparent altitude is that which appears by sensible observations made at any place on the surface of the earth. And the true altitude is that which results by correcting the apparent, on account of refraction and parallax.

The quantity of the refraction is different at different altitudes; and the quantity of the parallax is different according to the distance of the different luminaries: in the fixed stars this is too small to be observed; in the sun it is but about  $8\frac{1}{4}$  seconds; but in the moon it is about 52 minutes.

Altitudes are observed by a quadrant, or sextant, or by the shadow of a gnomon or high pole, and by various other ways, as may be seen in most books of astronomy.

Meridian ALTITUDE, is an arch of the meridian intercepted between any point in it and the horizon. So if HO be the horizon, and HEZO the meridian; then the arch HE, or the angle HCE, is the meridian altitude of an object in the meridian at the point E.



ALTITUDE, or elevation, of the Pole, is the angle OCP, or arch OP of the meridian, intercepted between the horizon and pole P.

This is equal to the latitude of the place; and it may be found by observing the meridian altitude of the pole star, when it is both above and below the pole, and taking half the sum, when corrected on account of refraction. Or the same may be found by the declination and meridian altitude of the sun.

ALTITUDE, or elevation, of the equator, is the angle HCE, or arch HE of the meridian, between the horizon and the equator at E; and it is equal to ZP the co-latitude of the place.

ALTITUDE of the Tropics, the same as what is otherwise called the solstitial altitude of the sun, or his meridian altitude when in the solstitial points.

ALTITUDE, or height, of the horizon, or of stars &c seen in it, is the quantity by which it is raised by refraction.

Refraction of ALTITUDE, is an arch of a vertical circle, by which the true altitude of the moon, or a star, or other object, is increased by means of the refraction; and



and is different at different altitudes, being nothing in the zenith, and greatest at the horizon, where it is about  $33'$ .

*Parallax of ALTITUDE*, is an arch of a vertical circle, by which the true altitude, observed at the centre of the earth, exceeds that which is observed on the surface; or the difference between the angles  $\angle LM$  and  $\angle IK$  of altitude there; and is equal to the angle  $\angle I \angle L$  formed at the moon or other body, and subtended by the radius  $IL$  of the earth.

It is evident that this angle is less, as the luminary is farther distant from the earth; and also less, for any one luminary, as it is higher above the horizon; being greatest there, and nothing in the zenith.

*ALTITUDE of the Nonagesimal*, is the altitude of the 90th degree of the ecliptic, counted upon it from where it cuts the horizon, or of the middle or highest point of it which is above the horizon, at any time; and is equal to the angle made by the ecliptic and horizon where they intersect at that time.

*ALTITUDE of the cone of the earth's or moon's shadow*, the height of the shadow of the body, made by the sun, and measured from the centre of the body. To find it, say, As the tangent of the angle of the sun's apparent semidiameter is to radius, so is 1 to a 4th proportional, which will be the height of the shadow, in semidiameters of the body.

So, the greatest height of the earth's shadow, is  $217.8$  semidiameters of the earth, when the sun is at his greatest distance, or his semidiameter subtends an angle of about  $15' 47''$ ; and the height of the same is  $210.7$  semidiameters of the earth, when the sun is nearest the earth, or when his semidiameter is about  $16' 19''$ : And proportionally between these limits for the intermediate distances or semidiameters of the sun.

The altitudes of the shadows of the earth and moon, are nearly as 11 to 3, the proportion of their diameters.

*ALTITUDE*, or *exaltation*, in astrology, denotes the second of the five essential dignities, which the planets acquire by virtue of the signs they are found in.

*ALTITUDE of motion*, is a term used by Dr. Wallis, for the measure of any motion, estimated in the line of direction of the moving force.

*ALTITUDE*, in speaking of fluids, is more frequently expressed by the term *depth*. The pressure of fluids, in every direction, is in proportion to their altitude or depth.

*ALTITUDE of the mercury*, in the *barometer* and *thermometer*, is marked by degrees, or equal divisions, placed by the side of the tube of those instruments.

The altitude of the barometer, or of the mercury in its tube, at London, is usually comprised between the limits of 28 and 31 inches; and the mean height, for every day in several years, is nearly  $29.87$  inches.


*ALTITUDE of the pyramids in Egypt*, was measured so long since as the time of Thales, which he effected by means of their shadow, and that of a pole set upright beside them, making the altitudes of the pole and pyramid proportional to the lengths of their shadows. Plutarch has given an account of the manner of this operation, which is one of the first geometrical observations we have an exact account of.

*ALTITUDE*, *circles of*, *parallels of*, *quadrant of*, &c. See the respective words.

*Equal ALTITUDE Instrument*, is an instrument used to observe a celestial object, when it has the same or an equal altitude, on both sides of the meridian, or before and after it passes the meridian: an instrument very useful in adjusting clocks &c, and for comparing equal and apparent time.

*AMBIENT*, encompassing round about; as the bodies which are placed about any other body, are called ambient bodies, and sometimes *circum-ambient* bodies; and the whole mass of the air or atmosphere, because it encompasses all things on the face of the earth, is called the *ambient air*.

*AMBIGENAL Hyperbola*, a name given by Newton, in his *Enumeratio linearum tertii ordinis*, to one of the triple hyperbolas  $EGF$  of the second-order, having one of its infinite legs, as  $EG$ , falling within the angle  $ACD$ , formed by the asymptotes  $AC$ ,  $CD$ , and the other leg  $GF$  falling without that angle.



*AMBIT*, of a figure, in Geometry, is the perimeter, or line, or sum of the lines, by which the figure is bounded.

*AMBLIGON*, or *AMBLIGONAL*, in Geometry, signifies obtuse-angular, as a triangle which has one of its angles obtuse, or consisting of more than 90 degrees.

*AMICABLE numbers*, denote pairs of numbers, of which each of them is mutually equal to the sum of all the aliquot parts of the other. So the first or least pair of amicable numbers are 220 and 284; all the aliquot parts of which, with their sums, are as follow, viz, of 220, they are 1, 2, 4, 5, 10, 11, 20, 22, 44, 55,

110, their sum ————— 284;

of 284, they are 1, 2, 4, 71, 142, and their sum is 220.

The 2d pair of amicable numbers are 17296 and 18416, which have also the same property as above.

And the 3d pair of amicable numbers are 9363584 and 9437056.

These three pairs of amicable numbers were found out by F. Schooten, sect. 9 of his *Exercitationes Mathematicæ*, who I believe first gave the name of *amicable* to such numbers, though such properties of numbers it seems had before been treated of by Rudolphus, Descartes, and others.

To find the first pair, Schooten puts  $4x$  and  $4yz$ , or  $a^2x$  and  $a^2yz$  for the two numbers where  $a = 2$ ; then making each of these equal to the sum of the aliquot parts of the other, gives two equations, from which are found the values of  $x$  and  $z$ , and consequently, assuming a proper value for  $y$ , the two amicable numbers themselves  $4x$  and  $4yz$ .

In like manner for the other pairs of such numbers; in which he finds it necessary to assume  $16x$  and  $16yz$  or  $a^4x$  and  $a^4yz$  for the 2d pair, and  $128x$  and  $128yz$  or  $a^7x$  and  $a^7yz$  for the 3d pair.



Schooten then gives this practical rule, from Descartes, for finding amicable numbers, viz; Assume the number 2, or some power of the number 2, such that if unity or 1 be subtracted from each of these three following quantities; viz;

from 3 times the assumed number,

also from 6 times the assumed number,

and from 18 times the square of the assumed number, the three remainders may be all prime numbers; then the last prime number being multiplied by double the assumed number, the product will be one of the amicable numbers sought, and the sum of its aliquot parts will be the other.

That is, if  $a$  be put = the number 2, and  $n$  some integer number, such that  $3a^n - 1$ , and  $6a^n - 1$ , and  $18a^{2n} - 1$  be all three prime numbers; then is  $18a^{2n} - 1 \times 2a^n$  one of the amicable numbers; and the sum of its aliquot parts is the other.

AMONTONS (WILLIAM), an ingenious French experimental philosopher, was born in Normandy the 31st of August 1663. While at the grammar school, he by sickness contracted a deafness that almost excluded him from the conversation of mankind. In this situation he applied himself to the study of geometry and mechanics; with which he was so delighted that it is said he refused to try any remedy for his disorder, either because he deemed it incurable, or because it increased his attention to his studies. Among other objects of his study, were the arts of drawing, of land-surveying, and of building; and shortly after he acquired some knowledge of those more sublime laws by which the universe is regulated. He studied with great care the nature of barometers and thermometers; and wrote his treatise of *Observations and Experiments concerning a new Hour-glass, and concerning Barometers, Thermometers, and Hygrosopes*; as also some pieces in the *Journal des Savans*. In 1687, he presented a new hygroscope to the Academy of Sciences, which was much approved. He found out a method of conveying intelligence to a great distance in a short space of time: this was by making signals from one person to another, placed at as great distances from each other as they could see the signals by means of telescopes. When the Royal Academy was new regulated in 1699, Amontons was chosen a member of it, as an eleve under the third Astronomer; and he read there his *New Theory of Friction*, in which he happily cleared up an important object in mechanics. In fact he had a particular genius for making experiments: his notions were just and delicate: and he knew how to prevent the inconveniences of his new inventions, and had a wonderful skill in executing them. He died of an inflammation in his bowels, the 11th of October 1705, being only 42 years of age.

The eloge of Amontons may be seen in the volume of the *Memoirs of the Academy of Sciences* for the year 1705, Hist. pa. 150. And his pieces contained in the different volumes of that work, which are pretty numerous, and upon various subjects, as the air, action of fire, barometers, thermometers, hygrometers, friction, machines, heat, cold, rarefactions, pumps, &c, may be seen in the volumes for the years 1696, 1699, 1702, 1703, 1704, and 1705.

AMPHISCII, or AMPHISCIAANS, are the people who inhabit the torrid zone; which are so called, because

VOL. I.

they have their shadow at noon turned sometimes one way, and sometimes another, namely, at one time of the year towards the north, and at the other towards the south.

AMPLITUDE, in *gunnery*, the range of the projectile, or the right line upon the ground subtending the curvilinear path in which it moves.

AMPLITUDE, in *astronomy*, is an arch of the horizon, intercepted between the true east or west point, and the centre of the sun or a star at its rising or setting: so that the amplitude is of two kinds; *ortive* or eastern, and *occiduous* or western. Each of these amplitudes is also either northern or southern, according as the point of rising or setting is in the northern or southern part of the horizon: and the complement of the amplitude, or the arch of distance of the point of rising or setting, from the north or south point of the horizon, is the azimuth.

The amplitude is of use in navigation, to find the variation of the compass or magnetic needle. And the rule to find it is this: As the cosine of the latitude is to radius, so is the sine of the sun's or star's declination, to the sine of the amplitude. So in the latitude of London, viz,  $51^{\circ} 31'$ , when the sun's declination is  $23^{\circ} 28'$ ; then

cos. $51^{\circ} 31'$ the lat.	- - -	9.7939907
fin. $23^{\circ} 28'$ the decl.	-	+ 9.6001181

fin. $39^{\circ} 47'$ the ampl.	9.8061274
---------------------------------	-----------

That is, the sun then rises or sets  $39^{\circ} 47'$  from the east or west point, to the north or south according as the declination is north or south.

Magnetical AMPLITUDE, is an arch of the horizon, contained between the sun or star, at the rising or setting, and the magnetical east or west point of the horizon, pointed out by the magnetical compass, or the amplitude or azimuth compass. And the difference between this magnetical amplitude, so observed, and the true amplitude, as computed in the last article, is the variation of the compass.

So if, for instance, the magnetical amplitude be observed, by the compass, to be  $61^{\circ} 47'$ , at the time when it is computed to be

-	39 47,
---	--------

then the difference - - 22 0 is the variation west.

ANABIBAZON, a name sometimes given to the dragon's tail, or northern node of the moon.

ANACAMPTICS, or the science of the reflections of sounds, frequently used in reference to echoes, which are said to be sounds produced *anacamptically*, or by reflection. And in this sense it was used by the ancients for that part of optics which is otherwise called Catoptrics.

ANACHRONISM, in Chronology, an error in computation of time, by which an event is placed earlier than it really happened. Such is that of Virgil, who makes Dido to reign at Carthage in the time of Æneas, though, in reality, she did not arrive in Africa till 300 years after the taking of Troy.

An error on the other side, by which a fact is placed later, or lower than it should be, is called a *parachronism*. But in common use, this distinction, though proper, is not attended to; and the word *anachronism* is used indifferently for the mistake on both sides.

ANACLASTICS, or ANACLATICS, an ancient name for that part of Optics which considers refracted light;

P

being



being the same as what is more usually called *dioptrics*. See the Compendium of *Ambrosius Rhodius*, lib. 3. *Optica*, pa. 384 & seq.

**ANACLASTIC Curves**, a name given by M. de Mairan to certain apparent curves formed at the bottom of a vessel full of water, to an eye placed in the air; or the vault of the heavens, seen by refraction through the atmosphere.

M. de Mairan determines these curves by a principle not admitted by all authors; but Dr. Barrow, at the end of his *Optics*, determines the same curves by other principles.

**ANALEMMA**, a planisphere, or projection of the sphere, orthographically made on the plane of the meridian, by perpendiculars from every point of that plane, the eye supposed to be at an infinite distance, and in the east or west point of the horizon. In this projection, the solstitial colure, and all its parallels, are projected into concentric circles, equal to the real circles in the sphere; and all circles whose planes pass through the eye, as the horizon and its parallels, are projected into right lines equal to their diameters; but all oblique circles are projected into ellipses, having the diameter of the circle for the transverse axis.

This instrument, having the furniture drawn on a plate of wood or brass, with an horizon fitted to it, is used for resolving many astronomical problems; as the time of the sun's rising and setting, the length and hour of the day, &c. It is also useful in dialling, for laying down the signs of the zodiac, with the lengths of days, and other matters of furniture, upon dials.

The oldest treatise we have on the analemma, was written by Ptolemy, which was printed at Rome in 1562, with a commentary by F. Commandine. Pappus also treated of the same. Since that time, many other authors have treated very well of the analemma; as Aguilonius, Taquet, Dechales, Witty, &c.

**ANALOGY**, the same as proportion, or equality, or similitude of ratios. Which see.

**ANALYSIS**, is, generally, the resolution of any thing into its component parts, to discover the thing or the composition. And in mathematics it is properly the method of resolving problems, by reducing them to equations. *Analysis* may be distinguished into the *ancient* and the *modern*.

The *ancient analysis*, as described by Pappus, is the method of proceeding from the thing sought as *taken* for granted, through its consequences, to something that is *really* granted or known; in which sense it is the reverse of synthesis or composition, in which we lay that down *first* which was the *last* step of the analysis, and tracing the steps of the analysis back, making that antecedent here which was consequent there, till we arrive at the thing sought, which was taken or assumed as granted in the first step of the analysis. This chiefly respected geometrical enquiries.

The principal authors on the ancient analysis, as recounted by Pappus, in the 7th book of his *Mathematical Collections*, are Euclid in his *Data*, *Porismata*, & *de Locis ad Superficiem*; Apollonius *de Sectione Rationis*, *de Sectione Spatii*, *de Tactionibus*, *de Inclinationibus*, *de Locis Planis*, & *de Sectionibus Conicis*; Aristæus, *de Locis Solidis*; and Eratosthenes, *de Mediis Proportionalibus*: from which

Pappus gives many examples in the same book. To these authors we may add Pappus himself. The same sort of analysis has also been well cultivated by many of the moderns; as Fermat, Viviani, Getaldus, Snellius, Huygens, Simson, Stewart, Lawson, &c, and more especially Hugo d'Omerique, in his *Analysis Geometrica*, in which he has endeavoured to restore the Analysis of the ancients. And, on this head, Dr. Pemberton tells us "that Sir Isaac Newton used to censure himself for not following the ancients more closely than he did; and spoke with regret of his mistake, at the beginning of his mathematical studies, in applying himself to the works of Descartes, and other algebraical writers, before he had considered the Elements of Euclid with that attention so excellent a writer deserves: that he highly approved the laudable attempt of Hugo d'Omerique to restore the ancient analysis."


In the application of the ancient analysis in geometrical problems, every thing cannot be brought within strict rules; nor any invariable directions given, by which we may succeed in all cases; but some previous preparation is necessary, a kind of mental contrivance and construction, to form a connexion between the *data* and *quæsitæ*, which must be left to every one's fancy to find out; being various, according to the various nature of the problems proposed: Right lines must be drawn in particular directions, or of particular magnitudes; bisecting perhaps a given angle, or perpendicular to a given line; or perhaps tangents must be drawn to a given curve, from a given point; or circles described from a given centre, with a given radius, or touching given lines, or other given circles; or such-like other operations. Whoever is conversant with the works of Archimedes, Apollonius, or Pappus, well knows that they founded their analysis upon some such previous operations; and the great skill of the analyst consists in discovering the most proper affections on which to found his analysis: for the same problem may often be effected in many different ways: of which it may be proper to give here an example or two. Let there be taken, for instance, this problem, which is the 155th prop. of the 7th book of Pappus.

From the extremities of the base A, B, of a given segment of a circle, it is required to draw two lines AC, BC, meeting at a point C in the circumference, so that they shall have a given ratio to each other, suppose that of F to G.

The solution of this problem, as given by Pappus, is thus.

#### ANALYSIS.

Suppose the thing done, and that the point C is found: then suppose CD is drawn a tangent to the circle at C, and meeting the line AB produced in the point D. Now by the hypothesis  $AC : BC :: F : G$ , and also  $AC^2 : BC^2 :: DA : DB$ , as may be thus proved.



Since DC touches the circle, and BC cuts it, the angle BCD is equal to BAC by Euc. iii. 32; also the angle D is



D is common to both the triangles DCA, DCB; these are therefore similar, and so, by vi 4,  $DA : DC :: DC : DB$ , and hence  $DA^2 : DC^2 :: DA : DB$  by cor. vi 20. But also, by vi 4,  $DA : AC :: DC : CB$ , and by permutation  $DA : DC :: AC : BC$ , or  $DA^2 : DC^2 :: AC^2 : BC^2$ ; and hence, by equality,  $AC^2 : BC^2 :: DA : DB$ .

But the ratio of  $AC^2$  to  $BC^2$  is given by prop. LVII of Simson's edition of the *Data*, because the ratio of AC to BC is given, and consequently that of DA to DB is given. Now since the ratio of DA to DB is given, therefore also, by Data vi, that of DA to AB, and hence, by Data ii, DA is given in magnitude.

And here the analysis properly ends. For it having been shewn that DA is given, or that a point D may be found in AB produced, such, that a tangent being drawn from it to the circumference, the point of contact will be the point sought; we may now begin the composition, or synthetical demonstration; which must be done by finding the point D, or laying down the line AD, which, it was affirmed, was given, in the last step of the analysis.

## SYNTHESIS.

*Construction.* Make as  $F^2 : G^2 :: AD : DB$ , (which may be done, since AB is given, by making it as  $F^2 : G^2 :: AB : DB$ , and then by composition it will be as  $F^2 : G^2 :: AD : DB$ ); and then from the point D, thus found, draw a tangent to the circle, and from the point of contact C drawing CA and CB, the thing is done.

*Demonstration.* Since, by the constr.  $F^2 : G^2 :: AD : DB$ , and also  $AD : DB :: AC^2 : BC^2$ , which has been already demonstrated in the analysis, and might be here proved in the same manner. Therefore  $F^2 : G^2 :: AC^2 : BC^2$ , and consequently  $F : G :: AC : BC$ . Q. E. D.

Here we see an instance of the method of *resolution* and *composition*, as it was practised by the ancients, the solution here given being that of Pappus himself. But as the method of referring and reducing every thing to the *Data*, and constantly quoting the same, may appear now to be tedious and troublesome: and indeed it is unnecessary to those who have already made themselves masters of the substance of that valuable book of Euclid, and have by practice and experience acquired a facility of reasoning in such matters: I shall therefore now shew how we may abate something of the rigour and strict form of the ancient method of solution, without diminishing any part of its admirable elegance and perspicuity. And this may be done by the instance of another solution, of the many more which might be given, of the same problem, as follows.

## ANALYSIS.

Let us again suppose that the thing is done, viz  $AC : BC :: F : G$ , and let there be drawn BH making the angle ABH equal to the angle ACB, and meeting AC produced in H. Then, the angle A being also common, the two triangles ABC and ABH are equiangular, and therefore, by vi 4,  $AC : BC :: AB : BH$ , in a given ratio; and, AB being given, therefore BH is given in position and magnitude.

## SYNTHESIS.

*Construction.* Draw BH making the angle ABH equal to that which may be contained in the given segment, and take AB to BH in the given ratio of F to G. Draw ACH, and BC.

*Demonstration.* The triangles ABC, ABH are equiangular, therefore, vi 4,  $AC : CB :: AB : BH$ , which is the given ratio by construction.

*Modern ANALYSIS*, consists chiefly of algebra, arithmetic of infinites, infinite series, increments, fluxions, &c; of each of which a particular account may be seen under their respective articles.

These form a kind of arithmetical and symbolical analysis, depending partly on modes of arithmetical computation, partly on rules peculiar to the symbols made use of, and partly on rules drawn from the nature and species of the quantities they represent; or from the modes of their existence or generation.

The modern analysis is a general instrument by which the finest inventions and the greatest improvements have been made in mathematics and philosophy, for near two centuries past. It furnishes the most perfect examples of the manner in which the art of reasoning should be employed; it gives to the mind a wonderful skill for discovering things unknown, by means of a small number that are given; and by employing short and easy symbols for expressing ideas, it presents to the understanding things which otherwise would seem to lie above its sphere. By this means geometrical demonstrations may be greatly abridged: a long train of arguments, in which the mind cannot, without the greatest effort of attention, discover the connection of ideas, is converted into visible symbols; and the various operations which they require, are simply effected by the combination of those symbols. And, what is still more extraordinary, by this artifice, a great number of truths are often expressed in one line only: instead of which, by following the ordinary way of explanation and demonstration, the same truths would occupy whole pages or volumes. And thus, by the bare contemplation of one line of calculation, we may understand in a short time whole sciences, which otherwise could hardly be comprehended in several years.

It is true that Newton, who best knew all the advantages of analysis in geometry and other sciences, laments, in several parts of his works, that the study of the ancient geometry is abandoned or neglected. And indeed the method employed by the ancients in their geometrical writings, is commonly regarded as more rigorous, than that of the modern analysis: and though it be greatly inferior to that of the moderns, in point of dispatch and facility of invention; it is nevertheless highly useful in strengthening the mind, improving the reasoning faculties, and in accustoming the young mathematician to a pure, clear, and accurate mode of investigation and demonstration, though by a long and laboured process, which he would with difficulty have submitted to if his taste had before been vitiated, as it were, by the more piquant sweets of the modern analysis. And it is principally on this that the complaints of Newton are founded, who feared lest by the too early and frequent use of the modern analysis, the science of geometry should lose that rigour and purity which characterise its investigations, and the mind become debilitated by the facility



facility of our analysis. This great man was therefore well founded, in recommending, to a certain extent, the study of the ancient geometricians: for, their demonstrations being more difficult, give more exercise to the mind, accustom it to a closer application, give it a greater scope, and habituate it to patience and resolution, so necessary for making discoveries. But this is the only or principal advantage from it; for if we should look no farther than the method of the ancients, it is probable that, even with the best genius, we should have made but few or small discoveries, in comparison of those obtained by means of the modern analysis. And even with regard to the advantage given to investigations made in the manner of the ancients, namely of being more rigorous, it may perhaps be doubted whether this pretension be well founded. For to instance in those of Newton himself, although his demonstrations be managed in the manner of the ancients; yet at the same time it is evident that he investigates his theorems by a method different from that employed in the demonstrations, which are commonly analytical calculations, disguised by substituting the name of lines for their algebraical value: and though it be true that his demonstrations are rigorous, it is no less so that they would be the same when translated and delivered in algebraic language; and what difference can it make in this respect, whether we call a line AB, or denote it by the algebraic character  $a$ ? Indeed this last designation has this peculiarity, that when all the lines are denoted by algebraic characters, many operations can be performed upon them, without thinking of the lines or the figure. And this circumstance proves of no small advantage: the mind is relieved, and spared as much as possible, that its whole force may be employed in overcoming the natural difficulty of the problem alone.

Upon the whole therefore the state of the comparison seems to be this; That the method of the ancients is fittest to begin our studies with, to form the mind and to establish proper habits; and that of the moderns to succeed, by extending our views beyond the present limits, and enabling us to make new discoveries and improvements.

*Analysis* is divided, with respect to its object, into that of *finites*, and that of *infinites*.

*Analysis of finite quantities*, is what is otherwise called *algebra*, or *specious arithmetic*.

*Analysis of infinites*, called also the *new analysis*, is that which is concerned in calculating the relations of quantities which are considered as infinite, or infinitely little; one of its chief branches being the *method of fluxions*, or the *differential calculus*. And the great advantage of the modern mathematicians over the ancients, arises chiefly from the use of this modern analysis.

*ANALYSIS of powers*, is the same as resolving them into their roots, and is otherwise called *evolution*.

*ANALYSIS of curve lines*, shews their constitution, nature and properties, their points of inflexion, station, retrogradation, variation, &c.

*ANALYST*, a person who analyses something, or makes use of the analytical method. In mathematics, it is a person skilled in algebra, or in the mathematical analysis in general.

*ANALYST*, the title of an ingenious, though sophistical book, written by the celebrated Dr. Berkeley, against the doctrine of fluxions.


*ANALYTIC*, or *ANALYTICAL*, something belonging to, or partaking of, the nature of analysis; or performed by the method of analysis.

Thus we say *analytical demonstration*, *analytical enquiry*, *analytical table or scheme*, *analytical method*, &c. The *analytical* stands opposed to the *synthetical*, or that which proceeds by the way of *synthesis*.

*ANALYTICS*, the science, or doctrine, and use of analysis.

*ANAMORPHOSIS*, in perspective and painting, a monstrous projection; or a representation of some image, either on a plane or curve surface, deformed or distorted; but which in a certain point of view shall appear regular, and drawn in just proportion.

*To construct an Anamorphosis, or monstrous projection, on a plane.*—Draw the square ABCD (fig. 1), of any size at pleasure, and divide it by crossing lines into a number of areolæ or smaller squares: and then in this square, or reticle, called also the *craticular prototype*, draw the regular image which is to be distorted.—Or, about any image, proposed to be distorted, draw a reticle of small squares.



Then draw the line  $ab$  (fig. 2.) equal to  $AB$ , dividing it into the same number of equal parts, as the side of the prototype  $AB$ ; and on its middle point  $E$  erect the perpendicular  $EV$ , and also  $VS$  perpendicular to  $EV$ , making  $EV$  so much the longer, and  $VS$  so much the shorter, as it is intended the image shall be more distorted. From each of the points of division draw right lines to the point  $V$ , and draw the right line  $aS$ . Lastly through the points  $c, e, f, g$ , &c, draw lines parallel to  $ab$ : So shall  $abcd$  be the space upon which the monstrous projection is to be drawn; and is called the *craticular ectype*.

Then, in every areola, or small trapezium, of the space  $abcd$ , draw what appears contained in the corresponding areola of the original space  $ABCD$ : so shall there be produced a deformed image in the space  $abcd$ , which yet will appear in just proportion to an eye distant from




from it the length of EV, and raised above it by a height equal to VS.

It will be amusing to contrive it so, that the deformed image may not represent a mere chaos, but some certain figure: thus, a river with soldiers, waggons, and other objects on the side of it, have been so drawn and distorted, that when viewed by an eye at S, it appeared like the face of a satyr.

An image may also be distorted mechanically, by perforating through in several places with a fine pin; then, placing it against a candle or lamp, observe where the rays, which pass through these small holes, fall on any plane or curve superficies; for they will give the correspondent points of the image deformed, and by means of which the deformation may be completed.

To draw an *Anamorphosis* upon the convex surface of a cone. It appears from the construction above, that we have only to make a craticular ectype upon the surface of the cone, which may appear equal to the craticular prototype, to an eye placed at a proper height above the vertex of the cone. Hence,

Let the base, or circumference, ABCD, of the cone (fig. 3) be divided by radii into any number of equal parts; and let some one radius be likewise divided into equal parts; then through each point of division draw concentric circles: so shall the craticular prototype be formed.



With double the diameter AB, as a radius, describe the quadrant EFG (fig. 4) so as the arch EG be equal to the whole periphery; then this quadrant, being plied or bent round, will form the superficies of a cone, whose base is the circle.

Next divide the arch EG into the same number of equal parts as the craticular prototype is divided into; and draw radii from all the points of division. Produce GF to I, so that FI be equal to FG; and from the centre I, with the radius IF, describe the quadrant FKH; and draw the right line IE. Then divide the arch KH into the same number of equal parts as the radius of the craticular prototype is divided into; and from the centre I draw radii through all the points of division, meeting EF in 1, 2, 3, &c. Lastly, from the centre F, with the radii F1, F2, F3, &c, describe concentric circles. So will the craticular ectype be formed, whose areolas will appear equal to each other.

Hence, what is delineated in every arcola of the craticular prototype, being transferred into the areolas of the craticular ectype, the images will be distorted or deformed; and yet they will appear in just proportion

to an eye elevated above the vertex at a height equal to the height of the cone itself.

If the chords of the quadrants be drawn in the craticular prototype, and chords of each of the 4th parts in the craticular ectype, every thing else remaining the same, there will be obtained the craticular ectype in a quadrangular pyramid.

And hence it will be easy to deform an image, in any other pyramid, whose base is any regular polygon.

Because the illusion is more perfect when the eye, by the contiguous objects, cannot estimate the distance of the parts of the deformed image, it is therefore proper to view it through a small hole.

Anamorphoses, or monstrous images, may also be made to appear in their natural shape and just proportions, by means of mirrors of certain shapes, from which those images are reflected again; and then they are said to be reformed.

For farther particulars, see Wolfius's *Catoptrics and Dioptrics*, and some other optical authors.

ANAPHORA, in *Astrology*, the second house, or that part of the heavens which is 30 degrees from the horoscope.

The term *anaphora* is also sometimes applied promiscuously to some of the succeeding houses, as the 5th, the 8th, and the 11th. In this sense *anaphora* is the same as *epanaphora*, and stands opposed to *cataphora*.

ANASTROUS signs, in *Astronomy*, a name given to the *duodecatemoria*, or the 12 portions of the ecliptic, which the signs possessed anciently, but have since deserted by the precession of the equinox.

ANAXAGORAS, one of the most celebrated philosophers among the ancients. He was born at Clazomene in Ionia, about the 70th Olympiad. He was a disciple of Anaximenes; and he gave up his patrimony, to be more at leisure for the study of philosophy, giving lectures in that science at Athens. Being persecuted in this place, and at last banished from it, he opened a school at Lampacum, where he was greatly honoured during his life, and still more after his death, statues having been erected to his memory. It is said he made some predictions relative to the phenomena of nature, as earthquakes &c., upon which he wrote some treatises. His principal tenets may be reduced to the following:—All things were in the beginning confusedly mixed together, without order and without motion. The principle of things is at the same time one and multiplex, which had the name of *homæmeries*, or similar particles, deprived of life. But there is beside this, from all eternity, another principle, an infinite and incorporeal spirit, who gave motion to these particles; in virtue of which, such as are homogeneous united, and such as were heterogeneous separated according to their different kinds. All things being thus put into motion by the spirit, and every thing being united to such as are similar, those that had a circular motion produced heavenly bodies, the lighter particles ascending, while those that were heavier descended. The rocks of the earth, being drawn up by the whirling force of the air, took fire, and became stars, beneath which the sun and moon took their stations. —It was said he also wrote upon the *Quadrature of the Circle*; the treatise upon which, Plutarch says, he composed during his imprisonment at Athens.

ANAX.



ANAXIMANDER, a very celebrated Greek philosopher, was born at Miletus in the 42d olympiad; for, according to Apollodorus, he was 64 years of age in the 2d year of the 58th olympiad. He was one of the first who publicly taught philosophy, and wrote upon philosophical subjects. He was the kinsman, companion, and disciple of Thales. He wrote also upon the sphere and geometry, &c. And he carried his researches into nature very far, for the time in which he lived. It is said that he discovered the obliquity of the zodiac; that he first published a geographical table; that he invented the gnomon, and set up the first sun-dial in an open place at Lacedæmon. He taught, that infinity of things was the principal and universal element; that this infinite always preserved its unity, but that its parts underwent changes; that all things came from it; and that all were about to return to it. By this obscure and indeterminate principle he probably meant the chaos of other philosophers. He held that the worlds are infinite; that the stars are composed of air and fire, which are carried about in their spheres, and that these spheres are gods; and that the earth is placed in the midst of the universe, as in a common centre. Farther, that infinite worlds were the produce of infinity; and that corruption proceeded from separation.

ANAXIMENES, an eminent Greek philosopher, born at Miletus, the friend, scholar, and successor of Anaximander. He diffused some degree of light upon the obscurity of his master's system. He made the first principle of things to consist in the air, which he considered as infinite or immense, and to which he ascribed a perpetual motion; that this air was the same as spirit or God, since the divine power resided in it, and agitated it. The stars were as fiery nails in the heavens; the sun a flat plate of fire; the earth an extended flat surface, &c.

ANDERSON (ALEXANDER), one of the brightest ornaments of the mathematical world, who flourished about 200 years ago. He was born at Aberdeen in Scotland, it would seem towards the latter part of the 16th century, as he was professor of mathematics at Paris in the early part of the 17th, where he published several ingenious works in geometry and algebra, both of his own, and of his friend Vieta's. Thus he published his "Supplementum Apollonii Redivivi; (of Ghetaldus) five analysis problematis hæcenus desiderati ad Apollonii Pergæi doctrinam περι νεύσεως, a Marino Ghetaldo Patritio Ragusino hujusque, non ita pridem restitutam. In qua exhibetur mechanice æqualitatum tertii gradus five solidarum, in quibus magnitudo omnino data, æquatur homogeneæ sub altero tantum coefficiente ignoto. Huic subnexa est variorum problematum practice." Paris, 1612, in 4to.

"Αιτιολογία: Pro Zetetico Apolloniani problematis a se jam pridem edito in supplemento Apollonii Redivivi. Ad clarissimum & ornatissimum virum Marinum Ghetaldum Patritium Ragusinum. In qua ad ea quae obiter mihi perstrinxit Ghetaldus respondetur, & analytices clarius detegitur." Paris, 1615, in 4to.

He published also,

"Francisci Vietæ Fontenacensis de Aequationum Recognitione & Emendatione Tractatus duo." Paris, 1615, in 4to; with a Dedication, Preface, and an Appendix, by Anderson.

And Vieta's Angulares Sectiones, with the Demonstrations by Anderson.

Alexander was cousin german to a Mr. David Anderson, of Finshaugh, a gentleman who also possessed a singular turn for mathematical and mechanical knowledge. This mathematical genius was hereditary in the family of the Andersons, and from them it seems to have been transmitted to their descendants of the name of Gregory in the same country: the daughter of the said David Anderson having been the mother of the celebrated mathematician James Gregory, and who herself first instructed her son James in the elements of the Mathematics, upon her observing in him, while yet a child, a strong propensity to those sciences.

The time either of the birth or death of our author Alexander, has not come to my knowledge.

ANDROGYNOUS, an appellation given, by astrologers, to such of the planets as are sometimes hot, and sometimes cold; as mercury, which is accounted hot and dry when near the sun, and cold and moist when near the moon.

ANDROMEDA, in *Astronomy*, a constellation of the northern hemisphere, representing the figure of a woman almost naked, her feet at a distance from each other, and her arms extended and chained; being one of the original 48 asterisms, or figures under which the ancients comprehended the stars, as derived to us from the Greeks, who probably had them from the Egyptians or Indians, and who, it is suspected, altered their names, and accompanied them with fabulous stories of their own. According to them, Cepheus, the father of Andromeda, was obliged to give her up to be devoured by a monster, to preserve his kingdom from the plague; but that she was delivered by Perseus, who slew the monster, and espoused her. And the family were all translated by Minerva to heaven, the mother being the constellation Cassiopeia.

She is sometimes called, in Latin, *Persea, Mulier catenata, Virgo devota*, &c. The Arabians, whose religion did not permit them to draw the figure of the human body on any occasion whatever, have changed this constellation into the figure of a sea-calf. Schickard has changed the name for that of the scripture name *Abigail*. And Schiller has also changed the figure of the constellation, for that of a sepulchre, and calls it the *Holy Sepulchre*.

This constellation contains about 27 stars that are visible to the naked eye; of which the principal are,  $\alpha$  *Andromeda's head*;  $\beta$  in the girdle, and called *mirach* or *mizar*;  $\gamma$  on the south foot, and named *alamak*, and sometimes *alhames*.

The number of stars placed in this constellation by the catalogue of Ptolemy is 23, by that of Tycho Brahe also 23, by that of Hevelius 47, and by that of Flamsteed 66.

ANEMOMETER, an instrument for measuring the force of the wind.

An instrument of this sort, it seems, was first invented by Wolfius in the year 1708, and first published in his *Arcometry* in 1709, also in the *Acta Eruditorum* of the same year; afterwards in his *Mathematical Dictionary*, and in his *Elem. Matheseos*. He says he tried the goodness of it, and observes that the internal struc-



ture may be preserved, so as to measure the force of running water, or that of men or horses when they draw or pull. The machine consists of sails, A, B, C, like those of a wind-mill, against which the wind blows, and by turning them about, raises an arm K with a weight L upon it, to different angles of elevation, shewn by the index M, according to the force of the wind. (Plate III. fig. 3)

In the *Philos. Transf.* another anemometer is described, in which the wind being supposed to blow directly against a flat side, or board, which moves along the graduated arch of a quadrant, the number of degrees it advances shews the comparative force of the wind.

In the same *Transactions*, for the year 1766, Mr. Alex. Brice describes a method, successfully practised by him, of measuring the velocity of the wind, by means of that of the shadow of clouds passing over a plane upon the earth.

Also in the same *Transactions*, for the year 1775, Dr. Lind gives a description of a very ingenious portable *Wind-Gauge*, by which the force of the wind is easily measured; a brief description of the principal parts of which here follows. This simple instrument consists of two glass tubes, AB, CD, (Plate III. fig. 4.) which should not be less than 8 or 9 inches long, the bore of each being about  $\frac{1}{8}$  of an inch diameter, and connected together by a small bent glass tube *ab*, only of about  $\frac{1}{8}$  of an inch diameter, to check the undulations of the water caused by a sudden gust of wind. On the upper end of the leg AB is fitted a thin metal tube, which is bent perpendicularly outwards, and having its mouth open to receive the wind blowing horizontally into it. The two tubes, or rather the two branches of the tube, are connected to a steel spindle KL by slips of brass near the top and bottom, by the sockets of which at *e* and *f* the whole instrument turns easily about the spindle, which is fixed into a block by a screw in its bottom, by the wind blowing in at the orifice at F. When the instrument is used, a quantity of water is poured in, till the tubes are about half full; then exposing the instrument to the wind, by blowing in at the orifice F, it forces the water down lower in the tube AB, and raises it so much higher in the other tube; and the distance between the surfaces of the water in the two tubes, estimated by a scale of inches and parts HI, placed by the sides of the tubes, will be the height of a column of water whose weight is equal to the force or momentum of the wind blowing or striking against an equal base. And as a cubic foot of water weighs 1000 ounces, or  $62\frac{1}{2}$  pounds, the 12th part of which is  $5\frac{1}{4}$  or  $5\frac{1}{2}$  pounds nearly, therefore for every inch the surface of the water is raised, the force of the wind will be equal to so many times  $5\frac{1}{2}$  pounds on a square foot. Thus, suppose the water stand 3 inches higher in the one tube, than in the other; then 3 times  $5\frac{1}{2}$  or  $15\frac{3}{2}$  pounds is equal to the pressure or force of the wind on the surface of a foot square.

This instrument of Dr. Lind's, measures only the force or momentum of the wind, but not its velocity. However the velocity of the wind may be deduced from its force so obtained, by help of some experiments performed by me at the Royal Military Academy, in the years 1786, 1787, and 1788; from which experiments it appears that a plane surface of a square foot suffers a

resistance of 12 ounces from the wind, when blowing with a velocity of 20 feet per second; and that the force is nearly as the square of the velocity. Hence then, taking the force of  $15\frac{3}{2}$  pounds, above found for the force of the wind when it sustains 3 inches of water, and taking the square roots of the forces, it will be, as  $\sqrt{12} : \sqrt{15\frac{3}{2}} :: 20 : 22\frac{4}{5}$  the 4th proportional, that is a velocity of  $22\frac{4}{5}$  feet per second, or  $15\frac{1}{2}$  miles per hour, is the rate or velocity at which the wind blows, when it raises the water 3 inches higher in the one tube than the other. And farther, as the said height is as the force, and the force as the square of the velocity, we shall have the force and velocity, corresponding to several heights of the water in the one tube, above that in the other, as in the following table.

*Table of the corresponding height of water, force on a square foot, and velocity of wind.*

Height of water.	Force of wind.	Velocity of wind per hour.
Inches.	Pounds.	Miles.
$0\frac{1}{2}$		4.5
$0\frac{1}{2}$		6.4
1	5.2	9.0
2	10.4	12.7
3	15.6	15.5
4	20.8	18.0
5	26.0	20.1
6	31.25	22.0
7	36.5	23.8
8	41.7	25.4
9	46.9	27.0
10	52.1	28.4
11	57.3	29.8
12	62.5	31.0

In one instance Dr. Lind found that the force of the wind was such as to be equal  $34\frac{9}{10}$  pounds, on a square foot; and this by proportion, in the following table, will be found to answer to a velocity of  $23\frac{1}{4}$  miles per hour.

Mr. Leutmann improved upon Wolfius's anemometer, by placing the sails horizontal, instead of vertical, which are easier to move, and turn what way soever the wind blows.

Mr. Benjamin Martin also (Plate III. fig. 5) improved upon the same. He made the axis like the fusee of a watch, having a cord winding upon it, with two weights at the ends which make always a balance to the force of the wind on the sails. See his *Philos. Britan.*

And M. D'Ons-en-Bray invented a new anemometer, which of itself expresses on paper, not only the several winds that have blown during the space of 24 hours, and at what hour each began and ended, but also the different strength or velocity of each. See *Mem. Acad. Scienc. an. 1734.* See also the article *Wind-Gauge*.


ANEMOSCOPE, is sometimes used to denote a machine invented to foretell the changes of the wind, or weather; and sometimes for an instrument shewing by an index what the present direction of the wind is. Of this latter sort, it seems, was that used by the ancients, and described by Vitruvius; and we have many of them at present in large or public buildings, where an index within a room or hall, points to the name of the quarter from whence the wind blows without; which is simply effected by connecting an index to the lower end of the spindle of a weather-cock.



It has been observed that hygrosopes made of catgut, or such like, prove very good anemoscopes; seldom failing, by the turning of the index, to foretell the shifting of the wind. See accounts of two different anemoscopes; one by Mr. Pickering, vol. 43 *Philos. Transf.* the other by Mr. B. Martin, vol. 2 of his *Philos. Britan.*

Otto Gueric also gave the title anemoscope to a machine invented by him to foretell the change of the weather, as to rain and fair. It consisted of the small wooden figure of a man, which rose and fell in a glass tube, as the atmosphere was more or less heavy. Which was only an application of the common barometer, as shewn by M. Couriers in the *Acta Eruditorum* for 1684.

ANGLE, *Angulus*, in *Geometry*, the opening or mutual inclination of two lines, or two planes, or three planes, meeting in a point called the vertex or angular point. Such as the angle formed by, or between, the two lines AB and AC, at the vertex or angular point A.—Also the two lines AB and AC, are called the *legs* or the *sides* of the angle.



Angles are sometimes denoted, or named, by the single letter placed at the angular point, as the angle A; and sometimes by three letters, placing always that of the vertex in the middle. The former method is used when only one angle has the same vertex; and the latter method it is necessary to use when several angles have the same vertex, to distinguish them from one another.

The measure of an angle, by which its quantity or magnitude is expressed, is an arch, as DE described from the centre A, with any radius at pleasure, and contained between its legs AB and AC.—Hence angles are compared and distinguished by the ratio of the arcs which subtend them, to the whole circumference of the circle; or by the number of degrees contained in the arc DE by which they are measured, to 360, the number of degrees in the whole circumference of the circle. And thus an angle is said to be of so many degrees, viz, as are contained in the arc DE.

Hence it matters not, with what radius the arc is described, by which an angle is measured, when great or small, as AD, or Ad, or any other: for the arcs DE, de, being similar, have the same ratio as their respective radii or circumferences, and therefore they contain the same number of degrees.—Hence it follows, that the quantity or magnitude of the angle remains still the same, though the legs be ever so much increased or diminished.—And thus, in similar figures, the like or corresponding angles are equal.

The taking or measuring of angles, is an operation of great use and extent in surveying, navigation, geography, astronomy, &c. And the instruments chiefly used for this purpose, are quadrants, sextants, octants,

theodolites, circumferentors, &c. Mr. Hadley invented an excellent instrument for taking the larger sort of angles, where much accuracy is required, or where the motion of the object, or any circumstance causing an unsteadiness in the common instruments, renders the observations difficult, or uncertain. And Mr. Dollond contrived an instrument for measuring small angles. See *Hadley's Quadrant, Micrometer*, and the *Philos. Transf.* Numbers 420, 425, and vol. 48.

#### To measure the Quantity of an Angle.

1. *On paper.* Apply the centre of a protractor to the vertex A of the angle, so that the radius may coincide with one leg, as AB; then the degree on the arch that is cut by the other leg AC, will give the measure of the angle required.

Or thus, by a line of chords. Take off the chord of 60 with a pair of compasses; and with that radius, from the centre A, describe an arc as DE. Then take this arc DE between the compasses, and apply the extent to the scale of chords, which will give the degrees in the angle as before.

M. De Lagny gave, in several memoirs of the Royal Academy of Sciences, a new method of measuring angles, which he called *Goniometry*. The method consists in measuring, with a pair of compasses, the arc which subtends the proposed angle, not by applying its extent to a pre-constructed scale, like chords, but in the following manner: From the angular point as a centre, with a pretty large radius, describe a circle, producing one leg of the angle backwards to cut off a semicircle; then search out what part of the semicircle the arc is which measures the given angle, in this manner; viz, take the extent of this arc with a very fine pair of compasses, and apply it several times to the arc of the semicircle, to find how often it is contained, with a small part remaining over; in the same manner take the extent of this small part, and apply it to the first arc, to find how often it is contained in it; and what remains this 2d time, apply in like manner to the first remainder; then the 3d remainder apply to the 2d, and so on, always counting how often the last remainder is contained in the next foregoing, till nothing remain, or till the remainder is insensible, and too small to be measured: Then, beginning at the last, and returning backwards, make a series of fractions of which the numerators are always 1, and the denominators are the number of times each remainder is contained in its next remainder, with the fractional part more, as derived from the following remainder; then the last fraction, thus obtained, will shew what part the given angle is of 180° or the semicircle; and being turned into degrees &c, will be the measure of the angle, and nearer, it is asserted, than it can be obtained by any other means; whether it be measuring, or calculating by trigonometrical tables.—Thus, if it be required to measure the angle GFH: With a large radius describe the semicircle GHI, meeting the leg FG produced in I; then take the extent of the arc GH in the compasses, and applying it from G upon the semicircle, suppose it contains 4 times to the point 4, and the part 4 I over; take 4 I and apply it from H to 1, so that HG contains 4 I once, and 1 G over; also apply this remainder to the former 4 I, and it contains 5 times, from 4 to 5, and 5 I over; and



and lastly the remainder 5 I is just two times contained in the former remainder 1 G or 1 2, without any remainder. Here then, the series of quotients, or numbers of times contained, are 4, 1, 5, 2; therefore, beginning at the last, the first fraction is  $\frac{1}{2}$ , or the last remainder is half the preceding one; and the 2d fraction is  $\frac{1}{5\frac{1}{2}}$  or  $\frac{2}{11}$ ; the 3d is  $\frac{1}{1\frac{2}{5}}$  or  $\frac{5}{7}$ ; and the fourth is

$\frac{1}{4\frac{1}{5}}$  or  $\frac{5}{21}$ ; that is, the arc GH is  $\frac{1}{2}\frac{1}{3}$  of a semicircle,

or the angle GFH is  $\frac{1}{2}\frac{1}{3}$  of two right angles, or of  $180^\circ$ ,

which is equivalent to  $37\frac{1}{2}$  degrees, or  $37^\circ 8' 34''\frac{2}{7}$ .

2. *On the ground.* Place a surveying instrument with its centre over the angular point to be measured, turning the instrument about till o, the beginning of its arch, fall in the line or direction of one leg of the angle; then turn the index about to the direction of the other leg, and it will cut off from the arch the degrees answering to the given angle.

To plot or lay down any given angle, either on paper or on the ground, may be performed in the same manner; and the method is farther explained under the articles PLOTTING and PROTRACTING, and under the names of the several instruments.

To bisect a given angle, as suppose the angle LKM. From the centre K, with any radius, describe the arc LM; then with the centres L and M, describe two arcs intersecting in N; and draw the line KN, which will bisect the given angle LKM, dividing it into the two equal angles LKN, MKN.



To trisect an angle, see TRISECTION.

Pappus, in his Mathematical Collections, book 4, treats of angular sections, but particularly and more largely, of trisections. He also treats of any section in general, in the 36th and following propositions.

ANGLES are of various kinds and denominations. With regard to the form of their legs, they are divided into *rectilinear*, *curvilinear*, and *mixed*.

*Rectilinear*, or *right-lined* ANGLE, is that whose legs are both right lines; as the foregoing angle CAB.

*Curvilinear* ANGLE, is that whose legs are both of them curves.

*Mixed*, or *mixtilinear* ANGLE, is that of which one leg is a right line, and the other a curve.

With regard to their *magnitude*, angles are again divided into *right* and *oblique*, *acute* and *obtuse*.

*Right* ANGLE, is that which is formed by one line perpendicular to another; or that which is subtended by a quadrant of a circle. As the angle BAC.—Therefore the measure of a right angle is a quadrant of a circle, or  $90^\circ$ ; and consequently all right angles are equal to each other.

*Oblique* ANGLE, is a common name for any angle that is not a right one; and it is either *acute* or *obtuse*.

VOL. I.


*Acute* ANGLE, is that which is less than a right angle, or less than 90 degrees; as the angle BAD. And

*Obtuse* ANGLE, is greater than a right angle, or whose measure exceeds 90 degrees; as the angle BAE.

With regard to their situation in respect of each other, angles are distinguished into *contiguous*, *adjacent*, *vertical*, *opposite*, and *alternate*.


*Contiguous* ANGLES, are such as have the same vertex, and one leg common to both. As the angles BAD, CAD, which have AD common.

*Adjacent* ANGLES, are those of which a leg of the one produced forms a leg of the other: as the angles GFH and IFH, which have the legs IF and FG in a straight line.—Hence adjacent angles are supplements to each other, making together 180 degrees. And therefore if one of these be given, the other will be known by subtracting the given one from 180 degrees. Which property is useful in surveying, to find the quantity of an inaccessible angle; viz, measure its adjacent accessible one, and subtract this from 180 degrees.



*Vertical* or *opposite* ANGLES, are such as have their legs mutually continuations of each other; as the two angles *a* and *b*, or *c* and *d*.—The property of these is, that the vertical or opposite angles are always equal to each other, viz,  $\angle a = \angle b$ , and  $\angle c = \angle d$ . And hence the quantity of an inaccessible angle of a field, &c, may be found, by measuring its accessible opposite angle.

*Alternate* ANGLES, are those made on the opposite sides of a line cutting two parallel lines; so, the angles *e* and *f*, or *g* and *h*, are alternates. And these are always equal to each other; viz, the  $\angle e = \angle f$ , or  $\angle g = \angle h$ .



*External* ANGLES, are the angles of a figure made without it, by producing its sides outwards; as the angles *i*, *k*, *l*, *m*. All the external angles of any right-lined figure, taken together, are equal to 4 right angles; and the external angle of a triangle is equal to both the internal opposite ones taken together; also any external angle of a trapezium inscribed in a circle, is equal to the internal opposite angle.

*Internal* ANGLES, are the angles within any figure, made by the sides of it; as the angles *n*, *o*, *p*, *q*.—In any right-lined figure, an internal angle as *n*, and its adjacent external angle *k*, together make two right angles, or 180 degrees; and all the internal angles *n*, *o*, *p*, *q*, taken



taken together, make twice as many right angles, wanting 4 right angles; also any two opposite internal angles of a trapezium inscribed in a circle, taken together, make two right angles, or 180 degrees.


*Homologous, or like ANGLES*, are such angles in two figures, as retain the same order from the first, in both figures.

*ANGLE out of the centre*, as  $G$ , is one whose vertex is not in the centre of the circle.—And its measure is


half the sum  $\frac{a+b}{2}$  of the arcs intercepted by its legs

when it is within the circle, or half the difference  $\frac{a-b}{2}$

when it is without.



*ANGLE at the centre*, is an angle whose vertex is in the centre; as the angle  $AFC$ , formed by two radii  $AF$ ,  $FC$ , and measured by the arc  $ADC$ .—An angle at the centre, as  $AFC$ , is always double of the angle  $ABC$  at the circumference, standing upon the same arc  $ADC$ ; and all angles at the centre are equal that stand upon the same or equal arcs: also all angles at the centre, are proportional to the arcs they stand upon; and so also are all angles at the circumference.



*ANGLE at the circumference*, is an angle whose vertex is somewhere in the circumference of a circle; as the angle  $ABC$ .


*ANGLE in a segment*, is an angle whose legs meet the extremities of the base of the segment, and its vertex is anywhere in its arch; as the angle  $B$  is in the segment  $ABC$ , or standing upon the supplemental segment  $ADC$ ; and is comprehended between two chords  $AB$  and  $BC$ .—An angle at the circumference is measured by half the arc  $ADC$  upon which it stands; and all the angles  $ABC$ ,  $AEC$ , in the same segment, are equal to each other.

*ANGLE in a semicircle* is an angle at the circumference contained in a semicircle, or standing upon a semicircle, or on a diameter.—An angle in a semicircle, is always a right angle; in a greater segment, the angle is less, and in a less segment the angle is greater than a right angle.

*ANGLE of a segment*, is that made by a chord with a tangent, at the point of contact. So  $IHK$  is the

angle of the less segment  $IMH$ , and  $IHL$ , the angle of the greater segment  $INH$ .—And the measure of each of these angles, is half the alternate or supplemental segment, or equal to the angle in it; viz, the  $\angle IHK = \angle INH$ , and the  $\angle IHL = \angle IMH$ .

*ANGLE of a semicircle*, is the angle which the diameter of a circle makes with the circumference. And Euclid demonstrates that this is less than a right angle, but greater than any acute angle.



*ANGLE of contact*, is that made by a curve line and a tangent to it, at the point of contact; as the angle  $IHK$ . It is proved by Euclid, that the angle of contact between a right line and a circle, is less than any right-lined angle whatever; though it does not therefore follow that it is of no magnitude or quantity. This has been the subject of great disputes amongst geometers, in which Peletarius, Clavius, Taquet, Wallis, &c, bore a considerable share; Peletarius and Wallis contending that it is no angle at all, against Clavius, who rightly asserts that it is not absolutely nothing in itself, but only of no magnitude in comparison with a right-lined angle, being a quantity of a different kind or nature; like as a line in respect to a surface, or a surface in respect to a solid, &c. And since his time, it has been proved by Sir I. Newton, and others, that angles of contact can be compared to each other, though not to right-lined angles, and what are the proportions which they bear to each other. Thus, the circular angles of contact  $IHK$ ,  $IHL$ , are to each other in the reciprocal subduplicate ratio of the diameters  $HM$ ,  $HN$ . And hence the circular angle of contact may be divided, by describing intermediate circles, into any number of parts, and in any proportion. And if, instead of circles, the curves be parabolas, and the point of contact  $H$  the common vertex of their axes; the angles of contact would then be reciprocally in the subduplicate ratio of their parameters. But in such elliptical and hyperbolic angles of contact, these will be reciprocally in the subduplicate of the ratio compounded of the ratios of the parameters, and the transverse axes. Moreover, if  $TOQ$  be a common parabola, to the axis  $OP$ , and tangent  $VOW$ , and whose equation is  $1 \times x = y^2$ , or  $x = y^2$ , where  $x$  is the absciss  $OP$ , and  $y$  the ordinate  $PQ$ , the parameter being 1: and if  $OR$ ,  $OS$ , &c, be other parabolas to the same axis, tangent, and parameter, their ordinate  $y$  being  $PR$ , or  $PS$ , &c, and their equations  $x = y^3$ ,  $x = y^4$ ,  $x = y^5$ , &c: then the series of angles of contact will be in succession infinitely greater than each other, viz, the angle of contact  $WOQ$  infinitely greater than  $WOR$ , and this infinitely greater than  $WOS$ , and so on infinitely.

And farther, between the angles of contact of any two of this kind, may other angles of contact be found *ad infinitum*, which shall infinitely exceed each other, and



and yet the greatest of them be infinitely less than the smallest right-lined angle. So also  $x^2 = y^2$ ,  $x^3 = y^4$ ,  $x^4 = y^5$ , &c, denote a series of curves, of which every succeeding one makes an angle with its tangent, infinitely greater than the preceding one; and the least of these, viz, that whose equation is  $x^2 = y^3$ , or the semicubical parabola, is infinitely greater than any circular angle of contact.

ANGLES are again divided into *plane*, *spherical*, and *solid*.

*Plane ANGLES*, are all those above treated of; which are defined by the inclination of two lines in a plane, meeting in a point.

*Spherical ANGLE*, is an angle formed on the surface of a sphere by the intersection of two great circles; or, it is the inclination of the planes of the two great circles.

The measure of a spherical angle, is the arc of a great circle of the sphere, intercepted between the two planes which form the angle, and which cuts the said planes at right angles. For their properties, &c, see SPHERE; SPHERICAL, and SPHERICAL TRIGONOMETRY.

*Solid ANGLE*, is the mutual inclination of more than two planes, or plane angles, meeting in a point, and not contained in the same plane; like the angles or corners of solid bodies. For their measure, properties, &c, see SOLID Angle.

Angles of other less usual kinds and denominations, are also to be found in some books of Geometry. As,

*Horned ANGLE*, *angulus cornutus*, that which is made by a right line, whether a tangent or secant, with the circumference of a circle.

*Lunular ANGLE*, *angulus lunularis*, is that which is formed by the intersection of two curve lines, the one concave, and the other convex.

*Cissoïd ANGLE*, *angulus cissoïdes*, the inner angle made by two spherical convex lines intersecting each other.

*Sistroïd ANGLE*, *angulus sistroïdes*, is that which is in form of a sistrum.

*Pelecoïd ANGLE*, *angulus pelecoides*, is that in form of a hatchet.

ANGLE, in *Trigonometry*. See TRIANGLE, TRIGONOMETRY, SINE, TANGENT, &c.


ANGLE, in *Mechanics*.—*Angle of Direction*, is that which is comprehended between the lines of direction of two conspiring forces.

ANGLE of *Elevation*, is that which is comprehended between the line of direction and any plane upon which the projection is made, whether horizontal or oblique.

ANGLE of *Incidence*, is that made by the line of direction of an impinging body, at the point of impact. As the angle ABC.

ANGLE of *Reflection*, is that made by the line of direction of the reflected body, at the point of impact. As the angle DBE.

Instead of the angles of incidence and reflection being estimated from the plane on which the body impinges, sometimes the complements of these are understood, viz, as estimated from a perpendicular to the reflecting plane; as the two angles ABF and DBF.



ANGLE in *Optics*.—*Visual* or *Optic Angle*, is the angle included between the two rays drawn from the two extreme points of an object to the centre of the pupil of the eye: as the angle HGI. The apparent magnitude of objects is greater or less, according to the angle under which they appear.—Objects seen under the same or an equal angle, always appear equal.—The least *visible* angle, or least angle under which a body can be seen, according to Dr. Hook, is one minute; but Dr. Jurin shews, that at the time of his debate with Hevelius on this subject, the latter could probably discover a single star under so small an angle as 20''. But bodies are visible under smaller angles as they are more bright or luminous. Dr. Jurin states the grounds of this controversy, and discusses the question at large, in his Essay upon distinct and indistinct Vision, published in Smith's Optics, pa. 148, & seq.

ANGLE of the *interval*, of two places, is the angle subtended by two lines directed from the eye to those places.

ANGLE of *incidence*, or *reflection*, or *refraction*, &c. See the respective words INCIDENCE, REFLECTION, REFRACTION, &c.

ANGLE in *Astronomy*.—*Angle of Commutation*. See COMMUTATION.

ANGLE of *elongation*, or *Angle at the Earth*. See ELONGATION.

*Parallactic ANGLE*, or the *parallax*, is the angle made at the centre of a star, the sun, &c, by two lines drawn, the one to the centre of the earth, and the other to its surface. See PARALLACTIC, and PARALLAX.

ANGLE of the *position of the sun*, of the *sun's apparent semi-diameter*, &c. See the respective words.

ANGLE at the *sun*, is the angle under which the distance of a planet from the ecliptic, is seen from the sun.

ANGLE of the *East*. See NONAGESIMAL.

ANGLE of *obliquity*, of the ecliptic, or the angle of inclination of the axis of the earth, to the axis of the ecliptic, is now nearly 23° 28'. See OBLIQUITY, and ECLIPTIC.

ANGLE of *longitude*, is the angle which the circle of a star's longitude makes with the meridian, at the pole of the ecliptic.

ANGLE of *right ascension*, is the angle which the circle of a star's right ascension makes with the meridian at the pole of the equator.

ANGLE in *Navigation*. ANGLE of the *rhumb*, or *loxodromic angle*. See RHUMB and LOXODROMIC.


ANGLES, in *Fortification*, are understood of those formed by the several lines used in fortifying, or making a place defensible.

These are of two sorts; *real* and *imaginary*.—*Real angles* are those which actually exist and appear in the works. Such as the *flanked angle*, the *angle of the epaule*, *angle of the flank*, and the *re-entering angle of the counterscarp*. *Imaginary*, or *occult angles*, are those which are only subservient to the construction, and which exist no more after the fortification is drawn. Such as the *angle of the centre*, *angle of the polygon*, *flanking angle*, *salient angle of the counterscarp*, &c.

ANGLE of, or at, the *centre*, is the angle formed at the centre of the polygon, by two radii drawn from the centre to two adjacent angles, and subtended by a side of



of it, as the angle ACB. This is found by dividing 360 degrees by the number of sides in the regular polygon.



**ANGLE of the Polygon**, is the angle intercepted between two sides of the polygon; as DAB, or ABE. This is the supplement of the angle at the centre, and is therefore found by subtracting the angle C from 180 degrees.

**ANGLE of the Triangle**, is half the angle of the polygon; as CAB or CBA; and is therefore half the supplement of the angle C at the centre.

**ANGLE of the Bastion**, is the angle FAG made by the two faces of the bastion. And is otherwise called the flanked angle.

**Diminished ANGLE**, is the angle BAG made by the meeting of the exterior side of the polygon with the face AG of the bastion.

**ANGLE of the curtain, or of the flank**, is the angle GHI made between the curtain and the flank.

**ANGLE of the epaule, or shoulder**, is the angle AGH made by the flank and the face of the bastion.

**ANGLE of the tenaille, or exterior flanking angle**, is the angle AKB made by the two rasant lines of defence, or the faces of two bastions produced.

**ANGLE of the counterscarp**, is the angle made by the two sides of the counterscarp, meeting before the middle of the curtain.

**ANGLE, flanking inward**, is the angle made by the flanking line with the curtain.

**ANGLE forming the flank**, is that consisting of one flank and one demigorge.

**ANGLE forming the face**, is that composed of one flank and one face.

**ANGLE of the moat**, is that made before the curtain, where it is intersected.

**Re-entering, or re-entrant ANGLE**, is that whose vertex is turned inwards, towards the place; as H or I.

**Salient, or fortant ANGLE**, is that turned outwards, advancing its point towards the field; as A or G.

**Dead ANGLE**, is a re-entering angle, which is not flanked or defended.

**ANGLE of a wall, in Architecture**, is the point or corner where the two sides or faces of a wall meet.

**ANGLES, in Astrology**, denote certain houses of a figure, or scheme of the heavens. So the horoscope of the first house, is termed the *angle of the east*.


**ANGUINEAL Hyperbola**, a name given by Sir

I. Newton to four of his curves of the second order, viz, species 33, 34, 35, 36, expressed by the equation  $xy^2 + ey = -ax^3 + bx^2 + cx + d$ ; being hyperbolas of a serpentine figure. See CURVES.

**ANGULAR**, something relating to, or that hath angles.

At a distance, angular bodies appear round; the angles and small inequalities disappearing at a much less distance than the bulk of the body.

**ANGULAR Motion**, is the motion of a body which moves circularly about a point; or the variation in the angle described by a line, or radius, connecting a body with the centre about which it moves.—Thus, a pendulum has an angular motion about its centre of motion; and the planets have an angular motion about the sun.—Two moveable points M and O, of which the one describes the arc MN, and the other the arc OP, in the same time, have an equal, or the same angular motion, although the real motion of the point O be much greater than that of the point M, viz, as the arc OP is greater than the arc MN. The angular motions of revolving bodies, as of the planets about the sun, are reciprocally proportional to their periodic times. And they are also, as their real or absolute motions directly, and as their radii of motion inversely.



**ANGULAR motion**, is also a kind of compound motion composed of a circular and a rectilinear motion; like the wheel of a coach, or other vehicle.

**ANIMATED needle**, a needle touched with a magnet or load-stone.

**ANNUAL**, in *Astronomy*, something that returns every year, or which terminates with the year.

**ANNUAL motion of the earth**. See EARTH.

**ANNUAL argument of longitude**. See ARGUMENT.

**ANNUAL epacts**. See EPACT.

**ANNUAL equation of the mean motion of the sun and moon**, and of the moon's apogee and nodes. See EQUATION.

**ANNUITIES**, a term for any periodical income, arising from money lent, or from houses, lands, salaries, pensions, &c; payable from time to time; either annually, or at other intervals of time.

Annuities may be divided into such as are *certain*, and such as depend on some *contingency*, as the continuance of a life, &c.

Annuities are also divided into annuities in *possession*, and annuities in *reversion*; the former meaning such as have commenced; and the latter such as will not commence till some particular event has happened, or till some given period of time has elapsed.

Annuities may be farther considered as payable either *yearly*, or *half yearly*, or *quarterly*, &c.

The *present value* of an annuity, is that sum, which, being improved at interest, will be sufficient to pay the annuity.

The *present value* of an annuity *certain*, payable yearly, is calculated in the following manner.—Let the annuity be 1, and let  $r$  denote the amount of 1*l.* for a year, or 1*l.* increased by its interest for one year. Then, 1 being the present value of the sum  $r$ , and having to find the present value of the sum 1, it will be, by proportion thus,



thus,  $r : 1 :: 1 : \frac{1}{r}$  the present value of 1*l.* due a year hence. In like manner  $\frac{1}{r^2}$  will be the present value of 1*l.* due 2 years hence; for  $r : 1 :: \frac{1}{r} : \frac{1}{r^2}$ . In like manner  $\frac{1}{r^3}, \frac{1}{r^4}, \frac{1}{r^5}$ , &c, will be the present value of 1*l.* due at the end of 3, 4, 5, &c, years respectively; and in general,  $\frac{1}{r^n}$  will be the value of 1*l.* to be received after the expiration of  $n$  years. Consequently the sum of all these, or  $\frac{1}{r} + \frac{1}{r^2} + \frac{1}{r^3} + \frac{1}{r^4} + \dots$  continued to  $n$  terms, will be the present value of all the  $n$  years annuities. And the value of the perpetuity, is the sum of the series continued *ad infinitum*.

But this series, it is evident, is a geometrical progression, whose first term and common ratio are each  $\frac{1}{r}$ , and the number of its terms  $n$ ; and therefore the sum  $s$  of all the terms, or the present value of all the annual payments, will be  $s = \frac{1}{r-1} - \frac{1}{r-1} \times \frac{1}{r^n}$ .

When the annuity is a perpetuity, it is plain that the last term  $\frac{1}{r^n}$  vanishes, and therefore  $\frac{1}{r-1} \times \frac{1}{r^n}$  also vanishes; and consequently the expression becomes barely  $s = \frac{1}{r-1}$ ; that is, any annuity divided by its interest for one year, is the value of the perpetuity.

So, if the rate of interest be 5 per cent; then  $\frac{100}{5} = 20$  is the value of the perpetuity at 5 per cent. Also  $\frac{100}{4} = 25$  is the value of the perpetuity at 4 per cent. And  $\frac{100}{3} = 33\frac{1}{3}$  is the value of the perpetuity at 3 per cent. interest. And so on.

If the annuity is not to be entered on immediately, but after a certain number of years, as  $m$  years; then the present value of the reversion is equal to the difference between two present values, the one for the first term of  $m$  years, and the other for the end of the last term  $n$ : that is, equal to the difference between

$$\frac{1}{r-1} - \frac{1}{r-1} \times \frac{1}{r^m} \text{ and } \frac{1}{r-1} - \frac{1}{r-1} \times \frac{1}{r^n}, \text{ or } = \frac{1}{r-1} \times \frac{1}{r^m} - \frac{1}{r-1} \times \frac{1}{r^n}.$$

Annuities certain differ in value, as they are made payable yearly, half-yearly, or quarterly. And by proceeding as above, using the interest or amount of a half year, or a quarter, as those for the whole year were

used, the following set of theorems will arise; where  $r$  denotes, as before, the amount of 1*l.* and its interest for a year, and  $n$  the number of years, during which, any annuity is to be paid; also  $P$  denotes the perpetuity

$\frac{1}{r-1}$ ,  $Y$  denotes  $\frac{1}{r-1} - \frac{1}{r-1} \times \frac{1}{r^n}$  the value of the annuity supposed payable yearly,  $H$  the value of the same when it is payable half-yearly, and  $Q$  the value when payable quarterly; or universally,  $M$  the value when it is payable every  $m$  part of a year.

$$\text{THEOR. 1. } Y = P - P \times \left(\frac{1}{r}\right)^n$$

$$\text{THEOR. 2. } H = P - P \times \left(\frac{2}{r+1}\right)^{2n}$$

$$\text{THEOR. 3. } Q = P - P \times \left(\frac{4}{r+3}\right)^{4n}$$

$$\text{THEOR. 4. } M = P - P \times \left(\frac{m}{r+m-1}\right)^{mn}$$

#### Example 1:

Let the rate of interest be 4 per cent, and the term 5 years; and consequently  $r = 1.04$ ,  $n = 5$ ,  $P = 25$ ; also let  $m = 12$ , or the interest payable monthly in theorem 4: then the present value of such annuity of 1*l.* a year, for 5 years, according as it is supposed payable 1*l.* yearly, or  $\frac{1}{2}$ *l.* every half year, or  $\frac{1}{4}$ *l.* every quarter, or  $\frac{1}{12}$ *l.* every month or  $\frac{1}{12}$ th part of a year, will be as follows:

$$\begin{aligned} Y &= 25 - 25 \times .821928 = 4.4518 \\ H &= 25 - 25 \times .820348 = 4.4913 \\ Q &= 25 - 25 \times .819543 = 4.5114 \\ M &= 25 - 25 \times .818996 = 4.5251 \end{aligned}$$

EXAMPLE 2. Supposing the annuity to continue 25 years, the rate of interest and every thing else being as before; then the values of the annuities for 25 years will be

$$\begin{aligned} Y &= 25 - 25 \times .375118 = 15.6221 \\ H &= 25 - 25 \times .371527 = 15.7118 \\ Q &= 25 - 25 \times .369709 = 15.7573 \\ M &= 25 - 25 \times .368477 = 15.7881 \end{aligned}$$

EXAMPLE 3. And if the term be 50 years, the values will be

$$\begin{aligned} Y &= 25 - 25 \times .140713 = 21.4822 \\ H &= 25 - 25 \times .138032 = 21.5492 \\ Q &= 25 - 25 \times .136685 = 21.5829 \\ M &= 25 - 25 \times .135775 = 21.6056 \end{aligned}$$

EXAMPLE 4. Also if the term be 100 years, the values will be

$$\begin{aligned} Y &= 25 - 25 \times .019800 = 24.5050 \\ H &= 25 - 25 \times .019053 = 24.5237 \\ Q &= 25 - 25 \times .018683 = 24.5329 \\ M &= 25 - 25 \times .018435 = 24.5391 \end{aligned}$$

Hence the difference in the value by making periods of payments smaller, for any given term of years, is the more as the intervals are smaller, or the periods more frequent. The same difference is also variable, both as the rate of interest varies, and also as the whole term of years  $n$  varies; and, for any given rate of interest, it



is evident that the difference, for any periods  $m$  of payments, first increases from nothing as the term  $n$  increases, when  $n$  is 0, to some certain finite term or value of  $n$ , when the difference  $D$  is the greatest or a maximum; and that afterwards, as  $n$  increases more, that difference will continually decrease to nothing again, and vanish when  $n$  is infinite: also the term or value of  $n$ , for the maximum of the difference, will be different according to the periods of payment, or value of  $m$ . And the general value of  $n$ , when the difference is a maximum between the yearly payments and the payments of  $m$  times in a year, is expressed by this formula, viz,

$$n = \frac{l. \frac{m+r-1}{m}}{l. r - \frac{m+r-1}{m}}, \text{ where } l. \text{ denotes the log-arithm of the quantity following it. Hence, taking the different values of } m, \text{ viz, 2 for half years, 4 for quarters, 12 for monthly payments, \&c, and substituting in the general formula, the term or value of } n \text{ for each case, when the difference in the present worths of the annuities, will be as follows, reckoning interest at 4 per cent, viz,}$$

$$n = \frac{l. \frac{r+1}{2}}{l. r - \frac{r+1}{2}} = 25.3777 \text{ for half-yearly payments,}$$

$$n = \frac{l. \frac{r+3}{4}}{l. r - \frac{r+3}{4}} = 25.3200 \text{ for quarterly payments,}$$

$$n = \frac{l. \frac{r+11}{12}}{l. r - \frac{r+11}{12}} = 25.2643 \text{ for monthly payments.}$$

Annuities may also be considered as in arrears, or as forborn, for any number of years; in which case each payment is to be considered as a sum put out to interest for the remainder of the term after the time it becomes due. And as 1*l.* due at the end of 1 year, amounts to  $r$  at the end of another year, and to  $r^2$  at the end of the 3d year, and to  $r^3$  at the end of the 4th year, and so on; therefore by adding always the last year's annuity, or 1, to the amounts of all the former years, the sum of all the annuities and their interests, will be the sum of the following geometrical series,  $1 + r + r^2 + r^3 + r^4 + \dots$  to  $r^{n-1}$ , continued till the last term be  $r^{n-1}$ , or till the number of terms be  $n$ , the number of years the annuity is forborn.

But the sum of this geometrical progression is  $\frac{r^n - 1}{r - 1}$ ,

which therefore is the amount of 1*l.* annuity forborn for  $n$  years. And this quantity being multiplied by any other annuity  $a$ , instead of 1, will produce the amount for that other annuity.

But the amounts of annuities, or their present values, are easiest found by the two following tables of numbers for the annuity of 1*l.* ready computed from the foregoing principles.

TABLE I.  
The Amount of an Annuity of 1*l.* at Comp. Interest.

Yrs.	at 3 per cent.	3½ per cent.	4 per cent.	4½ per cent.	5 per cent.	6 per cent.
1	1.000000	1.000000	1.000000	1.000000	1.000000	1.000000
2	2.030000	2.035000	2.040000	2.045000	2.050000	2.060000
3	3.090900	3.106230	3.121600	3.137030	3.152500	3.183600
4	4.183630	4.214940	4.246460	4.278190	4.310130	4.374620
5	5.309140	5.362470	5.416320	5.470710	5.525630	5.637090
6	6.468410	6.550150	6.632980	6.716890	6.801910	6.975320
7	7.662460	7.779410	7.898290	8.019150	8.142010	8.393840
8	8.892340	9.051690	9.214230	9.380010	9.549110	9.897470
9	10.159110	10.368500	10.582800	10.802110	11.026560	11.491320
10	11.463880	11.731390	12.006110	12.288210	12.577890	13.180790
11	12.807800	13.141990	13.486350	13.841180	14.206790	14.971640
12	14.192030	14.601960	15.025810	15.464030	15.917130	16.869940
13	15.617790	16.113030	16.626840	17.159910	17.712980	18.882140
14	17.086320	17.676990	18.291910	18.932110	19.598630	21.015070
15	18.598910	19.295680	20.323590	20.784050	21.578560	23.275970
16	20.156880	20.971030	21.824530	22.719340	23.657490	25.672530
17	21.761590	22.705020	23.697510	24.741710	25.840370	28.212880
18	23.414440	24.499690	25.645410	26.855080	28.132380	30.905650
19	25.116870	26.357180	27.671230	29.063560	30.539000	33.759990
20	26.870370	28.279680	29.778080	31.371420	33.065950	36.785590
21	28.676490	30.269470	31.969200	33.783140	35.719250	39.992730
22	30.536780	32.328900	34.247970	36.303380	38.505210	43.392290
23	32.452880	34.460410	36.617890	38.937030	41.430480	46.995830
24	34.426470	36.666530	39.082600	41.689200	44.502000	50.815580
25	36.459260	38.949860	41.645910	44.565210	47.727100	54.864510
26	38.553040	41.313100	44.311740	47.570640	51.113450	59.156380
27	40.709630	43.759060	47.084210	50.711320	54.669130	63.705770
28	42.930920	46.290630	49.967580	53.993330	58.402580	68.528110
29	45.218850	48.910800	52.966290	57.423030	62.322710	73.639800
30	47.575420	51.622680	56.084940	61.007070	66.438850	79.058190
31	50.002680	54.429470	59.328340	64.752390	70.760790	84.801680
32	52.502760	57.334500	62.701470	68.666250	75.298830	90.889780
33	55.077840	60.341210	66.209530	72.756230	80.063770	97.343160
34	57.730180	63.453150	69.857910	77.030260	85.066960	104.183750
35	60.462080	66.674010	73.652220	81.496620	90.320310	111.434780
36	63.275940	70.007600	77.598310	86.163970	95.836320	119.120870
37	66.174220	73.457870	81.702250	91.041340	101.628140	127.268120
38	69.159450	77.028890	85.970340	96.138200	107.709550	135.904210
39	72.234230	80.724910	90.409150	101.464420	114.095020	145.058460
40	75.401260	84.550280	95.025520	107.030320	120.799770	154.761970
41	78.663300	88.509540	99.826540	112.846690	127.839760	165.047680
42	82.023200	92.607370	104.819600	118.924790	135.231750	175.950540
43	85.483890	96.848630	110.012380	125.276400	142.993340	187.507580
44	89.048410	101.238330	115.412880	131.913840	151.143010	199.758030
45	92.719860	105.781670	121.029390	138.849970	159.700160	212.743510
46	96.501460	110.484030	126.870570	146.098210	168.685160	226.508120
47	100.396500	115.350970	132.945390	153.672630	178.119420	241.098610
48	104.408400	120.388260	139.263210	161.587900	188.025390	256.564530
49	108.540650	125.601850	145.833730	169.859360	198.426660	272.958400
50	112.796870	130.997910	152.667080	178.503030	209.348000	290.335900
51	117.180770	136.582840	159.773770	187.535660	220.815400	308.756060
52	121.696200	142.363240	167.164720	196.974770	232.856170	328.281420
53	126.347080	148.345950	174.851310	206.838630	245.498970	348.978310
54	131.137500	154.538060	182.845360	217.146370	258.773920	370.917010



TABLE II.

*The present Value of an Annuity of 1l.*

Yrs.	at 3 per cent.	3½ per cent.	4 per cent.	4½ per cent.	5 per cent.	6 per cent.
1	0.97087	0.96618	0.96154	0.95694	0.95238	0.94340
2	1.91347	1.89966	1.88610	1.87267	1.85941	1.83339
3	2.82861	2.80164	2.77509	2.74896	2.72325	2.67301
4	3.71710	3.67308	3.62990	3.58753	3.54595	3.46511
5	4.57971	4.51505	4.45182	4.38998	4.32948	4.21236
6	5.41719	5.32855	5.24214	5.15787	5.07569	4.91732
7	6.23028	6.11454	6.00205	5.89270	5.78637	5.53238
8	7.01969	6.87396	6.73274	6.59589	6.46321	6.20979
9	7.78611	7.60769	7.43533	7.26879	7.10782	6.80169
10	8.53020	8.31661	8.11090	7.91272	7.72173	7.36009
11	9.25262	9.00155	8.76048	8.52892	8.30541	7.88687
12	9.95400	9.66333	9.38507	9.11858	8.86325	8.38384
13	10.63496	10.30274	9.98565	9.68285	9.39357	8.85268
14	11.29607	10.92052	10.56312	10.22283	9.89864	9.29498
15	11.93794	11.51741	11.11839	10.73955	10.37966	9.71225
16	12.56110	12.09412	11.65230	11.23402	10.83777	10.10590
17	13.16612	12.65132	12.16567	11.70719	11.27407	10.47726
18	13.75351	13.18968	12.65930	12.15999	11.68959	10.82760
19	14.32380	13.70984	13.13394	12.59329	12.08532	11.15812
20	14.87747	14.21240	13.59033	13.00794	12.46221	11.46992
21	15.41502	14.69797	14.02916	13.40472	12.82115	11.76408
22	15.93692	15.16712	14.45112	13.78442	13.16300	12.04158
23	16.44361	15.62041	14.85684	14.14777	13.48857	12.30338
24	16.93554	16.05837	15.24696	14.49548	13.79864	12.55036
25	17.41315	16.48151	15.62208	14.82821	14.09294	12.78336
26	17.87684	16.89035	15.98277	15.14661	14.37519	13.00317
27	18.32703	17.28536	16.32959	15.45130	14.64303	13.21053
28	18.76411	17.66702	16.66306	15.74287	14.89813	13.40616
29	19.18845	18.03577	16.98371	16.02189	15.14107	13.58072
30	19.60044	18.39205	17.29203	16.28889	15.37245	13.76483
31	20.00043	18.73628	17.58849	16.54439	15.59281	13.92909
32	20.38877	19.06887	17.87355	16.78839	15.80268	14.08404
33	20.76579	19.39021	18.14765	17.02286	16.00255	14.23023
34	21.13184	19.70068	18.41120	17.24676	16.19290	14.36814
35	21.48722	20.00066	18.66461	17.46101	16.37419	14.49825
36	21.83225	20.29049	18.90828	17.66604	16.54685	14.62099
37	22.16724	20.57053	19.14258	17.86224	16.71129	14.73678
38	22.49246	20.84109	19.36786	18.04999	16.86789	14.84602
39	22.80822	21.10250	19.58448	18.22966	17.01704	14.94907
40	23.11477	21.35507	19.79277	18.40158	17.15909	15.04630
41	23.41240	21.59910	19.99305	18.56611	17.29437	15.13802
42	23.70136	21.83488	20.18563	18.72355	17.42321	15.22454
43	23.98190	22.06269	20.37079	18.87421	17.54591	15.30617
44	24.25427	22.28279	20.54884	19.01838	17.66277	15.38318
45	24.51871	22.49545	20.72004	19.15635	17.77407	15.45583
46	24.77545	22.70092	20.88465	19.28837	17.88007	15.52437
47	25.02471	22.89944	21.04294	19.41471	17.98102	15.58903
48	25.26671	23.09124	21.19513	19.53561	18.07716	15.65003
49	25.50166	23.27656	21.34147	19.65130	18.16872	15.70757
50	25.72976	23.45562	21.48218	19.76201	18.25593	15.76186
51	25.95123	23.62862	21.61749	19.86795	18.33898	15.81308
52	26.16624	23.79576	21.74758	19.96933	18.41807	15.86139
53	26.37499	23.95726	21.87267	20.06634	18.49340	15.90697
54	26.57766	24.11330	21.99296	20.15918	18.56515	15.94998

## THE USE OF TABLE I.

To find the Amount of an annuity forborn any number of years. Take out the amount from the 1st table, for the proposed years and rate of interest; then multiply it by the annuity in question; and the product will be its amount for the same number of years, and rate of interest.

And the converse to find the rate or time;

*Exam. 1.* To find how much an annuity of 50l. will amount to in 20 years at  $3\frac{1}{2}$  per cent. compound interest.—On the line of 20 years, and in the column of  $3\frac{1}{2}$  per cent, stands 28.27968, which is the amount of an annuity of 1l. for the 20 years; and therefore 28.27968 multiplied by 50, gives 1413.984l. or 1413l. 19s. 8d. for the answer.

*Exam. 2.* In what time will an annuity of 20l. amount to 1000l. at 4 per cent. compound interest?—Here the amount of 1000l. divided by 20l. the annuity, gives 50, the amount of 1l. annuity for the same time and rate. Then, the nearest tabular number in the column of 4 per cent. is 49.96758, which standing on the line of 28, shews that 28 years is the answer.

*Exam. 3.* If it be required to find at what rate of interest an annuity of 20l. will amount to 1000l. forborn for 28 years.—Here 1000 divided by 20 gives 50 as before. Then looking along the line of 28 years, for the nearest to this number 50, I find 49.96758 in the column of 4 per cent. which is therefore the rate of interest required.

## THE USE OF TABLE II.

*Exam. 1.* To find the present value of an annuity of 50l. which is to continue 20 years, at  $3\frac{1}{2}$  per cent.—By the table, the present value of 1l. for the same rate and time, is 14.21240; therefore  $14.2124 \times 50 = 710.62$ l. or 710l. 12s. 4d. is the present value sought.

*Exam. 2.* To find the present value of an annuity of 20l. to commence 10 years hence, and then to continue for 40 years, or to terminate 50 years hence, at 4 per cent. interest.—In such cases as this, it is plain we have to find the difference between the present values of two equal annuities, for the two given times; which therefore will be effected by subtracting the tabular value of the one term from that of the other, and multiplying by the annuity. Thus,

tabular value for 50 years 21.48218

tabular value for 10 years 8.11090

the difference 13.37128

mult. by 20

gives 267.4256

or 267l. 8s. 6d. the answer.

The foregoing observations, rules, and tables, contain all that is important in the doctrine of *annuities certain*. And for farther information, reference may be had to arithmetical writings, particularly Malcolm's Arithmetic, page 595; Simpson's Algebra, sect. 16; Dodson's Mathematical Repository, page 298, &c; Jones's Synopsis, ch. 10; Philos. Transf. vol. lxvi, page 109.

For what relates to the doctrine of *annuities on lives*, see ASSURANCE, COMPLEMENT, EXPECTATION, LIFE ANNUITIES, REVERSIONS, &c.

ANNULETS, in *Architecture*, are small square members, in the Doric capital, placed under the quarter round.

*Annulet* is also used for a narrow flat moulding, common to other parts of a column, as well as to the capital; and so called, because it encompasses the column around.



around. In which sense annulet is frequently used for *baguette*, or little *astragal*.

ANNULUS, a species of *Voluta*. See also *RING*.

ANOMALISTICAL Year, in *Astronomy*, called also *periodical year*, is the space of time in which the earth, or a planet, passes through its orbit. The anomalistical, or common year, is somewhat longer than the tropical year; by reason of the precession of the equinox.

And the apses of all the planets have a like progressive motion; by which it happens that a longer time is necessary to arrive at the aphelion, which has advanced a little, than to arrive at the same fixed star. For example, the tropical revolution of the sun, with respect to the equinox, is  $365^{\text{da}} 5^{\text{h}} 48^{\text{m}} 45^{\text{s}}$ ; but the sidereal, or return to the same

star,  $365 6 9 11$ ; and the anomalistic revolution is  $365 6 15 20$ , because the sun's apogee advances each year  $65''\frac{1}{2}$  with respect to the equinoxes, and the sun cannot arrive at the apogee till he has passed over the  $65''\frac{1}{2}$  more than the revolution of the year answering to the equinoxes.

To find the anomalistic revolution, say, As the whole secular motion of a planet *minus* the motion of its aphe-  
lion, is to 100 years or 3155760000 seconds, so is  $360^\circ$ , to the duration of the anomalistic revolution.


ANOMALOUS, is something irregular, or that deviates from the ordinary rule and method of other things of the same kind.

ANOMALY, in *Astronomy*, is an irregularity in the motion of a planet, by which it deviates from the aphelion or apogee; or it is the angular distance of the planet from the aphelion or apogee; that is, the angle formed by the line of the apses, and another line drawn through the planet.

Kepler distinguishes three kinds of anomaly; *mean*, *eccentric*, and *true*.

*Mean* or *Simple* ANOMALY, in the ancient astronomy, is the distance of a planet's mean place from the apogee. Which Ptolemy calls the angle of the mean motion.

But in the modern astronomy, in which a planet P is considered as describing an ellipse APB about the sun S, placed in one focus, it is the time in which the planet moves from its aphelion A, to the mean place or point of its orbit P.



Hence, as the elliptic area ASP is proportional to the time in which the planet describes the arc AP, that area may represent the mean anomaly.—Or, if PD be drawn perpendicular to the transverse axis AB, and meet the circle in D described on the same axis; then the mean anomaly may also be represented by the cir-

cular trilineal ASD, which is always proportional to the elliptic one ASP, as is proved in my *Menfuration*, pr. 8, page 296, second edition.—Or, drawing SG perpendicular to the radius DC produced; then the mean anomaly is also proportional to SG + the circular arc AD, as is demonstrated by Keil in his *Lect. Astron.*—Hence, taking DH = SG, the arc AH, or angle ACH will be the mean anomaly in practice, as expressed in degrees of a circle, the number of those degrees being to  $360^\circ$ , as the elliptic trilineal area ASP, is to the whole area of the ellipse; the degrees of mean anomaly, being those in the arc AH, or angle ACH.

*Eccentric* ANOMALY, or of the centre, in the modern astronomy, is the arc AD of the circle ADB intercepted between the apsis A and the point D determined by the perpendicular DPE to the line of the apses, drawn through the place P of the planet. Or it is the angle ACD at the centre of the circle.—Hence the eccentric anomaly is to the mean anomaly, as AD to AD + SG, or as AD to AH, or as the angle ACD to the angle ACH.

*True* or *Equated* ANOMALY, is the angle ASP at the sun, which the planet's distance AP from the aphelion, appears under; or the angle formed by the radius vector or line SP drawn from the sun to the planet, with the line of the apses.

The true anomaly being given, it is easy from thence to find the mean anomaly. For the angle ASP, which is the true anomaly, being given, the point P in the ellipse is given, and thence the proportion of the area ASP to the whole ellipse, or of the mean anomaly to  $360^\circ$  degrees. And for this purpose, the following easy rules for practice are deduced from the properties of the ellipse, by M. de la Caille in his *Elements of Astronomy*, and M. de la Lande, art. 1240 &c of his astronomy: 1st, As the square root of SB the perihelion distance, is to the square root of SA the aphelion distance, so is the tangent of half the true anomaly ASP, to the tangent of half the eccentric anomaly ACD. 2nd, The difference DH or SG between the eccentric and mean anomaly, is equal to the product of the eccentricity CS, by the sine of SCG the eccentric anomaly just found. And in this case, it is proper to express the eccentricity in seconds of a degree, which will be found by this proportion, as the mean distance 1: the eccentricity :: 206264.8 seconds, or  $57^\circ 17' 44''\frac{1}{2}$ , in the arch whose length is equal to the radius, to the seconds in the arc which is equal to the eccentricity CS; which being multiplied by the sine of the eccentric anomaly, to radius 1, as above, gives the seconds in SG, or in the arc DH, being the difference between the mean and eccentric anomalies. 3d, To find the radius vector SP, or distance of the planet from the sun, say either, as the sine of the true anomaly is to the sine of the eccentric anomaly, so is half the less axis of the orbit, to the radius vector SP; or as the sine of half the true anomaly is to the sine of half the eccentric anomaly, so is the square root of the perihelion distance SB, to the square root of the radius vector or planet's distance SP.

But the mean anomaly being given, it is not so easy to find the true anomaly, at least by a direct process: Kepler, who first proposed this problem, could not find a direct



a direct way of resolving it, and therefore made use of an indirect one, by the rule of false position, as may be seen page 695 of Kepler's *Epitom. Astron. Copernic.* See also § 628 Wolfius *Elem. Astron.* Now the easiest method of performing this operation, would be to work first for the eccentric anomaly, viz, assume it nearly, and from it so assumed compute what would be its mean anomaly by the rule above given, and find the difference between this result and the mean anomaly given; then assume another eccentric anomaly, and proceed in the same way with it, finding another computed mean anomaly, and its difference from the given one; and treating these differences as in the rule of position for a nearer value of the eccentric anomaly: repeating the operation till the result comes out exact. Then, from the eccentric anomaly, thus found, compute the true anomaly by the 1st rule above laid down.

Of this problem, Dr. Wallis first gave the geometrical solution by means of the protracted cycloid; and Sir Isaac Newton did the same at prop. 31 lib. 1 *Princip.* But these methods being unfit for the purpose of the practical astronomer, various series for approximation have been given, viz, several by Sir Isaac Newton in his *Fragmenta Epistolarum*, page 26, as also in the Schol. to the prop. above-mentioned, which is his best, being not only fit for the planets, but also for the comets, whose orbits are very eccentric. Dr. Gregory, in his *Astron.* lib. 3, has also given the solution by a series, as well as M. Reyneau, in his *Analyse Démontrée*, page 713, &c. And a better still for converging is given by Keil in his *Prælect. Astron.* page 375; he says, if the arc AH be the mean anomaly, calling its sine  $e$ , cosine  $f$ , the eccentricity  $g$ , also putting  $z = ge$ , and  $a = 1 + fg$ ; then the eccentric anomaly AD will be

$$= \frac{rz}{a} \times \left( 1 - \frac{z^2}{a^2} \&c \right), \text{ supposing } r = 57.29578 \text{ de-}$$

grees; of which the first term  $\frac{rz}{a}$  is sufficient for all the

planets, even for Mars itself, where the error will not exceed the 200th part of a degree; and in the orbit of the earth, the error is less than the 10000th part of a degree.

Dr. Seth Ward, in his *Astronomia Geometrica*, takes the angle AFP at the other focus, where the sun is not, for the mean anomaly, and thence gives an elegant solution. But this method is not sufficiently accurate when the orbit is very eccentric, as in that of the planet Mars, as is shewn by Bullialdus, in his defence of the *Philolaic. Astron.* against Dr. Ward. However, when Newton's correction is made, as in the Schol. above-mentioned, and the problem resolved according to Ward's hypothesis, Sir Isaac affirms that, even in the orbit of Mars, there will scarce ever be an error of more than one second.

ANSÆ, ANSES, in *Astronomy*, those seemingly prominent parts of the ring of the planet Saturn, discovered in its opening, and appearing like handles to the body of the planet; from which appearance the name *ansæ* is taken.

ANSER, in *Astronomy*, a small star, of the 5th or 6th magnitude, in the milky-way, between the eagle and swan, first brought into order by Hevelius.

VOL. I.

ANTARCTIC *pole*, denotes the southern pole, or southern end of the earth's axis.—The stars near the antarctic pole never appear above our horizon in these latitudes.

ANTARCTIC *circle*, is a small circle parallel to the equator, at the distance of  $23^{\circ} 28'$  from the antarctic or south pole.—At one time of the year the sun never rises above the horizon of any part within this circle; and at other times he never sets.

ANTARES, in *Astronomy*, the scorpion's heart; a fixed star of the first magnitude, in the constellation *Scorpio*.

ANTECANIS is used by some astronomers, to denote the constellation otherwise called *canis minor*, or the star *procyon*. It is so called, as preceding, or being the forerunner of the *canis major*, and rising a little before it.

ANTECEDENT, of a ratio, denotes the first of the two terms of the ratio, or that term which is compared with the other. Thus, if the ratio be 2 to 3, or  $a$  to  $b$ ; then 2 or  $a$  is the antecedent.

ANTECEDENTIAL *Method*, is a branch of general geometrical proportion, or universal comparison, and is derived from an examination of the Antecedents of ratios, having given consequents, and a given standard of comparison, in the various degrees of augmentation and diminution, which they undergo by composition and decomposition. This is a method invented by Mr. James Glenie, and published by him in 1793; a method which he says he always used instead of the fluxional and differential methods, and which is totally unconnected with the ideas of motion and time. See the author's treatise above-mentioned, and also his *Doctrine of Universal Comparison, or General Proportion*, 1789, upon which it is founded.

ANTECEDENTIA, a term used by astronomers when a planet &c moves westward, or contrary to the order of the signs aries, taurus, &c.—Like as when it moves eastward, or according to the order of the signs aries, taurus, &c, it is then said to move in *consequentia*.

ANTECIANS, or ANTOECI, in *Geography*, the inhabitants of the earth which occupy the same semicircle of the same meridian, but equally distant from the equator, the one north and the other south; as Peloponnesus and the Cape of Good Hope.

These have their noon, or midnight, or any other hour at the same time; but their seasons are contrary, being spring to the one, when it is autumn with the other; and summer with the one, when it is winter with the other; also the length of the day to the one, is equal to the length of night to the other.

ANTES, in *Architecture*, are small pilastres placed at the corners of buildings.

ANTICS, in *Architecture*, figures of men, beasts, &c, placed as ornaments to buildings.

ANTICUM, in *Architecture*, a porch before a door; also that part of a temple, which is called the outer temple, and lies between the body of the temple and the portico.

ANTILOGARITHM, the complement of the logarithm of a sine, tangent, secant, &c, to that of the radius. This is found by beginning at the left hand, subtracting each figure from 9, and the last figure from 10.

R

ANTINOUS,



ANTINOUS, in *Astronomy*, a part of the constellation *aquila*, or the eagle.

ANTIOCHIAN *Scē*, or *Academy*, a name given to the fifth academy or branch of academics. It took its name from being founded by Antiochus, a philosopher contemporary with Cicero; and it succeeded the Philonian academy. Though Antiochus was really a stoic, and only nominally an academic.

ANTIOCHIAN *epocha*, a method of computing time from the proclamation of liberty granted to the city of Antioch, about the time of the battle of Pharsalia.

ANTIPARALLELS, in *Geometry*, are those lines which make equal angles with two other lines, but contrary ways; that is, calling the former pair the 1st and 2d lines, and the latter pair the 3d and 4th lines, if the angle made by the 1st and 3d lines be equal to the angle made by the 2d and 4th, and contrariwise the angle made by the 1st and 4th equal to the angle made by the 2d and 3d; then each pair of lines are antiparallels with respect to each other, viz, the first and 2d, and the 3d and 4th. So, if AB and AC be any two lines, and FC and FE be two others, cutting them so,


that the angle B is equal to the angle E,

and the angle C is equal to the angle D;

then BC and DE are antiparallels with respect to AB and AC; also these latter are antiparallels with regard to the two former.—See also SUBCONTRARY.

It is a property of these lines, that each pair cuts the other into proportional segments, taking them alternately,

viz  $AB : AC :: AE : AD :: DB : EC$ ,  
and  $FE : FC :: FB : FD :: DE : BC$ .



ANTIPODES, in *Geography*, are the inhabitants of two places on the earth which lie diametrically opposite to each other, or that walk feet to feet; that is, if a line be continued down from our feet, quite through the centre of the earth, till it arrive at the surface on the other side, it will fall on the feet of our Antipodes, and *vice versa*.—Antipodes are 180 degrees distant from each other every way on the surface of the globe; they have equal latitudes, the one north and the other south, but they differ by 180 degrees of longitude: they have therefore the same climates or degrees of heat and cold, with the same seasons and length of days and nights; but all of these at contrary times, it being day to the one, when it is night to the other, summer to the one when it is winter to the other, &c: they have also the same horizon, the one being as far distant on the one side, as the other on the other side, and therefore when the sun, &c, rises to the one, it sets to the other. The Antipodes to London are a part a little south of New Zealand.

It has been said that Plato first started the notion of Antipodes, and gave them the name; which is likely enough, as he conceived that the earth was of a glo-

bular figure. But there have been many disputes upon this point, and the fathers of the church have greatly opposed it, especially Lactantius and Augustine, who laughed at it, and were greatly perplexed to think how men and trees should hang pendulous in the air with their feet uppermost, as he thought they must do, in the other hemisphere.

ANTISCIANS, or ANTISCII, in *Geography*, are people who dwell in the opposite hemispheres of the earth, as to north and south, and whose shadows at noon fall in contrary directions. This term is more general than *antæci*, with which it is often confounded. The *Antiscians* stand contradistinguished from *Periscians*.

ANTISCII is also used sometimes, among *Astrologers*, for two points of the heavens equally distant from the tropics. Thus the signs Leo and Taurus are accounted *antiscii* to each other.

ANTÆCI, see ANTECIANS.

APERTURE, in *Geometry*, is used for the space left between two lines which mutually incline towards each other, to form an angle.

APERTURE, in *Optics*, is the hole next the object-glass of a telescope or microscope, through which the light and the image of the object come into the tube, and are thence conveyed to the eye.

Aperture is also understood of that part of the object-glass itself which covers the former, and which is left pervious to the rays.

A great deal depends upon having a just aperture.—To find it experimentally: apply several circles of dark paper, of various sizes, upon the face of the glass, from the breadth of a straw, to such as leave only a small hole in the glass; and with each of these, separately, view some distinct objects, as the moon, stars, &c; then that aperture is to be chosen through which they appear the most distinctly.

Huygens first found the use of apertures to conduce much to the perfection of telescopes; and he found by experience (*Dioptr.* prop. 56.) that the best aperture for an object-glass, for example of 30 feet, is to be determined by this proportion, as 30 to 3, so is the square root of 30 times the distance of the focus of any lens, to its proper aperture: and that the focal distances of the eye-glasses are proportional to the apertures. And M. Auzout says he found, by experience, that the apertures of telescopes ought to be nearly in the subduplicate ratio of their lengths. It has also been found by experience that object-glasses will admit of greater apertures, if the tubes be blacked within side, and their passage furnished with wooden rings.

It is to be noted, that the greater or less aperture of an object-glass, does not increase or diminish the visible area of the object; all that is effected by this, is the admittance of more or fewer rays, and consequently the more or less bright the appearance of the object. But the largeness of the aperture or focal distance, causes the irregularity of its refractions. Hence, in viewing Venus through a telescope, a much less aperture is to be used than for the moon, or Jupiter, or Saturn, because her light is so bright and glaring. And this circumstance somewhat invalidates and disturbs Auzout's proportion, as is shewn by Dr. Hook, *Philos. Trans.* No. 4.

APHELION, or APHELIUM, in *Astronomy*, that point



point in the orbit of the earth, or a planet, in which it is at the greatest distance from the sun. Which is the point A (in the fig. to the art. ANOMALY) or extremity of the transverse axis, of the elliptic orbit, farthest from the focus S, where the sun is placed; and diametrically opposite to the perihelion B, or nearer extremity of the same axis. In the Ptolemaic system, or in the supposition that the sun moves about the earth, the aphelion becomes the *apogee*.

The times of the aphelia of the primary planets, may be known by their apparent diameter appearing the smallest, and also by their moving slowest in a given time. Calculations and methods of finding them have been given by many astronomers, as Ricciolli, *Almag. Nov. lib. 7, sect. 2 and 3*; Wolfius, *Elem. Astron. § 659*; Dr. Halley, *Philos. Transf. No. 128*; Sir I. Newton, *Princip. lib. 3, prop. 14*; Dr. Gregory, *Astron. lib. 3, prop. 14*; Keil, *Astron. Lect.*; De la Lande, *Memoires de l'Acad. 1755, 1757, 1766*, and in his *Astron. liv. 22*; also in the writings of MM. Euler, D'Alembert, Clairaut, &c, upon attraction.

The aphelia of the planets are not fixed; for their mutual actions upon one another keep those points of their orbits in a continual motion, which is greater or less in the different planets. This motion is made in consequentia, or according to the order of the signs; and Sir I. Newton shews that it is in the sesquiplicate ratio of the distance of the planet from the sun, that is, as the square root of the cube of the distance.

The quantities of this motion, as well as the place of the aphelion for a given time, are variously given by different authors. Kepler states them, for the year 1700 as in the following table.

Planets.	Aphelion.	Annual Motion.
Mercury	♊ 8° 25' 30"	1' 45"
Venus	♊ 3 24 27	1 18
Mars	♊ 0 51 29	1 7
Jupiter	♊ 8 10 40	0 47
Saturn	♊ 28 3 48	1 10
The Earth	♊ 8 25 30	

By De la Hire they are given as follows, for the same year 1700.

Planets.	Aphelion.	Annual Motion.
Mercury	♊ 13° 3' 40"	1' 39"
Venus	♊ 6 56 10	1 26
Mars	♊ 0 35 25	1 7
Jupiter	♊ 10 17 14	1 34
Saturn	♊ 29 14 41	1 22

And De la Lande states them as follows, for the year 1750.

Planets.	Aphelion.	Secular Motion.
Mercury	8 <sup>s</sup> 13° 33'	1° 57' 40"
Venus	10 8 13	4 10 0
Mars	5 1 28	1 51 40
Jupiter	6 10 22	1 43 20
Saturn	8 29 53	2 23 20
The Earth	9 8 38	1 49 10

Of the new planet, Herschel, or Georgium Sidus, the aphelion for 1790 was 11<sup>s</sup> 23° 29' 42", and its annual motion 50' 3". See *Connoissance des Temps*, 1786 and 1787.

APHRODISIUS, in Chronology, denotes the eleventh month in the Bythinian year, commencing on the 25th of July in ours.

APIAN or APPIAN (PETER), called in German *Bienerwitz*, a celebrated astronomer and mathematician, was born at Leisnig or Leipstick in Misnia, 1495, and made professor of mathematics at Ingolstadt, in 1524, where he died in the year 1552, at 57 years of age.

Apian wrote treatises upon many of the mathematical sciences, and greatly improved them; more especially astronomy and astrology, which in that age were much the same thing; also geometry, geography, arithmetic, &c. He particularly enriched astronomy with many instruments, and observations of eclipses, comets, &c. His principal work was the *Astronomicum Cæsareum*, published in folio at Ingolstadt in 1540, and which contains a number of interesting observations, with the descriptions and divisions of instruments. In this work he predicts eclipses, and constructs the figures of them in plano. In the 2d part of the work, or the *Meteoroscopium Planum*, he gives the description of the most accurate astronomical quadrant, and its uses. To it are added observations of five different comets, viz, in the years 1531, 1532, 1533, 1538, and 1539; where he first shews that the tails of comets are always projected in a direction from the sun.

Apian also wrote a treatise on *Cosmography*, or *Geographical Instruction*, with various mathematical instruments. This work Vossius says he published in 1524, and that Gemma Frisius republished it in 1540. But Weidler says he wrote it only in 1530, and that Gemma Frisius published it at Antwerp in 1550 and 1584, with observations of many eclipses. The truth may be, that perhaps all these editions were published.

In 1533 he made, at Norimberg, a curious instrument, which from its figure he called *Folium Populi*; which, by the sun's rays, shewed the hour in all parts of the earth, and even the unequal hours of the Jews.

In 1534 he published his *Inscriptiones Orbis*.

In 1540, his *Instrumentum Sinuum, sive Primi Mobilis*, with 100 problems.

Beside these, Apian was the author of many other works: among which may be mentioned the *Ephemerides* from the year 1534 to 1570: Books upon *Shadows: Arithmetical Centilogues*: Books upon *Arithmetic*, with the *Rule of Cos* (Algebra) demonstrated: Upon *Gauging: Almanacs*, with Astrological directions: A book upon *Conjunctions*: Ptolemy, with very correct figures, drawn in a quadrangular form: Ptolemy's works in Greek: Books of Eclipses: the works of Azoph, a very ancient astrologer: the works of Gebre: the Perspective of Vitello: of Critical Days, and of the Rainbow: a new Astronomical and Geometrical Radius, with various uses of Sines and Chords: Universal Astrolabe of Numbers: Maps of the World, and of particular countries: &c, &c.

Apian left a son, who many years afterwards taught mathematics at Ingolstadt, and at Tubinga. Tycho has preserved (*Progygn. p. 643*) his letter to the Landgrave of Hesse, in which he gives an opinion on the new star in Cassiopeia, of the year 1572.



One of the comets observed by Apian, viz, that of 1532, had its elements nearly the same as of one observed 128 $\frac{1}{2}$  years after, viz, in 1661, by Hevelius and other Astronomers; from hence Dr. Halley judged that they were the same comet, and that therefore it might be expected to appear again in the beginning of the year 1789. But it was not found that it returned at this period, although the astronomers then looked anxiously for it; and it is doubtful whether the disappointment might be owing to its passing unobserved, or to any errors in the observations of Apian, or to its period being disturbed and greatly altered by the actions of the superior planets, &c.

APIS, *musca*, the *Bee*, or *Fly*, in Astronomy, one of the southern constellations, containing 4 stars.

APOCATASTASIS, in Astronomy, is the period of a planet, or the time employed in returning to the same point of the zodiac from whence it set out.

APOGEE, *Apogæum*, in Astronomy, that point in the orbit of the sun, moon, &c, which is farthest distant from the earth. It is at the extremity of the line of the apsidæ; and the point opposite to it is called the *perigee*, where the distance from the earth is the least.

The ancient astronomers, considering the earth as the centre of the system, chiefly regarded the apogee and perigee: but the moderns, placing the sun in the centre, change these terms for the *aphelion* and *perihelion*.—The apogee of the sun, is the same thing as the aphelion of the earth; and the perigee of the sun is the same as the perihelion of the earth.

The manner of finding the apogee of the sun or moon, is shewn by Ricciolus, *Almag. Nov.* lib. 3, cap. 24; by Wolfius in *Elem. Astr.* § 618; by Cassini, *De la Hire*, and many others: see also *Memoires de l'Academie*, the *Philos. Transf.* vol. 5, 47, &c.

The quantity of motion in the apogee may be found by comparing two observations of it made at a great distance of time; converting the difference into minutes, and dividing them by the number of years elapsed between the two observations; the quotient gives the annual motion of the apogee. Thus, from an observation made by Hipparchus in the year before Christ 140, by which the sun's apogee was found 5° 30' of  $\Pi$ ; and another made by Ricciolus, in the year of Christ 1646, by which it was found 7° 26' of  $\odot$ ; the annual motion of the apogee is found to be 1' 2". And the annual motion of the moon's apogee is about 1° 10' 39' 52".

But the moon's apogee moves unequally. When she is in the syzygy with the sun, it moves forwards; but in the quadratures, backwards; and these progressions and regressions are not equable, but it goes forward slower when the moon is in the quadratures, or perhaps goes retrograde; and when the moon is in the syzygy, it goes forward the fastest of all.—See also Newton's *Theory of the Moon* for more upon this subject.

APOLLODORUS, a celebrated architect, under Trajan and Adrian, was born at Damascus, and flourished about the year of Christ 100. He had the direction of the stone bridge which Trajan ordered to be built over the Danube in the year 104, which was esteemed the most magnificent of all the works of that

emperor. Adrian, one day as Trajan was discoursing with this architect upon the buildings he had raised at Rome, would needs give his judgment, in which he shewed that he knew nothing of the matter. Apollodorus turned upon him bluntly, and said to him, Go paint Citruls, for you are very ignorant of the subject we are talking upon. Adrian at this time boasted of his painting Citruls well. This was the first step towards the ruin of Apollodorus; a slip which he was so far from attempting to retrieve, that he even added a new offence, and that too after Adrian was advanced to the empire, upon the following occasion: Adrian sent to him the plan of a temple of Venus; and though he asked his opinion, yet to shew that he had no need of him, and that he did not mean to be directed by it, the temple was already built. Apollodorus wrote his opinion very freely, and remarked such essential faults in it, as the emperor could neither deny nor remedy. He shewed that it was neither high nor large enough; that the statues in it were disproportioned to its bulk: for, said he, if the goddesses should have a mind to rise and go out, they could not do it. This put Adrian into a great passion, and prompted him to the destruction of Apollodorus. He banished him at first; then under the pretext of certain supposed crimes, of which he had him accused, he at last put him to death.

APOLLONIUS, of Perga, a city in Pamphilia, was a celebrated geometrician who flourished in the reign of Ptolemy Euergetes, about 240 years before Christ; being about 60 years after Euclid, and 30 years later than Archimedes. He studied a long time in Alexandria under the disciples of Euclid; and afterwards he composed several curious and ingenious geometrical works, of which only his books of Conic Sections are now extant, and even these not perfect. For it appears from the author's dedicatory epistle to Eudemus, a geometrician in Pergamus, that this work consisted of 8 books; only 7 of which however have come down to us.

From the Collections of Pappus, and the Commentaries of Eutocius, it appears that Apollonius was the author of various pieces in geometry, on account of which he acquired the title of the Great Geometrician. His *Conics* was the principal of them. Some have thought that Apollonius appropriated the writings and discoveries of Archimedes; Heraclius, who wrote the life of Archimedes, affirms it; though Eutocius endeavours to refute him. Although it should be allowed a groundless supposition, that Archimedes was the first who wrote upon Conics, notwithstanding his treatise on Conics was greatly esteemed; yet it is highly probable that Apollonius would avail himself of the writings of that author, as well as others who had gone before him; and, upon the whole, he is allowed the honour of explaining a difficult subject better than had been done before; having made several improvements both in Archimedes's problems, and in Euclid. His work upon Conics was doubtless the most perfect of the kind among the ancients, and in some respects among the moderns also. Before Apollonius, it had been customary, as we are informed by Eutocius, for the writers on Conics to require three different sorts of cones to cut the three different sections from, viz, the parabola from a right angled cone, the ellipse from an acute, and the



the hyperbola from an obtuse cone; because they always supposed the sections made by a plane cutting the cones to be perpendicular to the side of them: but Apollonius cut his sections all from any one cone, by only varying the inclination or position of the cutting plane; an improvement that has been followed by all other authors since his time. But that Archimedes was acquainted with the same manner of cutting any cone, is sufficiently proved, against Eutocius, Pappus, and others, by Guido Ubaldus, in the beginning of his Commentary on the 2d book of Archimedes's *Equiponderantes*, published at Pisa in 1588.

The first four books of Apollonius's Conics only have come down to us in their original Greek language; but the next three, the 5th, 6th, and 7th, in an Arabic version; and the 8th not at all. These have been commented upon, translated, and published by various authors. Pappus, in his *Mathematical Collections*, has left some account of his various works, with notes and lemmas upon them, and particularly on the Conics. And Eutocius wrote a regular elaborate commentary on the propositions of several of the books of the Conics.

The first four books were badly translated by Joan. Baptista Memmius. But a better translation of these in Latin was made by Commandine, and published at Bononia in 1566.—Vossius mentions an edition of the Conics in 1650; the 5th, 6th, and 7th books being recovered by Golius.—Claude Richard, Professor of mathematics in the imperial college of his order at Madrid, in the year 1632, explained, in his public lectures, the first four books of Apollonius, which were printed at Antwerp in 1655, in folio.—And the Grand Duke Ferdinand the 2d, and his brother Prince Leopold de Medicis, employed a professor of the Oriental languages at Rome to translate the 5th, 6th, and 7th books into Latin. These were published at Florence in 1661, by Borelli, with his own notes, who also maintains that these books are the genuine production of Apollonius, by many strong authorities, against Mydorgius and others, who suspected that these three books were not the real production of Apollonius.

As to the 8th book, some mention is made of it in a book of Golius's, where he had written that it had not been translated into Arabic; because it was wanting in the Greek copies, from whence the Arabians translated the others. But the learned Mersenne, in the preface to Apollonius's Conics, printed in his *Synopsis of the Mathematics*, quotes the Arabic philosopher Aben Nedin for a work of his about the year 400 of Mahomet, in which is part of that 8th book, and who asserts that all the books of Apollonius are extant in his language, and even more than are enumerated by Pappus; and Vossius says he has read the same; *De Scientiis Mathematicis*, pa. 55.—A neat edition of the first four books in Latin was published by Dr. Barrow, in 4to, at London in 1675.—A magnificent edition of all the 8 books, was published in folio, by Dr. Halley, at Oxford in 1710; together with the Lemmas of Pappus, and the Commentaries of Eutocius. The first four in Greek and Latin, but the latter four in Latin only, the 8th book being restored by himself.

The other writings of Apollonius, mentioned by Pappus, are,

1. The Section of a Ratio, or Proportional Sections, two books.
2. The Section of a Space, in two books.
3. Determinate Section, in two books.
4. The Tangencies, in two books.
5. The Inclinations, in two books.
6. The Plane Loci, in two books.

The contents of all these are mentioned by Pappus, and many lemmas are delivered relative to them; but none, or very little of these books themselves have descended down to the moderns. From the account however that has been given of their contents, many restorations have been made of these works, by the modern mathematicians, as follow: viz,

*Vieta*, Apollonius Gallus. The Tangencies. Paris, 1600, in 4to.

*Snellius*, Apollonius Batavus. Determinate Section. Lugd. 1601, 4to.

*Snellius*, Sectio Rationis & Spatii. 1607.

*Ghetaldus*, Apollonius Redivivus. The Inclinations. Venice, 1607, 4to.

*Ghetaldus*, Supplement to the Apollonius Redivivus: Tangencies. 1607.

*Ghetaldus*, Apollonius Redivivus, lib. 2. 1613.

*Alex. Anderson*, Supplem. Apol. Redivivi. Inclination. Paris, 1612, 4to.

*Alex. Anderson*, Pro Zetetico Apolloniani problematis a se jam pridem edito in Supplemento Apollonii Redivivi. Paris, 1615, 4to.

*Schooten*, Loca Plana restituta. Lug. Bat. 1656.

*Fermat*, Loca Plana, 2 lib. Tolos. 1679, folio.

*Halley*, Apol. de Sectione Rationis libri duo ex Arabico MS. Latine versi duo restituti. Oxon. 1706, 8vo.

*Simson*, Loca Plana, libri duo. Glasg. 1749, 4to.

*Simson*, Sectio Determinat. Glasg. 1776, 4to.

*Horsley*, Apol. Inclination. libri duo. Oxon. 1770, 4to.

*Lawson*, The Tangencies, in two books. Lond. 1771, 4to.

*Lawson*, Determinate Section, two books. Lond. 1772, 4to.

*Wales*, Determinate Section, two books. Lond. 1772, 4to.

*Burrow*, The Inclinations. Lond. 1779, 4to.

APONO (PETER de), a learned astronomer and philosopher, was born at Apono near Padua, about the year 1250. He described the *Astrolabium Planum*, by which were shewn the equations of the celestial houses for any hour and minute, and for any part of the world: it was published at Venice in 1502. He acquired the name of the *Conciliator*, on account of a book of his, in which he reconciles the writings of the ancient philosophers and physicians: the book was published at Venice in 1483. He resided at Padua, where, from his practising medicine, and his skill in astronomy, he fell under the suspicion of magic. He died in 1316, at 66 years of age.

APOPHYGE, in Architecture, is a concave part or ring of a column, lying above or below the flat member; and it owes its origin to the ring by which the ends of wooden columns were hooped, to prevent them from splitting.

APOTOME, the remainder or difference between two lines or quantities which are only commensurable in power. Such is the difference between 1 and  $\sqrt{2}$ .



or the difference between the side of the square and its diagonal.

The term is used by Euclid; and a pretty full explanation of such quantities is given in the tenth book of his Elements, where he distinguishes six kinds of apotomes, and shews how to find them all geometrically.

*APOTOME Prima*, is when the greater term is rational, and the difference of the squares of the two is a square number; as the difference  $3 - \sqrt{5}$ .

*APOTOME Secunda*, is when the less number is rational, and the square root of the difference of the squares of the two terms, has to the greater term, a ratio expressible in numbers; such is  $\sqrt{18} - 4$ , because the difference of the squares 18 and 16 is 2, and  $\sqrt{2}$  is to  $\sqrt{18}$  as  $\sqrt{1}$  to  $\sqrt{9}$  or as 1 to 3.

*APOTOME Tertia*, is when both the terms are irrational, and, as in the second, the square root of the difference of their squares, has to the greater term, a rational ratio: as  $\sqrt{24} - \sqrt{18}$ ; for the difference of their squares 24 and 18 is 6, and  $\sqrt{6}$  is to  $\sqrt{24}$  as  $\sqrt{1}$  to  $\sqrt{4}$  or as 1 to 2.

*APOTOME Quarta*, is when the greater term is a rational number, and the square root of the difference of the squares of the two terms, has not a rational ratio to it: as  $4 - \sqrt{3}$ , where the difference of the squares 16 and 3 is 13, and  $\sqrt{13}$  has not a ratio in numbers to 4.

*APOTOME Quinta*, is when the less term is a rational number, and the square root of the difference of the squares of the two, has not a rational ratio to the greater: as  $\sqrt{6} - 2$ , where the difference of the squares 6 and 4 is 2, and  $\sqrt{2}$  to  $\sqrt{6}$  or  $\sqrt{1}$  to  $\sqrt{3}$  or 1 to  $\sqrt{3}$  is not a rational ratio.

*APOTOME Sexta*, is where both terms are irrational, and the square root of the difference of their squares has not a rational ratio to the greater: as  $\sqrt{6} - \sqrt{2}$ ; where the difference of the squares 6 and 2 is 4, and  $\sqrt{4}$  to  $\sqrt{6}$  or 2 to  $\sqrt{6}$ , is not a rational ratio.

The doctrine of apotomes, in lines, as delivered by Euclid in the tenth book, is a very curious subject, and has always been much admired and cultivated by all mathematicians who have rightly understood this part of the elements; and therefore Peter Ramus has greatly exposed his judgment by censuring that book. And the first algebraical writers in Europe commonly employed a considerable portion of their works on an algebraical exposition of that book, which led them to the doctrine of surd quantities; as Lucas de Burgo, Cardan, Tartalea, Stifelius, Peletarius, &c. See also Pappus, lib. 4, prop. 3, and the introduction to lib. 7. And Dr. Wallis's Algebra, pa. 109.

*APOTOME*, in *Musical*, is the difference between a greater and less semitone, being expressed by the ratio of 128 to 125.


*APPARENT*, that which is visible, or evident to the eye, or the understanding.

*APPARENT conjunction* of the planets, is when a right line, supposed to be drawn through their centres, passes through the eye of the spectator, and not through the centre of the earth.—And, in general, the apparent conjunction of any objects, is when they appear or are placed in the same right line with the eye.


*APPARENT Altitude, Diameter, Distance, Horizon, Magnitude, Motion, Place, Time, &c.* See the respective substantives, for the quantity and measure of it.

The apparent state of things, is commonly very different from their real state, either as to distance, figure, magnitude, position, &c. &c. Thus,

*APPARENT Diameter, or Magnitude*, as for example of the heavenly bodies, is not the real length of the diameter, but the angle which they subtend at the eye, or under which they appear. And hence, the angle, or apparent extent, diminishing with the distance of the object, a very small object, as AB, may have the same apparent diameter as a very large one FG; and indeed the objects have all the same apparent diameter, that are contained in the same angle FEG. And if these are parallel, the real magnitudes are directly proportional to their distances.



But the apparent magnitude varies not only by the distance, but also by the position of it. So, if the object CD be changed from the direct position to the oblique one Cd, its apparent magnitude would then be only the angle CED, instead of the angle CED.



If the eye E be placed between two parallels AB, CD, these parallels will appear to converge or come nearer and nearer to each other the farther they are continued out, and at last they will appear to coincide in that point where the sight terminates, which will happen when the optic angle BED becomes equal to about one minute of a degree, the smallest angle under which an object is visible.—Also the apparent magnitudes of the same object FG or BD, seen at different distances, that is the angles FEG, BED, are in a less ratio than the reciprocal ratio of the distances, or the distance increases in a greater ratio than the angle or apparent magnitude diminishes. But when the object is very remote, or the optic angle is very small, as one degree or thereabouts, the angle then varies nearly as the distance reciprocally.

But although the optic angle be the usual or sensible measure of the apparent magnitude of an object, yet habit, and the frequent experience of looking at distant objects, by which we know that they are larger than they appear, has so far prevailed upon the imagination and judgment, as to cause this too to have some share in our estimation of apparent magnitudes; so that these will be judged to be more than in the ratio of the optic angles.

The apparent magnitude of the same object, at the same distance, is different to different persons, and different animals, and even to the same person, when viewed in different lights, all which may be occasioned by



by the different magnitudes of the eye, causing the optic angle to differ as that is greater or less: and since, in the same person, the more light there comes from an object, the less is the pupil of the eye, looking at that object; therefore the optic angle will also be less, and consequently the apparent magnitude of the object. Every one must have experienced the truth of this, by looking at another person in a room, and afterwards abroad in the sunshine, when he always appears smaller than in a room where the light is less. So also, objects up in the air, having more light coming from them than when they are upon the ground, or near it, may appear less in the former case than in the latter; like as the ball of the cross on the top of St. Paul's church, which is 6 feet in diameter, appears less than an object of the same diameter seen at the same distance below, near the ground. And this may be the chief reason why the sun and moon appear so much larger when seen in the horizon, where their beams are weak, than when they are raised higher, and their light is more bright and glaring.

Again, if the eye be placed in a rare medium, and view an object through a denser, as glass or water, having plane surfaces; the object will appear larger than it is: and contrariwise, smaller. And hence it is that fishes, and other objects, seen in the water, by an eye in the air, always appear larger than in the air.—In like manner, an object will appear larger when viewed through a globe of glass or water, or any convex spherical segments of these; and, on the contrary, it will appear smaller when viewed through a concave of glass or water.

*APPARENT Distance*, is that distance which we judge an object to be from us, when seen afar off. This is commonly very different from the true distance; because we are apt to think that all very remote objects, whose parts cannot well be distinguished, and which have no other visible objects near them, are at the same distance from us; though perhaps they may be thousands or millions of miles off; as in the case of the sun and moon. The apparent distances of objects are also greatly altered by the refraction of the medium through which they are seen.

*APPARENT Figure*, is the figure or shape which an object appears under when viewed at a distance; and is often very different from the true figure. For a straight line, viewed at a distance, may appear but as a point; a surface, as a line; and a solid, as a surface. Also these may appear of different magnitudes, and the surface and solid of different figures, according to their situation with respect to the eye: thus, the arch of a circle may appear a straight line; a square, a trapezium, or even a triangle; a circle, an ellipsis; angular magnitudes, round; and a sphere, a circle. Also all objects have a tendency to roundness and smoothness, or appear less angular, as their distance is greater: for, as the distance is increased, the smaller angles and asperities first disappear, by subtending a less angle than one minute; after these, the next larger disappear, for the same reason; and so on continually, as the distance is more and more increased; the object seeming still more and more round and smooth. So, a triangle, or square, at a great distance, shews only as a round speck; and the edge of the moon appears round to the eye,


notwithstanding the hills and valleys on her surface. And hence it is also, that near objects, as a range of lamps, or such like, seen at a great distance, appear to be contiguous, and to form one uniform continued magnitude, by the intervals between them disappearing, from the smallness of the angles subtended by them.

*APPARENT Motion*, is either that motion which we perceive in a distant body that moves, the eye at the same time being either in motion or at rest; or that motion which an object at rest seems to have, while the eye itself only is in motion.

The motions of bodies at a great distance, though really moving equally, or passing over equal spaces in equal times, may appear to be very unequal and irregular to the eye, which can only judge of them by the mutation of the angle at the eye. And motions, to be equally visible, or appear equal, must be directly proportional to the distances of the objects moving. Again, very swift motions, as those of the luminaries, may not appear to be any motions at all, but like that of the hour hand of a clock, on account of the great distance of the objects: and this will always happen, when the space actually passed over in one second of time, is less than about the 14000th part of its distance from the eye; for the hour hand of a clock, and the stars about the earth, move at the rate of 15 seconds of a degree in one second of time, which is only the 13751 part of the radius or distance from the eye. On the other hand, it is possible for the motion of a body to be so swift, as not to appear any motion at all; as when through the whole space it describes there constantly appears a continued surface or solid as it were generated by the motion of the object, like as when any thing is whirled very swiftly round, describing a ring, &c.


Also, the more oblique the eye is to the line which a distant body moves in, the more will the apparent motion differ from the true one. So, if a body revolve with an equable motion in the circumference of the circle ABCD &c, and the eye be at E in the plane of the circle; as the body moves from A to B and C, it seems to move slower and slower along the line ALK, till when the body arrives at C, it appears at rest at K; then while it really moves from C by D to F, it appears to move quicker and quicker from K by L to A, where its motion is quickest of all; after this it appears to move slower and slower from A to N while the body moves from F to H: there becoming stationary again, it appears to return from N to A in the straight line, while it really moves from H by I to A in the circle. And thus it appears to move in the line KN by a motion continually varying between the least, or nothing, at the extremes K and N, and the greatest of all at the middle point A. Or, if the motion be referred to the concave side of the circle, instead of the line KN, the appearances will be the same.

If an eye move directly forwards from E to O, &c; any remote object at rest at P, will appear to move the contrary





contrary way, or from P to Q, with the same velocity. But if the object P move the same way, and with the same velocity as the eye; it will seem to stand still. If the object have a less velocity than the eye, it will appear to move back towards Q with the difference of the velocities; and if it move faster than the eye, it will appear to move forwards from Q, with the same difference of the velocities. And so likewise when the object P moves contrary to the motion of the eye, it appears to move backwards with the sum of the motions of the two. And the truth of all this is experienced by persons sitting in a boat moving on a river, or in a wheel-carriage when running fast, and viewing houses or trees, &c, on the shore or side of the road, or other boats or wheel-carriages also in motion.



APPARENT PLACE of an object, in Optics, is that in which it appears, when seen in or through glass, water, or other refracting mediums; which is commonly different from the true place. So, if an object be seen in or through glass, or water, either plane or concave, it will appear nearer to the eye than its true place; but when seen through a convex glass, it appears more remote from the eye than the real place of it.

APPARENT PLACE of the Image of an object, in Catoptrics, is that where the image of an object made by the reflexion of a speculum appears to be; and the optical writers, from Euclid downwards, give it as a general rule that this is where the reflected rays meet the perpendicular to the speculum drawn from the object: so that if the speculum be a plane, the apparent place of the image will be at the same distance behind the speculum as the eye is before it; if convex, it will appear behind the glass nearer to the same; but if concave, it will appear before the speculum. And yet in some cases there are some exceptions to this rule, as is shewn by Kepler in his *Paralipomena in Vitellionem*, prop. 18. See also Wolfius *Catoptr.* § 51, 188, 233, 234.

APPARENT Place of a Planet, &c, in Astronomy, is that point in the surface of the sphere of the world, where the centre of the luminary appears from the surface of the earth.

APPARITION, in Astronomy, denotes a star's or other luminary's becoming visible, which before was hid. So, the heliacal rising, is rather an apparition than a proper rising.

Circle of perpetual APPARITION. See CIRCLE of perpetual apparition.

APPEARANCE, in Perspective, is the representation or projection of a figure, body, or the like object, on the perspective plane.—The appearance of an objective right line, is always a right line. See PERSPECTIVE.—Having given the appearance of an opaque body, and of a luminary, to find the appearance of the shadow; see SHADOW.

APPEARANCE of a star or planet. See APPARITION.

APPEARANCES, in Astronomy, &c, are more usually called *phænomena* and *phases*.—In Optics, the term *direct appearance* is used for the view or sight of any object by direct rays; without either refraction or reflexion.

APPIAN. See APIAN.

APPLICATE, *APPLICATA*, *Ordinate* *APPLICATE*, in Geometry, is a right line drawn to a curve, and bisected by its diameter. This is otherwise called an *ORDINATE*, which see.

APPLICATE Number. See CONCRETE.

APPLICATION, the act of applying one thing to another, by approaching or bringing them nearer together. So a longer space as measured by the continual application of a less, as a foot or yard by an inch, &c. And motion is determined by a successive application of any thing to different parts of space.

APPLICATION is sometimes used, both in Arithmetic and Geometry, for the rule or operation of division, or what is similar to it in geometry. Thus 20 applied to, or divided by 4, gives 5. And a rectangle

$ab$ , applied to a line  $c$ , gives the 4th proportional  $\frac{ab}{c}$ , or

another line which, with the given line  $c$ , will contain another rectangle which shall be equal to the given rectangle  $ab$ . And this is the sense in which Euclid uses the term, *lib. 6, pr. 28*.

APPLICATION, in Geometry, is also used for the act or supposition of putting or placing one figure upon another, to find whether they be equal or unequal; which seems to be the primary way in which the mind first acquires both the idea and proof of equality. And in this way Euclid, and other geometers, demonstrate some of the first or leading properties in geometry. Thus, if two triangles have two sides in the one triangle equal to two sides in the other, and also the angle included by the same sides equal to each other; then are the two triangles equal in all respects: for by conceiving the one triangle placed on the other, it is proved that they coincide or exactly agree in all their parts. And the same happens if, of two triangles, one side and the two adjacent angles of the one triangle, are equal, respectively, to one side and the two corresponding angles of the other. Thus also it may be proved that the diameter of a circle divides it into two equal parts, as also that the diagonal of a square or parallelogram bisects or divides it into two equal parts.

APPLICATION of one science to another, as of Algebra to Geometry, is said of the use made of the principles and properties of the one for augmenting and perfecting the other. Indeed all arts and sciences mutually receive aid from each other. But the application here meant, is of a more express and immediate nature; as will appear by what follows.

APPLICATION of Algebra or of Analysis to Geometry. The first and principal applications of algebra, were to arithmetical questions and computations, as being the first and most useful science in all the concerns of human life. Afterwards algebra was applied to geometry and all the other sciences in their turn. The application of algebra to geometry, is of two kinds; that which regards the plane or common geometry, and that which respects the higher geometry, or the nature of curve lines.

The first of these, or the application of algebra to common geometry, is concerned in the algebraical solution of geometrical problems, and finding out theorems in geometrical figures, by means of algebraical investigations or demonstrations. This kind of application



cation has been made from the time of the most early writers on algebra, as Diophantus, Lucas de Burgo, Cardan, Tartalea, &c, &c, down to the present times. Some of the best precepts and exercises of this kind of application, are to be met with in Newton's *Universal Arithmetic*, and in Thomas Simpson's *Algebra and Select Exercises*. Geometrical Problems are commonly resolved more directly and easily by algebra, than by the geometrical analysis, especially by young beginners; but then the synthesis, or construction and demonstration, is most elegant as deduced from the latter method. Now it commonly happens that the algebraical solution succeeds best in such problems as respect the sides and other lines in geometrical figures, and on the contrary, those problems in which angles are concerned, are best effected by the geometrical analysis. Newton gives these, among many other remarks on this branch. Having any problem proposed; compare together the quantities concerned in it; and, making no difference between the known and unknown quantities, consider how they depend upon, or are related to, one another; that we may perceive what quantities, if they are assumed, will, by proceeding synthetically, give the rest, and that in the simplest manner. And in this comparison, the geometrical figure is to be feigned and constructed at random, as if all the parts were actually known or given, and any other lines drawn that may appear to conduce to the easier and simpler solution of the problem. Having considered the method of computation, and drawn out the scheme, names are then to be given to the quantities entering into the computation, that is, to some few of them, both known and unknown, from which the rest may most naturally and simply be derived or expressed, by means of the geometrical properties of figures, till an equation be obtained, by which the value of the unknown quantity may be derived by the ordinary methods of reduction of equations, when only one unknown quantity is in the notation; or till as many equations are obtained as there are unknown letters in the notation.

For example, suppose it were required to inscribe a square in a given triangle. Let ABC be the given triangle; and feign DEFG to be the required square; also draw the perpendicular BP of the triangle, which will be given, as well as all the sides of it. Then, considering that the triangles BAC, BEF are similar, it will be proper to make the notation as follows, viz, making the base AC =  $b$ , the perpendicular BP =  $p$ , and the side of the square DE or EF =  $x$ . Hence then BQ = BP - ED =  $p - x$ ; consequently, by the proportionality of the parts of those two similar triangles, viz, BP : AC :: BQ : EF, it is  $p : b :: p - x : x$ ; then, multiply extremes and means &c, there arises  $px = bp - bx$ , or  $bx + px = bp$ , and  $x = \frac{bp}{b + p}$  the side of the square sought; that is, a fourth proportional to the base and perpendicular, and the sum of the two, taking this sum for the first term, or AC + BP : BP :: AC : EF.

The other branch of the application of algebra to geo-

metry, was introduced by Descartes, in his *Geometry*, which is the new or higher geometry, and respects the nature and properties of curve lines. In this branch, the nature of the curve is expressed or denoted by an algebraic equation, which is thus derived: A line is conceived to be drawn, as the diameter or some other principal line about the curve; and upon any indefinite points of this line other lines are erected perpendicularly, which are called ordinates, whilst the parts of the first line cut off by them, are called abscissés. Then, calling any absciss  $x$ , and its corresponding ordinate  $y$ , by means of the known nature, or relations of the other lines in the curve, an equation is derived, involving  $x$  and  $y$ , with other given quantities in it. Hence, as  $x$  and  $y$  are common to every point in the primary line, that equation, so derived, will belong to every position or value of the absciss and ordinate, and so is properly considered as expressing the nature of the curve in all points of it; and is commonly called the equation of the curve.

In this way it is found that any curve line has a peculiar form of equation belonging to it, and which is different from that of every other curve, either as to the number of the terms, the powers of the unknown letters  $x$  and  $y$ , or the signs or coefficients of the terms of the equation. Thus, if the curve line HK be a circle, of which HI is part of the diameter, and IK a perpendicular ordinate: then put HI =  $x$ , IK =  $y$ , and  $p$  = the diameter of the circle, the equation of the circle will be  $px - x^2 = y^2$ . But if HK be an ellipse, an hyperbola, or parabola, the equation of the curve will be different, and for all the four curves, will be respectively as follows, viz,

for the circle - -  $px - x^2 = y^2$ ,

for the ellipse -  $px - \frac{p}{t}x^2 = y^2$ ,

for the hyperbola -  $px + \frac{p}{t}x^2 = y^2$ ,

for the parabola -  $px - - = y^2$ ;



where  $t$  is the transverse axis, and  $p$  its parameter. And, in like manner for other curves.

This way of expressing the nature of curve lines, by algebraic equations, has given occasion to the greatest improvement and extension of the geometry of curve lines; for thus, all the properties of algebraic equations, and their roots, are transferred and added to the curve lines, whose abscissés and ordinates have similar properties. Indeed the benefit of this sort of application is mutual and reciprocal, the known properties of equations being transferred to the curves they represent; and, on the contrary, the known properties of curves transferred to their representative equations. See CURVES.

APPLICATION of Geometry to Algebra. Besides the use and application of the higher geometry, namely, of curve lines, to detecting the nature and roots of equations, and to the finding the values of those roots by the geometrical construction of curve lines, even common geometry may be made subservient to the purposes of algebra. Thus, to take a very plain and simple instance, if it were required to square the binomial  $a + b$ ;


S

by





by forming a square, as in the annexed figure, whose side is equal to  $a + b$ , or the two lines or parts added together denoted by the letters  $a$  and  $b$ ; and then drawing two lines parallel to the sides, from the points where the two parts join, it will be immediately evident that the whole square of the compound quantity  $a + b$ , is equal to the squares of both the parts, together with two rectangles under the two parts, or  $a^2$  and  $b^2$  and  $2ab$ , that is the square of  $a + b$  is equal to  $a^2 + b^2 + 2ab$ , as derived from a geometrical figure or construction. And in this very manner it was, that the Arabians, and the early European writers on algebra, derived and demonstrated the common rule for resolving compound quadratic equations. And thus also, in a similar way, it was, that Tartalea and Cardan derived and demonstrated all the rules for the resolution of cubic equations, using cubes and parallelopipedons instead of squares and rectangles. And many other instances might be given of the use and application of geometry in algebra.



*APPLICATION of Algebra and Geometry to Mechanics.* This is founded on the same principles as the application of algebra to geometry. It consists principally in representing by equations the curves described by bodies in motion, by determining the equation between the spaces which the bodies describe, when actuated by any forces, and the times employed in describing them, &c. A familiar instance also of the application of geometry to mechanics, may be seen at the article ACCELERATION, where the perpendiculars of triangles represent the times, the bases the velocities, and the areas the spaces described by bodies in motion; a method first given by Galileo. In short, as velocities, times, forces, spaces, &c, may be represented by lines and geometrical figures; and as these again may be treated algebraically; it is evident how the principles and properties, of both algebra and geometry, may be applied to mechanics, and indeed to all the other branches of mixed mathematics.

*APPLICATION of Mechanics to Geometry.* This consists chiefly in the use that is sometimes made of the centre of gravity of figures, for determining the contents of solids described by those figures.

*APPLICATION of Geometry and Astronomy to Geography.* This consists chiefly in three articles. 1st, In determining the figure of the globe we inhabit, by means of geometrical and astronomical operations. 2d, In determining the positions of places, by observations of latitudes and longitudes. 3d, In determining, by geometrical operations, the positions of such places as are not far distant from one another.

Geometry and Astronomy are also of great use in Navigation.

*APPLICATION of Geometry and Algebra to Physics or Natural Philosophy.* This application we owe to Newton, whose philosophy may therefore be called the geometrical or mathematical philosophy; and upon this application are founded all the physico-mathematical sciences. Here a single observation or experiment will often produce a whole science: so when we know, as we do by experience, that the rays of light, in reflect-

ing, make the angle of incidence equal to the angle of reflexion; we thence deduce the whole science of catoptrics: for that experiment once admitted, catoptrics become a science purely geometrical, since it is reduced to the comparison of angles and lines given in position. And the same in many other sciences.

*APPLICATION of one thing to another*, in general, is employed to denote the use that is made of the former, to understand or to perfect the latter. Thus, the application of the cycloid to pendulums, means the use made of the cycloidal curve for improving the doctrine and use of pendulums.

*APPLY.* This term is used two different ways, in geometry.

1st, It signifies to transfer or place a given line, either in a circle or some other figure, so that the extremities of the line shall be in the perimeter of the figure.

2d, It is also used to express division in geometry, or to find one dimension of a rectangle, when the area and the other dimension are given. As the area  $ab$  applied to the line  $c$ , is  $\frac{ab}{c}$ .

*APPROACH*, the *curve of equable approach*. It was first proposed by Leibnitz, namely, to find a curve, down which a body descending by the force of gravity, shall make equal approaches to the horizon in equal portions of time. It has been found by Bernoulli and others, that the curve is the second cubical parabola, placed with its vertex uppermost, and which the descending body must enter with a certain determinate velocity.—Varignon rendered the question general for any law of gravity, by which a body may approach towards a given point by equal spaces in equal times. And Maupertuis also resolved the problem in the case of a body descending in a medium which resists as the square of the velocity. See *Hist. de l'Acad. des Sciences* for 1699 and 1730.

*Method of APPROACHES*, a name given by Dr. Wallis, in his Algebra, to a method of resolving certain problems relating to square numbers, &c. This is done by first assigning certain limits to the quantities required, and then approaching nearer and nearer till a coincidence is obtained.—In this sense, the method of Trial-and-error, or double rule of False Position, may be considered as a method of approaches.

*APPROACHES*, in *Fortification*, the several works made by the besiegers, for advancing or getting nearer to a fortress or place besieged. Such as the trenches, mines, saps, lodgments, batteries, galleries, epaulments, &c.

*APPROACHES*, or *Lines of APPROACH*, are particularly used for trenches dug in the ground, and the earth thrown up on the side next the place besieged; under the defence or shelter of which, the besiegers may approach without loss, as near as possible to the place, to raise batteries and plant guns &c, to batter it.—The lines of approach are commonly carried on, in a zig-zag way, parallel to the opposite faces of the besieged work, or nearly so, that they may not be enfiladed by the guns from the enemy's works. And they are also connected by parallels or lines of communication.—The besieged commonly make counter-approaches, to interrupt and defeat the approaches of the besiegers.

The



The ancients made their approaches towards the place besieged, much after the same manner as the moderns. Folard shews, that they had their trenches, their parallels, saps, &c.; which, though usually thought of modern invention, it appears, have been practised long before, by the Greeks, Romans, Asiatics, &c.

APPROXIMATION, a continual approach, still nearer and nearer, to a root or any quantity sought. — Methods of continual approximation for the square roots and cube roots of numbers, have been employed by algebraists and arithmeticians, from Lucas de Burgo down to the present time. And the later writers have given various approximations, not only for the roots of higher powers, or all simple equations, but for the roots of all sorts of compound equations whatever: especially Newton, Wallis, Raphson, Halley, De Lagny, &c, &c; all of them forming a kind of infinite series, either expressed or understood, converging nearer and nearer to the quantity sought, according to the nature of the process.

It is evident that if a number proposed be not a true square, then no exact square root of it can be found, explicable by rational numbers, whether integers or fractions: therefore, in such cases, we must be content with approximations, or coming continually nearer and nearer to the truth. In like manner, for the cube and other roots, when the proposed quantities are not exact cubes, or other powers.

The most easy and general method of approximation, is perhaps by the rule of Double Position, or, what is sometimes called, the Method of Trial-and-error; which method see under its own name. And among all the methods for the roots of pure powers, of which there are many, I believe the best is that which was discovered by myself, and given in the first volume of my Mathematical Tracts, in point of ease, both of execution and for remembering it. The method is this: if  $N$  denote any number, out of which is to be extracted the root whose index is denoted by  $r$ , and if  $n$  be the nearest root first taken; then shall  $\frac{r + 1 \cdot N + r - 1 \cdot n^r}{r - 1 \cdot N + r + 1 \cdot n^r} \times n$  be the required root of  $N$  very nearly; or as  $r - 1$  times the given number added to  $r + 1$  times the nearest power, is to  $r + 1$  times the given number added to  $r - 1$  times the nearest power, so is the assumed root  $n$ , to the required root, very nearly. Then this last value of the root, so found, if one still nearer is wanted, is to be used for  $n$  in the same theorem, to repeat the operation with it. And so on, repeating the operation as often as necessary. Which theorem includes all the rational formulæ of Halley and De Lagny.

For example, suppose it were required to double the cube, or to find the cube root of the number 2. Here  $r = 3$ ; consequently  $r + 1 = 4$ , and  $r - 1 = 2$ ; and therefore the general theorem becomes

$$\frac{4N + 2n^3}{2N + 4n^3} \times n \text{ or } \frac{2N + n^3}{N + 2n^3} \times n \text{ for the cube root of } N;$$

or as  $N + 2n^3 : 2N + n^3 :: n : \text{the root sought nearly}$ . Now, in this case,  $N = 2$ , and therefore the nearest root  $n$  is 1, and its cube  $n^3 = 1$  also: hence  $N + 2n^3 = 2 + 2 = 4$ , and  $2N + n^3 = 4 + 1 = 5$ ; therefore, as  $4 : 5 :: 1 : \frac{5}{4}$  or  $1\frac{1}{4} = 1.25$  the first approximation.

Again, taking  $r = \frac{5}{4}$ , and consequently  $r^3 = \frac{125}{64}$ ;

$$\text{hence } N + 2n^3 = 2 + \frac{250}{64} = \frac{378}{64},$$

$$\text{and } 2N + n^3 = 4 + \frac{125}{64} = \frac{381}{64};$$

therefore as  $378 : 381$ , or as  $126 : 127 :: \frac{5}{4} : \frac{635}{504} = 1.259921$ , which is the cube root of 2, true in all the figures. And by taking  $\frac{635}{504}$  for a new value of  $n$ , and

repeating the process again, a great many more figures may be found.

*Of the Roots of Equations by APPROXIMATION.* — Stevinus and Vieta gave methods for finding values, always nearer and nearer, of the roots of equations. And Oughtred and others pursued and improved the same. These however were very tedious and imperfect, and required a different process for every degree of equations. But Newton introduced, not only general methods for expressing radical quantities by approximating infinite series, but also for the roots of all sorts of compound equations whatever, which are both easy and expeditious: which will be more particularly described under each respective word or article. His method for approximating of roots, is in substance this: First take a value of the root as near as may be, by trials, either greater or less; then assuming another letter to denote the unknown difference between this and the true value, substitute into the equation the sum or difference of the approximate root and this assumed letter, instead of the unknown letter or root of the equation, which will produce a new equation having only the assumed small difference for its root or unknown letter; and, by any means, find, from this equation, a near value of this small assumed quantity. Assume then another letter for the small difference between this last value and the true one, and substitute the sum or difference of them into the last equation, by which will arise a third equation, involving the second assumed quantity; whose near value is found as before. Proceeding thus as far as we please, all the near values, connected together by their proper signs, will form a series approaching still nearer and nearer to the true value of the root of the first or proposed equation. The approximate values of the several small assumed differences, may be found in different ways: Newton's method is this: As the quantity sought is small, its higher powers decrease more and more, and therefore neglecting them will not lead to any great error, Newton therefore neglects all the terms having in them the 2d and higher powers, leaving only the 1st power and the absolute known term; from which simple equation he always finds the value of the assumed unknown letter nearly, in a very simple and easy manner. Halley's method of doing the same thing, was to neglect all the terms above the square or 2d power, and then to find the root of the remaining quadratic equation; which would indeed be a nearer value of the assumed letter than Newton's was, but then it is much more troublesome to perform. — Raphson has another way, which is a little varied from that of New-



ton's again, which is this: having found a near value of the first assumed small quantity or difference, by this he corrects the first approximation to the root of the proposed equation; and then, assuming another letter for the next, or smaller difference, he introduces it into the original equation in the same way as before. And thus he proceeds, from one correction to another, employing always the first proposed equation to find them, instead of the successive new equations used by Newton.

For example, let it be required to find the root of the equation  $x^2 - 5x = 31$ , or  $x^2 - 5x - 31 = 0$ :—Here the root  $x$ , it is evident, is nearly  $= 8$ ; for  $x$  therefore take  $8 + z$ , and substitute  $8 + z$  for  $x$  in the given equation, and the terms will be thus;

$$\begin{array}{r} x^2 = 64 + 16z + z^2 \\ -5x = -40 - 5z \\ -31 = -31 \end{array}$$

the sum is  $-7 + 11z + z^2 = 0$ .

Then, rejecting  $z^2$ , it is  $11z - 7 = 0$ , and  $z = \frac{7}{11} = .6363$ , &c, or  $= .6$  nearly.

Assume now  $z = .6 + y$ : then

$$\begin{array}{r} z^2 = .36 + 1.2y + y^2 \\ 11z = 6.6 + 11y \\ -7 = -7 \end{array}$$

the sum  $-.04 + 12.2y + y^2 = 0$ ,

where  $y = \frac{.04}{12.2} = .003278$  nearly.

Assume it  $y = .003278 - v$ : then

$$\begin{array}{r} y^2 = .000010745284 - .006556v + v^2 \\ 12.2y = .0399916 - 12.2v \\ -.04 = -.04 \end{array}$$

the sum  $.000002345284 - 12.206556v + v^2 = 0$ ,

where  $v = \frac{.000002345284}{12.206556} = .000000192133$ .

Hence, then, collecting all the assumed differences, with their signs, it is found that  $x = 8 + z + y - v = 8 + .6 + .003278 - .000000192133 = 8.603277807867$  the root of the equation required, by Newton's method.

*The same by Raphson's way.*

First  $x = 8 + z$ ;

then  $x^2 = 64 + 16z + z^2$

$$-5x = -40 - 5z$$

$$-31 = -31$$

the sum  $-7 + 11z + z^2 = 0$ ;

hence  $z = \frac{7}{11} = .6$  nearly,

and  $x = 8 + z = 8.6$  nearly.

Assume it  $x = 8.6 + y$ ;

then  $x^2 = 73.96 + 17.2y + y^2$

$$-5x = -43 - 5y$$

$$-31 = -31$$

the sum  $-.04 + 12.2y + y^2 = 0$ ;

hence  $y = \frac{.04}{12.2} = .003278$  nearly,

and  $x = 8.6 + y = 8.603278$  nearly.

Assume it  $x = 8.603278 - v$ ;

then  $x^2 = 74.016392345284 - 17.206556v + v^2$

$$-5x = -43.016390 + 5v$$

$$-31 = -31$$

the sum  $.000002345284 - 12.206556v + v^2 = 0$ ;

hence  $v = .000000192133$ ,

and conseq.  $x = 8.603277807867$ , as before.

EXAMPLE 2. Again, taking the cubic equation  $y^3 - 2y - 5 = 0$ ; Newton proceeds thus:

$y$  is nearly  $= 2$ ; take it therefore  $y = 2 + p$ ;

then  $y^3 = 8 + 12p + 6p^2 + p^3$

$$-2y = -4 - 2p$$

$$-5 = -5$$

the sum  $-1 + 10p + 6p^2 + p^3 = 0$ ;

hence  $p = \frac{1}{10} = .1$  nearly.

Assume it  $p = .1 + q$ ;

then  $p^3 = .001 + .03q + .3q^2 + q^3$

$$+ 6p^2 = .06 + 1.2q + 6q^2$$

$$+ 10p = 1 + 10q$$

$$-1 = -1$$

the sum  $.0061 + 11.23q + 6.3q^2 + q^3 = 0$ ;

hence  $q = -.00054$  nearly.

Assume it  $q = -.00054 + r$ ;

then  $q^3 = -.000000157464 + .000008748r$ , &c.

$$+ 6.3q^2 = +.000183708 - .06804r$$
, &c.

$$+ 11.23q = -.060642 + 11.23r$$

$$+ .0061 = +.0061$$

the sum  $+ .000541550536 + 11.16204748r$ ;

hence  $r = -.000048517$ , &c.

Hence,  $y = 2 + p + q + r$

$$= 2 + .1 - .00054 - .000048517$$

$$= 2.094551483,$$

the root of the equation  $y^3 - 2y = 5$ . And in the same manner Newton performs the approximation for the roots of literal equations, that is, equations having literal coefficients; so the root of this equation

$y^3 + axy + a^2y - x^3 - 2a^3 = 0$ , is

$$y = a - \frac{x}{4} + \frac{x^2}{64a} + \frac{131x^3}{512a^2} + \frac{509x^4}{16384a^3} \text{ \&c.}$$

See also a memoir on this method by the Marquis de Courtivron, in the *Memoires de l'Academie* for 1744.

*Other Methods of APPROXIMATION.* Besides the foregoing general methods, other particular ways of approximating, for various purposes, have been given by many other persons.—As for example, methods of approximating, by series, to the roots of cubic equations belonging to the irreducible case, by Nicole in the same *Memoirs*, by M. Clairaut in his *Algebra*, and by myself in the *Philos. Transf.* for 1780. See also several parts of Simpson's works, and my *Tracts* vol. 1. Also the methods of infinite series by Wallis, Newton, Gregory, Mercator, &c, may be considered as approximations, in quadratures, and other branches of the mathematics, many instances of which may be seen in Wallis's *Algebra*, and other books:—Likewise the method of exhaustions of the ancients, by which Archimedes and others have approximated to the quadrature and rectification of the circle, &c, which was performed by continually bisecting the sides of polygons, both inscribed in a circle and circumscribed about it; by which means the sum of the sides of the like polygons approach continually nearer and nearer together, and the circumference of the circle is nearly a mean between the two sums. See also EQUATIONS.

APPULSE, in Astronomy, means the actual contact of two luminaries, according to some authors; but others describe it as their near approach to each other, so as to be seen, for instance, within the same telescope.

The appulses of the planets to the fixed stars have always



always been very useful to astronomers, as serving to fix and determine the places of the former. The ancients, wanting an easy method of comparing the planets with the ecliptic, which is not visible, had scarce any other way of fixing their situations, but by observing their track among the fixed stars, and marking their appulses to some of those visible points. See Hist. Acad. Scienc. for 1710, pa. 417. And Philos. Transf. No. 369, where Dr. Halley has given a method of determining the places of the planets, by observing their near appulses to the fixed stars. See also Philos. Transf. No. 76, pa. 351, and Mem. Acad. Scienc. for 1708, where Flamsteed and De la Hire have given observations of the moon's appulses to the Pleiades. See also Flamsteed's *Historia Cœlestis*, where a multitude of observations of appulses, or small distances, of the moon and planets, from the fixed stars, are recorded. And Dr. Halley has published a map or planisphere of the starry zodiac, in which are accurately laid down all the stars to which the moon's appulse has ever been observed in any part of the world. See Philos. Transf. No. 369; or Abridg. vol. vi. pa. 170.

APRIL, the 4th month of the year according to the common computation, and the 2d from the vernal equinox.—The word is derived from *Aprilis*, of *aperio*, *I open*; because the earth, in this month, begins to open her bosom for the production of vegetables.—In this month the sun travels through part of the signs Aries and Taurus.

APRON, in Gunnery, a piece of thin or sheet lead, used to cover the vent or touch-hole of a cannon.

APSES, in Astronomy, are the two points in the orbits of planets, where they are at their greatest and least distance, from the sun or the earth. The point at the greatest distance being called the *higher apsis*, and that at the nearest distance the *lower apsis*. And the two apses are also called *auges*. Also the higher apsis is more particularly called the *aphelion*, or the *apogee*; and the lower apsis, the *perihelion*, or the *perigee*. The diameter which joins these two points, is called the *line of the apses* or of the *apsides*; and it passes through the centre of the orbit of the planet, and the centre of the sun or the earth; and in the modern astronomy this line makes the longer or transverse axis of the elliptical orbit of the planet. In this line is counted the excentricity of the orbit; being the distance between the centre of the orbit and the focus, where is placed the sun or the earth.

The foregoing definitions suppose the lines of the greatest and least distances to lie in the same straight line; which is not always precisely the case; as they are sometimes out of a right line, making an angle greater or less than 180 degrees, and the difference from 180 degrees is called the motion of the line of the apses: when the angle is less than 180 degrees, the motion of the apses is said to be contrary to the order of the signs; on the other hand, when the angle exceeds 180 degrees, the motion is according to the order of the signs.

Different means have been employed to determine the motion of the apses. Dr. Keil explains, in his *Astronomy*, the method used by the ancients, who supposed the orbits of the planets to be perfectly circular, and the sun out of the centre. But since it has been

discovered that they describe elliptical orbits, various other methods have been devised for determining it. Halley has given one, which supposes to be known only the time of the planet's revolution, or periodic time. Seth Ward has also given a determination from three different observations of a planet, in any three places of its orbit: but his method being founded on an hypothesis not strictly true, Euler has given one much more exact in vol. 7. of the *Petersburgh Commentaries*. See various ways explained in the *Astronomy* of Keil and Mounier.

Newton has also given, in the *Principia*, an excellent method of determining the same motion, on the supposition that the orbits of the planets differ but little from circles, which is the case nearly. That great philosopher shews, that if the sun be immovable, and all the planets gravitate towards him in the inverse ratio of the squares of their distances, then the apses will be fixed, or their motion nothing; that is, the lines of greatest and least distance will form one right line, and the apses will be directly opposite, or at 180 degrees distance from each other. But, because of the mutual tendency of the planets towards each other, their gravitation towards the sun is not precisely in that ratio; and hence it happens, that the apses are not always exactly in a right line with the sun. And Newton has given a very elegant method of determining the motion of the apses, on the supposition that we know the force which is thereby added to the gravitation of the planet towards the sun, and that this additional force is always in that direction.

APUS or APOUS, *Avis Indica*, in Astronomy, a constellation of the southern hemisphere, situated near the south pole, between the *triangulum australe* and the *chameleon*, and supposed to represent the bird of paradise. Also supposed to be one of the birds named *Apodes*, as having no feet.

The number of stars contained in this constellation, are 11 in the British Catalogue, in Bayer's Maps 12, and a still greater number in La Caille's Catalogue; the principal star being but of the 4th or 5th order of magnitude. See *Calum Australe Stelliferum*, and the *Memoires de l'Acad.* for 1752, pa. 569.

AQUARIUS, in Astronomy, one of the celestial constellations, being the eleventh sign in the zodiac, reckoning from Aries, and is marked by the character  $\text{♒}$ , representing part of a stream of water, issuing from the vessel of Aquarius, or the water-pourer. This sign also gives name to the eleventh part of the ecliptic, through which the sun moves in part of the months of January and February.

The poets feign that Aquarius was Ganymede, whom Jupiter ravished under the shape of an eagle, and carried away into Heaven to serve as a cup-bearer, instead of Hebe and Vulcan; whence the name. Others hold, that the sign was thus called, because that when it appears in the horizon, the weather commonly proves rainy.

The stars in the constellation Aquarius, are, in Ptolemy's Catalogue, 45; in Tycho's 41; in Hevelius's 47; and in Flamsteed's 108. See the article CONSTELLATION; also CATALOGUE.

AQUEDUCT, or AQUÆDUCT, as much as to say *ductus aquæ*, a conduit of water, is a construction of stone



stone or timber built on uneven ground, to preserve the level of water, and convey it, by a canal, from one place to another.—Some aqueducts are under ground, being conducted through hills, &c; and others are raised above ground, and supported on arches, to conduct the water over vallies, &c.

The Romans were very magnificent in their aqueducts; having some that extended a hundred miles, or more. Frontinus, a man of consular dignity, who had the direction of the aqueducts under the emperor Nerva, speaks of nine that emptied themselves through 13594 pipes, of an inch diameter. And it is observed by Vigenere, that in the space of 24 hours, Rome received from these aqueducts not less than 500000 hogheads of water. The chief aqueducts now in being, are these: 1st, that of the Aqua Virginia, repaired by pope Paul IV; 2d, the Aqua Felice, constructed by pope Sixtus V, and is called from the name he assumed before he was exalted to the papal throne; 3d, the Aqua Paulina, repaired by pope Paul V, in the year 1611; and 4thly, the aqueduct built by Lewis XIV, near Maintenon, to convey the river Bure to Versailles, which is perhaps the largest in the world; being 7000 fathoms long, elevated 2560 fathoms in height, and containing 242 arcades. See *Philos. Trans.* for 1685, No. 171; or *Abridg.* vol. 1. pa. 594.

**AQUEOUS HUMOUR**, or the *watry humour* of the eye, is the first or outermost, and the rarest of the three humours of the eye. It is transparent and colourless, like water; and it fills up the space that lies between the cornea tunica, and the crystalline humour.

**AQUILA**, *the Eagle*, or the *Vulture* as it is sometimes called, is a constellation of the northern hemisphere, usually joined with Antinous. It is one of the 48 old constellations, according to the division of which Hipparchus made his Catalogue of the Fixed Stars, and which are described by Ptolemy. The number of stars in Aquila, and those near it, now in the later-formed constellation Antinous, amount to 15 in Ptolemy's Catalogue, to 19 in Tycho's, to 42 in that of Hevelius, and to 71 in Flamsteed's. But in Aquila alone, Tycho counts only 12 stars, and Hevelius 23; the principal star being Lucida Aquila, and is between the 1st and 2d magnitude. The Greeks, as usual, relate various fables of this constellation, to make the science appear as of their own invention.

**ARA**, *the Altar*, one of the 48 old constellations, mentioned by the ancient astronomers, and is situated in the southern hemisphere; containing only 7 stars in Ptolemy's Catalogue, and 9 in that of Flamsteed; none of which exceed the 4th magnitude.

**ARATUS**, celebrated for his Greek poem intitled *Φαινόμενα*, the *Phenomena*, flourished about the 127th Olympiad, or near 300 years before Christ, while Ptolemy Philadelphus reigned in Egypt. Being educated under Dionysius Heracleotes, a Stoic philosopher, he espoused the principles of that sect, and became physician to Antigonus Gonatus, the son of Demetrius Poliorcetes, King of Macedon. The *Phenomena* of Aratus gives him a title to the character of an astronomer, as well as a poet. In this work he describes the nature and motion of the stars, and shews their various dispositions and relations; he describes the figures of

the constellations, their situations in the sphere, the origin of the names which they bear in Greece and in Egypt, the fables which have given rise to them, the rising and setting of the stars, and he indicates the manner of knowing the constellations by their respective situations.

The poem of Aratus was commented upon and translated by many authors: of whom among the ancients were Cicero, Germanicus Cæsar, and Festus Avienus, who made Latin translations of it; a part of the former of which is still extant. Aratus must have been much esteemed by the ancients, since we find so great a number of scholiasts and commentators upon him; among whom are Aristarchus of Samos, the Arystyli the geometricians, Apollonius, the Evæneti, Crates, Numenius the grammarian, Pyrrhus of Magnesia, Thales, Zeno, and many others, as may be seen in Vossius, p. 156. Suidas ascribes several other works to Aratus. Virgil, in his *Georgics*, has translated or imitated many passages from this author: Ovid speaks of him with admiration, as well as many others of the poets: And St. Paul has quoted a passage from him; which is in his speech to the Athenians (*Acts* xvii. 28.) where he tells them that some of their own poets have said, *For we are also his offspring*, these words being the beginning of the 5th line of the *Phenomena* of Aratus.

His modern editors are as follow: Henry Stephens published his poem at Paris in 1566, in his collection of the poets, in folio. Grotius published an edition of the *Phenomena* at Leyden in quarto, 1600, in Greek and Latin, with the fragments of Cicero's version, and the translations of Germanicus and Avienus; all which the editor has illustrated with curious notes. Also a neat and correct edition of Aratus was published at Oxford, 1672, in 8vo. with the Scholia.

**ARÆOMETER**, see **AREOMETER**.

**ARC**, or **ARCH**; which see.

**ARCADE**, in Architecture, denotes an opening in the wall of a building formed by an arch.

**ARC-BOUTANT**, is a kind of arched buttress, formed of a flat arch, or part of an arch, abutting against the feet or sides of another arch or vault, to support them and prevent them from bursting or giving way.

**ARCAS**, a name by which some of the old writers call the star Arcturus; a single and very bright star of the first magnitude, between the legs of the constellation Bootes. They say Arcas, the son of Calisto by Jupiter, when he was about to have killed his mother in the shape of a bear, was, together with her, snatched up into Heaven; where she was converted into the constellation of the Great Bear, near the north pole, and the youth into this single star.

**ARCH**, **ARC**, *Arcus*, in Geometry, a part of any curve line; as, of a circle, or ellipsis, or the like.

It is by means of circular arcs, or arches, that all angles are measured; the arc being described from the angular point as a centre. For this purpose, every circle is supposed to be divided into 360 degrees, or equal parts; and an arch, or the angle it subtends and measures, is estimated according to the number of those degrees it contains: thus, an arc, or angle, is said to be of 30 or 80, or 100 degrees.—Circular arcs are also of great use in finding of fluents.


*Concentric*



*Concentric ARCS*, are such as have the same centre.

*Equal ARCS*, are such arcs, of the same circle, or of equal circles, as contain the same number of degrees. These have also equal chords, sines, tangents, &c.

*Similar ARCS*, of unequal circles, &c, are such as contain the same number of degrees, or that are the like part or parts of their respective whole circles. Hence, in concentric circles, any two radii cut off, or intercept, similar arcs MN and OP.—Similar arcs are proportional to the radii LM, LO, or to the whole circumferences.—Similar arcs of other like curves, are also like parts of the wholes, or determined by like parts alike posited.



*Of the Length of Circular ARCS.* The lengths of circular arcs, as found and expressed in various ways, may be seen in my large Treatise on Mensuration, pa. 118, & seq. 2d edition: some of which are as follow. The radius of a circle being 1; and of any arc  $a$ , if the tangent be  $t$ , the sine  $s$ , the cosine  $c$ , and the versed sine  $v$ : then the arc  $a$  will be truly expressed by several series, as follow, viz, the arc

$$a = t - \frac{1}{3}t^3 + \frac{1}{5}t^5 - \frac{1}{7}t^7 + \frac{1}{9}t^9 \&c.$$

$$a = \frac{s}{c} - \frac{1}{3} \cdot \frac{s^3}{c^3} + \frac{1}{5} \cdot \frac{s^5}{c^5} - \frac{1}{7} \cdot \frac{s^7}{c^7} \&c.$$

$$a = s + \frac{1}{2 \cdot 3} s^3 + \frac{1 \cdot 3}{2 \cdot 4 \cdot 5} s^5 + \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6 \cdot 7} s^7 \&c.$$

$$a = \sqrt{2v} \times \left(1 + \frac{1}{2 \cdot 3} \cdot \frac{v}{2} + \frac{1 \cdot 3}{2 \cdot 4 \cdot 5} \cdot \frac{v^2}{2^2} + \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6 \cdot 7} \cdot \frac{v^3}{2^3} \&c.\right)$$

$$a = \frac{3 \cdot 14159}{180} \&c. \quad d = .01745329 \&c \times d; \text{ where}$$

$d$  denotes the number of degrees in the given arc. Also  $a = \frac{8c - C}{3}$  nearly; where  $C$  is the chord of the arc, and  $c$  the chord of half the arc; whatever the radius is.

*To investigate the length of the arc of any curve.* Put  $x$  = the absciss,  $y$  = the ordinate, of the arc  $z$ , of any curve whatever. Put  $\dot{z} = \sqrt{\dot{x}^2 + \dot{y}^2}$ ; then, by means of the equation of the curve, find the value of  $\dot{x}$  in terms of  $y$ , or of  $\dot{y}$  in terms of  $\dot{x}$ , and substitute that value instead of it in the above expression  $\dot{z} = \sqrt{\dot{x}^2 + \dot{y}^2}$ ; hence, taking the fluents, they will give the length of the arc  $z$ , in terms of  $x$  or  $y$ .

*ARCH*, in Astronomy. Of this, there are various kinds. Thus, the latitude, elevation of the pole, and the declination, are measured by an arch of the meridian; and the longitude, by an arch of a parallel circle, &c.

*Diurnal ARCH* of the sun, is part of a circle parallel to the equator, described by the sun in his course from his rising to the setting. And his *Nocturnal ARCH* is of the same kind; excepting that it is described from setting to rising.

*ARCH of Progression*, or *Direction*, is an arch of the ecliptic, which a planet seems to pass over, when its motion is direct, or according to the order of the signs.

*ARCH of Retrogradation*, is an arch of the ecliptic,

described while a planet is retrograde, or moves contrary to the order of the signs.

*ARCH between the Centres*, in eclipses, is an arch passing from the centre of the earth's shadow, perpendicular to the moon's orbit, meeting her centre at the middle of an eclipse.—If the aggregate of this arch and the apparent semi-diameter of the moon, be equal to the semi-diameter of the shadow, the eclipse will be total for an instant, or without any duration; and if that sum be less than the radius of the shadow, the eclipse will be total, with some duration; but if greater, the eclipse will be only partial.

*ARCH of Vision*, is that which measures the sun's depth below the horizon, when a star, before hid by his rays, begins to appear again.—The quantity of this arch is not always the same, but varies with the latitude, declination, right ascension, or descension, and distance, of any planet or star. Ricciol. Almag. v. 1, pa. 42. However, the following numbers will serve nearly for the stars and planets.

TABLE exhibiting the *Arch of Vision* of the PLANETS and FIXED STARS.

PLANETS.		FIXED STARS.	
		Magnitude.	
Mercury	- - 10° 0'	1	- - - - 12°
Venus	- - - 5 0	2	- - - - 13
Mars	- - 11 30	3	- - - - 14
Jupiter	- - 10 0	4	- - - - 15
Saturn	- - 11 0	5	- - - - 16
		6	- - - - 17

*ARCH*, in Architecture, is a concave structure, raised or turned upon a mould, called the centering, in form of the arch of a curve, and serving as the inward support of some superstructure. Sir Henry Wotton says, An arch is nothing but a narrow or contracted vault; and a vault is a dilated arch.

Arches are used in large intercolumnations of spacious buildings; in porticoes, both within and without temples; in public halls, as ceilings, the courts of palaces, cloisters, theatres, and amphitheatres. They are also used to cover the cellars in the foundations of houses, and powder magazines; also as buttresses and counter-forts, to support large walls laid deep in the earth; for triumphal arches, gates, windows, &c; and, above all, for the foundations of bridges and aqueducts. And they are supported by piers, buttments, imposts, &c.

Arches are of several kinds, and are commonly denominated from the figure or curve of them; as circular, elliptical, cycloidal, catenarian, &c, according as their curve is in the form of a circle, ellipse, cycloid, catenary, &c.

There are also other denominations of circular arches, according to the different parts of a circle, or manner of placing them. Thus,

*Semicircular ARCHES*, which are those that make an exact semicircle, having their centre in the middle of the span or chord of the arch; called also by the French builders, perfect arches, and *arches en plein centre*. The arches of Westminster Bridge are semicircular.

*Scheme ARCHES*, or *skene*, are those which are less than semicircles, and are consequently flatter arches; containing-



containing 120, or 90, or 60, degrees, &c. They are also called *imperfect* and *diminished* arches.


ARCHES of the *third and fourth point*, or *Gothic arches*; or, as the Italians call them, *di terzo* and *quarto acuto*, because they always meet in an acute angle at top. These consist of two excentric circular arches, meeting in an angle above, and are drawn from the division of the chord into three or four or more parts at pleasure. Of this kind are many of the arches in churches and other old Gothic buildings.

*Elliptical* ARCHES, usually consist of semi-ellipses; and were formerly much used instead of mantle-trees in chimnies; and are now much used, from their bold and beautiful appearance, for many purposes, and particularly for the arches of a bridge, like that at Black-Friars, both for their strength, beauty, convenience, and cheapness.

*Straight* ARCHES, are those which have their upper and under edges parallel straight lines, instead of curves. These are chiefly used over doors and windows; and have their ends and joints all pointing towards one common centre.

ARCH is particularly used for the space between the two piers of a bridge, intended for the passage of the water, boats, &c.

ARCH of *equilibration*, is that which is in equilibrium in all its parts, having no tendency to break in one part more than in another, and which is therefore safer and stronger than any other figure. Every particular figure of the extrados, or upper side of the wall above an arch, requires a peculiar curve for the under side of the arch itself, to form an arch of equilibration, so that the incumbent pressure on every part may be proportional to the strength or resistance there. When the arch is equally thick throughout, a case that can hardly ever happen, then the catenarian curve is the arch of equilibration; but in no other case: and therefore it is a great mistake in some authors to suppose that this curve is the best figure for arches in all cases; when in reality it is commonly the worst. This subject is fully treated in my *Principles of Bridges*, pr. 5, where the proper intrados is investigated for every extrados, so as to form an arch of equilibration in all cases whatever. It there appears that, when the upper side of the wall is a straight horizontal line, as in the annexed figure, the equation



of the curve is thus expressed,

$$y = b \times \frac{\log. \frac{a+x+\sqrt{2ax+xx}}{a}}{\log. \frac{a+r+\sqrt{2ar+rr}}{a}};$$

where  $x = DP$ ,  $y = PC$ ,  $r = DQ$ ,  $b = AQ$ , and  $a = DK$ . And hence, when  $a$ ,  $b$ ,  $r$ , are any given numbers, a table is formed for the corresponding values of  $x$  and  $y$ , by which the curve is constructed for any particular occasion. Thus supposing  $a$  or  $DK = 6$ ,  $b$  or  $AQ = 50$ , and  $r$  or  $DQ = 40$ ; then the corresponding values of  $KI$  and  $IC$ , or horizontal and vertical lines, will be as in this table.

Table for constructing the Curve of Equilibration.

Value of KI.	Value of IC.	Value of KI.	Value of IC.	Value of KI.	Value of IC.
0	6.000	21	10.381	36	21.774
2	6.035	22	10.858	37	22.948
4	6.144	23	11.368	38	24.190
6	6.324	24	11.911	39	25.505
8	6.580	25	12.489	40	26.894
10	6.914	26	13.106	41	28.364
12	7.330	27	13.761	42	29.919
13	7.571	28	14.457	43	31.563
14	7.834	29	15.196	44	33.299
15	8.120	30	15.980	45	35.135
16	8.430	31	16.811	46	37.075
17	8.766	32	17.693	47	39.126
18	9.168	33	18.627	48	41.293
19	9.517	34	19.617	49	43.581
20	9.934	35	20.665	50	46.000

The doctrine and use of arches are neatly delivered by Sir Henry Wotton, though he is not always mathematically accurate in the principles. He says; *First*, All matter, unless impeded, tends to the centre of the earth in a perpendicular line. *Secondly*; All solid materials, as bricks, stones, &c, in their ordinary rectangular form, if laid in numbers, one by the side of another, in a level row, and their extreme ones sustained between two supporters; those in the middle will necessarily sink, even by their own gravity, much more if forced down by any superincumbent weight. To make them stand, therefore, either their figure or their position must be altered.—*Thirdly*; Stones, or other materials, being figured *cuneatim*, or wedge-like, broader above than below, and laid in a level row, with their two extremes supported as in the last article, and pointing all to the same centre; none of them can sink, till the supporters or buttments give way, because they want room in that situation to descend perpendicularly. But this is a weak structure; because the supporters are subject to too much impulsion, especially where the line is long; for which reason the form of straight arches is seldom used, excepting over doors and windows, where the line is short and the side walls strong. In order to fortify the work, therefore, we must change not only the figure of the materials, but also their position.—*Fourthly*; If the materials be shaped wedge-wise, and be disposed in form of an arch, and pointing to some centre; in this case, neither the pieces of the said arch can sink downwards, for want of room to descend perpendicularly; nor can the supporters or buttments suffer much violence, as in the preceding flat form: for the convexity will always make the incumbent rather rest upon the supporters, than thrust or push



push them outwards. His reasoning, however, afterwards, on the effect of circular and other arches, is not accurate, as he attends only to the side pressure, without considering the effect of different vertical pressures.

The chief properties of arches of different curves, may be seen in the 2d sect. of my *Principles of Bridges*, above quoted. It there appears that none, except the mechanical curve of the arch of equilibration, can admit of a horizontal line at top: that this arch is of a form both graceful and convenient, as it may be made higher or lower at pleasure, with the same span or opening: that all other arches require extrados that are curved, more or less, either upwards or downwards: of these, the elliptical arch approaches the nearest to that of equilibration for equality of strength and convenience; and it is also the best form for most bridges, as it can be made of any height to the same span, its hanches being at the same time sufficiently elevated above the water, even when it is very flat at top: elliptical arches also look bolder and lighter, are more uniformly strong, and much cheaper than most others, as they require less materials and labour. Of the other curves, the cycloidal arch is next in quality to the elliptical one, for those properties, and, lastly, the circle. As to the others, the parabola, hyperbola, and catenary, they are quite inadmissible in bridges that consist of several arches; but may, in some cases, be employed for a bridge of one single arch which may be intended to rise very high, as in such cases as they are not much loaded at the hanches.

ARCH Mural. See MURAL arch.

ARCHER, or *Sagittarius*, one of the constellations of the northern hemisphere, and one of the twelve signs of the zodiac, placed between the Scorpion and Capricorn. See SAGITTARIUS.

ARCHIMEDES, one of the most celebrated mathematicians among the ancients, who flourished about 250 years before Christ, being about 50 years later than Euclid. He was born at Syracuse in Sicily, and was related to Hiero, who was then king of that city. The mathematical genius of Archimedes set him with such distinguished excellence in the view of the world, as rendered him both the honour of his own age, and the admiration of posterity. He was indeed the prince of the ancient mathematicians, being to them what Newton is to the moderns, to whom in his genius and character he bears a very near resemblance. He was frequently lost in a kind of reverie, so as to appear hardly sensible; he would study for days and nights together, neglecting his food; and Plutarch tells us that he used to be carried to the baths by force. Many particulars of his life, and works, mathematical and mechanical, are recorded by several of the ancients, as Polybius, Livy, Plutarch, Pappus, &c. He was equally skilled in all the sciences, astronomy, geometry, mechanics, hydrostatics, optics, &c, in all of which he excelled, and made many and great inventions.

Archimedes, it is said, made a sphere of glass, of a most surprising contrivance and workmanship, exhibiting the motions of the heavenly bodies in a very pleasing manner. Claudian has an epigram upon this invention, which has been thus translated:

When in a glass's narrow space confin'd,  
Jove saw the fabric of th' almighty mind;

VOL. I.

He smil'd, and said, Can mortals' art alone,  
Our heavenly labours mimic with their own?  
The Syracusan's brittle work contains  
Th' eternal law, that through all nature reigns.  
Fram'd by his art, see stars unnumber'd burn,  
And in their courses rolling orbs return:  
His sun through various signs describes the year;  
And every month his mimic moons appear.  
Our rival's laws his little planets bind,  
And rule their motions with a human mind.  
Salmoneus could our thunder imitate,  
But Archimedes can a world create.

Many wonderful stories are told of his discoveries, and of his very powerful and curious machines, &c. Hiero once admiring them, Archimedes replied, these effects are nothing, "But give me, said he, some other place to fix a machine on, and I shall move the earth." He fell upon a curious device for discovering the deceit which had been practiced by a workman, employed by the said king Hiero to make a golden crown. Hiero, having a mind to make an offering to the gods of a golden crown, agreed for one of great value, and weighed out the gold to the artificer. After some time he brought the crown home of the full weight; but it was afterwards discovered or suspected that a part of the gold had been stolen, and the like weight of silver substituted in its stead. Hiero, being angry at this imposition, desired Archimedes to take it into consideration, how such a fraud might be certainly discovered. While engaged in the solution of this difficulty, he happened to go into the bath; where observing that a quantity of water overflowed, equal to the bulk of his body, it presently occurred to him, that Hiero's question might be answered by a like method: upon which he leaped out, and ran homeward, crying out εὕρηκα! εὕρηκα! I have found it! I have found it! He then made two masses, each of the same weight as the crown, one of gold and the other of silver: this done, he filled a vessel to the brim with water, and put the silver mass into it, upon which a quantity of water overflowed equal to the bulk of the mass; then taking the mass of silver out he filled up the vessel again, measuring the water exactly, which he put in; this shewed him what measure of water answered to a certain quantity of silver. Then he tried the gold in like manner, and found that it caused a less quantity of water to overflow, the gold being less in bulk than the silver, though of the same weight. He then filled the vessel a third time, and putting in the crown itself, he found that it caused more water to overflow than the golden mass of the same weight, but less than the silver one; so that, finding its bulk between the two masses of gold and silver, and that in certain known proportions, he hence computed the real quantities of gold and silver in the crown, and so manifestly discovered the fraud.

Archimedes also contrived many machines for useful and beneficial purposes: among these, engines for launching large ships; screw pumps, for exhausting the water out of ships, marshes or overflowed lands, as Egypt, &c, which they would do from any depth.

But he became most famous by his curious contrivances, by which the city of Syracuse was so long defended, when besieged by the Roman consul Marcellus; showering upon the enemy sometimes long darts, and stones of vast weight and in great quantities; at other times lifting their ships up into the air, that had come near

T

the



the walls, and dashing them to pieces by letting them fall down again; nor could they find their safety in removing out of the reach of his cranes and levers, for there he contrived to fire them with the rays of the sun reflected from burning glasses.

However, notwithstanding all his art, Syracuse was at length taken by storm, and Archimedes was so very intent upon some geometrical problem, that he neither heard the noise, nor minded any thing else, till a soldier that found him tracing of lines, asked him his name, and upon his request to begone, and not disorder his figures, slew him. "What gave Marcellus the greatest concern, says Plutarch, was the unhappy fate of Archimedes, who was at that time in his museum; and his mind, as well as his eyes, so fixed and intent upon some geometrical figures, that he neither heard the noise and hurly of the Romans, nor perceived the city to be taken. In this depth of study and contemplation, a soldier came suddenly upon him, and commanded him to follow him to Marcellus; which he refusing to do, till he had finished his problem, the soldier, in a rage, drew his sword, and ran him through." Livy says he was slain by a soldier, not knowing who he was, while he was drawing schemes in the dust: that Marcellus was grieved at his death, and took care of his funeral; and made his name a protection and honour to those who could claim a relationship to him. His death it seems happened about the 142 or 143 Olympiad, or 210 years before the birth of Christ.

When Cicero was questor for Sicily, he discovered the tomb of Archimedes, all overgrown with bushes and brambles; which he caused to be cleared, and the place set in order. There was a sphere and cylinder cut upon it, with an inscription, but the latter part of the verses quite worn out.

Many of the works of this great man are still extant, though the greatest part of them are lost. The pieces remaining are as follow: 1. Two books on the Sphere and Cylinder.—2. The Dimension of the Circle, or proportion between the diameter and the circumference.—3. Of Spiral lines.—4. Of Conoids and Spheroids.—5. Of Equiponderants, or Centres of Gravity.—6. The Quadrature of the Parabola.—7. Of Bodies floating on Fluids.—8. Lemmata.—9. Of the Number of the Sand.

Among the works of Archimedes which are lost, may be reckoned the descriptions of the following inventions, which may be gathered from himself and other ancient authors. 1. His account of the method which he employed to discover the mixture of gold and silver in the crown, mentioned by Vitruvius.—2. His description of the Cochleon, or engine to draw water out of places where it is stagnated, still in use under the name of Archimedes's Screw. Athenæus, speaking of the prodigious ship built by the order of Hiero, says, that Archimedes invented the cochleon, by means of which the hold, notwithstanding its depth, could be drained by one man. And Diodorus Siculus says, that he contrived this machine to drain Egypt, and that by a wonderful mechanism it would exhaust the water from any depth.—3. The Helix, by means of which, Athenæus informs us, he launched Hiero's great ship.—4. The Trispaston, which, according to Tzetzes and Oribasius, could draw the most stupendous weights.—5. The machines, which, according to Polybius, Livy, and Plutarch,

he used in the defence of Syracuse against Marcellus, consisting of Tormenta, Balistæ, Catapults, Sagittarii, Scorpions, Cranes, &c.—6. His Burning Glasses, with which he set fire to the Roman galleys.—7. His Pneumatic and Hydrostatic engines, concerning which subjects he wrote some books, according to Tzetzes, Pappus, and Tertullian.—8. His Sphere, which exhibited the celestial motions. And probably many others.

A whole volume might be written upon the curious methods and inventions of Archimedes, that appear in his mathematical writings now extant only. He was the first who squared a curvilinear space; unless Hypocrates must be excepted on account of his lunes. In his time the conic sections were admitted into geometry, and he applied himself closely to the measuring of them, as well as other figures. Accordingly he determined the relations of spheres, spheroids, and conoids, to cylinders and cones; and the relations of parabolas to rectilinear planes whose quadratures had long before been determined by Euclid. He has left us also his attempts upon the circle: he proved that a circle is equal to a right-angled triangle, whose base is equal to the circumference, and its altitude equal to the radius; and consequently, that its area is equal to the rectangle of half the diameter and half the circumference; thus reducing the quadrature of the circle to the determination of the ratio between the diameter and circumference; which determination however has never yet been done. Being disappointed of the *exact* quadrature of the circle, for want of the rectification of its circumference, which all his methods would not effect, he proceeded to assign an useful approximation to it: this he effected by the numeral calculation of the perimeters of the inscribed and circumscribed polygons: from which calculation it appears that the perimeter of the circumscribed regular polygon of 192 sides, is to the diameter, in a less ratio than that of  $3\frac{1}{7}$  or  $3\frac{1}{7}\frac{9}{10}$  to 1; and that the perimeter of the inscribed polygon of 96 sides, is to the diameter, in a greater ratio than that of  $3\frac{1}{7}\frac{9}{10}$  to 1; and consequently that the ratio of the circumference to the diameter, lies between these two ratios. Now the first ratio, of  $3\frac{1}{7}$  to 1, reduced to whole numbers, gives that of 22 to 7, for  $3\frac{1}{7} : 1 :: 22 : 7$ ; which therefore is nearly the ratio of the circumference to the diameter. From this ratio between the circumference and the diameter, Archimedes computed the approximate area of the circle, and he found that it is to the square of the diameter, as 11 is to 14. He determined also the relation between the circle and ellipse, with that of their similar parts. And it is probable that he likewise attempted the hyperbola; but it is not to be expected that he met with any success, since approximations to its area are all that can be given by the various methods that have since been invented.

Beside these figures, he determined the measures of the spiral, described by a point moving uniformly along a right line, the line at the same time revolving with a uniform angular motion; determining the proportion of its area to that of the circumscribed circle, as also the proportion of their sectors.

Throughout the whole works of this great man, we every where perceive the deepest design, and the finest invention. He seems to have been, with Euclid, exceedingly careful of admitting into his demonstrations nothing



nothing but principles perfectly geometrical and unexceptionable: and although his most general method of demonstrating the relations of curved figures to straight ones, be by inscribing polygons in them; yet to determine those relations, he does not increase the number, and diminish the magnitude, of the sides of the polygon *ad infinitum*; but from this plain fundamental principle, allowed in Euclid's Elements, (viz, that any quantity may be so often multiplied, or added to itself, as that the result shall exceed any proposed finite quantity of the same kind,) he proves that to deny his figures to have the proposed relations, would involve an absurdity. And when he demonstrated many geometrical properties, particularly in the parabola, by means of certain progressions of numbers, whose terms are similar to the inscribed figures; this was still done without considering such series as continued *ad infinitum*, and then collecting or summing up the terms of such infinite series.

There have been various editions of the existing writings of Archimedes. The whole of these works, together with the commentary of Eutocius, were found in their original Greek language, on the taking of Constantinople, from whence they were brought into Italy; and here they were found by that excellent mathematician John Muller, otherwise called Regiomontanus, who brought them into Germany: where they were, with that Commentary, published long afterwards, viz, in 1544, at Basil, being most beautifully printed in folio, both in Greek and Latin, by Hervagius, under the care of Thomas Gechauff Venatorius.—A Latin translation was published at Paris 1557, by Pascalius Hamellius.—Another edition of the whole, in Greek and Latin, was published at Paris 1615, in folio, by David Rivaltus, illustrated with new demonstrations and commentaries: a life of the author is prefixed; and at the end of the volume is added some account, by way of restoration, of our author's other works, which have been lost; viz, The Crown of Hiero; the Cochleon or Water Screw; the Helicon, a kind of endless screw; the Trispaſton, consisting of a combination of wheels and axles; the Machines employed in the defence of Syracuse; the Burning Speculum; the Machines moved by Air and Water; and the Material Sphere.—In 1675, Dr. Isaac Barrow published a neat edition of the works, in Latin, at London, in 4to; illustrated, and succinctly demonstrated in a new method.—But the most complete of any, is the magnificent edition, in folio, lately printed at the Clarendon press, Oxford, 1792. This edition was prepared ready for the press by the learned Joseph Torelli, of Verona, and in that state presented to the University of Oxford. The Latin translation is a new one. Torelli also wrote a preface, a commentary on some of the pieces, and notes on the whole. An account of the life and writings of Torelli is prefixed, by Clemens Sibiliati. And at the end a large appendix is added, in two parts; the first being a Commentary on Archimedes's paper upon Bodies that float on Fluids, by the Rev. Abram Robertson of Christ Church College; and the latter is a large collection of various readings in the Manuscript works of Archimedes, found in the library of the late king of France, and of another at Florence, as collated with the Basil edition above mentioned.

There are also extant other editions of certain parts

of the works of Archimedes. Thus, Commandine published, in 4to, at Bologna 1565, the two books concerning Bodies that Float upon Fluids, with a Commentary. Commandine published also a translation of the Arenarius. And Borelli published, in folio, at Florence 1661, Archimedes's *Liber Assumptorum*, translated into Latin from an Arabic manuscript copy. This is accompanied with the like translation, from the Arabic, of the 5th, 6th, and 7th books of Apollonius's Conics. Mr. G. Anderson published (in 8vo. Lond. 1784) an English translation of the Arenarius of Archimedes, with learned and ingenious notes and illustrations. Dr. Wallis published a translation of the Arenarius. And there may be other editions beside the above, but these are all that I have got, or know of.

ARCHIMEDES's *Screw*. See *SCREW of Archimedes*.

ARCHIMEDES's *Burning-glass*. See *BURNING-glass*.

ARCHITECT, a person skilled in architecture, or the art of building; who forms plans and designs for edifices, conducts the work, and directs the various artificers employed in it.

The most celebrated architects are, Vitruvius, Palladio, Scamozzi, Serlio, Vignola, Barbaro, Cataneo, Alberti, Viola, Inigo Jones, De Lorme, Perrault, S. Le Clerc, Sir Christopher Wren, and the Earl of Burlington.

ARCHITECTURE, *Architectura*, the art of planning and building or erecting any edifice, so as properly to answer the end proposed, for solidity, convenience, and beauty; whether houses, temples, churches, bridges, halls, theatres, &c, &c.—Architecture is divided into civil, military, and naval or marine.

*Civil* ARCHITECTURE, is the art of designing and erecting edifices of every kind for the uses of civil life in every capacity; as churches, palaces, private houses, &c; and it has been divided into five orders or manners of building, under the names of the *Tuscan*, *Doric*, *Ionian*, *Corinthian*, and *Composite*.

There were many authors on architecture among the Greeks and Romans, before Vitruvius; but he is the first whose work is entire and extant. He lived in the reigns of Julius Cæsar and Augustus, and composed a complete system of architecture, in ten books, which he dedicated to this prince. The principal authors on architecture since Vitruvius, are Philander, Barbarus, Salmasius, Baldus, Alberti, Gauricus, Demoniosius, Perrault, De l'Orme, Rivius, Wotton, Serlio, Palladio, Strada, Vignola, Scamozzi, Dénissart, Catanei, Freard, De Cambray, Blondel, Goldman, Sturmy, Wolfius, De Rossi, Desgodetz, Baratteri, Mayer, Gulielmus, Ware, &c, &c. See also ARCHITECT.

*Military* ARCHITECTURE, otherwise more usually called *Fortification*, is the art of strengthening and fortifying places, to screen them from the insults or attacks of enemies, and the violence of arms; by erecting forts, castles, and other fortresses, with ramparts, bastions, &c.—The authors who have chiefly excelled in this art, are Coehorn, Pagan, Vauban, Scheiter, Blondel, and Montalembert.

*Naval* ARCHITECTURE, or *Ship-building*, is the art of constructing ships, galleys, and other vessels proper to float on the water.

ARCHITRAVE, is that part of a column which bears immediately upon the capital. It is the lowest member



member of the entablature, and is supposed to represent the principal beam in timber buildings, in which it is sometimes called the *reason piece*, or *master-piece*. Also, in chimneys it is called the *mantle-piece*; and the *hyperthyron* over the jaumbs of doors, or lintels of windows.

ARCHITRAVE *Corniche*. See CORNICHE.

ARCHITRAVE *doors*, are those which have an architrave on the jaumbs, and over the door; upon the cap piece if straight; or on the arch, if the top be curved.

ARCHITRAVE *windows*, of timber, are usually an ogee raised out of the solid timber, with a list over it: though sometimes the mouldings are struck, and laid on; and sometimes they are cut in brick.

ARCHIVOLT, the contour of an arch; or a band or frame adorned with mouldings, running over the faces of the voussiors or arch-stones, and bearing upon the imposts.

ARCHYTAS, of Tarentum, a celebrated mathematician, cosmographer, and Pythagorean philosopher, whom Horace calls

——— Maris ac Terræ, numeroque carentis Arenæ  
Menforem.

He flourished about 400 years before Christ; and was the master of Plato, Eudoxus, and Philolaus. He gave a method of finding two mean proportionals between two given lines, and thence the Duplication of the Cube, by means of the conic sections. His skill in Mechanics was such, that he was said to be the inventor of the crane and the screw; and he made a wooden pigeon that could fly about, when it was once set off, but it could not rise again of itself, after it rested. He wrote several works, though none of them are now extant, particularly a treatise *περὶ τῆς Παντός, de Universo*, cited by Simplicius in *Aristot. Categ.* It is said he invented the ten categories. He acquired great reputation both in his legislative and military capacity; having commanded an army seven times without ever being defeated. He was at last shipwrecked, and drowned in the Adriatic sea.

ARCTIC *Circle*, is a lesser circle of the sphere, parallel to the equator, and passing through the north pole of the ecliptic, or distant from the north or arctic pole, by a quantity equal to the obliquity of the ecliptic, which was formerly estimated at  $23^{\circ} 30'$ , but its mean quantity is now  $23^{\circ} 28'$  nearly. This, and its opposite, the antarctic circle, are also called the polar circles, where the longest day and longest night are 24 hours, and within all the space of these circles, at one time of the year, the sun never sets, and at the opposite season he never rises for some days, more or less according as the place is nearer the pole.

ARCTIC *Pole*, the north pole of the world, and so called from *αρκτος, ursa*, the *bear*, from its proximity to the constellation of that name.

ARCTOPHYLAX, a constellation otherwise called BOOTES. Which see.

ARCTURUS, a fixed star of the first magnitude, between the thighs of the constellation Bootes. So called from *αρκτος, bear*, and *οψα, tail*; as being near the *bear's tail*.

This star is twice mentioned in the book of Job, viz, ix. 9, and xxxviii. 32, by the name *Aish* if the translation

be right; and by many of the ancients under its Greek name *Arcturus*. The Greeks gave the fabulous history of this star, or constellation, to this purport: Calisto, who was afterwards, in form of the great bear, raised up into a constellation, they tell us, brought forth a son to Jupiter, whom they called Arcas. That Lyacon, when Jupiter afterwards came to visit him, cut the boy in pieces, and served him up at table. Jupiter, in revenge, as well as by way of punishment, called down lightning to consume the palace, and turned the monarch into a wolf. The limbs of the boy were gathered up, to which the god gave life again, and he was taken and educated by some of the people. His mother, who was all this time a bear in the woods, fell in his way: he chased her, ignorant of the fact, and, to avoid him, she threw herself into the temple of Jupiter: he followed her thither to destroy her; and this being death by the laws of the country, Jupiter took them both up into heaven, to prevent the punishment, making her the constellation of the great bear, and converting the youth into this single star behind her.


Dr. Hornsby, the Savilian Professor of Astronomy, concludes that Arcturus is the nearest star to our system visible in the northern hemisphere, because the variation of its place, in consequence of a proper motion of its own, is more remarkable than that of any other of the stars; and by comparing a variety of observations respecting both the quantity and direction of the motion of this star, he infers, that the obliquity of the ecliptic decreases at the rate of  $58''$  in 100 years; a quantity which nearly corresponds to the mean of the computations framed by Euler and De la Lande, upon the principles of attraction. *Philos. Trans.* v. 63.

ARCTUS, a name given by the Greeks to two constellations of the northern hemisphere; by the Latins called *URSA major* and *minor*, and by us the *greater* and *lesser BEAR*.

AREA, in general, denotes any plain surface to walk upon; and derived from *arere, to be dry*.

AREA, in *Architecture*, denotes the space or scite of ground on which an edifice stands. It is also used for inner courts, and such like portions of ground.

AREA, in *Geometry*, denotes the superficial content of any figure. The areas of figures are estimated in squares and parts of squares. Thus, suppose a rectangle EFGH have its length EH equal to 4 inches, or feet, or yards, &c, and its breadth EF equal to 3; its area will then be 3 times 4, or 12 squares, each side of which is respectively one inch, or foot, or yard, &c. The areas of other particular figures may be seen under their respective names.



The areas of all similar figures, are in the duplicate ratio, or as the squares of their like sides, or of any like linear dimensions.—Also the law by which the planets move round the sun, is regulated by the areas described by a line connecting the sun and planet; that is, the time in which the planet describes, or passes over, any arc of its elliptic orbit, is proportional to the elliptic area described in that time by the said line, or the sector contained by the said arc and two radii drawn



drawn from its extremities to the focus in which the sun is placed.

AREA, in *Optics*. See FIELD.

ARENARIUS, the name of a book of Archimedes, in which he demonstrated, that not only the sands of the earth, but even a greater quantity of particles than could be contained in the immense sphere of the fixed stars, might be expressed by numbers, in a way by him invented and described. This notation proceeds by certain geometrical progressions; and in denoting and producing certain very distant terms of the progression, he here first of any one makes use of a property similar to that of logarithms, viz, adding the indices of the terms, to find the index of the product of them. See ARCHIMEDES.

AREOMETER, ARÆOMETRUM, an instrument to measure the density or gravity of fluids.

The areometer, or water-poise, is commonly made of glass; consisting of a round hollow ball, which terminates in a long slender neck, hermetically sealed at top; having first as much running mercury put into it, as will serve to balance or keep it swimming in an erect position. The stem, or neck, is divided into degrees or parts which are numbered, to shew, by the depth of its descent into any liquor, the lightness or density of it: for that fluid is heaviest in which it sinks least, and lightest in which it sinks deepest.

Another instrument of this kind is described by Homberg of Paris, in the Memoirs of the Acad. of Sciences for the year 1699; also in the Philos. Transf. N° 262, where a table of numbers is given, expressing the density of various fluids, as determined by this instrument both in summer and winter. By this table it appears that the density, or specific gravity of quicksilver and distilled water, in the two seasons, were as follow, viz,

in summer as	-	13.61 to 1,
in winter as	-	13.53 to 1;

and the medium of these two is as 13.57 to 1.

See also the Philos. Transf. vol. 36, or Abridg. vol. 6, for the description and use of another new areometer.

AREOMETRY, the science of measuring the lightness and density of fluids.—See the Philos. Transf. vol. 68, for an essay on areometry, &c.

AREOSTYLE, in Architecture, a sort of intercolumniation in which the columns were placed at a great distance from one another.

ARGENTICOMUS, among Ancient Astrologers, denotes a kind of silver-haired comet, of uncommon lustre, supposed to be the cause of great changes in the planetary system.

ARGETENAR, a star of the fourth magnitude, in the flexure of the constellation Eridanus.

ARGO NAVIS, or *the ship*, is a constellation of fixed stars, in the southern hemisphere, being one of the 48 old constellations. The number of stars in this constellation, are, in Ptolemy's catalogue 45, in Tycho Brahe's 11, in Flamsteed's 64.

The Greeks tell us, that this was the famous ship in which the Argonauts performed that celebrated expedition, which has been so famous in all their history.

ARGUMENT, in Astronomy, is an arch given, by which another arch is found in some proportion to it. Hence,

ARGUMENT of *Inclination*, or ARGUMENT of *Latitude*, of any planet, is an arch of a planet's orbit, intercepted between the ascending node, and the place of the planet from the sun, numbered according to the succession of the signs.

*Menstrual ARGUMENT of Latitude*, is the distance of the moon's true place from the sun's true place.—By this is found the quantity of the real obscuration in eclipses, or how many digits are darkened in any place.

*Annual ARGUMENT of the moon's apogee*, or simply, *Annual Argument*, is the distance of the sun's place from the place of the moon's apogee; that is, the arc of the ecliptic comprised between those two places.

ARIES, or the *Ram*, in *Astronomy*, one of the constellations of the northern hemisphere, and the first of the old twelve signs of the zodiac, and marked ♈ in imitation of a ram's head. It gives name to a twelfth part of the ecliptic, which the sun enters commonly about the 20th of March.—The stars of this constellation in Ptolemy's catalogue are 18, in Tycho Brahe's 21, in Hevelius's 27, and in Flamsteed's 66: but they are mostly very small, only one being of the 2d magnitude, two of the 3d magnitude, and all the rest smaller.

The fabulous account of this constellation, as given by the Greeks, is to this effect. That Nephele gave Phryxus, her son, a ram, which bore a golden fleece, as a guard against the greatest dangers. Juno, the step-mother both of him and Helle, laid designs against their lives; But Phryxus, remembering the admonition of his mother, took his sister with him, and getting upon the back of the ram, they were carried to the sea. The ram plunged in, and the youth was carried over; but Helle dropped off, and was drowned, and so gave name to the Hellespont. When he arrived in Colchis, Æeta, the king, received him kindly; and, sacrificing the ram to Jupiter, dedicated the fleece to the god; which was afterwards carried off by Jason. The animal itself, they say, Jupiter snatched up into the heavens, and made of it the constellation Aries. They have other fables also to account for its origin. But it is most probable that the inventors of this sign, placed it there as the father of those animals which are brought forth about the time the sun approaches to that part of the heavens, and so marking the beginning of spring.

ARIES also denotes a *battering ram*; being a military engine with an iron head, much used by the ancients, to batter and beat down the walls of places besieged. See RAM, and BATTERING ram.

ARISTARCHUS, a celebrated Greek philosopher and astronomer, and a native of the city of Samos; but of what date is not exactly known; it must have been however before the time of Archimedes, as some parts of his writings and opinions are cited by that author, viz, in his *Arenarius*; he probably flourished about 420 years before Christ. He held the opinion of Pythagoras as to the system of the world, but whether before or after him, is uncertain, teaching that the sun and stars were fixed in the heavens, and that the earth moved in a circle about the sun, at the same time that it revolved about its own centre or axis. He taught also, that the annual orbit of the earth, compared with the distance of the fixed stars, is but as a point. On this head Archimedes



chimedæ says, "Aristarchus, the Samian, confuting the notions of astrologers, laid down certain positions, from whence it follows, that the world is much larger than is generally imagined; for he lays it down, that the fixed stars and the sun are immoveable; and that the earth is carried round the sun in the circumference of a circle." On which account, although he might not suffer persecution and imprisonment like Galileo, yet he did not escape censure for his supposed impiety; for it is said Cleanthus was of opinion, that Greece ought to have tried Aristarchus for irreligion, for endeavouring to preserve the regular appearance of the heavenly bodies, by supposing that the heavens themselves stood still; but that the earth revolved in an oblique circle, and at the same time turned round its own axis.

Aristarchus invented a peculiar kind of sun-dials, mentioned by Vitruvius. There is extant of his works only a treatise upon the magnitude and distance of the sun and moon: this was translated into Latin, and commented upon by Commandine, who first published it with Pappus's Explanations, in 1572. Dr. Wallis afterwards published it in Greek, with Commandine's Latin version, in 1688, and which he inserted again in the 3d volume of his Mathematical Works, printed in folio at Oxford, 1699. The piece was animadverted upon by Mr. Foster, in his Miscellanies. There is another piece which has gone under the name of Aristarchus, of the *Mundane System*, its parts, and motions, published in Latin by Robervale, and by Merfenne, in his *Mathematical Synopsis*. But this piece is censured, by Menagius (in Diog. Laert.), and Descartes, in his *Epistles*, as a fictitious piece of Robervale's, and not the genuine work of Aristarchus.

ARISTOTELIAN, something that relates to the philosopher *Aristotle*. Thus we say, an *Aristotelian* dogma, the *Aristotelian* school, &c.

ARISTOTELIAN *Philosophy*, the philosophy taught by *Aristotle*, and maintained by his followers. It is otherwise called the *peripatetic* philosophy, from their practice of teaching while they were walking.—The principles of Aristotle's philosophy, the learned agree, are chiefly laid down in the four books *de Cælo*. Instead of the more ancient systems, he introduced *matter*, *form*, and *privation*, as the principles of all things; but it does not seem that he derived much benefit from them in natural philosophy. And his doctrines are, for the most part, so obscurely expressed, that it has not yet been satisfactorily ascertained, what were his sentiments on some of the most important subjects. He attempted to refute the Pythagorean doctrine concerning the two-fold motion of the earth; and pretended to demonstrate, that the matter of the heavens is ungenerated, incorruptible, and not subject to any alteration: and he supposed that the stars were carried round the earth in solid orbs.

ARISTOTELIANS, a sect of philosophers, so called, from their leader *Aristotle*, and are otherwise called *Peripatetics*.—The Aristotelians and their dogmata prevailed for a long while, in the schools and universities; even in spite of all the efforts of the Cartesian, Newtonians, and other corpuscularians. But the systems of the latter have at length gained the ascendancy; and the Newtonian philosophy in particular is now very generally received.

ARISTOTLE, a Grecian philosopher, the son of Nicomachus, physician to Amyntas king of Macedonia, was born 384 years before Christ, at Stagira, a town of Macedonia, or, as others say, of Thrace; from which he is also called the Stagirite. Not succeeding in the profession of arms, to which it seems he first applied himself, he turned his views to philosophy, and at 17 years of age entered himself a disciple of Plato, and attended in the academy till the death of that philosopher. Aristotle then retired to Atarna, where the prince Hermias gave him his daughter to wife. Repairing afterwards to the court of king Philip, he became preceptor to his son, Alexander the Great, whose education he attended for the space of 8 years; and by the magnificent encouragement of this prince he was afterwards enabled to procure all sorts of animals, from the inspection of which to write the history of them. On his quitting Macedon, he settled at Athens, where he established his school, having the Lyceum assigned him, by the magistrates, for the place of his instruction or disputation; where he became the head and founder of the sect called after his name, as also Peripatetics, from the circumstance of his giving instructions while walking. But being here accused of impiety by Eurymedon, priest of Ceres, and fearing the fate of Socrates, he retired to Chalcis, where he died at 63 years of age, and 322 years before Christ. Some say that he poisoned himself, others that he died of a cholic, and others again pretend that he threw himself into the sea for grief that he could not discover the cause of the flux and reflux of the waters. Laertius, in his life of Aristotle, estimates his books at the number of 4000; of which however scarce 20 have come down to us: these may be comprised under five heads; the first, relating to poetry and rhetoric; the second, to logics; the third, to ethics and politics; the fourth, to physics; and the fifth, to metaphysics. In the schools, Aristotle has been called the philosopher, and the prince of philosophers. And such was the veneration paid to him, that his opinion was allowed to stand on a level with reason itself: nor was any appeal from it admitted, the parties, in every dispute, being obliged to shew, that their conclusions were no less conformable to the doctrine of Aristotle than to truth.

ARITHMETIC, the art and science of numbers; or, that part of mathematics which considers their powers and properties, and teaches how to compute or calculate truly, and with ease and expedition. It is by some authors also defined the science of discrete quantity. Arithmetic consists chiefly in the four principal rules or operations of Addition, Subtraction, Multiplication, and Division; to which may perhaps be added involution and evolution, or raising of powers and extraction of roots. But besides these, for the facilitating and expediting of computations, mercantile, astronomical, &c, many other useful rules have been contrived, which are applications of the former, such as, the rules of proportion, progression, alligation, false position, fellowship, interest, barter, rebate, equation of payments, reduction, tare and tret, &c. Besides the doctrine of the curious and abstract properties of numbers.

Very little is known of the origin and invention of arithmetic. In fact it must have commenced with mankind, or as soon as they began to hold any sort of commerce



merce together; and must have undergone continual improvements, as occasion was given by the extension of commerce, and by the discovery and cultivation of other sciences. It is therefore very probable that the art has been greatly indebted to the Phœnicians or Tyrians; and indeed Proclus, in his commentary on the first book of Euclid, says, that the Phœnicians, by reason of their traffic and commerce, were accounted the first inventors of Arithmetic. From Asia the art passed into Egypt, whither it was carried by Abraham, according to the opinion of Josephus. Here it was greatly cultivated and improved; inasmuch that a considerable part of the Egyptian philosophy and theology seems to have turned altogether upon numbers. Hence those wonders related by them about unity, trinity, with the numbers 4, 7, 9, &c. In effect, Kircher, in his *Oedip. Ægypt.* shews, that the Egyptians explained every thing by numbers; Pythagoras himself affirming, that the nature of numbers pervades the whole universe; and that the knowledge of numbers is the knowledge of the deity.

From Egypt arithmetic was transmitted to the Greeks, by means of Pythagoras and other travellers; amongst whom it was greatly cultivated and improved, as appears by the writings of Euclid, Archimedes, and others: with these improvements it passed to the Romans, and from them it has descended to us.

The nature of the arithmetic however that is now in use, is very different from that above alluded to; this art having undergone a total alteration by the introduction of the Arabic notation, about 800 years since, into Europe: so that nothing now remains of use from the Greeks, but the theory and abstract properties of numbers, which have no dependence on the peculiar nature of any particular scale or mode of notation. That used by the Hebrews, Greeks, and Romans, was chiefly by means of the letters of their alphabets. The Greeks, particularly, had two different methods; the first of these was much the same with the Roman notation, which is sufficiently well known, being still in common use with us, to denote dates, chapters and sections of books, &c. Afterwards they had a better method, in which the first nine letters of their alphabet represented the first numbers, from one to nine, and the next nine letters represented any number of tens, from one to nine, that is, 10, 20, 30, &c, to 90. Any number of hundreds they expressed by other letters, supplying what they wanted with some other marks or characters: and in this order they went on, using the same letters again, with some different marks, to express thousands, tens of thousands, hundreds of thousands, &c: In which it is evident that they approached very near to the more perfect decuple scale of progression used by the Arabians, and who acknowledge that they had received it from the Indians. Archimedes also invented another peculiar scale and notation of his own, which he employed in his *Arenarius*, to compute the number of the sands. In the 2d century of christianity lived Cl. Ptolemy, who, it is supposed, invented the sexagesimal division of numbers, with its peculiar notation and operations: a mode of computation still used in astronomy &c, for the subdivisions of the degrees of circles. Those notations however were ill adapted to the practical operations of arithmetic: and hence it is that the art ad-

vanced but very little in this part; for, setting aside Euclid, who has given many plain and useful properties of numbers in his *Elements*, and Archimedes, in his *Arenarius*, they mostly consist in dry and tedious distinctions and divisions of numbers; as appears from the treatises of Nicomachus, supposed to be written in the 3d century of Rome, and published at Paris in 1538; as also that of Boethius, written at Rome in the 6th century of Christ. A compendium of the ancient arithmetic, written in Greek, by Psellus, in the 9th century, was published in Latin by Xylander, in 1556. A similar work was written soon after in Greek by Jodocus Willichius; and a more ample work of the same kind was written by Jordanus, in the year 1200, and published with a comment by Faber Stapulensis in 1480.

Since the introduction of the Indian notation into Europe, about the 10th century, arithmetic has greatly changed its form, the whole algorithm, or practical operations with numbers, being quite altered, as the notation required; and the authors of arithmetic have gradually become more and more numerous. This method was brought into Spain by the Moors or Saracens; whither the learned men from all parts of Europe repaired, to learn the arts and sciences of them. This, Dr. Wallis proves, began about the year 1000; particularly that a monk, called Gilbert, afterwards pope, by the name of Sylvester II, who died in the year 1003, brought this art from Spain into France, long before the date of his death: and that it was known in Britain before the year 1150, where it was brought into common use before 1250, as appears by the treatise of arithmetic of Johannes de Sacro Bosco, or Halifax, who died about 1256. Since that time, the principal writers on this art have been, Barlaam, Lucas de Burgo, Tostall, Aventinus, Purbach, Cardan, Scheubelius, Tartalia, Faber, Stifelius, Recorde, Ramus, Maurolycus, Hemischi, Peletarius, Stevinus, Xylander, Kersey, Snellius, Tacquet, Clavius, Metius, Gemma Frisius, Buteo, Ursinus, Romanus, Napier, Ceulen, Wingate, Kepler, Briggs, Ulacq, Oughtred, Cruger, Van Schooten, Wallis, Dee, Newton, Morland, Moore, Jeake, Ward, Hatton, Malcolm, &c, &c; the particular inventions or excellencies of whom, will be noticed under the articles of the several species or kinds of arithmetic here following, which may be included under these heads, viz, *theoretical, practical, instrumental, logarithmical, numerous, specious, universal, common or decadal, fractional, radical or of surds, decimal, duodecimal, sexagesimal, dynamical or binary, tetractylal, political, &c.*

*Theoretical* ARITHMETIC, is the science of the properties, relations, &c, of numbers, abstractedly considered; with the reasons and demonstrations of the several rules. Such is that contained in the 7th, 8th, and 9th books of Euclid's *Elements*; the *Logistics* of Barlaam the monk, published in Latin by J. Chambers, in 1600; the *Summa Arithmetica* of Lucas de Burgo, printed 1494, who gives the several divisions of numbers from Nicomachus, and their properties from Euclid, with the algorithm, both in integers, fractions, extraction of roots, &c; Malcolm's *New System of Arithmetic, theoretical and practical*, in 1730, in which the subject is very completely treated, in all its branches, &c.

*Practical*



*Practical ARITHMETIC*, is the art or practice of numbering or computing; that is, from certain numbers given, to find others which shall have any proposed relation to the former. As, having the two numbers 4 and 6 given; to find their sum, which is 10; or their difference, which is 2; or their product, 24; or their quotient,  $1\frac{1}{2}$ ; or a third proportional to them, which is 9; &c.—Lucas de Burgo's works contain the whole practice of arithmetic, then used, as well as the theory. Tunstall gave a neat practical treatise of Arithmetic in 1526; as did Stifelius, in 1544, both on the practical and other parts. Tartalea gave an entire body of practical arithmetic, which was printed at Venice in 1556, consisting of two parts; the former, the application of arithmetic to civil uses; the latter, the grounds of Algebra. And most of the authors in the list before enumerated, joined the practice of arithmetic with the theory.

*Binary or Dyadic ARITHMETIC*, is that in which only two figures are used, viz 1 and 0. See BINARY. —Leibnitz and De Lagny both invented an arithmetic of this sort, about the same time: and Dancicourt, in the Miscel. Berol. gives a specimen of the use of it in arithmetical progressions; where he shews, that the laws of progression may be more easily discovered by it, than by any other method where more characters are used.

*Common or Vulgar ARITHMETIC*, is that which is concerning integers and vulgar fractions.

*Decimal or Decadal ARITHMETIC*, is that which is performed by a series of ten characters or figures, the progression being ten-fold, or from 1 to 10's, 100's, &c; which includes both integers and decimal fractions, in the common scale of numbers; and the characters used are the ten Arabic or Indian figures 0, 1, 2, 3, 4, 5, 6, 7, 8, 9. This method of arithmetic was not known to the Greeks and Romans; but was borrowed from the Moors while they possessed a great part of Spain, and who acknowledge that it came to them from the Indians. It is probable that this method took its origin from the ten fingers of the hands, which were used in computations before arithmetic was brought into an art. The Eastern missionaries assure us, that to this day the Indians are very expert at computing on their fingers, without any use of pen and ink. And it is asserted, that the Peruvians, who perform all computations by the different arrangements of grains of maize outdo any European, both for certainty and dispatch, with all his rules.

*Duodecimal ARITHMETIC*, is that which proceeds from 12 to 12, or by a continual subdivision according to 12. This is greatly used by most artificers, in calculating the quantity of their work; as Bricklayers, Carpenters, Painters, Tilers, &c.

*Fractional ARITHMETIC*, or of fractions, is that which treats of fractions, both vulgar and decimal.

*Harmonical ARITHMETIC*, is so much of the doctrine of numbers, as relates to the making the comparisons, reductions, &c of musical intervals.

*ARITHMETIC of Infinites*, is the method of summing up a series of numbers, of which the numbers of terms is infinite. This method was first invented by Dr. Wallis, as appears by his treatise on that subject; where he shews its uses in geometry, in finding the areas of

superficies, the contents of solids, &c. But the method of fluxions, which is a kind of universal arithmetic of infinites, performs all these more easily; as well as a great many other things, which the former will not reach.

*Instrumental ARITHMETIC*, is that in which the common rules are performed by instruments, or some sort of tangible or palpable substance. Such are the methods of computing by the ten fingers and the grains of maize, by the East Indians and Peruvians, above-mentioned; by the Abacus or Shwanpan of the Chinese; the several sorts of scales and sliding rules; Napier's bones or rods; the arithmetical machine of Pascal, and others; Sir Samuel Morland's instrument, described in 1666; that of Leibnitz, described in the Miscell. Berol.; that of Polenus, published in the Venetian Miscellany, 1709; and that of Dr. Saunderson, of Cambridge, described in the introduction to his algebra.

*Integral ARITHMETIC*, or of integers, is that which respects integers, or whole numbers.

*Literal or Algebra ARITHMETIC*, is that which is performed by letters, which represent any numbers indefinitely.

*Logarithmical ARITHMETIC* is performed by the tables of logarithms. These were invented by baron Napier; and the best treatise on the subject, is Briggs's *Arithmetica Logarithmica*, 1624.

*Logistical ARITHMETIC*. See LOGISTICAL.

*Numerous or Numeral ARITHMETIC*, is that which teaches the calculus of numbers, or of abstract quantities; and is performed by the common numeral or Arabic characters.

*Political ARITHMETIC*, is the application of arithmetic to political subjects; such as, the strength and revenues of nations, the number of people, births, burials, &c. See POLITICAL Arithmetic. To this head may also be referred the doctrine of Chances, Gaming, &c.

*ARITHMETIC of Radicals, Rationals, and Irrationals*. See RADICAL, &c.

*Sexagesimal or Sexagenary ARITHMETIC*, is that which proceeds by sixties; or the doctrine of sexagesimal fractions: a method which, it is supposed, was invented by Ptolemy, in the 2d century; at least they were used by him. In this notation, the integral numbers from 1 to 59 were expressed in the common way, by the alphabetical letters: then sixty was called a *sexagena prima*, and marked with a dash to the character, thus I'; twice sixty, or 120, thus II'; and so on to 59 times 60, or 3540, which is LIX'. Again, 60 times 60, or 3600, was called *sexagena secunda*, and marked with two dashes, thus I''; twice 3600, thus II''; and ten times 3600, thus X''; &c. And in this way the notation was continued to any length. But when a number less than sixty was to be joined with any of the sexagesimal integers, their proper expression was annexed without the dash: thus 4 times 60 and 25, is IV'XXV; the sum of twice 60 and 10 times 3600 and 15, is X''II'XV. So near did the inventor of this method approach to the Arabic notation: instead of the sexagesimal progression, he had only to substitute decimal; and to make the signs of numbers, from 1 to 9, simple characters, and to introduce another character, which should signify nothing by itself, but serving only to fill up places. —The *sexagena integrorum* were soon laid aside, in ordinary



dinary calculations, after the introduction of the Arabic notation; but the sexagesimal fractions continued till the invention of decimals, and indeed are still used in the subdivisions of the degrees of circular arcs and angles.

Sam. Reyher has invented a kind of sexagenal rods, in imitation of Napier's bones, by means of which the sexagesimal arithmetic is easily performed.

*Specious ARITHMETIC*, is that which gives the calculus of quantities as designed by the letters of the alphabet: a method which was more generally introduced into algebra, by Vieta; being the same as literal arithmetic, or algebra.—Dr. Wallis has joined the numeral with the literal calculus; by which means he has demonstrated the rules for fractions, proportions, extraction of roots, &c; of which a compendium is given by himself, under the title of *Elementa Arithmetice*, in the year 1698.

*Tabular ARITHMETIC*, is that in which the operations of multiplication, division, &c, are performed by means of tables calculated for that purpose: such as those of Herwart, in 1610; and my tables of powers and products, published by order of the Commissioners of Longitude, in 1781.

*Tetraëtic ARITHMETIC*, is that in which only the four characters 0, 1, 2, 3 are used. A treatise of this kind of arithmetic is extant, by Erhard or Echard Weigel. But both this, and binary arithmetic, are little better than curiosities, especially with regard to practice; as all numbers are much more compendiously and conveniently expressed by the common decuple scale.

*Vulgar, or Common ARITHMETIC*, is that which is conversant about integers and vulgar fractions.

*Universal ARITHMETIC*, is the name given by Newton to the science of algebra; of which he left at Cambridge an excellent treatise, being the text-book drawn up for the use of his lectures, while he was professor of Mathematics in that University.

*ARITHMETICAL*, something relating to or after the manner of arithmetic.

*ARITHMETICAL Complement*, of a logarithm, is what the logarithm wants of 10.00000 &c; and the easiest way to find it is, beginning at the left hand, to subtract every figure from 9, and the last from 10. So, the arithmetical complement of 8.2501396 is 1.7498604.—It is commonly used in trigonometrical calculations, when the first term of a proportion is not radius; in that case, adding all together, the logarithms of the 3d, 2d, and arithmetical complement of the 1st term.

*ARITHMETICAL Instruments, or Machines*, are instruments for performing arithmetical computations; such as Napier's bones, scales, sliding rules, Pascal's machine, &c.

*ARITHMETICAL Mean, or Medium*, is the middle term of three quantities in arithmetical progression; and is always equal to half the sum of the extremes. So, an arithmetical mean between 3 and 7, is 5; and between  $a$  and  $b$ , is  $\frac{a+b}{2}$ , or  $\frac{1}{2}a + \frac{1}{2}b$ .

*ARITHMETICAL Progression*, is a series of three or more quantities that have all the same common difference: as 3, 5, 7, &c, which have the common difference 2; and  $a, a+d, a+2d, \&c$ , which have all the same difference  $d$ .

In an arithmetical progression, the chief properties are these: 1st, The sum of any two terms, is equal to the sum of every other two that are taken at equal distances from the two former, and equal to double the middle term when there is one equally distant between those two: so, in the series 0, 1, 2, 3, 4, 5, 6, &c,  $0+6=1+5=2+4=$  twice 3 or 6.—2d, The sum of all the terms of any arithmetical progression, is equal to the sum of as many terms of which each is the arithmetical mean between the extremes; or equal to half the sum of the extremes multiplied by the number of terms: so, the sum of these ten terms 0, 1, 2, 3, 4,

5, 6, 7, 8, 9, is  $\frac{0+9}{2} \times 10$ , or  $9 \times 5$ , which is 45:

and the reason of this will appear by inverting the terms, setting them under the former terms, and adding each two together, which will make double the same series;

thus      0, 1, 2, 3, 4, 5, 6, 7, 8, 9,  
inverted 9, 8, 7, 6, 5, 4, 3, 2, 1, 0,  
sums    9, 9, 9, 9, 9, 9, 9, 9, 9, 9;

where the double series being the same number of 9's, or sum of the extremes, the single series must be the half of that sum.—3d, The last, or any term, of such a series, is equal to the first term, with the product added of the common difference multiplied by 1 less than the number of terms, when the series ascends or increases; or the same product subtracted when the series descends or decreases: so, of the series 1, 2, 3, 4, &c, whose common difference is 1, the 50th term is  $1+1 \times 49$ , or  $1+49$ , that is 50; and of the series 50, 49, 48, &c, the 50th term is  $50-1 \times 49$ , or  $50-49$ , which is 1. Also, if  $a$  denote the least term,

$z$  the greatest term,

$d$  the common difference,

$n$  the number of the terms,

and  $s$  the sum of them all;

then the principal properties are expressed by these equations, viz,

$$z = a + d \cdot n - 1,$$

$$a = z - d \cdot n + 1,$$

$$s = a + z \cdot \frac{1}{2}n,$$

$$s = z - \frac{1}{2}d \cdot n + 1 \cdot n,$$

$$s = a + \frac{1}{2}d \cdot n - 1 \cdot n.$$

Moreover, when the first term  $a$ , is 0 or nothing; the theorems become  $z = d \cdot n - 1$ .

$$\text{and } s = \frac{1}{2}zn.$$

*ARITHMETICAL Proportion*, is when the difference between two terms, is equal to the difference between other two terms. So, the four terms, 2, 4, 10, 12, are in arithmetical proportion, because the difference between 2 and 4, which is 2, is equal to the difference between 10 and 12.—The principal property, besides the above, and which indeed depends upon it, is this, that the sum of the first and last, is equal to the sum of the two means: so  $2+12$ , or the sum of 2 and 12, is equal  $4+10$ , or the sum of 4 and 10, which is 14.



**ARITHMETICAL Ratio**, is the same as the difference of any two terms: so, the arithmetical ratio of the series 2, 4, 6, 8, is 2; and the arithmetical ratio of  $a$  and  $b$ , is  $a - b$ .

**ARITHMETICAL Scales**, a name given by M. de Buffon, in the *Memoirs of the Acad.* for 1741, to different progressions of numbers, according to which, arithmetical computations might be made. It has already been remarked above, that our common decuple scale of numbers was probably derived from the number of fingers on the two hands, by means of which the earliest and most natural mode of computation was performed; and that other scales of numbers, formed in a similar way, but of a different number of characters, have been devised; such as the binary and tetractic scales of arithmetic. In the memoir above cited, Buffon gives a short and simple method to find, at once, the manner of writing down a number given in any scale of numbers whatever; with remarks on different scales. The general effect of any number of characters, different from ten, is, that by a smaller number of characters, any given number would require more places of figures to express or denote it by, but then arithmetical calculations, by multiplication and division, would be easier, as the small numbers 2, 3, 4, &c, are easier to use than the larger 7, 8, 9; and by employing more than ten characters, although any given number would be expressed by fewer of them, yet the calculations in arithmetic would be more difficult, as by the larger numbers 11, 12, 13, &c. It is therefore concluded, upon the whole, that the ordinary decuple scale is a good convenient medium amongst them all, the numbers expressed being tolerably short and compendious, and no single character representing too large a number. The same might also be said, and perhaps more, of a duodecimal scale, by twelve characters, which would express all numbers in a more compendious way than the decuple one, and yet no single character would represent a number too large to compute by; as is confirmed by the now common practice of extending the multiplication table, in school books, to 12 numbers or dimensions, each way, instead of 10; and every person is taught, with sufficient ease, to multiply and divide by 11 and 12 as easily as by 8 or 9 or 10. Another convenience might be added, namely, that the number 12 admitting of more submultiples than the number 10, there would be fewer expressions of interminate fractions in that way than in decimals. So that on all accounts, it is very probable that the duodecimal would be the best of any scale of numbers whatever.

**ARITHMETICAL Triangle.** See *Arithmetical TRIANGLE*.

**ARMED.** A magnet or loadstone is said to be armed, when it is capped, cased, or set in iron or steel; to make it take up a greater weight; and also readily to distinguish its poles.

It is surprising, that a little iron fastened to the poles of a magnet, should so greatly improve its power, as to make it even 150 times stronger, or more, than it is naturally, or when unarmed. The effect however, it seems, is not uniform; but that some magnets, by arming, gain much more, and others much less, than one would expect; and that some magnets even lose some of their efficacy by arming. In general, however,

the thickness of the iron armour ought to be nearly proportioned to the natural strength of the magnet; giving thick irons to a strong magnet, and to the weaker ones thinner: so that a magnet may easily be over-loaded.

The usual armour of a load-stone, in form of a right-angled parallelopipedon, consists of two thin pieces of iron or steel, of a square figure, and of a thickness proportioned to the goodness of the stone; the proper thickness being found by trials; always filing it thinner and thinner, till the effect be found to be the greatest possible.—The armour of a spherical load-stone, consists of two steel shells, fastened together by a joint, and covering a good part of the convexity of the stone. This also is to be filed away, till the effect is found to be the greatest.

Kircher, in his book *de Magnete*, says, that the best way to arm a load-stone, is to drill a hole through the stone, from pole to pole, in which is to be placed a steel rod of a moderate length: this rod, he asserts, will take up more weight at the end, than the stone itself when armed in the common way. And Gassendus and Cabæus prescribe the same method of arming. But Muschenbroëk found, by repeated trials, that the usual armour already mentioned, is preferable to Kircher's; and he gives the following directions for preparing it. When, by means of steel filings and a small needle, the poles of a magnet have been discovered, he directs that the adjacent parts should be rubbed or ground into parallel planes, without shortening the polar axis; and the magnet may be afterwards shaped into the figure of a cube or parallelopipedon, or any other figure that may be more convenient. Plates of the softest iron are then prepared, of the same length and breadth with the whole polar sides of the magnet: the thickness of which plates, so as that they may admit and convey the greatest quantity of the magnetic virtue, is to be previously determined by experiment, in a manner which he prescribes for the purpose. A thicker piece of iron is to be annexed at right angles to these plates, which is called *pes armature*, the foot or base of the armour: then the plates, nicely smoothed and polished, are to be firmly attached to each of the polar sides, whilst the thicker part or base is brought into close contact with the lower part of the magnet. In this way, he says, almost all the magnetic virtue issuing from the poles, enters into the armour, is directed to the base, and condensed by means of its roundness, so as to sustain the greatest weight of iron. *Phys. Exper. and Geom. Dissert.* 1729, pa. 131.

**ARMILLARY Sphere**, a name given to the artificial sphere, composed of a number of circles of metal, wood, or paper, which represent the several circles of the system of the world, put together in their natural order. It serves to assist the imagination to conceive the disposition of the heavens, and the motion of the celestial bodies.

This sphere is represented at Plate II, Fig. 6, where P and Q represent the poles of the world, AD the equator, EL the ecliptic and zodiac, PAGD the meridian, or the solstitial colure, T the earth, FG the tropic of cancer, HT the tropic of capricorn, MN the arctic circle, OV the antarctic, N and O the poles of the ecliptic, and RS the horizon.

The Armillary sphere constructed not long since by Dr. Long, in Pembroke-hall, Cambridge, is 18 feet in diameter;



diameter; and will contain more than 30 persons sitting within it, to view, as from a centre, the representation of the celestial spheres. The lower part of the sphere, which is not visible to England, is cut off; and the whole apparatus is so contrived, that it may be turned round with as little labour as is employed to wind up a common jack.

See also Mr. Ferguson's sphere in his Lectures, p. 194.

*ARMILLARY Trigonometer*, an instrument first contrived by Mr. Mungo Murray, and improved by Mr. Ferguson, consisting of five semicircles; viz, meridian, vertical circle, horizon, hour circle, and equator; so adapted to each other by joints and hinges, and so divided and graduated, as to serve for expeditiously resolving many problems in astronomy, dialling, and spherical trigonometry. The drawing, description, and method of using it, may be seen in Ferguson's Tracts, pa. 80, &c.

*ARTIFICIAL Numbers, Sines, Tangents, &c.*, are the same as the Logarithms of the natural numbers, sines, tangents, &c.

*ARTILLERY*, the heavy equipage of war; comprehending all sorts of large fire-arms, with their appurtenances; as cannon, mortars, howitzers, balls, shells, petards, musquets, carbines, &c; being what is otherwise called *Ordnance*. The term is also applied to the larger instruments of war used by the ancients, as the catapult, balista, battering ram, &c.

The term *Artillery*, or *Royal Artillery*, is also applied to the persons employed in that service; and likewise to the art or science itself; and formerly it was used for what is otherwise called *pyrotechnia*, or the art of fire-works, with the apparatus and instruments belonging to the same.

There have been many authors on the subject of artillery; the principal of which are, Bucherius, Braunius, Tartalea, Collado, Sardi, Ufano, Hanzelet, Digges, Moretti, Simienowitz, Mieth, d'Avelour, Manesson, Mallet, St. Julien; and the later authors, of still more consequence, are Belidor, St. Remy, le Blond, Valiere, Morogue, Puget, Coudray, Robins, Muller, Euler, Antoni, Tignola, Scheele; to which may be added the extensive and accurate experiments published in my 1st vol. of Tracts, and in the Philos. Trans. for 1778.

*Park of ARTILLERY*, is that place in a camp which is set apart for the Artillery, or large fire arms.

*Traile or Train of ARTILLERY*, a number of pieces of ordnance, mounted on carriages, with all their furniture fit for marching. To this commonly belong mortars, cannon, balls, shells, &c.—There are trains of Artillery in most of the royal magazines; as in the Tower, at Portsmouth, Plymouth, &c, but, above all, at Woolwich, from whence the ships commonly receive their ordnance, and where they are all completely proved before they are received into the public service.

The officers and men of the artillery were formerly called also the Train of Artillery, but are now called the Royal Regiment of Artillery; consisting at present of four battalions, besides a battalion of invalids, and four troops of Horse or Cavalry Artillery.

*ASCENDANT*, in *Astrology*, denotes the horoscope; or the degree of the ecliptic which rises upon the horizon, at the time of the birth of any one.—This,

it is supposed, has an influence on the person's life and fortune, by giving him a bent and propensity to one thing more than other.—In the science of *Astrology*, this is called the first house, the Oriental angle, or angle of the East, and the significator of life: and the astrologer says, such a planet ruled in his Ascendant, or Jupiter was in his Ascendant, &c.

*ASCENDING*, in Astronomy, a term used to denote any star, or degree, or other point of the heavens, rising above the horizon.

*ASCENDING Latitude*, is the latitude of a planet when going towards the north.

*ASCENDING Node*, is that point of a planet's orbit where it crosses the ecliptic, in proceeding northward. It is otherwise called the Northern Node, and is denoted by this character ♊, representing a node, or knot, with the larger part upwards; like as the same character reversed is used to denote the opposite, or Descending Node.

*ASCENDING Signs*, are such as are upon their ascent, or rise, from the nadir or lowest point of the heavens, towards the zenith, or highest point.

*ASCENSION-DAY*, otherwise called *Holy Thursday*, is a festival of the church, held 10 days before Whitsunday, in memory of our Saviour's Ascension.

*ASCENSION*, in Astronomy, is either Right or Oblique.

*Right ASCENSION* of the sun, or of a star, is that degree of the equinoctial, accounted from the beginning of Aries, which rises with them, in a right sphere.—Or, *Right Ascension*, is that point of the equinoctial, counted as before, which comes to the meridian with the sun or star, or other point of the heavens. And the reason of thus referring it to the meridian, is, because this is always at right angles to the equinoctial; whereas the horizon is so only in a right or direct sphere.—The right ascension, stands opposed to the right descension; and is similar to the longitude of places on the earth. All the fixed stars, &c, which have the same right-ascension, that is, which are at the same distance from the first point of Aries, or, which comes to the same thing, which are in the same meridian, rise at the same time in a right sphere, namely to the people who live at the equator. And if they be not in the same meridian, the difference between their times of rising, or of coming to the meridian of any place, is the precise difference of their right ascension.—But, in an oblique sphere, where the horizon cuts all the meridians obliquely, different points of the same meridian never rise or set together: so that two or more stars on the same meridian, or having the same right ascension, never rise or set at the same time in an oblique sphere; and the more oblique the sphere is, the greater is the interval of time between them.

To find the right ascension of the sun, stars, &c, by Trigonometry, say, As radius is to the cosine of the sun's greatest declination, or obliquity of the ecliptic, so is the tangent of the sun's or star's longitude, to the tangent of the right ascension.

*RIGHT ASCENSION of the Mid-heaven*, often used by astronomers, especially in calculating eclipses by means of the nonagesimal degree, is the right ascension of that point of the equator which is in the meridian; and it is equal to the sum of the sun's right ascension and



the horary angle or true time reduced to degrees, or to the sum of the mean longitude and mean time.

*Oblique ASCENSION*, is an arch of the equator intercepted between the first point of Aries, and that point of the equator which rises together with the star, &c, in an oblique sphere.—The Oblique Ascension is counted from west to east; and is greater or less, according to the various obliquity of the sphere.—To find the Oblique Ascension of the sun, see *ASCENSIONAL* and *GLOBE*.

The *Arch of Oblique Ascension*, is an arch of the horizon intercepted between the beginning of Aries, and the point of the equator which rises with a star or planet, in an oblique sphere; and it varies with the latitude of the place.

*Refraction of ASCENSION and Descension*. See *REFRACTION*.

*ASCENSIONAL DIFFERENCE*, is the difference between the right and oblique ascension of the same point on the surface of the sphere. Or it is the time the sun rises or sets before or after 6 o'clock.

To find the Ascensional Difference, having given the sun's declination and the latitude of the place, say, As radius is to the tangent of the latitude, so is the tangent of the sun's declination to the sine of the Ascensional Difference sought. The sun's Ascensional Difference converted into time, shews how much he rises before, or sets after, 6 o'clock. When the sun has north declination, the right ascension is greater than the oblique; but the contrary, when the sun has south declination; and the difference, in either case, is the Ascensional Difference.

*ASCENT*, the motion of a body from below tending upwards; or the continual recess of a body from the earth, or from some other centre of force. And it is opposed to *descent*, or motion downwards.

The Peripatetics attributed the spontaneous ascent of bodies to a principle of levity, inherent in them. But the moderns deny that there is any such thing as spontaneous levity; and they shew, that whatever ascends, does so by virtue of some external impulse or extrusion. Thus it is that smoke, and other rare bodies, ascend in the atmosphere; and oil, light woods, &c, in water: not by any inherent principle of levity; but by the superior gravity, or tendency downwards of the medium in which they ascend and float.

The ascent of light bodies in heavy mediums, is produced after the same manner as the ascent of the lighter scale of a balance. It is not that such scale has an internal principle, by which it immediately tends upwards; but it is impelled upwards by the preponderancy of the other scale; the excess of the weight in the one having the same effect, by augmenting its impetus downwards, as so much real levity in the other: because the tendencies mutually oppose each other, and action and reaction are always equal.—See this farther illustrated under the articles *FLUID*, and *SPECIFIC GRAVITY*.

*ASCENT of Bodies on Inclined Planes*. See the doctrine and laws of them under *INCLINED PLANE*.

*ASCENT of Fluids*, is particularly understood of their rising above their own level, between the surfaces of nearly contiguous bodies, or in slender capillary glass tubes, or in vessels filled with sand, ashes, or the like porous substances. Which is an effect that takes place as well *in vacuo*, as in the open air, and in crooked,

as well as straight tubes. Indeed some fluids ascend swifter than others, as spirit of wine, and oil of turpentine; and some rise after a different manner from others. The phenomenon, with its causes, &c, in the instance of capillary tubes, will be treated more at large under *CAPILLARY Tube*.

As to planes: Two smooth polished plates of glass, metal, stone, or other matter, being placed almost contiguous, have the effect of several capillary tubes, and the fluid rises in them accordingly: the like may be said of a vessel filled with sand, &c; the various small interstices of which form, as it were, a kind of capillary tubes. So that the same principle accounts for the appearance in them all. And to the same cause may probably be ascribed the ascent of the sap in vegetables. And on this subject Sir I. Newton says, "If a large pipe of glass be filled with sifted ashes, well pressed together, and one end dipped into stagnant water, the fluid will ascend slowly in the ashes, so as in the space of a week or fortnight, to reach the height of 30 or 40 inches above the stagnant water. This ascent is wholly owing to the action of those particles of the ashes which are upon the surface of the elevated water; those within the water attracting as much downwards as upwards: it follows, that the action of such particles is very strong; though being less dense and close than those of glass, their action is not equal to that of glass, which keeps quicksilver suspended to the height of 60 or 70 inches, and therefore acts with a force which would keep water suspended to the height of above 60 feet. By the same principle, a sponge sucks in water, and the glands in the bodies of animals, according to their several natures and dispositions, imbibe various juices from the blood." *Optics*, pa. 367.

Again, if a drop of water, oil, or other fluid, be dropped upon a glass plane, perpendicular to the horizon, so as to stand without breaking, or running off; and another plane touching it at one end, be gradually inclined towards the former, till it touch the drop; then will the drop break and move along towards the touching end of the planes; and it will move the faster in proportion as it proceeds farther, because the distance between the planes is constantly diminishing. And after the same manner, the drop may be brought to any part of the planes, either upward or downward, or sideways, by altering the angle of inclination.

Lastly, if the same perpendicular planes be so placed, as that two of their sides meet, and form a small angle, the other two being only kept apart by the interposition of some thin body; and thus immersed in a fluid, tinged with some colour to render it visible; the fluid will ascend between the planes, and that the highest where the planes are nearest; so as to form a curve line which is found to be a true hyperbola, of which one of the asymptotes is the line of the fluid, the other being a line drawn along the touching sides.

And the physical cause of all these phenomena, is the same power of attraction.

*ASCENT of Vapour*. See *CLOUD* and *VAPOUR*.

*ASCENT*, in Astronomy, &c. See *ASCENSION*.

*ASCII*, are those inhabitants of the globe, who, at certain times of the year, have no shadow. Such are the inhabitants of the torrid zone, who twice a year having the sun at noon in their zenith, have then no shadow.

ASELLI,



**ASELLI**, two fixed stars of the fourth magnitude, in the constellation Cancer.

**ASH-WEDNESDAY**, the first day of Lent, supposed to have been so called from a custom in the church, of sprinkling ashes that day on the heads of penitents then admitted to penance.

**ASPECT**, is the situation of the stars and planets in respect of each other. Or, in Astrology, it denotes a certain configuration and mutual relation between the planets, arising from their situations in the zodiac, by which it is supposed that their powers are mutually either increased or diminished, as they happen to agree or disagree in their active or passive qualities. Though such configurations may be varied and combined a thousand ways, yet only a few of them are considered. Hence, **Wolffius** more accurately defines aspect to be, the meeting of luminous rays emitted from two planets to the earth, either posited in the same right line, or including an angle which is an aliquot part, or some number of aliquot parts, of four right angles, or of 360 degrees.

The doctrine of aspects was introduced by the astrologers, as the foundation of their predictions. And hence **Kepler** defines aspect to be, an angle formed by the rays of two planets meeting on the earth, capable of exciting some natural power or influence.

The ancients reckoned five aspects, viz, *conjunction*, *sextile*, *quartile*, *trine*, and *opposition*.

*Conjunction* is denoted by this character  $\odot$ , and is when the planets are in the same sign and degree, or have the same longitude.

*Sextile* is denoted by  $\ast$ , and is when the planets are distant by the 6th part of a circle, or 2 signs, or 60 degrees.

*Quartile* is denoted by  $\square$ , and is when the planets are distant  $\frac{1}{4}$  of the circle, or 90 degrees, or 3 signs.

*Trine* is denoted by  $\triangle$ , and is when the planets are distant by  $\frac{1}{3}$  of the circle, or 4 signs, or 120 degrees. And

*Opposition* is denoted by  $\oslash$ , and is when the planets are in opposite points of the circle, or differ by  $\frac{1}{2}$  the circle, or 6 signs, or 180 degrees of longitude.

Or their characters and distances are as in this following tablet.

NAME.	CHARACTER.	DISTANCE.
Conjunction -	$\odot$	0°
Sextile - - -	$\ast$	60
Quartile - - -	$\square$	90
Trine - - - -	$\triangle$	120
Opposition -	$\oslash$	180

These intervals are reckoned according to the longitudes of the planets; so that the aspects are the same, whether the planet be in the ecliptic or out of it.

To the five ancient aspects, modern writers have added several more: as *decile*, for the 10th part of a circle; *tridecile*, or  $\frac{3}{10}$ ; and *biquintile*, or  $\frac{2}{5}$  of a circle. And **Kepler** adds others, from meteorological observations, as he tells us: as the *semisextile*, being  $\frac{1}{12}$ , or 30°; and *quincunx*, or  $\frac{1}{5}$ , or 150°. Lastly, to the astrological physicians, &c, we owe *octile*, or  $\frac{1}{8}$ ; *trioctile*, or  $\frac{3}{8}$ ; and *quintile*,  $\frac{1}{5}$  of the circle.

The aspects are divided with regard to their supposed influences, into *benign*, *malign*, and *indifferent*.

The trine and sextile aspects being esteemed benign or friendly; the quartile and opposition, malign or unfriendly; and the conjunction an indifferent aspect.

Aspects are also distinguished into *partile* and *platic*.

*Partile* ASPECT, is when the planets are just so many degrees distant, as are expressed above. And these only are the proper aspects.

*Platic* ASPECT, is when the planets do not regard each other exactly from these very degrees of distance; but the one exceeding as much as the other falls short. So that the one does not cast its rays immediately on the body of the other, but only on its orb or sphere of light.

**ASPERITY**, signifies the inequality or roughness of the surface of any body; by which some parts of it are more prominent than the rest, so as to hinder the hand, &c, from passing over it with ease and freedom; and thus producing what is called friction.—Asperity, or roughness, stands opposed to smoothness, evenness, politure, &c.

According to the relations of **Vermausen**, the blind man so celebrated for distinguishing colours by the touch, it seems that every colour has its peculiar degree and kind of asperity. He makes black the roughest, as it is the darkest of colours; but the others are not smoother in proportion as they are lighter; that is, the roughest do not always reflect the least light: for, according to him, yellow is two degrees rougher than blue, and as much smoother than green. See COLOURS.

**ASSIGNABLE** *Magnitude*, is used for any finite magnitude that can be expressed or denoted. And,

**ASSIGNABLE** *Ratio*, for any expressible ratio.

**ASSUMPTION**, a feast celebrated in the Romish church, in honour of the miraculous ascent of the Holy Virgin, as they describe it, body and soul, into heaven. It is kept on the 15th of August.

**ASSURANCE** *on Lives*, a compact by which security is granted for the payment of a certain sum of money on the expiration of the life on which the policy is granted, in consideration of such a previous payment made to the assurer as is accounted a sufficient compensation for the loss and hazard to which he exposes himself.

The sum at which this compensation should be valued, depends principally on these two circumstances, viz, 1st, On the rate of interest given for the use of money; and 2d, On the probability of the duration of the life assured, and the values of annuities. For, 1st, If the interest of money be high, the value of the assurance will be proportionally low, *Et è contra*; because the higher the rate of interest, the less will be the present value which amounts to a certain proposed sum in any given time. Also, if the probability of the duration of life be high, the value of the assurance will again be proportionally low, *Et è contra*; because the longer the time is, the less will be the principal which will amount to any assigned sum. Thus, if it be required to know the premium or present value, to be given for 100 pounds to be received at the end of any time, as suppose 10 years; then, if the interest of money be at the rate of 5 per cent. the answer, or present premium, would be 61l. 7s. 10d; but at four per cent. it would be 67l. 11s. 1d; and at 3 per cent. it would amount to 74l. 8s. 2d. Again, suppose it were required



required to assure 100l. on a life, for any time, for instance 1 year; that is, let 100l. be supposed to be payable a year hence, provided a life of a given age fails in that time: here it is evident that, whatever be the rate of interest, the less the probability of the life failing within the year, the less the risk is, and the less the premium ought to be. In effect, the rate of interest being 5 per cent, if it were sure that the life would fail in that year, the value of the assurance would be the same as the present value of 100l. payable at the end of the year, which is 95l. 4s. 9d. But, if it be an equal chance whether the life does or does not fail in the year, in which case the probability of failing is  $\frac{1}{2}$ ; then the value of the assurance will be but half the former value, or 47l. 12s. 4 $\frac{1}{2}$ d. Or if the odds against its failing be as 2 to 1, that is, if one person out of every 3 die at the age of the proposed life, the probability of dying being only  $\frac{1}{3}$ , the value of the assurance will be  $\frac{1}{3}$  of the first value, or 31l. 14s. 11d. And if the odds be 19 to 1, or one person die out of 20, of that age, the probability of dying will be  $\frac{1}{20}$ , and the value of the assurance will also be  $\frac{1}{20}$  of the first value, or 4l. 15s. 3d. nearly. Lastly, if only one person die out of 50 at the given age, the probability of dying will be  $\frac{1}{50}$ , and the value of the assurance will be accordingly only  $\frac{1}{50}$  of the first sum, or 1l. 18s. 1d: the interest of money being all along considered as after the rate of 5 per cent.—Now, according to Dr. Halley's table of observations, one person dies out of 3, at the age of 87; one in 20 at the age of 64; and one in 50 at the age of 39: It follows, therefore, that the value of the assurance of 100l. for one year, on a life aged 87, is 31l. 14s. 11d; on a life aged 64, it is 4l. 15s. 3d; and on a life aged 39, it is 1l. 18s. 1d: reckoning interest at 5 per cent. But if interest were rated at 3 per cent. these values would be 32l. 7s. 3d, and 4l. 17s. 1d, and 1l. 18s. 10d.

The assurances most commonly practised, are such as these, on single lives, and for single years. But many private assurers, and even some large assuring offices, either from ignorance or imposition, pay no regard to any difference of age, but demand 5l. from all ages indiscriminately, for the assurance of 100l. for one year: a practice very absurd and inequitable; for it appears that this is more than the value of the assurance of a life of 64 years of age, and even more than double the value of the assurance of a life of 39 years of age; allowing the assured to make 5 per cent. of the money he advances.

When a life is assured for any number of years; the premium or value may be paid, either in one single present payment; in consequence of which the sum assured will become payable without any farther compensation, whenever, within the given term, the life shall happen to drop: or the value may be paid in annual payments, to be continued till the failure of the life, should that happen within the term; or, if not, till the determination of the term. And the determination of the value of assurance, in all cases, is to be made out from the rules for computing annuities on lives; the principal writers on which are Halley, De Moivre, Simpson, Smart, Kerseboom, De Parcieux, Price, Morgan, and Maseres. See also LIFE ANNUITIES, REVERSION, &c.

Assurances may be made either on *single lives*; as above explained; or they may be made on any number of *joint lives*, or on the *longest* of any lives; that is, an assurer may bind himself to pay any sums at the extinction of any *joint lives*, or the *longest* of any lives, or at the extinction of any one or two of any number of lives. There are further assurances on survivorships; by which is meant an obligation, for the value received, to pay a given sum or annuity, provided a given life shall survive any other given life or lives. For which see SURVIVORSHIP.

The principal offices for making these insurances, in England, are the "London and the Royal Exchange Assurance Offices;" "the Amicable Society, incorporated for a perpetual Assurance Office;" "the Society for equitable Assurances on Lives and Survivorships;" and "the Westminster Society for granting Annuities and insuring Money on Lives."

The first two of these offices, having chiefly in view assurances on ships and houses, deal but little in the way of assurances on lives; and all the business they transact in this way, is at 5l. for every 100l. assured on a single life for a single year, without paying any regard to the ages of the lives assured.

The next, or Amicable Society at Serjeant's Inn, requires an annual payment of 5l. from every member during life, payable quarterly. The whole annual income, hence arising, is equally divided among the nominees, or heirs, of such members as die every year. But this society engages that the dividends shall not be less than 150l. to each claimant, though they may be more. No members are admitted whose ages are greater than 45, or less than 12; nor is any difference of contribution allowed on account of difference of age. The society has subsisted ever since the year 1706, and its credit and usefulness are well established.

The Equitable Society for Assurances on Lives and Survivorships, which meets at Black-Friars' Bridge, was established in the year 1762, in consequence of proposals which had been made, and lectures, recommending such a design, which had been read by Mr. Dodson, author of the Mathematical Repository. It assures any sums or reversionary annuities, on any life or lives for any number of years, as well as for the whole continuance of the lives; and in any manner that may be best adapted to the views of the persons assured: that is, either by making the assured sums payable certainly at the failure of any given lives; or on condition of survivorship; and also, either by taking the price of the assurance in one present payment, or in annual payments, during any single or joint lives, or any terms, less than the whole possible duration of the lives. In short, there are no kinds of assurances on lives and survivorships, which this society does not make.

In doing this, the Society follows the rules which have been given by the best mathematical writers on the doctrine of Life Annuities and Reversions, particularly Mr. Thomas Simpson, professor of Mathematics in the Royal Military Academy. It is to be observed however that the Society takes the advantage of making its calculations on the supposition that the interest of money is at so low a rate as 3 per cent, instead of the usual interest of 4 per cent; which consequently raises the insurance proportionally higher; and it also founds its calculations



on the tables of the probabilities and values of lives in London; another circumstance which secures a very advantageous profit to the Society, as experience has proved that the deaths are really in a much lower proportion than according to those tables, and even lower than those of Dr. Halley, which are founded on the bills of mortality of Breslaw. By these means the Society finding itself, by experience, well secured against future hazards, and being unwilling to take from the public an extravagant profit, have determined to reduce all the future payments for assurances, one tenth, and also generously to return, to the persons now assured, one tenth of all the payments they have made: and it seems there is reason to expect that this will be only a preparation to farther reduction.

From the foregoing account of this society, it is manifest that its business is such, that none but skilful mathematicians are qualified to conduct it. The interest of the society therefore requires, that it should make the places of those who manage its business sufficiently advantageous, to induce the ablest mathematicians to accept them: and this will render it the more necessary for the society to take care, in filling up any future vacancies, to pay no regard to any other considerations than the ability and integrity of the candidates. The consequence of granting good pay, will be a multitude of solicitations on every vacancy, from persons who, however unqualified, will hope for success from their connexions, and the interest they are able to make. And should the society, in any future time, be led by such causes to trust its business in the hands of persons not possessed of sufficient ability, as mathematicians and calculators, such mistakes may be committed, as may prove, in the highest degree, detrimental and dangerous. There is reason to believe, that at present the society is in no danger of this kind; and one of the great public advantages attending it, is, that it has established an office, where not only the business above described, is transacted with faithfulness and skill; but where also all persons, who want solutions of any questions relating to life annuities and reversions, may apply, and be sure of receiving just answers. The following is a

*Table of the rates of assurance on single lives in the Society for Equitable Assurances. The Sum assured 100l.*

Age.	For one year.			For seven years at an annual payment of			For the whole life at an annual payment of		
	l.	s.	d.	l.	s.	d.	l.	s.	d.
10	1	9	6	1	10	7	2	2	10
15	1	11	0	1	12	7	2	6	6
20	1	13	11	1	16	0	2	12	10
25	1	17	7	2	0	2	3	0	6
30	2	2	6	2	6	0	3	8	11
35	2	8	7	2	14	2	3	17	9
40	2	19	2	3	5	1	4	7	11
45	3	11	0	3	18	6	5	0	0
50	4	4	8	4	11	2	5	12	11
55	5	0	9	5	11	7	6	9	3
60	5	19	1	6	16	10	7	17	7
65	7	0	11	8	13	0	10	3	9

These rates are 10 per cent. lower than the true values, according to the decrements of life in London, reckoning interest at 3 per cent; but at the same time, it is to be observed that, for all ages under 50, they are near one third higher than all the true values, according to Dr. Halley's table of the decrements of life at Breslaw, and Dr. Price's tables of the decrements of life at Northampton and Norwich. But as the society has lately found that the decrements of life among its members have hitherto been lower than even those given in these last tables, it may reasonably be expected, that they will in time reduce their rates of assurance to the true values, as determined by these tables.

As to the *Westminster Society for granting Annuities, and insuring Money on Lives*, lately established, viz, in the year 1789, from the number and respectability of its members, the equitable terms upon which it proposes to deal, and the known ability and accuracy of the mathematicians and calculators employed in conducting it, there is every reason to expect an honourable and equitable treatment of the public, and a permanent continuance of its usefulness.

ASTERISM, the same with constellation; or a collection of many stars, which are usually represented on globes by some particular image or figure, to distinguish the stars which compose this constellation from those of others.

ASTRÆA, a name given by some to the sign Virgo, by others called Erigone, and sometimes Isis. The poets feign that Justice quitted heaven to reside on earth, in the golden age; but, growing weary of the iniquities of mankind, she left the earth, and returned to heaven, placing herself in that part of the zodiac called Virgo, where she became a constellation of stars, and from her orb still looks down on the ways of men. Ovid. Metam. lib. i. ver. 149.

ASTRAGAL, in Architecture, a small round moulding, which encompasses the top of the fust or shaft of a column, like a ring or bracelet. The shaft always terminates at top with an astragal, and at bottom with a fillet, which in this place is called *ozia*.

ASTRAGAL, in Gunnery, is a kind of ring or moulding on a piece of ordnance, at about half a foot distance from the muzzle or mouth; serving as an ornament to the gun, as the former does to a column.

ASTRAL, something belonging to or depending on the stars.

ASTRAL Year, or Sidereal Year. See YEAR.

ASTRODICTICUM, an astronomical instrument invented by M. Weighel, by means of which many persons shall be able to view the same star at the same time.

ASTROGNOSIA, the art of knowing the fixed stars, their names, ranks, situations in the constellations, and the like.

ASTROLABE, from *αστηρ*, star, and *λαβειν*, I take; alluding to its use in taking, or observing, the stars. The Arabians call it in their tongue *astharlab*; a word formed by corruption from the common Greek name.

This name was originally used for a system or assemblage of the several circles of the sphere, in their proper order and situation with respect to each other. And the ancient instruments were much the same as our armillary spheres.



The first, and most celebrated of this kind, was that of Hipparchus, which he made at Alexandria, the capital of Egypt, and lodged in a secure place, where it served for divers astronomical operations. Ptolemy made the same use of it: but as the instrument had several inconveniences, he contrived to change its figure, though perfectly natural, and agreeable to the doctrine of the sphere; and to reduce the whole Astrolabe to a plane surface, to which he gave the name of the *Planisphere*. Hence,

ASTROLABE is used among the moderns for a *Planisphere*, or a stereographic projection of the circles of the sphere upon the plane of one of the great circles; which is usually either the plane of the equinoctial, the eye being then placed in the pole of the world; or that of the meridian, the eye being supposed in the point of intersection of the equinoctial and horizon; or on that of the horizon.

The Astrolabe has been treated at large by Stöffler, Gemma Frisius, Clavius, &c. And for a farther account of the nature and kinds of it, see the article PLANISPHERE.

ASTROLABE, or SEA ASTROLABE, more particularly denotes an instrument chiefly used for taking altitudes at sea; as the altitude of the pole, the sun, or the stars.

The common Astrolabe, represented Plate II, *fig. 7*, consists of a large brass ring, about 15 inches in diameter, whose limb, or a convenient part of it, is divided into degrees and minutes. It is fitted with a moveable label or index, which turns upon the centre, and carries two sights; and having a small ring, at A, to hang it by in time of observation.

To make use of the Astrolabe in taking altitudes; suspend it by the ring A, and turn it to the sun, &c, so as that the rays may pass freely through both the sights F and G; then will the label cut or point out the altitude on the divided limb. There are many other uses of the Astrolabe; of which Clavius, Henrion, and others have written very largely.

The Astrolabe, though now grown into disuse, is by many esteemed equal to any other instrument for taking the altitude at sea; especially between the tropics, where the sun comes near the zenith.

ASTROLOGER, a person professing or practising the art of Astrology.

ASTROLOGICAL, something relating to Astrology.

ASTROLOGY, the art of foretelling future events, from the positions, aspects, and influences of the heavenly bodies.

The word is compounded of *αστηρ*, *star*, and *λογος*, *discourse*; whence, in the literal sense of the term, Astrology should signify no more than the *doctrine* or *science of the stars*; which indeed was its original acceptation, and constituted the ancient Astrology; which consisted formerly of both the branches now called Astronomy and Astrology, under the name of the latter only; and for the sake of making judiciary predictions it was that astronomical observations, properly so called, were chiefly made by the ancients. And though the two branches be now perfectly separated, and that of Astrology almost universally rejected by men of real learning, this has but lately been the case, as their union subsisted, in

some degree, from Ptolemy till Kepler, who had a strong bias towards the ancient astrology.

Astrology may be divided into two branches, *natural* and *judiciary*.

TO NATURAL ASTROLOGY belongs the predicting of natural effects; such as the changes of weather, winds, storms, hurricanes, thunder, floods, earthquakes, &c. But this art properly belongs to *Physiology*, or *Natural Philosophy*; and is only to be deduced, *à posteriori*, from phenomena and observations. And for this sort of Astrology it is that Mr. Boyle makes an apology in his History of the Air. Its foundation and merits may be gathered from what is said under the articles AIR, ATMOSPHERE, and WEATHER.

Judicial or Judiciary ASTROLOGY, which is what is commonly and properly called *Astrology*, is that which professes to foretel moral events, or such as have a dependence on the free will and agency of man; as if they were produced or directed by the stars.

The professors of this kind of Astrology maintain, "That the heavens are one great volume or book, wherein God has written the history of the world; and in which every man may read his own fortune, and the transactions of his time. The art, say they, had its rise from the same hands as Astronomy itself: while the ancient Assyrians, whose serene unclouded sky favoured their celestial observations, were intent on tracing the paths and periods of the heavenly bodies, they discovered a constant, settled relation of analogy, between them and things below; and hence were led to conclude these to be the *Parcæ*, the Destinies so much talked of, which preside at our births, and dispose of our future fate."

"The laws therefore of this relation being ascertained by a series of observations, and the share each planet has therein; by knowing the precise time of any person's nativity, they were enabled, from their knowledge in astronomy, to erect a scheme or horoscope of the situation of the planets, at that point of time; and hence, by considering their degrees of power and influence, and how each was either strengthened or tempered by some other, to compute what must be the result thereof."

Judicial Astrology, it is commonly said, was invented in Chaldaea, and from thence transmitted to the Egyptians, Greeks, and Romans; though some insist that it was of Egyptian origin, and ascribe the invention to Cham. But it is to the Arabs that we owe it. At Rome the people were so infatuated with it, that the astrologers, or, as they were then called, the mathematicians, maintained their ground in spite of all the edicts of the emperors to expel them out of the city. See GENETHLIACI.

Among the Indians, the Bramins, who introduced and practised this art in the East, have hereby made themselves the arbiters of good and evil hours, which, gives them great authority: they are consulted as oracles; and they have taken care always to sell their answers at good rates.

The same superstition has prevailed in more modern ages and nations. The French historians remark, that in the time of queen Catharine de Medicis, Astrology was so much in vogue, that the most inconsiderable thing was not to be done without consulting the stars. And in the reigns of king Henry III. and IV. of France, the predictions



predictions of Astrologers were the common theme of the court conversation. And this predominant humour in that court was well rallied by Barclay, in his *Argenis*, lib. 2, on occasion of an Astrologer, who had undertaken to instruct king Henry in the event of a war which was then threatened by the faction of the Guises.

**ASTROMETEOROLOGIA**, the art of foretelling the weather, and its changes, from the aspects and configurations of the moon and planets: a species of Astrology distinguished by some under the denomination of *meteorological astrology*.

**ASTRONOMICAL**, something relating to Astronomy.

**ASTRONOMICAL Calendar**, *Characters, Column, Horizon, Hours, Month, Quadrant, Ring-Dial, Sector, Tables, Telescope, Time, Year*. See the several substantives.

**ASTRONOMICAL Observations**. Of these there are records, or mention, in almost all ages. It is said that the Chinese have observations for a course of many thousand years. But of these, as well as those of the Indians, we have never yet had any benefit. But the observations of most of the other ancients, as Babylonians, Greeks, &c, amongst which those of Hipparchus make a principal figure, are carefully preserved by Ptolemy, in his *Almagest*.

About the year 880, Albategni, a Saracen, applied himself to the making of observations; in which he was followed by others of the same nation, as well as Persians and Tartars; among whom were Nassir-Eddin-Ettusi, Arzachel, who also constructed a table of sines, and Ulug Beigh. In 1457 Regiomontanus undertook the province at Norimberg; and his disciples, J. Werner and Ber. Walther, continued the same from 1475 to 1504. Their observations were published together in 1544.—In 1509, Copernicus, and after him the landgrave of Hesse, with his assistants Rothman and Byrge, observed; and after them Tycho Brahe, assisted by the celebrated Kepler, from 1582 to 1601.—All the foregoing observations, together with Tycho's apparatus of instruments, are contained in the *Historia Cœlestis*, published in 1672, by order of the emperor Ferdinand.—In 1651, was published at Bononia, by Ricciolus, *Almagestum Novum*, being a complete body of ancient and modern observations, which he so named after the work of the same nature by Ptolemy.—Soon after, Hevelius, with a magnificent and well-contrived apparatus of instruments, described in his *Machina Cœlestis*, began a course of observations. It has been objected to him, that he only used plain sights, and could never be brought to take the advantage of telescopic ones; which occasioned Dr. Hook to write animadversions on Hevelius's instruments, printed in 1674, in which he too rashly despises them, on account of their inaccuracy: but Dr. Halley, who at the instance of the Royal Society went over to Dantzick in the year 1679, to inspect his instruments, approved of their justness, as well as of the observations made with them. See **SIGHS**.—Our two countrymen Jer. Horrox and Will. Crabtree, are celebrated for their observations from the year 1635 to 1645, who first observed the transit of Venus over the sun in the year 1639.—They were followed by Flamsteed, Cassini the father and son, Halley, de la Hire, Roemer, and Kirchius.—The observations of the celebrated Dr. Bradley have not yet been published, though long expected. We have also now published,

from time to time, the accurate observations of the present British Astronomer Royal: as also those of the French and other observatories, with the observations of many ingenious private astronomers, which are to be found in the *Transactions and Memoirs* of the various Philosophical Societies.—There have been also observations of many other eminent astronomers; as, Galileo, Huygens, and our countryman Harriot, whose very interesting observations have lately been brought to light by the earl of Egremont, and count Bruhl, by whose means they may come to be published. Other publications of celestial observations, are those of Cassini, La Caille, Monnier, &c.—See farther under **CELESTIAL Observations, CATALOGUE, OBSERVATORY, &c.**

**ASTRONOMICAL Place** of a star or planet, is its longitude, or place in the ecliptic, reckoned from the beginning of Aries, *in consequentia*, or according to the order of the signs.

**ASTRONOMICALS**, a name used by some writers for sexagesimal fractions; on account of their use in astronomical calculations.

**ASTRONOMICUS Radius**. See **RADIUS**.

**ASTRONOMY**, the doctrine of the heavens, and their phenomena.

Astronomy is properly a mixed mathematical science, by which we become acquainted with the celestial bodies, their motions, periods, eclipses, magnitudes, distances, and other phenomena. Some, however, understand the term astronomy in a more extensive sense, as comprising in it the theory of the universe, with the primary laws of nature: in which sense it seems to be rather a branch of physics than of mathematics.

#### *History of Astronomy.*

The invention of astronomy has been variously given, and ascribed to several persons, several nations, and several ages. Indeed it is probable that mankind never existed without some knowledge of astronomy amongst them. For, besides the motives of mere curiosity, which are sufficient of themselves to have excited men to a contemplation of the glorious and varying celestial canopy, it is obvious that some parts of the science answer such essential purposes to mankind, as to make the cultivation of it a matter of indispensable necessity. Accordingly we find traces of it, in different degrees of improvement, among all nations.

Adam, in his state of innocence, it is supposed by some of the Jewish rabbins, was endowed with a knowledge of the nature, influence, and uses of the heavenly bodies; and Josephus ascribes to Seth and his posterity a considerable knowledge of astronomy: he speaks of two pillars, the one of stone and the other of brick, called the pillars of Seth, upon which they engraved the principles of the science; and he says that the former was still entire in his time. But be this as it may, it is evident that the great length of the antediluvian lives would afford such excellent opportunities for observing the heavenly bodies, that we cannot but suppose that the science of astronomy was considerably advanced before the flood. Indeed Josephus says that longevity was bestowed upon them for the very purpose of cultivating the sciences of geometry and astronomy; observing that the latter could not be learned in less than 600 years; “for that period, he adds, is the *grand year*.”



An expression remarkable enough; and by which it may be supposed is meant the period in which the sun and moon come again into the same situation in which they were at the beginning of it, with regard to the nodes, apogee of the moon, &c. "This period, says Cassini, of which we find no intimation in any monument of any other nation, is the finest period that ever was invented: for it brings out the solar year more exactly than that of Hipparchus and Ptolemy; and the lunar month within about one second of what is determined by modern astronomers." If the Antediluvians had such a period of 600 years, they must have known the motions of the sun and moon more exactly than their descendants knew them some ages after the flood.

On the building of the Tower of Babel, it is supposed that Noah retired with his children, born after the flood, to the north-eastern part of Asia, where his descendants peopled the vast empire of China. And this, says Dr. Long, "may perhaps account for the Chinese having so early cultivated the study of astronomy, &c." It is said that the Jesuit missionaries have found traditional accounts among the Chinese, of their having been taught this science by their first emperor Fo-hi, who is supposed to be the same with Noah; and Kempfer asserts that Fo-hi discovered the motions of the heavens, divided time into years and months, and invented the 12 signs into which they divide the zodiac, and which they distinguish by these names following; 1, the mouse; 2, the ox or cow; 3, the tiger; 4, the hare; 5, the dragon; 6, the serpent; 7, the horse; 8, the sheep; 9, the monkey; 10, the cock or hen; 11, the dog; and, 12, the boar. They divide the heavens into 28 constellations, or classes of stars, allotting 4 to each of the 7 planets; so that the year always begins with the same planet; and their constellations answer to the 28 lunar mansions used by the Arabian astronomers. These constellations however they do not mark with the figures of animals, like most other nations, but by connecting the stars by straight lines, and denoting the stars themselves by small circles: so, for instance, the great bear would be marked thus,



The Chinese themselves have many records and traditions of the high antiquity of their astronomy; though not without suspicion of great mistakes. But, on more certain authority, it is asserted by F. Gaubil, that at least 120 years before Christ, the Chinese had determined by observation the number and extent of their constellations as they now stand; the situation of the fixed stars with respect to the equinoctial and solstitial points; and the obliquity of the ecliptic; with the theory of eclipses: and that they were, long before that, acquainted with the true length of the solar year, the method of observing meridian altitudes of the sun by the shadow of a gnomon, and of deducing from thence his declination, and the height of the pole. The same missionary also says, that the Chinese have yet remaining some books of astronomy, which were written about 200 years before Christ; from which it appears,

that the Chinese had known the daily motion of the sun and moon, and the times of the revolutions of the planets, many years before that period.

Du Halde informs us, that Tcheou-cong, the most skilful astronomer that ever China produced, lived more than a thousand years before Christ; that he passed whole nights in observing the celestial bodies, and arranging them into constellations, &c. At present however, the state of astronomy is but very low in that country, although it be cultivated at Peking, by public authority, in like manner as in most of the capital cities of Europe.

The inhabitants of Japan, of Siam, and of the Mogul's empire, have also been acquainted with astronomy from time immemorial; and the celebrated observatory at Benares, is a monument both of the ingenuity of the people, and of their skill in that science.

According to Porphyry, astronomy must have been of very ancient standing in the East. He informs us that, when Babylon was taken by Alexander, there were brought from thence celestial observations for the space of 1903 years; which therefore must have commenced within 115 years after the flood, or within 15 years after the building of Babel.—Epigenes, according to Pliny, affirmed that the Babylonians had observations of 720 years engraven on bricks.—Again, Achilles Tatius ascribes the invention of astronomy to the Egyptians; and adds, that their knowledge of that science was engraven on pillars, and by that means transmitted to posterity.

M. Bailly, in his elaborate History of ancient and modern astronomy, endeavours to trace the origin of this science among the Chaldeans, Egyptians, Persians, Indians and Chinese, to a very early period. And thence he maintains, that it was cultivated in Egypt and Chaldea 2800 years before Christ; in Persia, 3209; in India, 3101; and in China, 2952 years before that æra. He also apprehends, that astronomy had been studied even long before this distant period, and that we are only to date its revival from thence.

In investigating the antiquity and progress of astronomy among the Indians, M. Bailly examines and compares four different sets of astronomical tables of the Indian philosophers, namely that of the Siamese, explained by M. Cassini in 1689; that brought from India by M. le Gentil of the Academy of Sciences; and two other manuscript tables, found among the papers of the late M. de Lisle; all of which he found to accord together, and all referring to the meridian of Benares, above-mentioned. It appears that the fundamental epoch of the Indian astronomy, is a conjunction of the sun and moon, which took place at the amazing distance of 3102 years before Christ: and M. Bailly informs us that, by our most accurate astronomical tables, such a conjunction did really happen at that time. He further observes that, at present, the Indians calculate eclipses by the mean motions of the sun and moon observed 5000 years since; and that their accuracy, with regard to the solar motion, far exceeds that of the best Grecian astronomers. They had also settled the lunar motions by computing the space through which that luminary had passed in 1,600,984 days, or a little more than 4383 years. M. Bailly also informs us, that they make use of the cycle of 19 years, the same as that ascribed



ascribed by the Greeks to Meton; that their theory of the planets is much better than Ptolemy's, as they do not suppose the earth in the centre of the celestial motions, and believe that Venus and Mercury move round the sun; and that their astronomy agrees with the most modern discoveries as to the decrease of the obliquity of the ecliptic, the acceleration of the motion of the equinoctial points, &c.

In the 2d vol. of the Transactions of the Royal Society of Edinburgh is also a learned and ingenious dissertation on the astronomy of the Brahmins of India, by Mr. Professor Playfair; in which the great accuracy and high antiquity of the science, among them, is reduced to the greatest probability. It hence appears that their tables and rules of computation, have peculiar reference to an epoch, and to observations, 3 or 4 thousand years before Christ; and many other instances are there adduced, of their critical knowledge in the other mathematical sciences, employed in their precepts and calculations.

Astronomy, it seems, too, was not unknown to the Americans; though in their division of time, they made use only of the solar, and not of the lunar motions. And that the Mexicans, in particular, had a strange predilection for the number 13, by means of which they regulated almost every thing: their shortest periods consisted of 13 days; their cycle of 13 months, each containing 20 days; and their century of 4 periods, of 13 years each: and this excessive veneration for the number 13, arose, according to Siguenza, from its being the number of their greater gods. And it is very remarkable, that the Abbé Clavigero asserts it as a fact, that, having discovered the excess of a few hours in the solar above the lunar year, they made use of intercalary days, to bring them to an equality, as established by Julius Cæsar in the Roman Calendar; but with this difference, that, instead of one day every 4 years, they interposed 13 days every 52 years, which produces the same effect.

Most authors however fix the origin of astronomy and astrology, either in Chaldea or in Egypt; and accordingly among the ancients we find the word Chaldean often used for astronomer, or, which was the same thing, astrologer. Indeed both of these nations pretended to a very high antiquity, and claimed the honour of producing the first cultivators of this science. The Chaldeans boasted of their temple or tower of Belus, and of Zoroaster, whom they placed 5000 years before the destruction of Troy; while the Egyptians boasted of their colleges of priests, where astronomy was taught, and of the monument of Osymandyas, in which, it is said, there was a golden circle of 365 cubits in circumference, and one cubit thick, divided into 365 equal parts according to the days of the year, &c.

It is, indeed, evident, that both Chaldea and Egypt were countries very proper for astronomical observations, on account of the extended flatness of the country, and the purity and serenity of the air. The tower of Belus, or of Babel itself, of a great height, was probably an astronomical observatory; and the lofty pyramids of Egypt, whatever they were originally designed for, might perhaps answer the same purpose; and at least they shew the skill of this people in practical astronomy, as they are all placed with their four fronts exactly

facing the cardinal points of the compass. The Chaldeans certainly began to make observations soon after the confusion of languages, as appears from the observations found there on the taking of Babylon by Alexander; and it is probable they began much earlier. It hence appears that they had determined, with tolerable exactness, the length both of a periodical and synodical month. They had also discovered, that the motion of the moon was not uniform; and they even attempted to assign those parts of the orbit in which the motion is quicker or slower. We are also assured by Ptolemy that they were not unacquainted with the motion of the moon's apogee and nodes, the latter of which they supposed made a complete revolution in  $6585\frac{1}{2}$  days, or a little more than 18 years, and contained 223 complete lunations, which period is called the Chaldean *Saros*. From Hipparchus, the same author also gives us several observations of lunar eclipses made at Babylon above 720 years before Christ. And Aristotle informs us, that they had many occultations of the planets and fixed stars by the moon; a circumstance which led them to conceive that eclipses of the sun were to be attributed to the same cause. They had also no inconsiderable share in arranging the stars into constellations. Nor had even those eccentric bodies the comets escaped their observation: for both Diodorus Siculus and Appollinus Myndicus, Seneca informs us, accounted these to be permanent bodies, having stated revolutions as well as the planets, but in much more extensive orbits: although others of them were of opinion, that the comets were only meteors raised very high in the air, which, blazing for a while, disappear when the matter of which they consist is consumed or dispersed. The branch of dialling was also practised among them long before the Greeks were acquainted with that science.

The Egyptians, it appears from various circumstances, were much of the same standing in Astronomy as the Chaldeans. Herodotus ascribes their knowledge in the science to Sesostris; probably not the same whom Newton makes contemporary with Solomon, as they were acquainted with astronomy at least many hundred years before that æra. We learn, from the testimony of some ancient authors, many particulars relative to the state of their knowledge in astronomy; such as, that they believed the figure of the earth was spherical; that the moon was eclipsed by passing through the earth's shadow, though it does not certainly appear that they had any knowledge of the true system of the universe; that they attempted to measure the magnitude of the earth and sun, though their methods of ascertaining the latter were very erroneous; and that they even pretended to foretel the appearance of comets, as well as earthquakes and inundations; and the same is also ascribed to the Chaldeans; though these must probably have been rather a kind of astrological predictions, than observations drawn from astronomy, properly so called.

This science however fell into great decay with the Egyptians, and in the time of the emperor Augustus, it was entirely extinct among them.

From Chaldea and Egypt the science of astronomy passed into Phenicia, which this people applied to the purposes of navigation, steering their course by the north polar star; and hence they became masters of the sea, and of almost all the commerce in the world.



The Greeks, it is probable, derived their astronomical knowledge chiefly from the Egyptians and Phenicians, by means of several of their countrymen who visited these nations, for the purpose of learning the different sciences. Newton supposes that most of the constellations were invented about the time of the Argonautic expedition; but it is more probable that they were, at least most part of them, of a much older date, and derived from other nations, though clothed in fables of their own invention or application. Several of the constellations are mentioned by Hesiod and Homer, the two most ancient writers among the Greeks, and who lived about 870 years before Christ. Their knowledge in this science however was greatly improved by Thales the Milesian, and other Greeks, who travelled into Egypt, and brought from thence the chief principles of the science. Thales was born about 640 years before Christ; and he, first of all among the Greeks, observed the stars, the solstices, the eclipses of the sun and moon, and predicted the same. And the same was farther cultivated and extended by his successors Anaximander, Anaximanes, and Anaxagoras; but most especially by Pythagoras, who was born 577 years before Christ, and having resided for several years in Egypt, &c, brought from thence the learning of these people, taught the same in Greece and Italy, and founded the sect of the Pythagoreans. He taught that the sun was in the centre of the universe; that the earth was round, and people had antipodes; that the moon reflected the rays of the sun, and was inhabited like the earth; that comets were a kind of wandering stars, disappearing in the further parts of their orbits; that the white colour of the milky-way was owing to the united brightness of a great multitude of small stars; and he supposed that the distances of the moon and planets from the earth, were in certain harmonic proportions to one another.

Philolaus, a Pythagorean, who flourished about 450 years before Christ, asserted the annual motion of the earth about the sun; and not long after, the diurnal motion of the earth on her own axis, was taught by Hicetas, a Syracusan. About the same time flourished at Athens, Meton and Euctemon, where they observed the summer solstice 432 years before Christ, and observed the risings and settings of the stars, and what seasons they answered to. Meton also invented the cycle of 19 years, which still bears his name.

Eudoxus the Cnidian lived about 370 years before Christ, and was accounted one of the most skilful astronomers and geometricians of antiquity, being accounted the inventor of many of the propositions in Euclid's Elements, and having introduced geometry into the science of astronomy. He travelled into Asia, Africa, Sicily, and Italy, for improvements in astronomy; and we are informed by Pliny, that he determined the annual year to contain 365 days 6 hours, that he determined also the periodical times of the planets, and made other important observations and discoveries.

Calippus flourished soon after Eudoxus, and his celestial sphere is mentioned by Aristotle; but he is better known by a period of 76 which he invented, containing 4 corrected Metonic periods, and which commenced at the summer solstice in the year 330 before Christ. About his time the knowledge of the Pythagorean system was

carried into Italy, Gaul, and Egypt, by certain colonies of Greeks.

However, the introduction of Astronomy into Greece is represented by Vitruvius in a manner somewhat different. He maintains, that Berofus, a Babylonian, brought it immediately from Babylon itself, and opened an astronomical school in the isle of Cos. And Pliny says, that in consideration of his wonderful predictions, the Athenians erected him a statue in the gymnasium, with a gilded tongue. But if this Berofus be the same with the author of the Chaldaic histories, he must have lived before Alexander.

After the death of this conqueror, the sciences flourished chiefly in Egypt, under the auspices of Ptolemy Philadelphus and his successors. He founded a school there, which continued to be the grand seminary of learning, till the invasion of the Saracens in the year of Christ 650. From the founding of that school, the science of astronomy advanced considerably. Aristarchus, about 270 years before Christ, strenuously asserted the Pythagorean system, and gave a method of determining the sun's distance by the dichotomy of the moon.—Eratothenes, who was born at Cyrene in the year 271 before Christ, measured the circumference of the earth by means of a gnomon; and being invited to Alexandria, from Athens, by Ptolemy Euergetes, and made keeper of the royal library there, he set up for that Prince those armillary spheres, which Hipparchus and Ptolemy the astronomer afterwards employed so successfully in observing the heavens. He also determined the distance between the tropics to be  $\frac{1}{4}$  of the whole meridian circle, which makes the obliquity of the ecliptic in his time to be  $23^{\circ} 51' \frac{1}{3}$ .—The celebrated Archimedes, too, cultivated astronomy, as well as geometry and mechanics: he determined the distances of the planets from one another, and constructed a kind of planetarium or orrery, to represent the phenomena and motions of the heavenly bodies.

To pass by several others of the ancients, who practised or cultivated astronomy, more or less, we find that Hipparchus, who flourished about 140 years before Christ, was the first who applied himself to the study of every part of astronomy, and, as we are informed by Ptolemy, made great improvements in it: he discovered that the orbits of the planets are eccentric, that the moon moved slower in the apogee than in her perigee, and, that there was a motion of anticipation of the moon's nodes: he constructed tables of the motions of the sun and moon, collected accounts of such eclipses, &c, as had been made by the Egyptians and Chaldeans, and calculated all that were to happen for 600 years to come: he discovered that the fixed stars changed their places, having a slow motion of their own from west to east: he corrected the Calippic period, and pointed out some errors in the method of Eratothenes for measuring the circumference of the earth: he computed the sun's distance more accurately than any of his predecessors: but his chief work is a catalogue which he made of the fixed stars, to the number of 1022, with their longitudes, latitudes, and apparent magnitudes; which, with most of his other observations, are preserved by Ptolemy in his *Almagest*.

There was but little progress made in astronomy from the time of Hipparchus to that of Ptolemy, who was born



born at Pelusium in Egypt, in the first century of christianity, and who made the greatest part of his observations at the celebrated school of Alexandria in that country. Profiting of those of Hipparchus and other ancient astronomers, he formed a system of his own, which, though erroneous, was followed for many ages by all nations. He compiled a great work, called the *Almagest*, which contained the observations and collections of Hipparchus and others his predecessors in astronomy, on which account it will ever be valuable to the professors of that science. This work was preserved from the grievous conflagration of the Alexandrine library by the Saracens, and translated out of Greek into Arabic in the year 827, and from thence into Latin in 1230. The Greek original was not known in Europe till the beginning of the 15th century, when it was brought from Constantinople, then taken by the Turks, by George, a monk of Trapezond, by whom it was translated into Latin; and various other editions have been since made.

During the long period from the year 800 till the beginning of the 14th century, the western parts of Europe were immersed in gross ignorance and barbarity, while the Arabians, profiting by the books they had preserved from the wreck of the Alexandrine library, cultivated and improved all the sciences, and particularly that of astronomy, in which they had many able professors and authors. The caliph Al Mansur first introduced a taste for the sciences into his empire. His grandson Al Mamun, who ascended the throne in 814, was a great encourager and improver of the sciences, and especially of astronomy. Having constructed proper instruments, he made many observations; determined the obliquity of the ecliptic to be  $23^{\circ} 35'$ ; and under his auspices a degree of the circle of the earth was measured a second time in the plain of Singar, on the border of the Red Sea. About the same time Alferganus wrote elements of astronomy; and the science was from hence greatly cultivated by the Arabians, but principally by Albategnius, who flourished about the year 880, and who greatly reformed astronomy, by comparing his own observations with those of Ptolemy: hence he computed the motion of the sun's apogee from Ptolemy's time to his own; settled the precession of the equinoxes at one degree in 70 years; and fixed the obliquity of the ecliptic at  $23^{\circ} 35'$ . The tables which he composed, for the meridian of Aracta, were long esteemed by the Arabians. After his time, though the Saracens had many eminent astronomers, several centuries elapsed without producing any very valuable observations, excepting those of some eclipses observed by Ebn Younis, astronomer to the caliph of Egypt, by means of which the quantity of the moon's acceleration since that time may be determined.

Other eminent Arabic astronomers, were, Arzachel a Moor of Spain, who observed the obliquity of the ecliptic: he also improved Trigonometry by constructing tables of sines, instead of chords of arches, dividing the diameter into 300 equal parts. And Alhazen, his contemporary, who wrote upon the twilight, the height of the clouds, the phenomenon of the horizontal moon, and who first shewed the importance of the theory of refractions in astronomy.

Ulug Beg, grandson of the celebrated Tartar prince Tamerlane, was a great proficient in practical astronomy;

he had very large instruments, particularly a quadrant of about 180 feet high, with which he made good observations. From these he determined the latitude of Samercand, his capital, to be  $39^{\circ} 37' 23''$ ; and composed astronomical tables for the meridian of the same so exact, that they differ very little from those constructed afterwards by Tycho Brahe; but his principal work was his catalogue of the fixed stars, made also from his own observations in the year 1437.

During this period, almost all Europe was immersed in gross ignorance. But the settlement of the Moors in Spain introduced the sciences into Europe; from which time they have continued to improve, and to be communicated from one people to another, to the present time, when astronomy and all the sciences, have arrived at a very eminent degree of perfection. The emperor Frederick II, about 1230, first began to encourage learning; restoring some decayed universities, and founding a new one in Vienna: he also caused the works of Aristotle, and Ptolemy's *Almagest*, to be translated into Latin; and from the translation of this work we may date the revival of astronomy in Europe. Two years after this, John de Sacro Bosco, that is, of Halifax, compiled, from Ptolemy, Albategnius, Alferganus, and other Arabic astronomers, his work *De Sphæra*, which was held in the greatest estimation for 300 years after, and was honoured with commentaries by Clavius and other learned men. In 1240, Alphonso, king of Castile, not only cultivated astronomy himself, but greatly encouraged others; and by the assistance of several learned men he corrected the tables of Ptolemy, and composed those which were denominated from him the Alphonine Tables. About the same time also, Roger Bacon, an English monk, wrote several tracts relative to astronomy, particularly of the lunar aspects, the solar rays, and the places of the fixed stars. And, about the year 1270, Vitello, a Polander, composed a treatise on optics, in which he shewed the use of refractions in astronomy.

Little other improvement was made in astronomy till the time of Purbach, who was born in 1423. He composed new tables of sines for every 10 minutes, making the radius 60, with four ciphers annexed. He constructed spheres and globes, and wrote several astronomical tracts; as, a commentary on Ptolemy's *Almagest*; some treatises on Arithmetic and Dialling, with tables for various climates; new tables of the fixed stars reduced to the middle of that century; and he corrected the tables of the planets, making new equations to them where the Alphonine tables were erroneous. In his solar tables, he placed the sun's apogee in the beginning of Cancer; but retained the obliquity of the ecliptic  $23^{\circ} 33 \frac{1}{2}'$ , as determined by the latest observations. He also observed some eclipses, made new tables for computing them, and had just finished a theory of the planets, when he died in 1462, being only 39 years of age.

Purbach was succeeded in his astronomical and mathematical labours by his pupil and friend, John Muller, commonly called Regiomontanus, from Montereio, or Konigsberg, a town of Franconia, where he was born. He completed the epitome of Ptolemy's *Almagest*, which Purbach had begun; and after the death of his friend, was invited to Rome, where he made many astronomical



astronomical observations. Being returned to Nuremberg in 1471, by the encouragement of a wealthy citizen named Bernard Walther, he made several instruments for astronomical observations, among which was an armillary astrolabe, like that used at Alexandria by Hipparchus and Ptolemy, with which he made many observations, using also a good clock, which was then but a late invention. He made ephemerides for 30 years to come, shewing the lunations, eclipses, &c; and, the art of printing having then been lately invented, he printed the works of many of the most celebrated ancient astronomers. He wrote the Theory of the Planets and Comets, and a treatise on triangles, still in repute for several good theorems; computing the table of sines for every single minute, to the radius 1000000, and introducing the use of tangents also into trigonometry. After his death, which happened at Rome in 1476, being only 40 years of age, Walther collected his papers, and continued the astronomical observations till his own death also. The observations of both were collected by order of the senate of Nuremberg, and published there in 1544 by John Schoner: they were also afterwards published in 1618 by Snellius, at the end of the observations made by the Landgrave of Hesse; and lastly with those of Tycho Brahe in 1666.

Walther was succeeded, as astronomer at Nuremberg, by John Werner, a clergyman. He observed the motion of the comet in 1530; and wrote several tracts on geometry, astronomy, and geography, in a masterly manner; the most remarkable of which, are those concerning the motion of the 8th sphere, or of the fixed stars; in this tract, by comparing his own observations, made in 1514, with those of Ptolemy, Alphonsus, and others, he shewed that the motion of the fixed stars, since called the precession of the equinoxes, is  $1^{\circ} 10'$  in 100 years. He made also the first star of Aries  $26^{\circ}$  distant from the equinoctial point, and the obliquity of the ecliptic only  $23^{\circ} 28'$ . He constructed a planetarium, representing the celestial motions according to the Ptolemaic hypothesis; and he published a translation of Ptolemy's Geography, with a commentary, in which he first proposed the method of finding the longitude at sea by observing the moon's distance from the fixed stars; now so successfully practised for that purpose. Werner died in 1528, at 60 years of age.

Nicolaus Copernicus was the next who made any considerable figure in astronomy, by whom indeed the old Pythagorean system of the world was restored, which had been till now set aside from the time of Ptolemy. About the year 1507 Copernicus conceived doubts of this system, and entertained notions about the true one, which he gradually improved by a series of astronomical observations, and the contemplation of former authors. By these he formed new tables, and completed his work in the year 1530, containing these, and his renovation of the true system of the universe, in which all the planets are considered as revolving about the sun, placed in the centre. But the work was only printed in 1543, under the care of Schoner and Osiander, by the title of *Revolutiones Orbium Caelestium*; and the author just received a copy of the work a few hours before his death, which happened on the 23d of May 1543, at 70 years of age.

After the death of Copernicus, the science and practice of astronomy were greatly improved by many other persons, as Schoner, Nonius, Appian, Gemma Frisius, Rothman, Byrgius, the Landgrave of Hesse, &c.—Schoner reformed and explained the calendar, improved the methods of making celestial observations, and published a treatise on cosmography; but he died 4 years after Copernicus.—Nonius wrote several works on mathematics, astronomy and navigation, and invented some useful and more accurate instruments than formerly; one of these was the astronomical quadrant, on which he divided the degrees into minutes by a number of concentric circles; the first of which was divided in 90 equal parts or degrees, the second into 89, the third into 88, and so on, to 46; so that, the index of the quadrant always falling upon or near one of the divisions, the minutes would be known by an easy computation.—The chief work of Appian, *The Cæsarean Astronomy*, was published at Ingoldstat in 1540; in which he shews, how to observe the places of the stars and planets by the astrolabe; to resolve astronomical problems by certain instruments; to predict eclipses, and to describe the figures of them; and the method of dividing and using an astronomical quadrant: at the end are added observations of 5 comets, one of which has been supposed the same with that observed by Hevelius, and if so, it ought to have returned again in the year 1789;—but it was not observed then. Gemma Frisius wrote a commentary on Appian's *Cosmography*, accompanied with many observations of eclipses: he also invented the astronomical ring, and several other instruments, useful in taking observations at sea; and he was the first who recommended a time-keeper for determining the longitude at sea.—Rheticus gave up his professorship of mathematics at Wittemberg, that he might attend the astronomical lectures of Copernicus; and, for improving astronomical calculations, he began a very extensive work, being a table of sines, tangents and secants, to a very large radius, and to every 10 seconds, or  $\frac{1}{6}$  of a minute; which was completed by his pupil Valentine Otho, and published in 1594.

About the year 1561, William IV, Landgrave of Hesse Cassel, applied himself to the study of astronomy, having furnished himself with the best instruments that could then be made: with these he made a great number of observations, which were published by Snellius in 1618, and which were preferred by Hevelius to those of Tycho Brahe. From these observations he formed a catalogue of 400 stars, with their latitudes and longitudes, adapted to the beginning of the year 1593.

Tycho Brahe, a noble Dane, began his observations about the same time with the Landgrave of Hesse, above-mentioned, and he observed the great conjunction of Jupiter and Saturn: but finding the usual instruments very inaccurate, he constructed many others much larger and exacter, with which he applied himself diligently to observe the celestial phenomena. In 1571 he discovered a new star in the chair of Cassiopeia; which induced him, like Hipparchus on a similar occasion, to make a new catalogue of the stars; which he composed to the number of 777, and adapted their places to the year 1600. In the year 1576, by favour of the king of Denmark, he built his new observatory, called Uraniburg, on the small island Huenna, opposite to



to Copenhagen, and which he very amply furnished with many large instruments, some of them so divided as to shew single minutes, and in others the arch might be read off to 10 seconds. One quadrant was divided according to the method invented by Nonius, that is, by 47 concentric circles; but most of them were divided by diagonals; a method of division invented by a Mr. Richard Chancellor, an Englishman. Tycho employed his time at Uraniburg to the best advantage, till the death of the king, when, falling into discredit, he was obliged to remove to Holstein; and he afterwards found means of introducing himself to the Emperor Rodolph, with whom he continued at Prague till the time of his death in 1601.—It is well known that Tycho was the inventor of a system of astronomy, a kind of Semi-Ptolemaic, which he vainly endeavoured to establish instead of the Copernican or true system. His works, however, which are very numerous, shew that he was a man of great abilities; and his discoveries, together with those of Purbach and Regiomontanus, were collected and published together in 1621, by Longomontanus, the favourite disciple of Tycho.

While Tycho resided at Prague with the emperor, he prevailed on Kepler to leave the university of Glatz, and to come to him, which he did with his family and library in 1600: but Tycho dying in 1601, Kepler enjoyed all his life the title of mathematician to the Emperor, who ordered him to finish the tables of Tycho Brahe, which he did accordingly, and published them in 1627, under the title of Rodolphine. He died about the year 1630 at Ratisbon, where he was soliciting the arrears of his pension. From his own observations, and those of Tycho, Kepler discovered several of the true laws of nature, by which the motions of the celestial bodies are regulated. He discovered that all the planets revolved about the sun, not in circular, but in elliptical orbits, having the sun in one of the foci of the ellipse; that their motions were not equable, but varying, quicker or slower as they were near to the sun or farther from him; but that this motion was so regulated, that the areas described by the variable line drawn from the planet to the sun, are equal in equal times, and always proportional to the times of describing them. He also discovered, by trials, that the cubes of the distances of the planets from the sun, were in the same proportion as the squares of their periodical times of revolution. By observations also on comets, he concluded that they are freely carried about among the orbits of the planets, in paths that are nearly rectilinear, but which he could not then determine. Besides many other discoveries, which are to be found in his writings.

In Kepler's time there were many other good proficient in astronomy; as Edward Wright, baron Napier, John Bayer, &c. Wright made several good meridional observations of the sun, with a quadrant of 6 feet radius, in the years 1594, 1595, and 1596; from which he greatly improved the theory of the sun's motion, and computed more accurately his declination, than any person had done before. In 1599 he published also an excellent work entitled "Certain Errors in Navigation discovered and detected," containing a method which has commonly, though erroneously, been ascribed to Mercator.—To Napier we owe some excellent theorems and improvements in spherics, besides the ever memo-

orable invention of logarithms, one of the most useful ever made in the art of numbering, and of the greatest use in all the other mathematical sciences.—Bayer, a German, published his *Uranometria*, being a complete celestial atlas, or the figures of all the constellations visible in Europe, with the stars marked on them, and the stars also accompanied by names, or the letters of the Greek alphabet; a contrivance by which the stars may easily be referred to with distinctness and precision.—About the same time too, astronomy was cultivated by many other persons; namely, abroad by Mercator, Maurolycus, Maginus, Homelius, Schultet, Stevin, Galileo, &c; and in England by Thomas and Leonard Digges, John Dee, Robert Flood, Harriot, &c.

The beginning of the 17th century was particularly distinguished by the invention of telescopes, and the application of them to astronomical observations; an invention to which we owe the most brilliant discoveries, and all the accuracy to which the science is now brought. The more distinguished early observations with the telescope, were made by Galileo, Harriot, Huygens, Hook, Cassini, &c. It is said that from report only, and before he had seen one, Galileo made for himself telescopes, by which he discovered inequalities in the moon's surface, Jupiter's satellites, and the ring of Saturn; also spots on the sun, by which he found out the revolution of that luminary on his axis; and he discovered that the nebulae and milky way were full of small stars. Harriot also, who has hitherto been known only as an algebraist, made much the same discoveries as Galileo, and as early, if not more so, as appears by his papers not yet printed, in the possession of the earl of Egremont.

Mr. Horrox, a young astronomer of great talents, made considerable discoveries and improvements. In 1633 he found out that the planet Venus would pass over the sun's disc on the 24th of November, 1639; an event which he announced only to his friend Mr. Crabtree; and these two were the only persons in the world that observed this transit, which was also the first time it had ever been seen by mortal eyes. Mr. Horrox made also many other useful observations, and had even formed a new theory of the moon, taken notice of by Newton; but his early death, in the beginning of the year 1640, put a stop to his useful and valuable labours.

About the same time flourished Hevelius, Burgomaster of Dantzic, who furnished an excellent observatory in his own house, where he observed the spots and phases of the moon, from which observations he compiled his *Selenographia*; and an account of his apparatus is contained in his work entitled *Machina Caelestis*, a book now very scarce, as most of the copies were accidentally burnt, with the whole house and apparatus, in 1679. Hevelius died in 1688, aged 76.

Dr. Hook, a contemporary of Hevelius, invented instruments with telescopic sights, and censured those of the latter; which occasioned a sharp dispute between them, and to settle which the celebrated Dr. Halley was sent over to Hevelius to examine his instruments. The two astronomers made several observations together, very much to their satisfaction, and amongst them was one of an occultation of Jupiter by the moon, when they determined the diameter of the latter to be 30' 33".

Before



Before the middle of the 17th century the construction of telescopes had been greatly improved, particularly by Huygens and Fontana. The former constructed one of 123 feet, with which he long observed the moon and planets, and discovered that Saturn was encompassed with a ring. With telescopes too, of 200 and 300 feet focus, Cassini saw five satellites of Saturn, with his zones or belts, and the shadows of Jupiter's satellites passing over his body. In 1666 Azout applied a micrometer to telescopes, to measure the diameters of the planets, and other small distances in the heavens: but an instrument of this kind had been invented before, by Mr. Gascoigne, though it was but little known abroad.

To obviate the difficulties of the great lengths of refracting telescopes and the aberration of the rays, it is said that Merfennus first started the idea of making telescopes of reflectors, instead of lenses, in a letter to Descartes; and in 1663 James Gregory of Aberdeen shewed how such a telescope might be constructed. After some time spent also by Newton, on the construction of both sorts of telescopes, he found out the great inconvenience which arises to refractors from the different refrangibility of the rays of light, for which he could not then find a remedy; and therefore, pursuing the other kind, in the year 1672 he presented to the Royal Society two reflectors, which were constructed with spherical speculums, as he could not procure other figures. The inconveniences however arising from the different refrangibility of the rays of light, have since been fully obviated by the ingenious Mr. Dollond. Towards the latter part of the 17th, and beginning of the 18th century, practical astronomy it seems rather languished. But at the same time the speculative part was carried to the highest perfection by the immortal Newton in his *Principia*, and by the Astronomy of David Gregory.

Soon after this however, great improvements of astronomical instruments began to take place, particularly in Britain. Mr. Graham, a celebrated mechanic and watchmaker, not only improved clocks and watch work, but also carried the accuracy of astronomical instruments to a surprising degree. He constructed the old 8 feet mural arch at the Royal Observatory Greenwich, and a small equatorial sector for making observations out of the meridian; but he is chiefly remarkable for contriving the zenith sector of 24 feet radius, and afterwards one of  $12\frac{1}{2}$  feet, with which Dr. Bradley discovered the aberration of the fixed stars. The reflecting telescope of Gregory and Newton, was greatly improved by Mr. Hadley, who presented a very powerful instrument of that kind to the Royal Society in 1719. The same gentleman has also immortalized his memory by the invention of the reflecting quadrant or sector, now called by his name, which he presented to the society in 1731, and which is now so universally useful at sea, especially where nice observations are required. It appears however that an instrument similar to this in its principles, had been invented by Newton; and a description with a drawing of it given by him to Dr. Halley, when he was preparing for his voyage in 1701, to discover the variation of the needle: it has also been asserted that a Mr. Godfrey of Philadelphia in America, made the same discovery, and the first instrument of this kind. About the middle of this century, the constructing and dividing of large astronomical instruments were carried to great

perfection by Mr. John Bird; and reflecting telescopes were not less improved by Mr. Short, who also first executed the divided object-glass micrometer, which had been proposed and described by M. Louville and others. Mr. Dollond also brought refracting telescopes to the greatest perfection, by means of his acromatic glasses; and lately the discoveries of Herschel are owing to the amazing powers of reflectors of his own construction.

Thus the astronomical improvements in the present century, have been chiefly owing to the foregoing inventions and improvements in the instruments, and to the establishment of regular observatories in England, France, and other parts of Europe. Roemer, a celebrated Danish Astronomer, first made use of a meridional telescope; and, by observing the eclipses of Jupiter's satellites, he first discovered the progressive motion of light, concerning which he read a dissertation before the Academy of Sciences at Paris in 1675.—Mr. Flamsteed was appointed the first Astronomer Royal at Greenwich in 1675. He observed, for 44 years, all the celestial phenomena, the sun, moon, planets, and fixed stars, of all which he gave an improved theory and tables, viz. a catalogue of 3000 stars with their places, to the year 1689; also new solar tables, and a theory of the moon according to Horrox; likewise, in Sir Jonas Moore's System of Mathematics, he gave a curious tract on the doctrine of the sphere, shewing how, geometrically to construct eclipses of the sun and moon, as well as occultations of the fixed stars by the moon. And it was upon his tables that were constructed, both Halley's tables, and Newton's theory of the moon.—Cassini also, the first French Astronomer Royal, very much distinguished himself, making many observations on the sun, moon, planets and comets, and greatly improved the elements of their motions. He also erected the gnomon, and drew the celebrated meridian line in the church of Petronia at Bologna.

In 1719 Mr. Flamsteed was succeeded by Dr. Halley, as Astronomer Royal at Greenwich. The Doctor had been sent at the early age of 21 to the island of St. Helena, to observe the southern stars, and make a catalogue of them, which was published in 1679. In 1705 he published his *Synopsis Astronomice Cometicæ*, in which he ventured to predict the return of a comet in 1758 or 1759. He was the first who discovered the acceleration of the moon, and he gave a very ingenious method for finding her parallax by three observed phases of a solar eclipse. He published, in the Philosophical Transactions, many learned papers, and amongst them some that were concerning the use that might be made of the next transit of Venus in determining the distance of the sun from the earth. He composed tables of the sun, moon, and all the planets, which are still in great repute; with which he compared the observations he made of the moon at Greenwich, amounting to near 1500, and noted the differences. He recommended the method of determining the longitude by the moon's distances from the sun and certain fixed stars; a method which had before been noticed, and which has since been carried into execution, more particularly at the instance of the present Astronomer Royal.

About this time a dispute arose concerning the figure of the earth. Sir Isaac Newton had determined, from a consideration of the laws of gravity, and the diurnal motion of the earth, that the figure of it was an oblate spheroid,



spheroid, and flattened at the poles: but Cassini had determined, from the measures of Picart, that the figure was an oblong spheroid, or lengthened at the poles. To settle this dispute, it was resolved, under Lewis XV, to measure two degrees of the meridian; one near the equator, and the other as near the pole as possible. For this purpose, the Royal Academy of Sciences sent to Lapland, Mefs. Maupertuis, Clairault, Camus, and Le Monier; being also accompanied by the Abbé Outhier, and by M. Celfus, professor of anatomy at Upsal. And on the southern expedition were sent Mefs. Godin, Condamine, and Bouguer, to whom the king of Spain joined Don George Juan and Don Antonio de Ulloa. These set out in 1735, and returned at different times in 1744, 1745, and 1746; but the former party, who set out only in 1736, returned the year following; having both fulfilled their commissions. Picart's measure was also revised by Cassini and De la Caille, which after his errors were corrected, was found to agree very well with the other two; and the result of the whole served to confirm the determination of the figure before laid down by Newton.—On the southern expedition, it was found that the attraction of the great mountains of Peru had a sensible effect on the plumb-line of one of their largest instruments, deflecting it 7 or 8 seconds from the true perpendicular.

On the death of Dr. Halley, in 1742, he was succeeded by Dr. Bradley, as Astronomer Royal at Greenwich. The accuracy of his observations enabled him to detect the smaller inequalities in the motions of the planets and fixed stars. The consequence of this accuracy was, the discovery of the aberration of light, the nutation of the earth's axis, and a much greater degree of perfection in the lunar tables. He also observed the places, and computed the elements of the comets which appeared in the years 1723, 1736, 1743, and 1757. He made new and accurate tables of the motions of Jupiter's satellites; and from a multitude of observations of the luminaries, he constructed the most accurate table of refractions yet extant. Also, with a very large transit instrument, and a new mural quadrant of 8 feet radius, constructed by Mr. Bird in 1750, he made an immense number of observations for settling the places of all the stars in the British catalogue, together with near 1500 places of the moon, the greater part of which he compared with Mayer's tables. Dr. Bradley died in 1762.

In the mean time the mathematicians and astronomers elsewhere were assiduous in their endeavours to promote the science of astronomy. The theory of the moon was particularly considered by Messrs Clairault, D'Alembert, Euler, Meyer, Simpson, and Walmfly, and especially Clairault, Euler, and Mayer, who computed complete sets of lunar tables; those of the last of these authors, for their superior accuracy, were rewarded with a premium of 3000 pounds by the Board of Longitude, who brought them into use in the computation of the Nautical Ephemeris, published by that Board.—The most accurate tables of the satellites of Jupiter were composed, from observations by Mr. Wargentin, an excellent Swedish astronomer.—Among the many French astronomers who contributed to the advancement of the science, it was particularly indebted to M. De la Caille, for an excellent set of solar tables. And in 1750 he went to the Cape of Good Hope to make observations in concert

with the most celebrated astronomers in Europe, for determining the parallax of Mars and the moon, and thence, that of the sun, which it was concluded did not much exceed 10 seconds. Here he re-examined and adjusted, with great accuracy, the stars about the southern pole; and also measured a degree of the meridian.—In Italy the science was assiduously cultivated by Bianchini, Boscovich, Frisi, Manfredi, Zanotti, and many others; in Sweden by Wargentin already mentioned, Blingenstern, Mallet, and Planman; and in Germany by the Eulers, Mayer, Lambert, Grischow, and others.

In the year 1760 all the learned Societies in Europe made preparations for observing the transit of Venus over the sun, which had been predicted by Dr. Halley more than 80 years before, and the use that might be made of it in determining the sun's parallax, and the distances of the planets from the sun. And the same exertions were repeated, to observe the transit in 1769, by sending observers to different parts of the world, for the more convenience in observing. And from the whole, Mr. Short computed that the sun's parallax was nearly  $8\frac{1}{2}$  seconds, and consequently the distance of the sun from the earth about 24114 of the earth's diameters, or 96 millions of miles.

Dr. Bradley was succeeded, in 1762, in his office of Astronomer Royal, by Mr. Bliss, Savilian professor of astronomy; who being in a declining state of health, did not long enjoy it. But, dying in 1765, was succeeded by the learned Nevil Maskelyne, D. D. the present Astronomer Royal, who has discharged the duties of that office with the greatest honour to himself, and benefit to the science. In January 1761 this gentleman was sent by the Royal Society, at a very early age, to the Island of St. Helena, to observe the transit of Venus over the sun, and the parallax of the star Sirius. The first of these objects partly failed, by clouds preventing the sight of the 2d internal contact; and the 2d also, owing to Mr. Short having suspended the plumbline by a loop from the neck of the central pin. However, our astronomer indemnified himself by many other valuable observations: Thus, he observed at St. Helena, the tides; the horary parallaxes of the moon; and the going of a clock, to find, by comparison with its previous going which had been observed in England, the difference of gravity at the two places: also, in going out and returning, he practised the method of finding the longitude by the lunar distances taken with a Hadley's Quadrant, making out rules for the use of seamen, and taught the method to the officers on board the ship; which he afterwards explained in a letter to the Secretary of the Royal Society, inserted in the Philos. Transf. for the year 1762, and still more fully afterwards, in the British Mariner's Guide, which he published in the year 1763. He returned from St. Helena in the spring of 1762, after a stay there of 10 months; and in September 1763 sailed for the island of Barbadoes, to settle the longitude of the place, and to compare Mr. Harrison's watch with the time there when this gentleman should bring it out: another object was also to try Mr. Irwin's marine chair, which he did in his way out. While at Barbadoes, he also made many other observations, and amongst them, many relating to the moon's horary parallaxes, not yet published. Returning to England in the latter part of the year 1764, he was



appointed in 1765 to succeed Mr. Bliss as Astronomer Royal, and immediately recommended to the Board of Longitude the lunar method of finding the longitude, and proposed to them the project of a Nautical Almanac, to be calculated and published to facilitate that method; this they agreed to, and the first vol. was published for 1767, and it has continued ever since under his direction, to the great benefit of navigation and universal commerce.

A multitude of other useful writings by this gentleman are inserted in the volumes of the Philos. Trans. and particularly in consequence of a proposal, made by him to the Royal Society, the noble project was formed of measuring very accurately the effect of some mountain on the plumb line, in deflecting it from the perpendicular; and the mountain of Schhallien, in Scotland, having been found the most convenient in this island for the purpose, at the request of the Society, he went into Scotland to conduct the business, which he performed in the most accurate manner, shewing that the sum of the deflections on the two opposite sides was about  $11\frac{1}{2}$  seconds, of a degree; and proving, to the satisfaction of the whole world, the universal attraction of all matter. From the data resulting from these measures I have computed the mean density of the whole matter in the earth, which I have found to be about  $4\frac{1}{2}$  times that of common water. Besides many learned and valuable papers in the Philosophical Transactions, and the most assiduous exertions in the duties of the observatory, as abundantly appears by the curious and voluminous observations which he has published, the world is particularly obliged to his endeavours with the Board of Longitude, for the publication of the Nautical Ephemeris, and the method of observing the longitude, by the distances of the moon and stars, now adopted by all nations, and by which the practice of Navigation has been brought to the greatest perfection.

Finally, the discoveries of Dr. Herschel form a new æra in astronomy. He first, in 1781, began with observations on the periodical star in *Collo Ceti*, and a new method of measuring the lunar mountains, none of which he made more than half a mile in height: and, having constructed telescopes vastly more powerful than any former ones, he proceeded to other observations, concerning which he has had several papers printed in the Philosophical Transactions; as, on the rotation of the planets round their axes; On the parallax of the fixed stars; Catalogues of double, triple, &c stars; On the proper motion of the sun and solar system; On the remarkable appearances of the solar regions of the planet Mars, &c. And, above all, his discovery of a new primary planet, on the 13th of March 1781, which he calls the *Georgian Planet*, but it is named the *Planet Herschel* by the French and other foreign astronomers; by which, and its two satellites, which he has also discovered since that time, he has greatly enlarged the bounds of the solar system, this new planet being more than twice as far from the sun as the planet Saturn.

Lists and historical accounts of the principal writings and authors on astronomy are contained in Weidler's History of Astronomy, which is brought down to the year 1737. There is also Bailly's History of astronomy, ancient and modern. For this purpose, consult also the following authors, viz, Adam, Vossius, Bayle, Chauffepié, Nicéron, Perraut, the chronological table of

Riccioli, and that of Sherburn, at the end of his edition of Manilius; also the first volume of De la Lande's astronomy. The more modern, and popular books on astronomy, are very numerous, and well known: as those of Ferguson, Long, Emerson, Vince, De la Lande, Leadbetter, Brent, Keil, Whiston, Wing, Street, &c, &c.

**ASTROSCOPE**, a kind of astronomical instrument, composed of two cones, on the surface of which are delineated the constellations, with their stars, by means of which these may easily be known in the heavens. The astroscope was invented by William Shukhard, professor of mathematics at Tubingen, upon which he published a treatise in 1698.

**ASTROSCOPIA**, the art of observing and examining the stars, by means of telescopes, to discover their nature and properties.

**ASTROTHEMATA**, the places or positions of the stars, in an astrological scheme of the heavens.

**ASTROTHESIA**, is used by some for a constellation or collection of stars in the heavens.


**ASTRUM**, or **ASTRON**, a constellation, or assemblage of stars: in which sense it is distinguished from *Aster*, which denotes a single star. Some apply the term, in a more particular sense, to the Great Dog, or rather to the large bright star in his mouth.

**ASYMMETRY**, the want of proportion, otherwise called *incommensurability*, or the relation of two quantities which have no common measure, as between 1 and  $\sqrt{2}$ , or the side and diagonal of a square.

**ASYMPTOTE**, is properly a right line, which approaches continually nearer and nearer to some curve, whose asymptote it is said to be, in such sort, that when they are both indefinitely produced, they are nearer together than by any assignable finite distance; or it may be considered as a tangent to the curve when conceived to be produced to an infinite distance. Two curves are also said to be asymptotical, when they thus continually approach indefinitely to a coincidence: thus, two parabolas, placed with their axes in the same right line, are asymptotes to one another.



Of lines of the second kind, or curves of the first kind, that is the conic sections, only the hyperbola has asymptotes, which are two in number. All curves of the second kind have at least one asymptote; but they may have three. And all curves of the third kind may have four asymptotes. The conchoid, cissoid, and logarithmic curve, though not reputed geometrical curves, have each one asymptote. And the branch or leg of a curve that has an asymptote, is said to be of the hyperbolic kind.

The nature of asymptotes will be easily conceived from the instance of the asymptote to the conchoid. Thus, if, ABC &c be part of a conchoid, and the line MN be so drawn that the parts FB, GC, HD, IE, &c, of right lines, drawn from the pole P, be equal to each other; then will the line MN be the asymptote of the curve: because the perpendicular Cc is shorter than FB, and Dd than Cc, &c; so that the two lines continually approach; yet the points Ee &c can never coincide.





ASYMPTOTES of the HYPERBOLA are thus described. Suppose a right line DE drawn to touch the curve in any point A, and equal to the conjugate *de* of the diamete ACB drawn to that point A, viz, AD or AE equal to the semiconjugate Cd or Ce; then the two lines CDF, CEH, drawn from the centre C, through the points D and E, are the two asymptotes of the curve.



The parts of any right line, lying between the curve of the common hyperbola and its asymptotes, are equal to one another on both sides, that is,  $gG = bH$ . In like manner, in hyperbolas of the second kind, if there be drawn any right line cutting both the curve and its three asymptotes in three points, the sum of the two parts of that right line extended in the same direction from any two of the asymptotes to two points of the curve, is equal to the third part which extends in the contrary direction from the third asymptote to the third point of the curve.

If AGK be an hyperbola of any kind, whose nature, with regard to the curve and asymptote, is expressed by this general equation  $x^m y^n = a^{m+n}$ , where  $x$  is  $= CF$ , and  $y = FG$  drawn any where parallel to the other asymptote CH; and the parallelogram CFGI be completed: Then  $m - n$  is to  $n$ , as this parallelogram CFGI is to the hyperbolic space FGK, contained under the determinate line FG, with the asymptote FK and the curve GK, both indefinitely continued towards K. So that, if  $m$  be greater than  $n$ , the said asymptotic space is finite and quadrable: but when  $m = n$ , as in the common or conic hyperbola, then  $m - n = 0$ , the ratio of that space to the said parallelogram, is as  $n$  to 0: that is, the hyperbolic space is infinitely great, in respect of the finite parallelogram: and when  $m$  is less than  $n$ , then,  $m - n$  being negative, the asymptotic space is to the determinate parallelogram, as a positive number is to a negative one, and is what Dr. Wallis calls more than infinite.

ASYMPTOTE of the LOGARITHMIC CURVE. If LMN be the logarithmic curve, QON an asymptote, LQ and MP ordinates, MO a tangent, and PO the subtangent, which in this curve is a constant quantity. Then the indeterminate space LMNQ is equal to  $LQ \times PO$ , the rectangle under the ordinate LQ and the constant subtangent PO; and the solid generated by the rotation of that curve space about the asymptote NQ, is equal to half the cylinder, whose altitude is the said constant subtangent PO, and the radius of its base is LQ.

ASYMPTOTES, by some, are distinguished into various orders. The asymptote is said to be of the first order, when it coincides with the base of the curvilinear figure: of the 2d order, when it is a right line parallel to the base: of the 3d order, when it is a right line ob-

lique to the base: of the 4th order, when it is the common parabola, having its axis perpendicular to the base: and, in general, of the  $n + 2$  order, when it is a parabola whose ordinate is always as the  $n$  power of the base. The asymptote is oblique to the base, when the ratio of the first fluxion of the ordinate to the fluxion of the base, approaches to an assignable ratio, as its limit; but it is parallel to the base, or coincides with it, when this limit is not assignable.

The doctrine and determination of the asymptotes of curves, is a curious part of the higher geometry. Fontenelle has given several theorems relating to this subject, in his *Geometrie de l'Infini*. See also Stirling's *Lineæ Tertii Ordinis*, prop. vi, where the subject of Asymptotes is learnedly treated; and Cramer's *Introduction à l'analyse des lignes courbes*, art. 147 & seq. for an excellent theory of asymptotes of geometrical curves and their branches. Likewise Maclaurin's *Algebra*, and his *Fluxions*, book i, chap. 10, where he has carefully avoided the modern paradoxes concerning infinites and infinitesimals. But the easiest way of determining asymptotes, it seems, is by considering them as tangents to the curves at an infinite distance from the beginning of the absciss, that is when the absciss  $x$  is infinite in the equation of the curve, and in the proportion of  $x$  to  $y$ , or in that of the subtangent to the ordinate.

The areas bounded by curves and their asymptotes, though indefinitely extended, have sometimes limits to which they may approach indefinitely near: and this happens in hyperbolas of all kinds, except the first or Apollonian, and in the logarithmic curve; as was observed above. But in the common hyperbola, and many other curves, the asymptotical area has no such limit, but is infinitely great.—Solids, too, generated by hyperbolic areas, revolving about their asymptotes, have sometimes their limits; and sometimes they may be produced till they exceed any given solid.—Also the surface of such solid, when supposed to be infinitely produced, is either finite or infinite, according as the area of the generating figure is finite or infinite.

ATLANTIDES, a name given to the Pleiades, or seven stars, sometimes also called Vergiliæ. They were thus called, as being supposed by the poets to have been the daughters either of Atlas or his brother Hesperus, who were translated to heaven.

ATMOSPHERE, a term used to signify the whole of the fluid mass, consisting of air, aqueous and other vapours, electric fluids, &c, which surrounds the earth to a considerable height, and partaking of all its motions, both annual and diurnal.

The composition of that part of our atmosphere properly called air, was till lately but very little known. Formerly it was supposed to be a simple, homogeneous, and elementary fluid. But the experiments of Dr. Priestley and others, have discovered, that even the purest kind of air, which they call dephlogisticated, is in reality a compound, and might be artificially produced in various ways. This dephlogisticated air, however, is but a small part of the composition of our atmosphere. By accurate experiments, the air we usually breathe, is composed of only one-fourth part of this dephlogisticated air, or perhaps less; the other three parts, or more, consisting of what Dr. Priestley calls *phlogisticated*, and M. Lavoisier *nephitic air*.



Beside these sorts of air, it is obvious that the whole mass of the atmosphere contains a great deal of water, together with a vast heterogeneous collection of particles raised from all bodies of matter on the surface of the earth, by effluvia, exhalations, &c; so that it may be considered as a chaos of the particles of all sorts of matter confusedly mingled together. And hence the atmosphere has been considered as a large chemical vessel, in which the matter of all kinds of sublunary bodies is copiously floating; and thus exposed to the continual action of that immense surface the sun; from whence proceed innumerable operations, sublimations, separations, compositions, digestions, fermentations, putrefractions, &c.

There is, however, one substance, namely the electrical fluid, which is very distinguishable in the mass of the atmosphere. To measure the absolute quantity of this fluid, either in the atmosphere or any other substance, is perhaps impossible: and all that we know on this subject is, that the electric fluid pervades the atmosphere; that it appears to be more abundant in the superior than the inferior regions; that it seems to be the immediate bond of connection between the atmosphere and the water which is suspended in it; and that by its various operations, the phenomena of hail, rain, snow, lightning, and the other kinds of meteors are occasioned. See those respective articles; and see also Beccaria's Essay on Atmospheric Electricity, annexed to the English translation of his Artificial Electricity.

*Uses of the ATMOSPHERE.*—The uses of the atmosphere are so many and great, that it seems indeed absolutely necessary, not only to the comfort and convenience of men, but even to the existence of all animal and vegetable life, and to the very constitution of all kinds of matter whatever, and without which they would not be what they are: for by it we live, breathe, and have our being; and by insinuating itself into all the vacuities of bodies, it becomes the great spring of most of the mutations here below; as generation, corruption, dissolution, &c; and without which none of these operations could be carried on. Without the atmosphere, no animal could exist, or indeed be produced; neither any plant, all vegetation ceasing without its aid; there would be neither rain nor dews to moisten the face of the ground; and though we might perceive the sun and stars like bright specks, we should be in utter darkness, having none of what we call day light or even twilight: nor would either fire or heat exist without it. In short, the nature and constitution of all matter would be changed and cease; wanting this universal bond and constituting principle.

By the mechanical force of the atmosphere too, as well as its chemical virtues, many necessary purposes are answered. We employ it as a moving power, in the motion of ships, to turn mills, and for other such uses. And it is one of the great discoveries of the modern philosophers, that the several motions attributed by the ancients to a *fuga vacui*, are really owing to the pressure of the atmosphere. Galileo, having observed that there was a certain standard altitude, beyond which no water could be elevated by pumping, took an occasion from thence to call in question the doctrine of the schools, which ascribed the ascent of water in pumps, to the

*fuga vacui*, and instead of it he happily substituted the hypothesis of the weight and pressure of the air. It was with him, indeed, little better than an hypothesis, since it had not then those confirmations from experiment, afterwards found out by his pupil Torricelli, and other succeeding philosophers, particularly Mr Boyle.

Nor have the attempts to fly or float through the air been altogether without success. F. De Lana thought he had contrived an aeronautic machine for navigating the atmosphere: and Sturmius, who examined it, asserted that it was not impracticable: though Dr Hook was of a different opinion, and detected the fallacy of the contrivance. Roger Bacon, long before, proposed something of the same kind. The great secret of this art is to contrive a machine so much lighter than the air, that it will rise up and float in the atmosphere, and together with itself, buoy up and carry men along with it. The principle on which it is to be effected is, by means of an air pump, to exhaust the air from a very thin and light, yet firm, metalline vessel. But the hopes of success in such an enterprize will appear very small if it be considered, that if a globe were formed of brass of the thickness only of  $\frac{1}{12}$  of an inch, such a globe would require to be about 277 feet in diameter to float in the air: and if, as De Lana supposes, the diameter of the globe were but 25 feet, the thickness of the metal could not exceed  $\frac{1}{3}$  of an inch. See Herman's *Phoronomia* pa. 158. However, what is not to be expected from metalline globes or shells, has now been successfully accomplished by the balloons of cloth, silk, or skin, of Montgolfier and others. See the article *Aerostation*, &c.

*Salubrity of the ATMOSPHERE.* On the tops of mountains the air is generally more salubrious than in pits or very deep places. Indeed dense air is always more proper for respiration, as to the mere quality of density only, than that which is rarer. But then the air on mountains, though rarer, is freer from phlogistic vapours than that of pits; and hence it has been found that people can live very well on the tops of mountains, even when the air is but about half the density of that below. But it would seem that at some intermediate height between the two extremes, the air is the most salubrious and proper for animal life; and this height, according to M. de Saussure, is about 500 or 600 yards above the level of the sea.

Besides the difference arising from the mere difference of altitude, the salubrity of the atmosphere is greatly affected by many other circumstances. The air, when confined or stagnant, is commonly more impure than when agitated and shifted: thus, all close places are unhealthy, and even the air in a bed chamber is less salubrious in a morning, after it has been slept in, than in the evening. Dr White, in vol. 68 *Philos. Trans.* gives an account of experiments on this quality of the air, and remarks one instance when the air was particularly impure, viz September 13, 1777; when the barometer stood at 30.35, the thermometer at 69°; the air being then dry and sultry, and no rain having fallen for more than two weeks. A slight shock of an earthquake was perceived that day. In vol. 70 of the same *Transactions* Dr. Ingenhousz gives an account of some experiments on this head, made in various places and situations:



situations: he finds," That the air at sea, and close to it, is in general purer, and fitter for animal life, than the air on the land:" but the Doctor did not find much difference between the air of the towns and of the country, nor between one town and another. The Abbé Fontana, made nearly the same conclusions, from accurate experiments, asserting, "that the difference between the air of one country and that of another, at different times, is much less than what is commonly believed; and yet that this difference in the purity of the air at different times, is much greater than the difference between the air of the different places observed by him." Finally M. Fontana concludes, that "Nature is not so partial as we commonly believe. She has not only given us an air almost equally good every where at every time, but has allowed us a certain latitude, or a power of living and being in health in qualities of air which differ to a certain degree. By this I do not mean to deny the existence of certain kinds of noxious air in some particular places; but only say, that in general the air is good every where, and that the small differences are not to be feared so much as some people would make us believe. Nor do I mean to speak here of those vapours and other bodies which are accidentally joined to the common air in particular places, but do not change its nature and intrinsic property. This state of the air cannot be known by the test of nitrous air; and those vapours are to be considered in the same manner as we should consider so many particles of arsenic swimming in the atmosphere. In this case it is the arsenic, and not the degenerated air, that would kill the animals who ventured to breathe it."

*Figure of the ATMOSPHERE.*—As the atmosphere envelops all parts of the surface of our globe, if they both continued at rest, and were not endowed with a diurnal motion about their common axis, then the atmosphere would be exactly globular, according to the laws of gravity; for all the parts of the surface of a fluid in a state of rest, must be equally removed from its centre. But as the earth and the ambient parts of the atmosphere revolve uniformly together about their axis, the different parts of both have a centrifugal force, the tendency of which is more considerable, and that of the centripetal less, as the parts are more remote from the axis; and hence the figure of the atmosphere must become an oblate spheroid; since the parts that correspond to the equator are farther removed from the axis, than the parts which correspond to the poles. Besides, the figure of the atmosphere must, on another account, represent a flattened spheroid, namely because the sun strikes more directly the air which encompasses the equator, and is comprehended between the two tropics, than that which pertains to the polar regions: for, from hence it follows, that the mass of air, or part of the atmosphere, adjoining to the poles, being less heated, cannot expand so much, nor reach so high. And yet, notwithstanding, as the same force which contributes to elevate the air, diminishes its gravity and pressure on the surface of the earth, higher columns of it about the equatorial parts, all other circumstances being the same, may not be heavier than those about the poles.

In the Transactions of the Royal Irish Academy for 1788 Mr Kirwin has an ingenious dissertation on the

figure, height, weight, &c, of the atmosphere. He observes that, in the natural state of the atmosphere, that is, when the barometer would every where, at the level of the sea, stand at 30 inches, the weight of the atmosphere, at the surface of the sea, must be equal all over the globe; and in order to produce this equality, as the weight proceeds from its density and height, it must be lowest where the density is greatest, and highest where the density is least; that is, highest at the equator and lowest at the poles, with several intermediate gradations.

Though the equatorial air however be less dense to a certain height than the polar, yet at some greater heights it must be more dense: for since an equatorial and polar column are equal in total weight or mass, the lower part of the equatorial column, being more expanded by heat &c than that of the polar, must have less mass, and therefore a proportionably greater part of its mass must be found in its superior section; so that the lower extremity of the superior section of the equatorial column is more compressed, and consequently denser, than the corresponding part of the polar column. The same thing is to be understood also of the extra-tropical columns with respect to each other, where differences of heat prevail.

Hence, in the highest regions of the atmosphere, the denser equatorial air, not being supported by the collateral extra-tropical columns, gradually flows over, and rolls down to the north and south.

These superior tides consist chiefly of inflammable air, as it is much lighter than any other, and is generated in great plenty between the tropics; it furnishes the matter of the auroræ borealis and australis, by whose combustion it is destroyed, else its quantity would in time become too great, and the weight of the atmosphere annually increased; but its combustion is the primary source of the greatest perturbations of the atmosphere.

*Weight or Pressure of the ATMOSPHERE.*—It is evident that the mass of the atmosphere, in common with all other matter, must be endowed with weight and pressure; and this principle was asserted by almost all philosophers, both ancient and modern. But it was only by means of the experiments made with pumps and the barometrical tube, by Galileo and Torricelli, that we came to the proof, not only that the atmosphere is endued with a pressure, but also what the measure and quantity of that pressure is. Thus, it is found that the pressure of the atmosphere sustains a column of quicksilver, in the tube of the barometer, of about 30 inches in height; it therefore follows, that the whole pressure of the atmosphere is equal to the weight of a column of quicksilver, of an equal base, and 30 inches height: and because a cubical inch of quicksilver is found to weigh nearly half a pound avoirdupois, therefore the whole 30 inches, or the weight of the atmosphere on every square inch of surface, is equal to 15 pounds. Again, it has been found that the pressure of the atmosphere balances, in the case of pumps &c, a column of water of about  $34\frac{1}{2}$  feet high; and, the cubical foot of water weighing just 1000 ounces, or  $62\frac{1}{2}$  pounds,  $34\frac{1}{2}$  times  $62\frac{1}{2}$ , or 2158lb, will be the weight of the column of water, or of the atmosphere on a base of a square foot; and consequently the 144th part of this, or 15lb, is the weight of the atmosphere on a square inch;



inch; the same as before. Hence Mr Cotes computed that the pressure of this ambient fluid on the whole surface of the earth, is equivalent to that of a globe of lead of 60 miles in diameter. And hence also it appears, that the pressure upon the human body must be very considerable; for as every square inch of surface sustains a pressure of 15 pounds, every square foot will sustain 144 times as much, or 2160 pounds; then, if the whole surface of a man's body he supposed to contain 15 square feet, which is pretty near the truth, he must sustain 15 times 2160, or 32400 pounds, that is nearly  $14\frac{1}{2}$  tons weight, for his ordinary load. By this enormous pressure we should undoubtedly be crushed in a moment, if all parts of our bodies were not filled either with air or some other elastic fluid, the spring of which is just sufficient to counterbalance the weight of the atmosphere. But whatever this fluid may be, it is certain that it is just able to counteract the weight of the atmosphere, and no more: for, if any considerable pressure be superadded to that of the air, as by going into deep water, or the like, it is always severely felt let it be ever so equable, at least when the change is made suddenly; and if, on the other hand, the pressure of the atmosphere be taken off from any part of the human body, as the hand for instance, when put over an open receiver, from whence the air is afterwards extracted, the weight of the external atmosphere then prevails, and we imagine the hand strongly sucked down into the glass.

The difference in the weight of the air which our bodies sustain at one time more than another, is also very considerable, from the natural changes in the state of the atmosphere. This change takes place chiefly in countries at some distance from the equator; and as the barometer varies at times from 28 to 31 inches, or about one tenth of the whole quantity, it follows that this difference amounts to about a ton and a half on the whole body of a man, which he therefore sustains at one time more than at another. On the increase of this natural weight, the weather is commonly fine, and we feel ourselves what we call braced and more alert and active; but, on the contrary, when the weight of the air diminishes, the weather is bad, and people feel a listlessness and inactivity about them. And hence it is no wonder that persons suffer very much in their health, from such changes in the atmosphere, especially when they take place very suddenly, for it is to this circumstance chiefly that a sensation of uneasiness and indisposition is to be attributed; thus, when the variations of the barometer and atmosphere are sudden and great, we feel the alteration and effect on our bodies and spirits very much; but when the change takes place by very slow degrees, and by a long continuance, we are scarcely sensible of it, owing, undoubtedly, to the power with which the body is naturally endowed, of accommodating itself to this change in the state of the air, as well as to the change of many other circumstances of life, the body requiring a certain interval of time to effect the alteration in its state, proper to that of the air &c. Thus, in going up to the tops of mountains, where the pressure of the atmosphere is diminished two or three times more than on the plain below, little or no inconvenience is felt from the rarity of the air, if it is not mixed with other noxious vapours &c; because

that, in the ascent the body has had sufficient time to accommodate itself gradually to the slow variation in the state of the atmosphere: but, when a person ascends with a balloon, very rapidly to a great height in the atmosphere, he feels a difficulty in breathing and an uneasiness of body; and the same is soon felt by an animal when inclosed in a receiver, and the air suddenly drawn or pumped out of it. So also, on the condensation of the air, we feel little or no alteration in ourselves, except when the change happens suddenly, as in very rapid changes in the weather, and in descending to great depths in a diving bell, &c. I have often heard the late unfortunate Mr. Spalding speak of his experience on this point: he always found it absolutely necessary to descend with the bell very slowly, and that only from one depth to another, resting a while at each depth before he began to descend farther: he first descended slowly for about 5 or 6 fathom, and then stopped a while; he felt an uneasiness in his head and ears, which increased more and more as he descended, till he was obliged to stop at the depth above mentioned, where the density of the air was nearly doubled; having remained there a while, he felt his ears give a sudden crack, and after that he was soon relieved from any uneasiness in that part, and it seemed as if the density of the air was not altered. He then descended other 5 fathoms or 30 feet more, with the same precaution and the same sensations as before, being again relieved, in the same manner, after remaining awhile stationary at the next stage of his descent, where the density of the air was tripled. And thus he continued proceeding to a great depth, always with the same circumstances, repeated at every 5 or 6 fathoms, and adding the pressure of one more atmosphere at every period of the progress.

It is not easy to assign the true reason for the variations that happen in the gravity of the atmosphere in the same place. One cause of it however, either immediate or otherwise, it seems, is the heat of the sun; for where this is uniform, the changes are small and regular; thus between the tropics it seems the change depends on the heat of the sun, as the barometer constantly sinks about half an inch every day, and rises again to its former station in the night time. But in the temperate zones the barometer ranges from 28 to near 31 inches, shewing, by its various altitudes, the changes that are about to take place in the weather. If we could know therefore, the causes by which the weather is influenced; we should also know those by which the gravity of the atmosphere is affected. These may perhaps be reduced to immediate ones, viz, an emission of latent heat from the vapour contained in the atmosphere, or of electric fluid from the same, or from the earth; as it is observed that they both produce the same effect with the solar heat in the tropical climates, viz, to rarefy the air, by mixing with it, or setting loose a lighter fluid, which did not before act in such large proportion in any particular place.

With regard to the alteration of heat and cold in the atmosphere, many reasons and hypotheses have been given, and many experiments made; as may be seen by consulting the authors upon this subject, viz, M. Bouguer's observations in Peru, Lambert, De Luc, Saussure's journeys on the Alps, Sex's and Darwin's experiments



experiments in vol. 78 *Philos. Transf.* This last gentleman hence infers, "There is good reason to conclude that in all circumstances where air is mechanically expanded, it becomes capable of attracting the fluid matter of heat from other bodies in contact with it. Now, as the vast region of air which surrounds our globe is perpetually moving along its surface, climbing up the sides of mountains, and descending into the valleys; as it passes along it must be perpetually varying the degree of heat according to the elevation of the country it traverses: for, in rising to the summits of mountains, it becomes expanded, having so much of the pressure of the superincumbent atmosphere taken away; and when thus expanded, it attracts or absorbs heat from the mountains in contiguity with it; and, when it descends into the valleys and is compressed into less compass, it again gives out the heat it has acquired to the bodies it comes in contact with. The same thing must happen to the higher regions of the atmosphere, which are regions of perpetual frost, as has lately been discovered by the aerial navigators. When large districts of air, from the lower parts of the atmosphere, are raised two or three miles high, they become so much expanded by the great diminution of the pressure over them, and thence become so cold, that hail or snow is produced by the precipitation of the vapour: and as there is, in these high regions of the atmosphere, nothing else for the expanded air to acquire heat from after it has parted with its vapour, the same degree of cold continues till the air, on descending to the earth, acquires its former state of condensation and of warmth. The Andes, almost under the line, rests its base on burning sands: about its middle height is a most pleasant and temperate climate covering an extensive plain, on which is built the city of Quito; while its forehead is encircled with eternal snow, perhaps coeval with the mountain. Yet, according to the accounts of Don Ulloa, these three discordant climates seldom encroach much on each other's territories. The hot winds below, if they ascend, become cooled by their expansion; and hence they cannot affect the snow upon the summit; and the cold winds that sweep the summit, become condensed as they descend, and of temperate warmth before they reach the fertile plains of Quito."

*Height and Density of the ATMOSPHERE.* Various attempts have been made to ascertain the height to which the atmosphere is extended all round the earth. These commenced soon after it was discovered by means of the Torricellian tube, that air is endued with weight and pressure. And had not the air an elastic power, but were it every where of the same density, from the surface of the earth to the extreme limit of the atmosphere, like water, which is equally dense at all depths it would be a very easy matter to determine its height from its density and the column of mercury which it would counterbalance in the barometer tube: for, it having been observed that the weight of the atmosphere is equivalent to a column of 30 inches or  $2\frac{1}{2}$  feet of quicksilver, and the density of the former to that of the latter, as 1 to 11040; therefore the height of the uniform atmosphere would be 11040 times  $2\frac{1}{2}$  feet, that is 27600 feet, or little more than 5 miles and a quarter. But the air, by its elastic quality, expands and contracts; and it being found by repeated experiments in

most nations of Europe, that the spaces it occupies, when compressed by different weights, are reciprocally proportional to those weights themselves; or, that the more the air is pressed, so much the less space it takes up; it follows that the air in the upper regions of the atmosphere must grow continually more and more rare, as it ascends higher; and indeed that, according to that law, it must necessarily be extended to an indefinite height. Now, if we suppose the height of the whole divided into innumerable equal parts; the quantity of each part will be as its density; and the weight of the whole incumbent atmosphere being also as its density; it follows, that the weight of the incumbent air, is every where as the quantity contained in the subjacent part; which causes a difference between the weights of each two contiguous parts of air. But, by a theorem in arithmetic, when a magnitude is continually diminished by the like part of itself, and the remainders the same, these will be a series of continued quantities decreasing in geometrical progression: therefore if, according to the supposition, the altitude of the air, by the addition of new parts into which it is divided, do continually increase in arithmetical progression, its density will be diminished, or, which is the same thing, its gravity decreased, in continued geometrical proportion. And hence, again, it appears that, according to the hypothesis of the density being always proportional to the compressing force, the height of the atmosphere must necessarily be extended indefinitely. And, farther, as an arithmetical series adapted to a geometrical one, is analogous to the logarithms of the said geometrical one; it follows therefore that the altitudes are proportional to the logarithms of the densities, or weights of air; and that any height taken from the earth's surface, which is the difference of two altitudes to the top of the atmosphere, is proportional to the difference of the logarithms of the two densities there, or to the logarithm of the ratio of those densities, or their corresponding compressing forces, as measured by the two heights of the barometer there. This law was first observed and demonstrated by Dr. Halley, from the nature of the hyperbola; and afterwards by Dr. Gregory, by means of the logarithmic curve. See *Philos. Transf.* No. 181, or *Abridg.* vol. 2, p. 13, and *Greg. Astron.* lib. v, prop. 3.

It is now easy, from the foregoing property, and two or three experiments, or barometrical observations, made at known altitudes, to deduce a general rule to determine the absolute height answering to any density, or the density answering to any given altitude above the earth. And accordingly, calculations were made upon this plan by many philosophers, particularly by the French; but it having been found that the barometrical observations did not correspond with the altitudes as measured in a geometrical manner, it was suspected that the upper parts of the atmospheric regions were not subject to the same laws with the lower ones, in regard to the density and elasticity. And indeed, when it is considered that the atmosphere is a heterogeneous mass of particles of all sorts of matter, some elastic, and others not, it is not improbable but this may be the case, at least in the regions very high in the atmosphere, which it is likely may more copiously abound with the electrical fluid. Be this however as it may, it has lately been discovered that the law

above



above given, holds very well for all such altitudes as are within our reach, or as far as to the tops of the highest mountains on the earth, when a correction is made for the difference of the heat or temperature of the air only, as was fully evinced by M. De Luc, in a long series of observations, in which he determined the altitudes of hills both by the barometer, and by geometrical measurement, from which he deduced a practical rule to allow for the difference of temperature. See his *Treatise on the Modifications of the Atmosphere*. Similar rules have also been deduced from accurate experiments, by Sir George Shuckburgh and General Roy, both concurring to shew, that such a rule for the altitudes and densities, holds true for all heights that are accessible to us, when the elasticity of the air is corrected on account of its density: and the result of their experiments shewed, that the difference of the logarithms of the heights of the mercury in the barometer, at two stations, when multiplied by 10000, is equal to the altitude in English fathoms, of the one place above the other; that is, when the temperature of the air is about 31 or 32 degrees of Fahrenheit's thermometer; and a certain quantity more or less, according as the actual temperature is different from that degree.

But it may here be shewn, that the same rule may be deduced independent of such a train of experiments as those above, merely by the density of the air at the surface of the earth alone. Thus, let  $D$  denote the density of the air at one place, and  $d$  the density at the other; both measured by the column of mercury in the barometrical tube: then the difference of altitude between the two places, will be proportional to the log.

of  $D$  — the log. of  $d$ , or to the log. of  $\frac{D}{d}$ . But as this formula expresses only the relation between different altitudes, and not the absolute quantity of them, assume some indeterminate, but constant quantity  $b$ , which multiplying the expression  $\log. \frac{D}{d}$ , may be equal to the real difference of altitude  $a$ , that is,  $a = b \times \log. \frac{D}{d}$ . Then, to determine the value of the general

quantity  $b$ , let us take a case in which we know the altitude  $a$  which corresponds to a known density  $d$ ; as for instance, taking  $a = 1$  foot, or 1 inch, or some such small altitude: then because the density  $D$  may be measured by the pressure of the whole atmosphere, or the uniform column of 27600 feet, when the temperature is 55°; therefore 27600 feet will denote the density  $D$  at the lower place, and 27599 the less density  $d$  at 1 foot above it; consequently  $1 = b \times \log. \frac{27600}{27599}$ , which, by the nature of logarithms, is nearly

$$= b \times \frac{.43429448}{27600} \text{ or } \frac{1}{63551} \text{ nearly; and hence we}$$

find  $b = 63551$  feet; which gives us this formula for any altitude  $a$  in general, viz,  $a = 63551 \times \log. \frac{D}{d}$ ,

$$\text{or } a = 63551 \times \log. \text{ of } \frac{M}{m} \text{ feet, or } 10592 \times \log. \text{ of}$$

$\frac{M}{m}$  fathoms; where  $M$  denotes the column of mercury

in the tube at the lower place, and  $m$  that at the upper. This formula is adapted to the mean temperature of the air 55°: but it has been found, by the experiments of Sir Geo. Shuckburgh and General Roy, that for every degree of the thermometer, different from 55°, the altitude  $a$  will vary by its 435th part; hence, if we would change the factor  $b$  from 10592 to 10000, because the difference 592 is the 18th part of the whole factor 10592, and because 18 is the 24th part of 435; therefore the change of temperature, answering to the change of the factor  $b$ , is 24°, which reduces the 55° to 31°. So that,  $a = 10000 \times \log. \text{ of } \frac{M}{m}$  fathoms, is

the easiest expression for the altitude, and answers to the temperature of 31°, or very nearly the freezing point: and for every degree above that, the result must be increased by so many times its 435th part, and diminished when below it.

From this theorem it follows, that, at the height of  $3\frac{1}{2}$  miles, the density of the atmosphere is nearly 2 times rarer than it is at the surface of the earth; at the height of 7 miles, 4 times rarer; and so on, according to the following table:

Height in miles.	Number of times rarer,
$3\frac{1}{2}$	2
7	4
14	16
21	64
28	256
35	1024
42	4096
49	16384
56	65536
63	262144
70	1048576

And, by pursuing the calculations in this table, it might be easily shewn, that a cubic inch of the air we breathe would be so much rarefied at the height of 500 miles, that it would fill a sphere equal in diameter to the orbit of Saturn.

Hence we may perceive how very soon the air becomes so extremely rare and light, as to be utterly imperceptible to all experience; and that hence, if all the planets have such atmospheres as our earth, they will, at the distances of the planets from one another, be so extremely attenuated, as to give no sensible resistance to the planets in their motion round the sun for many, perhaps hundreds or thousands of ages to come. Even at the height of about 50 miles, it is so rare as to have no sensible effect on the rays of light; for it was found by Kepler, and De la Hire after him, who computed the height of the sensible atmosphere from the duration of twilight, and from the magnitude of the terrestrial shadow in lunar eclipses, that the effect of the atmosphere to reflect and intercept the light of the sun, is only sensible to the altitude of between 40 and 50 miles: and at that altitude we may collect, from what has been already said, that the air is above 10000 times rarer than at the surface of the earth. It is well known that the twilight begins and ends when the centre of the sun is about 18 degrees below the horizon, or only 17° 27', by subtracting 33' for refraction,



refraction, which raises the sun so much higher than he would be. And a ray coming from the sun in that position, and entering the earth's atmosphere, is refracted and bent into a curve line in passing through it to the eye. M. de la Hire took great pains to demonstrate, that, supposing the density of the atmosphere proportional to its weight, this curve is a cycloid: he also says, that if the ray be a tangent to the atmosphere, the diameter of its generating circle will be the height of the atmosphere; and that this diameter increases, till at last, when the rays are perpendicular, it becomes infinite, or the circle degenerates into a right line. This reasoning supposes that the refracting surface of the atmosphere is a plane; but since it is in reality a curve, he observes that these cycloids become in fact epicycloids. But Herman detected the error of M. de la Hire, and shewed that this curve is infinitely extended, and has an asymptote. And it is observed by Dr. Brook Taylor, in his *Methodus Increm.* pa. 168, &c, that this curve is one of the most intricate and perplexed that can well be proposed. The same ingenious author computes, that the refractive power of the air is to the force of gravity at the surface of the earth, as 320 millions to 1.

Considering the extreme rarity of the atmosphere at only 40 or 50 miles in height, it seems to be surprising that some meteors should be enflamed at such great heights as they have been observed at. A very remarkable one of this kind was observed by Dr. Halley in the month of March 1719, the altitude of which he computed at between 69 and 73½ English miles; its diameter 2800 yards, or more than a mile and a half; and its velocity about 350 miles in a minute. Others, apparently of the same kind, but whose altitude and velocity were still greater, have been observed; particularly that very remarkable one, of August 18th 1783, whose distance from the earth could not be less than 90 miles, its diameter at least as large as the former, while its velocity was certainly not less than 1000 miles in a minute. Now, from analogy of reasoning, it seems very probable, that the meteors which appear at such great heights in the air, are not essentially different from those which are seen on or near the surface of the earth. The difficulty with regard to the former is, that at the great heights above-mentioned, the atmosphere ought not to have any density sufficient to support flame, or to propagate sound; and yet such meteors are commonly succeeded by one explosion or more, and it is said are even sometimes accompanied with a hissing noise as they pass over our heads. The meteor of 1719 was not only very bright, seeming for a short time to turn night into day, but was attended with an explosion heard over all the island of Britain, causing a violent concussion in the atmosphere, and seeming to shake the earth itself: And yet, in the regions in which this meteor moved, the air ought to have been 300 thousand times rarer than the air we breathe, or 1000 times rarer than the vacuum commonly made by a good air-pump. Dr. Halley offers a conjecture, indeed, that the vast magnitude of such bodies might compensate for the thinness of the medium in which they moved. But

appearances of this kind are, by some others, attributed to electricity; though the circumstances of them cannot be reconciled to that cause; for the meteors move with all different degrees of velocity; and though the electrical fire easily pervades the vacuum of an air-pump, yet it does not in that case appear in bright well-defined sparks, as in the open air, but rather in long streams resembling the aurora borealis; and from some late experiments it has been concluded that the electric fluid cannot even penetrate a perfect vacuum.

*Of the Refractive and Reflective Power of the Atmosphere.* It has been observed above, that the atmosphere has a refractive power, by which the rays of light are bent from the right lined direction, as in the case of the twilight; and many other experiments manifest the same virtue, which is the cause of many phenomena. Alhazen, the Arabian, who lived about the year 1100, it seems was more inquisitive into the nature of refraction than former writers. But neither Alhazen, nor his follower Vitello, knew any thing of its just quantity, which was not known, to any tolerable degree of exactness, till Tycho Brahe, with great diligence, settled it. But neither did Tycho nor Kepler discover in what manner the rays of light were refracted by the atmosphere. Tycho thought the refraction was chiefly caused by dense vapours, very near the earth's surface: while Kepler placed the cause wholly at the top of the atmosphere, which he thought was uniformly dense; and thence he determined its altitude to be little more than that of the highest mountains. But the true constitution of the density of the atmosphere, deduced afterwards from the Torricellian experiment, afforded a juster idea of these refractions, especially after it appeared, by a repetition of Mr. Lowthorp's experiment, that the refractive power of the air is proportional to its density. By this variation in the density and refractive power of the air, a ray of light, in passing through the atmosphere, is continually refracted at every point, and thereby made to describe a curve, and not a straight line, as it would have done were there no atmosphere, or were its density uniform.

The atmosphere, or air, has also a reflective power; and this power is the means by which objects are enlightened so uniformly on all sides. The want of this power would occasion a strange alteration in the appearance of things; the shadows of which would be so very dark, and their sides enlightened by the sun so very bright, that probably we could see no more of them than their bright halves; so that for a view of the other halves, we must turn them half round, or if immoveable, must wait till the sun could come round upon them. Such a pellucid unreflective atmosphere would indeed have been very commodious for astronomical observations on the course of the sun and planets among the fixed stars, visible by day as well as by night; but then such a sudden transition from darkness to light, and from light to darkness, immediately upon the rising and setting of the sun, without any twilight, and even upon turning to or from the sun at noon day, would have been very inconvenient and offensive to our eyes. However, though the atmosphere



be greatly assistant in the illumination of objects, yet it must also be observed that it stops a great deal of light. By M. Bouguer's experiments, it seems that the light of the moon is often 2000 times weaker in the horizon, than at the altitude of 66 degrees; and that the proportion of her light at the altitudes of 66 and 19 degrees, is about 3 to 2; and the lights of the sun must bear the same proportion to each other at those heights; which Bouguer made choice of, as being the meridian heights of the sun, at the summer and winter solstices, in the latitude of Croisic in France. Smith's optics. Rem. 95.

For the Atmosphere of the sun, moon, and planets, see the respective articles.

ATMOSPHERE of *Solid or Consistent Bodies*, is a kind of sphere formed by the effluvia, or minute corpuscles, emitted from them. Mr. Boyle endeavours to shew, that all bodies, even the hardest and most coherent, as gems, &c, have their atmospheres.

ATMOSPHERE, in *Electricity*, denotes that medium which is conceived to be diffused over the surface of electrified bodies, and to some distance around them, and consisting of effluvia issuing from them; by which, other bodies immersed in it become endued with an electricity contrary to that of the body to which the atmosphere belongs. This was first noticed at a very early period in the history of this science by Otto Guericke, and afterwards by the academicians *del Cimento*, who contrived to render the electric atmosphere visible, by means of smoke attracted by a piece of amber, and gently rising from it, but vanishing as the amber cooled. Dr. Franklin exhibited this electric atmosphere with greater advantage, by dropping rosin on hot iron plates held under electrified bodies, from which the smoke arose and encompassed the bodies, giving them a very beautiful appearance. But the theory of electric atmospheres was not well explained and understood for a considerable time; and the investigation led to many curious experiments and observations. The experiments of Mr. Canton and Dr. Franklin prepared the way for the conclusion that was afterwards drawn from them by Mess. Wilcke and Epinus, though they retained the common opinion of electric atmospheres, and endeavoured to explain the phenomena by it. The conclusion was, that the electric fluid, when there is a redundancy of it in any body, repels the electric fluid in any other body, when they are brought within the sphere of each other's influence, and drives it into the remote parts of the body, or quite out of it, if there be any outlet for that purpose.

By Atmosphere, M. Epinus says, no more is to be understood than the sphere of action belonging to any body, or the neighbouring air electrified by it. Sig. Beccaria agrees in the same opinion, that electrified bodies have no other atmosphere than the electricity communicated to the neighbouring air, and which goes with the air, and not with the electrified bodies. Mr. Canton also, having relinquished the opinion that electrical atmospheres were composed of effluvia from excited or electrified bodies, maintained that they only result from an alteration in the state of the electric fluid contained in it, or belonging to the air surrounding these bodies to a certain distance; for instance, that

excited glass repels the electric fluid from it, and consequently beyond that distance makes it more dense; whereas excited wax attracts the electric fluid existing in the air nearer to it, making it rarer than it was before. In the course of experiments that were performed on this occasion, Mess. Wilcke and Epinus succeeded in charging a plate of air, by suspending large boards of wood covered with tin, with the flat sides parallel to one another, and at some inches asunder: for they found, upon electrifying one of the boards positively, that the other was always negative; and a shock was produced by forming a communication between the upper and lower plates. Beccaria has largely considered the subject of electric atmospheres, in his *Artificial Electricity*, pa. 179 &c, Eng. edit. See also Dr Priestley's *Hist. of Electricity*, vol. ii. sect. 5. and Cavallo's *Electricity*, pa. 241.

ATMOSPHERE, *Magnetic, &c*, is understood of the sphere within which the virtue of the magnet, &c, acts.

ATOM, a particle of matter indivisible on account of its solidity, hardness, and impenetrability; which preclude all division, and leave no vacancy for the admission of any foreign force to separate or disunite its parts. As atoms are the first matter, it is necessary they should be indissoluble, that they may be incorruptible. Newton adds, it is also required that they be immutable, that the world may continue in the same state, and bodies be of the same nature now as formerly.

ATOMICAL PHILOSOPHY, or the doctrine of atoms, a system which accounted for the origin and formation of things, from the hypothesis that atoms are endued with weight and motion. This philosophy was first taught by Moschus, some time before the Trojan war: but it was most cultivated by Epicurus; whence it is called the Epicurean philosophy.

ATTRACTION, or ATTRACTIVE POWER, a general term used to denote the cause, power, or principle, by which all bodies mutually tend towards each other, and cohere, till separated by some other power. The laws, phenomena, &c, of attraction, form the chief subject of Newton's philosophy, being the principal agent of nature, in almost all her wonderful operations.

The principle of attraction, in the Newtonian sense of it, it seems was first surmised by Copernicus. "As for gravity, says he, I consider it as nothing more than a certain natural appetite (*appetentia*) that the Creator has impressed upon all the parts of matter, in order to their uniting or coalescing into a globular form, for their better preservation; and it is probable that the same power is also inherent in the sun and moon, and planets, that those bodies may constantly retain that round form in which we see them." *De Revol. Orb. Caelest.* lib. i, cap. 9. Kepler calls gravity a corporeal and mutual affection between similar bodies, in order to their union. *Ast. Nov. in Introd.* And he pronounced more positively that no bodies whatever were absolutely light, but only relatively so; and consequently that all matter was subjected to the power and law of gravitation. *Ibid.*

The first in this country who adopted the notion of attraction, was Dr. Gilbert, in his book *De Magnete*; and the next was the celebrated Lord Bacon, in his *Nov. Organ.* lib. ii, aphor. 36, 45, 48. *Sylv.* cent. i, exp.



exp. 33; also in his treatise *De Motu*, particularly under the articles of the 9th and the 13th sorts of Motion. In France it was received by Fermat and Roberval; and in Italy by Galileo and Borelli. But till Newton appeared, this principle was very imperfectly defined and applied.

It must be observed, that though this great author makes use of the word attraction, in common with the school philosophers, yet he very studiously distinguishes between the ideas. The ancient attraction was conceived to be a kind of quality inherent in certain bodies themselves, and arising from their particular or specific forms. But the Newtonian attraction is a more indefinite principle; denoting not any particular kind or mode of action; nor the physical cause of such action; but only a general tendency, a *conatus accedendi*, to whatever cause, physical or metaphysical, such effect be owing; whether to a power inherent in the bodies themselves, or to the impulse of an external agent. Accordingly, that author remarks, in his *Philos. Nat. Prin. Math.* "that he uses the words *attraction*, *impulse*, and *propension* to the centre, indifferently; and cautions the reader not to imagine that by attraction he expresses the modus of the action, or its efficient cause, as if there were any proper powers in the centres, which in reality are only mathematical points; or as if centres could attract." Lib. 1, pa. 5. So, he "considers centripetal powers as attractions, though, physically speaking, it were perhaps more just to call them impulses. Ib. pa. 147. He adds, "that what he calls attraction may possibly be effected by impulse, though not a common or corporeal impulse, or after some other manner unknown to us." *Optic.* p. 322.

Attraction, if considered as a quality arising from the specific forms of bodies, ought, together with sympathy, antipathy, and the whole tribe of occult qualities, to be exploded. But when these are set aside, there will remain innumerable phenomena of nature, and particularly the gravity or weight of bodies, or their tendency to a centre, that argue a principle of action seemingly distinct from impulse; where, at least, there is no sensible impulsion concerned. Nay, what is more, this action, in some respects, differs from all impulsion we know of; impulse being always found to act in proportion to the surfaces of bodies; whereas gravity acts according to their solid content, and consequently it must arise from some cause that penetrates or pervades the whole substance of it. This unknown principle, unknown we mean in respect of its cause, for its phenomena and effects are most obvious, with all its species and modifications, is called attraction; being a general name, under which may be ranged all mutual tendencies, where no physical impulse appears, and which consequently cannot be accounted for upon any known laws of nature.

And hence arise divers particular kinds of attraction; as *Gravity*, *Magnetism*, *Electricity*, &c, which are so many different principles, acting by different laws; and only agreeing in this, that we do not perceive any physical causes of them: but that, as to our senses, they may really arise from some power or efficacy in such bodies, by which they are enabled to act even upon distant bodies; though our reason absolutely disallows of any such action.

Attraction may be divided, with respect to the law it observes, into two kinds.

1. That which extends to a sensible distance. As the attraction of gravity, which is found in all bodies; and the attractions of magnetism and electricity, found only in particular bodies. The several laws and phenomena of each, see under their respective articles.

The attraction of gravity, called also among mathematicians the *centripetal force*, is one of the greatest and most universal principles of all nature. We see and feel it operate on bodies near the earth, and find by observation that the same power (i. e. a power which acts in the same manner, and by the same rules, viz, always proportionally to the quantities of matter, and inversely as the squares of the distances) does also obtain in the moon, and the other planets, both primary and secondary, as well as in the comets; and even that this is the very power by which they are all retained in their orbits, &c. And hence, as gravity is found in all the bodies which come under our observation, it is easily inferred, by one of the established rules of philosophizing, that it obtains in all others. And since it is found to be proportional to the quantity of matter in any body, it must exist in every particle of it: and hence it is proved that every particle in nature attracts every other particle.

From this attraction arises all the motion, and consequently all the mutation, in the great world. By this heavy bodies descend, and light ones are made to ascend: by this projectiles are directed, vapours and exhalations rise, and rains &c fall: by this rivers glide, the ocean swells, the air presses, &c. In short, the motions and forces arising from this principle, constitute the subject of that extensive branch of mathematics, called *mechanics* or *statics*, with the parts or appendages of it, as hydrostatics, pneumatics, hydraulics, &c.

2. That which does not extend to sensible distances. Such is found to obtain in the minute particles of which bodies are composed, attracting each other at or extremely near the point of contact, with forces often much superior to that of gravity, but which at any distance decrease much faster than the power of gravity. This power a late ingenious author calls the *attraction of cohesion*, as being that by which the atoms or insensible particles of bodies are united into sensible masses.

This kind of attraction owns Sir Isaac Newton for its discoverer; as the former does for its improver. The laws of motion, percussion, &c, in sensible bodies, under various circumstances, as falling, projected, &c, ascertained by the later philosophers, do not reach those more reclusive, intestine motions in the component particles of the same bodies, on which depend the changes in the texture, colour, properties, &c, of bodies. So that our philosophy, if it were only founded on the principle of gravitation, and even carried as far as this would lead us, would still be very deficient.

But besides the common laws of sensible masses, the minute parts they are composed of are found subject to some others, which have but lately been noticed, and are even yet imperfectly known. Newton himself, to whose happy penetration we owe the hint, limits himself with establishing that there are such motions in the



*minima nature*, and that they flow from certain powers or forces, not reducible to any of those in the great world. He shews that, by virtue of these powers, "the small particles act on one another even at a distance; and that many of the phenomena of nature result from it. Sensible bodies, we have already observed, act on one another divers ways; and as we thus perceive the tenor and course of nature, it appears highly probable that there may be other powers of the like kind; nature being very uniform and consistent with herself. Those just mentioned, reach to sensible distances, and so have been observed by vulgar eyes; but there may be others which reach to such small distances as have hitherto escaped observation; and it is probable electricity may reach to such distances, even without being excited by friction."

The great author just mentioned proceeds to confirm the reality of these suspicions from a great number of phenomena and experiments, which plainly argue such powers and actions between the particles, for example of salts and water, oil of vitriol and water, aquafortis and iron, spirit of vitriol and saltpetre. He also shews, that these powers, &c. are unequally strong between different bodies; stronger, for instance, between the particles of salt of tartar and those of aquafortis than those of silver, between aquafortis and lapis calaminaris than iron, between iron than copper, and copper than silver or mercury. So spirit of vitriol acts on water, but more on iron or copper, &c. And the other experiments are innumerable which countenance the existence of such principle of attraction in the particles of matter.

These actions, by virtue of which the particles of the bodies above-mentioned tend towards each other, the author calls by a general indefinite name, *attraction*; a name equally applicable to all actions by which bodies tend towards one another, whether by impulse, or by any other more latent power: and from hence he accounts for an infinity of phenomena, otherwise inexplicable, to which the principle of gravity is inadequate.

Thus, adds our author, "will nature be found very conformable to herself, and very simple; performing all the great motions of the heavenly bodies by the attraction of gravity, which intercedes those bodies, and almost all the small ones of their parts, by some other attractive power diffused through their particles. Without such principles, there never would have been any motion in the world; and without the continuance of it, motion would soon perish, there being otherwise a great decrease or diminution of it, which is only supplied by these active principles."

It need not be said how unjust it is in the generality of foreign philosophers to declare against a principle which furnishes so beautiful a view, for no other reason but because they cannot conceive how one body should act on another at a distance. It is indeed true, that philosophy allows of no action but what is by immediate contact and impulsion; for how can a body exert any active power where it does not exist? yet we see effects, without perceiving any such impulse; and where effects are observed, there must exist causes whether we see them or not. But we may contemplate such effects, without entering into the considera-

tion of the causes, as indeed it seems the business of a philosopher to do: for to exclude a number of phenomena which we do see, would be to leave a great chasm in the history of nature; and to argue about actions which we do not see, would be to raise castles in the air. It follows therefore, that the phenomena of attraction are matter of physical consideration, and as such intitled to a share in the system of physics; but that their causes will only become so when they become sensible, that is when they appear to be the effect of some other higher causes; for a cause is no otherwise seen than as itself is an effect, so that the first cause must needs be always invisible: we are therefore at liberty to suppose the causes of attractions what we please, without any injury to the effects. The illustrious author himself seems to be a little indetermined as to the causes; inclining sometimes to attribute gravity to the action of an immaterial cause (*Optics*, pa. 343 &c), and sometimes to that of a material one, *Ib.* pa. 325.

In his philosophy, the research into causes is the last thing, and never comes under consideration till the laws and phenomena of the effect be settled; it being to these phenomena that the cause is to be accommodated. The cause even of any, the grossest and most sensible action, is not adequately known. How impulse or percussion itself produces its effects, that is how motion is communicated from body to body, confounds the deepest philosophers; yet is impulse received not only into philosophy, but into mathematics: and accordingly the laws and phenomena of its effects make the chief part of common mechanics.

The other species of attraction, therefore, in which no impulse is observable, when their phenomena are sufficiently ascertained, have the same title to be promoted from physical to mathematical consideration; and this without any previous inquiry into their causes, to which our conceptions may not be proportionate.

Our great philosopher, then, far from adulterating science with any thing foreign or metaphysical, as many have reproached him with doing, has the glory of having thrown every thing of this kind out of his system, and of having opened a new source of sublimer mechanics, which, duly cultivated, might be of far greater extent than all the mechanics yet known. Hence it is alone that we must expect to learn the manner of the changes, productions, generations, corruptions, &c. of natural things; with all that scene of wonders opened to us by the operations of chemistry.

Some of our own countrymen have prosecuted the discovery with laudable zeal. Dr. Keil particularly has endeavoured to deduce some of the laws of this new action, and applied them in resolving several of the more general phenomena of matter, as cohesion, fluidity, elasticity, softness, fermentation, coagulation, &c: and Dr. Freind, seconding his endeavours, has made a farther application of the same principles, at once to account for almost all the phenomena that chemistry presents. So that some philosophers are inclined to think that the new mechanics should seem already raised to a complete science, and that nothing now can occur but what we have an immediate solution of, from the principles of attractive forces.

But



But this seems a little too precipitate: a principle so fertile should have been further explained; its particular laws, limits, &c, more industriously detected and laid down, before we had proceeded to the application. Attraction in the gross is so complex a thing, that it may solve a thousand different phenomena alike. The notion is but one degree more simple and precise than action itself; and, till its properties are more fully ascertained, it were better to apply it less, and study it more. It may be added, that some of Newton's followers have been charged with falling into that error which he industriously avoided, viz, of considering attraction as a cause or active property in bodies, not merely as a phenomenon or effect.

For the laws, properties, &c, of the different sorts of Attraction, see their particular articles COHESION, GRAVITY, MAGNETISM, &c.

ATTRACTION, *Centre of*, See CENTRE of Attraction.

ATTRACTION of Mountains, See MOUNTAINS.

ATTRITION, the striking or rubbing of bodies against one another, so as to throw off some of their superficial particles: such as the grinding and polishing of bodies. Or simply the act of rubbing: as when amber and other electric bodies are rubbed, to make them attract, or emit their electric force.

AVANT-FOSS, or Ditch of the Counterscarp, in Fortification, is a wet ditch surrounding the counterscarp, on the outer side, next to the country, at the foot of the glacis. It would not be proper to have such a ditch if it could be laid dry, as it would then serve as a lodgment for the enemy.

AVERROES, or ABEN-ROES, a very subtle Arabian philosopher, who flourished about the end of the 11th century, when the Moors had possession of part of Spain. He was the son of the high priest and chief judge of Corduba or Cordova in Spain: but he was educated in the university of Morocco, where he was professor, and where he died in 1206, having there studied natural philosophy, medicine, mathematics, law, and divinity. After the death of his father, he enjoyed his posts in Spain, to which was afterward added that of judge of Morocco and Mauritania, where having settled deputies, he returned to his duty in Spain. Notwithstanding he was very rich, and had a very great income, his liberality to men of letters in necessity, whether they were his friends or his enemies, kept him always in debt. He was afterwards stripped of all his posts, and thrown into prison, for heresy, by the instigations of bad men, his enemies; but the oppressions of the judge who succeeded him, caused him to be restored to his former employments.

He was excessively fat, though he eat but once a day, and spent most part of the night in the study of philosophy, when he was fatigued, amusing himself with reading poetry or history. He was never seen to play at any game, or to partake in any diversion. He was extremely fond of Aristotle's works, and wrote commentaries upon them; whence he was styled *the Commentator*, by way of eminence. He wrote many other pieces; among them a work on the Whole Art of Physics; an Epitome of Ptolemy's Almagest, which Vossius dates about the year 1149; also a Treatise of

Astrology, which was translated into Hebrew by R. Jacob Ben Samson, and said to be extant in the French king's library. He wrote also several poems, and many amorous verses, but these last he threw into the fire when he grew old. His other poems are lost, except a small piece, in which he says, "that when he was young, he acted against his reason; but that when he was in years, he followed its dictates;" upon which he utters this wish, "Would to God I had been born old, and that in my youth I had been in a state of perfection!" As to religion, his opinions were, that Christianity is absurd; Judaism, the religion of children; Mahometanism, the religion of swine.

AVICENA, AVICENNE, or AVICENES, has been accounted the prince of Arabian philosophers and physicians. He was born at Assena, near Bokhara, in 978; and died at Hamadan in 1036, being 58 years of age.

The first years of Avicena were employed on the study of the Belles Lettres, and the Koran, and at 10 years of age he was perfect master of the hidden senses of that book. Then applying to the study of logic, philosophy and mathematics, he quickly made a rapid progress. After studying under a master the first principles of logic, and the first 5 or 6 propositions of Euclid's elements, he became disgusted with the slow manner of the schools, applied himself alone, and soon accomplished all the rest by the help of the commentators only.

Possessed with an extreme avidity to be acquainted with all the sciences, he studied medicine also. Persuaded that this art consists as much in practice as in theory, he sought all opportunities of seeing the sick; and afterwards confessed that he had learned more from such experience than from all the books he had read. Being now in his 16th year, and already celebrated for being the light of his age, he determined to resume his studies of philosophy, which medicine, &c, had made him for some time neglect: and he spent a year and a half in this painful labour, without ever sleeping all this time a whole night together. At the age of 21, he conceived the bold design of incorporating, in one work, all the objects of human knowledge; and he carried it into execution in an Encyclopedia of 20 volumes, to which he gave the title of the *Utility of Utilities*.

Many wonderful stories are related of his skill in medicine, and the cures which he performed. Several princes had been taken dangerously ill, and Avicenes was the only one that could know their ailments, and cure them. His reputation increased daily, and all the princes of the east desired to retain him in their families, and in fact he passed through several of them. But the irregularities of his conduct sometimes lost him their favour, and threw him into great distresses. His excesses in pleasures, and his infirmities, made a poet say, who wrote his epitaph, that the profound study of philosophy had not taught him good morals, nor that of medicine the art of preserving his own health.

After his death however, he enjoyed so great a reputation, that till the 12th century he was preferred for the study of philosophy and medicine to all his predecessors. Even in Europe his works were the only writings in vogue in the schools. They were very numerous, and various, the titles of which are as follow:

1. Of



1. Of the Utility and Advantage of the Sciences, in 20 books.—2. Of Innocence and Criminality, 2 books.—3. Of Health and Remedies, 18 books.—4. On the Means of preserving Health, 3 books.—5. Canons of Physic, 14 books.—6. On Astronomical Observations, 1 book.—7. On Mathematical Sciences.—8. Of Theorems, or Mathematical and Theological Demonstrations, 1 book.—9. On the Arabic Language, and its Properties, 10 books.—10. On the Last Judgment.—11. On the Origin of the Soul, and the Resurrection of Bodies.—12. On the end we should propose to ourselves in Harangues and Philosophical Argumentations.—13. Demonstration of the Collateral Lines in the Sphere.—14. Abridgment of Euclid.—15. On Finiteness and Infinity.—16. On Physics and Metaphysics.—17. On Animals and Vegetables, &c.—18. Encyclopedie, 20 volumes.

AUGUST, the 8th month of the year, containing 31 days. In the antient Roman calendar this was called *sextilis*, as being the 6th month from March, with which their year began; but changed to its present name by the emperor Augustus, calling it after his own name on account of his having obtained many victories and honours in that month.

AVOIRDUPOIS *Weight*, a weight used in England for weighing all the larger and coarser sorts of goods; as groceries, cheese, butter, flesh, wool, salt, hops, &c, and all metals except gold and silver. Avoirdupois weight is thus divided, viz.

16 dr. or drams make	1 ounce, marked	oz.
16 oz. - - - -	1 pound, - -	lb.
112 lb. - - - -	1 hundred weight,	cwt.
20 cwt. - - -	1 ton - -	ton.

The Avoirdupois ounce is less than the Troy ounce, in the proportion of 700 to 768, but the Avoirdupois pound greater than the Troy pound in the proportion of 700 to 576;

for 1 lb Avoird. is = 7000 grains Troy,  
but 1 lb Troy is = 5760 grains Troy,  
also 1 oz Avoird. is =  $437\frac{1}{2}$  grains Troy,  
and 1 oz Troy is = 480 grains Troy.

AURIGA, the *Waggoner*, a constellation in the northern hemisphere, consisting of 14 stars in Ptolemy's catalogue; but in Tycho's, 27; in Hevelius's, 40; and in the Britannic catalogue, 66.

This is one of the 48 old asterisms, mentioned by all the most ancient astronomers. It is represented by the figure of an old man, in a posture somewhat like sitting, with a goat and her kids in his left hand, and a bridle in his right.

The Greeks probably received this and all their other constellations from the Egyptians; but, wanting to appear the inventors of them themselves, and not understanding the meaning of the figures, they have clothed them with some of their own fabulous dresses, to favour the deceit. They accordingly tell us that this figure of a waggoner was an honourable character, and Erichthonius, the inventor of coaches. Vulcan, say they, once fell in love with Minerva, and when he could not prevail with her to marry him, he would have obtained her upon less honourable terms. There was a struggle

between them, and some way or other Erichthonius was begotten, though it does not seem that Minerva had much share in it: she took care of the offspring however. Some have supposed it was only a serpent; but the graver authors say, Erichthonius was a man with legs only like the body of a serpent, and that to hide this monstrous part of his figure he invented coaches to carry him about. They add, that Jupiter, doing him honour for an invention that was, in some degree, imitating the sun's carriage on the earth, raised him up among the stars.

But others, ill satisfied with a story which so badly agreed with the figure, have said that it belonged to Myrtillus, a son of Mercury and Clytie, and charioteer to Aenomanus; they say, that at his death, his father Mercury, by permission of his superiors, raised him up into the skies. All this however does not at all account for the goat and her two kids in the hands of Auriga. To set this right, they afterward made Auriga to be Olenus, a son of Vulcan, and the father of Aega and Helice, two of the Cretan nymphs that nursed the infant Jupiter. They talk of a goat that was used for giving milk to the young deity, and they suppose that this creature, and its two young ones, were placed in the hands of the father of the virgins, to commemorate the creature they took into their service on that occasion.

Besides the Hædi, this constellation contains also another of those stars which the ancients honoured with peculiar names, the goat Capra, and Amalthæa Capra: this is the bright one near the shoulder, and supposed to be the mother of the Hædi, and the nurse of Jupiter.

Although the whole constellation of Auriga is not mentioned among those from which the ancients formed presages of the succeeding weather, the two stars in his arm were of the foremost in that rank. It is these they called by the name Hædi, and dreaded so extremely on account of the storms and tempests that succeeded their rising, that it is said they shut up the navigation of the sea for their season. And the day of their influence being over, we find, was celebrated as a festival with sports and games, under the name of Natalis Navigationis. Germanicus calls them unfriendly stars to mariners; and Virgil couples them with Arcturus, mentioning their setting and its rising as things of the most important presage. Horace also puts them together as the most formidable of all the stars to those who follow the traffic of the sea. And to the same purpose speak all the ancient writers, thus making a part of the constellation Auriga, if not the whole constellation, a thing to be observed with the utmost attention, and to be feared as much as the blazing Arcturus.

AURORA, the morning twilight; or that faint light which appears in the morning when the sun is within 18 degrees of the horizon.

AURORA BOREALIS, NORTHERN LIGHT, or *Streamers*; a kind of meteor appearing in the northern part of the heavens, mostly in the winter season, and in frosty weather. It is usually of a reddish colour, inclining to yellow, and sends out frequent coruscations of pale light, which seem to rise from the horizon in a pyramidal undulating form, and shooting with great velocity up to the zenith. It appears often in form of an arch, which



which is partly bright, and partly dark, but generally transparent. And the matter of it is not found to have any effect on the rays of light, which pass freely through it. Dr. Hamilton observes, that he could plainly discern the smallest speck in the Pleiades through the density of those clouds which formed part of the Aurora borealis in 1763, without the least diminution of its splendour, or increase of twinkling. *Philos. Essays*, pa. 106.

Sometimes it produces an Iris. Hence M. Godin judges, that most of the extraordinary meteors and phenomena in the skies, related as prodigies by historians, as battles, and the like, may probably enough be reduced to the class of Auroræ boreales. *Hist. Acad. R. Scienc.* for 1762, pa. 405.

This kind of meteor never appears near the equator; but, it seems, is frequent enough towards the south pole, like as towards the north, having been observed there by voyagers. See *Philos. Trans.* N° 461, and vol. 54; also Forster's account of his voyage round the world with Captain Cook, where he describes their appearance as observed for several nights together, in sharp frosty weather, which was much the same as those observed in the north, excepting that they were of a lighter colour.

It seems that meteors of this kind have appeared sometimes more frequently than others. They were so rare in England, or else so little regarded, that none are recorded in our annals since that remarkable one of Nov. 14, 1574, till the surprising Aurora borealis of March 6, 1716, which appeared for three nights successively, but by far more strongly on the first: except that five small ones were observed in the year 1707 and 1708. Hence it would seem, that the air, or earth, or both, are not at all times disposed to produce this phenomenon.

The extent of these appearances is also amazingly great. That in March 1716 was visible from the west of Ireland, to the confines of Russia and the east of Poland; extending at least near 30 degrees of longitude, and from about the 50th degree in latitude, over almost all the north of Europe; and in all places, at the same time, it exhibited the like wondrous appearances. Father Boscovich has determined the height of an aurora borealis, which was observed by the Marquis of Polini the 16th of December, 1737, and found it was 825 miles high; and Mr. Bergman, from a mean of 30 computations, makes the average height of the aurora borealis amount to 70 Swedish, or 469 English miles. But Euler supposes the height to be several thousands of miles; and Mairan also assigns to them a very elevated region.

Many attempts have been made to determine the cause of this phenomenon. Dr. Halley imagines that the watery vapours, or effluvia, exceedingly rarefied by subterraneous fire, and tinged with sulphureous streams, which many naturalists have supposed to be the cause of earthquakes, may also be the cause of this appearance: or that it is produced by a kind of subtile matter, freely pervading the pores of the earth, and which, entering into it nearer the southern pole, passes out again with some force into the æther, at the same distance from the northern. This subtile matter, by becoming more dense, or having its velocity increased, may perhaps be

capable of producing a small degree of light, after the manner of effluvia from electric bodies, which, by a strong and quick friction, emit light in the dark; to which sort of light this seems to have a great affinity. *Philos. Trans.* N° 347. See also Mr. Cotes's description of this phenomenon, and his method of explaining it, by streams emitted from the heterogeneous and fermenting vapours of the atmosphere, in *Smith's Optics*, pa. 69; or *Philos. Trans. abr.* vol. 6, part 2.

The celebrated M. de Mairan, in an express treatise on the Aurora Borealis, published in 1731, supposes its cause to be the zodiacal light, which, according to him, is no other than the sun's atmosphere: this light happening, on some occasions, to meet the upper parts of our atmosphere about the limits where universal gravity begins to act more forcibly towards the earth than towards the sun, falls into our air to a greater or less depth, as its specific gravity is greater or less, compared with the air through which it passes. See *Tract. Phys. et Hist. de l'Aurore Boreale. Suite des Memoires de l'Acad. R. des Scien.* 1731. Also *Philos. Trans.* N° 433, or *Abridg.* vol. 8, pa. 540.

However M. Euler thinks the cause of the aurora borealis not owing to the zodiacal light, as M. de Mairan supposes; but to particles of our atmosphere, driven beyond its limits by the impulse of the solar light. And on this supposition he endeavours to account for the phenomena observed concerning this light. He supposes the zodiacal light, and the tails of comets, to be owing to a similar cause.

But ever since the identity of lightning and the electric matter has been determined, philosophers have been naturally led to seek for the explication of aerial meteors in the principles of electricity; and there is now no doubt but most of them, and especially the aurora borealis, are electrical phenomena. Besides the more obvious and known appearances which constitute a resemblance between this meteor and the electric matter by which lightning is produced, it has been observed that the aurora occasions a very sensible fluctuation in the magnetic needle; and that when it has extended lower than usual in the atmosphere, the flashes have been attended with various sounds of rumbling and hissing, especially in Russia and the other more northern parts of Europe; as noticed by Sig. Beccaria and M. Messier. Mr. Canton, soon after he had obtained electricity from the clouds, offered a conjecture, that the aurora is occasioned by the dashing of electric fire positive towards negative clouds at a great distance, through the upper part of the atmosphere, where the resistance is least: and he supposes that the aurora which happens at the time when the magnetic needle is disturbed by the heat of the earth, is the electricity of the heated air above it: and this appears chiefly in the northern regions, as the alteration in the heat of the air in those parts is the greatest. Nor is this hypothesis improbable, when it is considered, that electricity is the cause of thunder and lightning; that it has been extracted from the air at the time of the aurora borealis; that the inhabitants of the northern countries observe it remarkably strong when a sudden thaw succeeds very cold severe weather; and that the tourmalin is known to emit and absorb the electric fluid only by the increase or diminution of its heat.



heat. Positive and negative electricity in the air, with a proper quantity of moisture to serve as a conductor, will account for this and other meteors, sometimes seen in a serene sky. Mr. Canton has since contrived to exhibit this meteor by means of the Torricellian vacuum, in a glass tube about 3 feet long, and sealed hermetically. When one end of the tube is held in the hand, and the other applied to the conductor, the whole tube will be illuminated from end to end, and will continue luminous without interruption for a considerable time after it has been removed from the conductor. If, after this, it be drawn through the hand either way, the light will be remarkably intense through the whole length of the tube. And though a great part of the electricity be discharged by this operation, it will still flash at intervals, when held only at one extremity, and kept quite still; but if, at the same time, it be grasped by the other hand in a different place, strong flashes of light will dart from one end to the other; and these will continue 24 hours or more, without a fresh excitation. Sig. Beccaria conjectures that there is a constant and regular circulation of the electric fluid from north to south; and he thinks that the aurora borealis may be this electric matter performing its circulation in such a state of the atmosphere as renders it visible, or approaching nearer than usual to the earth. Though probably this is not the mode of its operation, as the meteor is observed in the southern hemisphere, with the same appearances as in the northern. Dr. Franklin supposes, that the electric fire discharged into the polar regions, from many leagues of vaporised air raised from the ocean between the tropics, accounts for the aurora borealis; and that it appears first, where it is first in motion, namely in the most northern part; and the appearance proceeds southward, though the fire really moves northward. Franklin's *Exper. and Obs.* 1769, pa. 49. *Philos. Trans.* vol. 48, pa. 358, 784; *Ib.* vol. 51, pa. 403; *Lettere dell' Ellettricismo*, pa. 269; or Priestley's *Hist. of Electricity*. See also an ingenious solution of this phenomenon, on the same principles, by Dr. Hamilton, in his *Philos. Essays*. Mr. Kirwan (in the *Transactions of the Royal Irish Academy*, ann. 1788) has some ingenious remarks on the *auroræ borealis & australis*. He gives his reasons for supposing the rarefaction of the atmosphere in the polar regions to proceed from them, and these from a combustion of inflammable air caused by electricity. He observes, that after an aurora borealis the barometer commonly falls, and high winds from the south generally follow.

AURUM FULMINANS, a preparation from gold, which being thrown into the fire, it explodes with a violent noise, like thunder. The matter is produced by dissolving gold in aqua regia, and precipitating the solution by oil of tartar *per deliquium*, or volatile spirit of sal ammoniac. The powder being washed in warm water, and dried to the consistence of a paste, is afterwards formed into small grains of the size of hempseed.

It is inflammable, not only by fire, but also by a gentle warmth; and gives a report much louder than that of gunpowder. A single grain laid on the point

of a knife, and lighted at a candle, explodes with a greater report than a musquet: and a scruple of this powder, it is said, acts more loudly than half a pound of gunpowder; and yet it is said that, by mixture, it does not increase the elastic force of fired gunpowder. Dr. Black attributes the increase of weight, and also the explosive property of this powder, to adhering fixable air.—This is a very dangerous preparation, and should be used with great caution.

AUSTRAL, the same with *southern*. Thus, Austral signs, are the last 6 signs of the zodiac; and are so called because they are on the south side of the equinoctial.

AUSTRALIS *Corona*; see *CORONA Australis*.

AUSTRALIS PISCIS, the *Southern Fish*, is a constellation of the southern hemisphere. See *Piscis Australis*.

AUTOMATON, a seemingly self-moving machine; or one so constructed, by means of weights, levers, pulleys, springs, &c, as to move for a considerable time, as if it were endued with animal life. And according to this description, clocks, watches, and all machines of that kind, are automata.

It is said, that Archytas of Tarentum, 400 years before Christ, made a wooden pigeon that could fly; that Archimedes also made such-like automata; that Regiomontanus made a wooden eagle that flew forth from the city, met the emperor, saluted him, and returned; also that he made an iron fly, which flew out of his hand at a feast, and returned again after flying about the room; that Dr. Hook made the model of a flying chariot, capable of supporting itself in the air. Many other surprizing automata we have been eye-witnesses of, in the present age: thus, we have seen figures that could write, and perform many other actions in imitation of animals: M. Vaucanson made a figure that played on the flute; the same gentleman also made a duck, which was capable of eating, drinking, and imitating exactly the voice of a natural one; and, what is still more surprizing, the food it swallowed was evacuated in a digested state, or considerably altered on the principles of solution; also the wings, viscera, and bones were formed so as strongly to resemble those of a living duck; and the actions of eating and drinking shewed the strongest resemblance, even to the muddling the water with its bill. M. Le Droz of la Chaux de Fonds, in the province of Neuchatel, has also executed some very curious pieces of mechanism: one was a clock, presented to the king of Spain; which had, among other curiosities, a sheep that imitated the bleating of a natural one, and a dog watching a basket of fruit, that barked and snarled when any one offered to take it away; besides a variety of human figures, exhibiting motions truly surprizing. But all these seem to be inferior to M. Kempell's chess-player, which may truly be considered as the greatest master-piece in mechanics that ever appeared in the world. See also Baptista Porta's *Magia Nat.* c. 19, and Scaliger's *Subtil.* 326.

AUTUMN, the third season, when the harvest and fruits are gathered in. This begins at the descending equinox, which, in the northern hemisphere,



is when the sun enters the sign Libra, or about the 22d day of August; and it ends, when winter commences, about the same day in December.

**AUTUMNAL**, something belonging to autumn. Thus,

**AUTUMNAL Equinox**, the time when the sun enters the descending point of the ecliptic, where it crosses the equinoctial; and is so called, because the nights and days are then equal.

**AUTUMNAL Point**, the point of the ecliptic answering to the autumnal equinox.

**AUTUMNAL Signs**, are the signs Libra, Scorpio, Sagittary, through which the sun passes during the autumn.

**AXIOM**, a self-evident truth, or a proposition assented to by every person at first sight. Such as, that the whole is greater than its part; that a thing cannot both be and not be at the same time; and that from nothing, nothing can arise.

Some Axioms are in effect, strictly speaking, no other than identical propositions. Thus, to say that all right angles are equal to each other, is as much as to say, all right angles are right angles; such equality being implied in the very definition, or the very name or term itself.

**AXIOM** is also an established principle in some art or science. Thus, it is an axiom in physics, that nature does nothing in vain; that effects are proportional to their causes; &c. It is an axiom in geometry, that two things equal to the same third thing, are also equal to each other; that if to equal things equals be added, the sums will be equal. And it is an axiom in optics, that the angle of incidence is equal to the angle of reflection. In this sense also the general laws of motion are called axioms; as, that all motion is rectilinear, that action and reaction are equal, &c.

**AXE** or **AXIS**, in *Geometry*, the straight line in a plane figure, about which it revolves, to produce or generate a solid. Thus, if a semicircle be moved round its diameter at rest, it will generate a sphere, whose axis is that diameter. And if a right-angled triangle be turned about its perpendicular at rest, it will describe a cone, whose axis is that perpendicular.

**AXIS** is yet more generally used for a right line conceived to be drawn from the vertex of a figure to the middle of the base. So the

**Axis of a circle or sphere**, is any line drawn through the centre, and terminated at the circumference, on both sides.

**Axis of a cone**, is the line from the vertex to the centre of the base.

**Axis of a cylinder**, is the line from the centre of the one end to that of the other.

**Axis of a conic section**, is the line from the principal vertex, or vertices, perpendicular to the tangent at that point. The ellipse and hyperbola have each two axes, which are finite and perpendicular to each other; but the parabola has only one, and that infinite in length.

**Transverse Axis**, in the Ellipse and Hyperbola, is the diameter passing through the two foci, and the two principal vertices of the figure. In the hyperbola it is the shortest diameter, but in the ellipse it is the longest.

VOL. I.

**Conjugate Axis**, or **Second Axis**, in the Ellipse and Hyperbola, is the diameter passing through the centre, and perpendicular to the transverse axis; and is the shortest of all the conjugate diameters.

**Axis**, of a curve line, is still more generally used for that diameter which has its ordinates at right angles to it, when that is possible. For, like as in the conic sections, any diameter bisects all its parallel ordinates, making the two parts of them on both sides of it equal; and that diameter which has such ordinates perpendicular to it, is an **Axis**: So, in curves of the second order, if any two parallel lines each meeting the curve in three points; the right line which cuts these two parallels so, that the sum of the two parts on one side of the cutting line, between it and the curve, is equal to the third part terminated by the curve on the other side, then the said line will in like manner cut all other parallels to the former two lines, viz, so that, of every one of them, the sum of the two parts, or ordinates, on one side, will be equal to the third part or ordinate on the other side. Such cutting line then is a diameter; and that diameter whose parallel ordinates are at right angles to it, when possible, is an **Axis**. And the same for other curves of still higher orders. Newton, *Enumeratio Linearum Tertii Ordinis*, sect. 2, art. 1.

**Axis, in Astronomy**. As, the **Axis of the world**, is an imaginary right line conceived to pass through the centre of the earth, and terminating at each end in the surface of the mundane sphere. About this line, as an axis, the sphere, in the Ptolomaic system, is supposed daily to revolve.

**Axis of the Earth**, is the line connecting its two poles, and about which the earth performs its diurnal rotation, from west to east. This is a part of the axis of the world, and always remains parallel to itself during the motion of the earth in its orbit about the sun, and perpendicular to the plane of the equator.

**Axis of a Planet**, is the line passing through its centre, and about which the planet revolves.—The Sun, Earth, Moon, Jupiter, Mars, and Venus, it is known from observation, move about their several axes; and the like motion is easily inferred of the other three, Mercury, Saturn, and Georgian planet.

**Axis of the Horizon, Equator, Ecliptic, Zodiac**, &c, are right lines passing through the centres of those circles, perpendicular to their planes.

**Axis of a Magnet**, or **magnetical Axis**, is a line passing through the middle of a magnet, lengthwise; in such manner, that however the magnet be divided, provided the division be made according to a plane passing through that line, the magnet will then be cut into two loadstones. And the extremities of such lines are called the poles of the stone.

**Axis in Mechanics**.—The axis of a balance, is the line upon which it moves or turns.

**Axis of Oscillation**, is a line parallel to the horizon, passing through the centre about which a pendulum vibrates, and perpendicular to the plane in which it oscillates.


**Axis in Peritrochio**, or **wheel and axle**, is one of the five mechanical powers, or simple machines; contrived chiefly for the raising of weights to a consider-

A a

able



able height, as water from a well, &c. This machine



consists of a circle AB, concentric with the base of a cylinder, and moveable together with it about its axis CD. This cylinder is called the *axis*; and the circle, the *peritrochium*; and the radii, or spokes, which are sometimes fitted immediately into the cylinder, without any circle, the *scytalæ*. About the axis winds a rope, or chain, by means of which great weights are raised by turning the wheel.—The axis in peritrochio takes place in the motion of every machine, in which a circle may be conceived as described about a fixed axis, concentric with the plane of a cylinder about which it is placed; as in Crane wheels, Mill wheels, Capstans, &c.

The chief properties of the *Axe-in-peritrochio*, are as follow:

1. If the power F applied in the direction AF a tangent to the circumference, or perpendicular to the spoke, be to a weight E, as the radius of the axis Ce is to the radius of the wheel AD, or the length of the spoke; the power will just sustain the weight; that is, the power and the weight will be in equilibrio, when they are in the reciprocal proportion of their distances from the centre.

2. When the wheel moves, with the power and weight; the velocities of their motion, and the spaces passed over by them, will be both in the same proportion as above, namely, directly proportional to their distances from the centre, and reciprocally proportional to their own weights when they are in equilibrio.

3. A power, and a weight, being given to construct an axis-in-peritrochio, by which it shall be sustained and raised. Let the axis be taken large enough to support the weight and power without breaking: then, as the weight is to the power, so make the radius of the wheel to the radius of the axis. Hence, if the power be very small in respect of the weight, the radius of the wheel will be vastly great. For example, suppose the weight 4050, and the power only 50; then the radius of the wheel will be 81 times that of the axis; which would be a very inconvenient size. But this inconvenience is provided against by increasing the number of the wheels and axes; making one to turn another, by means of teeth or pinions. And to find the effect of a number of wheels and axes, thus turning one another, multiply together, all the radii of the axes, and all the radii of the wheels, and then it will be, as the product of the former is to the product of the latter, so is the power to the weight. So, if there be 4 wheels and axes, the radius of each axis being 1 foot, and the

radius of each wheel 3 feet; then the continual product of all the wheels is  $3 \times 3 \times 3 \times 3$  or 81 feet, and that of the axis only 1; therefore the effect is as 81 to 1, or the weight is 81 times the power. And, on the contrary, if it be required to find the diameter of each of four equal wheels, by which a weight of 4050lb shall be balanced by a power of 50lb, the diameter of each axis being 1 foot: dividing 4050 by 50, the quotient is 81; extract the 4th root of 81, or twice the square root, and it will give 3, for the diameter of the four wheels sought.

*Axis of a vessel*, is that quiescent right line passing through the middle of it, perpendicular to its base, and equally distant from its sides.

*Axis in Optics*.—*Optic Axis*, or *visual axis*, is a ray passing through the centre of the eye, or falling perpendicularly on the eye.

*Axis of a lens*, or *glass*, is the axis of the solid of which the lens is a segment. Or the axis of a glass, is the line joining the two vertices or middle points of the two opposite surfaces of the glass.

*Axis of Incidence*, in Dioptrics, is the line passing through the point of incidence, perpendicularly to the refracting surface.

*Axis of Refraction*, is the line continued from the point of incidence or refraction, perpendicularly to the refracting surface, along the further medium.

*AZIMUTH*, of the sun, or star, &c, is an arch of the horizon, intercepted between the meridian of the place, and the azimuth or vertical circle passing through the sun or star; and is equal to the angle at the zenith formed by the said meridian and vertical circle. Or it is the complement to the eastern or western amplitude.—The azimuth is thus found by trigonometry;

As radius is to the tangent of the latitude,  
So is the tangent of the altitude of the sun or star,  
To the cosine of the azimuth from the south, at  
the time of the equinox.

*AZIMUTH, magnetical*, an arch of the horizon contained between the magnetical meridian, and the azimuth or vertical circle of the object; or its apparent distance from the north or south point of the compass. This is found by observing the sun, or star, &c, with an azimuth compass, when it is 10 or 15 degrees high, either before or after noon.

*AZIMUTH COMPASS*, an instrument for finding either the magnetical azimuth or amplitude of a celestial object. The description and use of this instrument, see under the article COMPASS.

*AZIMUTH DIAL*, a dial whose stile or gnomon is perpendicular to the plane of the horizon.

*AZIMUTHS*, or *Vertical Circles*, are great circles of the sphere intersecting each other in the zenith and nadir, and cutting the horizon at right angles.—These azimuths are represented by the rhumbs on common sea charts; and on the globe by the quadrant of altitude, when screwed in the zenith. On these azimuths is counted the height of the sun or stars, &c, when out of the meridian.




## B.

B A C

**BACK-STAFF**, an instrument formerly used for taking the sun's altitude at sea; being so called because the back of the observer is turned towards the sun when he makes the observation. It was sometimes called Davis's quadrant, from its inventor captain John Davis, a Welchman, and a celebrated navigator, who produced it about the year 1590.

This instrument consists of two concentric arches of box-wood, and three vanes: the arch of the longer radius is of 30 degrees, and the other 60 degrees, making between them 90 degrees, or a quadrant: also the vane A at the centre is called the *horizon-vane*, that on the arch of 60° at B the *shade-vane*, and that on the other arch at C the *sight-vane*.



*To use the Back-Staff.* The shade-vane is to be set upon the 60 arch, at an even degree of some latitude, less by 10 or 15 degrees than you judge the complement of the sun's altitude will be; also the horizon-vane being put on at A, and the sight-vane on the 30 arch FG, the observer turns his back to the sun, lifts up the instrument, and looks through the sight-vane, raising or falling the quadrant, till the shadow of the upper edge of the shade vane fall on the upper edge of the slit in the horizon-vane; and then if he can see the horizon through the said slit, the observation is exact, and the vanes are right set: But if the sea appear instead of the horizon, the sight-vane must be moved downward towards F; or if the sky appear, it must be moved upward towards G; thus trying till it comes right: the observer then examines how many degrees and minutes are cut by that edge of the sight-vane that answers to the sight hole, and to them he adds the degrees cut by the upper edge of the shade-vane; then the sum is the sun's distance from the zenith, or the

B A C

complement of the altitude; that is, of his upper limb when the upper end of the shade-vane is used in the observation, or of his lower limb when the lower part of that vane is used; therefore in the former case add 16 minutes, the sun's semidiameter, and subtract 16 minutes in the latter case, to give the zenith distance or co-altitude of the sun's centre.

Mr. Flamsteed contrived a glass lens, or double convex, to be placed in the middle of the shade-vane, which throws a small bright spot on the slit of the horizon-vane, instead of the shade; which is a great improvement, if the glass be truly made; for by this means, the instrument may be used in hazy weather, and a much more accurate observation made at all times.

**BACON (ROGER)**, an English monk of the Franciscan order, an amazing instance of genius and learning, was born near Ilchester in Somersetshire, in the year 1214. He commenced his studies at Oxford; from whence he removed to the university of Paris, which at that time was esteemed the centre of literature; and where it seems he made such progress in the sciences, that he was esteemed the glory of that university, and was there greatly caressed by several of his countrymen, particularly by Robert Grossetest or Grouthet, afterwards bishop of Lincoln, his great friend and patron. Having taken the degree of doctor, he returned to England in 1240, and took the habit of the Franciscan order, being but about 26 years of age; but according to some he became a monk before he left France. He now pursued his favourite study of experimental philosophy with unremitting ardour and assiduity. In this pursuit, in experiments, instruments, and in scarce books, he informs us he spent, in the course of 20 years, no less than 2000*l*, an amazing sum in those days, and which sum it seems was generously furnished to him by some of the heads of the university, to enable him the better to pursue his noble researches. By such extraordinary talents, and amazing progress in the sciences, which in that ignorant age were so little known to the rest of mankind, while they raised the admiration of the more intelligent few, could not fail to excite the envy of his illiterate fraternity, whose malice he farther drew upon him by the freedom with which he treated the clergy in his writings, in which he spared neither their ignorance nor their want of morals: these therefore found no difficulty in possessing the vulgar with the notion of Bacon's dealing with the devil. Under this pretence he was restrained from reading lectures; his writings were confined to his convent; and at length, in 1278, he himself was imprisoned in his cell, at 64 years of age. However, being allowed the use of his books, he still proceeded in the rational pursuit of knowledge, correcting his former labours, and writing several curious pieces.



When Bacon had been 10 years in confinement, Jerom de Ascoli, general of his order, who had condemned his doctrine, was chosen pope by the name of Nicholas IV; and being reputed a person of great abilities, and one who had turned his thoughts to philosophical studies, Bacon resolved to apply to him for his discharge; and to shew both the innocence and the usefulness of his studies, addressed to him a treatise *On the means of avoiding the infirmities of old age*. What effect this had on the pope does not appear; it did not at least produce an immediate discharge: however, towards the latter end of his reign, by the interposition of some noblemen, Bacon obtained his liberty; after which he spent the remainder of his life in the college of his order, where he died in the year 1294, at 80 years of age, and was buried in the Franciscan church. Such are the few particulars which the most diligent researches have been able to discover concerning the life of this very extraordinary man.

Bacon's printed works are, 1. *Epistola Fratris Rogeri Baconis de Secretis Operibus Artis et Naturæ, et de Nullitate Magiæ*: Paris, 1542, in 4to. Basil, 1593, in 8vo. 2. *Opus Majus*: London, 1733, in fol. published by Dr. Jebb. 3. *Thesaurus Chemicus*: Francf. 1603 and 1620. These printed works of Bacon contain a considerable number of essays, which have been considered as distinct books in the catalogue of his writings by Bale, Pitts, &c; but there remain also in different libraries several manuscripts not yet published.

By an attentive perusal of his works, the reader is astonished to find that this great luminary of the 13th century was deeply skilled in all the arts and sciences, and in many of them made the most important inventions and discoveries. He was, says Dr. Peter Shaw, beyond all comparison the greatest man of his time, and he might perhaps stand in competition with the greatest that have appeared since. It is wonderful, considering the ignorant age in which he lived, how he came by such a depth of knowledge on all subjects. His writings are composed with that elegance, conciseness and strength, and adorned with such just and exquisite observations on nature, that, among all the chemists, we do not know his equal. In his chemical writings, he attempts to shew how imperfect metals may be ripened into perfect ones; making, with Geber, mercury the common basis of all metals, and sulphur the cement.

His other physical writings shew no less genius and force of mind. In his treatise *Of the Secret Works of Art and Nature*, he shews that a person perfectly acquainted with the manner observed by nature in her operations, would be able to rival, and even to surpass her. In another piece, *Of the Nullity of Magic*, he shews with great sagacity and penetration, whence the notion of it sprung, and how weak all pretences to it are. From a perusal of his works, adds the same author, we find Bacon was no stranger to many of the capital discoveries of the present and past ages. Gunpowder he certainly knew: thunder and lightning, he tells us, may be produced by art; for that sulphur, nitre and charcoal, which when separate have no sensible effect, yet when mixed together in due proportion, and closely confined, and fired, they yield a loud report. A more precise description of gunpowder cannot be

given in words. He also mentions a sort of unextinguishable fire prepared by art: which shews he was not unacquainted with phosphorus: and that he had a notion of the rarefaction of the air, and the structure of an air-pump, is past contradiction. He was the miracle, says Dr. Freind, of the age he lived in, and the greatest genius, perhaps, for mechanical knowledge, that ever appeared in the world since Archimedes. He appears likewise to have been master of the whole science of optics: he has accurately described the uses of reading-glasses, and shewn the way of making them. Dr. Freind adds, that he also describes the camera obscura, and all sorts of glasses, which magnify or diminish any object, or bring it nearer to the eye, or remove it farther off. Bacon says himself, that he had great numbers of burning-glasses: and that there were none ever in use among the Latins, till his friend Peter de Mahara Curia applied himself to the making of them. That the telescope was not unknown to him, appears from a passage where he says, that he was able to form glasses in such a manner, with respect to our sight and the objects, that the rays shall be refracted and reflected wherever we please, so that we may see a thing under what angle we think proper, either near or at a distance, and be able to read the smallest letters at an incredible distance, and to count the dust and sand, on account of the greatness of the angle under which we see the objects; and also that we shall scarce see the greatest bodies near us, on account of the smallness of the angle under which we view them. His skill in astronomy was amazing: he discovered that error which occasioned the reformation of the calendar; one of the greatest efforts, according to Dr. Jebb, of human industry: and his plan for correcting it was followed by pope Gregory the 13th, with this variation, that Bacon would have had the correction to begin from the birth of our Saviour, whereas Gregory's amendment reaches no higher than the Nicene council.

BACON (FRANCIS), baron of Verulam, viscount of St. Albans, and lord high chancellor of England under king James I. He was born in 1560, being son of Sir Nicholas Bacon lord keeper of the great seal in the reign of queen Elizabeth, by Anne daughter of Sir Anthony Cook, eminent for her skill in the Latin and Greek languages. He gave even in his infancy tokens of what he would one day become; and queen Elizabeth had many times occasion to admire his wit and talents, and used to call him her young lord keeper. He studied the philosophy of Aristotle at Cambridge; where he made such progress in his studies, that at 16 years of age he had run through the whole circle of the liberal arts as they were then taught, and even began to perceive those imperfections in the reigning philosophy, which he afterwards so effectually exposed, and thence not only overturned that tyranny which prevented the progress of true knowledge, but laid the foundation of that free and useful philosophy which has since opened a way to so many glorious discoveries. On his leaving the university, his father sent him to France; where, before he was 19 years of age, he wrote a general view of the state of Europe: but his father dying, he was obliged suddenly to return to England; where he applied himself to the study of the common law, at Gray's-inn. His merit at length raised him to the highest



highest dignities in his profession, attorney-general, and lord high chancellor. But being of an easy and liberal disposition, his servants took advantage of that temper, and their situation under him, by accepting presents in the line of his profession. Being abandoned by the king, he was tried by the house of lords, for bribery and corruption, and by them sentenced to pay a fine of 40,000*l.* and to remain prisoner in the Tower during the king's pleasure. The king however soon after remitted the fine and punishment: but his misfortunes had given him a distaste for public affairs, and he afterwards mostly lived a retired life, closely pursuing his philosophical studies and amusements, in which time he composed the greatest part of his English and Latin works. Though even in the midst of his honours and employments he forgot not his philosophy, but in 1620 published his great work *Novum Organum*. After some years spent in his philosophical retirement, he died in 1626, being 66 years of age.

The chancellor Bacon is one of those extraordinary geniuses who have contributed the most to the advancement of the sciences. He clearly perceived the imperfection of the school philosophy, and he pointed out the only means of reforming it, by proceeding in the opposite way, from experiments to the discovery of the laws of nature. Addison has said of him, That he had the sound, distinct, comprehensive knowledge of Aristotle, with all the beautiful light graces and embellishments of Cicero. Mr. Walpole calls him the *Prophet of Arts*, which Newton was afterwards to reveal; and adds, that his genius and his works will be universally admired as long as science exists. He did not yet, said another great man, understand nature, but he knew and pointed out all the ways that lead to her. He very early despised all that the universities called philosophy; and he did every thing in his power that they should not disgrace her by their quiddities, their horrors of a vacuum, their substantial forms, and such like impertinencies.

He composed two works for perfecting the sciences. The former *On the Dignity and Augmentation of the Sciences*. He here shews the state in which they then were, and points out what remains to be discovered for perfecting them; condemning the unnatural way of Aristotle, in reversing the natural order of things. He here also proposes his celebrated division of the sciences.

To remedy the faults of the common logic, Bacon composed his second work, the *New Organ of Sciences*, above-mentioned. He here teaches a new logic, the chief end of which is to shew how to make a good inference, as that of Aristotle's is to make a syllogism. Bacon was 18 years in composing this work, and he always esteemed it as the chief of his compositions.

The pains which Bacon bestowed upon all the sciences in general, prevented him from making any considerable applications to any one in particular: and as he knew that natural philosophy is the foundation of all the other sciences, he chiefly endeavoured to give perfection to it. He therefore proposed to establish a new system of physics, rejecting the doubtful principles of the ancients. For this purpose he took the resolution of composing every month a treatise on some

branch of physics; he accordingly began with that of the winds; then he gave that of heat; next that of motion; and lastly that of life and death. But as it was impossible that one man alone could so compose the whole circle of sciences with the same precision, after having given these patterns, to serve as a model to those who might choose to labour upon his principles, he contented himself with tracing in a few words the design of four other tracts, and with furnishing the materials, in his *Silva Silvarum*, where he has amassed a vast number of experiments, to serve as a foundation for his new physics. In fact, no one before Bacon understood any thing of the experimental philosophy; and of all the physical experiments which have been made since his time, there is scarcely one that is not pointed out in his works.

This great precursor of philosophy was also an elegant writer, an historian, and a wit. His moral essays are valuable, but are formed more to instruct than to please. There are excellent things too in his work *On the Wisdom of the Ancients*, in which he has moralized the fables which formed the theology of the Greeks and Romans. He wrote also *The History of Henry the VIIIth king of England*, by which it appears that he was not less a great politician than a great philosopher.

Bacon had also some other writings, published at different times; the whole of which were collected together, and published at Frankfort, in the year 1665, in a large folio volume, with an introduction concerning his life and writings. Another edition of his works was published at London in 1740; the enumeration of which is as below:

1. De Dignitate et Augmentis Scientiarum.
2. Novum Organum Scientiarum, sive Judicia vera de Interpretatione Naturæ; cum Parasceve ad Historiam Naturalem & Experimentalem.
3. Phænomena Universi, sive Historia Naturalis & Experimentalis de Ventis; Historia Denſi & Rari; Historia Gravis & Levis; Historia Sympathiæ & Antipathiæ Rerum; Historia Sulphuris, Mercurii, & Salis; Historia Vitæ & Mortis; Historia Naturalis & Experimentalis de Forma Calidi; De Motus, sive Virtutis activæ variis speciebus; Ratio inveniendi causas Fluxus & Refluxus Maris; &c. &c.
4. Silva Silvarum, sive Historia Naturalis.
5. Novus Atlas.
6. Historia Regni Henrici vii Angliæ Regis.
7. Sermones Fideles, Ethici, Politici, Oeconomici.
8. De Sapientia Veterum.

BACULÉ, in *Fortification*, a kind of portcullis, or gate, made like a pit-fall, with a counterpoise, and supported with two great stakes. It is usually made before the *corps de garde*, not far from the gate of a place.

BACULOMETRY, the art of measuring either accessible or inaccessible lines, by the help of *baculi*, staves, or rods. Schwenter has explained this art in his *Geometria Practica*; and the rules of it are delivered by Wolfius, in his *Elements*: Ozanam also gives an illustration of the principles of Baculometry.

BAILLY (JEAN SYLVAIN), a celebrated French astronomer, historiographer, and politician, was born at Paris the 15th of September 1736, and has figured as



one of the greatest men of the age, being a member of several academies, and an excellent scholar and writer. He enjoyed for several years the office of keeper of the king's pictures at Paris. He published, in 1766, a volume in 4to, *An Essay on the Theory of Jupiter's Satellites*, preceded by a History of the Astronomy of these Satellites. In the Journal Encyclopédique for May and July 1773, he addressed a letter to M. Bernoulli, Astronomer Royal at Berlin, upon some discoveries relative to these satellites, which he had disputed. In 1768 he published the Eulogy of Leibnitz, which obtained the prize at the Academy of Berlin, where it was printed. In 1770 he printed at Paris, in 8vo, the Eulogies of Charles the Vth, of de la Caille, of Leibnitz, and of Corneille. This last had the second prize at the Academy of Rouen, and that of Moliere had the same honour at the French Academy.

M. Bailly was admitted into the Academy as Adjunct the 29th of January 1763, and as Associate the 14th of July 1770.—In 1775 came out at Paris, in 4to, his *History of the Ancient Astronomy*, in 1 volume: In 1779 the *History of Modern Astronomy* in 2 volumes: and in 1787 the *History of the Indian and Oriental Astronomy*, being the 2d vol. of the Ancient Astronomy.

M. Bailly's memoirs published in the volumes of the Academy, are as follow:

Memoir upon the Theory of the Comet of 1759.

Memoir upon the Epoches of the Moon's motions at the end of the last century.

First, second, and third Memoirs on the Theory of Jupiter's Satellites, 1763.

Memoir on the Comet of 1762: vol. for 1763.

Astronomical Observations, made at Nollon: 1764.

On the Sun's Eclipse of the 1st of April, 1764.

On the Longitude of Polling; 1764.

Observations made at the Louvre from 1760 to 1764: 1765.

On the cause of the Variation of the Inclination of the Orbit of Jupiter's second Satellite; 1765.

On the Motion of the Nodes, and on the Variation of the Inclination of Jupiter's Satellites; 1766.

On the Theory of Jupiter's Satellites, published by M. Bailly, and according to the Tables of their Motions and of those of Jupiter, published by M. Jaurat; 1766.

Observations on the Opposition of the Sun and Jupiter; 1768.

On the Equation of Jupiter's Centre, and on some other Elements of the Theory of that Planet; 1768.

On the Transit of Venus over the Sun, the 3d of June 1769; and on the Solar Eclipse the 4th of June the same year; 1769.

In the beginning of the revolution in France, in 1789, M. Bailly took an active part in that business, and was so popular and generally esteemed, that he was chosen the first president of the States General, and of the National Assembly, and was afterwards for two years together the Mayor of Paris; in both which offices he conducted himself with great spirit, and gave general satisfaction.

He soon afterward however experienced a sad reverse of fortune; being accused by the ruling party of favouring the king, he was condemned for incivism and wanting to overturn the Republic, and died by

the Guillotine at Paris on the eleventh day of November, 1793, at 57 years of age.

BAINBRIDGE (JOHN), an eminent physician, astronomer, and mathematician. He was born in 1582, at Ashby de la Zouch, Leicestershire. He studied at Cambridge, where having taken his degrees of Bachelor and Master of Arts, he returned to Leicestershire, where for some years he kept a grammar-school, and at the same time practised physic; employing his leisure hours in studying mathematics, especially astronomy, which had been his favourite science from his earliest years. By the advice of his friends, he removed to London, to better his condition, and improve himself with the conversation of learned men there; and here he was admitted a fellow of the college of physicians. His description of the comet, which appeared in 1618, greatly raised his character, and procured him the acquaintance of Sir Henry Savile, who, in 1619, appointed him his first professor of astronomy at Oxford. On his removal to this university, he entered a master commoner of Merton college; the master and fellows of which appointed him junior reader of Linacer's lecture in 1631, and superior reader in 1635. As he resolved to publish correct editions of the ancient astronomers, agreeably to the statutes of the founder of his professorship, that he might acquaint himself with the discoveries of the Arabian astronomers, he began the study of the Arabic language when he was above 40 years of age. Before completing that work however he died, in the year 1643, at 61 years of age.

Dr. Bainbridge wrote many works, but most of them have never been published; those that were published, were the three following, viz:

1. An Astronomical Description of the late Comet, from the 18th of November 1618, to the 16th of December following; 4to, London, 1619.—This piece was only a specimen of a larger work, which the author intended to publish in Latin, under the title of *Cometographia*.

2. *Procli Sphæra, Ptolomæi de Hypothesibus Planetarum liber singularis*. To which he added Ptolomy's Canon Regnorum. He collated these pieces with ancient manuscripts, and gave a Latin version of them, illustrated with figures: printed in 4to, 1620.

3. *Canicularia*. A treatise concerning the Dog-star, and the Canicular Days: published at Oxford in 1648, by Mr. Greaves, together with a demonstration of the heliacal rising of Sirius, the dog-star, for the parallel of Lower Egypt. Dr. Bainbridge undertook this work at the request of archbishop Usher, but he left it imperfect; being prevented by the breaking out of the civil war, or by death.

There were also several dissertations of his prepared for and committed to the press the year after his death, but the edition of them was never completed. The titles of them are as follow:

1st, *Antiprognosticon, in quo Mavlixi Astrologiæ, Cœlestium Domorum, et Triplicitatum Commentis, magnisque Saturni et Jovis (cujusmodi anno 1623, et 1643, contigerunt, et vicesimo fere quoque deinceps anno, ratis naturæ legibus, recurrent) Conjunctionibus innixæ, Vanitas breviter detegitur*.

2nd, *De Meridionorum five Longitudinum Differentiis inveniendis Dissertatio*.

3d, *De*



3d, De Stella Veneris Diatriba.

Beside the foregoing, there were several other tracts, never printed, but left by his will to archbishop Usher; among whose manuscripts they are preserved in the library of the college of Dublin. Among which are the following: 1. A Theory of the Sun. 2. A Theory of the Moon. 3. A Discourse concerning the Quantity of the Year. 4. Two volumes of Astronomical Observations. 5. Nine or ten volumes of Miscellaneous Papers relating to Mathematical subjects.

BAKER (THOMAS), a mathematician of some eminence, was born at Ilton in Somersetshire, in 1625. He entered upon his studies at Oxford in 1640, where he remained seven years. He was afterwards appointed vicar of Bishop's-Nymmet in Devonshire, where he lived a studious and retired life for many years, chiefly pursuing the mathematical sciences; of which he gave a proof of his critical knowledge, in the book he published, concerning the general construction of bi-quadratic equations, by a parabola and a circle; the title of which book at full length is, "The Geometrical Key; or the Gate of Equations unlocked: or a new discovery of the Construction of all Equations, howsoever affected, not exceeding the 4th degree, viz, of Linears, Quadratics, Cubics, Biquadratics, and the finding of all their Roots, as well False as True, without the use of Mesolabe, Trisection of Angles, without Reduction, Depression, or any other previous preparations of equations by a circle, and any (and that one only) Parabole, &c": 1684, 4to, in English and Latin.

There is some account of this work in the Philos. Transf. an. 1684. And a little before his death, the Royal Society sent him some mathematical queries; to which he returned such satisfactory answers, as procured the present of a medal, with an inscription full of honour and respect.—Mr. Baker died at Bishop's-Nymmet, 1690, in the 65th year of his age.

BAKER (HENRY), an ingenious and diligent naturalist, was born in London about the beginning of the 18th century. He was brought up under an eminent bookseller, but being of a philosophical turn of mind, he quitted that line of business soon after the expiration of his apprenticeship, and took to the employment of teaching deaf and dumb persons to speak and write &c, in which occupation, in the course of his life he acquired a handsome fortune. For his amusement he cultivated various natural and philosophical sciences, particularly botany, natural history, and microscopical subjects, in which he especially excelled, having, in the year 1744, obtained the Royal Society's gold medal, for his microscopical experiments on the crystallizations and configurations of saline particles. He had various papers published in the Philos. Transf. of the Royal Society, of which he was a worthy member, as well as of the Societies of Antiquaries, and of Arts. He was author of many pieces, on various subjects, the principal of which were, his Treatise on the Water Polype, and two Treatises on the Microscope, viz, *The Microscope made easy*, and *Employment for the Microscope*, which have gone through several editions.

Mr. Baker married Sophia, youngest daughter of

the celebrated Daniel Defoe, by whom he had two sons, who both died before him. He terminated an honourable and useful life, at his apartments in the Strand on the 25th of November 1774, being then upwards of 70 years of age.

BAKER'S CENTRAL RULE, for the Construction of Equations, is a method of constructing all equations, not exceeding the 4th degree, by means of a given parabola and a circle, without any previous reduction of them, or first taking away their second term. See CENTRAL Rule.

BALANCE, one of the six simple powers in mechanics, chiefly used in determining the equality or difference of weight in heavy bodies, and consequently their masses or quantities of matter.

The balance is of two kinds, the *ancient* and *modern*. The ancient or Roman, called also *Statera Romana*, or *Steelyard*, consists of a lever or beam, moveable on a centre, and suspended near one of its extremities. The bodies to be weighed are suspended from the shorter end, and their weight is shewn by the division marked on the beam, where the power or constant weight, which is moveable along the lever, keeps the steelyard in equilibrio. This balance is still in common use for weighing heavy bodies.

The modern balance, now commonly used, consists of a lever or beam suspended exactly in the middle, and having scales suspended from the two extremities, to receive the weights to be weighed.

In either case the lever is called the *jugum* or the *beam*, and its two halves on each side the axis, the *brachia* or *arms*; also the line on which the beam turns, or which divides it in two, is called the *axis*; and when considered with regard to the length of the brachia, is esteemed only a point, and called the *centre of the balance*, or *centre of motion*: the extremities where the weights are applied, are the *points of application* or *suspension*; the handle by which the balance is held, or by which the whole apparatus is suspended, is called *trutina*; and the slender part perpendicular to the beam, by which is determined either the equilibrium or preponderancy of bodies, is called the *tongue of the balance*.


From these descriptions we easily gather the characteristic distinction between the Roman balance and the common one, viz, that in the Roman balance, there is one constant weight used as a counterpoise, the point where it is suspended being varied; but, on the contrary, in the common balance or scales, the points of suspension remain the same, and the counterpoise is varied. The principle of both of them may be easily understood from the general properties of the lever, and the following observations.

The beam ABC, the principal part of the balance, is a lever of the first kind; but instead of resting on a fulcrum, it is suspended by a handle, &c, fastened to its centre of motion B: and hence the mechanism of the balance depends on the same theorems as that of the lever. Consequently as the distance between the centre of motion and the place of the unknown weight, is to the distance between the same centre and the place of the known weight, so is the latter weight, to the former. So that the unknown weight is discovered by means of




of the known one, and their distances from the common centre of motion; viz, if the distances from the centre be equal, then the two weights will be equal also, as in the common balance; but if the distances be unequal, then the weights will also be unequal, and in the very same proportion, alternately, the less weight having so much the greater distance, as in the steelyard.

*The Common Balance or Scales.*



The two brachia AB, BC, should be exactly equal in length, and in weight also when their scales D and E are fixed on their ends; the beam should hang exactly level or horizontal in the case of an equipoise; and for this purpose the centre of gravity of the whole should fall a little below the centre of motion, and but a little, that the balance be sufficiently sensible to the least variation of weight: the friction on the centre should also be as small as possible.


*The Steel Yard.*



Having made a proper bar of steel AB, tapering at the longer end, and very strong at the other, suspend it by a centre C near the shorter or thicker end, so that it may exactly balance itself in equilibrio, and prepare a constant weight I to weigh with: then hang on any weight, as one pound for instance, at the shorter arm, and slide the constant weight backwards and forwards upon the longer arm, till it be just in equilibrio with the former; and there make a notch and number 1, for the place of 1 pound: take off the 1lb, and hang a two pound weight in its stead at the shorter arm; then slide the constant weight back on the longer arm, till the whole come again into equilibrio, making a notch at the place of the constant weight and the number 2, for the place of 2lb. Proceed in the same manner for all other weights 3, 4, 5, &c; as also for the intermediate halves and quarters, &c, if it be necessary; always sus-

pending the variable weights at the end of the shorter arm, shifting the constant weight so as to balance them, and marking and numbering the places on the longer arm where the constant weight always makes a counterpoise. The use of the Steelyard is hence very evident: the thing whose weight is required being suspended by a hook at the short end, move the constant weight backwards and forwards on the longer arm, till the beam is balanced horizontally: then look what notch the constant weight is placed at, and its number will shew the weight of the body that was required. DC is the handle and tongue; F the centre of motion; EG a scale sometimes hung on at the end by the hook H.

*The Bent-Lever Balance.*



This instrument operates by a fixed weight, C, increasing in power as it ascends along the arc FG of a circle, and pointing by an index to the number or division of the arc which denotes the weight of any body put into the scale at E. And thus one constant weight serves to weigh all others, by only varying the position of the arms of the balance, instead of varying the places or points of suspension in the arms themselves.

*The Deceitful Balance.* This operates in the same manner as the steelyard, and cheats or deceives by having one arm a little longer than the other; though the deception is not perceived, because the shorter arm is made somewhat heavier, so as to compensate for its shortness, by which means the beam of the balance, when no weights are in the scales, hangs horizontal in equilibrio. The consequence of this construction is, that any commodity put in the scale of the longer arm, requires a greater weight in the other scale to balance it; and so the body is fallaciously accounted heavier than it really is. But the trick will easily be detected by making the body and the weight change places, removing them to the opposite scales, when the weight will immediately be seen to preponderate.

*Affay-Balance.* This is a very nice balance, used in determining the exact weights of very small bodies. Its structure is but little different from the common sort; except



except that it is made of the best and hardest steel, and made to turn with the smallest weight.

*Hydrostatical Balance.* This is an instrument for determining the specific gravity of bodies. See *HYDRO-STATICAL*, and *SPECIFIC Gravity*.

*BALANCE*, in Astronomy, the same as *Libra*.

*BALANCE of a Clock or Watch*, is that part which, by its motion, regulates and determines the beats. The circular part of it is called the *rim*, and its spindle the *verge*; there belong to it also two pallets or nuts, that play in the fangs of the crown-wheel: in pocket watches, that strong stud in which the lower pivot of the verge plays, and in the middle of which one pivot of the crown-wheel runs, is called the *potence*: the wrought piece which covers the balance, and in which the upper pivot of the balance plays, is the *cock*; and the small spring in the new pocket watches, is called the *regulator*.

*BALCONY*, a projecture in the front of a house, or other building, commonly supported by pillars or consoles, and encompassed by a balustrade.

*BALL*, any spherical, globular, or round body.

*BALL*, in the military art, signifies all sorts of bullets for fire arms, from the pistol up to the largest cannon. Cannon balls are made of cast iron; but the musket and pistol balls of lead, as these are both heavier under the same bulk, and do not furrow the barrels of the pieces.

*BALL of a Pendulum*, is the weight at the bottom of it; and is sometimes, especially in shorter pendulums, called the *bob*.

*BALLS of Fire*, in Meteorology. See *FIRE balls*.

*BALLS*, in Electricity, invented by Mr. Canton, are two pieces of cork, or pith of elder tree, nicely turned in a lathe to the size of a small pea, and suspended by fine linen threads. They are used as electrometers, and are of excellent use to discover small degrees of electricity; and to observe its changes from positive to negative, or the reverse; as also to estimate the force of a shock before the discharge, so that the operator shall always be able to tell very nearly before hand what the explosion will be, by knowing how high he has charged his jars.

*BALLISTA*, a military engine much used by the ancients for throwing stones, darts, and javelins; and somewhat resembling our cross-bows, but much larger and stronger. The word is Latin, signifying a cross-bow; but is derived from the Greek *βαλλω*, to shoot, or throw.

Vegetius informs us, that the ballista discharged darts with such violence and rapidity, that nothing could resist their force: and Athenæus adds, that Agi-stratus made one of little more than 2 feet in length, that shot darts 500 paces. Authors have often confounded the ballista with the catapulta, attributing to the one what belongs to the other. According to Vitruvius, the ballista was made after divers manners, though all were used to the same purpose: one sort was framed with levers and bars; another with pulleys; some with a crane; and others again with a toothed wheel. Marcellinus describes the ballista thus; A round iron cylinder is fastened between two planks, from which reaches a hollow square beam placed cross-wise, fastened with cords, to which are added screws. At one end of this stands the engineer, who puts a wooden shaft, or arrow, with a large head, into the cavity of the beam; this done, two men bend the engine, by drawing some

wheels; when the top of the head is drawn to the utmost end of the cords, the shaft is driven out of the ballista, &c.

The ballista is ranked by the ancients in the sling kind, and its structure and effect reduced to the principles of that instrument; whence it is called by Hero and others, *funda*, and *fundibulus*. Gunther calls it *Balearica machina*, as a sling peculiar to the Balearic islands.—Perrault, in his notes on Vitruvius, gives a new contrivance of a like engine for throwing bombs without powder.

Fig. 1, Plate v, represents the ballista used in sieges, according to Folard: where 2, 2, denote the base of the ballista; 3, 4, upright beams; 5, 6, transverse beams; 7, 7, the two capitals in the upper transverse beam, (the lower transverse beam has also two similar capitals, which cannot be seen in this transverse figure); 9, 9, two posts or supports for strengthening the transverse beams; 10, 10, two skains of cords fastened to the capitals; 11, 11, two arms inserted between the two strands, or parts of the skains; 12, a cord fastened to the two arms; 13, darts which are shot by the ballista; 14, 14, curves in the upright beams, and in the concavity of which cushions are fastened, in order to break the force of the arms, which strike against them with great force when the dart is discharged; 16, the arbor of the machine, in which a straight groove or canal is formed to receive the darts, in order to their being shot by the ballista; 17, the nut of the trigger; 18, the roll or windlass, about which the cord is wound; 19, a hook, by which the cord is drawn towards the centre, and the ballista cocked; 20, a stage or table on which the arbor is in part sustained.

*BALLISTA*, in Practical Geometry, the same as the geometrical cross, called also *Jacob's Staff*. See *CROSS-Staff*.

*BALLISTIC PENDULUM*, an ingenious machine invented by Benjamin Robins, for ascertaining the velocity of military projectiles, and consequently the force of fired gun-powder. It consists of a large block of wood, annexed to the end of a strong iron stem, having a cross steel axis at the other end, placed horizontally, about which the whole vibrates together like the pendulum of a clock. The machine being at rest, a piece of ordnance is pointed straight towards the wooden block, or ball of this pendulum, and then discharged: the consequence is this; the ball discharged from the gun strikes and enters the block, and causes the pendulum to vibrate more or less according to the velocity of the projectile, or the force of the blow; and by observing the extent of the vibration, the force of that blow becomes known, or the greatest velocity with which the block is moved out of its place, and consequently the velocity of the projectile itself which struck the blow and urged the pendulum.

The more minute and particular description may be seen in my Tracts, vol. 1, where are given all the rules for using it, and for computing the velocities, with a multitude of accurate experiments performed with cannon balls, by means of which the most useful and important conclusions have been deduced in military projectiles and the nature of physics. I have also since that publication, made many other experiments of the same kind, by discharging cannon balls at various distances from the block; from which have resulted the



discovery of a complete series of the resistances of the air to balls passing through it with all degrees of velocity, from 0 up to 2000 feet in a second of time.

Other writers on this subject are Euler, Antoni, Le Roy, Darcy, &c. See also Robins's Mathematical Tracts.

**BALLISTICS**, is used by some for projectiles, or the art of throwing heavy bodies. Merfennus has published a treatise on the projection of bodies, under this title.

**BALLOON**, or **BALLON**, in a general sense, signifies any spherical hollow body. Thus, with chemists, it denotes a round short-necked vessel, used to receive what is distilled by means of fire: in architecture, a ball or globe on the top of a pillar, &c: and among engineers, a kind of bomb made of pasteboard, and played off in fireworks, in imitation of a real iron bomb-shell.

*Air-BALLOON*. See **AEROSTATION** and *Air-Balloon*.

**BALLUSTER**, a small kind of column or pillar, used for ballustrades.

**BALLUSTRADE**, a series or row of ballusters, joined by a rail; serving for a rest to the arms, or as a fence or inclosure to balconies, altars, staircases, &c.

**BAND**, in Architecture, denotes any flat low member, or moulding, that is broad, but not very deep. The word *lace* sometimes means the same thing.

**BANQUET**, or **BANQUETTE**, in Fortification, a little foot-bank, or elevation of earth, forming a path along the inside of a parapet, for the soldiers to stand upon to discover the counterescarp, or to fire on the enemy, in the moat, or in the covert way. It is commonly about 3 feet wide, and a foot and a half high.

**BARLOWE** (**WILLIAM**), an eminent mathematician and divine, in the 16th century. He was born in Pembroke-shire, his father (William Barlowe) being then bishop of St. David's. In 1560 he was entered commoner of Baliol college in Oxford; and in 1564, having taken a degree in arts, he left the university, and went to sea; but in what capacity is uncertain: however he thence acquired considerable knowledge in the art of navigation, as his writings afterwards shewed. About the year 1573, he entered into orders, and became prebendary of Winchester, and rector of Easton, near that city. In 1588 he was made prebendary of Litchfield, which he exchanged for the office of treasurer of that church. He afterwards was appointed chaplain to prince Henry, eldest son of king James the first; and, in 1614, archdeacon of Salisbury. Barlowe was remarkable, especially for having been the first writer on the nature and properties of the loadstone, 20 years before Gilbert published his book on that subject. He was the first who made the inclinatory instrument transparent, and to be used with a glass on both sides. It was he also who suspended it in a compass-box, where, with 2 ounces weight, it was made fit for use at sea. He also found out the difference between iron and steel, and their tempers for magnetical uses. He likewise discovered the proper way of touching magnetical needles; and of piecing and cementing of loadstones; and also why a loadstone, being double-capped, must take up so great a weight.

Barlowe died in the year 1625.—His works are as follow:

1. *The Navigator's Supply, containing many things of principal importance belonging to Navigation, and use of*

*diverse Instruments framed chiefly for that purpose.* Lond. 1597, 4to; dedicated to Robert earl of Essex.

2. *Magnetical Advertisement, or Diverse Pertinent Observations and improved Experiments concerning the nature and properties of the Loadstone.* Lond. 1616, 4to.

3. *A Brief Discovery of the idle Animadversions of Mark Ridley, M. D. upon a treatise entitled Magnetical Advertisements.* Lond. 1618, 4to.

In the first of these pieces, Barlowe gave a demonstration of Wright's or Mercator's division of the meridian line, as communicated by a friend; observing that "This manner of carte has been publickly extant in print these thirtie yeares at least [he should have said 28 only], but a cloude (as it were) and thicke miste of ignorance doth keepe it hitherto concealed: And so much the more, because some who were reckoned for men of good knowledge, have by glauncing speeches (but never by any one reason of moment) gone about what they could to disgrace it."

This work of Barlowe's contains descriptions of several instruments for the use of Navigation, the principal of which is an Azimuth Compass, with two upright sights; and as the author was very curious in making experiments on the loadstone, he treats well and fully upon the Sea-Compass. And he treated still farther on the same instrument in his second work, the Magnetical Advertisement.

**BAROMETER**, an instrument for measuring the weight or pressure of the atmosphere; and by that means the variations in the state of the air, foretelling the changes in the weather, and measuring heights or depths, &c.

This instrument is founded on what is called the Torricellian experiment, related below, and commonly consists of a glass tube, open at one end; which being first filled with quicksilver, and then inverted with the open end downwards into a basin of the same, the mercury descends in the tube till it remains at about the height of 29 or 30 inches, according to the weight or pressure of the atmosphere at the time, which is just equal to the weight of that column of the quicksilver. Hence it follows that, if by any means the pressure of the air be altered, it will be indicated by the rising or falling of the mercury in the tube; or if the barometer be carried to a higher station, the quicksilver will descend lower in the tube, but when carried to a lower place, it will rise higher in the tube, according to the difference in elevation between the two places.

*History of the Barometer.*—About the beginning of the last century, when the doctrine of a plenum was in vogue, it was a common opinion among philosophers, that the ascent of water in pumps was owing to what they called nature's abhorrence of a vacuum; and that thus fluids might be raised by suction to any height whatever. But an accident having just discovered, that water could not be raised in a pump unless the sucker reached to within 33 feet of the water in the well, it was conjectured by Galileo, who flourished about that time, that there might be some other cause of the ascent of water in pumps, or at least that this abhorrence was limited to the finite height of 33 feet. Being unable to satisfy himself on this head, he recommended the consideration of the difficulty to Torricelli, who had been his disciple. After some time Torricelli fell upon the suspicion that the pressure of the atmosphere was the cause of the ascent of water in pumps; that a column



of water 33 feet high was a just counterpoise to a column of air, of the same base, and which extended up to the top of the atmosphere; and that this was the true reason why the water did not follow the sucker any farther. And this suspicion was soon after confirmed by various experiments. Torricelli considered, that if a column of water 33 feet high were a counterpoise to a whole column of the atmosphere, then a column of mercury of about 2 feet and a half high would also be a counterpoise to it, since quicksilver is near 14 times heavier than water, and so the 14th part of the height, or near 2 feet and a half, would be as heavy as the column of water. This reasoning was soon verified; for having filled a glass tube with quicksilver, and inverted it into a basin of the same, the mercury presently descended till its height, above that in the basin, was about two feet and a half, just as he expected. And this is what has, from him, been called the Torricellian experiment.

The new opinion, with this confirmation of it, was readily acquiesced in by most of the philosophers, who repeated the experiment in various ways. Others however still adhered to the old doctrine, and raised several pretended objections against the new one; such as that there was a film or imperceptible *rope of mercury*, extended through the upper part of the tube, which suspended the column of mercury, and kept it from falling into that in the basin. This and other objections were however soon overcome by additional confirmations of the true doctrine, particularly by varying the elevation of the place. It was hinted by Descartes and Pascal, that if the mercury be sustained in the tube by the pressure of the atmosphere, by carrying it to a higher situation, it would descend lower in the tube, having a shorter column of the atmosphere to sustain it, and vice versa. And Pascal engaged his brother-in-law, M. Perier, to try that experiment for him, being more conveniently situated for that purpose than he was at Paris. This he accordingly executed, by observing the height of the quicksilver in the tube, first at the bottom of a mountain in Auvergne, and then at several stations, or different altitudes, in ascending, by which it was found that the mercury fell lower and lower all the way to the top of the mountain; and so confirming the truth of the doctrine relating to the universal pressure of the atmosphere, and the consequent suspension of the mercury in the tube of the barometer. Thus, by the united endeavours of Torricelli, Descartes, Pascal, Merfenne, Huygens, and others, the cause of the suspension of the quicksilver in the tube of the barometer, became pretty generally established.

It was some time however after this general consent, before it was known that the pressure of the air was various at different times, in the same place. This could not however remain long unknown, as the frequent measuring of the column of mercury, must soon shew its variations in altitude; and experience and observation would presently shew that those variations in the mercurial column, were always succeeded by certain changes in the weather, as to rain, wind, frosts, &c. Hence this instrument soon came into use as the means of foretelling the changes of the weather; and on this account it obtained the name of the *weather-glass*, as it did that of *barometer* from its being the measure of the weight or

pressure of the air. We may now proceed to take a view of its various forms and uses.

*The Common Barometer.* This is represented at fig. 1, plate iv, such as it was invented by Torricelli. AB is a glass tube, of  $\frac{1}{4}$ , or  $\frac{1}{3}$ , or  $\frac{1}{2}$  inch wide, the more the better, and about 34 inches long, being close at the top A, and the open end B immersed in a basin of quicksilver CD, which is the better the wider it is. To fill this, or any other barometer; take a clean new glass tube, of the dimensions as above, and pour into it well purified quicksilver, with a small funnel either of glass or paper, in a fine continued stream, till it wants about half an inch or an inch of being full; then stopping it close with the finger, invert it slowly, and the air in the empty part will ascend gradually to the other end, collecting into itself such other small air bubbles as unavoidably get into the tube among the mercury, in filling it with the funnel: and thus continue to invert it several times, turning the two ends alternately upwards, till all the air bubbles are collected, and brought up to the open end of the tube, and when the part filled shall appear, without speck, like a fine polished steel rod. This done, pour in a little more quicksilver, to fill the empty part quite full, and so exclude all air from the tube: then, stopping the orifice again with the finger, invert the tube, and immerse the finger and end, thus stopped, into a basin of like purified quicksilver; in this position withdraw the finger, so shall the mercury descend in the tube to some place as E, between 28 and 31 inches above that in the basin at F, as these are the limits between which it always stands in this country on the common surface of the earth. Then measure, from the surface of the quicksilver in the basin at F, 28 inches to G, and 31 inches to H, dividing the space between them into inches and tenths, which are marked on a scale placed against the side of the tube; and the tenths are subdivided into hundredth parts of an inch by a sliding index carrying a vernier or nonius. These 3 inches, between 28 and 31, so divided, will answer for all the ordinary purposes of a stationary or chamber barometer; but for experiments on altitudes and depths, it is proper to have the divisions carried on a little higher up, and a great deal lower down. In the proper filling and otherwise fitting up of the barometer, several circumstances are to be carefully noted; as, that the bore of the tube be pretty wide, to allow the freer motion of the quicksilver, without being impeded by an adhesion to the sides; that the basin below it be also pretty large, in order that the surface of the mercury at F may not sensibly rise or fall with that in the tube; that the bottom of the tube be cut off rather obliquely, that when it rests on the bottom of the basin there may be a free passage for the quicksilver; and that, to have the quicksilver very pure, it is best to boil it in the tube, which will expel all the air from it. This barometer is commonly fitted up in a neat mahogany case, together with a thermometer and hygrometer, as represented in plate 4, fig. 13.

As the scale of variation is but small, being included within 3 inches in the common barometer, several contrivances have been devised to enlarge the scale, or to render the motion of the quicksilver more sensible.

Descartes first suggested a method of increasing the sensibility, which was executed by Huygens. This



was effected by making the barometrical tube end in a pretty large cylindrical vessel at top, into which was inserted also the lower or open end of a much finer tube than the former, which was partly filled with water, to give little obstruction by its weight to the motion of the mercury, while it moved through a pretty long space of the very fine tube by a small variation of the mercury below it, and so rendered the small changes in the state of the air very sensible. But the inconvenience was this, that the air contained in the water gradually disengaged itself, and escaped through into the vacuum in the top of the small tube, till it was collected in a body there, and by its elasticity preventing the free rise of the fluids in the tubes, spoiled the instrument as a barometer. And this, it may be observed by-the-bye, is the reason why a water barometer cannot succeed. This barometer is here represented in fig. 2, where CD is the vessel, in which are united the upper or small water tube AC, with the lower or mercurial one CB.

To remedy this inconvenience, Huygens thought of placing the mercury at top, and the water at bottom, which he thus contrived. ADG (fig. 3) is a bent tube hermetically sealed at A, but open at G, of about one line in diameter, and passing through the two equal cylindrical vessels BC, EF, which are about 20 inches apart, and of 15 lines diameter, their length being 10. The mercury being put into the tube, will stand between the middle of the vessels EF and BC, the remaining space to A being void both of air and mercury. Lastly, common water, tinged with a 6th part of aqua regis, to prevent its freezing, is poured into the tube FG, till it rises a foot above the mercury in DF. To prevent the water from evaporating, a drop of oil of sweet almonds floats on the top of it. But the column of water will be sensibly affected by heat and cold, which spoils the accuracy of the instrument. For which reason other contrivances have been made, as below.

*The Horizontal or Rectangular Barometer*, fig. 4, was invented by J. Bernoulli and Cassini; where AB is a pretty wide cylindrical part at the top of the tube, which tube is bent at right angles at C, the lower part of it CD being turned into the horizontal direction, and close above at A, but open at the lower end D, where however the mercury cannot run out, being there opposed by the pressure of the atmosphere. This and the foregoing contrivance of Huygens are founded on the theorem in hydrostatics, that fluids of the same base press according to their perpendicular altitude, not according to the quantity of their matter; so that the same pressure of the atmosphere sustains the quicksilver that fills the tube ACD, and the cistern B, as would support the mercury in the tube alone. Hence, having fixed upon the size of the scale, as suppose the extent of 12 inches, instead of the 3, in the common barometer from 28 to 31, that is 4 times as long; then the area of a section of the cylinder AB must be 4 times that of the tube, and consequently its diameter double, since the areas of circles are as the squares of their diameters: then for every natural variation of an inch in the cylinder AB, there will be a variation of four inches in the tube CD.—But on account of the attrition of the mercury against the sides of the glass, and the great momentum from the quick motion in CD, the quicksilver is apt to break, and the rise and fall is no longer equa-

ble; and besides, the mercury is apt to be thrown out of the orifice at D by sudden motions of the machine.

*The Diagonal Barometer* of Sir Samuel Moreland, fig. 5, is another method of enlarging the natural scale of three inches perpendicular, or CD, by extending it to any length BC in an oblique direction. This is liable in some degree to the same inconvenience, from friction and breaking, as the horizontal one; and hence it is found that the diagonal part BC cannot properly be bent from the perpendicular more than in an angle of  $45^\circ$ , which only increases the scale nearly in the proportion of 7 to 5.

*Doctor Hook's Wheel Barometer*, fig. 6. This was invented about 1668, and is meant to render the alterations in the air more sensible. Here the barometer tube has a large ball AB at top, and is bent up at the lower or open end, where an iron ball G floats on the top of the mercury in the tube, to which is connected another ball H by a cord, hanging freely over a pulley, turning an index KL about its centre. When the mercury rises in the part FG, it raises the ball, and the other ball descends and turns the pulley with the index round a graduated circle from N towards M and P; and the contrary way when the quicksilver and the ball sink in the bent part of the tube. Hence the scale is easily enlarged 10 or 12 fold, being increased in proportion of the axis of the pulley to the length of the index KL. But then the friction of the pulley and axis is some obstruction to the free motion of the quicksilver. Contrivances to lessen the friction &c, may also be seen in the *Philos. Transf.* vol. 52, art. 29, and vol. 60, art. 10.

*The Steelyard Barometer*, for so that may be called which is represented by fig. 7, which enlarges the scale in the proportion of the shorter to the longer arm of a steelyard. AB is the barometer tube, close at A and open at B, immersed in a cylindrical glass cistern CD, which is but very little wider than the tube AB is. The barometer tube is suspended to the shorter arm of an index like a steelyard, moving on the fulcrum E, and the extremity of its longer arm pointing to the divisions of a graduated arch, with which index the tube is nearly in equilibrio. When the pressure of the atmosphere is lessened, the mercury descends out of the tube into the cistern which raises the tube and the shorter arm of the index, and consequently the extremity of the longer moves downwards, and passes over a part of the graduated arch. And on the contrary this moves upwards when the pressure of the atmosphere increases.

*The Pendant Barometer*, fig. 8, was invented by M. Amontons, in 1695. It consists of a single conical tube AB, hung up by a thread, the larger or open end downwards, and having no vessel or cistern, because the conical figure supplies that, and the column of mercury sustained is always equal to that in the common barometer tube; which is effected thus; when the pressure of the air is less, the mercury sinks down to a lower and wider part of the tube, and consequently the altitude of its column will be less; and on the contrary, by a greater pressure of the atmosphere the mercury is forced up to a higher and narrower part, till the length of the column CD be equal to that in the tube of the common barometer.—The inconvenience of this barometer is, that as the bore must be made very small, to prevent the mercury from falling out by an accidental shake, the friction



friction and adhesion to the sides of the tube prevent the free motion of the mercury.

*Mr. Rowning's Compound Barometers.* This gentleman has several contrivances for enlarging the scale, and that in any proportion whatever. One of these is described in the *Philos. Trans.* N° 427, and also in his *Nat. Philos.* part 2; and another in the same part, which is here represented at fig. 9. ABC is a compound tube, hermetically sealed at A, and open at C; empty from A to D, filled with mercury from thence to B, and from hence to E with water. Hence by varying the proportions of the two tubes AF and FC, the scale of variation may be changed in any degree.

*The Marine Barometer.* This was first invented by Dr. Hook, to be used on board of ship, being contrived so as not to be affected or injured by the motion of the ship. His contrivance consisted of a double thermometer, or a couple of tubes half filled with spirit of wine; the one sealed at both ends, with a quantity of air included; the other sealed at one end only. The former of these is affected only by the warmth of the air; but the other is affected both by the external warmth and by the variable pressure of the atmosphere. Hence, considering the spirit thermometer as a standard, the excess of the rise or fall of the other above it will shew the increase or decrease of the pressure of the atmosphere. This instrument is described by Dr. Halley, in the *Philos. Trans.* N° 269, where he says of it, "I had one of these barometers with me in my late southern voyage, and it never failed to prognosticate and give early notice of all the bad weather we had, so that I depended thereon, and made provision accordingly; and from my own experience I conclude, that a more useful contrivance hath not for this long time been offered for the benefit of navigation."

*Mr. Nairne*, an ingenious artist in London, has lately invented a new kind of *Marine Barometer*; which differs from the common barometer by having the lower part of the tube, for about 2 feet long, made very small, to check the vibrations of the mercury, which would otherwise arise from the motions of the ship. This is also assisted by being hung in gimbals, by a part which subjects it to be the least affected by such motions.

Another sort of *Marine Barometer* has also been invented by M. Passemente, an ingenious artist at Paris. This contrivance consists only in twisting the middle of the tube into a spiral of two revolutions; by which contrivance the impulses which the mercury receives from the motions of the ship are destroyed, by being transmitted in contrary directions.

*The Statical Baroscope, or Barometer*, of Mr. Boyle, &c. This consists of a large glass bubble, blown very thin, and then balanced by a small brass weight. Hence these two bodies being of unequal bulk, the larger will be very much affected by a change of the density of the medium, but the less not at all as to sense: So that, when the atmosphere becomes denser, the ball loses more of its weight, and the brass weight preponderates; and contrariwise when the air grows lighter.

*Mr. Caswell's Baroscope, or Barometer.* This is described in the *Philos. Trans.* vol. 24, and seems to be the most sensible and exact of any. It is thus described: Suppose ABCD, (fig. 10) is a bucket of water, in

which is the baroscope *xrezybism*, which consists of a body *xrism*, and a tube *exyo*, which are both concave cylinders, made of tin, or rather glass, and communicating with each other. The bottom of the tube *zy* has a leaden weight to sink it, so that the top of the body may just swim even with the surface of the water by the addition of some grain weights on the top. When the instrument is forced with its mouth downwards, the water ascends into the tube to the height *yu*. To the top is added a small concave cylinder, or pipe, to keep the instrument from sinking down to the bottom: *md* is a wire: and *ms*, *dc* are two threads oblique to the surface of the water, which perform the office of diagonals: for while the instrument sinks more or less by an alteration in the gravity of the air, where the surface of the water cuts the thread is formed a small bubble, which ascends up the thread while the mercury of the common baroscope ascends, and vice versa.

It appears from a calculation which the author makes, that this instrument shews the alterations in the air 1200 times more accurately than the common barometer. He observes, that the bubble is seldom known to stand still even for a minute; that a small blast of wind, which cannot be heard in a chamber, will sensibly make it sink; and that a cloud passing over it always makes it descend, &c.

While some have been increasing the sensibility of the barometer by enlarging the variations, others have endeavoured to make it more convenient by reducing the length of the tube. *M. Amontons*, in 1688, first proposed this alteration in the structure of barometers, by joining several tubes to one another, alternately filled with mercury and with air, or some other fluid; and the number of these tubes may be increased at pleasure: but the contrivance is perhaps more ingenious than useful.

*M. Mairan's reduced Barometer*, which is only 3 inches long, serves the purpose of a manometer, in shewing the dilatations of the air in the receiver of an air-pump; and instruments of this kind are now commonly applied to this use.

*The Portable Barometer*, is so contrived that it may be carried from one place to another without being disordered. The end of the tube is tied up in a leather bag not quite full of mercury; which being pressed by the air, forces the mercury into the tube, and keeps it suspended at its proper height: This bag is usually inclosed in a box, through the bottom of which passes a screw, by means of which the mercury may be forced up to the top of the tube, and prevented from breaking it by dashing against the top when the instrument is removed from one station to another. It seems Mr. Patrick first made a contrivance of this kind: but the portable barometer has received various improvements since; and the most complete of this kind has been described by M. De Luc, in his *Recherches*, vol. 2, pa. 5 &c, together with the apparatus belonging to it, the method of construction and use, and the advantages attending it. Improvements have also been suggested by Sir George Shuckburgh, and Col. Roy, which have been carried into execution, with farther improvements also, by Mr. Ramsden, and other ingenious artists in London.



Fig. 11 represents this instrument, as inclosed in its mahogany case by means of three metallic rings *a a a*. This case is a hollow cone, so shaped within as to contain steadily the body of the barometer, and is divided into three branches from *b* to *c*, forming three legs or supports for the instrument when observations are making, and sustaining it at the part *d* of the case, as it appears in Fig. 12, by an improved kind of gimbals, in which its own weight renders it sufficiently steady at any time. In the part of the frame *f g* where the barometer tube appears, is made a long slit or opening, that the column of mercury may be seen against the light, and the vernier piece *f* brought down to coincide very nicely with the edge of the mercury. When the instrument is fixed in its stand, the screw *l* is to be turned to let the mercury down to its proper station, and a peg at *i* must be loosened, to admit the external air to act upon the mercury contained in the box *k*. The proper adjustment, or mode of observing what is called the zero or 0 division of the column of mercury, is by observing it in the transparent part of the box *k*, which has a glass tube or reservoir for the quicksilver, and an edged piece of metal attached to the external part of it; with the edge of which the mercury is to be brought into contact by turning the screw *l* to the right or left as occasion requires. The vernier piece at *f*, which determines the altitude of the mercurial column, is first brought down by the hand to a near contact, and then accurately adjusted by turning the screw *e* at the top. The divisions annexed to the tube of this instrument may be of any sort, or of any degree of smallness, according to the purposes it is intended to serve. To accommodate it to the use of foreigners as well as natives, there are commonly added scales of both French and English inches, with their subdivisions to any extent required. It is usual to place the French scale of inches on the right side at *f g*, from 19 to 31 inches, measured from the zero or surface of the mercury in the box *k* below; each inch being divided into lines or 12th parts, and each line subdivided by the vernier into 10th parts, or 120th parts of inches; by means of which therefore the length of the mercurial column may be determined to the 120th part of a French inch. The other scale, which is placed on the left side of the instrument, is divided into English inches, and each inch into 20th parts, which by a vernier are subdivided into 25th parts, or 500th parts of inches; by this means shewing the height of the mercury to the 500th part of an English inch. But this vernier is figured double or each division is accounted 2, which reduces the measures to 1000ths of an inch for the conveniency of calculation, in measuring altitudes of hills &c.

A thermometer is always attached to the instrument, as a necessary appendage to it, being fastened to the body at *h*, and sunk into the surface of the frame, to preserve it from injury: the degrees of this thermometer are marked on two scales, one on each side of it, viz, the scale of Fahrenheit, and that of Reaumur; the freezing point of the former being at 32, and of the latter at 0. Also on the right hand side of these two scales there is a third, called a scale of *correction*,

placed oppositely to that of Fahrenheit, with the word *add* and *subtract* marked; which shews the necessary correction of the observed altitude of the mercury at any given temperature of the air, indicated by the thermometer.

There are several other pieces of mechanism about the instrument, which will be evident by inspection; and the manner of making the observations, with the necessary calculations, are fully explained in M. de Luc's *Recherches sur les Modifications de l'Atmosphere*, and the *Philos. Trans.* vol. 67 and 68, before cited.

The *Common Chamber Weatherglass*, is also usually fitted up in a neat mahogany frame, and other embellishments, to make it an ornamental piece of furniture. It consists of the common tube barometer, with a thermometer by the side of it, and an hygrometer at the top, as exhibited in fig. 13.

To the foregoing may be added a new sort of *Barometer*, or *Weather Instrument by the Sound of a Wire*. This is mentioned by M. Lazowski in his Tour through Switzerland: it is as yet but in an imperfect state, and was lately discovered there by accident. It seems that a clergyman, though near-sighted, often amused himself with firing at a mark, and contrived to stretch a wire so as to draw the mark to him to see how he had aimed. He observed that the wire sometimes sounded as if it vibrated like a musical cord; and that after such soundings, a change always ensued in the state of the atmosphere; from whence he came to predict rain or fine weather. On making farther experiments, it was found that the sounds were most distinct when extended in the plane of the meridian. And according to the weather which was to follow, it was found that the sounds were more or less soft, or more or less continued; also fine weather, it is said, was announced by the tones of counter-tenor, and rain by those of bass. It has been said that M. Volta mounted 15 chords in this way at Pavia, to bring this method to some precision, but no accounts have yet appeared of the success of his observations.

*The Phenomena and Observations of the Barometer* \*. The phenomena of the barometer are various; but authors are not yet agreed upon the causes of them; nor is the use of it, as a weather-glass, yet perfectly ascertained, though daily observations and experience lead us still nearer to precision. Mr. Boyle observes that the phenomena of the barometer are so precarious, that it is exceedingly difficult to form any certain general rules concerning the rise and fall of the mercury. Even that rule fails which seems to hold the most generally, viz, that the mercury is low in high winds. The best rules however that have been deduced by several authors are as follow.

*Dr Halley's Rules for judging of the Weather.*

1. In calm weather, when the air is inclined to rain, the mercury is commonly low.
2. In serene, good, and settled weather, the mercury is generally high.
3. Upon very great winds, though they be not accompanied with rain, the mercury sinks lowest of all, according to the point of the compass the wind blows from.

\* An ingenious author observes that, by means of barometers we may regain the knowledge that still resides, in brutes, and which we forfeited by not continuing in the open air, as they mostly do; and, by our intemperance, corrupting the *crasis* of our organs of sense.



4. The greatest heights of the mercury are found upon easterly or north-easterly winds, other circumstances alike.

5. In calm frosty weather, the mercury commonly stands high.

6. After very great storms of wind, when the mercury has been very low, it generally rises again very fast.

7. The more northerly places have greater alterations of the barometer than the more southerly, near the equator.

8. Within the tropics, and near them, there is little or no variation of the barometer, in all weathers. For instance, at St. Helena it is little or nothing, at Jamaica  $\frac{3}{10}$ ths of an inch, and at Naples the variation hardly ever exceeds an inch; whereas in England it amounts to 2 inches and a half, and at Peterburgh to  $3\frac{1}{2}$  nearly.

*Dr Beal*, who followed the opinion of M. Pascal, observes that, *ceteris paribus*, the mercury is higher in cold weather than in warm: and in the morning and evening usually higher than at mid-day.—That in settled and fair weather, the mercury is higher than either a little before or after, or in the rain; and that it commonly descends lower after rain than it was before it. And he ascribes these effects to the vapours with which the air is charged in the former case, and which are dispersed by the falling rain in the latter. If it chance to rise higher after rain, it is usually followed by a settled serenity. And that there are often great changes in the air, without any perceptible alteration in the barometer.

*Mr Patrick's Rules for judging of the Weather.* These are esteemed the best of any general rules hitherto made:

1. The rising of the mercury presages, in general, fair weather; and its falling, foul weather, as rain, snow, high winds, and storms.

2. In very hot weather, the falling of the mercury indicates thunder.

3. In winter, the rising presages frost: and in frosty weather, if the mercury falls 3 or 4 divisions, there will certainly follow a thaw. But in a continued frost, if the mercury rises, it will certainly snow.

4. When foul weather happens soon after the falling of the mercury, expect but little of it; and on the contrary, expect but little fair weather when it proves fair shortly after the mercury has risen.

5. In foul weather, when the mercury rises much and high, and so continues for 2 or 3 days before the foul weather is quite over, then expect a continuance of fair weather to follow.

6. In fair weather, when the mercury falls much and low, and thus continues for 2 or 3 days before the rain comes; then expect a great deal of wet, and probably high winds.

7. The unsettled motion of the mercury, denotes uncertain and changeable weather.

8. You are not so strictly to observe the words engraved on the plates, as the mercury's rising and falling; though in general it will agree with them. For if it stands at *much rain*, and then rises up to *changeable*, it presages fair weather; though not to continue so long as if the mercury had risen higher. And so, on

the contrary, if the mercury stood at *fair*, and falls to *changeable*, it presages foul weather; though not so much of it as if it had sunk lower.

Upon these rules of Mr Patrick, the following *Remarks* are made by *Mr Rowning*. That it is not so much the absolute height of the mercury in the tube that indicates the weather, as its motion up and down: wherefore, to pass a right judgment of what weather is to be expected, we ought to know whether the mercury is actually rising or falling; to which end the following rules are of use.

1. If the surface of the mercury is convex, standing higher in the middle of the tube than at the sides, it is a sign that the mercury is then rising.

2. But if the surface be concave, or hollow in the middle, it is then sinking. And,

3. If it be plain, or rather a very little convex, the mercury is stationary: for mercury being put into a glass tube, especially a small one, naturally has its surface a little convex, because the particles of mercury attract one another more forcibly than they are attracted by glass. Farther,

4. If the glass be small, shake the tube; then if the air be grown heavier, the mercury will rise about half a 10th of an inch higher than it stood before; but if it be grown lighter, it will sink as much. And, it may added, in the wheel or circular barometer, tap the instrument gently with the finger, and the index will visibly start forwards or backwards according to the tendency to rise or fall at that time. This proceeds from the mercury's sticking to the sides of the tube, which prevents the free motion of it till it be disengaged by the shock: and therefore when an observation is to be made with such a tube, it ought to be first shaken; for sometimes the mercury will not vary of its own accord, till the weather is present which it ought to have indicated.

And to the foregoing may be added the following additional rules, more accurately drawn from later and more close observation of the motions of the barometer, and the consequent changes in the air in this country.

1. In winter, spring, and autumn, the sudden falling of the mercury, and that for a large space, denotes high winds and storms; but in summer it denotes heavy showers, and often thunder: and it always sinks lowest of all for great winds, though not accompanied with rain; though it falls more for wind and rain together than for either of them alone. Also, if, after rain, the wind change into any part of the north, with a clear and dry sky, and the mercury rise, it is a certain sign of fair weather.

2. After very great storms of wind, when the mercury has been low, it commonly rises again very fast. In settled fair and dry weather, except the barometer sink much, expect but little rain; for its small sinking then, is only for a little wind, or a few drops of rain; and the mercury soon rises again to its former station. In a wet season, suppose in hay-time and harvest, the smallest sinking of the mercury must be minded; for when the constitution of the air is much inclined to showers, a little sinking in the barometer then denotes more rain, as it never then stands very high. And, if



in such a season, it rise suddenly, very fast, and high, expect not fair weather more than a day or two, but rather that the mercury will fall again very soon, and rain immediately to follow: the slow gradual rising, and keeping on for 2 or 3 days, being most to be depended on for a week's fair weather. And the unsettled state of the quicksilver always denoting uncertain and changeable weather, especially when the mercury stands any where about the word *changeable* on the scale.

3. The greatest heights of the mercury, in this country, are found upon easterly and north-easterly winds; and it may often rain or snow, the wind being in these points, and the barometer sink little or none, or it may even be in a rising state, the effect of those winds counteracting. But the mercury sinks for wind, as well as rain, in all the other points of the compass; but rises as the wind shifts about to the north or east, or between those points: but if the barometer should sink with the wind in that quarter, expect it soon to change from thence; or else, should the fall of the mercury be much, a heavy rain is then likely to ensue, as it sometimes happens.

*Cause of the Phenomena of the Barometer.*

To account for the foregoing phenomena of the barometer, many hypotheses have been framed, which may be reduced to two general heads, viz, *mechanical* and *chemical*. The chief writers upon these causes, are Pascal, Beal, Wallis, Garcin, Garden, Lister, Halley, Garsten, De la Hire, Mariotte, Le Cat, Woodward, Leibnitz, De Mairan, Hamberger, D. Bernoulli, Muschenbroek, Chambers, De Luc, Black, &c; and an account of most of their hypotheses may be seen at large in M. De Luc's *Recherches sur les Modifications de l'Atmosphere*, vol. 1. chap. 3; see also the *Philos. Transf.* and various other works on this subject. It may suffice to notice here slightly a few of the principal of them.

Dr. Lister accounts for the changes of the barometer from the alterations by heat and cold in the mercury itself; contracting by cold, and expanding by heat. But this, it is now well known, is quite insufficient to account for the whole of the effect.

The changes in the weight or pressure of the atmosphere must therefore be regarded as the principal cause of those in the barometer. But then, the difficulty will be to assign the cause of that cause, or whence arise those alterations that take place in the atmosphere, which are sometimes so great as to alter its pressure by the 10th part of the whole quantity. It is probable that the winds, as driven about in different directions, have a great share in them; vapours and exhalations, rising from the earth, may also have some share; and some perhaps the flux and reflux occasioned in the air by the moon; as well as some chemical causes operating between the different particles of matter.

Dr. Halley thinks the winds and exhalations sufficient; and on this principle gives a theory, the substance of which may be comprised in what follows:

1st, That the winds must alter the weight of the air in any particular country; and this, either by bringing together a greater quantity of air, and so load-

ing the atmosphere of any place; which will be the case as often as two winds blow from opposite parts, at the same time, towards the same point: or by sweeping away some part of the air, and giving room for the atmosphere to expand itself; which will happen when two winds blow opposite ways from the same point at the same time: or lastly by cutting off the perpendicular pressure of the air; which is the case when a single wind blows briskly any way; it being found by experience, that a strong blast of wind, even made by art, will render the atmosphere lighter; and hence the mercury in a tube below it, as well as in others more distant, will considerably subside. See *Philos. Transf.* N<sup>o</sup> 292.

2dly, That the cold nitrous particles, and even the air itself condensed in the northern regions, and driven elsewhere, must load the atmosphere, and increase its pressure.

3dly, That heavy dry exhalations from the earth must increase the weight of the atmosphere, as well as its elastic force; as we find the specific gravity of menstrooms increased by dissolved salts and metals.

4thly, That the air being rendered heavier by these and the like causes, is thence better able to support the vapours; which being likewise intimately mixed with it, make the weather serene and fair. Again, the air being made lighter from the contrary causes, it becomes unable to support the vapours with which it is replete; these therefore precipitating, are collected into clouds, the particles of which in their progress unite into drops of rain.

Hence he infers, it is evident that the same causes which increase the weight of the air, and render it more able to support the mercury in the barometer, do likewise produce a serene sky, and a dry season; and that the same causes which render the air lighter, and less able to support the mercury, do likewise generate clouds and rain.

But these principles, though well adapted to many of the particular cases of the barometer, seem however to fall short of some of the principal and most obvious ones, besides being liable to several objections.

Leibnitz accounted for the fall of the mercury before rain by another principle, viz, That as a body specifically lighter than a fluid, while it is sustained by it, adds more weight to that fluid than when, by being reduced in bulk, it becomes specifically heavier, and descends; so the vapour, after it is reduced into the form of clouds, and descends, adds less weight to the air than it did before; and hence the mercury sinks in the tube.—But here, granting that the drops of rain formed from the vapours always increasing in size as they fall lower, were continually accelerated also in their motion, and so the air suffer a continued loss of their weight as they descend; it may however be objected, that by the descent of the mercury the rain is foretold a much longer time before it comes, than the vapour can be supposed to take up in falling: that many times, and in different places, there falls a great deal of rain, without any sinking of the mercury at all; as also that there often happens a fall of the mercury without any rain ensuing: and that sometimes the mercury will suddenly sink, in a short space of



of time, half an inch or more, which answers to 7 inches of rain, or about one third of the whole quantity falling in the whole year.

Mr. *De Luc* supposes that the changes observed in the pressure of the atmosphere, are chiefly produced by the greater or less quantity of vapours floating in it: as others have attributed them to the same cause, but have given a different explanation of it. His opinion is, that vapours diminish the specific gravity, and consequently the absolute weight, of those columns of the atmosphere into which they are received, and which, notwithstanding this admixture, still remain of the same height with adjoining columns that consist of pure or dry air. He afterwards vindicates and more fully explains this theory, and applies it to the solution of the principal phenomena of the barometer, as depending on the varying density and weight of the atmosphere.

Dr. *James Hutton*, in his *Theory of Rain*, printed in the *Transactions of the Royal Society of Edinburgh*, vol. 1, gives ingenious and plausible reasons for thinking that the lessening the weight of the atmosphere, by the fall of rain, is not the cause of the fall of the barometer; but that the principal, if not the only cause, arises from the commotions in the atmosphere, which are chiefly produced by sudden changes of heat and cold in the air. "The barometer, says he, is an instrument necessarily connected with motions in the atmosphere; but it is not equally affected with every motion in that fluid body. The barometer is chiefly affected by those motions by which there are produced accumulations and abstractions of this fluid, in places or regions of sufficient extent to affect the pressure of the atmosphere upon the surface of the earth. But as every commotion in the atmosphere may, under proper conditions, be a cause for rain, and as the want of commotion in the atmosphere is naturally a cause of fair weather, this instrument may be made of great importance for the purpose of meteorological observations, although not in the certain and more simple manner in which it has been, with the increase of science, so successfully applied to the measuring of heights." See RAIN.

In the *Encyclopædia Britannica* there is another theory of the changes in the barometer, as depending on the heat in the atmosphere, not as producing commotions there, but as altering the specific gravity of the air by the changes of heat and cold. The principles of this theory are, 1st, That vapour is formed by an intimate union between the elements of fire and water, by which the fire or heat is so totally enveloped, and its action so perfectly suspended by the aqueous particles, that it not only loses its properties of burning and of giving light, but becomes incapable of affecting the most sensible thermometer, in which case it is said to be in a latent state: and 2d, That if the atmosphere be affected by any unusual degree of heat, it thence becomes incapable of supporting so long a column of mercury as before; for which reason it is that the barometer sinks.

From these axioms it would follow, that as vapour is formed by an union of fire with water, whether by attraction or a solution of the water in the fire, the vapour cannot be condensed till this union, attraction,

or solution, is at an end. Hence the beginning of the condensation of the vapour, or the first signs of approaching rain, must be the separation of the fire which is latent in the vapour. In the beginning, this may be either slow and partial, or it may be sudden and violent: in the first case, the rain will come on slowly, and after a considerable time; but in the other, it will come very quickly, and in a great quantity. But Dr. Black has proved, that when fire quits its latent state, however long it may have lain dormant and insensible, it always re-assumes its proper qualities, and affects the thermometer just the same as if it had never been absorbed. The consequence of this is, that in proportion as the latent heat is discharged from the vapour, those parts of the atmosphere into which it is discharged must be sensibly affected by it; and in proportion to the heat communicated to those parts, they will become specifically lighter, and the mercury will sink of course.

In the *Memoirs of the Literary Society of Manchester*, vol. 4, is also a curious paper on this subject, viz, *Meteorological Observations made on different Parts of the Western Coast of Great Britain: arranged by T. Garnett, M. D.* This paper is composed of materials furnished by several observers; those of Mr. Copland, surgeon at Dumfries, are of special importance. This gentleman is of opinion that the changes of the barometer indicate approaching hot and cold weather, with much more certainty than dry and wet. "Every remarkable elevation of the barometer, says he, where it is of any duration, is followed by very warm or dry weather, and moderate as to wind, or by all of them; but heat seems to have most influence and connexion; and when it is deficient, the continuance of the other two will be longer and more remarkable; therefore the calculation must be in a compound ratio of the excess and deficiency of the heat, and of the dryness of the weather in comparison of the medium of the season; and with regard to the want of strong wind, it appears to be intimately connected with the last, as they shew that no precipitation is going on in any of the neighbouring regions."

In his 14th and 15th remarks, he had said,

'14th, That the barometer being lower, and continuing so longer than what can be accounted for by immediate falls, or stormy weather, indicates the approach of very cold weather for the season; and also, cold weather, though dry, is always accompanied by a low barometer, till near its termination.'

'15th, That warm weather is always preceded and mostly accompanied by a high barometer; and the rising of the barometer in the time of broken or cold weather, is a sign of the approach of warmer weather: and also if the wind is in any of the cold points, a sudden rise of the barometer indicates the approach of a southerly wind, which in winter generally brings rain with it.'

In the two following remarks, Mr. Copland had explained certain phenomena from a principle similar to that on which Dr. Darwin has so much insisted: (Botanic Garden, I. notes p. 79, &c.)

'That the falling of the barometer may proceed from a decomposition of the atmosphere occurring a-



round or near that part of the globe where we are placed, which will occasion the electricity of the atmosphere to be repelled upwards in fine lambent portions; or driven downwards or upwards in more compacted balls of fire; or lastly, to be carried along with the rain, &c, in an imperceptible manner to the surface of the earth: the precipitation of the watery parts generally very soon takes place, which diminishes the real gravity of the atmosphere, and also by the decomposition of some of the more active parts, the air loses part of that elastic and repulsive power which it so eminently possessed, and will therefore press with less force on the mercury of the barometer than before, by which means a fall ensues.

'That the cause of the currents of air, or winds, may also be this way accounted for: and in very severe storms, where great decompositions of the atmosphere take place, this is particularly evident, such as generally occur in one or more of the West India islands at one time, a great loss of real gravity, together with a considerable diminution of the spring of the air immediately ensues; hence a current commences, first in that direction whence the air has most gravity, or is most disposed to undergo such a change; but it being soon relieved of its superior weight or spring on that side, by the decomposition going on as fast as the wind arrives on the island, it immediately veers to another point, which then rushes in mostly with an increase of force; thus it goes on till it has blown more than half way round the points of the compass during the continuation of the hurricane. For in this manner the West India phenomena, as well as the alteration of the wind during heavy rains in this country, can only be properly accounted for.' See remark No. 4.

Mr. C.'s 4th aphorism is, 'That the heaviest rains, when of long continuance, generally beign with the wind blowing easterly, when it gradually veers round to the south; and that the rain does not then begin to cease till the wind has got to the west, or rather a little to the northward of it, when, it may be added, it commonly blows with some violence.'

Many other observations on the barometer, the weather, &c. may be seen in various parts of the *Philos. Trans.* And for other curious papers on the same, and other subjects connected with the barometer, see the *Gentleman's Magazine* for 1789, p. 317; also *Greus Journal of Nat. Philos.* printed at Leipzig 1792, for the influence of the sun and moon upon the barometer.

*The Barometer applied to the measuring of Altitudes.*

The secondary character of the barometer, namely as an instrument for measuring accessible heights or depths, was first proposed by Pascal, and Descartes, as has been before observed; and succeeding philosophers have been at great pains to ascertain the proportion between the fall of the barometer and the height to which it is carried; as Halley, Mariotte, Maraldi, Schenchzer, J. Cassini, D. Bernoulli, Horrebow, Bouguer, Shuckburgh, Roy, and more especially by De Luc, who has given a critical and historical detail of most of the attempts that have at different times been made for applying the motion of the mercury in the barometer to the measurement of accessible heights.

And for this purpose serves the portable barometer, before described, (fig. 11 and 12, plate 4,) which should be made with all the accuracy possible. Various rules have been given by the writers on this subject, for computing the height ascended from the given fall of the mercury in the tube of the barometer, the most accurate of which was that of Dr. Halley, till it was rendered much more accurate by the indefatigable researches of De Luc, by introducing into it the corrections of the columns of mercury and air, on account of heat. And other corrections and modifications of the same may be seen inserted under the article *ATMOSPHERE*, where the most correct rule is deduced from one single experiment only. This rule is as follows:

*The Rule for Computing Altitudes, is this,*

Viz,  $10000 \times \log. \text{ of } \frac{M}{m}$  is the altitude in fathoms,

in the mean temperature of  $31^{\circ}$ ; and for every degree of the thermometer above that, the result must be increased by so many times its 435th part, and diminished when below it: in which theorem *M* denotes the length of the column of mercury in the barometer tube at the bottom, and *m* that at the top of the hill, or other eminence; which lengths may be expressed in any one and the same sort of measures, whether feet, or inches, or tenths, &c, and either English, or French, or of any other nation; but the result is always in fathoms, of 6 English feet each.

And the *Precepts*, in words, for the practice of measurements by the barometer, are these following:

1st, Observe the height of the barometer at the bottom of any height or depth, proposed to be measured; together with the temperature of the mercury by means of the thermometer attached to the barometer, and also the temperature of the air in the shade by another thermometer which is detached from the barometer.

2dly, Let the same thing be done also at the top of the said height or depth, and as near to the same time with the former as may be. And let those altitudes of mercury be reduced to the same temperature, if it be thought necessary, by correcting either the one or the other, viz, augmenting the height of the mercury in the colder temperature, or diminishing that in the warmer, by its 9600th part for every degree of difference between the two; and the altitudes of mercury so corrected, are what are denoted by *M* and *m*, in the algebraic formula above.

3dly, Take out the common logarithms of the two heights of mercury, so corrected, and subtract the less from the greater, cutting off from the right hand side of the remainder three places for decimals; so shall those on the left be fathoms in whole numbers, the tables of logarithms being understood to be such as have 7 places of decimals.

4thly, Correct the number last found, for the difference of the temperature of the air, as follows: viz, Take half the sum of the two temperatures of the air, shewn by the detached thermometers, for the mean one; and for every degree which this differs from the standard temperature of  $31^{\circ}$ , take so many times the 435th part of the fathoms above found, and add them if the mean temperature



temperature be more than  $31^{\circ}$ , but subtract them if it be below  $31^{\circ}$ ; so shall the sum or difference be the true altitude in fathoms, or being multiplied by 6, it will give the true altitude in English feet.

*Example 1.* Let the state of the barometers and thermometers be as follows, to find the altitude: viz,

Thermometers.		Barometers.
detached.	attached.	
57	57	29.68 lower
42	43	25.28 upper
mean $49\frac{1}{2}$	diff. 14	
As 9600	: 14	: : 29.68 : .04
		cor. .04 logs.
mean $49\frac{1}{2}$		M = 29.64 - 4718782
stand. 31		m = 25.28 - 4027771
diff. $18\frac{1}{2}$		As 435 : $18\frac{1}{2}$ :: 691.011 : 29.388
		29.388
	the altitude {	720.399 fath.
	sought is {	or 4322.394 feet.

*Example 2.* To find the altitude of a hill, when the state of the barometer and thermometer, as observed at the bottom and top of it, is as follows; viz,

Thermometers.		Barometers.
detached.	attached.	
35	41	29.45
31	38	26.82
mean 33	diff. 3	
As 9600	: 3	: : 29.45 : .01
		.01 logs.
mean 33		M = 29.44 - 4689378
stand. 31		m = 26.82 - 4284588
diff. 2		As 435 : 2 :: 404.790 : 1.86
		1.86
	the altitude {	406.65 fathoms.
	sought is {	or 2439.90 feet.

See this rule investigated under the article PNEUMATICS, at the end.

N. B. The mean height of the barometer in London, upon an average of two observations in every day of the year, kept at the house of the Royal Society, for many years past, is 29.88; the medium temperature, or height of the thermometer, according to the same, being  $58^{\circ}$ . But the medium height at the surface of the sea, according to Sir Geo. Shuckburgh (Philos. Transf. 1777, p. 586) is 30.04 inches, the heat of the barometer being  $55^{\circ}$ , and of the air  $62^{\circ}$ .

BAROSCOPE, a machine for shewing the alterations in the weight or pressure of the atmosphere. See BAROMETER.

BARREL, an English vessel or cask, containing 36 gallons of beer measure, or 32 gallons of ale measure. The barrel of beer, vinegar, or of liquor preparing for vinegar, ought to contain 34 gallons, according to the standard of the ale quart.

BARREL, in Clock-work, is the cylinder about which the spring is wrapped.

BARRICADE, or BARRICADO, a military term for a fence, or retrenchment, hastily made with vessels, or baskets of earth, carts, trees, stakes, or the like, to preserve an army from the shot or assault of an enemy.

BARRIER, a kind of fence made at a passage, retrenchment, gate, or such like, to stop it up against an enemy.

BARROW (ISAAC), a very eminent mathematician and divine of the 17th century, was born at London in October, 1630, being the son of Thomas Barrow, then a linen-draper of that city, but descended from an ancient family in Suffolk. He was at first placed at the Charter-house school for two or three years; where his behaviour afforded but little hopes of success in the profession of a scholar, being fond of fighting, and promoting it among his school-fellows: but being removed to Felsted in Essex, his disposition took a different turn; and having soon made a great progress in learning, he was first admitted a pensioner of Peter House in Cambridge; but when he came to join the university, in Feb. 1645, he was entered at Trinity college. He now applied himself with great diligence to the study of all parts of literature, especially natural philosophy. He afterward turned his attention to the profession of physic, and made a considerable progress in anatomy, botany, and chemistry: he next studied divinity; then chronology, astronomy, geometry, and the other branches of the mathematics; with what success, his writings afterwards most eminently shewed.

When Dr. Dupont resigned the chair of Greek professor, he recommended his pupil Mr. Barrow for his successor, who, in his probation exercise, shewed himself equal to the character that had been given him by this gentleman; but being suspected of favouring Arminianism, he was not preferred. This disappointment it seems determined him to quit the college, and visit foreign countries; but his finances were so low, that he was obliged to dispose of his books, to enable him to execute that design.

He left England in June 1655, and visited France, Italy, Turkey, &c. At several places, in the course of this tour, he met with kindness and liberal assistance from the English ambassadors, &c, which enabled him to benefit the more from it, by protracting his stay, and prolonging his journey. He spent more than a year in Turkey, and returned to England by way of Venice, Germany, and Holland, in 1659. At Constantinople he read over the works of St. Chrysostom, once bishop of that see, whom he preferred to all the other fathers.

On his return home Barrow was episcopally ordained by bishop Brownrig; and in 1660, he was chosen to the Greek professorship at Cambridge. In July 1662, he was elected professor of geometry in Gresham college: in which station, he not only discharged his own duty, but supplied likewise the absence of Dr. Pope the astronomy professor. Among his lectures, some were upon the projection of the sphere and perspective, which are lost; but his Latin oration, previous to his lectures, is still extant. About this time Mr. Barrow was offered a good living; but the condition annexed, of teaching the patron's son, made him refuse it, as thinking it too



like a simonial contract. Upon the 20th of May 1663 he was elected a fellow of the Royal Society, in the first choice made by the council after their charter. The same year the executors of Mr. Lucas having, according to his appointment, founded a mathematical lecture at Cambridge, they selected Mr. Barrow for the first professor; and though his two professorships were not incompatible with each other, he chose to resign that of Gresham-college, which he did May the 20th, 1664. In 1669 he resigned the mathematical chair to his learned friend Mr. Isaac Newton, being now determined to quit the study of mathematics for that of divinity. On quitting his professorship, he had only his fellowship of Trinity-college, till his uncle gave him a small sinecure in Wales, and Dr. Seth Ward bishop of Salisbury conferred upon him a prebend in his church. In the year 1670 he was created doctor in divinity by mandate; and, upon the promotion of Dr. Pearson master of Trinity college to the see of Chester, he was appointed to succeed him by the king's patent bearing date the 13th of February 1672: upon which occasion the king was pleased to say, "he had given it to the best scholar in England." In this, his majesty did not speak from report, but from his own knowledge; the doctor being then his chaplain, he used often to converse with him, and, in his humorous way, to call him an "unfair preacher," because he exhausted every subject, and left no room for others to come after him. In 1675 he was chosen vice-chancellor of the university; and he omitted no endeavours for the good of that society, nor in the line of his profession as a divine, for the promotion of piety and virtue; but his useful labours were abruptly terminated by a fever on the 4th of May 1677, in the 47th year of his age. He was interred in Westminster abbey, where a monument, adorned with his bust, was soon after erected, by the contribution of his friends.

Dr. Barrow's works are very numerous, and indeed various, mathematical, theological, poetical, &c, and such as do honour to the English nation. They are principally as follow:

1. *Euclidis Elementa*. Cantab. 1655, in 8vo.
2. *Euclidis Data*. Cantab. 1657, in 8vo.
3. *Lectiones Opticæ* xviii, Lond. 1669, 4to.
4. *Lectiones Geometricæ* xiii, Lond. 1670, 4to.
5. *Archimedis Opera*, Apollonii Conicorum libri iv, Theodosii Sphericorum lib. iii; nova methodo illustrata, et succincte demonstrata. Lond. 1675, in 4to.

The following were published after his decease, viz:

6. *Lectio*, in qua theoremata Archimedis de sphaera et cylindro per methodum indivisibilium investigata, ac breviter investigata, exhibentur. Lond. 1678, 12mo.
7. *Mathematicæ Lectiones habitæ in scholis publicis academici Cantabrigienfis*, an. 1664, 5, 6, &c. Lond. 1683.
8. All his English works in 3 volumes, Lond. 1683, folio.—These are all theological, and were published by Dr. John Tillotson.
9. *Isaaci Barrow Opuscula*, viz, *Determinationes*, *Conciones ad Clerum*, *Orationes*, *Poemata*, &c. volumen quartum. Lond. 1687, folio.

Dr. Barrow left also several curious papers on mathematical subjects, written in his own hand, which were

communicated by Mr. Jones to the author of "The Lives of the Gresham Professors," a particular account of which may be seen in that book, in the Life of Barrow.

Several of his works have been translated into English, and published; as the *Elements* and *Data* of Euclid; the *Geometrical Lectures*, the *Mathematical Lectures*. And accounts of some of them were also given in several volumes of the *Philos. Transf.*

Dr. Barrow must ever be esteemed, in all the subjects which exercised his pen, a person of the clearest perception, the finest fancy, the soundest judgment, the profoundest thought, and the closest and most nervous reasoning. "The name of Dr. Barrow (says the learned Mr. Granger) will ever be illustrious for a strength of mind and a compass of knowledge that did honour to his country. He was unrivalled in mathematical learning, and especially in the sublime geometry; in which he has been excelled only by his successor Newton. The same genius that seemed to be born only to bring hidden truths to light, and to rise to the heights or descend to the depths of science, would sometimes amuse itself in the flowery paths of poetry, and he composed verses both in Greek and Latin. He at length gave himself up entirely to divinity; and particularly to the most useful part of it, that which has a tendency to make men wiser and better."

Several good anecdotes are told of Barrow, as well of his great integrity, as of his wit, and bold intrepid spirit and strength of body. His early attachment to fighting when a boy is some indication of the latter; to which may be added the two following anecdotes: In his voyage between Leghorn and Smyrna the ship was attacked by an Algerine pirate, which after a stout resistance they compelled to sheer off, Barrow keeping his post at the gun assigned him to the last. And when Dr. Pope in their conversation asked him, "Why he did not go down into the hold, and leave the defence of the ship to those, to whom it did belong?" He replied, "It concerned no man more than myself: I would rather have lost my life, than to have fallen into the hands of those merciless infidels."

There is another anecdote told of him, which shewed not only his intrepidity, but an uncommon goodness of disposition, in circumstances where an ordinary share of it would have been probably extinguished. Being once on a visit at a gentleman's house in the country, where the necessary was at the end of a long garden, and consequently at a great distance from the room where he lodged; as he was going to it before day, for he was a very early riser, a fierce mastiff, that used to be chained up all day, and let loose at night for the security of the house, perceiving a strange person in the garden at that unusual time, set upon him with great fury. The doctor caught him by the throat, grappled with him, and, throwing him down, lay upon him: once he had a mind to kill him; but he altered his resolution, on recollecting that this would be unjust, since the dog did only his duty, and he himself was in fault for rambling out of his room before it was light. At length he called out so loud, that he was heard by some of the family, who came presently out, and freed the doctor and the dog from the danger they both had been in.

Among



Among other instances of his wit and vivacity, they relate the following rencontre between him and that wicked wit lord Rochester. These two meeting one day at the court, while the doctor was king's chaplain in ordinary, Rochester, thinking to banter him, with a flip-pant air, and a low formal bow, accosted him with, "Doctor, I am yours to my shoe-tie:" Barrow perceiving his drift, and determined upon defending himself, returned the salute, with, "My lord, I am yours to the ground." Rochester, on this, improving his blow, quickly returned it, with, "Doctor, I am yours to the centre;" which was as smartly followed up by Barrow, with, "My lord, I am yours to the antipodes." Upon which, Rochester, disdaining to be foiled by a musty old piece of divinity, as he used to call him, exclaimed, "Doctor, I am yours to the lowest pit of hell;" upon which Barrow, turning upon his heel, with a sarcastic smile, archly replied, "There, my lord, I leave you."

**BARS**, in Music, are the spaces quite through any composition, separated by upright lines drawn across the five horizontal lines, each of which either contains the same number of notes of the same kind, or so many other notes as will make up a like interval of time; for all the bars, in any piece, must be of the same length, and played in the same time.

**BARTER**, or **TRUCK**, is the exchanging of one commodity for another; and forms a rule in the commercial part of arithmetic, by which the commodities are properly calculated and equalled, by computing first the value of the commodity which is given, and then the quantity of the other which will amount to the same sum.

**BASE**, **BasIS**, in Architecture, denotes the lower part of a column or pedestal.

**BASE**, in Geometry, the lowest side of any figure. Any side of a figure may be considered as its base, according to the position in which it may be conceived as standing; but commonly it is understood of the lowest side: as the base of a triangle, of a cone, cylinder, &c.

**BASE LINE**, in Perspective, denotes the common section of the picture and the geometrical plane.

**BASE RING**, of a Cannon, is the great ring next behind the vent or touch-hole.

**BASE**, *alternate*. See **ALTERNATE**.

**BASEMENT**, in Architecture, a continued base, extended a considerable length, as about a house, a room, or other piece of building.

**BASILIC**, in the ancient Architecture, was a large hall, or court of judicature, where the magistrates sat to administer justice.

**BASILICA**, or **BASILICUS**, the same as *Regulus*, or *Cor Leonis*, being a fixed star of the first magnitude in the constellation *Leo*.

**BASILISK**, in the older Artillery, was a large piece of ordnance so called from its resemblance to the supposed serpent of that name. It threw an iron ball of 200 pounds weight; and was in great repute in the time of Solyman emperor of the Turks, in the wars of Hungary; but it is now grown out of use in most parts of Europe. Paulus Jovius relates the terrible slaughter made in a Spanish ship by a single ball from one of these basilisks; after passing through the beams and planks in the ship's head, it killed upwards of 30 men. And Maffeus speaks of basilisks made of brass, each of which

required 100 yoke of oxen to draw them.—More modern writers also give the name basilisk to a much smaller and sizeable piece of ordnance, made of 15 feet long by the Dutch, but of only 10 by the French, and carrying a ball of 48 pounds. The largest size of cannon now used by the English, are the 32 pounders.

**BASIS**, in Geometry, the same as **BASE**.

**BASS**, the lowest in the four parts of music; by some esteemed the basis and principal part of all, and by others as scarcely necessary in some tunes.

**BASSANTIN** (**JAMES**), a Scotch astronomer of the 16th century, born in the reign of James the 4th of Scotland. He was a son of the Laird of Bassantin in the Merse. After finishing his education at the university of Glasgow, he travelled through Germany and Italy, and then settled in the university of Paris, where he taught mathematics with great applause. Having acquired some property in this employment, he returned to Scotland in 1562, where he died 6 years after.

From his writings it appears he was no inconsiderable astronomer, for the age he lived in; but, according to the fashion of the times, he was not a little addicted to judicial astrology. It was doubtless to our author that Sir James Melvil alludes in his Memoirs, when he says that his brother Sir Robert, when he was using his endeavours to reconcile the two queens Elizabeth and Mary, met with one Bassantin a man learned in the high sciences, who told him, "that all his travel would be in vain; for, said he, they will never meet together; and next, there will never be any thing but dissembling and secret hatred for a while, and at length captivity and utter wreck to our queen from England." He added, "that the kingdom of England at length shall fall, or right, to the crown of Scotland: but it shall cost many bloody battles; and the Spaniards shall be helpers, and take a part to themselves for their labour." A prediction in which Bassantin partly guessed right, which it is likely he was enabled to do from a judicious consideration of probable circumstances and appearances.

Bassantin's works are,

1. *Astronomia Jacobi Bassantini Scoti, opus absolutissimum, &c; ter. edit. Latine et Gallice.* Genev. 1599, fol. This is the title given it by Tornæsius, who translated it into Latin from the French, in which language it was first published.

2. *Paraphrase de l'Astrolabe, avec une amplification de l'usage de l'Astrolabe.* Lyons 1555. Paris 1617, 8vo.

3. *Mathematica Genethliaca.*

4. *Arithmetica.*

5. *Musica secundum Platonem.*

6. *De Mathefi in Genere.*


**BASSOON**, a musical instrument of the wind kind, serving for a bass to the haut-boy. It is blown with a reed, and furnished with eleven holes.

**BASS-VIOLE**, a bass to the viol.

**BASTION**, in the modern fortification, a large mass of earth at the angles of a work, connecting the curtains to each other; and answers to the bulwark of the ancients. It is formed by two faces, two flanks, and two demigorges. The two faces form the salient angle, or angle of the bastion; the two flanks form with the faces, the *epaules* or shoulders; and the union of the



the other two ends of the flanks with the curtains forms the two angles of the flanks.



**Solid BASTION**, are those that are entirely filled up with earth to the height of the rampart, without any void space towards the centre.

**Void or Hollow BASTION**, has the rampart and parapet ranging only round the flanks and spaces, so that a void space is left within towards the centre, where the ground is so low that if the rampart be taken, no retrenchment can be made in the centre, but what will lie under the fire of the besieged.

**Regular BASTION**, is that which has its due proportion of faces, flanks, and gorges.

**Deformed or Irregular BASTION**, is when the irregularity of the lines and angles throws the bastion out of shape: as when it wants one of the demigorges, one side of the interior polygon being too short, &c.

**Demi BASTION**, or *Half bastion*, also otherwise called an *Epaulment*, has but one face and flank.

**Double BASTION**, is when one bastion is raised within, and upon the plane of another bastion.

**Flat BASTION**, is one built in the middle of the curtain, when it is too long to be defended by the usual bastions at the extremities.

**Composed BASTION**, is when the two sides of the interior polygon are very unequal, which makes the gorges also unequal.

**Cut BASTION**, is that which has a re-entering angle at the point, and is sometimes called a *BASTION with a Tenaille*, whose point is cut off, making an angle inwards, and two points outwards. This is used when the salient angle would be too sharp, or when water or some other impediment prevents it from being carried out to its full extent.

**BASTON**, or **BATOON**, in Architecture, a moulding in the base of a column, called also a *Tore* or *Torus*.

**BATTEN**, a name given by workmen to a scantling or piece of wooden stuff, about an inch thick, and from 2 to 4 inches broad; of a considerable but indeterminate length.

**BATTERING**, the attacking a place, work, or the like, with heavy artillery.

**BATTERING-RAM**, a military engine used for beating down walls, before the invention of gunpowder and the modern artillery. It was no other than a long heavy beam of timber, armed with an iron head, something like the head of a ram. This being pushed violently with constant successive blows against a wall, gradually shakes it with a vibratory motion, till the stones are

disjointed and the wall falls down. There were several kinds of the battering-ram, the first rude and plain, which the soldiers carried in their arms by main force, and so struck the head of it against the wall. The second was slung by a rope about the middle to another beam lying across upon a couple of posts; which was the kind described by Josephus as used at the siege of Jerusalem. A third fort was covered over with a shell or screen of boards, to defend the men from the stones and darts of the besieged upon the walls, and thence called *testudo arietaria*. And Felibien describes a fourth sort of battering-ram, which ran upon wheels; and was the most perfect and effectual of any.

Vitruvius affirms, that the battering-ram was first invented by the Carthaginians, while they laid siege to Cadiz: yet Pliny assures us, that the ram was invented or used at the siege of Troy; and that it was this that gave occasion to the fable of the wooden horse. In fact there can be no doubt but that the use of some sort of a battering-ram is as old as the art of war itself. And it has even been suspected that the walls of Jericho, mentioned in the book of Joshua, were beaten down by this instrument, the rams horns there mentioned, by means of which they were overthrown, being no other than the horns of the battering-rams. Pephasthenos, a Tyrian, afterwards contrived to suspend it with ropes; and lastly, Polydus, the Theffalian, mounted it on wheels, at the siege of Byzantium, under Philip of Macedon.

Plutarch relates, that Marc Anthony, in the Parthian war, made use of a ram 80 feet long: and Vitruvius affirms that they were sometimes 106, and even 120 feet long; which must have given an amazing force to this engine. The ram required 100 soldiers to work and manage it at one time; who being exhausted, another century relieved them; by which means it was kept playing continually without intermission. See fig. 2, plate V, which represents the battering-ram suspended in its open frame; in which 3 denotes the form of the head, fastened to the enormous beam 2, by three or four bands (4) of iron, of about four feet in breadth. At the extremity of each of these bands was an iron chain (5), one end of which was fastened to a hook (6), and to the last link at the other extremity was firmly bound a cable, which ran the whole length of the beam to the end of the ram 7, where these cables were bound all together as fast as possible with small ropes. To the end of these cables was fastened another, that consisted of several strong cords platted together to a certain length, and then running single (8), at each of which were placed several men, to balance and work the machine. 10 Is the chain or cable by which the ram was hung to the cross beam (11), fixed on the top of the frame; and 12 is the base of the machine.

The unsuspended ram differed from this only in the manner of working it; as it moved on small wheels upon another large beam, instead of being slung by a chain or cable.

**BATTERY**, in the Military Art, a place raised to plant cannon upon, to play with more advantage upon the enemy. It consists of an epaulment or a breastwork, of about 8 feet high, and 18 or 20 feet thick.

In all batteries, the open spaces through which the muzzles



muzzles of the cannon are pointed, are called *Embrasures*, and the distances between the embrasures, *merlons*. The guns are placed upon a platform of planks &c, ascending a little from the parapet, to check the recoil, and that the gun may be the easier brought back again to the parapet: they are placed from 12 to 16 feet distant from one another, that the parapet may be strong, and the gunners have room to work.

*Mortar BATTERIES* differ from the others, in that the slope of the parapet is inwards, and it is without embrasures, the shells being fired quite over the parapet, commonly at an angle of 45 degrees elevation.

*Open BATTERY*, is nothing more than a number of cannon, commonly field-pieces, ranged in a row abreast of one another, perhaps on some small natural elevation of the ground, or an artificial bank a little raised for the purpose.

*Covered or Masked BATTERY*, is when the cannon and gunners are covered by a bank or breast-work, commonly made of brush-wood, faggots, and earth, called a fascine battery.

*Sunk or Buried BATTERY*, is when its platform is sunk, or let down into the ground, so that trenches must be cut in the earth opposite the muzzles of the guns, to serve as embrasures to fire through. This is mostly used on the first making of approaches in besieging and battering a place.

*Cross BATTERIES*, are two batteries playing athwart each other upon the same object, forming an angle there, and battering to more effect, because what one battery shakes, the other beats down.

*BATTERY d'Enfilade*, is one that scours or sweeps the whole length of a straight line.

*BATTERY en Echarpe*, is one that plays obliquely.

*BATTERY de Reverse, or Murdering Battery*, is one that plays upon the enemy's back.

*Camerade or Joint BATTERY*, is when several guns play upon one place at the same time.

*BATTERY, in Electricity*, is a combination of coated surfaces of glass, commonly jars, so connected together that they may be charged at once, and discharged by a common conductor. Mr. Galvani, a German electrician, first contrived to increase the shock by charging several phials at the same time.—Dr Franklin, having analysed the Leyden phial, and found that it lost at one surface the electric fire received at the other, constructed a battery of eleven large panes of sash window glass, coated on both sides, and so connected that the whole might be charged together, and with the same labour as one single pane; then by bringing all the *giving* sides into contact with one wire, and all the *receiving* sides with another, he contrived to unite the force of all the plates, and to discharge them at once.—Dr. Priestley describes a still more complete battery. This consists of 64 jars, each 10 inches long, and 2½ inches in diameter, all coated within an inch and a half of the top, forming in the whole about 32 square feet of coated surface. A piece of very fine wire is twisted about the lower end of the wire of each jar, to touch the inside coating in several places; and it is put through a pretty large piece of cork, within the jar, to prevent any part of it from touching the side, by which a spontaneous discharge might be made. Each wire is turned round so as to make a loop at the

upper end; and through these loops passes a pretty thick brass rod with knobs, each rod serving for one row of the jars; and these rods are made to communicate together by a thick chain laid over them, or as many of them as may be wanted. The jars stand in a box, the bottom of which is covered with a tin plate; and a bent wire touching the plate passes through the box, and appears on the outside. To this wire is fastened any conductor designed to communicate with the outside of the battery; and the discharge is made by bringing the brass knob to any of the knobs of the battery. When a very great force is required, the size or number of the jars may be increased, or two or more batteries may be used.—But the largest and most powerful battery of all, is that employed by Dr. Van Marum, to the amazing large electrical machine, lately constructed for Teyler's museum at Haarlem. This grand battery consists of a great number of jars coated as above, to the amount of about 130 square feet; and the effects of it, which are truly astonishing, are related by Dr. Van Marum in his description of this machine, and of the experiments made with it, at Haarlem 1785, &c. See also Franklin's *Exper. and Observ.* and Priestley's *History of Electricity*.

*BATTLEMENTS*, in Architecture, are notches or indentures in the top of a wall or other building, like embrasures, to look through.

*BAY*, in Geography, denotes a small gulph, or an arm of the sea stretching up into the land; being larger in the middle within, than at its entrance, which is called the mouth of the bay.

*BAYER (JOHN)*, a German lawyer and astronomer of the latter part of the 16th and beginning of the 17th century, but in what particular year or place he was born, is not certainly known: however, his name will be ever memorable in the annals of astronomy, on account of that great and excellent work which he first published in the year 1603, under the title of *Uranometria*, being a complete celestial atlas, or large folio charts of all the constellations, with a nomenclature collected from all the tables of astronomy, ancient and modern, with the useful invention of denoting the stars in every constellation by the letters of the Greek alphabet, in their order, and according to the order of magnitude of the stars in each constellation. By means of these marks, the stars of the heavens may, with as great facility, be distinguished and referred to, as the several places of the earth are by means of geographical tables; and as a proof of the usefulness of this method, our celestial globes and atlases have ever since retained it; and hence it is become of general use through all the literary world; astronomers, in speaking of any star in the constellation, denoting it by saying it is marked by Bayer,  $\alpha$ , or  $\beta$ , or  $\gamma$ , &c.

Bayer lived many years after the first publication of this work, which he greatly improved and augmented by his constant attention to the study of the stars. At length, in the year 1627, it was republished under a new title, viz, *Caelum Stellatum Christianum*, that is, the *Christian Stelled Heaven*, or the *Starry Heavens Christianized*: for in this work, the heathen names and characters, or figures of the constellations, were rejected, and others, taken from the scriptures, were inserted



serted in their stead, to circumscribe the respective constellations. This was the project of one Julius Schiller, a civilian of the same place. But this attempt was too great an innovation, to find success, or a general reception, which might occasion great confusion. And, we even find in the later editions of this work, that the ancient figures and names were restored again; at least so I find them in two editions, of the years 1654, and 1661, which are now before me.

**BEAD**, in Architecture, is a round moulding, carved in short embossments, like beads in necklaces: and sometimes an astragal is thus carved. There is also a sort of plain bead often set on the edge of each fascia of an architrave; as also sometimes on the lining board of a door case, the upper edge of skirting boards, &c.

**BEAD**, in assaying, the small ball or mass of pure metal separated from the *scoria*, and seen distinct and pure in the middle of the coppel while in the fire.

**BEAM**, in Architecture, a large timber laid across a building, into which the principal rafters are framed. Several ingenious authors have considered the force or strength of beams, as supporting their own weight and any other additional weight; particularly Varignon, and Parent in the *Memoir. Acad. R. Scien. an. 1708*, and Mr. Emerson, on the Strength and Strefs of Timber, in his *Mechanics*. Mr. Parent makes the proportion of the depth to the breadth of a beam to be as 7 to 5 when it is strongest.

**BEAMS of a ship**, are the large, main, cross timbers, stretched from side to side, to support the decks, and keep the sides of the ship from falling together.

**BEAM of a balance**, is the horizontal piece of wood or iron supported on a pivot in the middle, and at the extremities of which the two scales are suspended, for weighing any thing.

**BEAM-Compass**, an instrument consisting of a wooden or brass square beam, having sliding sockets carrying steel or pencil points; and are used for describing large circles, the radii of which are beyond the extent of the common compasses.

**BEAR**, in Astronomy, a name given to two constellations, called the *greater* and the *lesser bear*, or *URSA major* and *minor*. The pole star is in the tail of the little bear, and is within less than 2 degrees of the north pole. See *URSA, major* and *minor*.

**BEARD**, of a Comet, the rays which it emits in the direction in which it moves; as distinguished from the tail, or the rays emitted or left behind it as it moves along, being always in a direction from the sun.

**BEARER**, in Architecture, a post or brick wall, trimmed up between the two ends of a piece of timber, to shorten its bearing, or to prevent its bearing with the whole weight at the ends only.

**BEARING**, in Geography and Navigation, the situation of one place from another, with regard to the points of the compass; or an arch of the horizon between the meridian of a place and a line drawn through this and another place, or the angle formed by a line drawn through the two places and their meridians.—The bearings of places on the ground are usually determined by the magnetic needle.

**BEATS**, in a Clock or Watch, are the strokes made by the fangs or pallets of the spindle of the balance; or of the pads in a royal pendulum. For the number and use of the beats, see *Derham's Artificial Clock Maker*, pa. 14 and seq.

**BED, of a Great Gun**, a plank laid between the cheeks of the carriage, on the middle transum, for the breech of the gun to rest upon.

**BED, or Stool**, of a mortar, a thick and strong planking on which a mortar is placed, hollowed a little to receive the breech and trunions.

**BED-MOULDING**, in Architecture, a term used by workmen for those members in a cornice which are placed below the coronet, or crown. It usually consists of these four members, an ogee, a list, a large boutine, and another list under the coronet.

**BELIDOR (BERNARD FOREST DE)**, an engineer in the service of France, but born in Catalonia in 1698. He was professor in the new school of artillery at la Fere, where he published his course of mathematics for the use of the artillery and engineers. He was the first who seriously considered the quantity of gunpowder proper for charges, and reduced it to 2-3ds the quantity. He was named Associate in the Academy of Sciences in 1751; and died Sept. 8, 1761, at 63 years of age.

His works that have been published, are:

1. *Sommaire d'un Cours d'Architecture militaire, civile & hydraulique*, in 12mo, 1720.
2. *Nouveau Cours de Mathematiques, &c.* in 4to, 1725.
3. *La Science des Ingenieurs*, in 4to, 1729.
4. *Le Bombardier Francois*, in 4to, 1734.
5. *Architecture Hydraulique*, 4 vols. in 4to, 1737.
6. *Dictionnaire portatif de l'Ingenieur*, in 8vo.
7. *Traité des Fortifications*, 4 vols. in 4to.

Besides several pieces inserted in the volumes of the *Memoirs of the Academy of Sciences*, for the years 1737, 1750, 1753, and 1756.

**BELLA'TRIX**, in Astronomy, a ruddy, glittering star of the 2d magnitude, in the left shoulder of Orion. Its name is from the Latin *bellum*, as being anciently supposed to have great influence in kindling wars, and forming warriors.

**BELTS, Fasciæ**, in Astronomy, two zones or girdles surrounding the planet Jupiter, brighter than the rest of his body, and terminated by parallel lines. They are observed however to be sometimes broader and sometimes narrower, and not always occupying exactly the same part of the disc. Jupiter's belts were first observed and described by Huygens, in his *Syst. Saturn*. Dark spots have often been observed on the belts of Jupiter; and M. Cassini observed a permanent one on the northern side of the most southern belt, by which he determined the length of Jupiter's days, or the time in which the planet revolves upon its axis, which is 9h. 56m. Some astronomers suppose that these belts are seas, which alternately cover and leave bare large tracts of the planet's surface: and that the spots are gulphs in those seas, which are sometimes dry, and sometimes full. But Azout conceived that the spots are protuberances of the belts; and others again



again are of opinion that the transparent and moveable spots are the shadows of Jupiter's satellites.

Cassini also speaks of the belts of Saturn; being three dark, straight, parallel bands, or *fasciæ*, on the disc of that planet. But it does not appear that Saturn's belts adhere to his body, as those of Jupiter do; but rather that they are large dark rings surrounding the planet at a distance. Some imagine that they are clouds in the atmosphere of Saturn; though it would seem that the middlemost is the shadow of his ring.

**BENDING**, the reducing a body to a curved or crooked form. The bending of boards, planks, &c, is effected by means of heat, whether by boiling or otherwise, by which their fibres are so relaxed that they may be bent into any figure. Bernoulli has a discourse on the bending of springs, or elastic bodies. And Amontons gives several experiments concerning the bending of ropes. The friction of a rope bent or wound about an immovable cylinder, is sufficient, with a very small power, to sustain very great weights.

**BERENICE's Hair**; see *COMA Berenices*.

**BERKELEY** (GEORGE), the virtuous and learned bishop of Cloyne in Ireland, was born in that kingdom, at Kilerin, the 12th of March 1684. After receiving the first part of his education at Kilkenny school, he was admitted a pensioner of Trinity College, Dublin, at 15 years old; and chosen fellow of that college in 1707.

The first public proof he gave of his literary abilities was, *Arithmetica absque Algebra aut Euclide demonstrata*; which from the preface it appears he wrote before he was 20 years old, though he did not publish it till 1707. It is followed by a Mathematical Miscellany, containing observations and theorems inscribed to his pupil Samuel Molineux.

In 1709 came out the *Theory of Vision*; which of all his works it seems does the greatest honour to his sagacity; being, it has been observed, the first attempt that ever was made to distinguish the immediate and natural objects of sight, from the conclusions we have been accustomed from infancy to draw from them. The boundary is here traced out between the ideas of sight and touch; and it is shewn, that though habit hath so connected these two classes of ideas in the mind, that they are not without a strong effort to be separated from each other, yet originally they have no such connection; inasmuch, that a person born blind, and suddenly made to see, would at first be utterly unable to tell how any object that affected his sight would affect his touch; and particularly would not from sight receive any idea of distance, or external space, but would imagine all objects to be in his eye, or rather in his mind.

In 1710 appeared *The Principles of Human Knowledge*; and in 1713 *Dialogues between Hylas and Philonous*: the object of both which pieces is, to prove that the commonly received notion of the existence of matter, is false; that sensible material objects, as they are called, are not external to the mind, but exist in it, and are nothing more than impressions made upon it by the immediate act of God, according to certain rules termed laws of nature.

VOL. I.

Acuteness of parts and beauty of imagination were so conspicuous in Berkeley's writings, that his reputation was now established, and his company courted; men of opposite parties concurred in recommending him. For Steele he wrote several papers in the *Guardian*, and at his house became acquainted with Pope, with whom he always lived in friendship. Swift recommended him to the celebrated earl of Peterborough, who being appointed ambassador to the king of Sicily and the Italian States, took Berkeley with him as chaplain and secretary in 1713, with whom he returned to England the year following.

His hopes of preferment expiring with the fall of queen Anne's ministry, he some time after embraced an offer made him by Ashe, bishop of Clogher, of accompanying his son in a tour through Europe. In this he employed four years; and besides those places which fall within the grand tour, he visited some that are less frequented, and with great industry collected materials for a natural history of those parts, but which were unfortunately lost in the passage to Naples. He arrived at London in 1721; and being much affected with the miseries of the nation, occasioned by the South-sea scheme in 1720, he published the same year *An Essay towards preventing the ruin of Great Britain*: reprinted in his *Miscellaneous Tracts*.

His way was now open into the very first company. Pope introduced him to lord Burlington, by whom he was recommended to the duke of Grafton, then appointed lord-lieutenant of Ireland, who took Berkeley over as one of his chaplains in 1721. The latter part of this year he accumulated the degrees of bachelor and doctor in divinity: and the year following he had a very unexpected increase of fortune from the death of Mrs. Vanhomrigh, the celebrated Vanessa, to whom he had been introduced by Swift: this lady had intended Swift for her heir; but perceiving herself to be slighted by him, she left her fortune, of 8000l. between her two executors, of whom Berkeley was one. In 1724 he was promoted to the deanery of Derry, worth 1100l. a year.

In 1725 he published, and it has since been reprinted in his *Miscellaneous Tracts*, *A Proposal for converting the savage Americans to Christianity, by a college to be erected in the Summer Isles, otherwise called the Isles of Bermuda*. The proposal was well received, at least by the king; and he obtained a charter for founding the college, with a parliamentary grant of 20,000l. toward carrying it into execution: but he could never get the money, it being otherwise employed by the minister; so that after two years stay in America on this business, he was obliged to return, and the scheme dropped.

In 1732 he published *The Minute Philosopher*, in 2 volumes 8vo, against Freethinkers. In 1733 he was made bishop of Cloyne; and might have been removed in 1745, by lord Chesterfield, to Clogher; but declined it. He resided constantly at Cloyne, where he faithfully discharged all the offices of a good bishop; yet continued his studies with unabated attention.

About this time he engaged in a controversy with the mathematicians, which made a good deal of noise in the literary world; and the occasion of it was this:

D d

Addison



Addison had given the bishop an account of the behaviour of their common friend Dr. Garth in his last illness, which was equally unpleasing to both these advocates of revealed religion. For when Addison went to see the doctor, and began to discourse with him seriously about another world, "Surely, Addison, replied he, I have good reason not to believe those trifles, since my friend Dr. Halley, who has dealt so much in demonstration, has assured me, that the doctrines of christianity are incomprehensible, and the religion itself an imposture." The bishop therefore took up arms against Halley, and addressed to him, as to an Infidel Mathematician, a discourse called *The Analyst*; with a view of shewing that mysteries in faith were unjustly objected to by mathematicians, who he thought admitted much greater mysteries, and even falsehoods in science, of which he endeavoured to prove that the doctrine of Fluxions furnished a clear example. This attack gave occasion to *Robins's Discourse concerning the Method of Fluxions*, to *Maclaurin's Fluxions*, and to other smaller works upon the same subject; but the direct answers to *The Analyst* were made by a person under the name of Philalethes Cantabrigiensis, but commonly supposed to be Dr. Jurin, whose first piece was, *Geometry no Friend to Infidelity*, 1734. To this the bishop replied in *A Defence of Freethinking in Mathematics; with an Appendix concerning Mr. Walton's Vindication*, 1735; which drew a second answer the same year from Philalethes, under the title of *The Minute Mathematician, or the Freethinker no just Thinker*, 1735. Other writings in this controversy, beside those before mentioned, were

1. A Vindication of Newton's Principles of Fluxions against the objections contained in the *Analyst*, by J. Walton, Dublin, 1735.

2. The Catechism of the Author of the Minute Philosopher fully answered, by J. Walton, Dublin, 1735.

3. Reasons for not replying to Mr. Walton's Full Answer, in a letter to P. T. P. by the author of the Minute Philosopher, Dublin, 1735.

4. An Introduction to the Doctrine of Fluxions, and Defence of the Mathematicians against the objections of the author of the *Analyst*, &c. Lond. 1736.

5. A new Treatise of Fluxions; with answers to the principal objections in the *Analyst*, by James Smith, A. M. Lond. 1737.

6. Mr. Robins's Discourse of Newton's Methods of Fluxions, and of Prime and Ultimate Ratios, 1735.

7. Mr. Robins's Account of the preceding Discourse, in the *Repub. of Letters*, for October 1735.

8. Philalethes's Considerations upon some passages contained in two letters to the author of the *Analyst* &c, in *Repub. of Letters*, Novemb. 1735.

9. Mr. Robins's Review of some of the principal objections that have been made to the doctrine of Fluxions &c. *Repub. of Letters* for Decem. 1735.

10. Philalethes's Reply to ditto, in the *Repub. of Letters*, Jan. 1736.

11. Mr. Robins's Dissertation, shewing that the account of the doctrines of Fluxions &c, is agreeable to the real sense and meaning of their great Inventor, &c, *Repub. of Letters*, April 1736.

12. Philalethes's Considerations upon ditto, in *Repub. of Letters*, July 1736.

13. Mr. Robins's Remarks on ditto, in *Repub. of Letters*, Aug. 1736.

14. Mr. Robins's Remainder of ditto, in an Appendix to the *Repub. of Letters*, Sept. 1736.

15. Philalethes's Observations upon ditto, in an Appendix to the *Repub. of Letters*, Nov. 1736.

16. Mr. Robins's Advertisement in *Repub. of Letters*, Decemb. 1736.

17. Philalethes's Reply to ditto, in an Appendix to the *Repub. of Letters* for Decem. 1736.

18. Some Observations on the Appendix to the *Repub. of Letters* for Decem. 1736, by Dr. Pemberton, in the *Works of the Learned* for Feb. 1737. With some smaller pieces in the same.

In 1736 bishop Berkeley published *The Querist*, "a discourse addressed to magistrates, occasioned by the enormous licence and irreligion of the times;" and many other things afterward of a smaller kind. In 1744 came out his celebrated and curious book, "*Siris*; a Chain of Philosophical Reflections and Inquiries concerning the virtues of Tar-water." It had a second impression, with additions and emendations, in 1747; and was followed by *Farther Thoughts on Tar-water*, in 1752. In July the same year he removed, with his lady and family, to Oxford, partly to superintend the education of a son, but chiefly to indulge the passion for learned retirement, which had always strongly possessed him. He would have resigned his bishoprick for a canonry or headship at Oxford; but it was not permitted him. Here he lived highly respected, and collected and printed the same year all his smaller pieces in 8vo. But this happiness did not long continue, being suddenly cut off by a palsy of the heart Jan. 14, 1753, in the 69th year of his age, while listening to a sermon that his lady was reading to him.

The excellence of Berkeley's moral character is conspicuous in his writings: he was certainly a very amiable as well as a very great man; and it is thought that Pope scarcely said too much, when he ascribed

"To Berkeley every virtue under heaven."

BERME, in Fortification, a small space of ground, 4 or 5 feet wide, left without the rampart, between it and the side of the moat, to receive the earth that rolls down from the rampart, and prevent its falling into the ditch and filling it up.—Sometimes, for greater security, the berme is palisadoed.

BERNARD (Dr. EDWARD), a learned astronomer, critic and linguist, was born at Perry St. Paul, near Towcester, the 2d of May 1638, and educated at Merchant-Taylor's school, and at St. John's college Oxford. Having laid in a good fund of classical learning at school, in the Greek and Latin languages, he applied himself very diligently at the university to the study of history, the eastern languages, and mathematics under the celebrated Dr. Wallis. In 1668 he went to Leyden to consult some Oriental manuscripts left to that university by Joseph Scaliger and Levin Warner, and especially the 5th, 6th, and 7th books of Apollonius's Conics, the Greek text of which is lost, and this Arabic version having been brought from the east by the celebrated Golius,



Golius, a transcript of which was thence taken by Bernard, and brought with him to Oxford, with intent to publish it there with a Latin translation; but he was obliged to drop that design for want of encouragement. This however was afterwards carried into effect by Dr. Halley in 1710, with the addition of the 8th book, which he supplied by his own ingenuity and industry.

At his return to Oxford, Bernard examined and collated the most valuable manuscripts in the Bodleian library. In 1669, the celebrated Christopher Wren, Savilian professor of astronomy at Oxford, having been appointed surveyor-general of his majesty's works, and being much detained at London by this employment, obtained leave to name a deputy at Oxford, and pitched upon Mr. Bernard, which engaged the latter in a more particular application to the study of astronomy. But in 1673 he was appointed to the professorship himself, which Wren was obliged to resign, as, by the statutes of the founder, Sir Henry Saville, the professors are not allowed to hold any other office either ecclesiastical or civil.

About this time a scheme was set on foot at Oxford, of collecting and publishing the ancient mathematicians. Mr. Bernard, who had first formed the project, collected all the old books published on that subject since the invention of printing, and all the manuscripts he could discover in the Bodleian and Savilian libraries, which he arranged in order of time, and according to the matter they contained; of this he drew up a synopsis or view; and as a specimen he published a few sheets of Euclid, containing the Greek text, and a Latin version, with Proclus's commentary in Greek and Latin, and learned scholia and corollaries. The synopsis itself was published by Dr. Smith, at the end of his life of our author, under the title of *Veterum Mathematicorum Græcorum, Latino-rum, et Arabum, Synopsis*. And at the end of it there is a catalogue of some Greek writers, whose works are supposed to be lost in their own language, but are preserved in the Syriac or Arabic translations of them.

Mr. Bernard undertook also an edition of the *Parva Syntaxis Alexandrina*; in which, besides Euclid, are contained the small treatises of Theodosius, Menelaus, Aristarchus, and Hipficles; but it never was published.

In 1676 he was sent to France, as tutor to the dukes of Grafton and Northumberland, sons to king Charles the 2d by the dutchess of Cleveland, who then lived with their mother at Paris: but the simplicity of his manners not suiting the gaiety of the dutchess's family, he returned about a year after to Oxford, and pursued his studies; in which he made great proficiency, as appears by his many learned and critical works. In 1691, being presented to the rectory of Brightwell in Berkshire, he quitted his professorship at Oxford, in which he was succeeded by David Gregory, professor of mathematics at Edinburgh.

Toward the latter end of his life he was much afflicted with the stone; yet notwithstanding this, and other infirmities, he undertook a voyage to Holland, to attend the sale of Golius's manuscripts, as he had once before done at the sale of Heinsius's library. On his return to England, he fell into a languishing consumption, which put an end to his life the 12th of January 1696, in the 58th year of his age.

Beside the works of his before mentioned, he was

author of many other compositions. He composed tables of the longitudes, latitudes, right-ascensions, &c. of the fixed stars: he wrote Observations on the Obliquity of the Ecliptic; and other pieces inserted in the Philosophical Transactions. He wrote also,

1. *A Treatise of the Ancient Weights and Measures.*
2. *Chronologiæ Samaritanæ Synopsis*, in two tables.
3. *Testimonies of the Ancients concerning the Greek Version of the Old Testament by the Seventy.*

And several other learned works. Besides a great number of valuable manuscripts left at his death.

BERNARD (Dr. JAMES), professor of philosophy and mathematics, and minister of the Walloon church at Leyden, was born September the 1st 1658, at Nions in Dauphine. Having studied at Geneva, he returned to France in 1679, and was chosen minister of Venterol, a village in Dauphine; but some time after he was removed to the church of Vinsobres in the same province. To avoid the persecutions against the protestants in France, he went into Holland, where he was appointed one of the pensionary ministers of Ganda. He here published several political and historical works. And in 1699 he began the *Nouvelles de la Republique des Lettres*, which continued till December 1710. In 1705 he was chosen minister of the Walloon church at Leyden; and about the same time, Mr. de Volder, professor of philosophy and mathematics at Leyden, having resigned, Mr. Bernard was appointed his successor; upon which occasion the university also presented him with the degrees of doctor of philosophy and master of arts. In 1716 he published a supplement to Moreri's dictionary in 2 volumes folio. The same year he resumed his *Nouvelles de la Republique des Lettres*; which he continued till his death, which happened the 27th of April 1718, in the 60th year of his age.

BERNOULLI (JAMES), a celebrated mathematician, born at Basil the 27th of December 1654. Having taken his degrees in that university, he applied himself to divinity at the entreaties of his father, but against his own inclination, which led him to astronomy and mathematics. He gave very early proofs of his genius for these sciences, and soon became a geometri-cian, without a preceptor, and almost without books; for if one by chance fell into his hands, he was obliged to conceal it, to avoid the displeasure of his father, who designed him for other studies. This situation induced him to choose for his device, Phaeton driving the chariot of the sun, with these words, *Invito patre sidera verso*, "I traverse the stars against my father's will:" alluding particularly to astronomy, to which he then chiefly applied himself.

In 1676 he began his travels. When he was at Geneva, he fell upon a method to teach a young girl to write who had been blind from two months old. At Bourdeaux he composed universal gnomonic tables; but they were never published. He returned from France to his own country in 1680. About this time there appeared a comet, the return of which he foretold; and wrote a small treatise upon it. Soon after this he went into Holland, where he applied himself to the study of the new philosophy. Having visited Flanders and Brabant, he passed over to England; where he formed an acquaintance with the most eminent men in the sciences, and was frequent at their philosophical meetings. He



returned to his native country in 1682; and exhibited at Basil a course of experiments in natural philosophy and mechanics, which consisted of a variety of new discoveries. The same year he published his Essay on a new System of Comets; and the year following, his Dissertation on the weight of the air. About this time Leibnitz having published, in the *Acta Eruditorum* at Leipzig, some essays on his new *Calculus Differentialis*, but concealing the art and method of it, Mr. Bernoulli and his brother John discovered, by the little which they saw, the beauty and extent of it: this induced them to endeavour to unravel the secret; which they did with such success, that Leibnitz declared that the invention belonged to them as much as to himself.

In 1687 James Bernoulli succeeded to the professorship of mathematics at Basil; a trust which he discharged with great applause; and his reputation drew a great number of foreigners from all parts to attend his lectures. In 1699 he was admitted a foreign member of the Academy of Sciences of Paris; and in 1701 the same honour was conferred upon him by the Academy of Berlin: in both of which he published several ingenious compositions, about the years 1702, 3, and 4. He wrote also several pieces in the *Acta Eruditorum* of Leipzig, and in the *Journal des Sçavans*. His intense application to study brought upon him the gout, and by degrees a slow fever, which put a period to his life the 16th of August 1705, in the 51st year of his age. — Archimedes having found out the proportion of a sphere and its circumscribing cylinder, ordered them to be engraven on his monument: In imitation of him, Bernoulli appointed that a logarithmic spiral curve should be inscribed on his tomb, with these words, *Eadem mutata resurgo*; in allusion to the hopes of the resurrection, which are in some measure represented by the properties of that curve, which he had the honour of discovering.

James Bernoulli had an excellent genius for invention and elegant simplicity, as well as a close application. He was eminently skilled in all the branches of the mathematics, and contributed much to the promoting the new analysis, infinite series, &c. He carried to a great height the theory of the quadrature of the parabola; the geometry of curve lines, of spirals, of cycloids and epicycloids.

His works, that had been published, were collected, and printed in 2 volumes 4to, at Geneva in 1744. At the time of his death he was occupied on a great work entitled *De Arte Conjectandi*, which was published in 4to, in 1713. It contains one of the best and most elegant introductions to Infinite Series, &c. This posthumous work is omitted in the collection of his works above mentioned.

BERNOULLI (JOHN), the brother of James, last mentioned, and a celebrated mathematician, was born at Basil the 7th of August 1667. His father intended him for trade; but his own inclination was at first for the Belles-Lettres, which however, like his brother, he left for mathematics. He laboured with his brother to discover the method used by Leibnitz, in his essays on the Differential Calculus, and gave the first principles of the Integral Calculus. Our author, with Messieurs Huygens and Leibnitz, was the first who gave the solution of the problem proposed by James

Bernoulli, concerning the catenary, or curve formed by a chain suspended by its two extremities.

John Bernoulli had the degree of doctor of physic at Basil, and two years afterward was named professor of mathematics in the university of Groningen. It was here that he discovered the mercurial phosphorus or luminous barometer; and where he resolved the problem proposed by his brother concerning Isoperimetricals.

On the death of his brother James, the professor at Basil, our author returned to his native country, against the pressing invitations of the magistrates of Utrecht to come to that city, and of the university of Groningen, who wished to retain him. The Academic Senate of Basil soon appointed him to succeed his brother, without assembling competitors, and contrary to the established practice: an appointment which he held during his whole life.

In 1714 was published his treatise on the management of ships; and in 1730, his memoir on the elliptical figure of the planets gained the prize of the Academy of Sciences. The same academy also divided the prize, for their question concerning the inclination of the planetary orbits, between our author and his son Daniel.

John Bernoulli was a member of most of the academies of Europe, and received as a foreign associate of that of Paris in 1699. After a long life spent in constant study and improvement of all the branches of the mathematics, he died full of honours the 1st of January 1748, in the 81st year of his age. Of five sons which he had, three pursued the same sciences with himself. One of these died before him; the two others, Nicolas and Daniel, he lived to see become eminent and much respected in the same sciences.

The writings of this great man were dispersed through the periodical memoirs of several academies, as well as in many separate treatises. And the whole of them were carefully collected and published at Lausanne and Geneva, 1742, in 4 volumes, 4to.

BERNOULLI (DANIEL), a celebrated physician and philosopher, and son of John Bernoulli last mentioned, was born at Groningen Feb. the 9th, 1700, where his father was then professor of mathematics. He was intended by his father for trade, but his genius led him to other pursuits. He passed some time in Italy; and at 24 years of age he declined the honour offered him of becoming president of an academy intended to have been established at Genoa. He spent several years with great credit at Petersburg; and in 1733 returned to Basil, where his father was then professor of mathematics; and here our author successively filled the chair of physic, of natural and of speculative philosophy.

In his work *Exercitationes Mathematicæ*, 1724, he took the only title he then had, viz, "Son of John Bernoulli," and never would suffer any other to be added to it. This work was published in Italy, while he was there on his travels; and it classed him in the rank of inventors. In his work, *Hydrodynamica*, published in 4to at Argentoratum or Strasbourg, in 1738, to the same title was also added that of *Med. Prof. Basil.*

Daniel Bernoulli wrote a multitude of other pieces, which have been published in the Mem. Acad. of Sciences at Paris, and in those of other Academies. He gained and divided ten prizes from the Academy of Sciences,



Sciences, which were contended for by the most illustrious mathematicians in Europe. The only person who has had similar success of the same kind, is Euler, his countryman, disciple, rival, and friend. His first prize he gained at 24 years of age. In 1734 he divided one with his father; which hurt the family union; for the father considered the contest itself as a want of respect; and the son did not sufficiently conceal that he thought (what was really the case) his own piece better than his father's. And besides, he declared for Newton, against whom his father had contended all his life. In 1740 our author divided the prize, "On the Tides of the Sea," with Euler and Maclaurin. The Academy at the same time crowned a fourth piece, whose chief merit was that of being Cartesian; but this was the last public act of adoration paid by the Academy to the authority of the author of the Vortices, which it had obeyed but too long. In 1748 Daniel Bernoulli succeeded his father John in the Academy of Sciences, who had succeeded his brother James; this place, since its first erection in 1699, having never been without a Bernoulli to fill it.

Our author was extremely respected at Basil; and to bow to Daniel Bernoulli, when they met him in the streets, was one of the first lessons which every father gave every child. He was a man of great simplicity and modesty of manners. He used to tell two little adventures, which he said had given him more pleasure than all the other honours he had received. Travelling with a learned stranger, who, being pleased with his conversation, asked his name; "I am Daniel Bernoulli," answered he with great modesty; "And I," said the stranger (who thought he meant to laugh at him), "am Isaac Newton." Another time having to dinner with him the celebrated Koenig the mathematician, who boasted, with some degree of self-complacency, of a difficult problem he had resolved with much trouble, Bernoulli went on doing the honours of his table, and when they went to drink coffee he presented Koenig with a solution of the problem more elegant than his own.

After a long, useful, and honourable life, Daniel Bernoulli died the 17th of March 1782, in the 83d year of his age.

BETELGEUSE, a fixed star of the first magnitude in the right shoulder of Orion.

BEZOUT (STEPHEN), a celebrated French mathematician, Member of the Academies of Sciences and the Marine, and Examiner of the Guards of the Marine and of the Eleves of Artillery, was born at Nemours the 31st of March 1730. In the course of his studies he met with some books of geometry, which gave him a taste for that science; and the *Eloges of Fontenelle*, which shewed him the honours attendant on talents and the love of the sciences. His father in vain opposed the strong attachment of young Bezout to the mathematical sciences. April 8, 1758, he was named adjoint-mechanician in the French academy of sciences; having before that sent them two ingenious memoirs on the integral calculus, and given other proofs of his proficiency in the sciences. In 1763, he was named to the new office of Examiner to the Marine, and appointed to compose a Course of Mathematics for their use; and in 1768, on the death of

M. Camus, he succeeded as Examiner of the Artillery Eleves.

Bezout fixed his attention more particularly to the resolution of algebraic equations; and he first found out the solution of a particular class of equations of all degrees. This method, different from all former ones, was general for the cubic and biquadratic equations, and just became particular only at those of the 5th degree. Upon this work of finding the roots of equations, our author laboured from 1762 till 1779, when he published it. He composed two courses of mathematics; the one for the Marine, the other for the Artillery: The foundation of these two works was the same; the applications only being different, according to the two different objects: these courses have every where been held in great estimation. In his office of examiner he discharged the duties with great attention, care, and tenderness; a trait of his justice and zeal is remarkable in the following instance: During an examination which he held at Toulon, he was told that two of the pupils could not be present, being confined by the small-pox: he himself had never had that disease, and he was greatly afraid of it; but as he knew that if he did not see these two young men, it would much impede their improvement; he ventured therefore to their bed-sides, to examine them, and was happy to find them so deserving of the hazard he put himself into for their benefit.

Mr. Bezout lived thus several years beloved of his family and friends, and respected by all, enjoying the fruits and the credit of his labours. But the trouble and fatigues of his offices, with some personal chagrins, had reduced his strength and constitution; he was attacked by a malignant fever, of which he died Sept. 27, 1783, in the 54th year of his age, regretted by his family, his friends, the young students, and by all his acquaintance in general.

The books published by him, were:

1. Course of Mathematics for the use of the Marine, with a Treatise on Navigation; 6 vols. in 8vo, Paris, 1764.

2. Course of Mathematics for the Corps of Artillery; 4 vols. in 8vo, 1770.

3. General Theory of Algebraic Equations; 1779.

His papers printed in the volumes of the *Memoirs of the Academy of Sciences*, are:

1. On curves whose rectification depends on a given quantity; in the vol. for 1758.

2. On several classes of equations that admit of an algebraic solution; 1762.

3. First vol. of a course of mathematics, 1764.

4. On certain equations, &c; 1764.

5. General resolution of all equations; 1765.

6. Second vol. of a course of mathematics; 1765.

7. Third vol. of the same; 1766.

8. Fourth vol. of the same; 1767.

9. Integration of differentials, &c. vol. 3, Sav. Etr.

10. Experiments on cold; 1777.

BIANCHINI (FRANCIS), a very learned Italian philosopher and mathematician of the 17th century, was born at Verona the 13th of December 1662. He was much esteemed by the learned, and was a member of several academies; and was even the founder of that at Verona, called the Academy of Aletofilii, or Lovers



Lovers of Truth. He went to Rome in 1684; and was made librarian to cardinal Ottoboni, who was afterwards Pope by the name of Alexander the 8th. He entered into the church, and became canon of St. Mary de la Rotondo, and afterward of St. Lawrence in Damaso.

Bianchini was author of several learned and ingenious dissertations. In 1697 he published *La Istoria universale provata con monumenti, & figurata con simboli de gli antichi*. In 1701 pope Clement the 11th named him secretary of the conferences for the reformation of the calendar; and he published in 1703, *De Calendario & Cyclo-Cæsaris, ac de Canone Paschali sancti Hyppoliti, Martyris, Dissertationes duæ*. Bianchini was employed on the construction of the large gnomon in the church of the Chartreux at Rome, upon which he published an ample dissertation intitled, *De Nummo & Gnomone Clementino*. The research concerning the parallax and the spots of Venus occupied him a long time; but his most remarkable discovery is that of the parallelism of the axis of Venus in her orbit. He proposed to trace a meridian line through the whole extent of Italy. He was admitted a foreign Associate in the Paris Academy of Sciences, in 1706; and he had many astronomical dissertations inserted in their Memoirs, particularly in those of the years 1702, 1703, 1704, 1706, 1707, 1708, 1713, and 1718.—Bianchini died the 2d of March 1729, in the 67th year of his age.

**BIMEDIAL Line**, is the sum of two Medials. Euclid reckons two of these bimedials, in pr. 38 and 39 lib. x; the first is when the rectangle is rational, which is contained by the two medial lines whose sum makes the bimedral; and the second when that rectangle is a medial, or contained under two lines that are commensurable only in power.

**BINARY Number**, that which is composed of two units.

**BINARY Arithmetic**, that in which two figures or characters, viz, 1 and 0, only are used; the cipher multiplying every thing by 2, as in the common arithmetic by ten: thus, 1 is one, 10 is 2, 11 is 3, 100 is 4, 101 is 5, 110 is 6, 111 is 7, 1000 is 8, 1001 is 9, 1010 is ten; being founded on the same principles as common arithmetic.—This sort of arithmetic was invented by Leibnitz, who pretended that it is better adapted than the common arithmetic, for discovering certain properties of numbers, and for constructing tables; but he does not venture to recommend it, for ordinary use, on account of the great number of places of figures requisite to express all numbers, even very small ones. Jos. Pelican of Prague has more largely explained the principles and practice of the binary arithmetic, in a book entitled *Arithmeticus Perfectus, qui tria numerare nescit*; 1712. And De Lagni proposed a new system of logarithms, on the plan of the binary arithmetic; which he finds shorter, and more easy and natural than the common ones.

**BINOCLE**, or **BINOCULAR TELESCOPE**, is one by which an object is viewed with both eyes at the same time. It consists of two tubes, each furnished with glasses of the same power, by which means it has been said to shew objects larger and more clearly than a mo-

nocular or single telescope; though this is probably only an illusion, occasioned by the stronger impression which two equal images, alike illuminated, make upon the eyes; and they have been found more embarrassing than useful in practice. This telescope has been chiefly treated of by the fathers Reita and Cherubin of Orleans.—There are also microscopes of the same kind, though but little used; being subject to the same inconveniences as the telescopes.

**BINOMIAL**, a quantity consisting of two terms or members connected by the sign + or −, viz, plus or minus; as  $a + b$ , or  $3a - 2c$ , or  $a^2 + b$ , or  $x^2 - 2\sqrt{c}$ , &c; denoting the sum or the difference of the two terms. But the difference is also sometimes named a *residual*, and by Euclid an *apotome*. The term binomial was first introduced by Robert Recorde; see his algebra, pa. 46<sup>2</sup>.

**BINOMIAL Line**, or *Surd*, is that in which at least one of the parts is a surd. Euclid enumerates six kinds of binomial lines or surds, in the 10th book of his Elements, which are exactly similar to the six residuals or apotomes there treated of also, and of which an account is given under the art. **APOTOME**, which see. Those apotomes become binomials by only changing the sign of the latter term from minus to plus, which therefore are as below.

*Euclid's 6 Binomial Lines.*

- First binomial  $3 + \sqrt{5}$ ,
- 2d binomial  $\sqrt{18} + 4$ ,
- 3d binomial  $\sqrt{24} + \sqrt{18}$ ,
- 4th binomial  $4 + \sqrt{3}$ ,
- 5th binomial  $\sqrt{6} + 2$ ,
- 6th binomial  $\sqrt{6} + \sqrt{2}$ .

*To extract the Square Root of a Binomial*, as of  $a + \sqrt{b}$ , or  $\sqrt{c} + \sqrt{b}$ . Various rules have been given for this purpose. The first is that of Lucas De Burgo, in his *Summa de Arith.* &c, which is this: When one part, as  $a$ , is rational, divide it into two parts such, that their product may be equal to  $\frac{1}{4}$ th of the number under the radical  $b$ ; then shall the sum of the roots of those parts be the root of the binomial sought: or their difference is the root when the quantity is residual. That is, if  $c + e = a$ , and  $ce = \frac{1}{4}b$ ; then is  $\sqrt{c} + \sqrt{e} = \sqrt{a + \sqrt{b}}$  the root sought. As if the binomial be  $23 + \sqrt{448}$ ; then the parts of 23 are 16 and 7, and their product is 112, which is  $\frac{1}{4}$ th of 448; therefore the sum of their roots  $4 + \sqrt{7}$  is the root sought of  $23 + \sqrt{448}$ .

De Burgo gives also another rule for the same extractions, which is this: The given binomial being, for ex-

ample,  $\sqrt{c} + \sqrt{b}$ , its root will be  $\sqrt{\frac{1}{2}\sqrt{c} + \frac{1}{2}\sqrt{c} - b} + \sqrt{\frac{1}{2}\sqrt{c} - \frac{1}{2}\sqrt{c} - b}$ .—So in the foregoing example,  $23 + \sqrt{448}$ , here  $\sqrt{c} = 23$ , and  $\sqrt{b} = \sqrt{448}$ ; hence  $\frac{1}{2}\sqrt{c} = 11\frac{1}{2}$ , and  $\frac{1}{2}\sqrt{c} - b = \frac{1}{2}\sqrt{23^2 - 448} = \frac{1}{2}\sqrt{81} = 4\frac{1}{2}$ ;

theref.  $\sqrt{\frac{1}{2}\sqrt{c} + \frac{1}{2}\sqrt{c} - b} = \sqrt{11\frac{1}{2} + 4\frac{1}{2}} = \sqrt{16} = 4$ ,

and  $\sqrt{\frac{1}{2}\sqrt{c} - \frac{1}{2}\sqrt{c} - b} = \sqrt{11\frac{1}{2} - 4\frac{1}{2}} = \sqrt{7}$ ; conseq.  $4 + \sqrt{7}$  is the root sought, as before. Again, if the binomial be  $\sqrt{18} + \sqrt{10}$ ; here  $c = 18$ ,  
and



and  $b = 10$ ; theref.  $\frac{1}{2}\sqrt{c} = \frac{1}{2}\sqrt{18} = \frac{3}{2}\sqrt{2}$ , and  $\frac{1}{2}\sqrt{c-b} = \frac{1}{2}\sqrt{8} = \sqrt{2}$ ; hence,

$$\sqrt{\frac{1}{2}\sqrt{c} + \frac{1}{2}\sqrt{c-b}} = \sqrt{\frac{3}{2}\sqrt{2} + \sqrt{2}} = \sqrt{\frac{5}{2}\sqrt{2}} = \sqrt[4]{\frac{25}{2}}, \text{ and}$$

$$\sqrt{\frac{1}{2}\sqrt{c} - \frac{1}{2}\sqrt{c-b}} = \sqrt{\frac{3}{2}\sqrt{2} - \sqrt{2}} = \sqrt{\frac{1}{2}\sqrt{2}} = \sqrt[4]{\frac{1}{2}};$$

conseq.  $\sqrt[4]{\frac{25}{2}} + \sqrt[4]{\frac{1}{2}}$  or  $\frac{\sqrt{5+1}}{\sqrt[4]{2}}$  is the root of  $\sqrt{18} + \sqrt{10}$  sought. And this latter rule has been used by all other authors, down to the present time.

*To extract the Cubic and other higher Roots of a Binomial.* This is useful in resolving cubic and higher equations, and was introduced with the resolution of those equations by Tartalea and Cardan. The rules for such extractions are in great measure tentative; and some of the principal ones are the following.

*Tartalea's Rule for the Cube Root of a Binomial  $p+q$ .*

This rule is given in his 9th book of Miscellaneous Questions, quest. 40; and it is made out from either of the terms,  $p$  or  $q$ , of the binomial, taken singly, in this manner: Separate either term, as  $p$ , into two such parts that the one of them may be a cubic number, and the other part divisible by 3 without a remainder; then the cube root of the said cubic part will be one term of the root, and the other term will be the square root of the quotient arising from dividing the afore-said third part by the first term just found. So if  $p$  be divided into  $r^3 + 3s$ , then the root is  $r + \sqrt{\frac{s}{r}}$ . For

example, to extract the cube root of  $\sqrt{108} + 10$ . Suppose the part 10 be taken: this separates into the parts 1 and 9, the former of which is a cube, and the latter divisible by 3; that is  $r^3 = 1$ , and  $3s = 9$ ;

hence  $r = 1$ , and  $s = 3$ ; consequently  $r + \sqrt{\frac{s}{r}}$

$= 1 + \sqrt{3}$  is the cubic root of  $\sqrt{108} + 10$  sought. Again, to use the other term  $\sqrt{108}$ : this divides into  $\sqrt{27} + \sqrt{27}$ , of which the former is a cube, and the latter divisible by 3; that is,  $r^3 = \sqrt{27}$ , and  $3s = \sqrt{27}$ ; therefore  $r = \sqrt{3}$ , and  $s = \frac{1}{3}\sqrt{27}$

$= \sqrt{3}$  also: hence  $r + \sqrt{\frac{s}{r}} = \sqrt{3} + \sqrt{\frac{\sqrt{3}}{\sqrt{3}}} =$

$\sqrt{3} + 1$  the cube root, the same as before.

*Bombelli's Rule for the Cube Root of the Binomial*

$a + \sqrt{-b}$ . First find  $\sqrt[3]{a^2 + b}$ ; then, by trials, search out a number  $c$ , and a square root  $\sqrt{d}$ , such that the sum of their squares  $c^2 + d$  be  $= \sqrt[3]{a^2 + b}$ , and also  $c^3 - 3cd$  be  $= a$ ; then shall  $c + \sqrt{-d}$  be the cube root of  $a + \sqrt{-b}$  sought. For example, to find the cube root of  $2 + \sqrt{-121}$ : here  $\sqrt[3]{a^2 + b} = \sqrt[3]{125} = 5$ ; then taking  $c = 2$ , and  $d = 1$ , it is  $c^2 + d = 5 = \sqrt[3]{a^2 + b}$ , and  $c^3 - 3cd = 8 - 6 = 2 = a$ , as it ought; therefore  $2 + \sqrt{-1}$  is the cube root of  $2 + \sqrt{-121}$  sought.—Bombelli gave also a rule for the cube root of the binomial  $a + \sqrt{b}$ , but it is good for nothing.

*Albert Girard's Rule for the Cube Root of a Binomial.* This is given in his *Invention Nouvelle en l'Algebre*, and is explained by him thus: Let  $72 + \sqrt{5120}$  be the given binomial whose cubic root is sought.

The square of 72 the greater term is 5184  
and of the less term is 5120  
their difference 64  
its cube root 4,

which 4 must be the difference between the squares of the two terms of the root sought; and as the rational part 72 of the given binomial is the greater term, therefore the rational part of the required root will be the greater part also; consequently  $2 + \sqrt{0}$  the cubic root sought must be one of the  $3 + \sqrt{5}$  binomials here set in the margin, where  $4 + \sqrt{12}$  the difference of the squares of the terms  $5 + \sqrt{20}$  is always 4, as required; and to find out which of them it must be, proceed thus: The first,  $2 + \sqrt{0}$  must be rejected, because one term of it is 0 or nothing; also because 5 exceeds the cube root of 72, or  $\sqrt{20}$  exceeds the cube root of  $\sqrt{5120}$ , therefore  $5 + \sqrt{20}$ , and all after it must be rejected too; so that the root must be either  $3 + \sqrt{5}$  or  $4 + \sqrt{12}$ , if the given quantity has a binomial root: to know which of these is to be taken, it must be considered that the rational term of the root must measure the rational term given; and also the irrational term of the root must measure the irrational term given; then, on examination it is found that both 3 and 4 measure or divide the 72 without a remainder, but that only the  $\sqrt{5}$ , and not  $\sqrt{12}$ , measures  $\sqrt{5120}$ ; consequently none but  $3 + \sqrt{5}$  can be the cube root of the given quantity  $72 + \sqrt{5120}$ ; which is found to answer, by cubing the said root  $3 + \sqrt{5}$ .

*Dr. Wallis's Rule for the Cube Root of a Binomial  $a \pm m\sqrt{b}$  or  $a \pm m\sqrt{-b}$ .* In these forms the greatest rational part  $m$  is extracted out of the radical part, leaving only  $b$  the least radical part possible under the radical sign. He then observes that if the given quantity have a binomial root, it must be of this form  $c \pm n\sqrt{b}$ , with the same radical  $b$ . Then to find the value of  $c$  and  $n$ , he raises this root to the 3d power, which gives  $c^3 + 3cn^2b \pm 3c^2n + n^3b \cdot \sqrt{b}$ , which must be  $= a \pm m\sqrt{b}$  the given quantity; hence putting the rational part of the one quantity equal to that of the other, and also the radical part of the one equal to that of the other, gives  $c^3 + 3cn^2b = a$ , and  $3c^2n + n^3b = m$ . Then assuming several values of  $n$ , from the last equation he finds the value of  $c$ ; hence if these values of  $c$  and  $n$ , substituted in the first equation, make it just, they are right; but if not, another value of  $n$  is assumed, and so on, till the first equation hold true. And it is to be noted that  $n$  is always an integer or else the half of an integer. For example, if the cube root of  $135 \pm \sqrt{1825}$  be required, or  $135 \pm 78\sqrt{3}$ ; here  $a = 135$ ,  $m = 78$ , and  $b = 3$ ; hence  $3c^2n + n^3b = m$  is  $3c^2n + 3n^3 = 78$ , or  $c^2n + n^3 = 26$ ; then assuming  $n = 1$ , this last equation becomes  $c^2 + 1 = 26$ , from whence  $c$  is found  $= 5$ ; which values of  $c$  and  $n$  being substituted in the first equation  $c^3 + 3cn^2b = a$ , makes  $5^3 + 3 \cdot 5 \cdot 3 = 170$ , but ought to be 135, shewing that  $c$  is too great, and consequently  $n$  taken too little. Let  $n$  therefore be assumed  $= 2$ , so shall  $2c^2 + 8 = 26$ , and  $c = 3$ ; and the first equation becomes  $3^3 + 3 \cdot 3 \cdot 2^2 \cdot 3 = 27 \cdot 5 = 135 = a$  as it ought, which shews that the true value of  $n$  is 2, and that of  $c$  is 3; hence then the cube root of  $135 \pm 78\sqrt{3}$  or  $c \pm n\sqrt{b}$  is



$3 \pm 2\sqrt{3}$  or  $3 \pm \sqrt{12}$ . And in like manner is the process instituted when the number in the radical is negative, as the cube root of  $81 \pm 30\sqrt{-3}$ , which is  $\frac{9}{2} \pm \frac{5}{2}\sqrt{-3}$ .

Another rule for extracting the cube root of an imaginary binomial was also given by Demoivre, at the end of Saunderson's Algebra, by means of the trisection of an arc or angle.

*Sir I. Newton's Rule for any Root of a Binomial*  $a \pm b$ . In his Universal Arith. is given a rule for the square root of a binomial, which is the same as the 2d of Lucas de Burgo, before given; and also a general rule for any root of a binomial, which I have not met with elsewhere; and it is this: Of the given quantity  $a \pm b$ , let  $a$  be the greater term, and  $c$  the index of the root to be extracted. Seek the least number  $n$  whose power  $n^c$  can be divided by  $aa - bb$  without a remainder, and let the quotient be  $q$ ; Compute

$\sqrt[n]{a + b} \cdot \sqrt[n]{q}$  in the nearest integer number, which call  $r$ ; divide  $a\sqrt[n]{q}$  by its greatest rational divisor, calling the quotient  $s$ ; and let the nearest integer number

above  $\frac{r + \frac{n}{2}}{2s}$  be  $t$ : so shall  $\frac{ts \pm \sqrt{t^2 s^2 - n}}{2\sqrt[n]{q}}$  be the root

sought, if the root can be extracted. And this rule is demonstrated by s'Gravesande in his commentary on Newton's Arithmetic. And many numeral examples, illustrating this rule, are given in s'Gravesande's Algebra, above-mentioned, pa. 160, as also in Newton's Univers. Arith. pa. 53 2d edit. and in Maclaurin's Algebra pa. 118. Other rules may be found in Schooten's Commentary on the Geometry of Descartes, and elsewhere.

*Impossible or Imaginary BINOMIAL*, is a binomial which has one of its terms an impossible or an imaginary quantity; as  $a + \sqrt{-b}$ .

*BINOMIAL Curve*, is a curve whose ordinate is expressed by a binomial quantity; as the curve whose ordinate is  $x^2 \times b + dx^2$ . Stirling, Method. Diff. pa. 58.

*BINOMIAL Theorem*, is used to denote the celebrated theorem given by Sir I. Newton for raising a binomial to any power, or for extracting any root of it by an approximating infinite series. It was known by Stifelius, and others, about the beginning of the 16th century, how to raise the integral powers, not barely by a continued multiplication of the binomial given, but Stifelius formed also a table of numbers by a continued addition, which shewed by inspection the coefficients of the terms of any power of the binomial, contained within the limits of the table; but still they could not independent of a table, and of any of the lower powers, raise any power of a binomial at once, by determining its terms one from another only, viz, the 2d term from the 1st, the 3d from the 2d, and so on as far as we please, by a general rule; and much less could they extract general algebraic roots in infinite series by any rule whatever.

For although the nature and construction of that table, which is a table of figurate numbers, was so early known, and employed in the raising of powers, and the extracting the roots of pure numbers; yet it was only by raising the numbers one from another by continual additions, and then taking them from the table

when wanted, till Mr. Briggs first pointed out the way of raising any line in the table by itself, without any of the preceding lines; and thus teaching to raise the terms of any power of a binomial, independent of any of the other powers; and so gave the substance of the binomial theorem in words, wanting only the algebraic notation in symbols; as is shewn at large at pa. 75 of the historical introduction to my Mathematical Tables. Whatever was known however of this matter, related only to pure or integral powers, no one before Newton having thought of extracting roots by infinite series. He happily discovered that, by considering powers and roots in a continued series, roots being as powers having fractional exponents, the same binomial series would equally serve for them all, whether the index should be fractional or integral, or whether the series be finite or infinite. The truth of this method however was long known only by trial in particular cases, and by induction from analogy; nor does it appear that even Newton himself ever attempted any direct proof of it: however, various demonstrations of the theorem have since been given by the more modern mathematicians, some of which are by means of the doctrine of fluxions, and others, more legally, from the pure principles of algebra only: for a full account of which, see pa. 71 &c, of my Mathematical Tracts, vol. 1.

This theorem was first discovered by Sir I. Newton in 1669, and sent in a letter of June 13, 1676, to Mr. Oldenburgh, Secretary to the Royal Society, to be by him communicated to Mr. Leibnitz; and it was in this

form:  $\overline{p + pq}^{\frac{m}{n}} = p^{\frac{m}{n}} + \frac{m}{n} aq + \frac{m-n}{2n} bq + \frac{m-2n}{3n} cq + \frac{m-3n}{4n} dq + \&c$ : where  $p + pq$  signifies the

quantity whose root, or power, or root of any power, is to be found;  $p$  being the first term of that quantity;  $q$  the quotient of all the rest of the terms divided by that first term; and  $\frac{m}{n}$  the numeral index of the power

or root of the quantity  $p + pq$ , whether it be integral or fractional, positive or negative; and lastly  $a, b, c, d, \&c$ , are assumed to denote the several terms in their order as they are found, viz,  $a$  = the first term  $p^{\frac{m}{n}}$ ,  $b$  = the 2d term  $\frac{m}{n} aq$ ,  $c$  = the 3d term  $\frac{m-n}{2n} bq$ , and so on. As Newton's general notation of indices was not commonly known, he takes this occasion to explain it; and then he gives many examples of the application of this theorem, one of which is the following.

*Ex. 1.* To find the value of  $\sqrt{c^2 + x^2}$  or  $c^2 + x^2$   $^{\frac{1}{2}}$ , that is, to extract the square root of  $c^2 + x^2$  in an infinite series. Here  $p = c^2$ ,  $q = \frac{x^2}{c^2}$ ,  $m = 1$ , and

$n = 2$ ; therefore  $a = p^{\frac{m}{n}} = \overline{cc}^{\frac{1}{2}} = c$ ,  $b = \frac{m}{n} aq$

$= \frac{x^2}{2c}$ ,  $c = \frac{m-n}{2n} bq = \frac{-x^4}{8c^3}$ , &c; and therefore

the root sought is  $c + \frac{x^2}{2c} - \frac{x^4}{8c^3} + \frac{x^6}{16c^5} - \frac{5x^8}{128c^7} + \&c$ .

A variety



A variety of other examples are also given in the same place, by which it is shewn that the theorem is of universal application to all sorts of quantities whatever.

This theorem is sometimes represented in other forms, as

$$p^{\frac{m}{n}} \times : 1 + \frac{m}{n} q + \frac{m}{n} \cdot \frac{m-n}{2n} q^2 + \frac{m}{n} \cdot \frac{m-n}{2n} \cdot \frac{m-2n}{3n} q^3 + \&c; \text{ which comes to the same thing. Or}$$

$$\text{also thus } p^{\frac{m}{n}} + \frac{m}{n} p^{\frac{m-n}{n}} r + \frac{m}{n} \cdot \frac{m-n}{2n} p^{\frac{m-2n}{n}} r^2 + \&c;$$

where the binomial is  $p + r$ .

In another letter to Mr. Oldenburgh, of Oct. 24, 1676, Newton explains the train of reasoning by which he obtained the said theorem, as follows: "In the beginning of my mathematical studies, when I was perusing the works of the celebrated Dr. Wallis (see his Arith. of Infinites, prop. 118, and 121, also his Algebra chap. 82), and considering the series by the interpolation of which he exhibits the area of the circle and hyperbola; for instance, in this series of curves, whose common base or axis is  $x$ , and the ordinates respectively  $\sqrt{1-xx}$ ,  $\sqrt{1-xx^2}$ ,  $\sqrt{1-xx^3}$ ,  $\sqrt{1-xx^4}$ ,  $\&c$ ; I perceived that if the areas of the alternate curves, which

are  $x$ ,

$$x - \frac{1}{3}x^3,$$

$$x - \frac{2}{5}x^3 + \frac{1}{5}x^5,$$

$$x - \frac{3}{7}x^3 + \frac{3}{5}x^5 - \frac{1}{7}x^7,$$

&c; could be interpolated, we should obtain the areas of the intermediate ones; the first of which, or

$\sqrt{1-xx}$ , is the area of the circle: now in order to this, it appeared that in all the series the first term was  $x$ ; that the 2d terms  $\frac{1}{3}x^3$ ,  $\frac{2}{5}x^3$ ,  $\frac{3}{7}x^3$ ,  $\&c$ , were in arithmetical progression; and consequently that the first two terms of all the series to be interpolated would be

$$x - \frac{1}{3}x^3, \quad x - \frac{2}{5}x^3, \quad x - \frac{3}{7}x^3, \quad \&c.$$

"Now for the interpolation of the rest, I considered that the denominators 1, 3, 5, 7, &c, were in arithmetical progression; and that therefore only the numeral coefficients of the numerators were to be investigated. But these in the alternate areas, which are given, were the same with the figures of which the several powers of 11 consist, viz, of 11<sup>0</sup>, 11<sup>1</sup>, 11<sup>2</sup>, 11<sup>3</sup> &c; that is, the first 1,

the second 1, 1

the third 1, 2, 1

the fourth 1, 3, 3, 1

the fifth 1, 4, 6, 4, 1

&c.

"I enquired therefore how, in these series, the rest of the terms may be derived from the first two being given; and I found that by putting  $m$  for the 2d figure or term,

VOL. I.

the rest would be produced by the continued multiplication of the terms of this series,

$$\frac{m-0}{1} \times \frac{m-1}{2} \times \frac{m-2}{3} \times \frac{m-3}{4} \times \frac{m-4}{5} \&c.$$

"For instance, if the 2d term  $m=4$ ; then shall  $4 \times \frac{m-1}{2}$ , or 6, be the 3d term; and  $6 \times \frac{m-2}{3}$ ,

or 4, the 4th term; and  $4 \times \frac{m-3}{4}$ , or 1, the 5th

term; and  $1 \times \frac{m-4}{5}$ , or 0, the 6th; which shews

that in this case the series terminates.

"This rule therefore I applied to the series to be interpolated. And since, in the series for the circle, the

2d term was  $\frac{1}{2}x^3$  I put  $m = \frac{1}{2}$ , which produced

the terms  $\frac{1}{2} \times \frac{\frac{1}{2}-1}{2}$  or  $-\frac{1}{8}$ ;  $-\frac{1}{8} \times \frac{\frac{1}{2}-2}{3}$  or

$+\frac{1}{16}$ ;  $+\frac{1}{16} \times \frac{\frac{1}{2}-3}{4}$  or  $-\frac{5}{128}$ ; and so on *ad infinitum*. And hence I found that the required area of

the circular segment is

$$x - \frac{\frac{1}{2}x^3}{3} - \frac{\frac{1}{8}x^5}{5} - \frac{\frac{1}{16}x^7}{7} - \frac{\frac{5}{128}x^9}{9} - \&c.$$

"And in the same manner might be produced the interpolated areas of the other curves: as also the area of the hyperbola and the other alternates in this series

$$\sqrt{1+xx}, \sqrt{1+xx^2}, \sqrt{1+xx^3}, \sqrt{1+xx^4}, \&c.$$

And in the same way also may other series be interpolated, and that too if they should be taken at the distance of two or more terms.

"This was the way then in which I first entered upon these speculations; which I should not have remembered, but that in turning over my papers a few weeks since, I chanced to cast my eyes on those relating to this matter.

"Having proceeded so far, I considered that the terms  $\sqrt{1-xx^{\frac{1}{2}}}$ ,  $\sqrt{1-xx^{\frac{2}{5}}}$ ,  $\sqrt{1-xx^{\frac{3}{7}}}$ ,  $\sqrt{1-xx^{\frac{4}{9}}}$ ,  $\&c$ , that is, 1

$$1 - x^2$$

$$1 - 2x^2 + x^4$$

$$1 - 3x^2 + 3x^4 - x^6, \&c, \text{ might be interpolated}$$

in the same manner as the areas generated by them; and for this, nothing more was required but to omit the denominators 1, 3, 5, 7, &c, in the terms expressing the areas; that is, the coefficients of the terms of the

quantity to be interpolated,  $\sqrt{1-xx^{\frac{1}{2}}}$ , or  $\sqrt{1-xx^{\frac{2}{5}}}$ , or

generally  $\sqrt{1-xx^{\frac{m}{n}}}$ , will be produced by the continued multiplication of the terms of this series

$$m \times \frac{m-1}{2} \times \frac{m-2}{3} \times \frac{m-3}{4} \&c.$$

"Thus, for example, there would be found

$$\sqrt{1-xx^{\frac{1}{2}}} = 1 - \frac{1}{2}x^2 - \frac{1}{8}x^4 - \frac{1}{16}x^6 \&c.$$

$$\sqrt{1-xx^{\frac{2}{5}}} = 1 - \frac{2}{5}x^2 + \frac{3}{5}x^4 + \frac{1}{125}x^6 \&c.]$$

$$\sqrt{1-xx^{\frac{3}{7}}} = 1 - \frac{3}{7}x^2 - \frac{1}{7}x^4 - \frac{5}{2401}x^6 \&c.$$

E c

Thus



"Thus then I discovered a general method of reducing radical quantities into infinite series, by the theorem which I sent in the beginning of the former letter, before I knew the same by the extraction of roots.

"But having discovered that way, this other could not long remain unknown: for, to prove the truth of those operations, I multiplied

$1 - \frac{1}{2}x^2 - \frac{1}{8}x^4 - \frac{1}{16}x^6 \&c$ , by itself, and the product is  $1 - x^2$ , all the rest of the terms vanishing after these, *in infinitum*. In like manner,

$1 - \frac{1}{3}x^2 - \frac{1}{9}x^4 - \frac{5}{81}x^6 \&c$ , twice multiplied by itself, produced  $1 - x^2$ . But as this was a certain proof of those conclusions, so I was naturally led to try conversely whether these series, which were thus known to be the roots of the quantity  $1 - x^2$ , could not be extracted out of it after the manner of arithmetic; and upon trial I found it to succeed. The process for the square root is here set down

$$1 - x^2 \quad (1 - \frac{1}{2}x^2 - \frac{1}{8}x^4 - \frac{1}{16}x^6 \&c$$

1

$$0 - x^2$$

$$-x^2 + \frac{1}{4}x^4$$

$$-\frac{1}{4}x^4$$

$$-\frac{1}{4}x^4 + \frac{1}{8}x^6 + \frac{1}{16}x^8$$

$$-\frac{1}{8}x^6 - \frac{1}{16}x^8$$

"These methods being found, I laid aside the other way by interpolation of series, and used these operations only as a more genuine foundation. Neither was I ignorant of the reduction by division, which is so much easier." See Collins's *Commercium Epistolicum*.

And this is all the account that Newton gives of the invention of this theorem, which is engraved on his monument in Westminster Abbey, as one of his greatest discoveries.

**BIPARTIENT**, is a number that divides another into two equal parts without a remainder. So 2 is a bipartient to 4, and 5 a bipartient to 10.

**BIPARTITION**, is a division into two equal parts.

**BIQUADRATE**, or **BIQUADRATIC Power**, is the squared square, or 4th power of any number or quantity. Thus 16 is the biquadrate or 4th power of 2, or it is the square of 4 which is the 2d power of 2.

**BIQUADRATIC Root**, of any quantity, is the square root of the square root, or the 4th root of that quantity. So the biquadratic root of 16 is 2, and the biquadratic root of 81 is 3.

**BIQUADRATIC Equation**, is that which rises to 4 dimensions, or in which the unknown quantity rises to the 4th power; as  $x^4 + ax^3 + bx^2 + cx + d = 0$ .

Any biquadratic equation may be conceived to be generated or produced from the continual multiplication of four simple equations,

$$\text{as } x - p \times x - q \times x - r \times x - s = 0;$$

or from that of two quadratic equations,

$$\text{as } x^2 + px + q \times x^2 + rx + s = 0;$$

or, lastly, from that of a cubic and a simple equation,

as  $x - p \times x^3 + qx^2 + rx + s = 0$ : which was the invention of Harriot. And, on the contrary, a biquadratic equation may be resolved into four simple equations, or into two quadratics, or into a cubic and a simple equation, having all the same roots with it.

### 1. Ferrari's Method for Biquadratic Equations:

The first resolution of a biquadratic equation was given in Cardan's Algebra, chap. 39, being the invention of his pupil and friend Lewis Ferrari, about the year 1540. This is effected by means of a cubic equation, and is indeed a method of depressing the biquadratic equation to a cubic, which Cardan demonstrates, and applies in a great variety of examples. The principle is very general, and consists in completing one side of the equation up to a square, by the help of some multiples or parts of its own terms and an assumed unknown quantity; which it is always easy to do; and then the other side is made to be a square also, by assuming the product of its 1st and 3d terms equal to the square of half the 2d term; for it consists only of three terms, or three different denominations of the original letter; then this equality will determine the value of the assumed quantity by a cubic equation: other circumstances depend on the artist's judgment. But the method will be farther explained by the following examples, extracted from Cardan's book.

*Ex. 1.* Given  $x^4 + 6x^2 + 36 = 60x$ , to be resolved. Add  $6x^2$  to both sides of the equation, so shall

$$x^4 + 12x^2 + 36 \text{ or } (x^2 + 6)^2 = 6x^2 + 60x.$$

Assume  $y$ , and add  $x^2 + 6 \cdot 2y + y^2$  to both sides, then is  $(x^2 + 6)^2 + x^2 + 6 \cdot 2y + y^2 = 6x^2 + 60x + x^2 + 6 \cdot 2y + y^2$ , or  $(x^2 + 6 + y)^2 = 6 + 2y \cdot x^2 + 60x + 12y + y^2$ .

Make now the 1st  $\times$  3d term  $= \frac{1}{4}$  sq. 2d,

this gives  $6 + 2y \cdot 12y + y^2 \cdot x^2 = 900x^2$ ,

or  $y^3 + 15y^2 + 36y = 450$ ; and hence

$$y = \sqrt[3]{287\frac{1}{2}} + \sqrt{80449\frac{1}{4}} + \sqrt[3]{287\frac{1}{2}} - \sqrt{80449\frac{1}{4}} - 5.$$

From which  $x$  may be found by a quadratic equation.

*Ex. 2.* Given  $x^4 = x + 2$ .

Before applying Ferrari's method to this example, Cardan resolves it by another way as follows: subtract 1, then is  $x^4 - 1 = x + 1$ ; divide by  $x + 1$ , then is  $x^3 - x^2 + x - 1 = 1$ , or  $x^3 + x = x^2 + 2$ ; and hence

$$x = \sqrt[3]{\sqrt{\frac{2241}{2916}} + \frac{47}{54}} - \sqrt[3]{\sqrt{\frac{2241}{2916}} - \frac{47}{54}} + \frac{1}{3}.$$

But to resolve it by Ferrari's rule:

Because  $x^4 = x + 2$ , therefore

$x^4 + 2yx^2 + y^2$  or  $(x^2 + y)^2 = 2yx^2 + x + 2 + y^2$ ; hence  $2y^3 + 4y = \frac{1}{4}$ , or  $y^3 + 2y = \frac{1}{8}$ ; and the root is

$$y = \sqrt[3]{\sqrt{\frac{2075}{6912}} + \frac{1}{16}} - \sqrt[3]{\sqrt{\frac{2075}{6912}} - \frac{1}{16}};$$

by means of which  $x$  is found by a quadratic equation.

*Ex. 3.* Given  $x^4 + 32x^2 + 16 = 48x$ .—Add 240, then  $x^4 + 32x^2 + 256$  or  $(x^2 + 16)^2 = 48x + 240$ ; complete square again, then

$$(x^2 + 16)^2 + 2y \cdot x^2 + 16 + y^2 = 2yx^2 + 48x + y^2 + 32y + 240;$$

make the last side a sq. by the rule, which gives

$$y^3 + 32y^2 + 240y = 24 \times 12 = 288.$$

Put now  $z = y + 10\frac{2}{3}$ , and the last transforms to  $z^3 = 101\frac{1}{3}z + 420\frac{1}{27}$ ; then the value of  $z$  found from this, gives the value of  $y$ , and hence the value of  $x$ , as before.



## 2. Descartes's Rule for Biquadratic Equations.

Another solution was given of biquadratic equations by Descartes, in the 3d book of his Geometry. In this solution he resolved the given biquadratic equation into two quadratics, by means of a cubic equation, in this manner: First, let the 2d term or 3d power be taken away out of the equation, after which it will stand thus,

$$x^4 + px^2 + qx + r = 0. \text{ Find } y \text{ in this}$$

cubic equation  $y^6 + 2py^4 + \frac{p^2}{4r}y^2 - q^2 = 0$ ; and then

the values of  $x$  in these two quadratics.

$$\begin{cases} x^2 - yx + \frac{1}{2}y^2 + \frac{1}{2}p + \frac{q}{2y} = 0 \\ x^2 + yx + \frac{1}{2}y^2 + \frac{1}{2}p - \frac{q}{2y} = 0, \end{cases}$$

and these values of  $x$  will be the roots of the given biquadratic equation.

*Ex.* Let the equ. be  $x^4 - 17x^2 - 20x - 6 = 0$ . Hence  $p = -17$ ,  $q = -20$ , &  $r = -6$ ; and the cubic equ. is  $y^6 - 34y^4 + 313y^2 - 400 = 0$ , the root of which is  $y^2 = 16$ , or  $y = 4$ ; hence the two quadratics are  $\begin{cases} x^2 - 4x - 3 = 0 \\ x^2 + 4x + 2 = 0, \end{cases}$  the four roots of which are  $2 \pm \sqrt{7}$  and  $-2 \pm \sqrt{2}$ .

## 3. Euler's Method for Biquadratic Equations.

The celebrated Leonard Euler gave, in the 6th volume of the Petersburg Ancient Commentaries, for the year 1738, an ingenious and general method of resolving equations of all degrees, by means of the equation of the next lower degree, and among them of the biquadratic equation by means of the cubic; and this last was also given more at large in his treatise of Algebra, translated from the German into French in 1774, in 2 volumes 8vo. The method is this: Let  $x^4 - ax^2 - bx - c = 0$ , be the given biquadratic equation, wanting the 2d term. Take  $f = \frac{1}{2}a$ ,  $g = \frac{1}{16}aa + \frac{1}{4}c$ , and  $h = \frac{1}{8}bb$ ; with which values of  $f, g, h$ , form the cubic equation  $z^3 - fz^2 + gz - h = 0$ . Find the three roots of this cubic equation, and let them be called  $p, q, r$ . Then shall the four roots of the proposed biquadratic be these following, viz,

When $\frac{1}{8}b$ is positive:	When $\frac{1}{8}b$ is negative:
1st. $\sqrt{p} + \sqrt{q} + \sqrt{r}$	$\sqrt{p} + \sqrt{q} - \sqrt{r}$ ,
2d. $\sqrt{p} + \sqrt{q} - \sqrt{r}$	$\sqrt{p} - \sqrt{q} + \sqrt{r}$ ,
3d. $\sqrt{p} - \sqrt{q} + \sqrt{r}$	$-\sqrt{p} + \sqrt{q} + \sqrt{r}$ ,
4th. $\sqrt{p} - \sqrt{q} - \sqrt{r}$	$-\sqrt{p} - \sqrt{q} - \sqrt{r}$ .

*Ex.* Let the eq. be  $x^4 - 25x^2 + 60x - 36 = 0$ . Here  $a = 25$ ,  $b = -60$ , and  $c = 36$ ;

theref.  $f = \frac{25}{2}$ ,  $g = \frac{625}{16} + 9 = \frac{769}{16}$ , and  $h = \frac{225}{4}$ .

Conseq. the cubic equation will be

$$z^3 - \frac{25}{2}z^2 + \frac{769}{16}z - \frac{225}{4} = 0.$$

The three roots of which are

$$z = \frac{9}{4} = p, \text{ and } z = 4 = q, \text{ and } z = \frac{25}{4} = r;$$

the roots of which are  $\sqrt{p} = \frac{3}{2}$ ,  $\sqrt{q} = 2$  or  $\frac{4}{2}$ ,  $\sqrt{r} = \frac{5}{2}$ .

Hence, as the value of  $\frac{1}{8}b$  is negative, the four roots are

$$\begin{aligned} 1\text{st. } x &= \frac{3}{2} + \frac{4}{2} - \frac{5}{2} = 1, \\ 2\text{d. } x &= \frac{3}{2} - \frac{4}{2} + \frac{5}{2} = 2, \\ 3\text{d. } x &= -\frac{3}{2} + \frac{4}{2} + \frac{5}{2} = 3, \\ 4\text{th. } x &= -\frac{3}{2} - \frac{4}{2} - \frac{5}{2} = -6. \end{aligned}$$

## 4. Simpson's Rule for Biquadratic Equations.

Mr. Simpson gave also a general rule for the solution of biquadratic equations, in the 2d edit. of his Algebra, pa. 150, in which the given equation is also resolved by means of a cubic equation, as well as the two former ways; and it is investigated on the principle, that the given equation is equal to the difference between two squares; being indeed a kind of generalization of Ferrari's method.

Thus, he supposes the given equation, viz,

$x^4 + px^3 + qx^2 + rx + s = x^2 + \frac{1}{2}px + A^2 - Bx + C^2$ ; then from a comparison of the like terms, the values of the assumed letters are found, and the final equation becomes

$$A^3 - \frac{1}{2}qA^2 + kA - \frac{1}{2}l = 0,$$

where  $k = \frac{1}{2}pr - s$ , and  $l = \frac{1}{4}r^2 + s \cdot \frac{1}{4}p^2 - q$ .

The value of  $A$  being found in this cubic equation, from it will be had the values of  $B$  and  $C$ , which have these general values, viz,  $B = \sqrt{2A + \frac{1}{4}p^2 - q}$ , and  $C = \frac{pA - r}{2B}$ . Hence, finally, the root  $x$  will be ob-

tained from the assumed equation  $x^2 + \frac{1}{2}px + A^2 - Bx + C^2 = 0$ , or  $x^2 + \frac{1}{2}px + A = \pm Bx \pm C$ , in four several values.

*Ex.* Given the equ.  $x^4 - 6x^3 - 58x^2 - 114x - 11 = 0$ .

Here  $p = -6$ ,  $q = -58$ ,  $r = -114$ , and  $s = -11$ , whence

$k$  or  $\frac{1}{2}pr - s = 182$ ,  $l$  or  $\frac{1}{4}r^2 + s \cdot \frac{1}{4}p^2 - q = 2512$ ; and therefore the cubic equation becomes

$$A^3 + 29A^2 + 182A - 1256 = 0,$$

the root of which is  $A = 4$ .

Hence then  $B$  or  $\sqrt{2A + \frac{1}{4}p^2 - q} = \sqrt{75}$ , and  $C$  or  $\frac{pA - r}{2B} = \sqrt{27}$ :

and the quadratic equation becomes

$x^2 - 3x + 4 = \pm x\sqrt{75} \pm \sqrt{27}$ , the four roots of which are  $x = 1\frac{1}{2} \pm 2\frac{1}{2}\sqrt{3} + \sqrt{17} \pm 11\frac{1}{2}\sqrt{3}$ ,

$$\text{and } x = 1\frac{1}{2} \pm 2\frac{1}{2}\sqrt{3} - \sqrt{17} \pm 11\frac{1}{2}\sqrt{3}.$$

Mr. Simpson here subjoins an observation which it has since been found is erroneous, viz, that "The value of  $A$ , in this equation, will be *commensurate* and *rational* (and therefore the easier to be discovered), not only when all the roots of the given equation are *commensurate*, but when they are *irrational* and even *impossible*; as will appear from the examples subjoined." This is a strange reason for Simpson to give for the proof of a proposition; and it is wonderful that he fell upon no examples that disprove it, as the instances in which it holds true, are very few indeed, in comparison with the number of those in which it fails.

*Note.* In any biquadratic equation having all its terms, if  $\frac{3}{8}$  of the square of the coefficient of the 2d term be greater than the product of the coefficients of the 1st and 3d terms, or  $\frac{3}{8}$  of the square of the coefficient



of the 4th term be greater than the product of the coefficients of the 3d and 5th terms, or  $\frac{1}{2}$  of the square of the coefficient of the 3d term greater than the product of the coefficients of the 2d and 4th terms; then all the roots of that equation will be real and unequal; but if either of the said parts of those squares be less than either of those products, the equation will have imaginary roots.

For the construction of biquadratic equations, see *Construction*. See also *Descartes's Geometry*, with the *Commentaries of Schooten* and others; *Baker's Geometrical Key*; *Slusius's Mesolabium*; *L'Hospital's Conic Sections*; *Wolffius's Elementa Matheseos*; &c.


**BIQUADRATIC Parabola**, a curve of the 3d order, having two infinite legs, and expressed by one of these three equations, viz,

$$a^3x = y^4, \text{ as in fig. 1,}$$

$$a^3x = y^4 - a^2y^2, \text{ as in fig. 2,}$$

$$a^3x = y^4 - a + b \cdot y^3 - aby^2, \text{ as in fig. 3;}$$

where  $x = AP$  the absciss,  $y = PQ$  the ordinate,  $b = AB$ ,  $c = AC$ , and  $a =$  a certain given quantity.




But the most general equation of this curve is the following, which belongs to fig. 4, viz,



$$ay = x^4 + bx^3 + cx^2 + dx + e;$$

where  $x = Ap$  or  $AP$  the absciss, and  $-y$  or  $+y$  is the ordinate  $pm$  or  $PM$ , also  $a, b, c, d, e$ , are constant quantities; the beginning of the absciss being at any point  $A$  in the indefinite line  $AP$ .

But if the beginning of the absciss  $A$  be where this line intersects the curve, as in fig. 5, then the nature of the curve will be defined by this equation  $a \times pm = Ap \times Bp \times Cp \times Dp$ , wherever the point  $p$  is taken in the infinite line  $RS$ .



When the curve has no serpentine part, as fig. 6, the equation is more simple, being in this case barely

$a \times pm = Ap^2 \times pB^2$ . See *Curve Lines*, and *GEOMETRICAL Lines*.

**BIQUINTILE**, an aspect of the planets when the distance between them is 144 degrees, or twice the 5th part of 360 degrees.

**BISECTION**, or **BISSECTION**, the division of a quantity into two equal parts, otherwise called *bipartition*.

**BISSEXTILE**, or *Leap-year*, a year consisting of 366 days, and happening every 4th year, by the addition of a day in the month of February, which that year consists of 29 days. And this is done to recover the 6 hours which the sun takes up nearly in his course, more than the 365 days commonly allowed for it in other years.

The day thus added was by Julius Cæsar appointed to be the day before the 24th of February, which among the Romans was the 6th of the calends, and which on this occasion was reckoned twice; whence it was called the *bissextile*.

By the statute *De anno bissextile*, 21 Hen. III, to prevent misunderstandings, the intercalary day and that next before it are to be accounted as one day.

To find what year of the period any given year is; divide the given year by 4; then if 0 remains, it is leap year; but if any thing remain, the given year is so many after leap year.

But the astronomers concerned in reforming the calendar in 1582, by order of pope Gregory XIII, observing that in 4 years the bissextile added 44 minutes more than the sun spent in returning to the same point of the ecliptic; and computing that in 133 years these supernumerary minutes would form a day; to prevent any changes being thus insensibly introduced into the seasons, directed, that in the course of 400 years there should be three sextiles retrenched; so that every centesimal year, which is a leap year according to the Julian account, is a common year in the Gregorian account, unless the number of centuries can be divided by 4 without a remainder. So 1600 and 2000 are bissextile; but 1700, 1800, and 1900 are common years.

The Gregorian computation has been received in most foreign countries ever since the reformation of the calendar in 1582; excepting some northern countries, as Russia, &c. And by act of parliament, passed in 1751, it commenced in all the dominions under the crown of Great Britain in the year following; it being ordered by that act that the natural day next following the 2d of September, should be accounted the 14th; omitting the intermediate 11 days of the common calendar. The supernumerary day, in leap years, being added at the end of the month February, and called the 29th of that month.

**BLACK**, a colour so called, or rather a privation of all colour. This, it seems, arises from such a peculiar texture and situation of the superficial parts of a black body, that they absorb all or most of the rays of light, reflecting little or none to the eye: and hence it happens that black bodies, thus imbibing the rays, are always found to be hotter than those of a lighter colour. Dr. Franklin observes that black cloaths heat more, and dry sooner in the sun than white cloaths; that therefore black is a bad colour for cloaths in hot climates; but a fit colour for the linings of ladies' summer hats;



and that a chimney painted black, when exposed to the sun, will draw more strongly. Franklin's *Experim. &c.*—Dr. Watson, the present bishop of Landaff, covered the bulb of a thermometer with a black coating of Indian ink, and the consequence was that the mercury rose 10 degrees higher. *Philos. Transf.* vol. 63, pa. 40.—And a virtuoso of unsuspected credit assured Mr. Boyle, that in a hot climate by blacking the shells of eggs, and exposing them to the sun, he had seen them thus well roasted in a short time.

**BLACKNESS**, the quality of a black body, as to colour; arising from its stifling or absorbing the rays of light, and reflecting little or none. In which sense it stands directly opposed to whiteness; which consists in such a texture of parts, as indifferently reflects all the rays thrown upon it, of whatever colour they may be.

Descartes, it seems, first rightly distinguished these causes of black and white, though he might be mistaken with respect to the general nature of light and colours.—Sir Isaac Newton shews, in his *Optics*, that to produce black colours, the corpuscles must be smaller than for exhibiting the other colours; because, where the sizes of the component particles of a body are greater, the light reflected is too much for constituting this colour: but when they are a little smaller than is requisite to reflect the white, and very faint blue of the first order, they will reflect so little light, as to appear intensely black; and yet they may perhaps reflect it variously to and fro within them so long, till it be stifled and lost.

And hence, it appears, why fire, or putrefaction, by dividing the particles of substances, turn them black: why small quantities of black substances impart their colours very freely, and intensely, to other substances; to which they are applied; the minute particles of these, on account of their very great number, easily overspreading the gross particles of others. Hence it also appears, why glass ground very elaborately on a copper-plate with sand, till it be well polished, makes the sand, with what is rubbed off from the copper and glass, become very black: also why blacks commonly incline a little towards a blueish colour; as may be seen by illuminating white paper with light reflected from black substances, when the paper usually appears of a blueish white; the reason of which is, that black borders on the obscure blue of the first order of colours, and therefore reflects more rays of that colour than of any other: and lastly why black substances do sooner than others become hot in the sun's light, and burn; an effect which may proceed partly from the multitude of refractions in a little space, and partly from the easy commotion among such minute particles.

**BLAGRAVE** (JOHN), an eminent mathematician, who flourished in the 16th and 17th centuries. He was the second son of John Blagrove, of Bulmarsh-court near Sunning in Berkshire, descended from an ancient family in that country. From a grammar school at Reading he was sent to St. John's college in Oxford, where he applied himself chiefly to the study of mathematics. From hence he retired to his paternal seat of Southcote-lodge near Reading, where he spent the rest of his life, in a retired manner, without marrying, that he might have more leisure to pursue his favourite studies; which he did with great

application and success. After a life thus spent in study, and in acts of benevolence to all around him, he died in the year 1611; and was buried at Reading in the church of St. Lawrence, where a sumptuous monument was erected to his memory.

He left the bulk of his fortune to the posterity of his three brothers, which were very numerous. There have been mentioned various acts of his beneficence in private life, for the encouragement of learning, the reward of merit, and the relief of distress. Some of these were the result of a quaint, humorous disposition, discovered chiefly in his legacies: One of these was 10 pounds left to be annually disposed of in the following manner: On Good-friday, the churchwardens of each of the three parishes of Reading send to the town-hall "one virtuous maid who has lived five years with her master:" there, in the presence of the magistrates these three virtuous maids throw dice for the ten pounds. The year following the two losers are returned with a fresh one, and again the third year, till each has had three chances. He also left an annuity to 80 poor widows, who should attend annually on Good-friday also, and hear a sermon, for the preaching of which he left ten shillings to the minister. He took care also for the maintenance of his servants, rewarding their diligence and fidelity, and providing amply for their support. Thus it appears he was not more remarkable for his scientific knowledge, than for his generosity and philanthropy. His works are,

1. *A Mathematical Jewel.* Lond. 1585, folio.
2. *Of the Making and Use of the Familiar Staff.* Lond. 1590, 4to.
3. *Astrolabium Uranicum Generale.* Lond. 1596, 4to.
4. *The Art of Dyalling.* Lond. 1609, 4to.

**BLAIR** (JOHN), an eminent chronologist, was educated at Edinburgh. Afterward, coming to London, he was for some time usher of a school in Hedge-lane. In 1754 he first published "The Chronology and History of the World, from the Creation to the year of Christ 1753; illustrated in 56 tables. In 1755 he was elected a fellow of the Royal Society, and in 1761 of the Society of Antiquaries. In 1756 he published a 2d edition of his *Chronological Tables*; and in 1768 an improved edition of the same with the addition of 14 maps of Ancient and Modern Geography, for illustrating the *Tables of Chronology and History*; to which is prefixed a *Dissertation on the Progress of Geography*. In 1757 he was appointed chaplain to the Princess Dowager of Wales, and Mathematical Tutor to the Duke of York; whom he attended in 1763 in a tour to the continent, from which they returned the year after. Dr. Blair had successively several good church livings: as, a prebendal stall at Westminster, the vicarage of Hinckley, and the rectory of Burton Coggles in Lincolnshire, all in 1761; the vicarage of St. Bride's in London, in 1771, in exchange for that of Hinckley; the rectory of St. John the Evangelist in Westminster, in 1776, in exchange for the vicarage of St. Bride's; in the same year the rectory of Horton near Colebrooke, Bucks. Dr. Blair died the 24th of June, 1782.

**BLIND**, an epithet applied to an animal deprived of the use of eyes; or one from whom light, colours, and



and all the glorious objects of the visible creation, are intercepted by some natural or accidental cause.

**BLINDNESS**, a privation of the sense of sight. The ordinary causes of blindness are, some external violence, vicious conformation, growth of a cataract, gutta serena, small-pox, &c; or a decay of the optic nerve; an instance of which we have in the Academy of Sciences, where, upon opening the eye of a person long blind, it was found that the optic nerve was extremely shrunk and decayed, and without any medulla in it. The more extraordinary causes of blindness are malignant stenches, poisonous juices dropped into the eye, baneful vermin, long confinement in the dark, or the like.

We find various recompenses for blindness, or substitutes for the use of eyes, in the wonderful sagacity of many blind persons, related by Zahnius in his *Oculus Artificialis*, and others. In some, the defect has been supplied by a most excellent gift of remembering what they had seen before; others by a delicate nose or the sense of smelling; others by a very nice ear; and others again by an exquisite touch, or sense of feeling, which they have had in such perfection, that as it has been said of some, they learned to hear with their eyes, it may be said of these that they taught themselves to see with their hands.

Some have been able to perform all sorts of curious works in the nicest and most dexterous manner. Aldrovandus speaks of a sculptor, who had become blind at 20 years of age, and yet 10 years afterwards he made a perfect marble statue of Cosmo II de Medicis, and another of clay like pope Urban VIII. Bartholin speaks of a blind sculptor in Denmark, who, by mere touch, distinguished perfectly well all sorts of wood, and even colours; and father Grimaldi relates an instance of the same kind; besides the blind organist, lately living in Paris, who it was said did the same thing. What seems more extraordinary still, we are told, by authors of good report, of a blind guide, who used to conduct the merchants through the sands and deserts of Arabia: and a not less marvellous instance is now existing in this country, in one John Metcalf near Manchester, who became quite blind at a very early age; and yet passed many years of his life as a waggoner, and occasionally, as a guide in different roads during the night, or when the paths were covered with snow; and, what is stranger still, his present occupation is that of surveyor and projector of highways in difficult and mountainous parts, particularly about Buxton, and the Peak in Derbyshire.

There are also many instances of blind men who have been highly distinguished for their mental and literary talents, not to speak of the poets Homer, Milton, Ossian, &c; of which we have a remarkable instance in the late Dr. Sanderfon, professor of mathematics in the university of Cambridge, and in the present Dr. Henry Moyes, public lecturer in philosophy, who both of them lost their sight by the small pox at an age before they had any recollection; these men were well skilled in all branches of the mathematics, philosophy, and optics, &c, which they taught with the greatest reputation; besides the monument of fame which the former has left behind him in his mathematical and philosophical works.

The effects of a sudden recovery of sight in such as have been born blind, are also very remarkable: devoid of the experience of distance and figure arising from sight, they are liable to the greatest mistakes in this respect, in so much it has been said that they could not distinguish by the mere sight which was a cube and which a globe, without first touching them. Mr. Boyle mentions a gentleman of this sort, who having been restored to sight at eighteen years of age, was near going distracted with the joy: see Boyle's works abridg. vol. 1. pa. 4. See also a remarkable case of this kind in the Tatler, N<sup>o</sup> 55, vol. 1. And the gentleman couched by Mr. Cheselden had no ideas of colour, shape, or distance: though he knew the colours afunder in a good light during his blind state; yet when he saw them after he had been couched, the faint ideas he had of them before, were not sufficient for him to know them by afterwards: as to distance, his ideas were so deficient, that he thought all the objects he saw touched his eyes, as what he felt did his skin; and it was a considerable time before he could remember which was the dog and which the cat, though often informed, without feeling them.

**BLINDS**, or **BLINDES**, in Fortification, a kind of defence usually made of osiers or branches interwoven, and laid across between two rows of stakes, about a man's height, and 4 or 5 feet asunder. They are used particularly at the heads of trenches, when these are extended in front towards the glacis; serving to defend the workmen, and prevent the enemy from overlooking them.

**BLOCKADE**, is the blocking up a place, by posting troops all about it at the avenues, to prevent supplies of men and provisions from getting into it; and thus starving it out, without forming any regular siege or attacks.

**BLONDEL** (FRANCIS), a celebrated French mathematician and military engineer. He was born at Ribemond in Picardy in 1617. While he was yet but young, he was chosen Regius Professor of Mathematics and Architecture at Paris. Not long after he was appointed governor to Lewis-Henry de Lomenix, Count de Brienne, whom he accompanied in his travels from 1652 to 1655, of which he published an account. He enjoyed many honourable employments, both in the navy and army; and was entrusted with the management of several negotiations with foreign princes. He arrived at the dignity of marshal de camp, and counsellor of state, and had the honour to be appointed mathematical preceptor to the Dauphin. He was a member of the Royal Academy of Sciences, director of the Academy of Architecture, and lecturer to the Royal College: in all which he supported his character with dignity and applause. Blondel was no less versed in the knowledge of the Belles Lettres than in the mathematical sciences, as appears by the comparison he published between Pindar and Horace. He died at Paris the 22d of February 1686, in the 69th year of his age. His chief mathematical works were,

1. Cours d'Architecture. Paris, 1675, in folio.
2. Resolution des quatre principaux problemes d'Architecture. Paris, 1676, in folio.
3. Histoire du Calendrier Romain. Paris, 1682, in 4to.

4. Cours



4. Cours de Mathematiques. Paris, 1683, in 4to.
5. L'Art de jetter des Bombes. La Haye, 1685, in 4to.

Besides a New Method of fortifying Places, and other works.

Blondel had also many ingenious pieces inserted in the Memoirs of the French Academy of Sciences, particularly in the year 1666.

BLOW, in a general sense, denotes a stroke given either with the hand, a weapon, or an instrument. The effect of a blow is estimated like the force of percussion, and so is expressed by the velocity of the body multiplied by its weight.

BLOWING of a Fire-arm, is when the vent or touch-hole is run or gullied, and becomes wide, so that the powder will flame out.

BLUE, one of the seven primitive colours of the rays of light, into which they are divided when refracted through a glass prism. See Newton's Optics &c. See also CHROMATICS.

BLUENESS, that quality of a body, as to colour, from whence it is called blue; depending on such a size and texture of the parts that compose the surface of a body, as disposes them to reflect only the blue or azure rays of light to the eye.

The blueness of the sky is thus accounted for by De la Hire, after Da Vinci; viz, that a black body viewed through a thin white one, gives the sensation of blue, like the immense expanse viewed through the air illuminated and whitened by the sun. For the same reason he says it is, that foot mixed with white, makes a blue; for that white bodies, being always a little transparent, when mixing with a black behind, give the perception of blue. From the same principle too he accounts for the blueness of the veins on the surface of the skin, though the blood they are filled with be a deep red.

In the same manner was the blueness of the sky accounted for by many other of the earlier writers, as Fromondus, Funceius, Otto Guericke, and many others, together with several of the more modern writers, as Wolfius, Muschenbroek, &c. But in the explication of this phenomenon, Newton observes that all the vapours, when they begin to condense and coalesce into natural particles, become first of such a magnitude as to reflect the azure rays, before they can constitute clouds of any other colour. This being therefore the first colour they begin to reflect, must be that of the finest and most transparent skies, in which the vapours are not yet arrived at a grossness sufficient to reflect other colours.

Bouguer however ascribes this blueness of the sky to the constitution of the air itself, being of such a nature that these fainter-coloured rays are incapable of making their way through any considerable tract of it. And as to the blue shadows which were first observed by Buffon in the year 1742, he accounts for them by the aerial colour of the atmosphere, which enlightens these shadows, and in which the blue rays prevail; whilst the red rays are not reflected so soon, but pass on to the remoter regions of the atmosphere. And the Abbé Mazeas accounts for the phenomenon of blue shadows by the diminution of light; observing that,

of two shadows which were cast upon a white wall from an opaque body, illuminated by the moon and by a candle at the same time, that from the candle was reddish, while the other from the moon was blue. See Newton's Optics pa. 228, Bouguer Traité d'Optique pa. 368, Edinb. Ess. vol. 2 pa. 75, or Priestley's Hist. of Vision &c, pa. 436.

BOB of a Pendulum, the same as the ball, which see.

BODY, in Geometry, is a figure conceived to be extended in all directions, or what is usually said to consist of length, breadth, and thickness; being otherwise called a *Solid*. A body is conceived to be formed or generated by the motion of a surface; like as a surface by the motion of a line, and a line by the motion of a point.—Similar bodies, or solids, are in proportion to each other, as the cubes of their like sides, or linear dimensions.

BODY, in Physics, or Natural Philosophy, a solid, extended, palpable substance; of itself merely passive, being indifferent either to motion or rest, and capable of any sort of motion or figure.

Body is composed, according to the Peripatetics, of matter, form, and privation; according to the Epicureans and Corpuscularians, of an assemblage of hooked, heavy atoms; according to the Cartesians, of a certain extension; and according to the Newtonians, of a system or association of solid, massy, hard, impenetrable, moveable particles, ranged or disposed in this or that manner; from which arise bodies of this or that form, and distinguished by this or that name. These elementary or component particles of bodies, they assert, must be perfectly hard, so as never to wear or break in pieces; which, Newton observes, is necessary, in order to the world's persisting in the same state, and bodies continuing of the same nature and texture in several ages.

BODY of a piece of Ordnance, the part contained between the centre of the trunnions and the cascabel. This should always be more fortified or stronger than the rest. See CANNON.

BODY of the Place, in Fortification, denotes either the buildings inclosed, or more generally the inclosure itself. Thus, to construct the body of the place, is to fortify or inclose the place with bastions and curtains.

BODY of a Pump, the thickest part of the barrel or pipe of a pump, within which the piston moves.


BODIES, Regular or Platonic, are those which have all their sides, angles, and planes, similar and equal. Of these there are only 5; viz, the *tetraedron*, contained by 4 equilateral triangles; the *hexaedron* or *cube*, by 6 squares; the *octaedron*, by 8 triangles; the *dodecaedron*, by 12 pentagons; and the *icosaedron*, by 20 triangles.

To form the five Regular Bodies.


Let the annexed figures be exactly drawn on paste-board, or stiff paper, and cut out from it by the extreme or bounding lines: then cut the others, or internal lines, only half through, so that the parts may be turned up by them, and then glued or otherwise fastened together with paste, sealing-wax, &c; so shall they form the respective body marked with the corresponding number;






number; viz, N<sup>o</sup> 1 the tetraedron, N<sup>o</sup> 2 the hexaedron or cube, N<sup>o</sup> 3 the octaedron, N<sup>o</sup> 4 the dodecaedron, and N<sup>o</sup> 5 the icosaedron.




1






2




3



4



5



To find the Superficies or Solidity of the Regular Bodies.

1. Multiply the proper tabular area (taken from the following table) by the square of the linear edge of the solid, for the superficies.

2. Multiply the tabular solidity by the cube of the linear edge, for the solid content.

Table of the Surfaces and Solidities of the five Regular Bodies, the linear edge being 1.			
No. of Faces	Names	Surfaces	Solidities
4	Tetraedron	1.73205	0.11785
6	Hexaedron	6.00000	1.00000
8	Octaedron	3.46410	0.47140
12	Dodecaedron	20.64573	7.66312
20	Icosaedron	8.66025	2.18169

For more particular properties, see each respective word. See also my large *Mensuration*, pa. 248, edit. 2.

These bodies were called *plato*, because they were said to have been invented, or first treated of, by *Plato*, who conceived certain mysteries annexed to them.

BOFFRAND (GERMAIN), a celebrated French architect and engineer, was born at Nantes in Bretagne in 1667. He was brought up under Harduin Mansard, who trusted him with conducting his greatest works. Boffrand was admitted into the French Academy of Architecture in 1709. Many German princes chose him for their architect, and raised considerable edifices on his plans. His manner of building approached that of Palladio; and there was much of grandeur in all his designs. As engineer and inspector-general of bridges and highways, he directed and constructed a number of canals, sluices, bridges, and other mechanical works. He published a curious and useful book, containing the general principles of his art; with an account of the plans, profiles, and elevations of the principal works, which he executed in France and other countries. Boffrand died at Paris in 1755, dean of the Academy of Architecture, first engineer and inspector-general of the bridges and highways, architect and administrator of the general hospital.

BOILING, or EBULLITION, the bubbling up of any fluid, by the application of heat. This is, in general, occasioned by the discharge of an elastic vapour through the fluid that boils; whether that be common air, fixed air, or steam, &c. It is proved by Dr. Hamilton of Dublin, in his Essay on the ascent of vapour, that the boiling of water is occasioned by the lowermost particles of it being heated and rarefied into vapour, or steam; in consequence of this diminution of their specific gravity, they ascend through the surrounding heavier fluid with great velocity, lacerating and throwing up the body of water in the ascent, and so giving it the tumultuous motion called *boiling*.

That this is occasioned by elastic steam, and not by particles of fire or air, as some have imagined, is easily proved by the following simple experiment: Take a common drinking glass filled with hot water, and invert it into a vessel of the same: then, as soon as the water in the vessel begins to boil, large bubbles will be seen to ascend in the glass, by which the water in it will be displaced, and there will soon be a continued bubbling from under its edge: but if the glass be then drawn up, so that its mouth may just touch the water, and a cloth wetted in cold water be applied to the outside, the elastic steam within it will be instantly condensed, upon which the water will ascend so as nearly to fill it again. Some small parts of air &c, that may happen to be lodged in the fluid, may also perhaps be expelled, as well as the rarefied steam. And this is particularly recommended as a method of purifying quicksilver, for making more accurately barometers and thermometers.

We commonly annex the idea of a certain very great degree of heat to the boiling of liquids, though often without reason; for different liquids boil with different degrees of heat; and any one given liquid also, under different pressures of the atmosphere. Thus, a vessel of tar being set over the fire till it boils, it is said a person may then put his hand into it without injury:



injury : and by putting water under the receiver of an air-pump, and applying the flame of a candle or lamp under it, by gradually exhausting, the water is made to boil with always less and less degrees of heat ; and without applying any heat at all, the water, or even the moisture about the bottom or edges of the receiver, will rise in an elastic vapour up into it, when the exhaustion is near completed.

Spirit of wine boils still sooner in vacuo than water. And Dr. Freind gives a table of the different times required to make several fluids boil by the same heat. See also Philos. Transf. N<sup>o</sup> 122.

BOMB, in Artillery, a shell, or hollow ball of cast-iron, having a large vent, by which it is filled with gun-powder, and which is fitted with a fuze or hollow plug to give fire by, when thrown out of a mortar, &c : about the time when the shell arrives at the intended place, the composition in the pipe of the fuze sets fire to the powder in the shell, which blows it all in pieces, to the great annoyance of the enemy, by killing the people, or firing the houses, &c. They are now commonly called *shells* simply, in the English artillery.

These shells, or bombs, are of various sizes, from that of 17 or 18 inches diameter downwards. The very large ones are not used by the English, that of 13 inches diameter being the highest size now employed by them ; the weight, dimensions, and other circumstances of them, and the others downwards, are as in the following table.

Diameter of the Shell.	Weight of the Shell.	Powder to fill them.		Powder to burst them into most pieces.	
		lb.	oz.	lb.	oz.
13 inch	195	9	4½	7	8
10	89	4	14½	3	4
8	46	2	3½	2	0
5½ Royal	14½	1	1½	0	14
4½ Cohorn	7½	0	8	0	7

Mr. Muller gives the following proportion for all shells. Dividing the diameter of the mortar into 30 equal parts, then the other dimensions, in 30ths of that diameter, will be thus :

Diameter of the bore, or mortar	-	-	30
Diameter of the shell,	-	-	29½
Diameter of the hollow sphere	-	-	21
Thickness of metal at the fuze hole	-	-	3½
Thickness at the opposite part	-	-	5
Diameter of the fuze hole	-	-	4
Weight of shell empty	-	-	$\frac{1}{11}d$
Weight of powder to fill it	-	-	$\frac{2}{11}d$

where  $d$  denotes the cube of the diameter of the bore in inches.—But shells have also lately been made with the metal all of the same thickness quite around.

In general, the windage, or difference between the diameter of the shell and mortar, is  $\frac{1}{10}$  of the latter ; also the diameter of the hollow part of the shell is  $\frac{7}{10}$  of the same.

Bombs are thrown out of mortars, or howitzers ; but they may also be thrown out of cannon ; and a very small fort are thrown by the hand, which are called granados : and the Venetians at the siege of Candia,

when the Turks had possessed themselves of the ditch used large bombs without any piece of ordnance, but barely rolled them down upon the enemy along a plank set aslope, with ledges on the sides to keep the bomb right forwards.

Mr. Blondel, in his *Art de jetter des Bombes*, says the first bombs were those thrown into the city of Watchtendonch in Guelderland, in 1588 ; and they are described by our countryman Lucar, in his book on Artillery published this same year 1588 ; though it is pretended by others that they were in use near a century before, namely at the siege of Naples in 1495. They only came into common use, however, in 1634, and then only in the Dutch and Spanish armies. It is said that one Malthus, an English engineer, was sent for from Holland by Lewis the 14th, who used them for him with much success, particularly at the siege of Cohoure in 1642.

The art of throwing bombs, or shells, forms a principal branch of Gunnery, founded on the theory of projectiles, and the quantities and laws of force of gunpowder. And the principal writers on this art are Mess. Blondel, Guisnée, De Resson, De La Hire, &c.

BOMB-CHEST, is a kind of chest usually filled with bombs, and sometimes only with gunpowder, placed under ground, to blow it up into the air with those who stand upon it ; being set on fire by means of a faucissee fastened at one end. But they are now much out of use.

BOMBARD, an ancient piece of ordnance, now out of use. It was very short and thick, with a large mouth ; some of which it is said threw balls of 300 pounds weight, requiring the use of cranes to load them. The Bombard is by some called *bafilisk*, and by the Dutch *donderbus*.

To BOMBARD, is to attack by throwing of bombs, or shells.

BOMBARDIER, a person employed about throwing bombs or shells. He adjusts the fuze, and loads and fires the mortar.

BONES, *Napier's*. See NAPIER.

BONING, in Surveying and Levelling, &c, is the placing three or more rods or poles, all of the same length, in or upon the ground, in such a manner that the tops of them be all in one continued straight line, whether it be horizontal or inclined, so that the eye can look along the tops of them all, from one end of the line to the other.

BONNET, in Fortification, a small work of two faces, having only a parapet, with two rows of palisadoes at about 10 or 12 feet distance. It is commonly placed before the salient angle of the counterscarp, and having a communication with the covered way, by means of a trench cut through the glacis, and palisadoes on each side.

BONNET à Prêtre, or *Priest's Cap*, is an outwork, having three salient angles at the head, besides two inwards. It differs from the double tenaille only in this, that its sides, instead of being parallel, grow narrower, or closer, at the gorge, and opening at the front ; from whence it is called *queue d'aronde*, or swallow's tail.

BOOTES, a constellation of the northern hemisphere, and one of the 48 old ones ; having 23 stars in Ptolemy's



Ptolemy's catalogue, 28 in Tycho's, 34 in Bayer's, 52 in Hevelius's, and 54 in Flamsteed's; of which one, in the skirt of his coat, is of the first magnitude, and called *Arcturus*.

Bootes is represented as a man in the posture of walking; his right hand grasping a club, and his left extended upwards, and holding the cord of the two dogs which seem barking at the Great Bear.

The Greeks, contrary to their usual custom, do not give any certain account of the origin of this constellation. Those who in very early days made the stars which were afterwards formed into the great bear represent a waggon drawn by oxen, made this Bootes the driver of them, from whence he was called the waggoner: others continued the office when the waggon was destroyed, and made a celestial bearward of Bootes, making it his office to drive the two bears round about the pole: and some, when the greater waggon was turned into the greater bear, were still for preserving the form of that machine in those stars which constitute Bootes.

This constellation is called by various other names; as *Arcas*, *Arctophylax*, *Arcturus-Minor*, *Bubulcus*, *Bubulus*, *Canis-Latrans*, *Clamator*, *Icarus*, *Lycaon*, *Philometus*, *Plaustrum-Custos*, *Plorans*, *Thegnis*, and *Vociferator*; by Hesychius it is called *Orion*, and by the Arabs *Aramech*, or *Archamech*. Schiller, instead of Bootes, makes the figure of St. Sylvester; Schickhard, that of Nimrod; and Weigelius, the three Swedish crowns. See Wolf. *Lex Math.* p. 266.

BORE, of a gun, or other piece of ordnance, is the chase, cylinder, or hollow part of the piece.

BOREAL SIGNS, are the first six signs of the Zodiac, or those on the northern side of the equinoctial; viz, the signs ♈ aries, ♉ taurus, ♊ gemini, ♋ cancer, ♌ leo, ♍ virgo.

BOREALIS, AURORA. See *AURORA Borealis*.

BORELLI (JOHN ALPHONSO), a celebrated philosopher and mathematician, born at Naples the 28th of January 1608. He was professor of philosophy and mathematics in some of the most celebrated universities of Italy, particularly at Florence and Pisa, where he became highly in favour with the princes of the house of Medici. But having been concerned in the revolt of Messina, he was obliged to retire to Rome, where he spent the remainder of his life under the protection of Christina queen of Sweden, who honoured him with her friendship, and by her liberality towards him softened the rigour of his hard fortune. He continued two years in the convent of the regular clergy of St. Pantaleon, called the *Pious Schools*, where he instructed the youth in mathematical studies. And this study he prosecuted with great diligence for many years afterward, as appears by his correspondence with several ingenious mathematicians of his time, and the frequent mention that has been made of him by others, who have endeavoured to do justice to his memory. He wrote a letter to Mr. John Collins, in which he discovers his great desire and endeavours to promote the improvement of those sciences: he also speaks of his correspondence with, and great affection for, Mr. Henry Oldenburgh, Secretary of the Royal Society; of Dr. Wallis; of the then late learned Mr. Boyle, and lamented the loss sustained by his

death to the commonwealth of learning. Mr. Baxter, in his *Enquiry into the Nature of the Human Soul*, makes frequent use of our author's book *De Motu Animalium*, and tells us, that he was the first who discovered that the force exerted within the body prodigiously exceeds the weight to be moved without, or that nature employs an immense power to move a small weight. But he acknowledges that Dr. James Keil had shewn that Borelli was mistaken in his calculation of the force of the muscle of the heart; but that he nevertheless ranks him with the most authentic writers, and says he is seldom mistaken: and, having remarked that it is so far from being true, that great things are brought about by small powers, that, on the contrary, a stupendous power is manifest in the most ordinary operations of nature, he observes that the ingenious Borelli first observed this in animal motion; and that Dr. Stephen Hales, by a course of experiments in his *Vegetable Statics*, had shewn the same in the force of the ascending sap in vegetables.

After a course of unceasing labours, Borelli died at Pantaleon of a pleurisy, the 31st of December 1679, at 72 years of age.

Beside several books on physical subjects, Borelli published the following mathematical ones: viz.

1. *Apollonii Pergæi Conicorum Lib. 5, 6, & 7.* Floren. 1661, fol.
2. *Theoriæ Medicorum Planetarum ex causis physicis deductæ.* Flor. 1666, 4to.
3. *De Vi Percussionis.* Bologna, 1667, 4to.—This piece was reprinted, with his celebrated treatise *De Motu Animalium*, and that other *De Motionibus Naturalibus*, in 1686.
4. *Euclides Restitutus, &c.* Pisa, 1668, 4to.
5. *Observatione intorno alla vista ineguale degli Occhi.*—This piece was inserted in the *Journal of Rome*, for the year 1669.
6. *De Motionibus Naturalibus de Gravitate pendentibus.* Regio Julio, 1670, 4to.
7. *Meteorologia Aetnea, &c.* Regio Julio, 1670, 4to.
8. *Observatione dell' Ecclissi Lunare, 11 Gennaro 1675.*—Inserted in the *Journal of Rome* 1675, p. 34.
9. *Elementa Conica Appollonii Pergæi, & Archimedis Opera, nova & breviori methodo demonstrata.*—Printed at Rome in 1679, in 12mo, at the end of the 3d edition of his *Euclides Restitutus*.
10. *De Motu Animalium.* Pars prima in 1680, and Pars altera in 1681, 4to.—These were reprinted at Leyden 1685: revised and purged from many errors; with the addition of John Bernoulli's *Mathematical Meditations concerning the Motion of the Muscles*.
11. At Leyden, 1686, in 4to, a more correct and accurate edition, revised by J. Broen, M. D. of Leyden, of his two pieces, *De Vi Percussionis*, & *De Motionibus de Gravitate pendentibus*.

BOUGUER (PETER), a celebrated French mathematician, was born at Croisic, in Lower Bretagne, the 10th of February 1698. He was the son of John Bouguer, Professor Royal of Hydrography, a tolerably good mathematician, and author of *A complete Treatise on Navigation*. Young Bouguer was accustomed to learn mathematics from his father, from the

time



time he was able to speak, and thus became a proficient in those sciences while he was yet a child. He was sent very early to the Jesuits' college at Vannes, where he had the honour to instruct his regent in the mathematics, at eleven years of age.

Two years after this he had a public contest with a professor of mathematics, upon a proposition which the latter had advanced erroneously; and he triumphed over him; upon which the professor, unable to bear the disgrace, left the country.

Two years after this, when young Bouguer had not yet finished his studies, he lost his father; whom he was appointed to succeed in his office of hydrographer, after a public examination of his qualifications; being then only 15 years of age; an occupation which he discharged with great respect and dignity at that early age.

In 1727, at the age of 29, he obtained the prize proposed by the Academy of Sciences, for the best way of masting of ships. This first success of Bouguer was soon after followed by two others of the same kind; he successively gained the prizes of 1729 and 1731; the former, for the best manner of observing at sea the height of the stars, and the latter, for the most advantageous way of observing the declination of the magnetic needle, or the variation of the compass.

In 1729, he gave an *Optical Essay upon the Gradation of Light*; a subject quite new, in which he examined the intensity of light, and determined its degrees of diminution in passing through different pellucid mediums, and particularly that of the sun in traversing the earth's atmosphere. Mairan gave an extract of this first essay in the *Journal des Savans*, in 1730.

In this same year, 1730, he was removed from the port of Croisic to that of Havre, which brought him into a nearer connection with the Academy of Sciences, in which he obtained, in 1731, the place of associate geometrician, vacant by the promotion of Maupertuis to that of pensioner; and in 1735 he was promoted to the office of pensioner-astronomer. The same year he was sent on the commission to South America, along with Messieurs Godin, Condamine, and Jeussieu, to determine the measure of the degrees of the meridian, and the figure of the earth. In this painful and troublesome business, of 10 years duration, chiefly among the lofty Cordelier mountains, our author determined many other new circumstances, beside the main object of the voyage; such as the expansion and contraction of metals and other substances, by the sudden and alternate changes of heat and cold among those mountains; observations on the refraction of the atmosphere from the tops of the same, with the singular phenomenon of the sudden increase of the refraction, when the star can be observed below the line of the level; the laws of the density of the air at different heights, from observations made at different points of these enormous mountains; a determination that the mountains have an effect upon a plummet, though he did not assign the exact quantity of it; a method of estimating the errors committed by navigators in determining their route; a new construction of the log for measuring a ship's way; with several other useful improvements.

Other inventions of Bouguer, made upon different occasions, were as follow: The heliometer, being a telescope with two object glasses, affording a good method of measuring the diameters of the larger planets with ease and exactness: his researches on the figure in which two lines or two long ranges of parallel trees appear: his experiments on the famous reciprocation of the pendulum: and those upon the manner of measuring the force of the light: &c, &c.

The close application which Bouguer gave to study, undermined his health, and terminated his life the 15th of August 1758, at 60 years of age.—His chief works, that have been published, are,

1. The Figure of the Earth, determined by the observations made in South America; 1749, in 4to.

2. Treatise on Navigation and Pilotage; Paris, 1752, in 4to. This work has been abridged by M. La Caille, in 1 volume, 8vo, 1768.

3. Treatise on Ships, their Construction and Motions; in 4to, 1756.

4. Optical Treatise on the Gradation of Light; first in 1729; then a new edition in 1760, in 4to.

His papers that were inserted in the *Memoirs of the Academy*, are very numerous and important: as, in the *Memoirs for 1726*, Comparison of the force of the solar and lunar light with that of candles.—1731, Observations on the curvilinear motion of bodies in mediums.—1732, Upon the new curves called the *lines of pursuit*.—1733, To determine the species of conoid, to be constructed upon a given base which is exposed to the shock of a fluid, so that the impulse may be the least possible.—Determination of the orbit of comets.—1734, Comparison of the two laws which the earth and the other planets must observe in the figure which gravity causes them to take.—On the curve lines proper to form the arches in domes.—1735, Observations on the equinoxes.—On the length of the pendulum.—1736, On the length of the pendulum in the torrid zone.—On the manner of determining the figure of the earth by the measure of the degrees of latitude and longitude.—1739, On the astronomical refractions in the torrid zone.—Observations on the lunar eclipse, of the 8th of September 1737, made at Quito.—1744, Short account of the voyage to Peru, by the members of the Royal Academy of Sciences, to measure the degrees of the meridian near the equator, and from thence to determine the figure of the earth.—1745, Experiments made at Quito and divers other places in the torrid zone, on the expansion and contraction of metals by heat and cold.—On the problem of the masting of ships.—1746, Treatise on ships, their structure and motions.—On the impulse of fluids upon the fore parts of pyramidoids having their base a trapezium.—Continuation of the short account given in 1744, of the voyage to Peru for measuring the earth.—1747, On a new construction of the log, and other instruments for measuring the run of a ship.—1748, Of the diameters of the larger planets. The new instrument called a *heliometer*, proper for determining them; with observations of the sun.—Observation of the eclipse of the moon the 8th of August 1748.—1749, Second memoir on astronomical refractions, observed in the torrid zone, with remarks on the manner of constructing the tables of them.—



Figure of the earth determined by MM. Bouguer and Condamine, with an abridgment of the expedition to Peru.—1750, Observation of the lunar eclipse of the 13th of December 1750.—1751, On the form of bodies most proper to turn about themselves, when they are pushed by one of their extremities, or any other point.—On the moon's parallax, with the estimation of the changes caused in the parallaxes by the figure of the earth.—Observation of the lunar eclipse, the 2d of December 1751.—1752, On the operations made by seamen, called *Corrections*.—1753, Observation of the passage of Mercury over the sun, the 6th of May 1753.—On the dilatations of the air in the atmosphere.—New treatise of navigation, containing the theory and practice of pilotage, or working of ships.—1754, Operations, &c, for distinguishing, among the different determinations of the degree of the meridian near Paris, that which ought to be preferred.—On the direction which the string of a plummet takes.—Solution of the chief problems in the working of ships.—1755, On the apparent magnitude of objects.—Second memoir on the chief problems in the working of ships.—1757, Account of the treatise on the working of ships.—On the means of measuring the light.—1758, His Eulogy.

In the volumes of the prizes given by the academy, are the following pieces by Bouguer:

In vol. 1, on the masting of ships.—Vol. 2, On the method of exactly observing at sea the height of the stars; and the variation of the compass. Also on the cause of the inclination of the planets' orbits.

BOULTINE, in Architecture, a convex moulding, of a quarter of a circle, and placed next below the plinth in the Tuscan and Dorick capital.

BOW, an offensive weapon made of wood, horn, steel, or other elastic matter, by which arrows are thrown with great force. This instrument was of very ancient and general use, and is still found among all savage nations who have not the use of fire arms, by which it has been superseded among us. There are two species of the Bow, the *Long*, and the *Cross* Bow.

The *Long Bow* is simply a bow, or a rod, with a string fastened to each end of it, to the middle of which the end of an arrow being applied, and then drawn by the hand, on suddenly quitting the hold, the bow returns by means of its elasticity, and impels the arrow from the string with great violence. The old English archers were famous for the long bow, by means of which they gained many victories in France and elsewhere.

The *Cross Bow*, called also *arbalest* or *arbalet*, is a bow strung and set in a shaft of wood, and furnished with a trigger; serving to throw bullets, darts, and large arrows, &c. The ancients had large machines for throwing many arrows at once, called *arbalets*, or *balistæ*.

The force of a bow may be calculated on this principle, that its spring, *i. e.* the power by which it restores itself to its natural position, is always proportional to the space or distance it is bent or removed from it.

Bow, a mathematical instrument formerly used at sea for taking the sun's altitude. It consisted of a large arch divided into 90 degrees, fixed on a staff,

and furnished with three vanes, viz, a side vane, a sight vane, and a horizon vane.

Bow-Compass, an instrument for drawing arches of very large circles, for which the common compasses are too small. It consists of a beam of wood or brass, with three long screws that govern or bend a lath of wood or steel to any arch.

BOX AND NEEDLE, the small compass of a theodolite, circumferentor, or plain-table.

BOYAU, in Fortification, a ditch covered by a parapet, and serving as a communication between two trenches. It runs parallel to the works of the body of the place; and serves as a line of contravallation, both to hinder the sallies of the besieged, and to secure the miners. When it is a particular cut running from the trenches, to cover some spot of ground, it is drawn so as not to be enfiladed or scoured by the enemy's shot.

BOYLE (ROBERT), one of the greatest philosophers, as well as best men, that any country has ever produced, was the 7th son and the 14th child of Richard earl of Cork, and was born at Lismore in the province of Munster in Ireland, the 25th of January, 1626-7; the very year of the death of the learned Lord Bacon, whose plans of experimental philosophy our author afterwards so ably seconded. While very young, he was instructed in his father's house to read and write, and to speak French and Latin. In 1635, when only 8 years old, he was sent over to England, to be educated at Eton school. Here he soon discovered an extraordinary force of understanding, with a disposition to cultivate and improve it to the utmost.

After remaining at Eton between 3 and 4 years, his father sent our author and his brother Francis, in 1638, on their travels upon the continent. They passed through France to Geneva, where they settled for some time to pursue their studies: here our author resumed his acquaintance with the elements of the mathematics, which he had commenced at Eton when 10 years old, on occasion of an illness which prevented his other usual studies.

In the autumn of 1641, he quitted Geneva, and travelled through Switzerland and Italy to Venice, from whence he returned again to Florence, where he spent the winter, studying the Italian language and history, and the works of the celebrated astronomer Galileo, who died in a village near this city during Mr. Boyle's residence here.

About the end of March 1642, he set out from Florence, visited Rome and other places in Italy; then returned to the south of France. At Marseilles, in May 1642, Mr. Boyle received letters from his father, which informed him that the rebellion had broken out in Ireland, and with how much difficulty he had procured 250l. then remitted to help him and his brother home. This remittance however never reached them, and they were obliged to return to Geneva with their governor Mr. Marcombes, who contrived on his own credit, and by selling some jewels, to raise money enough to send them to England, where they arrived in 1644. On their arrival they found that their father was dead, and had left our author the manor of Stalbridge in England, with some other considerable estates in Ireland.



From this time Mr. Boyle's chief residence, for some years at least, was at his manor of Stalbridge, from whence he made occasional excursions to Oxford, London, &c; applying himself with great industry to various kinds of studies, but especially to philosophy and chemistry; and seizing every opportunity of cultivating the acquaintance of the most learned men of his time. He was one of the members of that small but learned body of men who, when all academical studies were interrupted by the civil wars, secreted themselves about the year 1645; and held private meetings, first in London, afterwards at Oxford, to cultivate subjects of natural knowledge upon that plan of experiment which Lord Bacon had delineated. They styled themselves then *The Philosophic College*; but after the restoration, when they were incorporated, and distinguished openly, they took the name of the *Royal Society*.

In the summer of 1654 he retired to settle at Oxford, the Philosophical Society being removed from London to that place, that he might enjoy the conversation of the other learned members, his friends, who had retired thither, such as Wilkins, Wallis, Ward, Willis, Wren, &c. It was during his residence here that he improved that admirable engine the air-pump; and by numerous experiments was enabled to discover several qualities of the air, so as to lay a foundation for a complete theory. He declared against the philosophy of Aristotle, as having in it more of words than things; promising much, and performing little; and giving the inventions of men for indubitable proofs, instead of building upon observation and experiment. He was so zealous for this true method of learning by experiment, and so careful about it, that though the Cartesian philosophy then made a great noise in the world, yet he could never be persuaded to read the works of Descartes, for fear he should be amused and led away by plausible accounts of things founded on conjecture, and merely hypothetical. But philosophy, and enquiries into nature, though they engaged his attention deeply, did not occupy him entirely; as he still continued to pursue critical and theological studies. He had offers of preferment to enter into holy orders, by the government, after the restoration. But he declined the offer, choosing rather to pursue his studies as a layman, in such a manner as might be most effectual for the support of religion; and began to communicate to the world the fruits of these studies. These fruits were very numerous and important, as well as various: the principal of which, as well as of some other memorable occurrences of his life, were nearly in the following order.

In 1660 came out, 1. New experiments, physico-mechanical, touching the spring of the air and its effects.—2. Seraphic love; or some motives and incentives to the love of God, pathetically discoursed of in a letter to a friend. A work which it has been said was owing to his courtship of a lady, the daughter of Cary earl of Monmouth; though our author was never married.—3. Certain physiological essays and other tracts, in 1661.—4. Sceptical chemist, 1662; reprinted about the year 1679, with the addition of divers experiments and notes on the producibleness of chemical principles.

In the year 1663, the Royal Society being incorporated by king Charles the 2d, Mr. Boyle was named one of the council; and as he might justly be reckoned among the founders of that learned body, so he continued one of the most useful and industrious of its members during the whole course of his life. His next publications were, 5. Considerations touching the usefulness of experimental natural philosophy, 1663.—6. Experiments and considerations upon colours; to which was added a letter, containing Observations on a diamond that shines in the dark, 1663. This treatise is full of curious and useful remarks on the hitherto unexplained doctrine of light and colours; in which he shews great judgment, accuracy, and penetration; and which may be said to have led the way to Newton, who made such great discoveries in that branch of physics.—7. Considerations on the style of the holy scriptures, 1663. This was an extract from a larger work, intitled An essay on scripture; which was afterwards published by Sir Peter Pett, a friend of Mr. Boyle's.

In 1664 he was elected into the company of the royal mines; and was all this year occupied in prosecuting various good designs, which was probably the reason that he did not publish any works in this year. Soon after came out, 8. Occasional reflections upon several subjects, 1665. This piece exposed our author to the censure of the celebrated Dean Swift, who, to ridicule these discourses, wrote *A pious meditation upon a broomslick, in the style of the honourable Mr. Boyle*.—9. New experiments and observations upon cold, 1665.—10. Hydrostatical paradoxes made out by new experiments, for the most part physical and easy, 1666.—11. The origin of forms and qualities, according to the corpuscular philosophy, 1666.—Both in this and the former year, our author communicated to his friend Mr. Oldenburgh, then secretary to the Royal Society, several curious and excellent short pieces of his own, upon a great variety of subjects, and others transmitted to him by his learned friends, which are printed in the *Philos. Transf.*

In the year 1668 Mr. Boyle resolved to settle in London for life; and for that purpose he removed to the house of his sister, the lady Ranelagh, in Pall-Mall. This removal was to the great benefit of the learned in general, and particularly of the Royal Society, to whom he gave great and continual assistance, as abundantly appears by the several pieces communicated to them from time to time, and printed in their *Transactions*. To avoid improper waste of time, he had set hours in the day appointed for receiving such persons as wanted to consult him, either for their own assistance, or to communicate new discoveries to him: And he besides kept up an extensive correspondence with the most learned men in Europe; so that it is wonderful how he could bring out so many new works as he did. His next publications were, 12. A continuation of new experiments touching the weight and spring of the air; to which is added, A discourse of the atmosphere of consistent bodies, 1669.—13. Tracts about the cosmical qualities of things; cosmical suspensions; the temperature of the subterraneous regions; the bottom of the sea; to which is prefixed an introduction to the history of particular qualities, 1669.—14. Considerations on the usefulness of experimental and natural philosophy,



osophy, the 2d part, 1671.—15. A collection of tracts upon several useful and important points of practical philosophy, 1671.—16. An essay upon the origin and virtues of gems, 1672.—17. A collection of tracts upon the relation between flame and air; and several other useful and curious subjects, 1672. Besides furnishing, in this and the former year, a number of short dissertations upon a great variety of topics, addressed to the Royal Society, and inserted in their Transactions.—18. Essays on the strange subtilty, great efficacy, and determinate nature, of effluvia; with a variety of experiments on other subjects, 1673.—19. The excellency of theology compared with philosophy, 1673. This discourse was written in the year 1665, while our author, to avoid the great plague which then raged in London, was forced to go from place to place in the country, having little or no opportunity of consulting his books.—20. A collection of tracts upon the saltiness of the sea, the moisture of the air, the natural and preternatural state of bodies; to which is prefixed a dialogue concerning cold, 1674.—21. A collection of tracts containing suspicions about hidden qualities of the air; with an appendix touching celestial magnets; animadversions upon Mr. Hobbes's problem about a vacuum; a discourse of the cause of attraction and suction, 1674.—22. Some considerations about the reasonableness of reason and religion; by T. E. (the final letters of his names). To which is annexed a discourse about the possibility of the resurrection; by Mr. Boyle, 1675. The same year several papers communicated to the Royal Society, among which were two upon quicksilver growing hot with gold.—23. Experiments and notes about the mechanical origin or production of particular qualities, in several discourses on a great variety of subjects, and among the rest on electricity, 1676.—He then communicated to Mr. Hook a short memorial of some observations made upon an artificial substance that shines without any preceding illustration; published by Hook in his *Lectiones Cutlerianæ*.—24. Historical account of a degradation of gold made by an anti-elixir.—25. Aerial noctiluca; or some new phænomena, and a process of a factitious self-shining substance, 1680. This year the Royal Society, as a proof of the just sense of his great worth, and of the constant and particular services which through a course of many years he had done them, made choice of him for their president; but he being extremely, and, as he says, peculiarly tender in point of oaths, declined that honour.—26. Discourse of things above reason; inquiring, whether a philosopher should admit any such, 1681.—27. New experiments and observations upon the icy noctiluca; to which is added a chemical paradox, grounded upon new experiments, making it probable that chemical principles are transmutable, so that out of one of them others may be produced, 1682.—28. A continuation of new experiments, physico-mechanical, touching the spring and weight of the air, and their effects, 1682.—29. A short letter to Dr. Beale, in relation to the making of fresh water out of salt, 1683.—30. Memoirs for the natural history of human blood, especially the spirit of that liquor, 1684.—31. Experiments and considerations about the porosity of bodies, 1684.—32. Short memoirs for the natural experimental history of mineral waters, &c, 1685.—

33. An essay on the great effects of even languid and unheeded motion, &c, 1685.—34. Of the reconcileableness of specific medicines to the corpuscular philosophy, &c, 1685.—35. Of the high veneration man's intellect owes to God, peculiarly for his wisdom and power, 1685.—36. Free inquiry into the vulgarly received notion of nature, 1686.—37. The martyrdom of Theodora and Didymia, 1687. A work he had drawn up in his youth.—38. A disquisition about the final causes of natural things, and about vitiated light, 1688.

Mr. Boyle's health declining very much, he abridged greatly his time given to conversations and communications with other persons, to have more time to prepare for the press some others of his papers, before his death, which were as follow:—39. *Medicina Hydrostatica*, &c, 1690.—40. The Christian virtuoso, &c, 1690. 41. *Experimenta et Observationes Physicæ*, &c, 1691; which is the last work that he published.

Mr. Boyle died on the last day of December of the same year 1691, in the 65th year of his age, and was buried in St. Martin's church in the Fields, Westminster; his funeral sermon being preached by Dr. Gilbert Burnet bishop of Salisbury; in which he displayed the excellent qualities of our author, with many circumstances of his life, &c. But as the limits of this work will not allow us to follow the bishop in the copious and eloquent account he has given of this great man's abilities, we must content ourselves with adding the following short eulogium by the celebrated physician, philosopher, and chemist, Dr. Boerhaave; who, after having declared lord Bacon to be the father of experimental philosophy, asserts, that "Mr. Boyle, the ornament of his age and country, succeeded to the genius and inquiries of the great chancellor Verulam. Which, says he, of Mr. Boyle's writings shall I recommend? All of them. To him we owe the secrets of fire, air, water, animals, vegetables, fossils: so that from his works may be deduced the whole system of natural knowledge."

Mr. Boyle left also several papers behind him, which have been published since his death. Beautiful editions of all his works have been printed at London, in 5 volumes folio, and 6 volumes 4to. Dr. Shaw also published in 3 volumes 4to, the same works "abridged, methodized, and disposed under the general heads of Physics, Statics, Pneumatics, Natural History, Chymistry, and Medicine;" to which he has prefixed a short catalogue of the philosophical writings, according to the order of time when they were first published, &c, as follows:

Physico-mechanical experiments on the spring	
and weight of the air	1661
The Sceptical Chymist	1661
Physiological Essays	1662
History of Colours	1663
Usefulness of Experimental Philosophy	1663
History of Cold	1665
Historical Paradoxes	1666
Origin of Forms and Qualities	1666
Cosmical Qualities	1670
The admirable Rarefaction of the air	1670
The Origin and Virtues of Gems	1672
The Relation betwixt Flame and Air	1672
Efflu-	



Effluvia	-	-	1673
Saltiness of the Sea	-	-	1674
Hidden Qualities in the Air	-	-	1674
The Excellence &c of the Mechanical Hypothesis	-	-	1674
Considerations on the Resurrection	-	-	1675
Particular Qualities	-	-	1676
Aerial Noctiluca	-	-	1680
Icy Noctiluca	-	-	1680
Things above Reason	-	-	1681
Natural History of Human Blood	-	-	1684
Porosity of Bodies	-	-	1684
Natural History of Mineral Waters	-	-	1684
Specific Medicines	-	-	1685
The High Veneration due to God	-	-	1685
Languid Motion	-	-	1685
The Notion of Nature	-	-	1685
Final Causes	-	-	1688
Medicina Hydrostatica	-	-	1690
The Christian Virtuoso	-	-	1690
Experimenta & Observationes Physicæ	-	-	1691
Natural History of the Air	-	-	1692
Medicinal Experiments	-	-	1718

BRADLEY (DR. JAMES), a celebrated English astronomer, the third son of William Bradley, was born at Sherborne in Gloucestershire in the year 1692. He was fitted for the university at Northleach in the same county, at the boarding school of Mr. Egles and Mr. Brice. From thence he was sent to Oxford, and admitted a commoner of Baliol college March 15, 1710; where he took the degree of bachelor the 14th of October 1714, and of master of arts the 21st of January 1716. His friends intending him for the church, his studies were regulated with that view; and as soon as he was of a proper age to receive holy orders, the bishop of Hereford, who had conceived a great esteem for him, gave him the living of Bridstow, and soon after he was inducted to that of Landewy Welfry in Pembrokeshire.

He was nephew to Mr. Pound, a gentleman well known in the learned world, by many excellent astronomical and other observations, and who would have enriched it much more, if the journals of his voyages had not been burnt at Pulo Condor, when the place was set on fire, and the English who were settled there cruelly massacred, Mr. Pound himself very narrowly escaping with his life. With this gentleman, at Wanstead, Mr. Bradley passed all the time that he could spare from the duties of his function; being then sufficiently acquainted with the mathematics to improve by Mr. Pound's conversation. It may easily be imagined that the example and conversation of this gentleman did not render Bradley more fond of his profession, to which he had before no great attachment: he continued however as yet to fulfil the duties of it, though at this time he had made such observations as laid the foundation of those discoveries which afterward distinguished him as one of the greatest astronomers of his age. These observations gained him the notice and friendship of the lord chancellor Macclesfield, Mr. Newton afterward Sir Isaac, Mr. Halley, and many other members of the Royal Society, into which he was soon after elected a member.

Soon after, the chair of Savilian professor of astro-

nomy at Oxford became vacant, by the death of the celebrated Dr. John Keil; and Mr. Bradley was elected to succeed him on the 31st of October 1721, at 29 years of age: his colleague being Mr. Halley, who was professor of geometry on the same foundation. Upon this appointment, Mr. Bradley resigned his church livings, and applied himself wholly to the study of his favourite science. In the course of his observations, which were innumerable, he discovered and settled the laws of the alterations of the fixed stars, from the progressive motion of light, combined with the earth's annual motion about the sun, and the nutation of the earth's axis, arising from the unequal attraction of the sun and moon on the different parts of the earth. The former of these effects is called the *aberration* of the fixed stars, the theory of which he published in 1727; and the latter the *nutation* of the earth's axis, the theory of which appeared in 1737: so that in the space of about 10 years, he communicated to the world two of the finest discoveries in modern astronomy; which will for ever make a memorable epoch in the history of that science. See ABERRATION and NUTATION.

In 1730 our author succeeded Mr. Whiteside, as lecturer in astronomy and experimental philosophy in the Museum at Oxford: which was a considerable emolument to him, and which he held till within a year or two of his death; when the ill state of his health induced him to resign it.

Our author always preserved the esteem and friendship of Dr. Halley; who, being worn out by age and infirmities, thought he could not do better for the service of astronomy, than procure for Mr. Bradley the place of regius professor of astronomy at Greenwich, which he himself had many years possessed with the greatest reputation. With this view he wrote many letters, desiring Mr. Bradley's permission to apply for a grant of the reversion of it to him, and even offered to resign it in his favour, if it should be thought necessary: but Dr. Halley died before he could accomplish this kind object. Our author however obtained the place, by the interest of lord Macclesfield, who was afterward president of the Royal Society; and upon this appointment the university of Oxford sent him a diploma of doctor of divinity.

The appointment of astronomer royal at Greenwich, which was dated the 3d of February 1741-2, placed our author in his proper element; and he pursued his observations with unwearied diligence. However numerous the collection of astronomical instruments at that observatory, it was impossible that such an observer as Dr. Bradley should not desire to increase them, as well to answer those particular views, as in general to make observations with greater exactness. In the year 1748 therefore, he took the opportunity of the visit of the Royal Society to the observatory, annually made to examine the instruments and receive the professor's observations for the year, to represent so strongly the necessity of repairing the old instruments, and providing new ones, that the society thought proper to make application to the king, who was pleased to order 1000 pounds for that purpose. This sum was laid out under the direction of our author, who, with the assistance of the late celebrated Mr. Graham and Mr. Bird, furnished the observatory with as complete a collection



lection of astronomical instruments, as the most skilful and diligent observer could desire. Dr. Bradley, thus furnished with such assistance, pursued his observations with great assiduity during the rest of his life; an immense number of which was found after his death, in 13 folio volumes, and were presented to the university of Oxford in the year 1776, on condition of their printing and publishing them; but which however, unfortunately for the improvement of astronomy, now after a lapse of almost 20 years, has never yet been done.

During Dr. Bradley's residence at the Royal Observatory, the living of the church at Greenwich became vacant, and was offered to him: upon his refusing to accept it, from a conscientious scruple, "that the duty of a pastor was incompatible with his other studies and necessary engagements," the king was pleased to grant him a pension of 250l. over and above the astronomer's original salary from the Board of Ordnance, "in consideration (as the sign manual, dated the 15th Feb. 1752, expresses it) of his great skill and knowledge in the several branches of astronomy and other parts of the mathematics, which have proved so useful to the trade and navigation of this kingdom." A pension which has been regularly continued to the astronomers royal ever since.

About 1748, our author became entitled to bishop Crew's benefaction of 30l. a year, to the lecture reader in experimental philosophy at Oxford. He was elected a member of the Academy of Sciences at Berlin, in 1747; of that at Paris, in 1748; of that at Petersburg, in 1754; and of that at Bologna, in 1757. He was married in the year 1744, but never had more than one child, a daughter.

By too close application to study and observations, Dr. Bradley became afflicted, for near two years before his death, with a grievous oppression on his spirits; which interrupted his useful labours. This distress arose chiefly from an apprehension that he should outlive his rational faculties: but this so much dreaded evil never came upon him. In June 1762 he was seized with a suppression of urine, occasioned by an inflammation in the reins, which terminated his existence the 13th of July following. His death happened at Chalfont in Gloucestershire, in the 70th year of his age; and he was interred at Minchinhampton in the same county.

As to his character, Dr. Bradley was remarkable for a placid and gentle modesty, very uncommon in persons of an active temper and robust constitution. Although he was a good speaker, and possessed the rare but happy art of expressing his ideas with the utmost precision and clearness, yet no man was a greater lover of silence, for he never spoke but when he thought it absolutely necessary. Nor was he more inclined to write than to speak, as he has published very little: he had a natural diffidence, which made him always afraid that his works might injure his character; so that he suppressed many which might have been worthy of publication.

His papers, which have been inserted in the *Philos. Transf.* are,

1. Observations on the comet of 1703. Vol. 33, p. 41.

2. The longitude of Lisbon and of the fort of New York from Wansled and London determined by the eclipse of the first satellite of Jupiter. Vol. 34, p. 85.

3. An account of a new discovered Motion of the Fixed Stars. Vol. 35, p. 637.

4. On the Going of Clocks with Isochronal Pendulums. Vol. 38, p. 302.

5. Observations on the Comet of 1736-7. Vol. 40, p. 111.

6. On the apparent Motion of the fixed Stars. Vol. 45, p. 1.

7. On the Occultation of Venus by the Moon, the 15th of April 1751. Vol. 46, p. 201.

8. On the Comet of 1757. Vol. 50, p. 408.

9. Directions for using the Common Micrometer. Vol. 62, p. 46.

BRADWARDIN (THOMAS), archbishop of Canterbury, was born at Hartfield in Suffex, about the close of the 13th century. He was educated at Merton College, Oxford, where he took the degree of doctor of divinity; and acquired the reputation of a profound scholar, a skilful mathematician, and consummate divine. It has been said he was professor of divinity at Oxford; that he was chancellor of the diocese of London, and confessor to Edward the 3d, whom he constantly attended during his war with France. After his return from the war, he was made prebendary of Lincoln, and afterward archbishop of Canterbury. He died at Lambeth in the year 1349, forty days after his consecration. His works are, 1. *De Causa Dei*, printed London 1618, published by J. H. Savil.—2. *De Geometria speculativa*, &c. Paris, 1495, 1512, 1530.—3. *De Arithmetica practica*, Paris, 1502, 1512.—4. *De Proportionibus*, Paris, 1495. Venice, 1505, folio.—5. *De Quadratura Circuli*, Paris, 1495, folio.

BRAHE (TYCHO), a celebrated astronomer, descended from a noble family originally of Sweden but settled in Denmark, was born the 14th of December 1546, at Knudstorp in the county of Schonen, near Helsingbourg. He was taught Latin when 7 years old, and studied 5 years under private tutors. His father dying while our author was very young, his uncle, George Brahe, having no children, adopted him, and sent him, in 1559, to study philosophy and rhetoric at Copenhagen. The great eclipse of the sun, on the 21st of August 1560, happening at the precise time the astronomers had foretold, he began to consider astronomy as something divine; and purchasing the tables of Stadius, he gained some notion of the theory of the planets. In 1562 he was sent by his uncle to Leipzig to study the law, where his acquirements gave manifest indications of extraordinary abilities. His natural inclination however was to the study of the heavens, to which he applied himself so assiduously, that, notwithstanding the care of his tutor to keep him close to the study of the law, he made use of every means in his power for improving his knowledge of astronomy; he purchased with his pocket money whatever books he could meet with on the subject, and read them with great attention, procuring assistance in difficult cases from Bartholomew Scultens his private tutor; and having procured a small celestial globe, he took opportunities, when his tutor was in bed, and when the weather was clear, to examine the constellations in the heavens, to learn their



names from the globe, and their motions from observation.

After a course of 3 years study at Leipzig, his uncle dying, he returned home in 1565. In this year a difference arising between Brahe and a Danish nobleman, they fought, and our author had part of his nose cut off by a blow; a defect which he so artfully supplied with one made of gold and silver, that it was not perceivable. About this time he began to apply himself to chemistry, proposing nothing less than to obtain the philosopher's stone. But becoming greatly disgusted to see the liberal arts despised, and finding his own relations and friends uneasy that he applied himself to astronomy, as thinking it a study unsuitable to a person of his quality, he went to Wirtemberg in 1566, from whence the breaking out of the plague soon occasioned his removal to Rostock, and in 1569 to Augsburg, where he was visited by Peter Ramus, then professor of astronomy at Paris, and who greatly admired his uncommon skill in this science.

In 1571 he returned to Denmark; and was favoured by his maternal uncle, Steno Billes, a lover of learning, with a convenient place at his castle of Herritzvad near Knudstorp, for making his observations, and building a laboratory. And here it was he discovered, in 1573, a new star in the constellation Cassiopeia. But soon after, his marrying a country girl, beneath his rank, occasioned so violent a quarrel between him and his relations, that the king was obliged to interpose to reconcile them.

In 1574, by the king's command, he read lectures at Copenhagen on the theory of the planets. The year following he began his travels through Germany, and proceeded as far as Venice. He then resolved to remove his family, and settle at Basil; but Frederic the 2d, king of Denmark, being informed of his design, and unwilling to lose a man who was capable of doing so much honour to his country, he promised to enable him to pursue his studies, and bestowed upon him for life the island of Huen in the Sound, and promised that an observatory and laboratory should be built for him, with a supply of money for carrying on his designs: and accordingly the first stone of the observatory was laid the 8th of August 1576, under the name of Uranibourg: The king also gave him a pension of 2000 crowns out of his treasury, a fee in Norway, and a canonry of Roskilde, which brought him in 1000 more. This situation he enjoyed for the space of about 20 years, pursuing his observations and studies with great industry: here he kept always in his house ten or twelve young men, who assisted him in his observations, and whom he instructed in astronomy and mathematics. Here it was that he received a visit from James the 6th, king of Scotland, afterward James the 1st of England, having come to Denmark to espouse Anne, daughter of Frederick the 2d: James made our author some noble presents, and wrote a copy of Latin verses in his praise.

Brahe's tranquillity however in this happy situation was at length fatally interrupted. Soon after the death of king Frederick, by the aspersions of envious and malevolent ministers, he was deprived of his pension, fee, and canonry, in 1596. Being thus rendered incapable of supporting the expences of his establish-

ment, he quitted his favourite Uranibourg, and withdrew to Copenhagen, with some of his instruments, and continued his astronomical observations and chemical experiments in that city, till the same malevolence procured from the new king, Charles the 4th, an order for him to discontinue them. This induced him to fall upon means of being introduced to the emperor Rodolphus, who was fond of mechanism and chemical experiments: and to smooth the way to an interview, Tycho now published his book, *Astronomia inflammata Mechanica*, adorned with figures, and dedicated it to the emperor. That prince received him at Prague with great civility and respect; gave him a magnificent house, till he could procure one for him more fit for astronomical observations; he also assigned him a pension of 3000 crowns; and promised him a fee for himself and his descendants. Here then he settled in the latter part of 1598, with his sons and scholars, and among them the celebrated Kepler, who had joined him. But he did not long enjoy this happy situation; for, about 3 years after, he died, on the 24th of October 1601, of a retention of urine, in the 55th year of his age, and was interred in a very magnificent manner in the principal church at Prague, where a noble monument was erected to him; leaving, beside his wife, two sons and four daughters. On the approach of death, he enjoined his sons to take care that none of his works should be lost; exhorted the students to attend closely to their exercises; and recommended to Kepler the finishing of the Rudolphine tables he had constructed for regulating the motion of the planets.

Brahe's skill in astronomy is universally known; and he is famed for being the inventor of a new system of the planets, which he endeavoured, though without success, to establish on the ruins of that of Copernicus. He was very credulous with regard to judicial astrology and presages: If he met an old woman when he went out of doors, or a hare upon the road on a journey, he would turn back immediately, being persuaded that it was a bad omen: Also, when he lived at Uranibourg, he kept at his house a madman, whom he placed at his feet at table, and fed himself; for as he imagined that every thing spoken by mad persons presaged something, he carefully observed all that this man said; and because it sometimes proved true, he fancied it might always be depended on. He was of a very irritable disposition: a mere trifle put him in a passion; and against persons of the first rank, whom he thought his enemies, he openly discovered his resentment. He was very apt to rally others, but highly provoked when the same liberty was taken with himself.—The principal part of his writings, according to Cassendus, are,

1. An account of the New Star, which appeared Nov. 11th 1572, in Cassiopeia; Copenh. 1573, in 4to.
- 2. An Oration concerning the Mathematical Sciences, pronounced in the university of Copenhagen, in the year 1574: published by Conrad Aslac, of Bergen in Norway.—3. A treatise on the Comet of the year 1577, immediately after it disappeared. Nine years afterward, he revised it, and added a 10th chapter. Printed at Uranibourg, 1589.—4. Another treatise on the New Phenomena of the heavens. In the first part of which he treats of the Restitution, as he calls it, of the sun, and of the fixed stars. And in the 2d part, of



a New Star, which then had made its appearance.—5. A collection of Astronomical Epistles: printed in 4to, at Uranibourg in 1596; Nuremberg in 1602; and at Franckfort in 1610. It was dedicated to Maurice landgrave of Hesse; because there are in it a considerable number of letters of the landgrave William his father, and of Christopher Rothmann, the mathematician of that prince, to Tycho, and of Tycho to them.—6. The Mechanical Principles of Astronomy restored: Wandersburg, 1598, in folio.—7. An Answer to the Letter of a certain Scotchman, concerning the comet, in the year 1577.—8. On the composition of an Elixir for the Plague; addressed to the emperor Rodolphus.—9. An elegy upon his Exile: Rostock, 1614, 4to.—10. The Rudolphine Tables; which he had not finished when he died; but were revised, and published by Kepler, as Tycho had desired.—11. An accurate Enumeration of the Fixed Stars: addressed to the emperor Rodolphus.—12. A complete Catalogue of 1000 of the Fixed Stars; which Kepler has inserted in the Rudolphine Tables.—13. *Historia Caelestis*; or a History of the Heavens; in two parts: The 1st contains the Observations he had made at Uranibourg, in 16 books: The latter contains the Observations made at Wandersburg, Wittenberg, Prague, &c; in 4 books.—14. Is an Epistle to Caster Pucer; printed at Copenhagen 1668.

BRANCKER, or BRANKER (THOMAS), an eminent mathematician of the 17th century, son of Thomas Brancker some time bachelor of arts in Exeter College, Oxford, was born in Devonshire in 1636, and was admitted butler of the said college Nov. 8, 1652, in the 17th year of his age. In 1655, June the 15th, he took the degree of bachelor of arts, and was elected probationary fellow the 30th of the same month. In 1658, April the 22d, he took the degree of master of arts, and became a preacher; but after the restoration, refusing to conform to the ceremonies of the church of England, he quitted his fellowship in 1662, and retired to Chester: But not long after, he became reconciled to the service of the church, took orders from a bishop, and was made a minister of Whitegate. He had however, for some time, enjoyed great opportunity and leisure for pursuing the bent of his genius in the mathematical sciences; and his skill both in the mathematics and chemistry procured him the favour of lord Brereton, who gave him the rectory of Tilston. He was afterward chosen master of the well-endowed school at Macclesfield, in that county, where he spent the remaining years of his life, which was terminated by a short illness in 1676, at 40 years of age; and he was interred in the church at Macclesfield.

Brancker wrote a piece on the Doctrine of the Sphere, in Latin, which was published at Oxford in 1662; and in 1668, he published at London, in 4to, a translation of Rhonius's Algebra, with the title of *An Introduction to Algebra*; which treatise having communicated to Dr. John Pell, he received from him some assistance towards improving it; which he generously acknowledges in a letter to Mr. John Collins; with whom, and some other gentlemen, proficient in this science, he continued a correspondence during his life.

BREACH, in Fortification, a gap or opening made in any part of the works of a town, by the cannon or

mines of the besiegers, with intent to storm or attack the place.

BREAKING *Ground*, in Military Affairs, is beginning the works for carrying on the siege of a place; more especially the beginning to dig trenches, or approaches.

BREECH *of a Gun*, the hinder part, from the cascabel to the lower part of the bore.

BREREWOOD (EDWARD), a learned mathematician and antiquary, was the son of Robert Brerewood, a reputable tradesman, who was three times mayor of Chester. Our author was born in that city in 1565, where he was educated in grammar learning at the free school; and was afterward admitted, in 1581, of Brazen-nose College, Oxford; where he soon acquired the character of a hard student; as he has shewn by the commentaries he wrote upon Aristotle's Ethics, which were written by him about the age of 21.

In the year 1596 he was chosen the first Professor of Astronomy in Gresham College, being one of the two who, at the desire of the electors, were recommended to them by the university of Oxford. He loved retirement, and wholly devoted himself to the pursuit of knowledge. And though he never published any thing himself, yet he was very communicative, and ready to impart what he knew, to others, either in conversation or in writing. His retired situation at Gresham College being agreeable, it did not appear that he had any other views, but continued there the remainder of his life, which was terminated by a fever the 4th of November 1613, at 48 years of age, in the midst of his pursuits, and before he had taken proper care to collect and digest his learned labours; which however were not lost; being reduced to order, and published after his death. These were little or nothing mathematical, being of a miscellaneous nature, upon the several subjects of Weights, Money, Languages, Religion, Logic, the Sabbath, Meteors, the Eye, Ethics, &c.

BRIDGE, a work of carpentry or masonry, built over a river, canal, or the like, for the convenience of passing from one side to the other; and may be considered as a road over water, supported by one or more arches, and these again supported by proper piers or buttments. Besides these essential parts, may be added the paving at top, the banquet, or raised footway, on each side, leaving a sufficient breadth in the middle for horses and carriages, also the parapet wall either with or without a balustrade, or other ornamental and useful parts. The breadth of a bridge for a great city should be such, as to allow an easy passage for three carriages and two horsemen abreast in the middle way, and for 3 foot passengers in the same manner on each banquet: but for other smaller bridges, a less breadth.

Bridges are commonly very difficult to execute, on account of the inconvenience of laying foundations and walling under water. The earliest rules and instructions for building of bridges are given by Alberti, in his *Archit.* 1, 8. Other rules were afterwards laid down by Palladio, Serlio, and Scamozzi, which are collected by Blondel, in his *Cours d'Archit.* pa. 629 &c. The best of these rules were also given by Gold-man, Baukhurst, and in Hawkesmoor's History of London



London Bridge. M. Gautier has a considerable volume expressly on bridges, ancient and modern. See also Riou's Short Principles for the Architecture of Stone Bridges; as also Emerson, Muller, Labelye, and my own Principles of Bridges.

The conditions required in a bridge are, That it be well designed, commodious, durable, and suitably decorated. It should be of such a height as to be quite convenient for the passage over it, and yet easily admitting through its arches the vessels that navigate upon it, and all the water, even at high tides and floods: the neglect of this precept has been the ruin of many bridges. Bridges are commonly continued in a straight direction perpendicular to the stream; though some think they should be made convex towards the stream, the better to resist floods, &c. And bridges of this sort have been executed in some places, as Pont St. Esprit near Lyons. Again, a bridge should not be made in too narrow a part of a navigable river, or one subject to tides or floods: because the breadth being still more contracted by the piers, this will increase the depth, velocity, and fall of the water under the arches, and endanger the whole bridge and navigation. There ought to be an uneven number of arches, or an even number of piers; both that the middle of the stream or chief current may flow freely without the interruption of a pier; and that the two halves of the bridge, by gradually rising from the ends to the middle, may there meet in the highest and largest arch; and also, that by being open in the middle, the eye in viewing it may look directly through there. When the middle and ends are of different heights, their difference however ought not to be great in proportion to the length, that the ascent and descent may be easy; and in that case also it is more beautiful to make the top in one continued curve, than two straight lines forming an angle in the middle. Bridges should rather be of few and large arches, than of many smaller ones, if the height and situation will possibly allow of it; for this will leave more free passage for the water and navigation, and be a great saving in materials and labour, as there will be fewer piers and centres, and the arches &c will require less materials; a remarkable instance of which appears in the difference between the bridges of Westminster and Blackfriars, the expence of the former being more than double the latter.

For the proper execution of a bridge, and making an estimate of the expence, &c, it is necessary to have three plans, three sections, and an elevation. The three plans are so many horizontal sections, viz, first a plan of the foundation under the piers, with the particular circumstances attending it, whether of gratings, planks, piles, &c; the 2d is the plan of the piers and arches; and the 3d is the plan of the superstructure, with the paved road and banquet. The three sections are vertical ones; the first of them, a longitudinal section from end to end of the bridge, and through the middle of the breadth; the 2d a transverse one, or across it, and through the summit of an arch; and the 3d also across, but taken upon a pier. The elevation is an orthographic projection of one side or face of the bridge, or its appearance as viewed at a distance, shewing the exterior aspect of the materials, with the manner in which they are disposed &c.

For the figure of the arches, some prefer the semi-circle, though perhaps without knowing any good reason why; others the elliptical form; as having many advantages over the semi-circular; and some talk of the catenarian arch, though its pretended advantages are only chimerical; but the arch of equilibration is the only perfect one, so as to be equally strong in every part: see my Principles of Bridges. The piers are of diverse thickness, according to the figure, span, and height of the arches; as may be seen in the work above mentioned.

With the Romans, the repairing and building of bridges were committed to the priests, thence named *pontifices*; next to the censores, or curators of the roads; but at last the emperors took the care of the bridges into their own hands. Thus, the Pons Janiculensis was built of marble by Antoninus Pius; the Pons Cestius was restored by Gordian; and Arian built a new one which was called after his own name. In the middle age, bridge-building was counted among the acts of religion; and, toward the end of the 12th century, St. Benezet founded a regular order of hospitallers, under the name of *pontifices*, or bridge-builders, whose office was to assist travellers, by making bridges, settling ferries, and receiving strangers into hospitals, or houses, built on the banks of rivers. We read of an hospital of this kind at Avignon, where the hospitallers resided under the direction of their first superior St. Benezet: and the Jesuit Raynaldus has a treatise on St. John the bridge-builder.

Among the bridges of antiquity, that built by Trajan over the Danube, it is allowed, is the most magnificent. It was demolished by his next successor Adrian, and the ruins are still to be seen in the middle of the Danube, near the city Warhel in Hungary. It had 20 piers, of square stone, each of which was 150 feet high above the foundation, 60 feet in breadth, and 170 feet distant from one another, which is the span or width of the arches; so that the whole length of the bridge was more than 1530 yards, or one mile nearly.

In France, the Pont de Garde is a very bold structure; the piers being only 13 feet thick, yet serving to support an immense weight of a triplicate arcade, and joining two mountains. It consists of three bridges, one over another; the uppermost of which is an aqueduct.

The bridge of Avignon, which was finished in the year 1188, consists of 18 arches, and measures 1340 paces, or about 1000 yards in length.

The famous bridge at Venice, called the Rialto, passes for a master piece of art, consisting of only one very flat and bold arch, near 100 feet span, and only 23 feet high above the water: it was built in 1591.—Poulet also mentions a bridge of a single arch, in the city of Munster in Bothnia, much bolder than that of the Rialto at Venice. Yet these are nothing to a bridge in China, built from one mountain to another, consisting only of a single arch, 400 cubits long, and 500 cubits high, whence it is called the flying bridge; and a figure of it is given in the Philos. Transf. Kircher also speaks of a bridge in the same country 360 perches long without any arch, but supported by 300 pillars.

There are many bridges of considerable note in our



own country. The triangular bridge at Crowland in Lincolnshire, it is said, is the most ancient Gothic structure remaining intire in the kingdom; and was erected about the year 860.

London bridge is on the old Gothick structure, with 20 small locks or arches, each of only 20 feet wide; but there are now only 18 open, two having lately been thrown into one in the centre, and another next one side is concealed or covered up. It is 900 feet long, 60 high, and 74 wide; the piers are from 25 to 34 feet broad, with starlings projecting at the ends; so that the greatest water-way, when the tide is above the starlings, was 450 feet, scarce half the breadth of the river; and below the starlings, the water-way was reduced to 194 feet, before the late opening of the centre. London bridge was first built with timber between the years 993 and 1016; and it was repaired, or rather new built with timber, 1163. The stone bridge was begun in 1176, and finished in 1209. It is probable there were no houses on this bridge for upwards of 200 years; since we read of a tilt and tournament held on it in 1395. Houses it seems were erected on it afterwards; but being found of great inconvenience and nuisance, they were removed in 1758, and the avenues to it enlarged and the whole made more commodious; the two middle arches were then thrown into one, by removing the pier from between them; the whole repairs amounting to above 80,000l.

There are still some more bridges in England built in the old manner of London bridge; as the bridge at Rochester, and some others; also the late bridge at Newcastle upon Tyne, which was broken down by a great flood in the year 1771, for want of a sufficient quantity of water-way through the arches.

The longest bridge in England is that over the Trent at Burton, built in the 12th century, of squared free-stone, and is strong and lofty; it contains 34 arches, and the whole length is 1545 feet. But this falls far short of the wooden bridge over the Drave, which according to Dr. Brown is at least 5 miles long.

But one of the most singular bridges in Europe is that built over the Taaf in Glamorganshire, by William Edward, a poor country mason, in the year 1756. This remarkable bridge consists of only one stupendous arch, which, though only 8 feet broad, and 35 feet high, is no less than 140 span, being part of a circle of 175 feet diameter.

There is also a remarkable bridge of one arch, built at Colebrook Dale in 1779, of cast iron: and another still larger of the same metal, is now raising over the river Wear, at Sunderland, the arch being of 240 feet span.

Of modern bridges, perhaps the two finest in Europe, are the Westminster and Blackfriars bridges over the river Thames at London. The former is 1220 feet long, and 44 feet wide, having a commodious broad foot path on each side for passengers. It consists of 13 large and two small arches, all semicircular, with 14 intermediate piers. The arches all spring from about 2 feet above low-water mark; the middle arch is 76 feet wide, and the others on each side decrease always by 4 feet at a time. The two middle piers are each 17 feet thick at the springing of the arches; and the others decrease equally on each side by one foot at

a time; every pier terminating with a saliant right angle against either stream. This bridge is built of the best materials, and in a neat and elegant taste, but the arches are too small for the quantity of masonry contained in it. This bridge was begun in 1738, and opened in 1750; and the whole sum of money granted and paid for the erection of this bridge, with the purchase of houses to take down, and widening the avenues, &c, amounted to 389,500l.

Blackfriars bridge, nearly opposite the centre of the city of London, was begun in 1760, and was completed in 10 years and three quarters; and is an exceeding light and elegant structure, but the materials unfortunately do not seem to be of the best, as many of the arch stones are decaying. It consists of 9 large, elegant, elliptical arches; the centre arch being 100 feet wide, and those on each side decreasing in a regular gradation, to the smallest, at each extremity, which is 70 feet wide. The breadth of the bridge is 42 feet, and the length from wharf to wharf 995. The upper surface is a portion of a very large circle, which forms an elegant figure, and is of convenient passage over it. The whole expence was 150,840l.

There are various sorts of bridges, of stone, wood, or metal, of boats or floats, pendant or hanging bridges, draw bridges, flying bridges, &c, &c, and even natural bridges, or such as are found formed by nature, of which kind a most wonderful one is described by Mr. Jefferson, in his *State of Virginia*; and another, but smaller, is described by Don Ulloa, in the province of Angaraez in South America.

BRIGGS (HENRY), one of the greatest mathematicians in the 16th and 17th centuries, was born at Warleywood, near Halifax, in Yorkshire, in 1556. From a grammar school in that country he was sent to St. John's College, Cambridge, 1579; where after taking both the degrees in arts, he was chosen fellow of his college in 1588. He applied himself chiefly to the study of the mathematics, in which he greatly excelled; in consequence in 1592 he was made examiner and lecturer in that faculty; and soon after, reader of the physic lecture, founded by Dr. Linacer.

Upon the settlement of Gresham College, in London, he was chosen the first professor of geometry there, in 1596. Soon after this, he constructed a table, for finding the latitude, from the variation of the magnetic needle being given. In the year 1609 he contracted an acquaintance with the learned Mr. James Usher, afterwards archbishop of Armagh, which continued many years after by letters, two of Mr. Briggs being still extant in the collection of Usher's letters that were published: in the former of these, dated August 1610, he writes among other things, that he was engaged in the subject of eclipses; and in the latter, dated the 10th of March 1615, that he was wholly taken up and employed about the noble invention of logarithms, which had come out the year before, and in the improvement of which he had afterwards so great a concern. For Briggs immediately set himself to the study and improvement of them; expounding them also to his auditors in his lectures at Gresham college. In these lectures he proposed the alteration of the scale of logarithms, from the hyperbolic form which Napier had



had given them, to that in which I should be the logarithm of the ratio of 10 to 1; and soon after he wrote to Napier to make the same proposal to himself. In the year 1616 Briggs made a visit to Napier at Edinburgh, to confer with him upon this change; and the next year he did the same also. In these conferences, the alteration was agreed upon accordingly, and upon Briggs's return from his second visit, in 1617, he published the first chiliad, or 1000 of his logarithms. See the Introduction to my Logarithms.

In 1619 he was made the first Savilian professor of geometry; and resigned the professorship of Gresham college the 25th of July 1620. At Oxford he settled himself at Merton college, where he continued a most laborious and studious life, employed partly in the duties of his office as geometry lecturer, and partly in the computation of the logarithms, and in other useful works. In the year 1622 he published a small tract on the "North-west passage to the South Seas, through the continent of Virginia and Hudson's Bay;" the reason of which was probably, that he was then a member of the company trading to Virginia. His next performance was his great and elaborate work, the *Arithmetica Logarithmica* in folio, printed at London in 1624; a stupendous work for so short a time! containing the logarithms of 30 thousand natural numbers, to 14 places of figures beside the index. Briggs lived also to complete a table of logarithmic sines and tangents for the 100th part of every degree, to 14 places of figures beside the index; with a table of natural sines for the same 100th parts to 15 places, and the tangents and secants for the same to ten places; with the construction of the whole. These tables were printed at Gouda in 1631, under the care of Adrian Vlacq, and published in 1633, with the title of *Trigonometria Britannica*. In the construction of these two works, on the logarithms of numbers, and of sines and tangents, our author, beside extreme labour and application, manifests the highest powers of genius and invention; as we here for the first time meet with several of the most important discoveries in the mathematics, and what have hitherto been considered as of much later invention; such as the Binomial Theorem; the Differential Method and Construction of Tables by Differences; the Interpolation by Differences; with Angular Sections, and several other ingenious compositions: a particular account of which may be seen in the Introduction to my Mathematical Tables.

This truly great man terminated his useful life the 26 of January 1630, and was buried in the choir of the chapel of Merton College. As to his character, he was not less esteemed for his great probity and other eminent virtues, than for his excellent skill in mathematics. Doctor Smith gives him the character of a man of great probity; easy of access to all; free from arrogance, moroseness, envy, ambition and avarice; a contemner of riches, and contented in his own situation; preferring a studious retirement to all the splendid circumstances of life. The learned Mr. Thomas Gataker, who attended his lectures when he was reader of mathematics at Cambridge, represents him as highly esteemed by all persons skilled in mathematics, both at home and abroad; and says, that desiring him once to give his judgment concerning judicial astrology, his

answer was, "that he conceived it to be a mere system of groundless conceits." Oughtred calls him the mirror of the age, for his excellent skill in geometry. And one of his successors at Gresham college, the learned Dr. Isaac Barrow, in his oration there upon his admission, has drawn his character more fully; celebrating his great abilities, skill, and industry, particularly in perfecting the invention of logarithms, which, without his care and pains, might have continued an imperfect and useless design.

His writings were more important than numerous: some of them were published by other persons: the list of the principal part of them as follows.

1. *A Table to find the Height of the Pole; the Magnetical Declination being given.* This was published in Mr. Thomas Blundevile's *Theoriques of the Seven Planets*: London 1602, 4to.

2. *Tables for the improvement of Navigation.* These consist of, A table of declination of every minute of the ecliptic, in degrees, minutes and seconds: A table of the sun's prosthaphaereses: A table of equations of the sun's ephemerides: A table of the sun's declination: Tables to find the height of the pole in any latitude, from the height of the pole star. These tables are printed in the 2d edition of Edward Wright's treatise, intitled, *Certain Errors in Navigation detected and corrected*; London 1610, 4to.

3. *A description of an Instrumental Table to find the Part Proportional, devised by Mr. Edward Wright.* This is subjoined to Napier's table of logarithms, translated into English by Mr. Wright, and after his death published by Briggs, with a preface of his own: Lond. 1616 and 1618, 12mo.

4. *Logarithmorum chiliad prima.* Lond. 1617, 8vo.

5. *Lucubrationes & Annotationes in opera posthuma J. Neperi*: Edinb. 1619, 4to.

6. *Euclidis Elementorum vi libri priores &c.* Lond. 1620, folio. This was printed without his name to it.

7. *A treatise of the North-west passage to the South Sea &c.* By H. B. Lond. 1622, 4to. This was reprinted in Purchas's *Pilgrims*, vol. 3, p. 852.

8. *Arithmetica Logarithmica, &c.* Lond. 1624, folio.

9. *Trigonometria Britannica, &c.* Goudæ 1633, folio.

10. *Two letters to archbishop Usher*.

11. *Mathematica ab antiquis minus cognita*.—This is a summary account of the most observable inventions of modern mathematicians, communicated by Mr. Briggs to Dr. George Hakewill, and published by him in his *Apologie*; Lond. folio.

Beside these publications, Briggs wrote some other pieces, that have not been printed: as,

(1). *Commentaries on the Geometry of Peter Ramus.*

(2). *Dux Epistolæ ad celeberrimum virum, Chr. Sever. Longomontanum.* One of these letters contained some remarks on a treatise of Longomontanus, about squaring the circle; and the other a defence of arithmetical geometry,

(3). *Animadversiones Geometricæ*: 4to.

(4). *De eodem Argumento*: 4to.—These two were in the possession of the late Mr. Jones. They both contain a great variety of geometrical propositions, concerning the properties of many figures, with several arithme-



arithmetical computations, relating to the circle, angular sections, &c.—The two following were also in possession of Mr. Jones.

(5). *A treatise of Common Arithmetic*; folio.

(6). *A letter to Mr. Clarke of Gravesend*, dated 25 Feb. 1606; with which he sends him the description of a ruler, called Bedwell's ruler, with directions how to use it.

BRIGGS (WILLIAM), an eminent physician in the latter part of the 17th century, was born at Norwich, for which town his father was four times member of parliament. He studied at the university of Cambridge. He afterwards travelled into France, where he attended the lectures of the famous anatomist Vicussens, at Montpellier. Upon his return he published his *Ophthalmographia*, in 1676. The year following he was made doctor of medicine at Cambridge, and soon after fellow of the college of physicians at London. In 1682 he resigned his fellowship to his brother; and the same year his *Theory of Vision* was published by Hook. The ensuing year he sent to the Royal Society a continuation of that discourse, which was published in their Transactions; and the same year he was appointed physician to St. Thomas's hospital. In 1684 he communicated to the Royal Society two remarkable cases relating to vision, which were likewise printed in their Transactions; and in 1685 he published a Latin version of his *Theory of Vision*, at the desire of Mr. Newton, afterwards Sir Isaac, then professor of mathematics at Cambridge, with a recommendatory epistle from him prefixed to it. He was afterwards made physician in ordinary to king William, and continued in great esteem for his skill in his profession till he died the 4th of September 1704.

BRIGGS's *Logarithms*, that species of them in which 1 is the logarithm of the ratio of 10 to 1, or the logarithm of 10. See LOGARITHMS.

BROKEN Number, the same as *Fraction*; which see.

BROKEN Ray, or *Ray of Refraction*, in Dioptrics, is the line into which an incident ray is refracted or broken, in crossing the second medium.

BROUNCKER, or BROUNKER, (WILLIAM), lord viscount of Castle Lyons in Ireland, son of Sir William Brounker, afterwards made viscount in 1645, was born about the year 1620. He very early discovered a genius for mathematics, in which he afterwards became very eminent. He was made doctor of physic at Oxford June 23, 1646. In 1657 and 1658, he was engaged in a correspondence by letters on mathematical subjects with Dr. John Wallis, who published them in his *Commercium Epistolicum*, printed 1658, at Oxford. He was one of the nobility and gentry who signed the remarkable declaration concerning king Charles the 2d, published in April 1660.

After the restoration, lord Brounker was made chancellor and keeper of the great seal to the queen consort, one of the commissioners of the navy, and master of St. Katherine's hospital near the tower of London. He was one of those great men who first formed the Royal Society, of which he was by the charter appointed the first president in 1662: which office he held, with great advantage to the Society, and honour to himself, till the anniversary election, Nov. 30, 1677. He died at his house in St. James's street, Westminster,

after, the 5th of April 1684; and was succeeded in his title by his younger brother Harry, who died in Jan. 1687.

Lord Brounker had several papers inserted in the Philosophical Transactions, the chief of which were, 1. Experiments concerning the Recoiling of Guns.—2. A Series for the Quadrature of the Hyperbola; which was the first series of the kind upon that subject.—3. Several of his letters to archbishop Usher were also printed in Usher's letters; as well as some to Dr. Wallis, in his *Commercium Epistolicum*, above mentioned.

BROWN (Sir WILLIAM), a noted physician and miscellaneous writer, of the 18th century. He was settled originally at Lynn in Norfolk, where he published a translation of Dr. Gregory's Elements of Catoptrics and Dioptrics; to which he added, 1. A Method for finding the Foci of all Specula and Lenses universally; as also Magnifying or Lessening a given object by a given Speculum or lens, in any assigned proportion.—2. A Solution of those Problems which Dr. Gregory has left undemonstrated.—3. A particular account of Microscopes and Telescopes, from Mr. Huygens; with the discoveries made by Catoptrics and Dioptrics.

Having acquired a competence by his profession, he removed to Queen's Square, Ormond Street, London, where he resided till his death, in 1774, at 82 years of age; leaving by his will two prize-medals to be annually contended for by the Cambridge poets.

Sir William Brown was a very facetious man; and a great number of his lively essays, both in prose and verse, were printed and circulated among his friends. The active part taken by him in the contest with the licentiates, in 1768, occasioned his being introduced by Mr. Foote in his *Devil upon Two Sticks*.—Upon Foote's exact representation of him with his identical wig and coat, tall figure, and glass stiffly applied to his eye, he sent him a card complimenting him on having so happily represented him; but as he had forgot his muff, he had sent him his own.—This good-natured way of resenting disarmed Foote.—He used to frequent the annual ball at the ladies boarding-school, Queen Square, merely as a neighbour, a good-natured man, and fond of the company of sprightly young folks. A dignitary of the church being there one day to see his daughter dance, and finding this upright figure stationed there, told him he believed he was Hermippus redivivus who lived *anhelitu puellarum*.—When he lived at Lynn, a pamphlet was written against him; which he nailed up against his house-door.—At the age of 80, on St. Luke's day 1771, he came to Batson's coffee-house in his laced coat and band, and fringed white gloves, to shew himself to Mr. Crosby, then lord mayor. A gentleman present observing that he looked very well, he replied, he had neither wife nor debts.

BULLIALD (ISMAEL), an eminent astronomer and mathematician, was born at Laon in the Isle of France in 1605. He travelled in his youth, for the sake of improvement, and gave very early proofs of his astronomical genius; and his riper years rendered him beloved and admired. Riccioli styled him, *Astronomus profunde indaginis*. He first published his dissertation



tation intituled, *Philolaus, sive de vero Systemate Mundi*; or his true system of the world, according to Philolaus, an ancient philosopher and astronomer. Afterward, in the year 1645, he set forth his *Astronomia Philolaica*, grounded upon the hypothesis of the earth's motion, and the elliptical orbit described by the planet's motion about a cone. To which he added tables intituled, *Tabula Philolaica*: a work which Riccioli says ought to be attentively read by all students of astronomy.—He considered the hypothesis, or approximation of bishop Ward, and found it not to agree with the planet Mars; and shewed in his defence of the Philolaic astronomy against the bishop, that from four observations made by Tycho on the planet Mars, that planet in the first and third quarters of the mean anomaly, was more forward than it ought to be according to Ward's hypothesis; but in the 2d and 4th quadrant of the same, the planet was not so far advanced as that hypothesis required. He therefore set about a correction of the bishop's hypothesis, and made it to answer more exactly to the orbits of the planets, which were most eccentric, and introduced what is called, by Street in his *Caroline Tables*, the Variation: for these tables were calculated from this correction of Bulliald's, and exceeded all in exactness that went before. This correction is, in the judgment of Dr. Gregory, a very happy one, if it be not set above its due place; and be accounted no more than a correction of an approximation to the true system: For by this means we are enabled to gather the coequate anomaly *a priori* and directly from the mean, and the observations are well enough answered at the same time; which, in Mercator's opinion, no one had effected before.—It is remarkable that the ellipsis which he has chosen for a planet's motion, is such a one as, if cut out of a cone, will have the axis of the cone passing through one of its foci, viz, that next the aphelion.

In 1657 was published his treatise *De Lineis Spiraliibus, Exerc. Geom. & Astron.* Paris, 4to.—In 1682 came out at Paris, in folio, his large work intituled, *Opus novum ad Arithmetica Infinitorum*: A work which is a diffuse amplification of Dr. Wallis's Arithmetic of Infinites, and which Wallis treats of particularly in the 80th chapter of his historical treatise of Algebra.—He wrote also two Admonitions to Astronomers. The first, concerning a new star in the neck of the Whale, appearing at some times, and disappearing at others. The 2d, concerning a nebulous star in the northern part of Andromeda's girdle, not discovered by any of the ancients. This star also appeared and disappeared by turns. And as these phenomena appeared new and surprizing, he strongly recommended the observing them to all that might be curious in astronomy.

**BURNING**, the action of fire on some pabulum or fuel, by which its minute parts are put into a violent motion, and some of them, assuming the nature of fire, fly off *in orbem*, while the rest are dissipated in vapour or reduced to ashes..

**BURNING-Glass**, or *Burning-Mirror*, a machine by which the sun's rays are collected into a point; and by that means their force and effect are extremely heightened, so as to burn objects placed in it.

Burning glasses are of two kinds, *convex* and *concave*. The convex ones are lenses, which acting according to the laws of refraction, incline the rays of light towards the axis, and unite them in a point or focus. The concave ones are mirrors or reflectors, whether made of polished metal or silvered glass, and which acting by the laws of reflection, throw the rays back into a point or focus before the glass.

The use of burning glasses it appears is very ancient, many of the old authors relating some effects of them. Diodorus Siculus, Lucian, Dion, Zonaras, Galen, Anthemius, Eustatius, Tzetzes, and others, relate that by means of them Archimedes set fire to the Roman fleet at the siege of Syracuse. Tzetzes is so particular in his account of this matter, that his description suggested to Kircher the method by which it was probably accomplished. That author says that "Archimedes set fire to Marcellus's navy by means of a burning glass composed of small square mirrors, moving every way upon hinges; which when placed in the sun's rays, directed them upon the Roman fleet, so as to reduce it to ashes at the distance of a bow-shot." And the burning power of reflectors is mentioned in Euclid's Optics, theor. 31. Again, Aristophanes, in his comedy of The Clouds, introduces Socrates as examining Strepsiades about a method he had discovered of getting clear of his debts. He replies, that "he thought of making use of a burning-glass which he had hitherto used in kindling his fire; for should they bring a writ against me, I'll immediately place my glass in the sun at some little distance from it, and set it on fire." Pliny and Lactantius have also spoken of glasses that burn by refraction. The former calls them *balls* or *globes* of *crystal* or *glass*, which being exposed to the sun, transmit a heat sufficient to set fire to cloth, or corrode the dead flesh of those patients who stand in need of caustics; and the latter, after Clemens Alexandrinus, observes that fire may be kindled by interposing glasses filled with water between the sun and the object, so as to transmit the rays to it.

Among the ancients the most celebrated burning mirrors were those of Archimedes and Proclus; by the former was burnt the fleet of Marcellus, as above mentioned; and by the latter, the navy of Vitellius, besieging Byzantium, according to Zonaras was burnt to ashes.

Among the moderns, the most remarkable burning-glasses, are those of Mazine of 20 inches diameter: of Sepatala of Milan, near 42 inches diameter, and which burnt at the distance of 15 feet; of Settala of Villette, of Tschirnhausen, of Buffon, of Trudaine, and of Parker.

Villette, a French artist at Lyons, made a large mirror, which was bought by Tavernier, and presented to the king of Prussia; a second, bought by the king of Denmark; a third, presented to the Royal Academy by the king of France; and a 4th came to England, and was publicly shewn. This mirror is 47 inches wide, being a segment of a sphere of 76 inches radius; so that its focus is about 38 inches from the vertex; and its substance is a composition of tin, copper, and tin-glass. Some of its effects were as follow:

A silver



A silver sixpence melted in	sec.
A George the 1st's halfpenny in	7½
and runs with a hole in	16
Tin melts in	34
Cast iron in	3
Slate in	16
A fossil shell calcines in	3
Piece of Pompey's pillar vitrifies, the black part in	7
the white part in	50
Copper ore in	54
Bone calcines in 4, and vitrifies in	8
	33

An emerald melts into a substance like a torquois stone; a diamond weighing 4 grains loses  $\frac{7}{8}$  of its weight: the asbestos vitrifies; as all other bodies will do if kept long enough in the focus; but when once vitrified, the mirror can go no farther with them. *Philos. Transf.* vol. iv. pa. 198.

Tschirnhausen's reflecting mirrors produced equally surprizing effects; as they may be seen described in the *Acta Erudit.* for 1687, pa. 52. And other persons have made very good ones of wood, straw, paper, ice, and other substances capable of taking a proper form and polish.

Every lens, whether convex, plano-convex, or convexo-convex, collects the sun's rays, dispersed over its convexity, into a point by refraction; and it is therefore a burning-glass. The most considerable of this kind is that made by Tschirnhausen, and described in the same *Acta Erudit.* The diameters of his lenses are from 3 to 4 feet, having the focus at the distance of 12 feet, and its diameter an inch and a half. To make the focus more vivid, the rays are collected a second time, by a second lens parallel to the first, and placed at such a distance that the diameter of the cone of rays formed by the first lens is equal to the diameter of the second; so that it receives them all; and the focus is reduced from an inch and a half to half the quantity, and consequently its force is quadrupled. This glass vitrifies tiles, slates, pumice-stones &c. in a moment. It melts sulphur, pitch, and all resins, under water; the ashes of vegetables, woods and other matters, are transmuted into glass; and every thing applied to its focus is either melted, changed into a calx, or into fumes. The author observes that it succeeds best when the matter applied is laid on a hard charcoal well burnt. —But though the force of the solar rays be thus found so surprizing, yet the rays of the full moon, collected by the same burning-glass, do not shew the least increase of heat.

Sir Isaac Newton presented a burning-glass to the Royal Society, consisting of 7 concave glasses, so placed that all their foci join in one physical point. Each glass is about 11½ inches diameter: six of them are placed contiguous to, and round the seventh, forming a kind of spherical segment, whose subtense is about 34½ inches: the common focus is about 22½ inches distant, and about an inch in diameter. This glass vitrifies brick or tile in 1 second, and melts gold in 30 seconds.

M. Buffon also made a variety of very powerful burning-glasses, both as mirrors and as lenses; but at length concluded with one which is probably of the same nature with that of Archimedes, and consisted of 400 mirrors

reflecting their rays all to one point, and with which he could melt lead and tin at the distance of 140 feet; and with others he consumed substances at the distance of 210 feet. See *Philos. Transf.* vol. 44; or Buffon's *Histoire Naturelle*, Suppl. vol. 1; or Montucla's *Histoire des Math.* vol. i. pa. 246.

It would seem there is no substance capable of resisting the efficacy of modern burning-glasses; though water &c. are not affected by them at all. Thus, Messrs Macquer and Baumé have succeeded in melting small portions of platina by means of a concave glass, 22 inches diameter, and 28 inches focus; though this metal is not fusible by the strongest fires that can be excited in furnaces, or sustained by any chemical apparatus. Yet it was long since observed, by the Academicians del Cimento, that spirit of wine could not be fired by any burning-glass which they used; and notwithstanding the great improvements these instruments have since received, M. Nollet has not been able, by the most powerful burning mirrors, to set fire to any inflammable liquors whatever.

However, a large burning lens, for fusing and vitrifying such substances as resist the fires of furnaces, and especially for the application of heat in vacuo, and in certain other circumstances in which heat cannot be applied by other means, has long been a desideratum with persons concerned in philosophical experiments: and this it appears is now in a great measure accomplished by Mr. Parker, an ingenious glass manufacturer in Fleet-street, London. His lens is made of flint glass, and is 3 feet in diameter, but when fixed in its frame exposes a surface of 32 inches in the clear; the length of the focus is 6 feet 8 inches, and its diameter one inch. The rays from this large lens are received and transmitted through a smaller, of 13 inches diameter in the clear within the frame, its focal length 29 inches, and diameter of its focus 3-8ths of an inch: so that this second lens increases the power of the former more than 7 times, or as the square of 8 to the square of 3.

From a great number of experiments made with this lens, the following are selected to serve as specimens of its powers:

Substances fused; with their weight, and time of fusion.	Time in sec.	Weight in grs.
Scoria of wrought iron	2	12
Common slate	2	10
Silver, pure	3	20
Platina, pure	3	10
Nickell	3	16
Cast Iron, a cube	3	10
Kearsh	3	10
Gold, pure	4	20
Cryстал pebble	6	7
Cauk, or terra ponderosa	7	10
Lava	7	10
Asbestos	10	10
Bar Iron, a cube	12	10
Steel, a cube	12	10
Garnet	17	10
Copper, pure	20	33
Onyx	20	10

Zeolites



## SUBSTANCES FUSED, &amp;c.

	Time in sec.	Wgt. in grs.
Zeolites - - - - -	23	10
Pumice Stone - - - - -	24	10
Oriental Emerald - - - - -	25	2
Jasper - - - - -	25	10
White-Agate - - - - -	30	10
Flint, oriental - - - - -	30	10
Topaz, or chrysolite - - - - -	45	3
Common Limestone - - - - -	55	10
White rhomboidal Spar - - - - -	60	10
Volcanic Clay - - - - -	60	10
Cornish Moorstone - - - - -	60	10
Rough Cornelian - - - - -	75	0
Rotten Stone - - - - -	80	10

BURNING Zone, or *Torrid Zone*, the space within  $23\frac{1}{2}$  degrees of the equator, both north and south.

**BUSHEL**, a measure of capacity for dry goods; as grain, pulse, fruits, &c; containing 4 pecks, or 8 gallons, or  $\frac{1}{4}$  of a quarter. By act of Parliament, made in 1697, it was ordained that "Every round bushel with a plain and even bottom, being made  $18\frac{1}{2}$  inches wide throughout, and 8 inches deep, shall be esteemed a Legal Winchester Bushel, according to the standard in his majesty's exchequer." Now a bushel being thus made will contain 2150.42 cubic inches, and consequently the corn gallon contains only 268 $\frac{4}{5}$  cubic inches.

**BUTMENTS**, are those supporters, or props, by which the feet of arches, or the extremities of bridges are supported; and should be made very strong and firm.

**BUTTRESS**, is an arch, or a mass of masonry, serving to support the sides of a building, wall, or the like, on the outside. See **ARCH**, and **ARCH-BOUTANT**.

## C.

## C A I

**CAILLE** (NICHOLAS LEWIS DE LA), an eminent French mathematician and astronomer, was born at Rumigny in the diocese of Rheims in 1713. His father having quitted the army, in which he had served, amused himself in his retirement with studying mathematics and mechanics, in which he proved the happy author of several inventions of considerable use to the public. From this example of his father, our author almost in his infancy took a fancy to mechanics, which proved of signal service to him in his maturer years. At school he discovered early tokens of genius. He next came to Paris in 1729; where he studied the classics, philosophy and mathematics. He afterwards studied divinity in the college de Navarre, with the view of embracing the ecclesiastical life: however he never entered into priest's orders, apprehending that his astronomical studies, to which he had become much devoted, might too much interfere with his religious duties. His turn for astronomy soon connected him with the celebrated Cassini, who procured him an apartment in the observatory; where, assisted by the counsels of this master, he soon acquired a name among the astronomers. In 1739 he was joined with M. Cassini de Thury, son to M. Cassini, in verifying the meridian through the whole extent of France: and in the same year he was named professor of mathematics in the college of Mazarine. In 1741 our author was admitted into the Academy of Sciences as an adjoint member for astronomy; and had many excellent papers inserted in their memoirs; beside which he published several useful treatises, viz, Elements of Geometry, Astronomy, Mechanics, and Optics. He also carefully computed all the eclipses of the sun and moon that had happened

Vol. I.

## C A I

since the christian era, which were printed in the work entitled *L'Art de verifier les dates*, &c, Paris, 1750, in 4to. He also compiled a volume of astronomical ephemerides for the years 1745 to 1755; another for the years 1755 to 1765; and a third for the years 1765 to 1775: as also the most correct solar tables of any; and an excellent work entitled *Astronomia fundamenta novissimis solis & stellarum observationibus stabilita*.

Having gone through a seven years series of astronomical observations in his own observatory in the Mazarine college, he formed the project of going to observe the southern stars at the Cape of Good Hope: being countenanced by the court, he set out upon this expedition in 1750, and in the space of two years he observed there the places of about 10 thousand stars in the southern hemisphere that are not visible in our latitudes, as well as many other important elements, viz, the parallaxes of the sun, moon, and some of the planets, the obliquity of the ecliptic, the refractions, &c. Having thus executed the purpose of his voyage, and no present opportunity offering for his return, he thought of employing the vacant time in another arduous attempt; no less than that of taking the measure of the earth, as he had already done that of the heavens. This indeed had been done before by different sets of learned men both in Europe and America; some determining the quantity of a degree at the equator, and others at the arctic circle: but it had not as yet been decided whether in the southern parallels of latitude the same dimensions obtained as in the northern. His labours were rewarded with the satisfaction he wished for; having determined a distance of 4108 $\frac{1}{4}$  feet from a place called *Klip-Fontyn* to the

H h

Cape,



Cape, by means of a base of 38802 feet, three times actually measured: whence he discovered a new secret of nature, namely, that the radii of the parallels in south latitude are not the same length as those of the corresponding parallels in north latitude. About the 23d degree of south latitude he found a degree on the meridian to contain 342222 Paris feet. The court of Versailles also sent him an order to go and fix the situation of the Isles of France and of Bourbon. While at the Cape too he observed a wonderful effect of the atmosphere in some states of it: Although the sky at the Cape be generally pure and serene, yet when the southeast wind blows, which is pretty often, it is attended with some strange and even terrible effects: the stars look larger, and seem to dance; the moon has an undulating tremor; and the planets have a sort of beard like comets.

M. de la Caille returned to France in the autumn of 1754, after an absence of about 4 years; loaded, not with the spoils of the east, but with those of the southern heavens, before then almost unknown to astronomers. Upon his return, he first drew up a reply to some strictures which the celebrated Euler had published relative to the meridian: after which he settled the results of the comparison of his observations for the parallaxes, with those of other astronomers: that of the sun he fixed at  $9\frac{1}{2}''$ ; of the moon at  $56'56''$ ; of Mars in his opposition,  $36''$ ; of Venus  $38''$ . He also settled the laws by which astronomical refractions are varied by the different density or rarity of the air, by heat or cold, and by dryness or moisture. And lastly he shewed an easy and practicable method of finding the longitude at sea by means of the moon. His fame being now celebrated every where, M. de la Caille was soon elected a member of most of the academies and societies of Europe, as London, Bologna, Petersburg, Berlin, Stockholm, and Gottingen.

In 1760 our author was attacked with a severe fit of the gout; which however did not interrupt the course of his studies; for he then planned out a new and large work, no less than a history of astronomy through all ages, with a comparison of the ancient and modern observations, and the construction and use of the instruments employed in making them. Towards the latter part of 1761, his constitution became greatly reduced; though his mind remained unaffected, and he resolutely persisted in his studies to the last; death only putting an end to his labours the 21st of March 1762, at 49 years of age; after having committed his manuscripts to the care and discretion of his esteemed friend M. Maraldi.

Beside the publications before mentioned, and perhaps some others also, he had a vast number inserted in the volumes of the Memoirs of the French Academy of Sciences, much too numerous indeed, though very important, to be here all mentioned particularly; suffice it therefore just to distinguish the years of those volumes in which his pieces are to be found, by the following list of them, viz, 1741, 1742, 1743, 1744, 1745, 1746, 1747, 1748, 1749, 1750, 1751, 1752, 1753, 1754, 1755, 1756, 1757, 1758, 1759, 1760, 1761, 1763; in all or most of which years there are two or three or more of his papers.

CAISSON, in Architecture, a kind of chest or flat-

bottomed boat, in which the pier of a bridge is built, then sunk to the bottom of the water, and the sides loosened and taken off from the bottom, by a contrivance for that purpose; the bottom of the caisson being left under the pier as a foundation to it. The caisson is kept afloat till the pier is built to above the height of low-water mark; and for that purpose, its sides are either made of more than that height at first, or else gradually raised to it as it sinks by the weight of the work, so as always to keep its top above water. Mr. Labéye tells us, that the caissons in which he built some of the piers of Westminster bridge, contained above 150 load of fir timber, of 40 cubic feet each, and that it was of more tonnage or capacity than a 40 gun ship of war.

CAISSON, in Military Affairs, is sometimes used for a chest; and in particular for a bomb or shell chest, and is used as a superficial mine, or fourneau. This is done by filling a chest either with gun powder and loaded shells, or else with shells alone, and burying it in a spot where an enemy, besieging a place, is expected to come, and then firing it by a train to blow the men up.

CALCULATION, the act of computing several sums, by adding, subtracting, multiplying, dividing, &c. From *calculus*, in allusion to the practice of the ancients, who used *calculi*, or little stones, in making computations, in taking suffrages, and in keeping accounts, &c; as we now use counters, figures, &c. Calculation is more particularly used to signify the computations in astronomy, trigonometry, &c, for making tables of astronomy, of logarithms, ephemerides, finding the times of eclipses, and such like.

CALCULATOR, a person who makes or performs calculations.—It is also the name given by Mr. Ferguson to a machine in the shape of an orrery, which he constructed for exhibiting the motions of the earth and moon, and resolving a variety of astronomical problems. See his *Astron.* 4to pa. 265, or 8vo pa. 393.

CALCULATOIRES, were anciently accountants who reckoned their sums by *calculi*, or little stones, or counters.—In ancient canons too we find a sort of diviners or enchanters, censured under the denomination of *calculatores*.

CALCULUS denotes primarily a small stone, pebble, or counter, used by the ancients in making calculations or computations, taking of suffrages, playing at tables, and the like.

CALCULUS denotes now a certain way of performing mathematical investigations and resolutions. Thus, we say the Arithmetical or Numeral Calculus, the Algebraical Calculus, the Differential Calculus, the Exponential Calculus, the Fluxional Calculus, the Integral Calculus, the Literal or Symbolical Calculus, &c; for which, see each respective word.

Arithmetical or Numeral CALCULUS, is the method of performing arithmetical computations by numbers. See ARITHMETIC, and NUMBER.

Algebraical, Literal, or Symbolical CALCULUS, is the method of performing algebraical calculations by letters or other symbols. See ALGEBRA.

Differential CALCULUS, is the arithmetic of the indefinitely small differences of variable quantities; a mode of computation much used by foreign mathematicians, and introduced by Leibnitz, as similar to the method



method of Fluxions of Newton. See DIFFERENTIAL &c.

*Exponential* CALCULUS, is the applying the fluxional or differential methods to exponential quantities; such as  $a^x$ , or  $x^x$ , or  $ay^x$ , &c. See EXPONENTIAL.

*Fluxional* CALCULUS, is the method of fluxions, invented by Newton. See FLUXIONS.

*Integral* CALCULUS, or *Summatorius*, is a method of integrating, or summing up differential quantities; and is similar to the finding of fluents. See INTEGRAL and FLUENT.

CALCULUS *Literalis*, or *Literal Calculus*, is the same with algebra, or specious arithmetic, so called from its using the letters of the alphabet; in contradistinction to numeral arithmetic, in which figures are used.

CALENDAR, or KALENDAR, a distribution of time as accommodated to the uses of life; or an Almanac, or table, containing the order of days, weeks, months, feasts, &c, occurring in the course of the year: being so called from the word *Calendæ*, which among the Romans denoted the first days of every month, and anciently was written in large characters at the head of each month. See ALMANAC, CALENDs, MONTH, TIME, YEAR, &c.

In Calendars the days were originally divided into octoades, or eights; but afterwards, in imitation of the Jews, they were divided into hebdomades, or sevens, for what we now call a week: which custom, Scaliger observes, was not in use among the Romans till after the time of Theodosius.

Divers calendars are established in different countries, according to the different forms of the year, and distributions of time: As the Persian, the Roman, the Jewish, the Julian, the Gregorian, &c, calendars. — The ancient Roman Calendar is given by Ricciolus, Struvius, Danet, and others; in which we perceive the order and number of the Roman holy-days and work-days. — The Jewish calendar was fixed by Rabbi Hillel, about the year 360; from which time the days of their year may be reduced to those of the Julian calendar. — The three Christian calendars are given by Wolfius in his Elements of Chronology; as also the Jewish and Mohamedan calendars. Other writers on the calendars are Vieta, Clavius, Scaliger, Blondel, &c.

The Roman CALENDAR was first formed by Romulus, who distributed time into several periods for the use of his followers and people. He divided the year into 10 months, of 304 days; beginning on the first of March, and ending with December.

Numa reformed the calendar of Romulus. He added the months of January and February, making it to commence on the first of January, and to consist of 355 days. But as this was evidently deficient of the true year, he ordered an intercalation of 45 days to be made every 4 years, in this manner, viz, Every 2 years an additional month of 22 days, between February and March; and at the end of each two years more, another month of 23 days; the month thus interposed, being called Marcedonius, or the intercalary February.

Julius Cæsar, with the aid of Sosigenes, a celebrated astronomer of those times, farther reformed the Roman calendar, from whence arose the Julian calendar, and the Julian or old style. Finding that the sun per-

formed his annual course in 365 days and a quarter nearly, he divided the year into 365 days, but every 4th year 366 days, adding a day that year before the 24th of February, which being the 6th of the calends, and being thus reckoned twice, gave occasion to the name *bissextile*, or what we also call leap-year.

This calendar was farther reformed by order of the pope Gregory XIII, from whence arose the term Gregorian calendar and style, or what we also call the new style, which is now observed by almost all European nations. The year of Julius was too long by nearly 11 minutes, which amounts to about 3 days in 400 years; the pope therefore, by the advice of Clavius and Ciaccinius, ordained that there should be omitted a day in every 3 centuries out of 4; so that every century, which would otherwise be a bissextile year, is made to be only a common year, excepting only such centuries as are exactly divisible by 4, which happens once in 4 centuries. See BISSEXTILE. This reformation of the calendar, or the new style, as we call it, commenced in the countries under the popish influence, on the 4th of October 1582, when 10 days were omitted at once, which had been over-run since the time of the council of Nice, in the year 325, by the surplus of 11 minutes each year. But in England it only commenced in 1752, when 11 days were omitted at once, the 3d of September being accounted the 14th that year; as the surplus minutes had then amounted to 11 days.

*Julian Christian* CALENDAR, is that in which the days of the week are determined by the letters A, B, C, D, E, F, G, by means of the solar cycle; and the new and full moons, particularly the paschal full moon, with the feast of Easter, and the other moveable feasts depending upon it, by means of golden numbers, or lunar cycles, rightly disposed through the Julian year. See CYCLE, and GOLDEN NUMBER.

In this calendar, it is supposed that the vernal equinox is fixed to the 21st day of March; and that the golden numbers, or cycles of 19 years, constantly indicate the places of the new and full moons; though both are erroneous; and from hence arose a great irregularity in the time of Easter.

*Gregorian* CALENDAR, is that which, by means of Epacts, rightly disposed through the several months, determines the new and full moons, with the time of Easter, and the moveable feasts depending upon it, in the Gregorian year. This differs therefore from the Julian calendar, both in the form of the year, and in as much as epacts are substituted instead of golden numbers. See EPACT.

Though the Gregorian calendar be more accurate than the Julian, yet it is not without imperfections, as Scaliger and Calvisius have fully shewn; nor is it perhaps possible to devise any one that shall be quite perfect. Yet the Reformed Calendar, and that which is ordered to be observed in England, by act of Parliament made the 24th of George II, come very near to the point of accuracy: For, by that act it is ordered that "Easter-day, on which the rest depend, is always the first Sunday after the full moon, which happens upon, or next after the 21st day of March; and if the full-moon happens upon a Sunday, Easter-day is the Sunday after."

*Reformed, or Corrected*, CALENDAR is that which, rejecting



rejecting all the apparatus of golden numbers, epacts, and dominical letters, determines the equinox, and the paschal full-moon, with the moveable feasts depending upon it, by computation from astronomical tables. This calendar was introduced among the protestant states of Germany in the year 1700, when 11 days were omitted in the month of February, to make the corrected style agree with the Gregorian. This alteration in the form of the year, they admitted for a time; in expectation that, the true quantity of the tropical year being at length more accurately determined by observation, the Romanists would agree with them on some more convenient intercalation.

*French New CALENDAR*, is a quite new form of calendar that commenced in France on the 22d of September 1792. At the time of printing this (viz, in July 1794), it does not certainly appear whether this new calendar will be made permanent or not; but merely as a curiosity in the science of chronology, a very brief notice of it may here be added, as follows.

The year, in this calendar, commences at midnight the beginning of that day in which falls the true autumnal equinox for the observatory of Paris. The year is divided into 12 equal months, of 30 days each; after which 5 supplementary days are added, to complete the 365 days of the ordinary year: these 5 days do not belong to any month. Each month is divided into three decades of 10 days each; distinguished by 1st, 2d, and 3d decade. All these are named according to the order of the natural numbers, viz, the 1st, 2d, 3d, &c, month, or day of the decade, or of the supplementary days. The years which receive an intercalary day, when the position of the equinox requires it, which we call embolismic or bissextile, they call olympic; and the period of four years, ending with an olympic year, is called an olimpiade; the intercalary day being placed after the ordinary five supplementary days, and making the last day of the olympic year. Each day, from midnight to midnight, is divided into 10 parts, each part into 10 others, and so on to the last measurable portion of time.

In this calendar too the months and days of them have new names. The first three months of the year, of which the autumn is composed, take their etymology, the first from the vintage which takes place from September to October, and is called *vendemaire*; the second, *brumaire*, from the mists and low fogs, which shew as it were the transudation of nature from October to November: the third, *frimaire*, from the cold, sometimes dry and sometimes moist, which is felt from November to December. The three winter months take their etymology, the first, *nivose*, from the snow which whitens the earth from December to January; the second, *pluviose*, from the rains which usually fall in greater abundance from January to February; the third, *ventose*, from the wind which dries the earth from February to March. The three spring months take their etymology, the first, *germinal*, from the fermentation and development of the sap from March to April; the second, *floreale*, from the blowing of the flowers from April to May: the third, *prairial*, from the smiling fecundity of the meadow crops from May to June. Lastly, the three summer months take their

etymology, the first, *messidor*, from the appearance of the waving ears of corn and the golden harvests which cover the fields from June to July; the second, *thermidor*, from the heat, at once solar and terrestrial, which inflames the air from July to August; the third, *fructidor*, from the fruits gilt and ripened by the sun from August to September. Thus, the whole 12 months are,

AUTUMN.	SPRING.
Vendemaire	Germinal
Brumaire	Floreale
Frimaire.	Prairial.
WINTER.	SUMMER.
Nivose	Messidor
Pluviose	Thermidor
Ventose.	Fructidor.

From these denominations it follows, that by the mere pronounciation of the name of the month, every one readily perceives three things and all their relations, viz, the kind of season, the temperature, and the state of vegetation: for instance, in the word *germinal*, his imagination will easily conceive, by the termination of the word, that the spring commences; by the construction of the word, that the elementary agents are busied; and by the signification of the word, that the buds unfold themselves.

As to the names of the days of the week, or decade of 10 days each, which they have adopted instead of seven, as these bear the stamp of judicial astrology and heathen mythology, they are simply called from the first ten numbers; thus,

Primidi	Sextidi
Duodi	Septidi
Tridi	Octidi
Quartidi	Nonidi
Quintidi	Decadi.

In the almanac, or annual calendar, instead of the multitude of fables, one for each day of the year, as in the popish calendars, they annex to every day the name of some animal, or utensil, or work, or fruit, or flower, or vegetable, &c, appropriate and most proper to the times.

*Astronomical CALENDAR*, an instrument engraven upon copper-plates, printed on paper, and pasted on board, with a brass slider which carries a hair, and shews by inspection, the sun's meridian, altitude, right ascension, declination, rising, setting, amplitude, &c, to a greater exactness than can be shewn by the common globes.

*CALENDS*, *Calendæ*, in the Roman Chronology, denoted the first days of each month; being so named from *καλεω*, *calo*, *I call*, or *proclaim*; because that, before the publication of the Roman *Fasts*, and counting their months by the motion of the moon, a priest was appointed to observe the first appearance of the new moon; who, having seen her, gave notice to the president of the sacrifices to offer one; and calling the people together, he proclaimed unto them how they should reckon the days until the nones; pronouncing the word *Caleo* 5 times if the nones should happen on the 5th day, or seven times if they happened on the 7th day of the month.

The calends were reckoned backwards, or in a retrograde



etrograde order: thus, for example, the first of May being the calends of May; the last or 30th day of April, was the *pridie calendarum*, or 2d of the calends of May; the 29th of April, the 3d of the calends, or before the calends: and so back to the 13th, where the ides commence; which are likewise numbered backwards to the 5th, where the nones begin; which are also reckoned after the same manner to the first day of the month, which is the calends of April.

Hence comes this rule to find the day of the calends answering to any day of the month, viz, Consider how many days of the month are yet remaining after the day proposed, and to that number add 2, for the number of or from the calends. For example, suppose it were the 23d day of April, it would then be the 9th of the calends of May: for April containing 30 days, from which 23 being taken, there remains 7; to which 2 being added, makes the sum 9. And the reason for this addition of the constant number 2, is because the last day of the month is called the 2d of the calends of the month following.

CALIBER, or CALIPER, is the thickness or diameter of a round body, particularly the bore or width of a piece of ordnance, or that of its ball.

CALIBER-Compasses, or CALIPER-Compasses, or simple CALIPERS, a sort of compasses made with bowed or arched legs, the better to take the diameter of any round body; as the diameters of balls, or the bores of guns; or the diameter, and even the length of casks, and such like. The best sort of calipers usually contain the following articles, viz, 1st, the measure of convex diameters in inches &c; 2d, of concave diameters: 3d, the weight of iron shot of given diameters; 4th, the weight of iron shot for given gun bores; 5th, the degrees of a semicircle; 6th, the proportion of troy and averdupois weight; 7th, the proportion of English and French feet and pounds weight; 8th, factors used in circular and spherical figures; 9th, tables of the specific gravities and weights of bodies; 10th, tables of the quantity of powder necessary for the proof and service of brass and iron guns; 11th, rules for computing the number of shot or shells in a complete pile; 12th, rules for the fall or descent of heavy bodies; 13th, rules for the raising of water; 14th, rules for firing artillery and mortars; 15th, a line of inches; 16th, logarithmic scales of numbers, sines, versed sines, and tangents; 17th, a sectoral line of equal parts, or the line of lines; 18th, a sectoral line of planes and superficies; and 19th, a sectoral line of solids.

CALIPPIC Period, in Chronology, a period of 76 years, continually recurring; at every repetition of which, it was supposed, by its inventor Calippus, an Athenian astronomer, that the mean new and full moons would always return to the same day and hour.

About a century before, the golden number, or cycle of 19 years, had been invented by Meton, which Calippus finding to contain 19 of Nabonassar's year, 4 days and  $\frac{3}{4}$ , to avoid fractions he quadrupled it, and so produced his period of 76 years, or 4 times 19; after which he supposed all the lunations &c would regularly return to the same hour. But neither is this exact, as it brings them too late by a whole day in 225 years.

CAMBER-BEAM, a piece of timber cut arch-wise, or with an obtuse angle in the middle. They are commonly used in platforms, as for church-roofs, and other occasions where long timbers are wanted to lie at a small slope. A camber-beam is much stronger than another of the same dimensions; for being laid with the hollow side downwards, and having good butments at the ends, they serve for a kind of arch.

CAMELEON, one of the constellations of the southern hemisphere, near the south pole, and invisible in our latitude. There are 10 stars marked in this constellation in Sharp's catalogue.


CAMELOPARDALUS, a new constellation of the northern hemisphere, formed by Hevelius, consisting of 32 stars first observed by him. It is situated between Cepheus, Cassiopeia, Perseus, the Two Bears, and Draco; and it contains 58 stars in the British catalogue.

CAMERA *Æolia*, a name given by Kircher to a contrivance for blowing the fire, for the fusion of ores, without bellows. This is effected by means of water falling through a funnel into a close vessel, which sends from it so much air or vapour, as continually blows the fire. See Hook's Philos. Coll. n° 3, pa. 8c.

CAMERA *Lucida*, a contrivance of Dr. Hook to make the image of any thing appear on a wall in a light room, either by day or night. See Philos. Trans. n° 38, pa. 741.

CAMERA *Obscura*, or *Dark Chamber*, an optical machine or apparatus, representing an artificial eye, by which the images of external objects, received through a double convex glass, are shewn distinctly, and in their native colours, on a white ground placed within the machine, in the focus of the glass. The first invention of the camera obscura is ascribed to John Baptista Porta. See his *Magia Naturalis*, lib. 17, cap. 6, where he largely describes the effects of it. See also the end of s'Gravefande's *Perspective*, and other optical writers, for the construction and uses of various sorts of camera obscuras.

This machine serves for many useful and entertaining purposes. For example, it is very useful in explaining the nature of vision, representing a kind of artificial eye: it exhibits very diverting sights or spectacles; shewing images perfectly like their objects, clothed in their natural colours, but more intense and vivid, and at the same time accompanied with all their motions; an advantage which no art can imitate: and by this instrument, a person unacquainted with painting, or drawing, may delineate objects with the greatest accuracy of drawing and colouring.



Theory of the Camera Obscura. The theory and principle



principle of this instrument may be thus explained. If any object *AB* radiate through a small aperture *L*, upon a white ground opposite to it, within a darkened room, or box, &c; the image of the object will be painted on that ground in an inverted situation. For, by the smallness of the aperture, the rays from the object will cross each other there, the image of the point *A* being at *a*, and that of *B* at *b*; so that the whole object *AB* will appear inverted, as at *ab*. And as the corresponding rays make equal angles on both sides of the aperture, if the ground be parallel to the object, their heights will be to each other directly as their distances from the aperture.


*Construction of a Camera Obscura*, by which the images of external objects shall be represented distinctly, and in their genuine colours. 1st, Darken a chamber that has one of its windows looking towards a place containing various objects to be viewed; leaving only a small aperture open in one shutter. 2d, In this aperture fit a proper lens, either plano-convex, or convex on both sides; the convexity forming a small portion of a large sphere. But note, that if the aperture be made very small, as of the size of a pea, the objects will be represented even without any lens at all. 3d, At a proper distance to be determined by trials, stretch a paper or white cloth, unless there be a white wall at that distance, to receive the images of the objects: or the best way is to have some plaister of Paris cast on a convex mould, so as to form a concave smooth surface, and of a curvature and size adapted to the lens, to be placed occasionally at the proper distance. 4th, If it be rather desired to have the objects appear erect, instead of inverted, this may be done either by placing a concave lens between the centre and the focus of the first lens; or by reflecting the image from a plane speculum inclined to the horizon in an angle of 45 degrees; or by having two lenses included in a draw-tube, instead of one.

That the images be clear and distinct, it is necessary that the objects be illuminated by the sun's light shining upon them from the opposite quarter: so that, in a western prospect the images will be best seen in a forenoon, an eastern prospect the afternoon, and a northern prospect about noon; a southern aspect is the least eligible of any. But the best way of any is, if the lens be fixed in a proper frame, on the top of a building, and made to move easily round in all directions, by a handle extended to the person who manages the instrument; the images being then thrown down into a dark room immediately below it, upon a horizontal round plaister of Paris ground: for thus a view of all the objects quit around may easily be taken in the space of a few minutes; as is the case of the excellent camera obscura placed on the top of the Royal Observatory at Greenwich.


The objects will be seen brighter, if the spectator first wait a few minutes in the dark. Care should also be taken, that no light escape through any chinks; and that the ground be not too much illuminated. It may further be observed, that the greater distance there is between the aperture and the ground, the larger the images will be; but then at the same time the brightness is weakened more and more with the increase of distance.

*To construct a Portable Camera Obscura.* 1st, Provide a small box or chest of dry wood, and of about 10 inches broad, and 2 feet long or more, according to the size of the lenses. 2d, In one side of it, as *BD*, fit a sliding tube *EF* with two lenses; or, to have the image at a less distance from the tube, with three lenses, convex on both sides; the diameter of the two outer ones to be about 7 inches, but that of the inner to be less, as  $4\frac{3}{4}$  or 5 inches. 3d, At a proper distance, within the box, set up perpendicularly an oiled paper *GH*, so that images thrown upon it may be seen through. 4th, In the opposite side, at *I*, make a round hole, for a person to look conveniently through with both eyes. Then if the tube be turned towards the objects, and the lenses be placed by trials at the proper distance, by sliding the tube in and out, the objects will be seen delineated on the paper, erect as before.

The machine may be better accommodated for drawing, by placing a mirror to pass from *G* to *C*; for this will reflect the image upon a rough glass plane, or an oiled paper, placed horizontally at *AB*; and a copy of it may there be sketched out with a black-lead pencil.



*Another Portable Camera Obscura* is thus made. 1st, On the top of a box or chest raise a little turret *HI*, open towards the object *AB*. 2d, Behind the aperture, incline a small mirror *ab* at an angle of 45 degrees, to reflect the rays *Aa* and *Bb* upon a lens *G* convex on both sides, and included in a tube *GL*. Or the lens may be fixed in the aperture. 3d, At the dis-



tance of the focus place a table, or board *EF*, covered with



with a white paper, to receive the image *ab*. Lastly, In MN make an oblong aperture to look through; and an opening may also be made in the side of the box, for the convenience of drawing.

This sort of camera is easily changed into a show-box, for viewing prints, &c: placing the print at the bottom of the box, with its upper part inwards, where it is enlightened through the front, left open for this purpose, either by day or candle-light; and the print may be viewed through the aperture in HI.—A variety of contrivances for this purpose may be seen described in Harris's Optics, b. ii. sect. 4. — Mr. Storer has also procured a patent for an instrument of this sort, which he calls a delineator; being formed of two double convex lenses and a plane mirror, fitted into a proper box. One lens is placed close to the mirror, making with it an angle of 45 degrees; the other being placed at right angles to the former, and fixed in a moveable tube. If the moveable lens be directed towards the object, which is to be viewed or copied, and moved nearer to or farther from the mirror, till the image is distinctly formed on a greyed glass, laid upon that surface of the upper lens which is next the eye, it will be found more sharp and vivid than those formed in the common instruments; because the image is taken up so near the upper lens. And by increasing the diameter and curvature of the lenses, the effect will be much heightened.

CAMUS, (*Charles-Stephen-Lewis*), a celebrated French mathematician, Examiner of the royal Schools of Artillery and Engineers, Secretary and Professor of the Royal Academy of Architecture, Honorary member of that of the Marine, and fellow of the Royal Society of London, was born at Cressy en Brie, the 25th of August 1699. His early ingenuity in mechanics and his own intreaties induced his parents to send him to study at a college in Paris, at 10 years of age; where in the space of two years his progress was so great, that he was able to give lessons in mathematics, and thus to defray his own expences at the college, without any farther charge to his parents. By the assistance of the celebrated Varignon, young Camus soon ran through the course of the higher mathematics, and acquired a name among the learned. He made himself more particularly known to the Academy of Sciences in 1727, by his memoir upon the subject of the prize which they had proposed for that year, viz, 'To determine the most advantageous way of masting ships;' in consequence of which he was named that year Adjoint-Mechanician to the Academy; and in 1730 he was appointed professor of Architecture. In less than three years after, he was honoured with the secretaryship of the same; and the 18th of April 1733, he obtained the degree of Associate in the Academy, where he distinguished himself greatly by his memoirs upon living forces, or bodies in motion acted upon by forces, on the figure of the teeth of wheels and pinions, on pump work, and several other ingenious memoirs.

In 1736 he was sent, in company with Messieurs Clairaut, Maupertuis, and Monnier, upon the celebrated expedition to measure a degree at the north polar circle; in which he rendered himself highly useful, not only as a mathematician, but also as a mechanician and an artist, branches for which he had a remarkable talent.

In 1741 Camus had the honour to be appointed Pensioner-Geometrician in the Academy; and the same year he invented a gauging-rod and sliding-rule proper at once to gauge all sorts of casks, and to calculate their contents. About the year 1747 he was named Examiner of the Schools of Artillery and Engineers: and, in 1756, one of the eight mathematicians appointed to examine by a new measurement, the base which had formerly been measured by Picard, between Villejuive and Juvisy; an operation in which his ingenuity and exactness were of great utility. In 1765 M. Camus was elected a fellow of the Royal Society of London; and died the 4th of May 1768, in the 69th year of his age; being succeeded by the celebrated d'Alembert in his office of Geometrician in the French Academy; and leaving behind him a great number of manuscript treatises on various branches of the mathematics.

The works published by M. Camus, are:

1. Course of Mathematics for the use of the Engineers, 4 vols. in 8vo.

2. Elements of Mechanics.

3. Elements of Arithmetic.

And his memoirs printed in the volumes of the Academy, are:

1. Of Accelerated Motions by living forces: vol. for 1728.

2. Solution of a Geometrical Problem of M. Cramer: 1732.

3. On the figure of the teeth and pinions in clocks: 1733.

4. On the action of a Musket ball, piercing a pretty thick piece of wood, without communicating any considerable velocity to it: 1738.

5. On the best manner of employing buckets for raising water: 1739.

6. A Problem in Statics: 1740.

7. On an Instrument for gauging of vessels: 1741.

8. On the Standard of the Ell Measure: 1746.

9. On the Tangents of Points common to several branches of the same curve: 1747.

10. On the Operations in measuring the distance between the centres of the pyramids of Villejuive and Juvisy, to discover the best measure of the degree about Paris: 1754.

11. On the Masting of Ships: Prize Tom. 2.

12. The Manner of working Oars: Mach. tom. 2.

13. A Machine for moving many Colters at once: Mach. tom. 2.

CANCER, the *Crab*, one of the twelve signs of the zodiac, usually drawn on the globe in the form of a crab, and in books of Astronomy denoted by a character resembling the number sixty-nine, turned sideways, thus ☊.

This is one of the 48 old constellations; and, from the hieroglyphic mode of writing among the Egyptians &c, it is probable that they gave the name and figure to this constellation from the following circumstance, viz, that as the crab is an animal that goes sideling backwards, so the sun, in his annual course through the zodiac, when he arrives at this part of the ecliptic, having reached his utmost limit northwards, begins there to return back again towards the south. But the Greeks, who adapted some fable of their own to every thing of this kind, pretend that when Hercules was fighting with



the Lernaean hydra, there was a crab upon the marsh which seized his foot. The hero crushed the reptile to pieces under his heel; but Juno, in gratitude for the offered service, little as it was, raised the creature into the heavens.

The number of stars in the sign cancer, Ptolemy makes 13, Tycho 15, Bayer and Hevelius 29, and Flamsteed 83.

*Tropic of CANCER*, a little circle of the sphere parallel to the equinoctial, and passing through the beginning of the sign cancer.

CANDLEMAS, or the *Purification*, a feast of the church, held on the 2d of February, in memory of the purification of the Virgin; taking its name of Candlemas, either from the number of lighted candles used by the Romish church, in the processions of this day, or because that the church then consecrated candles for the whole year.

CANES *Venatici*, the *Hounds*, or the *Greyhounds*, one of the new constellations of the northern hemisphere, which Hevelius has formed out of the unformed stars of the old catalogues. These two dogs are farther distinguished by the names of *asterion* and *chara*. They contain 23 stars according to Hevelius, but 25 in the British catalogue.

CANICULA, a name given by many of the earlier astronomers to the constellation which we call the Lesser Dog, and Canis Minor, but some Procyon and Antecanis. See CANIS MINOR.

It is also used for one of the stars of the constellation Canis Major; called also simply the Dog-star; and by the Greeks Σείριος, Sirius. It is situated in the mouth of the constellation, and is the largest and brightest of all the stars in the heavens. From the heliacal rising of this star, that is, its emersion from the sun's rays, which now happens with us about the 11th of August, the ancients reckoned their *dies caniculares*, or dog-days.

The Egyptians and Ethiopians began their year at the heliacal rising of Canicula; reckoning to its rise again the next year, which is called the *Annus Canarius*.

CANICULAR *Days*, or *Dog-days*, denote a certain number of days, before and after the heliacal rising of canicula, or the dog-star, in the morning. The ancients imagined that this star, so rising, occasioned the sultry weather usually felt in the latter part of the summer, or dog-days; with all the distempers of that sickly season: Homer's *Il.* lib. 5, v. 10, and Virgil's *Æn.* lib. 10, v. 270. Some authors say, from Hippocrates and Pliny, that the day this star first rises in the morning, the sea boils, wine turns sour, dogs begin to grow mad, the bile increases and irritates, and all animals grow languid; also that the diseases it usually occasions in men, are burning fevers, dysenteries, and phrensies. The Romans too sacrificed a brown dog every year to Canicula at his first rising, to appease its rage. All this however arose from a groundless idea that the dog-star, so rising, was the occasion of the extreme heat and the diseases of that season; for the star not only varies in its rising, in any one year, as the latitude varies, but it is always later and later every year in all latitudes; so that in time the star may, by the same rule, come to be charged with bringing frost and snow, when he comes to rise in winter.

The dog-days were commonly counted for about 40 days, viz, 20 days before and 20 days after the heliacal rising; and almanac-makers have usually set down the dog-days in their almanacs to the changing time of the star's rising, by which means they had at length fallen considerably after the hottest time of the year, till of late we have observed an alteration of them in the almanacs, and very properly, from July 3 to August 11. For, by the dog-days, the ancients meant to express the hottest time of the year, which is commonly during the month of July, about which month the dog-star rose heliacally in the time of the most ancient astronomers that we know of: but the precession of the equinoxes has carried this heliacal rising into a much later and cooler part of the year; and because Hesiod tells us that the hot time of the year ends on the 50th day after the summer solstice, which brings us to about August 10 or 11, therefore the above alteration seems to be very proper.

CANICULAR *Year*, denotes the Egyptian natural year, which was computed from one heliacal rising of canicula, to the next. This year was also called *annus canarius*, and *annus cynicus*; and by the Egyptians themselves the *Sethic year*, from *Seth*, by which name they called Sirius. Some call it also the *heliacal year*. This year consisted ordinarily of 365 days, and every 4th year of 366; by which means it was accommodated to the civil year, like the Julian account. And the reason why they chose this star, in preference to others, to compute their time by, was not only the superior brightness of that star, but because that in Egypt its heliacal rising was a time of very singular note, as coinciding with the greatest augmentation of the Nile, the reputed father of Egypt. Ephesius adds, that from the aspect of canicula, its colour &c, the Egyptians drew prognostics concerning the rise of the Nile; and, according to Florus, predicted the future state of the year. So that it is no wonder the first rising of this star was observed with great attention. Bainbrigge, *Canicul.* cap. 4. p. 26.

CANIS *Major*, the *Great Dog*, a constellation of the southern hemisphere, below the feet of Orion, and one of the old 48 constellations. The Greeks, as usual, have many fables of their own about the exaltation of the dog into the skies; but the origin of this constellation, as well as its other name Sirius, lies more probably among the Egyptians, who carefully watched the rising of this star, and by it judged of the swelling of the Nile, calling the star the sentinel and watch of the year; and hence, according to their manner of hieroglyphic writing, represented it under the figure of a dog. They also called the Nile *Siris*; and hence their *Osiris*.

The stars in this constellation, Ptolemy makes 29; Tycho however observed only 13, and Hevelius 21; but in Flamsteed's catalogue they are 31.

CANIS *Minor*, a constellation of the northern hemisphere, just below Gemini, and is one of the 48 old constellations. The Greeks fabled that this is one of Orion's hounds; but the Egyptians were most probably the inventors of this constellation, and they may have given it this figure to express a little dog, or watchful creature, going before as leading in the larger, or rising before it: and hence the Latins have called it Antecanis, the star before the dog.



The stars in this constellation are, in Ptolomy's catalogue 2, the principal of which is the star Procyon; in Tycho's 5, in Hevelius's 13; and in Flamsteed's 14.

CANNON, in Military Affairs, a long round hollow engine, made of iron or brass, &c, for throwing balls, &c, by means of gunpowder. The length is distinguished into three parts; the first reinforce, the second reinforce, and the chase: the inside hollow where the charge is lodged, being also called the chase, or bore. But for the several parts and members of a cannon, see ASTRAGAL, BASE-RING, BORE, BREECH, CASCABEL, CHASE, MUZZLE, OGEE, REINFORCE-RING, TRUNNIONS, &c. See also GUN, and GUNNERY.

Cannon were first made of several bars of iron adapted to each other lengthways, and hooped together with strong iron rings. They were employed in throwing stones and metal of several hundred weight. Others were made of thin sheets of iron rolled up, and hooped: and on emergencies they have been even made of leather, with plates of iron or copper. They are now made of cast iron or brass; being cast solid, and the tube bored out of the middle of the solid metal.

Larrey makes brass cannon the invention of J. Owen; and asserts that the first known in England, were in 1535; and farther that iron cannon were first cast here in 1547. He acknowledged that cannon were known before; and remarks that at the battle of Cressi, in 1346, there were 5 pieces of cannon in the English army, which were the first ever seen in France. Mezeray also observes that king Edward struck terror into the French army, by 5 or 6 pieces of cannon; it being the first time they had met such thundering machines.

In the list of aids raised for the redemption of king John of France, in 1368, mention is made of an officer in the French army called *master of the king's cannon*, and of his providing 4 large cannon for the garrison of Harfleur. But father Daniel, in his life of Philip of Valois, produces a proof from the records of the chamber of accounts at Paris, that cannon and gunpowder were used in the year 1338. And Du-Cange even finds mention of the same engines in Froissart, and other French historians, some time earlier.

The Germans carry the invention of cannon farther back, and ascribe it to Albertus Magnus, a Dominican monk, about the year 1250. But Isaac Vossius finds cannon in China upwards of 1700 years ago; being used by the emperor Kitey, in the year of Christ 85. The ancients too, of Europe and Asia, had their fiery tubes, or *cannæ*, which being loaden with pitch, stones, and iron balls, were exploded with a vehement noise, smoke, and great effect.

Cannon were formerly made of a very great length, which rendered them exceedingly heavy, and their use very troublesome and confined. But it has lately been found by experiment that there is very little added to the force of the ball by a great length of the cannon, and therefore they have very properly been much reduced both in their length and weight, and rendered easily manageable upon all occasions. They were formerly distinguished by many hard and terrible names, but are now only named from the weight of their ball; as a 6 pounder, a 12 pounder, a 24 pounder, or a 42 pounder, which is the largest size now used by the English for battering.

CANON, in Algebra, Arithmetic, Geometry, &c,

VOL. I.

is a general rule for resolving all cases of a like nature with the present enquiry. Thus the last step of every equation is such a canon, and if turned into words, becomes a rule to resolve all cases or questions of the same kind with that proposed.

Tables of sines, tangents, &c, whether natural or artificial, are also called canons.

CANON, in Ancient Music, is a rule or way of determining the intervals of musical notes. Ptolomy, rejecting the Aristoxenian way of measuring the intervals in music by the magnitude of a tone, formed by the difference between a diapente and a diatessaron, thought that they should be distinguished by the ratios which the sounds terminating those intervals bear to one another, when considered according to their degree of acuteness or gravity; which, before Aristoxenus, was the old Pythagorean way. He therefore made the diapason consist in a double ratio; the diapente consist in a sesquialterate; the diatessaron, in a sesquitercian; and the tone itself, in a sesquioctave; and all the other intervals, according to the proportion of the sounds that terminate them: wherefore, taking the canon, as it is called, for a determinate line of any length, he shews how this is to be cut, that it may represent the respective intervals: and this method answers exactly to experiments in the different lengths of musical chords. From this canon, Ptolomy and his followers have been called *Canonici*; as those of Aristoxenus were called *Musici*.

Pascal CANON, a table of the moveable feasts, shewing the day of Easter, and the other feasts depending upon it, for a cycle or period of 19 years. It is said that the Pascal Canon was the calculation of Eusebius of Cæsarea, and that it was made by order of the council of Nice.

CANOPUS, a name given by some of the old astronomers to a star under the 2d bend of Eridanus. These writers say that the river in the heavens is not the Eridanus, but the Nile, and that this star commemorates an island made by that river, which was called by the same name.

CANOPUS is also the name of a bright star of the first magnitude in the rudder of the ship Argo, one of the southern constellations. Its situation, as given by several authors, at different times, is as follows:

Authors	Dates	Longit.	Lat.	Rt. Ascen.	Declin. So.
F. Thomas	Jan. 1682	8° 52'	75° 15'	93° 32' 20"	52° 31' 33"
J. Noel	1697			93 54	52 29
Dr. Halley	1700	10 52	72 49		
F. Feuille	Mar. 1709				52 30 4

CANTALIVERS, in Architecture, are the same with modillions, except that the former are plain, and the latter carved. They are both a kind of cartouches, set at equal distances under the corona of the cornice of a building.

CANTON (JOHN), an ingenious natural philosopher, was born at Stroud, in Gloucestershire, in 1718; and was placed, when young, under the care of Mr Davis, an able mathematician of that place, with whom he had learned both vulgar and decimal arithmetic before he was quite 9 years of age. He next proceeded to higher parts of the mathematics, and particularly to algebra and astronomy, in which he had made a considerable progress when his father took him from school, and set him to learn his own business, which was that



of a broad cloth weaver. This circumstance was not able to damp his zeal for acquiring knowledge. All his leisure time was devoted to the assiduous cultivation of astronomical science; by which he was soon able to calculate lunar eclipses and other phenomena, and to construct various kinds of sun-dials, even at times when he ought to have slept, being done without the knowledge and consent of his father, who feared that such studies might injure his health. It was during this prohibition, and at these hours, that he computed, and cut upon stone, with no better an instrument than a common knife, the lines of a large upright sun-dial, on which, beside the hour of the day, were shewn the sun's rising, his place in the ecliptic, and some other particulars. When this was finished and made known to his father, he permitted it to be placed against the front of his house, where it excited the admiration of several neighbouring gentlemen, and introduced young Canton to their acquaintance, which was followed by the offer of the use of their libraries. In the library of one of these gentlemen he found Martin's Philosophical Grammar, which was the first book that gave him a taste for natural philosophy. In the possession of another gentleman he first saw a pair of globes; a circumstance that afforded him great pleasure, from the great ease with which he could resolve those problems he had hitherto been accustomed to compute.

Among other persons with whom he became acquainted in early life, was Dr. Henry Miles of Tooting; who perceiving that young Canton possessed abilities too promising to be confined within the narrow limits of a country town, prevailed on his father to permit him to come up to London. Accordingly he arrived at the metropolis the 4th of March 1737, and resided with Dr. Miles at Tooting till the 6th of May following; when he articulated himself, for the term of 5 years, as a clerk to Mr. Samuel Watkins, master of the academy in Spital Square. In this situation his ingenuity, diligence, and prudence, were so distinguished, that on the expiration of his clerkship in May 1742, he was taken into partnership with Mr. Watkins for 3 years; which gentleman he afterward succeeded in the school, and there continued during his whole life.

Towards the end of 1745, electricity received a great improvement by the discovery of the famous Leyden phial. This event turned the thoughts of most of the philosophers of Europe to that branch of natural philosophy; and our author, who was one of the first to repeat and to pursue the experiment, found his endeavours rewarded by many notable discoveries.—Towards the end of 1749, he was engaged with his friend, the late ingenious Benjamin Robins, in making experiments to determine the height to which rockets may be made to ascend, and at what distance their light may be seen.—In 1750 was read at the Royal Society, Mr. Canton's "Method of making Artificial Magnets, without the use of, and yet far superior to, any natural ones." This paper procured him the honour of being elected a member of the Society, and the present of their gold medal. The same year he was complimented with the degree of M. A. by the university of Aberdeen. And in 1751 he was chosen one of the council of the Royal Society; an honour which was twice repeated afterwards.

In 1752, our philosopher was so fortunate as to be the first person in England who, by attracting the electric fire from the clouds during a thunder-storm, verified Dr. Franklin's hypothesis of the similarity of lightning and electricity. Next year his paper intitled "Electrical Experiments, with an attempt to account for their several phenomena," was read at the Royal Society. In the same paper Mr. Canton mentioned his having discovered, by many experiments, that some clouds were in a positive, and some in a negative state of electricity: a discovery which was also made by Dr. Franklin in America much about the same time. This circumstance, together with our author's constant defence of the doctor's hypothesis, induced that excellent philosopher, on his arrival in England, to pay Mr. Canton a visit, and gave rise to a friendship which ever after continued between them.—In the Ladies' Diary for 1756, our author answered the prize query that had been proposed in the preceding year, concerning the meteor called shooting stars. The solution, though only signed A. M. was so satisfactory to his friend, the excellent mathematician Mr. Thomas Simpson, who then conducted that ingenious and useful little work, that he sent Mr. Canton the prize, accompanied with a note, in which he said he was sure that he was not mistaken in the author of it, as no one besides, that he knew of, could have given that answer.—Our philosopher's next communication to the public, was a letter in the Gentleman's Magazine for September 1759, on the electrical properties of the tourmalin, in which the laws of that wonderful stone are laid down in a very concise and elegant manner. On the 13th of December in the same year was read at the Royal Society, "An attempt to account for the Regular Diurnal Variation of the Horizontal Magnetic Needle; and also for its Irregular Variation at the time of an Aurora Borealis." A complete year's observations of the diurnal variations of the needle are annexed to the paper.—Nov. 5, 1761, our author communicated to the Royal Society an account of the Transit of Venus of the 6th of June that year, observed in Spital Square. His next communication to the Society, was a Letter, read the 4th of Feb. 1762, containing some remarks on Mr. Delaval's electrical experiments. On the 16th of Dec. the same year, another curious addition was made by him to philosophical knowledge, in a paper, intitled, "Experiments to prove that Water is not Incompressible." And on Nov. 8, the year following, were read before the Society, his farther "Experiments and Observations on the Compressibility of Water, and some other fluids." These experiments are a complete refutation of the famous Florentine experiment, which so many philosophers have mentioned as a proof of the incompressibility of water. For this communication he had a second time the Society's prize gold medal.

Another communication was made by our author to the Society, on Dec. 22, 1768, being "An easy method of making a phosphorus that will imbibe and emit light like the Bolognian Stone; with experiments and observations." When he first shewed to Dr. Franklin the instantaneous light acquired by some of this phosphorus from the near discharge of an electrified bottle, the doctor immediately exclaimed, "And God said let there be light, and there was light."



The Dean and Chapter of St. Paul's having, in a letter, dated March 6, 1769, requested the opinion of the Royal Society relative to the best method of fixing electrical conductors to preserve that cathedral from damage by lightning, Mr. Canton was one of the committee appointed to take the letter into consideration, and to report their opinion upon it. The gentlemen joined with him in this business were, Mr. Delaval, Dr. Franklin, Dr. Watson, and Mr. Wilson. Their report was made on the 8th of June following: and the mode recommended by them has been carried into execution.—Our author's last communication to the Royal Society, was a paper read Dec. 21, 1769, containing "Experiments to prove that the Luminousness of the Sea arises from the Putrefaction of its animal Substances."

Besides the papers above mentioned, Mr. Canton wrote a number of others, both in the earlier and the later parts of his life, which appeared in several publications, and particularly in the Gentleman's Magazine.—He died of a dropsy, the 22d of March 1772, in the 51th year of his age.

CAPACITY, is the solid content of any body. Also our hollow measures for corn, beer, wine, &c, are called measures of capacity.

CAPE, or *Promontory*, is any high land, running out with a point into the sea; as Cape Verde, Cape Horn, the Cape of Good Hope, &c.

CAPELLA, a bright star of the first magnitude, in the left shoulder of Auriga.

CAPILLARY Tubes, in Physics, are very small pipes, whose canals are exceedingly narrow; being so called from their resemblance to a hair in smallness. Their usual diameter may be from  $\frac{1}{20}$  to  $\frac{1}{30}$  of an inch: though Dr. Hook assures us that he drew tubes in the flame of a lamp much smaller, and resembling a spider's thread.

The *Ascent of Water &c*, in capillary tubes, is a noted phenomenon in philosophy. Take several small glass tubes, of different diameters, and open at both ends; immerse them a little way into water, and the fluid will be seen to stand higher in the tubes than the surface of the water without, and higher as the tube is smaller, almost in the reciprocal ratio of the diameter of the tube; and that both in open air, and in vacuo. The greatest height to which Dr. Hook ever observed the water to stand, in the smallest tubes, was 21 inches above the surface in the vessel.

This does not however happen uniformly the same in all fluids; some standing higher than others; and in quicksilver the contrary takes place, as that fluid stands lower within the tube than its surface in the vessel, and the lower as the tube is smaller. See *Philos. Trans.* N° 355, or *Abr.* vol. 4, pa. 423, &c, or *Cotes's Hydr. and Pneum. Lect.* pa. 265.

Another phenomenon of these tubes is, that such of them as would only naturally discharge water by drops, when electrified, yield a continued and accelerated stream; and the acceleration is proportional to the smallness of the tube: indeed the effect of electricity is so considerable, that it produces a continued stream from a very small tube, out of which the water had not before been able to drop. *Priestley's Hist. Electr.* 8vo. vol. 1, pa. 171, ed. 3d.

This ascent and suspension of the water in the tube, is by Dr. Jurin, Mr. Hawksbee, and other philosophers, ascribed to the attraction of the periphery of the concave surface of the tube, to which the upper surface of the water is contiguous and adheres.

CAPITAL, in Architecture, the uppermost part of a column or pilaster, serving as a head or crowning to it; being placed immediately over the shaft, and under the entablature. It is made differently in the different orders, and is that indeed which chiefly distinguishes the orders themselves.

CAPITAL of a Bastion, is an imaginary line dividing any work into two equal and similar parts; or a line drawn from the angle of the polygon to the point of the bastion, or from the point of the bastion to the middle of the gorge.

CAPONIERE, or CAPONNIERE, in Fortification, is a passage made from one work to another, of 10 or 12 feet wide, and about 5 feet deep, covered on each side by a parapet, terminating in a glacis or slope. Sometimes it is covered with planks and earth.

CAPRA, or the *She-goat*, a name given to the star Capella, on the left shoulder of Auriga; and sometimes to the constellation Capricorn. Some again represent Capra as a constellation in the northern hemisphere, consisting of 3 stars, comprised between the 45th and 55th degree of latitude.

The poets fable her to be Amalthea's goat, which suckled Jupiter in his infancy.

CAPRICORN, the *Goat*, a southern constellation, and the 10th sign of the zodiac, as also one of the 48 original constellations received by the Greeks from the Egyptians. The figure of this sign is drawn as having the fore part of a goat, but the hinder part of a fish; and sometimes simply under the form of a goat. In writing, it is denoted by a character representing the crooked horns of a goat's head, thus ♊.

As to the figure of this constellation, the Greeks pretend that Pan, to avoid the terrible giant Typhon, threw himself into the Nile, and was changed into the figure here drawn; in commemoration of which exploit, Jupiter took it up to heaven. But it is probable, as Macrobius observes, that the Egyptians marked the point of the ecliptic appropriated to this sign, where the sun begins again to ascend up towards the north with the figure of a goat, an animal which is always climbing the sides of mountains.

The stars in this constellation, in Ptolemy's and Tycho's catalogue, are 28; in that of Hevelius 29: though it is to be remarked that one of those in the tail, of the 6th magnitude, marked the 27th in Tycho's book, was lost in Hevelius's time. Flamsteed gives 51 stars to this sign.

Tropic of CAPRICORN, a little circle of the sphere, parallel to the equator, passing through the beginning of Capricorn, or the winter solstice, or the point of the sun's greatest south declination.

CAPSTAN, a large massy column shaped like a truncated cone; being set upright on the deck of a ship, and turned by levers or bars, passing through holes in its upper extremity. The capstan is a kind of perpetual lever, or an axis-in-peritrochio, which, by means of a strong rope or cable passed round, serves to raise very great weights; such as to hoist sails, to weigh  
the



the anchors, to draw the vessels on shore, and hoist them up to be refitted, &c.

**CAPUT DRACONIS**, or dragon's head, a name given by some to a fixed star of the first magnitude, in the head of the constellation Draco.

**CARACT**, or **CARAT**, a name given to the weight which expresses the degree of goodness or fineness of gold. The whole quantity of metal is considered as consisting of 24 parts, which are the carats, so that the carat is the 24th part of the whole; this carat is divided into 4 equal parts, called grains of a carat, and the grain into halves and quarters.

When gold is purified to the utmost degree possible, so that it loses no more by farther trials, it is considered as quite pure, and said to be 24 carats fine; if it lose 1 carat, or 1—24th in purifying, it was of 23 carats fine; and if it lose 2 carats, it was 22 carats fine; and so on.

**CARCASS**, is a hollow case formed of ribs of iron, and covered over with pitched cloth &c, about the size of bomb-shells; or sometimes made all of iron except two or three holes for the fire to blaze through. These are filled with various matters and combustibles, to fire houses, when thrown out of mortar pieces into besieged places.

**CARCAVI** (**PETER DE**), was born at Lyons, but in what year is not known. He was Counsellor to the Parliament of Toulouse, afterward Counsellor to the Grand Council, and Keeper of the King's Library. He was appointed Geometrician to the French Academy of Sciences in 1666; and died at Paris in 1684. There are extant some letters of his, printed among those of Descartes.

**CARDAN** (**HIERONYMUS**, or **JEROM**), one of the most extraordinary geniuses of his age, was born at Pavia, in Italy, Sept. 24, 1501. At 4 years old he was carried to Milan, his father being an advocate and physician in that city: at the age of 20 he went to study in the university of the same city; and two years afterward he explained Euclid. In 1524, he went to Padua: the same year he was admitted to the degree of master of arts; and the year following, that of doctor of physic. He married about the year 1531; and became professor of mathematics, and practised medicine at Milan about 1533. In 1539 he was admitted a member of the college of physicians at Milan: in 1543 he read public lectures in medicine there; and the same at Pavia the year following; but he discontinued them because he could not get payment of his salary, and returned to Milan.

In 1552 he went into Scotland, having been sent for by the archbishop of St. Andrews, to cure him of a grievous disorder, after trying the physicians of the king of France and of the emperor of Germany, without benefit. He began to recover from the day that Cardan prescribed for him: our author took his leave of him at the end of six weeks and three days, leaving him prescriptions which in two years wrought a complete cure. Upon this visit Cardan passed through London, and calculated king Edward's nativity; for he was famous for his knowledge in astrology, as well as those of mathematics and medicine. Returning to Milan, after four months absence, he remained there till the beginning of Oct. 1552; and then went to

Pavia, from whence he was invited to Bologna in 1562. He taught in this last city till the year 1570; at which time he was thrown into prison; but some months after he was sent home to his own house. He quitted Bologna in 1571; and went to Rome, where he lived for some time without any public employment. He was however admitted a member of the college of physicians, and received a pension from the Pope, till the time of his death, which happened at Rome on the 21st of September 1575.

Cardan, at the same time that he was one of the greatest geniuses and most learned men of his age, in all the sciences, was one of the most eccentric and fickle in conduct of all men that ever lived: despising all good principles and opinions, and without one friend in the world. The same capriciousness that was remarkable in his outward conduct, is also observable in the composition of his numerous and elaborate works. In many of his treatises the reader is stopped almost every moment by the obscurity of his text, or by digressions from the point in hand. In his arithmetical writings there are several discourses on the motions of the planets, on the creation, on the Tower of Babel, and such like. And the apology which he made for these frequent digressions is, that he might by that means enlarge and fill up his book, his bargain with the bookseller being at so much per sheet; and that he worked as much for his daily support as for fame. The Lyons edition of his works, printed in 1663, consists of no less than 10 volumes in folio.

In fact, when we consider the transcendent qualities of Cardan's mind, it cannot be denied that he cultivated it with every species of knowledge, and made a greater progress in philosophy, in the medical art, in astronomy, in mathematics, and the other sciences, than the most part of his contemporaries who had applied themselves but to one only of those sciences. In particular, he was perhaps the very best algebraist of his time, a science in which he made great improvements; and his labours in cubic equations especially have rendered his name immortal, the rules for resolving them having ever since borne his name, and are likely to do so as long as the science shall exist, although he received the first knowledge of them from another person; the account of which, and his disputes with Tartalea, have been given at large under the article **ALGEBRA**.

Scaliger affirms, that Cardan, having by astrology predicted and fixed the time of his death, abstained from all food, that his prediction might be fulfilled, and that his continuance to live might not discredit his art. It is farther remarkable, that Cardan's father also died in this manner, in the year 1524, having abstained from sustenance for nine days. Our author too informs us that his father had white eyes, and could see in the night-time.

**CARDINAL Points**, in Geography, are the east, west, north, and south points of the horizon.

**CARDINAL Points of the Heavens**, or of a Nativity, are the rising and setting of the sun, the zenith and nadir.

**CARDINAL Signs**, are those at the four quarters, or the equinoxes and solstices, viz, the signs Aries, Libra, Cancer, and Capricorn.


CARDINAL



**CARDINAL Winds**, are those that blow from the four cardinal points, viz, the east, west, north, and south winds.

**CARDIOIDE**, the name of a curve so called by Castilliani.—But it was first treated of by Koersma, and by Carré. See *Philos. Trans.* 1741, and *Memoires de l'Acad.* 1705.

*The Cardioide is thus generated.* APB is a circle, and AB its diameter. Through one extremity A of the diameter draw a number of lines APQ, cutting the circle in P; upon these set off always PQ equal to the diameter AB; so shall the points Q be always in the curve of the cardioide.



From this generation of the curve, its chief properties are evident, viz, that,

everywhere  $PQ = AB$ ,

$CQ$ , or  $QQ$  is  $= Aa$  or  $2 AB$ ,

$AQ = AB \pm AP$ ,

P always bisects  $QQ$ .

The cardioide is an algebraical curve, and the equation expressing its nature is thus:

Put  $a = AB$  the diameter,

$x = aD$  perp. AB,

$y = DQ$  perp. AD; then is

$$\left. \begin{aligned} y^4 - 6ay^3 + 2x^2y^2 - 6ax^2y + x^4 \\ + 12a^2y^2 - 8a^2y + 3a^2x^2 \end{aligned} \right\} = 0,$$

which is the equation of the curve.

Many properties of the cardioide may be seen in the places above cited.

**CARRÉ (Lewis)**, was born in the year 1663, in the province of Brie in France. His father, a substantial farmer, intended him for the church. But young Carré, after going through the usual course of education for that purpose, having an utter aversion to it, he refused to enter upon that function; by which he incurred his father's displeasure. His resources being thus cut off, he was obliged to quit the university, and look out into the world for some employment. In this exigency he had the good fortune to be engaged as an amanuensis by the celebrated father Malebranche; by which he found himself transported all at once from the mazes of scholastic darkness, to the source of the most brilliant and enlightened philosophy. Under this great master he studied mathematics and the most sublime metaphysics. After seven years spent in this excellent school; M. Carré found it necessary, in order to procure himself some less precarious establishment, to teach mathematics and philosophy in Paris; but especially that philosophy which, on account of its tendency to improve our morals, he valued more than all the mathematics in the world. And accordingly his greatest care was to make geomet-

try serve as an introduction to his well-beloved metaphysics.

Most of M. Carré's pupils were of the fair sex. The first of these, who soon perceived that his language was rather the reverse of elegant and correct, told him pleasantly that, as an acknowledgement for the pains he took to teach her philosophy, she would teach him French; and he ever after owned that her lessons were of great service to him. In general he seemed to set more value upon the genius of women than that of men.

M. Carré, although he gave the preference to metaphysics, did not neglect mathematics; and while he taught both, he took care to make himself acquainted with all the new discoveries in the latter. This was all that his constant attendance on his pupils would allow him to do, till the year 1697, when M. Varignon, so remarkable for his extreme scrupulousness in the choice of his elevés, took M. Carré to him in that station. Soon after, viz. in the year 1700, our author thinking himself bound to do something that might render him worthy of that title, published the first complete work on the Integral Calculus, under the title of "A method of measuring Surfaces and Solids, and finding their Centres of Gravity, Percussion, and Oscillation." He afterwards discovered some errors in the work, and was candid enough to own and correct them in a subsequent edition.

In a little time M. Carré became Associate, and at length one of the Pensioners of the Academy. And as this was a sufficient establishment for one, who knew so well how to keep his desires within just bounds, he gave himself up entirely to study; and as he enjoyed the appointment of Mechanician, he applied himself more particularly to mechanics. He took also a survey of every branch relating to music; such as the doctrine of sounds, the description of musical instruments; though he despised the practice of music, as a mere sensual pleasure. Some sketches of his ingenuity and industry in this way may be seen in the *Memoirs of the French Academy of Sciences*. M. Carré also composed some treatises on other branches of natural philosophy, and some on mathematical subjects; all which he bequeathed to that illustrious body; though it does not appear that any of them have yet been published. It is not unlikely that he was hindered from putting the last hand to them by a train of disorders proceeding from a bad digestion, which, after harassing him during the space of five or six years, at length brought him to the grave in 1711, at 48 years of age.

His memoirs printed in the volumes of the Academy, with the years of the volumes, are as below.

1. The Rectification of Curve Lines by Tangents: 1701.

2. Solution of a problem proposed to Geometricians, &c. 1701.

3. Reflections on the Table of Equations: 1701.

4. On the Cause of the Refraction of Light: 1702.

5. Why the Tides are always augmenting from Brest to St. Malo, and diminishing along the coasts of Normandy: 1702.

6. The Number and the Names of Musical Instruments: 1702.

7. On



7. On the Vinegar which causes small stones to roll upon an inclined plane : 1703.
8. On the Rectification &c. of the Causatics by reflection : 1703.
9. Method for the Rectification of Curves : 1704.
10. Observations on the Production of Sound : 1704.
11. On a Curve formed from a Circle : 1705.
12. On the Refraction of Musket-balls in water, and on the Resistance of that fluid : 1705.
13. Experiments on Capillary Tubes : 1705.
14. On the Proportion of Pipes to have a determinate quantity of water : 1705.
15. On the Laws of Motion : 1706.
16. On the Properties of Pendulums ; with some new properties of the Parabola : 1707.
17. On the Proportion of Cylinders that their sounds may form the musical chords : 1709.
18. On the Elasticity of the Air : 1710.
19. On Catoptrics : 1710.
20. On the Monochord : in the Machines, tom. 1. with some other pieces, not mathematical.

**CARRIAGE**, of a Cannon, is the machine upon which it is mounted ; serving to point or direct it for shooting, and to convey it from place to place.

**Wheel CARRIAGE**, one that is mounted and moved about upon wheels. Horses draw in general, to most advantage, when the direction of their draft is parallel to the ground, or rather a little upwards. A carriage also goes easiest when the centre of gravity is placed very high ; since then, when once put in motion, it continues it with very little labour to the horses.

**CARTES** (**RENE DES**), one of the most eminent philosophers and mathematicians of the 17th century, or indeed of any age whatever. He was descended of an ancient noble family in Touraine in France, being a younger son of a counsellor in the parliament of Rennes, and was born March 31, 1596. His father gave him a liberal education, and the more so as he observed in him the appearance of a promising genius, using to call him the philosopher, on account of his insatiable curiosity in asking the reasons of every thing that he did not understand.

Des Cartes was sent to the Jesuits college at La Fleche in 1604, and put under the tuition of Father Charlet. Here he made a great progress in the learned languages and polite literature ; but having passed through his course of philosophy without any great satisfaction to himself, he left the college in 1612, and began to learn military arts, to ride and fence, and other such like exercises. But notwithstanding his inclination to military achievements, the weakness of his constitution not permitting him early to expose himself to the fatigues of war, he was sent to Paris in 1613. Here he formed an acquaintance with several learned persons, who helped to reclaim him from his intention of declining his studies, particularly Father Merfenne, whose conversation revived in him a love for truth, and induced him to retire from the world to pursue his studies without interruption ; which he did for two years : but in May 1616, at the repeated solicitations of his relations, he set out for Holland, and entered as a volunteer under the Prince of Orange.

Whilst he lay in garrison at Breda, during the truce between the Spanish and Dutch, an unknown person

caused a problem in mathematics, in the Dutch language, to be fixed up in the streets : when Des Cartes, seeing a concourse of people stop to read it, desired one who stood near him to explain it to him in Latin or French. The person promised to satisfy him, upon condition that he would engage to resolve the problem ; and Des Cartes agreed to the condition with such an air, that though he little expected such a thing from a young military cadet, he gave him his address, desiring he would bring him the solution. Des Cartes next day visited Beekman, principal of the college of Dort, who was the person that had translated the problem to him. Beekman was surprised at his having resolved it in such a short time ; but his wonder was much increased, to find, in the course of conversation, that the young man's knowledge was much superior to his own in those sciences, in which he had employed his whole time for several years. During his stay at Breda, Des Cartes wrote in Latin, a treatise on Music, and laid the foundation of several of his other works.

In 1619, he entered himself a volunteer in the army of the duke of Bavaria. In 1621, he made the campaign in Hungary, under the count de Bucquoy ; but the loss of his general, who was killed at a siege that year, determined him to quit the army. He soon after began his travels into the north, and visited Silesia, Poland, Pomerania, the coasts of the Baltic, Brandenburg, Holftein, East Friesland, West Friesland ; in his passage to which last place he was in danger of being murdered. The sailors fancied he was a merchant, who had a large sum of money about him ; and perceiving that he was a foreigner who had little acquaintance in the country, and a man of a mild disposition, they resolved to kill him, and throw his body into the sea. They even discoursed of their design before his face, thinking he understood no language but French, in which he always spoke to his servant. Des Cartes suddenly started up ; and drawing his sword, spoke to them in their own language, in such a tone as struck terror into them : upon which they behaved very civilly. The year following he went to Paris, where he cleared himself from the imputation of having been received among the Rosicrucians, whom he considered as a company of visionaries and impostors.

Dropping the study of mathematics, he now applied himself again to ethics and natural philosophy. The same year he took a journey through Switzerland to Italy. Upon his return he settled at Paris ; but his studies being interrupted by frequent visits, he went in 1628 to the siege of Rochelle. He returned to Paris in November ; but in the following spring he repaired to Amsterdam ; and from thence to a place near Franeker in Friesland, where he began his *Metaphysical Meditations*, and spent some time in *Dioptrics* ; about this time too he wrote his thoughts upon *Meteors*. After about six months he returned to Amsterdam.

Des Cartes imagined that nothing could more promote the temporal felicity of mankind, than the union of natural philosophy with mathematics. But before he should set himself to relieve men's labours, or multiply the conveniencies of life by mechanics, he thought it necessary to discover some means of securing the hu-  
mar



man body from disease and debility: this led him to the study of anatomy and chemistry, in which he employed the winter at Amsterdam.

He now, viz, about 1630 or 1631, took a short tour to England, and made some observations near London on the variation of the compass. In the spring of 1633 he removed to Deventer, where he completed several works that were left unfinished the year before, and resumed his studies in astronomy. In the summer he put the last hand to his "Treatise of the World." The next year he returned to Amsterdam; but soon after took a journey into Denmark, and the lower parts of Germany. In autumn 1635 he went to Lewarden in Friesland, where he remained till 1637, and wrote his "Treatise of Mechanics." The same year he published his four treatises concerning Method, Dioptries, Meteors, and Geometry. About this time he received an invitation to settle in England from Sir Charles Cavendish, brother to the earl of Newcastle, with which he did not appear backward to comply, especially upon being assured that the king was a catholic in his heart: but the breaking out of the civil wars in this country prevented his journey.

At the end of 1641, Lewis the 13th, of France, invited him to his court, upon very honourable terms; but he could not be persuaded to quit his retirement. This year he published his Meditations concerning the Existence of God, and the Immortality of the Soul. In 1645 he again applied to anatomy; but was a little diverted from this study, by the question concerning the Quadrature of the Circle, which was at that time agitated. During the winter of the same year he composed a small tract against Gassendus's Institutes; and another on the Nature of the Passions. About this time he carried on an epistolary correspondence with the princess Elizabeth, daughter to Frederick the 5th, elector palatine, and king of Bohemia, who had been his pupil in Holland.

A dispute arising between Christina, queen of Sweden, and M. Chanut, the resident of France, concerning the following question; When a man carries love or hatred to excess, which of these two irregularities is the worst? The resident sent the question to Des Cartes, who upon that occasion drew up the dissertation upon Love, that is published in the first volume of his letters, which proved highly satisfactory to the queen. In June 1647 he took a journey to France, where the king settled on him a pension of 3000 livres; but he returned to Holland about the end of September. In November he received a letter from M. Chanut, in queen Christina's name, desiring his opinion of the sovereign good; which he accordingly sent her, with some letters upon the same subject formerly written to the princess Elizabeth, and his treatise on the Passions. The queen was so highly pleased with them, that she wrote him a letter of thanks with her own hand, and invited him to come to Sweden. He arrived at Stockholm in Oct. 1648. The queen engaged him to attend her every morning at five o'clock, to instruct her in his philosophy; and desired him to revise and digest all his unpublished writings, and to draw up from them a complete body of philosophy. She purposed also to fix him in Sweden, by allowing him a revenue of 3000 crowns a year, with an estate which should

descend to his heirs and assigns for ever; and to establish an academy, of which he was to be the director. But these designs were frustrated by his death, which happened the 11th of Feb. 1650, in the 54th year of his age. His body was interred at Stockholm: but 17 years after it was removed to Paris, where a magnificent monument was erected to him in the church of Genevieve du Mont.

As to the character of our author:

Dr. Barrow in his *Opuscula* tells us, that Des Cartes was doubtless a very ingenious man, and a real philosopher, and one who seems to have brought those assistances to that part of philosophy relating to matter and motion, which perhaps no one had done before: namely, a great skill in mathematics; a mind habituated, both by nature and custom, to profound meditation; a judgment exempt from all prejudices and popular errors, and furnished with a good number of certain and select experiments; a great deal of leisure; an entire disengagement, by his own choice, from the reading of useless books, and the avocations of life; with an incomparable acuteness of wit, and an excellent talent of thinking clearly and distinctly, and of expressing his thoughts with the utmost perspicuity.

Dr. Halley, in a paper concerning Optics, communicated to Mr. Wotton, and published by the latter in his "Reflections upon Ancient and Modern Learning," writes as follows: As to dioptries, though some of the ancients mention refraction, as a natural effect of transparent media; yet Des Cartes was the first, who in this age has discovered the laws of refraction, and brought dioptries to a science.

Dr. Keil, in the introduction to his "Examination of Dr. Burnet's Theory of the Earth," tells us, that Des Cartes was so far from applying geometry and observations to natural philosophy, that his whole system is but one continued blunder on account of his negligence in that point; which he could easily prove, by shewing that his theory of the vortices, upon which his system is founded, is absolutely false, for that Newton has shewn that the periodical times of all bodies, that swim in vortices, must be directly as the squares of their distances from the centre of them: but it is evident from observations, that the planets, in moving round the sun, observe a law quite different from this; for the squares of their periodical times are always as the cubes of their distances: and therefore, since they do not observe that law, which of necessity they must if they swim in a vortex, it is a demonstration that there are no vortices in which the planets are carried round the sun.

"Nature, says Voltaire, had favoured Des Cartes with a strong and clear imagination, whence he became a very singular person, both in private life, and in his manner of reasoning. This imagination could not be concealed even in his philosophical writings, which are every where adorned with very brilliant ingenious metaphors. Nature had almost made him a poet; and indeed he wrote a piece of poetry for the entertainment of Christina queen of Sweden, which however was suppressed in honour of his memory. He extended the limits of geometry as far beyond the place where he found them, as Newton did after him; and first taught the method of expressing curves by equations. He applied



plied this geometrical and inventive genius to dioptrics, which when treated by him became a new art ; and if he was mistaken in some things, the reason is, that a man who discovers a new tract of land, cannot at once know all the properties of the soil. Those who come after him, and make these lands fruitful, are at least obliged to him for the discovery." Voltaire acknowledges, that there are innumerable errors in the rest of Des Cartes' works ; but adds, that geometry was a guide which he himself had in some measure formed, and which would have safely conducted him through the several paths of natural philosophy : nevertheless he had at last abandoned this guide, and gave entirely into the humour of framing hypotheses ; and then philosophy was no more than an ingenious romance, fit only to amuse the ignorant.

It has been pretty generally acknowledged, that he borrowed his improvements in Algebra from Harriot's *Artis Analytica Praxis* ; which is highly probable, as he was in England about the time when Harriot's book was published, and as he follows the manner of Harriot, except in the method of noting the powers. Upon this head the following anecdote is told by Dr. Pell, in Wallis's Algebra, pa. 198. Sir Charles Cavendish, then resident at Paris, discoursing there with M. Roberval, concerning Des Cartes's Geometry, then lately published : I admire, said Roberval, that method in Des Cartes, of placing all the terms of the equation on one side, making the whole equal to nothing, and how he lighted upon it. The reason why you admire it, said Sir Charles, is because you are a Frenchman ; for if you were an Englishman, you would not admire it. Why so ? asked Roberval. Because, replied Sir Charles, we in England know whence he had it ; namely from Harriot's Algebra. What book is that ? says Roberval, I never saw it. Next time you come to my chamber, saith Sir Charles, I will shew it to you. Which a while after he did ; and upon perusal of it, Roberval exclaimed with admiration, *Il l'a vu ! il l'a vu !* He had seen it ! he had seen it ! finding all that in Harriot which he had before admired in Des Cartes, and not doubting but that Des Cartes had it from thence. See also Montucla's History of Mathematics.

The real improvements of Des Cartes in Algebra and Geometry, I have particularly treated of under the article ALGEBRA ; and his philosophical doctrines are displayed in the article CARTESIAN Philosophy, here following. He was never married, but had one natural daughter, who died when she was but five years old. There have been several editions of his works, and commentaries upon them ; particularly those of Schooten on his Geometry.

CARTESIAN Philosophy, or *Cartesianism*, the system of philosophy advanced by Des Cartes, and maintained by his followers, the Cartesians.

The Cartesian philosophy is founded on two great principles, the one metaphysical, the other physical. The metaphysical one is this : *I think, therefore I am, or I exist* : the physical principle is, that *nothing exists but substances*. Substance he makes of two kinds ; the one a substance that thinks, the other a substance extended : so that actual thought and actual extension make the essence of substance.

The essence of matter being thus fixed in extension,

Des Cartes concludes that there is no vacuum, nor any possibility of it in nature ; but that the universe is absolutely full : by this principle, mere space is quite excluded ; for extension being implied in the idea of space, matter is so too.

Des Cartes defines motion to be the translation of a body from the neighbourhood of others that are in contact with it, and considered as at rest, to the neighbourhood of other bodies : by which he destroys the distinction between motion that is absolute or real, and that which is relative or apparent. He maintains that the same quantity of motion is always preserved in the universe, because God must be supposed to act in the most constant and immutable manner. And hence also he deduces his three laws of motion. See MOTION.

Upon these principles Des Cartes explains mechanically how the world was formed, and how the present phenomena of nature came to arise. He supposes that God created matter of an indefinite extension, which he separated into small square portions or masses, full of angles : that he impressed too motions on this matter ; the one, by which each part revolved about its own centre ; and another, by which an assemblage, or system of them, turned round a common centre. From whence arose as many different vortices, or eddies, as there were different masses of matter, thus moving about common centres.

The consequence of these motions in each vortex, according to Des Cartes, is as follows : The parts of matter could not thus move and revolve amongst one another, without having their angles gradually broken ; and this continual friction of parts and angles must produce three elements : the first of these, an infinitely fine dust, formed of the angles broken off ; the second, the spheres remaining, after all the angular parts are thus removed ; and those particles not yet rendered smooth and spherical, but still retaining some of their angles, and hamous parts, from the third element.

Now the first or subtlest element, according to the laws of motion, must occupy the centre of each system, or vortex, by reason of the smallness of its parts : and this is the matter which constitutes the sun, and the fixed stars above, and the fire below. The second element, made up of spheres, forms the atmosphere, and all the matter between the earth and the fixed stars ; in such sort, that the largest spheres are always next the circumference of the vortex, and the smallest next its centre. The third element, formed of the irregular particles, is the matter that composes the earth, and all terrestrial bodies, together with comets, spots in the sun, &c.

He accounts for the gravity of terrestrial bodies from the centrifugal force of the ether revolving round the earth : and upon the same general principles he pretends to explain the phenomena of the magnet, and to account for all the other operations in nature.

CARY (ROBERT), a learned English chronologer and divine, was born at Cockington, in the county of Devon, about the year 1615. He took his degrees in arts, and LL.D. in Oxford. After returning from his travels he was presented to the rectory of Portlemouth, near Kingsbridge in Devonshire : but not long after he was drawn over by the presbyterian ministers to their party, and chosen moderator of that part of the second division of the county of Devon, which was appointed



pointed to meet at Kingsbridge. And yet, upon the restoration of Charles the 2d, he was one of the first to congratulate that prince upon his return, and soon after was preferred to the archdeaconry of Exeter; but from which he was however some time afterward ejected. He spent the rest of his days at his rectory at Portlemouth, and died in 1688, at 73 years of age.—He published *Palæologia Chronica*, a chronological account of ancient time, in three parts. 1, Didactical; 2, Apodidactical; 3, Canonical: in 1677.

CASATI (PAUL), a learned Jesuit, born at Placentia in 1617. He entered early among the Jesuits; and after having taught mathematics and divinity at Rome, he was sent into Sweden to queen Christina, whom he prevailed on to embrace the popish religion. His writings are as follow:

1. *Vacuum Proscriptum*.—2. *Terra Machinis mota*.—3. *Mechanicorum, libri octo*.—4. *De Igne Dissertationes*.—5. *De Angelis Disputatio Theolog.*—6. *Hydrostaticæ Dissertationes*.—7. *Opticæ Dissertationes*. It is remarkable that he wrote this treatise on optics at 88 years of age, and after he was blind. He was also author of several books in the Italian language.

CASCABEL, the knob or button of metal behind the breech of a cannon, as a kind of handle by which to elevate and direct the piece; to which some add the fillet and ogees as far as the base-ring.

CASEMATE, or CAZEMATE, in Fortification, a kind of vault or arch, of stone-work, in that part of the flank of a bastion next the curtain; serving as a battery, to defend the face of the opposite bastion, and the moat or ditch.

The casemate sometimes consists of three platforms, one above another; the highest being on the rampart; though it is common to withdraw this within the bastion.

The casemate is also called the low place, and low flank, as being at the bottom of the wall next the ditch; and sometimes the retired flank, as being the part of the flank nearest the curtain, and the centre of the bastion. It was formerly covered by an apaulement, or a massive body, either round or square, which prevented the enemy from seeing within the batteries; whence it was also called *covered flank*.

It is now seldom used, because the batteries of the enemy are apt to bury the artillery of the casemate in the ruins of the vault: beside, the great smoke made by the discharge of the cannon, renders it intolerable to the men. So that, instead of the ancient covered casemates, later engineers have contrived open ones, only guarded by a parapet, &c.

CASEMATE is also used for a well with several subterraneous branches, dug in the passage of the bastion, till the miner is heard at work, and air given to the mine.

CASERNS, or CAZERNS, in Fortification, small rooms, or huts, erected between the ramparts and the houses of fortified towns, or even on the ramparts themselves; to serve as lodgings for the soldiers on immediate duty, to ease the garrison.

CASE-SHOT, or CANNISTER-SHOT, are a number of small balls put into a round tin cannister, and so shot out of great guns. These have superseded, and been substituted instead of the grape-shot, which have been laid aside.

CASSINI (JOHN DOMINIC), an eminent astronomer.

VOL. I.

mer, was born of noble parents, at a town in Piedmont in Italy, June 8, 1625. After laying a proper foundation in his studies at home, he was sent to continue them in a college of Jesuits at Genoa. He had an uncommon turn for Latin poetry, which he exercised so very early, that some of his poems were published when he was but 11 years old. At length he met with books of astronomy, which he read, with great eagerness. Pursuing the bent of his inclinations in this way, in a short time he made so amazing a progress, that in 1650 the senate of Bologna invited him to be their public mathematical professor. Cassini was but 25 years of age when he went to Bologna, where he taught mathematics, and made observations upon the heavens, with great care and assiduity. In 1652 a comet appeared, which he observed with great accuracy; and he discovered that comets were not bodies accidentally generated in the atmosphere, as had been supposed, but of the same nature, and probably governed by the same laws, as the planets. The same year he resolved an astronomical problem, which Kepler and Bulliald had given up as insolvable; viz, to determine geometrically the apogee and eccentricity of a planet, from its true and mean place.—In 1653, when a church in Bologna was repaired and enlarged, he obtained leave of the senate to correct and settle a meridian line, which had been drawn by an astronomer in 1575.—In 1657 he attended, as an assistant, a nobleman, who was sent to Rome to compose some differences, which had arisen between Bologna and Ferrara, from the inundations of the Po; and he shewed so much skill and judgment in the management of the affair, that in 1663 the pope's brother appointed him inspector general of the fortifications of the castle of Urbino: and he had afterward committed to him the care of all the rivers in the ecclesiastical state.

Mean while he did not neglect his astronomical studies, but cultivated them with great care. He made several discoveries relating to the planets Mars and Venus, particularly the revolution of Mars upon his own axis: but the point he had chiefly in view, was to settle an accurate theory of Jupiter's satellites; which, after much labour and observation, he happily effected, and published it at Rome, among other astronomical pieces, in 1666.

Picard, the French astronomer, getting Cassini's tables of Jupiter's satellites, found them so very exact, that he conceived the highest opinion of his skill; and from that time his fame increased so fast in France, that the government desired to have him a member of the academy. Cassini however could not leave his station without leave of his superiors; and therefore the king, Lewis the 14th, requested of the pope and the senate of Bologna, that Cassini might be permitted to come into France. Leave was granted for 6 years; and he came to Paris in the beginning of 1669, where he was immediately made the king's astronomer. When this term of 6 years was near expiring, the pope and the senate of Bologna insisted upon his return, on pain of forfeiting his revenues and emoluments, which had hitherto been remitted to him: but the minister Colbert prevailed on him to stay, and he was naturalized in 1673; the same year also in which he was married.

The Royal Observatory of Paris had been finished

K k

some



some time. The occasion of its being built was this: In 1638, the celebrated Merfenne was the chief institutor and promoter of a society, where several ingenious and learned men met together to talk upon physical and astronomical subjects; among whom were Gassend, Descartes, Monmort, Thevenot, Bulliald, our countryman Hobbes, &c: and this society was kept up by a succession of learned men for many years. At length the government considering that a number of such men, acting in a body, would succeed much better in the promotion of science, than if they acted separately, each in his particular art or province, established under the direction of Colbert, in 1666, the Royal Academy of Sciences: and for the advancement of astronomy in particular, erected the Royal Observatory at Paris, and furnished it with all kinds of instruments that were necessary to make observations. The foundation of this noble pile was laid in 1667, and the building completed in 1670. Of this observatory, Cassini was appointed to be the first inhabitant; which he took possession of in Sept. 1671, when he set himself with fresh alacrity to attend the duties of his profession. In 1672 he endeavoured to determine the parallax of Mars and the sun: and in 1677 he proved that the diurnal rotation of Jupiter round his axis was performed in 9 hours 58 minutes, from the motion of a spot in one of his larger belts: also in 1684 he discovered four satellites of Saturn, besides that which Huygens had found out. In 1693 he published a new edition of his "Tables of Jupiter's Satellites," corrected by later observations. In 1695 he took a journey to Bologna, to examine the meridian line, which he had fixed there in 1655; and he shewed, in the presence of eminent mathematicians, that it had not varied in the least, during that 40 years. In 1700 he continued the meridian line through France, which Picard had begun, to the very southern limits of that country.

After our author had resided at the royal observatory for more than 40 years, making many excellent and useful discoveries, which he published from time to time, he died September the 14th, 1712, at 87 years of age; and was succeeded by his only son James Cassini. His publications were very numerous, far too much so, even to be enumerated in this place.

CASSINI (JAMES), a celebrated French astronomer, and member of the several Academies of Sciences of France, England, Prussia, and Bologna, was born at Paris Feb. 18, 1677, being the younger son of John-Dominic Cassini, above mentioned, whom he succeeded as astronomer at the royal observatory, the elder son having lost his life at the battle of La Hogue.

After his first studies in his father's house, in which it is not to be supposed that mathematics and astronomy were neglected, he was sent to study philosophy at the Mazarine college, where the celebrated Varignon was then professor of mathematics; from whose assistance young Cassini profited so well, that at 15 years of age he supported a mathematical thesis with great honour. At the age of 17 he was admitted a member of the Academy of Sciences; and the same year he accompanied his father in his journey to Italy, where he assisted him in the verification of the meridian at Bologna, and other measurements. On his return he made other similar operations in a journey into Holland, where he

discovered some errors in the measure of the earth by Snell, the result of which was communicated to the Academy in 1702. He made also a visit to England in 1696, where he was made a member of the Royal Society.—In 1712 he succeeded his father as astronomer royal at the observatory.—In 1717 he gave to the Academy his researches on the distance of the fixed stars, in which he shewed that the whole annual orbit, of near 200 million of miles diameter, is but as a point in comparison of that distance. The same year he communicated also his discoveries concerning the inclination of the orbits of the satellites in general, and especially of those of Saturn's satellites and ring.—In 1725 he undertook to determine the cause of the moon's libration, by which she shews sometimes a little towards one side, and sometimes a little on the other, of that half which is commonly behind or hid from our view.

In 1732 an important question in astronomy exercised the ingenuity of our author. His father had determined, by his observations, that the planet Venus revolved about her axis in the space of 23 hours: and M. Bianchini had published a work in 1729, in which he settled the period of the same revolution at 24 days 8 hours. From an examination of Bianchini's observations, which were upon the spots in Venus, he discovered that he had intermitted his observations for the space of 3 hours, from which cause he had probably mistaken new spots for the old ones, and so had been led into the mistake. He soon afterwards determined the nature and quantity of the acceleration of the motion of Jupiter, at half a second per year, and of that of the retardation of Saturn at two minutes per year; that these quantities would go on increasing for 2000 years, and then would decrease again.—In 1740 he published his Astronomical Tables; and his Elements of Astronomy; very extensive and accurate works.

Although astronomy was the principal object of our author's consideration, he did not confine himself absolutely to that branch, but made occasional excursions into other fields. We owe also to him, for example, Experiments on Electricity, or the light produced by bodies by friction. Experiments on the recoil of fire arms; Researches on the rise of the mercury in the barometer at different heights above the level of the sea; Reflections on the perfecting of burning-glasses; and other memoirs.

The French Academy had properly judged that one of its most important objects, was the measurement of the earth. In 1669 Picard measured a little more than a degree of latitude to the north of Paris; but as that extent appeared too small from which to conclude the whole circumference with sufficient accuracy, it was resolved to continue that measurement on the meridian of Paris to the north and the south, through the whole extent of the country. Accordingly, in 1683, the late M. de la Hire continued that on the north side of Paris, and the older Cassini that on the south side. The latter was assisted in 1700 in the continuation of this operation by his son our author. The same work was farther continued by the same Academicians; and finally the part left unfinished by de la Hire in the north, was finished in 1718 by our author, with the late Maraldi, and de la Hire the younger.

These operations produced a considerable degree of precision.



precision. It appeared also, from this measured extent of 6 degrees, that the degrees were of different lengths in different parts of the meridian; and in such sort that our author concluded, in the volume published for 1718, that they decreased more and more towards the pole, and that therefore the figure of the earth was that of an oblong spheroid, or having its axe longer than the equatorial diameter. He also measured the perpendicular to the same meridian, and compared the measured distance with the differences of longitude as before determined by the eclipses of Jupiter's satellites; from whence he concluded that the length of the degrees of longitude was smaller than it would be on a sphere, and that therefore again the figure of the earth was an oblong spheroid; contrary to the determination of Newton by the theory of gravity. In consequence of these assertions of our author, the French government sent two different sets of measurers, the one to measure a degree at the equator, the other at the polar circle; and the comparison of the whole determined the figure to be an oblate spheroid, contrary to Cassini's determination.

After a long and laborious life, our author James Cassini lost his life by a fall in April 1756, in the 80th year of his age, and was succeeded in the Academy and Observatory by his second son Cesar-François de Thury. He published, A Treatise on the Magnitude and Figure of the Earth; as also The Elements or Theory of the Planets, with Tables; beside an infinite number of papers in the Memoirs of the Academy, from the year 1699 to 1755.

CASSINI DE THURY (CESAR-FRANÇOIS), a celebrated French astronomer, director of the observatory, pensioner astronomer, and member of most of the learned societies of Europe, was born at Paris June 17, 1714, being the second son of James Cassini, whose occupations and talents our author inherited and supported, with great honour. He received his first lessons in astronomy and mathematics from MM. Maraldi and Camus. He was hardly 10 years of age when he calculated the phases of the total eclipse of the sun of 1727. At the age of 18 he accompanied his father in his two journies undertaken for drawing the perpendicular to the observatory meridian from Strasbourg to Brest. From that time a general chart of France was devised; for which purpose it was necessary to traverse the country by several lines parallel and perpendicular to the meridian of Paris, and our author was charged with the conduct of this business. He did not content himself with the measure of a degree by Picard: suspecting even that the measures which had been taken by his father and grandfather were not exempt from some errors, which the imperfections of their instruments at least would be liable to, he again undertook to measure the meridian of Paris, by means of a new series of triangles, of a smaller number, and more advantageously disposed. This great work was published in 1740, with a chart shewing the new meridian of Paris, by two different series of triangles, passing along the sea coasts, to Bayonne, traversing the frontiers of Spain to the Mediterranean and Antibes, and thence along the eastern limits of France to Dunkirk, with parallel and perpendicular lines described at the distance of 6000 toises from one another, from side to side of the country.—In

1735, he had been received into the academy as adjoint supernumerary at 21 years of age.

A tour which our author made in Flanders, in company with the king, about 1741, gave rise to the particular chart of France, at the instance of the king. Cassini published different works relative to these charts, and a great number of the sheets of the charts themselves.

In 1761, Cassini undertook an expedition into Germany; for the purpose of continuing to Vienna the perpendicular of the Paris meridian; to unite the triangles of the chart of France with the points taken in Germany; to prepare the means of extending into this country the same plan as in France; and thus to establish successively for all Europe a most useful uniformity. Our author was at Vienna the 6th of June 1761, the day of the transit of the planet Venus over the sun, of which he observed as much as the state of the weather would permit him to do, and published the account of it in his *Voyage en Allemagne*.

Finally, M. Cassini, always meditating the perfection of his grand design, profited of the late peace to propose the joining of certain points taken upon the English coast with those which had been determined on the coast of France, and thus to connect the general chart of the latter with that of the British isles, like as he had before united it with those of Flanders and Germany. The proposal was favourably received by the English government, and presently carried into effect, under the direction of the Royal Society, the execution being committed to the late General Roy; after whose death the business was for some time suspended; but it has lately been revived under the auspices of the duke of Richmond, Master General of the Ordnance, and the execution committed to the care of Col. Edward Williams and Capt. William Mudge, both respectable officers of the Artillery, and Mr. Isaac Dalby, who had before accompanied and assisted General Roy; from whose united skill and zeal the happiest prosecution of this business may be expected.

M. Cassini published in the volumes of Memoirs of the French Academy a prodigious number of pieces, chiefly astronomical, too numerous to particularize in this place, between the years 1735 and 1770; consisting of astronomical observations and questions; among which are observable, Researches concerning the Parallax of the sun, the moon, Mars, and Venus; On astronomical refractions, and the effect caused in their quantity and laws by the weather; Numerous observations on the obliquity of the ecliptic, and on the law of its variations. In short, he cultivated astronomy for 50 years, of the most important for that science that ever elapsed, for the magnitude and variety of objects, in which he commonly sustained a principal share.

M. Cassini was of a very strong and vigorous constitution, which carried him through the many laborious operations in geography and astronomy which he conducted. An habitual retention of urine however rendered the last 12 years of his life very painful and distressing, till it was at length terminated by the small-pox the 4th of September 1784, in the 71st year of his age; being succeeded in the academy, and as director of the observatory, by his only son the present count John-Dominic Cassini; who is the 4th in order by direct descent in that honourable station.



**CASSIOPEIA**, one of the 48 old constellations, placed near Cepheus, not far from the north pole. The Greeks probably received this figure, as they did the rest, from the Egyptians, and in their fables added it to the family in the neighbouring part of the heavens, making her the wife of Cepheus, and mother of Andromeda. They pretend she was placed in this situation, to behold the destruction of her favourite daughter Andromeda, who is chained just by her on the shore, to be devoured; and that as a punishment for her pride and vanity in presuming to stand the comparison of beauty with the Nereids.

In the year 1572 there burst out all at once in this constellation a new star, which at first surpassed Jupiter himself in magnitude and brightness; but it diminished by degrees, till it quite disappeared at the end of 18 months. This star alarmed all the astronomers of that age, many of whom wrote dissertations upon it; among the rest Tycho Brahe, Kepler, Maurolycus, Lycetus, Gramineus, and others. Beza, the Landgrave of Hesse, Rosa, and others, wrote to prove it a comet, and the same that appeared to the Magi at the birth of Christ, and that it came to declare his second coming: these were answered by Tycho.

The stars in the constellation Cassiopeia, are in Ptolemy's catalogue 13, in Hevelius's 37, in Tycho's 46, and in Flamsteed's 55.

**CASTOR**, a moiety of the constellation Gemini; called also Apollo. Also a star in this constellation, whose latitude, for the year 1700, according to Hevelius, was  $10^{\circ} 4' 20''$  north; and its longitude  $16^{\circ} 4' 14''$ .

**CASTOR and POLLUX**. See GEMINI.

**CASTOR and POLLUX**, in Meteorology, is a fiery meteor, which at sea appears sometimes adhering to a part of the ship, in the form of a ball, or even several balls. When one is seen alone, it is properly called Helena; but two are called Castor and Pollux, and sometimes Tyndaridæ.

By the Spaniards, Castor and Pollux are called San Elmo; by the French, St. Elme, St. Nicholas, St. Clare, St. Helene; by the Italians, Hermo; and by the Dutch, Vree Vuuren.

The meteor Castor and Pollux, it is commonly thought, denotes a cessation of the storm, and a future calm; as it is rarely seen till the tempest is nigh spent. But Helena alone portends ill weather, and denotes the severest part of the storm yet behind.


When the meteor adheres to the masts, yards, &c, it is concluded, from the air not having motion enough to dissipate this flame, that a profound calm is at hand; but if it flutter about, that it denotes a storm.

**CASTRAMETATION**, the art, or act, of encamping an army.

**CATACAUSTICS**, or *Catacaustic Curves*, in the Higher Geometry, are the species of caustic curves formed by reflection.


These curves are generated after the following manner: If there be an infinite number of rays  $AB$ ,  $AC$ ,  $AD$ , &c, proceeding from the radiating point  $A$ , and reflected at any given curve  $BCDH$ , so that the angles of incidence be still equal to the angles of reflection; then the curve  $BEG$ , to which the reflect-

ed rays  $BI$ ,  $CE$ ,  $DF$ , &c, are always tangents, as at the points  $I$ ,  $E$ ,  $F$ , &c, is the catacaustic, or caustic-by-reflection. - Or it is the same thing as to say, that a caustic curve is that formed by joining the points of concurrence of the several reflected rays.



Some properties of these curves are as follow. If the reflected ray  $IB$  be produced to  $K$ , so that  $AB = BK$ , and the curve  $KL$  be the evolute of the caustic  $BEG$ , beginning at the point  $K$ ; then the portion of the caustic  $BE$  is  $= AC - AB + CE - BI$ , that is, the difference of the two incident rays added to the difference of the two reflected rays.

When the given curve  $BCD$  is a geometrical one, the caustic will be so too, and will always be rectifiable. The caustic of the circle, is a cycloid, or epicycloid, formed by the revolution of a circle upon a circle.



Thus,  $ABD$  being a semicircle exposed to parallel rays; then those rays which fall near the axis  $CB$  will be reflected to  $F$ , the middle point of  $BC$ ; and those which fall at  $A$ , as they touch the curve only, will not be reflected at all; but any intermediate ray  $HI$  will be reflected to a point  $K$ , somewhere between  $A$  and  $F$ . And since every different incident ray will have a different focal point, therefore those various focal points will form a curve line  $AEF$  in one quadrant, and  $FGD$  in the other, being the cycloid above-mentioned. And this figure may be beautifully exhibited experimentally by exposing the inside of a smooth bowl, or glass, to the sun beams, or strong candle light; for then this curve  $AEFGD$  will appear plainly delineated on any white surface placed horizontally within the same, or on the surface of milk contained in the bowl.

The caustic of the common cycloid, when the rays are parallel to its axis, is also a common cycloid, described by the revolution of a circle upon the same base. The caustic of the logarithmic spiral, is the same curve.

The principal writers on the caustics, are l'Hôpital, Carré, &c. See *Memoires de l'Acad. an.* 1666 & 1703.

CATA-



**CATACOUSTICS**, or *Cataphonics*, is the science of reflected sounds; or that part of acoustics which treats of the properties of echoes.

**CATADIOPTICAL Telescope**, the same as Reflecting telescope; which see.

**CATALOGUE of the Stars**, is a list of the fixed stars, disposed according to some order; in their several constellations; with the longitudes, latitudes, right-ascensions, &c, of each.

Catalogues of the stars have usually been disposed, either as collected into certain figures called constellations, or according to their right ascensions, that is the order of their passing over the meridian. All the catalogues, from the most ancient down to Flamsteed's inclusively, were of the first of these forms, or in constellations: but most of the others since that have been of the latter form, as being much more convenient for most purposes. Indeed one has lately been disposed in classes according to zones or degrees of polar distance.

Hipparchus of Rhodes first undertook to make a catalogue of the stars, about 128 years before Christ; in which he made use of the observations of Timocharis and Aristyllus, for about 140 years before him. Ptolomy retained Hipparchus's catalogue, containing 1026 fixed stars in 48 constellations, though he himself made abundance of observations, with a view to a new catalogue, an. dom. 140. Albategni, a Syrian, brought the same down to his own time, viz, about the year of Christ 880. Anno 1437, Ulugh Beigh, or Beg, king of Parthia and India, made a new catalogue of 1022 fixed stars, or according to some 1016; since translated out of Persian into Latin by Dr. Hyde, in 1665. The third person who made a catalogue of stars from his own observations was Tycho Brahe, who determined the places of 777 stars for the end of the year 1600; which Kepler, from other observations of Tycho, afterwards increased to the number of 1000 in the Rudolphine tables; adding those of Ptolomy and other authors, omitted by Tycho; so that his catalogue amounts to above 1160. About the same time, William, landgrave of Hesse, with his mathematicians Byrgius and Rothman, determined the places of 400 stars from new observations, rectifying them for the year 1593; which Hevelius prefers to those of Tycho. Ricciolus, in his *Astronomia Reformata*, determined the places of 101 stars for the year 1700, from his own observations: for the rest he followed Tycho's catalogue; altering it where he thought fit. Anno 1667 Dr. Halley, in the island of St. Helena, observed 350 of the southern stars, not visible in our horizon. The same labour was repeated by Father Noel in 1710, who published a new catalogue of the same stars constructed for the year 1687. Also De la Caille, at the Cape of Good Hope, made accurate observations of about 10 thousand stars near the south pole, in the years 1751 and 1752; the catalogue of which was published in the *Memoirs of the French Academy of Sciences* for the year 1752, and in some of his own works, as more particularly noticed below.

Bayer, in his *Uranometria*, published in 1603 a catalogue of 1160 stars, compiled chiefly from Ptolomy and Tycho, in which every star is marked with some letter of the Greek alphabet; the brightest or princi-

pal star in any constellation being denoted by the first letter of the alphabet, the next star in order by the 2d letter, and so on; and when the number of stars exceeds the Greek alphabet, the remaining stars are marked by the letters of the Roman alphabet; which letters are preserved by Flamsteed in his catalogue, and by Senex on his globes, and indeed by most astronomers since that time.

In 1673, the celebrated John Hevelius, of Dantzick, published, in his *Machina Caelestis*, a catalogue of 1888 stars, of which 1553 were observed by himself; and their places set down for the end of the year 1660. But this catalogue, as it stands in Flamsteed's *Historia Caelestis* of 1725, contains only 1520 stars.

The most complete catalogue ever given from the labours of one man, was the Britannic catalogue, compiled from the observations of the accurate and indefatigable Mr. Flamsteed, the first Royal Astronomer at Greenwich; who for a long series of years devoted himself wholly to that business. As there was nothing wanting either in the observer or apparatus, his may be considered as a perfect work, so far as it goes. It is however to be regretted that the edition had not passed through his own hands: that now extant was published by authority, but without the author's consent, and contains 2734 stars. Another edition was published in 1725, pursuant to his testament, containing 3000 stars, with their places adapted to the beginning of the year 1689; to which is added Mr. Sharp's catalogue of the southern stars not visible in our hemisphere, set down for the year 1726. See vol. 3 of his *Historia Caelestis*, in which are printed the catalogues of Ptolomy, Ulugh Beigh, Tycho, the Prince of Hesse, and Hevelius; with an account of each of them in the Prolegomena.

The first catalogue we believe that was printed in the new or second form, according to the order of the right ascensions, is that of De la Caille, given in his *Ephemerides* for the 10 years between 1755 and 1765, and printed in 1755. It contains the right ascensions and declinations of 307 stars, adapted to the beginning of the year 1750.—In 1757, De la Caille published his *Astronomiae Fundamenta*, containing a catalogue of the right ascensions and declinations of 398 stars, likewise adapted to the beginning of 1750.—And in 1763, the year after his death, was published the *Calum Australe Stelliferum* of the same author; containing a catalogue of the places of 1942 stars, all situated to the southward of the tropic of Capricorn, and observed by him while he was at the Cape of Good Hope, in 1751 and 1752; their places being also adapted to the beginning of 1750.—In the same year was published his *Ephemerides* for the 10 years between 1765 and 1775; in the introduction to which are given the places of 515 zodiacal stars, all deduced from the observations of the same author; the places adapted to the beginning of the year 1765.

In the *Nautical Almanac* for 1773, is given a catalogue of 387 stars, in right ascension, declination, longitude, and latitude, derived from the observations of the late celebrated Dr. Bradley, and adjusted to the beginning of the year 1760. This small catalogue, and the results of about 1200 observations of the moon, are all that the public have yet seen of the multiplied labours of



of this most accurate and indefatigable observer, although he has now (1794) been dead upwards of 32 years.

In 1775 was published a thin volume entitled *Opera Inedita*, containing several papers of the late Tobias Mayer, and among them a catalogue of the right ascensions and declinations of 998 stars, which may be occulted by the moon and planets; the places being adapted to the beginning of the year 1756.

At the end of the first volume of 'Astronomical Observations made at the Royal Observatory at Greenwich,' published in 1776, Dr. Maskelyne, the present Astronomer Royal, has given a catalogue of the places of 34 principal stars, in right-ascension and north-polar distance, adapted to the beginning of the year 1770. These, being the result of several years' repeated observations, made with the utmost care, and the best instruments, it may be presumed are exceedingly accurate.

In 1782, M. Bode, of Berlin, published a very extensive catalogue of 5058 of the fixed stars, collected from the observations of Flamsteed, Bradley, Hevelius, Mayer, De la Caille, Messier, Monnier, D'Arquier, and other astronomers; all adapted to the beginning of the year 1780; and accompanied with a Celestial Atlas, or set of maps of the constellations, engraved in a most delicate and beautiful manner.

To these may be added, Dr. Herschel's catalogue of double stars, printed in the *Philos. Transf.* for 1782 and 1783; Messier's nebulae and clusters of stars, published in the *Connoissance des Temps* for 1784; and Herschel's catalogue of the same kind, given in the *Philos. Transf.* for 1786.

In 1789 Mr. Francis Wollaston published 'A Specimen of a General Astronomical Catalogue, in Zones of North-polar Distance, and adapted to Jan. 1, 1790.' These stars are collected from all the catalogues before-mentioned, from that of Hevelius downwards. This work contains five distinct catalogues; viz, Dr. Maskelyne's new catalogue of 36 principal stars; a general catalogue of all the stars, in zones of north-polar distance; an index to the general catalogue; a catalogue of all the stars, in the order in which they pass the meridian; and a catalogue of zodiacal stars, in longitude and latitude.

Finally, in 1792, Dr. Zach published at Gotha, *Tabula Motuum Solis*, to which is annexed a new catalogue of the principal fixed stars, from his own observations made in the years 1787, 1788, 1789, 1790. This catalogue contains the right ascensions and declinations of 381 principal stars, adapted to the beginning of the year 1800.

**CATAPULT**, *Catapulta*, a military engine, much used by the ancients for throwing huge stones, and sometimes large darts and javelins, 12, 15, or even 18 feet long, on the enemy. It is sometimes confounded with the Ballista, which is more peculiarly adapted for throwing stones; some authors making them the same, and others different.


The catapult, which it is said was invented by the Syrians, consisted of two huge timbers, like masts of ships, placed against each other, and bent by an engine for the purpose; these being suddenly unbent again by the stroke of a hammer, threw the javelins with prodigious force. Its structure and the manner of

working it are described by Vitruvius; and a figure of it is also given by Perrault. M. Folard asserts that the catapult made infinitely more disorder in the ranks than our cannon charged with iron balls. See Vitruv. Archit. lib. x. cap. 15 and 18; and Perr. notes on the same; also Rivius, pa. 597.

Josephus takes notice of the surprising effects of these engines, and says, that the stones thrown out of them, of a hundred weight or more, beat down the battlements, knocked the angles off the towers, and would level a whole file of men from one end to the other, were the phalanx ever so deep.

See plate V, fig. 3 and 4, for two forms of the catapult, the one for throwing darts and javelins, the other for stones.

**CATENARY**, a curve line which a chain, cord, or such like, forms itself into, by hanging freely from two points of suspension, whether these be in the same horizontal line or not; as the curve ACB, formed by a heavy flexible line suspended by any two points A and B.



The nature of this curve was sought after by Galileo, who thought it was the same with the parabola; but though Jungius detected this mistake, its true nature was not discovered till the year 1691, in consequence of M. John Bernoulli having published it as a problem in the *Acta Eruditorum*, to the mathematicians of Europe. In 1697 Dr. David Gregory published an investigation of the properties before discovered by Bernoulli and Leibnitz; in which he pretends that an inverted catenary is the best figure for an arch of a bridge &c. See *Philos. Transf. abr.* vol. 1. pa. 39; also Bernoulli *Opera*, vol. 1. pa. 48, and vol. 3. pa. 491; and Cotes's *Harmon. Mensur.* pa. 108.

The catenary is a curve of the mechanical kind, and cannot be expressed by a finite algebraical equation, in simple terms of its absciss and ordinate; but is easily expressed by means of fluxions; thus if AQ be its axis perpendicular to the horizon, and PQ an ordinate parallel to the same, or perp. to AQ; also pq another ordinate indefinitely near the former, and po parallel to AQ; then, a being some given or constant quantity, the fundamental property of the curve is this, viz,  $Po : op :: AP : a$ , or  $x : y :: z : a$ , that is, the fluxion of the axis, is to the fluxion of the ordinate, as the length of the curve is to the given quantity a; where  $x = AQ$ ,  $y = PQ$ , and  $z = AP$ . This, and the other properties of the curve, will easily appear from the following considerations: First, supposing the curve hung up by its two points B and C against a perpendicular or upright wall: then, every lower part of the curve being kept in its position by the tension of that which is immediately above it, the lower parts of the curve will retain the same position unvaried, by whatever points it is suspended



pended above; thus, if it were fixed to the wall by the point F, or G, &c, the whole curve CAB will remain just as it was; for the tensions at F and G have the same effect upon the other parts of the curve as when it is fixed by those points: and hence it follows that the tension of the curve at the point A, in the horizontal direction, is a constant quantity, whether the two legs or branches of the curve, on both sides of it, be longer or shorter: which constant tension at A let be denoted by the quantity  $a$ .

Now because any portion of the curve, as AP, is kept in its position by three forces, viz, the tensions at its extremities A and P, and its own weight, of which the tension at A acts in the direction AH or  $pa$ , and the tension at P acts in the direction Pp, and the wt. of the line acts in the perpendicular direction oP; that is, the three forces which retain the curve AP in its position, act in the directions of the sides of the elementary triangle opP; but, by the principles of mechanics, any three forces, keeping a body in equilibrio, are proportional to the three sides of a triangle drawn in the directions in which those forces act; therefore it follows that the forces keeping AP in its position, viz, the tension at A, the tension at P, and the wt. of AP, are respectively as op, pP, and oP, that is, as  $y$ ,  $z$ , and  $x$ .

But the tension at A is the constant quantity  $a$ , and the wt. of the uniform curve AP may be expounded by its length  $z$ ; therefore it follows that  $x : y :: z : a$ ; which was to be proved.

Also from this last proportion, by proper analogy, or similar combinations of the terms, there arises this other property,  $x : \sqrt{ax + yy}$  or  $z :: z : \sqrt{aa + zz}$ , or  $z\dot{z} = \dot{x}\sqrt{aa + zz}$ , or  $\dot{x} = \frac{z\dot{z}}{\sqrt{a^2 + z^2}}$ ; and the fluents of these give  $x = \sqrt{a^2 + z^2}$ . But at the vertex of the curve, where  $x = 0$ , and  $z = 0$ , this becomes  $0 = \sqrt{a^2 + 0} = a$ ; and therefore by correction the true equation of the fluents is  $x = \sqrt{a^2 + z^2} - a$ , or  $a + x = \sqrt{a^2 + z^2}$ : and hence also  $z = \sqrt{a + x^2} - a^2 = \sqrt{2ax + x^2}$  and  $a = \frac{z^2 - x^2}{2x}$ .

Any of which is the equation of the curve in terms of the arch and its absciss; in which it appears that  $a + x$  is the hypotenuse of a right-angled triangle whose two legs are  $a$  and  $z$ . So that, if in QA and HA produced, there be taken AD =  $a$ , and AE = the curve  $z$  or AP; then will the hypotenuse DE be =  $a + x$  or DQ. And hence, any two of these three,  $a$ ,  $x$ ,  $z$ , being given, the third is given also.

Again, from the first simple property, viz,  $x : y :: z : a$ , or  $ax = zy$ , by substituting the value of  $z$  above found, it becomes  $ax = y\sqrt{2ax + x^2}$ , or  $y = \frac{ax}{\sqrt{2ax + x^2}}$ ; and the fluent of this equation is

$y = 2a \times \text{hyp. log. of } \sqrt{x} + \sqrt{2a + x}$ . But at the vertex of the curve, where  $x = 0$  and  $y = 0$ , this becomes  $0 = 2a \times \text{hyp. log. of } \sqrt{2a}$ ; therefore the correct equation of the fluents is  $y = 2a \times \text{hyp. log. of}$

$\frac{\sqrt{x} + \sqrt{2a + x}}{\sqrt{2a}}$ ; an equation to the curve also, in

terms of  $x$  and  $y$ , but not in simple algebraic terms. This last equation however may be brought to much simpler terms in different ways; as first by squaring the logarithmic quantity and dividing its coef. by 2, then

$$y = a \times \text{hyp. log. of } \frac{a + x + \sqrt{2ax + x^2}}{a} = a \times$$

$\text{hyp. log. } \frac{a + x + z}{a}$ ; and 2d by multiplying both nu-

merator and denominator by  $\sqrt{2a + x} - \sqrt{x}$ , then squaring the product, and dividing the coef. by 2,

which gives  $y = a \times \text{hyp. log. } \frac{z + x}{\sqrt{z^2 - x^2}} = a \times$

$$\text{hyp. log. } \frac{z + x^2}{z^2 - x^2} = a \times \text{hyp. log. } \frac{z + x}{z - x}.$$

**CATHETUS**, in Geometry, a name by which the perpendicular leg of a right-angled triangle is sometimes called. Or it is in general any line or radius falling perpendicularly on another line, or surface.

**CATHETUS of Incidence**; in Catoptrics, is a right line drawn from a radiant point, or point of incidence, perpendicular to the reflecting line, or plane of the speculum.

**CATHETUS of Reflection**, or of the Eye, a right line drawn from the eye, perpendicular to the plane of reflection.

**CATHETUS**, in Architecture, denotes the axis of a column &c. In the Ionic Capital, it denotes a line passing perpendicularly through the eye or centre of the volute.


**CATOPTRICS**, the science of reflex vision; or that part of optics which explains the laws and properties of light reflected from mirrors, or specula.

The first treatise extant on catoptrics, is that which was composed by Euclid: this was published in Latin in 1604 by John Pena; it is also contained in Herigon's Course of Mathematics, and in Gregory's edition of the works of Euclid: though it is suspected by some that this piece was not the work of that great geometer, notwithstanding that it is ascribed to him by Proclus in lib. 2, and by Marinus in his Preface to Euclid's Data. Alhazen, an Arabian author, composed a large volume of optics about the year 1100, in which he treats pretty fully of catoptrics: and after him Vitello, a Polish writer, composed another about the year 1270. Tacquet, in his Optics, has very well demonstrated the chief propositions of plane and spherical speculums. And the same is very ably done by Dr. Barrow in his Optical Lectures. There are also Trabe's Catoptrics; David Gregory's Elements of Catoptrics; Wolfius's Elements of Catoptrics; and those of Dr. Smith, contained in his learned and very elaborate Treatise on Optics; and many others of less note.

As this subject is treated under the general term Optics, the less need be said of it here. The whole doctrine of Catoptrics depends upon this simple principle, that the angle of incidence is equal to the angle of reflection, that is, that the angle in which a ray of light falls upon any surface, called the angle of incidence, is




is equal to the angle in which it quits it when reflected from it, called the angle of reflection; though it is sometimes defined that the angles of incidence and reflection, are those which the incident and reflected rays make, not with the reflecting surface itself, but with a perpendicular to that surface, at the point of contact, which are the complements to the others: but it matters not by what name these angles are called, as to the truth and principles of the science; since, if the angles are equal, their complements are also equal. This principle of the equality between the angles of incidence and reflection, is mere matter of experience, being a phenomenon that has always been observed to take place, in every case that has fallen under observation, as near at least as mechanical measurements can ascertain; and hence it is inferred that it is a universal law of nature, and to be considered as matter of fact in all cases. Thus, let AC be an incident ray falling upon the reflecting surface DE, and CB the reflected ray, also CO perpendicular to DE; then is the angle  $\angle ACD = \angle BCE$ , or the angle  $\angle ACO = \angle BCO$ .



Of this law in nature, viz, the equality between the angles of incidence and reflection, it is remarkable, that in this way, the length or rout  $AC + CB$ , in a ray passing from any point A to another given point B, by being reflected from any surface DE, is the shortest possible, namely  $AC + CB$  is shorter than the sum  $AG + GB$  of any other two lines inflected at the line DE; and hence also the passage of the ray from A to B is performed in a shorter time than if it had passed by any other way.

From this simple principle, and the common properties of lineal geometry, the chief phenomena of catoptrics are easily deduced, and are as here following.



1. Rays of light reflected from a plane surface, have the same inclination to each other after reflection as they had before it. Thus, the rays AC, AI, AK, issuing from the radiant point A, and reflected by the surface DE into the lines CF, IL, KM; these latter lines will have the same inclination to each other as the former AC, AI, AK have. For draw ABG perpendicular to DE, and produce FC, LI, MK backwards to meet this perpendicular, so shall they all meet in the same point G, and AB will in every case be equal to BD: for the incident  $\angle ACB$  is equal to the re-


flected  $\angle FCE$ , which is equal to the opposite angle BCG; so that the two triangles ABC, GBC, have the angles at C equal, as also the right angles at B equal, and consequently the 3d angles at A and G equal; and having also the side BC common, they are equal in all respects, and so  $AB = BG$ . And the same for the other rays. Consequently the angles BGC, BGI, BGK are respectively equal to the angles BAC, BAI, BAK; that is, the reflected rays have the same inclinations as the incident ones have.

2. Hence it is that the image of an object, seen by reflection from a plain mirror, seems to proceed from a place G as far beyond, or on the other side of the reflecting plane DE, as the object A itself is before the plane. This is when the incident rays diverge from some point as A.

But if the case be reversed, and FC, LI, MK be considered as incident rays, issuing from points F, L, M, and converging to some point G beyond the reflecting plane; then CA, IA, KA will become the reflected rays, and they will converge to the point A as far before the plane, as the point G is beyond it.

So that universally, when the incident rays diverge from a point A, the reflected rays will also diverge from a point G; and when the incident rays converge towards a point G, the reflected ones will also converge to a point A; and in both cases these two points are at equal distances on the opposite sides of the reflecting plane DE.

3. Parallel rays reflected from a concave spherical surface, converge after reflection. For, let AF, CD, EB be three parallel rays falling upon the concave surface FB, whose centre is C. To the centre draw the perpendiculars FC and BC; also draw FM making the reflected angle CFM equal to the incident angle CFA; and in like manner BM to make the angle CBM = the angle CBE; so shall the rays AF and EB be reflected into the converging rays FM and BM. As to the ray CD, being perpendicular to the surface, it is reflected back again in the same line DC.



4. Converging rays falling upon the concave surface are made to converge more. Thus, let GB and HF be the incident rays: then because the incident angle HFC is larger than the angle AFC, therefore the equal reflected




reflected angle  $NFC$  is greater than the reflected angle  $MFC$ , and so the point  $N$  is below the point  $M$ , or the line  $FN$  below the line  $FM$ ; and in like manner  $BN$  is below  $BM$ ; that is, the reflected rays  $FN$  and  $BN$  are more converging in this case, than  $FM$  and  $BM$  in the other.

5. The focus to which all parallel rays, falling near the vertex  $D$ , are reflected, is in the middle of the radius  $M$ . For, because the  $\angle MFC = \angle AFC$  which is = the alternate  $\angle FCM$ , therefore the sides opposite these angles are also equal, namely the side  $FM = CM$ ; consequently when the point  $F$  is very near the vertex  $D$ , then the sum  $CM + MF$  is nearly =  $CD$ , and so  $CM$  nearly =  $MD$ , or the focus of the parallel rays is nearly in the middle of the radius.—But the focus of other reflected rays is either above or below that of the parallel rays; namely, below when the incident rays are converging, and above when they are diverging; as is evident by inspection; thus,  $N$  the reflected focus of the converging rays  $GB$  and  $HF$ , is below  $M$ ;  $I$  that of the diverging rays  $YB$  and  $YF$ , is above  $M$ .

6. Incident and reflected rays are reciprocal, or so that if the reflected rays be returned back, or considered as incident ones, they will be reflected back into what were before their incident rays. And hence it follows that diverging rays, after reflection from a concave spherical surface, become either parallel or less diverging than before. Thus the incident rays  $MF$  and  $MB$  are reflected into the parallel rays  $FA$  and  $BE$ , and the rays  $NF$  and  $NB$  are reflected into  $FH$  and  $BG$ , which are less diverging; also the rays  $IF$  and  $IB$  are reflected into  $FK$  and  $BL$ , which converge.—And hence all the phenomena of concave mirrors will be evident.

7. Rays reflected from a convex speculum, become quite contrary to those reflected from a concave one; so that the parallel rays become diverging, and the diverging rays become still more diverging; also converging rays will become either diverging, or parallel, or else less converging. Thus  $BDF$  being a spherical surface, whose centre is  $C$ , produce the radii  $CBV$  and  $CFT$  which are perpendicular to the surface; then it is evident that the parallel rays  $AF$  and  $EB$  will be reflected into the diverging ones  $FK$  and  $BL$ ; and the diverging rays  $YB$  and  $YF$  become  $BO$  and  $FP$  which are more diverging; also the converging rays  $HF$  and  $GB$  become  $FR$  and  $BS$  which diverge, or else  $KF$  and  $LB$  become  $FA$  and  $BE$  which are parallel, or else lastly  $PF$  and  $OB$  become  $FY$  and  $BY$  which are converging.



8. Hence, as in the concave speculum, so also in the convex one, of parallel incident rays  $AF$  and  $EB$ , the imaginary focus  $M$  of their reflected rays  $FL$  and  $BK$ , is in the middle of the radius when the speculum is a small segment of a sphere: but the reflected imaginary focus of other rays is either above or below the middle point  $M$ , viz  $N$  being that of the converging rays  $GB$  and  $HF$ , below  $M$ ; but  $I$ , that of the diverging rays  $YB$  and  $YF$ , above  $M$ .

9. When the speculum is the small segment of a sphere, either convex or concave, and the incident rays either converging or diverging, the distances of the foci, or points of concurrence, of the incident rays, and of the reflected rays, from the vertex of the speculum, are directly proportional to the distances of the same from the centre of it;

that is  $YD : ID :: YC : IC$ ,

and  $QD : ND :: QC : NC$ .

For because the radius  $CF$ , or the same produced, bisects the angle  $YFI$  in the concave speculum, or the external angle  $YFP$  in the convex one, therefore  $YF : IF :: YC : IC$ ; but when  $F$  is very near to  $D$ , then  $YF$  and  $IF$  become nearly  $YD$  and  $ID$ ; consequently  $YD : ID :: YC : IC$ .

In like manner, because  $CF$  bisects the angle  $QFN$  in the convex, or its external angle  $NFH$  in the concave speculum, therefore  $QF : FN :: QC : NC$ ; but when  $F$  is very near to  $D$ , then  $QF$  and  $FN$  become nearly  $QD$  and  $ND$ ; and therefore  $QD : ND :: QC : NC$ .

For example, suppose it were required to find the focal distance of diverging rays incident upon a convex surface, the radius of the sphere being 5 inches, and the distance of the radiant point from the surface 20 inches. Here then are given  $YD = 20$ , and  $CD = 5$ , to find  $ID$ : then

the theorem  $YD : ID :: YC : IC$ ,

in numbers is  $20 : ID :: 25 : 5 - ID$ ,

or by permutation  $20 : 25 :: ID : 5 - ID$ ,

and by composition  $45 : 20 :: 5 : ID = \frac{100}{45} = \frac{20}{9} = 2\frac{2}{9}$  the focal distance sought.

And if it should happen in any case that the value of  $ID$  in the calculation should come out a negative quantity, the focal distance must then be taken on the contrary side of the surface.

From the foregoing principles may be deduced and collected the following practical maxims, for plane and spherical mirrors, viz,

#### I. In a Plane Mirror,

- (1). The image will appear as far behind the mirror, as the object is before it.
- (2). The image will appear of the same size, and in the same position as the object.
- (3). Any plain mirror will reflect the image of an object of twice its own length and breadth.

#### II. In a Spherical Convex Mirror,

- (1). The image will always appear behind the mirror, or within the sphere.
- (2). The image will be in the same position, but less than the object.
- (3). The image will be curved, but not spherical, like the mirror.
- (4). Parallel rays falling on this mirror, will have the



the image at half the distance of the centre from the mirror.

(5). In converging rays, the distance of the object must be equal to half the distance of the centre, to make the image appear behind the mirror.

(6). Diverging rays will have their image at less than half the distance of the centre.

### III. In a Spherical Concave Mirror,

(1). Parallel rays have their focus, or the image, at half the distance of the centre.

(2). In the centre of the sphere the image appears of the same dimensions as the object.

(3). Converging rays form an image before the mirror.

(4). In diverging rays, if the object be at less than half the distance of the centre, the image will be behind the mirror, erect, curved, and magnified; but if the distance of the object be greater, the image will be before the mirror, inverted and diminished.

(5). The solar rays, being parallel, will be collected in a focus at half the distance of its centre, where their heat will be augmented in proportion as the surface of the mirror exceeds that of the focal spot.

(6). If a luminous body be placed in the focus of a concave mirror, its rays, being reflected in parallel lines, will strongly enlighten a space of the same dimensions with the mirror, at a great distance. If the luminous object be placed nearer than the focus, its rays will diverge, and so enlighten a larger space, but not so strongly. And upon this principle it is that reverberators are constructed.



**CATOPTRIC Dial**, a dial that exhibits objects by reflected rays. See *Reflecting DIAL*.

**CATOPTRIC Telescope**, a telescope that exhibits objects by reflection. See *Reflecting TELESCOPE*.

**CATOPTRIC Cistula**, a machine, or apparatus, by which small bodies are represented extremely large, and near ones extremely wide, and diffused through a vast space; with other very pleasing phenomena, by means of mirrors, disposed by the laws of catoptrics, in the concavity of a kind of chest.

There are various kinds of these machines, accommodated to the various intentions of the artificer: some multiply the objects, some magnify, some deform them, &c. The structure of one or two of them will suffice to shew how many more may be made.

*To make a Catoptric Cistula to represent several different scenes of objects, when viewed by different holes.*



Provide a polygonal cistula, or box, like the multangular prism ABCDEF, and divide its cavity by diagonal planes AD, BE, CF, intersecting in the centre, into as many triangular cells as the chest has sides. Line those diagonal partitions with plain mirrors; and in the sides of the box make round holes, through which the eye may peep within the cells of it. These holes are to be covered with plain glasses, ground within-side, but not polished, to prevent the objects in the cells from appearing too distinctly. In each cell are to be placed the different objects whose images are to be exhibited; then covering up the top of the chest with a thin transparent membrane, or parchment, to admit the light, the machine is complete.

For, from the laws of reflection, it follows, that the images of objects, placed within the angles of mirrors, are multiplied, and appear some more remote than others; by which the objects in one cell will appear to take up more room than is contained in the whole box. Therefore by looking through one hole only, the objects in one cell will be seen, but those multiplied, and diffused through a space much larger than the whole box. Thus every hole will afford a new scene; and according to the different angles the mirrors make with each other, the representations will be different: if they be at an angle greater than a right one, the images will be monstrous, &c.

*To make a Catoptric Cistula to represent the objects within it prodigiously multiplied, and diffused through a vast space.*

Make a polygonous cistula or box, as before, but without dividing the inner cavity into any apartments, or cells; line the insides CBHI, BHIA, ALMF, &c, with plane mirrors, and at the holes pare off the tin and quicksilver, to look through; place any object in the bottom MI, as a bird in a cage, &c.

Now by looking through the aperture *hi*, each object placed at the bottom will be seen vastly multiplied, and the images removed at equal distances from one another, like a great multitude of birds, or a large aviary.

**CAVALIER**, in Fortification, a mount of earth raised in a fortress higher than the other works, on which to place cannon &c for scouring the field, or opposing a commanding work. Cavaliers are of different shapes; and are bordered with a parapet, to cover the cannon mounted upon them; their situation is also various, either in the curtain, bastion, or gorge. The cavalier is sometimes called a double bastion, and its use is to overlook the enemy's batteries, and to scour their trenches.

**CAVALIERI (BONAVENTURA)**, an eminent Italian mathematician in the 17th century. He was a native of Milan, and a friar of the order of the Jesuati of St. Jerome. Cavalieri was a disciple of Galileo; and the friend of Torricelli. He was a very eminent mathematician, and was professor of that science at Bologna; where several of his books were published, and where he died in the year 1647. His works that have been published, as far as I can find, are as follow:

1. *Directorium Generale Uranometricum*; 4to, Bononia, 1632.—In this work the author treats of Trigonometry; and Logarithms, their construction, uses, and applications. The work includes also tables of logarithms.



of common numbers; with trigonometrical tables, of natural sines, and logarithmic sines, tangents, secants and versed sines.

2. *Lo Specchio Ustorio overo Trattato delle Settoni Coniche*: 4to, Bologna, 1632.—An ingenious treatise of conic sections.

3. *Geometria Indivisibilibus continuorum nova quadam ratione promota*: 4to, Bononia, 1635; and a 2d edition in 1653.—This is a curious original work in geometry, in which the author conceives the geometrical figures as resolved into their very small elements, or as made up of an infinite number of infinitely small parts, and on account of which he passes in Italy for the inventor of the infinitesimal calculus.

4. *Trigonometria Plana & Sphærica, Linearis, & Logarithmica*: 4to, Bononia, 1643.—A very neat and ingenious treatise on Trigonometry; with the tables of sines, tangents, and secants, both natural and logarithmical.

5. *Exercitationes Geometricæ Sex*: 4to, Bononia, 1647. This work contains Exercises on the method of Indivisibles; Answers to the objections of Guldini; The use of Indivisibles in cubic powers or algebra, and in considerations about gravity; with a miscellaneous collection of problems.

CAUDA *Capricorni*, a fixed star of the 4th magnitude, in the tail of Capricorn; called also by the Arabs, Dineb Algedi; and marked  $\gamma$  by Bayer.

CAUDA *Ceti*, a fixed star of the 3d magnitude; called also by the Arabs, Dineb Kactos; marked  $\beta$  by Bayer.

CAUDA *Cygni*, a fixed star of the 2d magnitude in the Swan's tail; called by the Arabs, Dineb Adigege, or Eldegiagich; and marked  $\alpha$  by Bayer.

CAUDA *Delphini*, a fixed star of the 3d magnitude, in the tail of the Dolphin; marked  $\epsilon$  by Bayer.

CAUDA *Draconis*, or Dragon's tail, the moon's southern or descending node.

CAUDA *Leonis*, a fixed star of the first magnitude in the Lion's tail; called also by the Arabs, Dineb Eleced; and marked  $\beta$  by Bayer. It is called also Lucida Cauda.

CAUDA *Ursæ Majoris*, a fixed star of the 3d magnitude, in the tip of the Great Bear's tail; called also by the Arabs, Alalioth, and Benenath; and marked  $\eta$  by Bayer.

CAUDA *Ursæ Minoris*, a fixed star of the 3d magnitude, at the end of the Lesser Bear's tail; called also the Pole Star, and by the Arabs, Alrukabah; and marked  $\alpha$  by Bayer.

CAVETTO, a hollow member or moulding, containing a quadrant of a circle, and having an effect just contrary to that of a quarter round. It is used as an ornament in cornices.

CAUSTIC CURVES. See *Catacaustics*, and *Diacausics*.

CAZEMATE. See CASEMATE.

CAZERN. See CASERN.

CEGINUS, a fixed star of the 3d magnitude, in the left shoulder of Bootes; and marked  $\gamma$  by Bayer.

CELERITY, is the velocity or swiftness of a body in motion; or that affection of a body in motion by which it can pass over a certain space in a certain time.

CELESTIAL GLOBE, &c. See GLOBE, &c.

CELLARIUS (CHRISTOPHER), a learned geographer and historiographer of the 17th century. He was born in 1638, at Smalcalde in Franconia, where his father was minister. Our author was successively rector of the colleges at Weymar, Zeitz, and Mersebourg, and professor of eloquence and history in the university founded by the king of Prussia at Hall in 1693, where he composed the greatest part of his works.

His great application to study hastened the infirmities of old age; for it has been said, he would spend whole days and nights together over his books, without any attention to his health, or even the calls of nature. He died in 1707, at 69 years of age.

Cellar was author of an amazing number of books, upon various subjects; but those on account of which he has a place here, are his geographical works, which are as follow:

1. *Notitia Orbis Antiqui*, 2 vols in 4to; and is esteemed the best work extant on the ancient geography.

2. *Atlas Cælestis*; in folio.

3. *Historia Antiqua*, 2 vols in 12mo; being an abridgement of universal history.

CENTAURUS, the *Centaur*, one of the 48 old constellations, being a southern one, and is in form half man and half horse. It is fabled by the Greeks that it was Chiron the Centaur, who was the tutor of Achilles and Esculapius. The stars of this constellation are, in Ptolemy's catalogue 37, in Tycho's 4, and in the Britannic catalogue, with Sharp's appendix, 35.

CENTER. See CENTRE.

CENTESM, the 100th part of any thing.

CENTRAL, something relating to a centre. Thus we say central eclipse, central forces, central rule, &c.

CENTRAL *Eclipse*, is when the centres of the luminaries exactly coincide, and come in a line with the eye.

CENTRAL *Forces*, are forces having a tendency directly towards or from some point or centre; or forces which cause a moving body to tend towards, or recede from, the centre of motion. And accordingly they are divided into two kinds, in respect to their different relations to the centre, and hence are called centripetal, and centrifugal.

The doctrine of central forces makes a considerable branch of the Newtonian philosophy, and has been greatly cultivated by mathematicians, on account of its extensive use in the theory of gravity, and other physico-mathematical sciences.

In this doctrine, it is supposed that matter is equally indifferent to motion or rest; or that a body at rest never moves itself, and that a body in motion never of itself changes either the velocity or the direction of its motion; but that every motion would continue uniformly, and its direction rectilinear, unless some external force or resistance should affect it, or act upon it. Hence, when a body at rest always tends to move, or when the velocity of any rectilinear motion is continually accelerated or retarded, or when the direction of a motion is continually changed, and a curve line is thereby described, it is supposed that these circumstances proceed from the influence of some power that acts incessantly; which power may be measured, in the first case, by the pressure of the quiescent body against the obstacle which prevents it from moving, or by the ve-




locity gained or lost in the second case, or by the flexure of the curve described in the 3d case: due regard being had to the time in which these effects are produced, and other circumstances, according to the principles of mechanics. Now the power or force of gravity produces effects of each of these kinds, which fall under our constant observation near the surface of the earth; for the same power which renders bodies heavy, while they are at rest, accelerates their motion when they descend perpendicularly; and bends the track of the motion into a curve line, when they are projected in a direction oblique to that of their gravity. But we can judge of the forces or powers that act on the celestial bodies by effects of the last kind only. And hence it is, that the doctrine of central forces is of so much use in the theory of the planetary motions.

Sir I. Newton has treated of central forces in lib. 1 sec. 2 of his Principia, and has demonstrated this fundamental theorem of central forces, viz, that the areas which revolving bodies describe by radii drawn to an immoveable centre, lie in the same immoveable planes, and are proportional to the times in which they are described. Prop. 1.

It is remarked by a late eminent mathematician, that this law, which was originally observed by Kepler, is the only general principle in the doctrine of centripetal forces; but since this law, as Newton himself has proved, cannot hold in cases where a body has a tendency to any other than one and the same point, there seems to be wanting some law that may serve to explain the motions of the moon and satellites which gravitate towards two different centres: the law he lays down for this purpose is, That when a body is urged by two forces tending constantly to two fixed points, it will describe, by lines drawn from the two fixed points, equal solids in equal times, about the line joining those fixed points. See Maclin, on the Laws of the Moon's Motion, in the Postscript. See also a demonstration of this law by Mr. William Jones, in the Philos. Transf. vol. 59. Very learned tracts have also been since given, when the motion respects, not two only, but several centres, by many ingenious authors, and practical rules deduced from them for computing the places &c of planets and satellites; as by La Grange, De la Place, Waring, &c, &c. See Berlin Memoirs; those of the Academy of Sciences at Paris; and the Philos. Transf. of London.

M. De Moivre gave elegant general theorems relating to central forces, in the Philos. Transf. and in his Miscel. Analyt. pa. 231.—Let MPQ be any given curve, in which a body moves: let P be the place of the body at any time; S the centre of force, or the point to which the central force acting on the body is always directed; PG the radius of curvature at the point P; and ST perpendicular to the tangent PT; then will the centripetal force be everywhere propor-

tional to the quantity  $\frac{SP}{GP \times ST^3}$ . Vid. ut supra.



M. Varignon has also given two general theorems on this subject in the Memoirs of the Acad. an. 1700, 1701; and has shewn their application to the motions of the planets. See also the same Memoirs, an. 1706, 1710.

Mr. MacLaurin has also treated the subject of central forces very ably and fully, in his Treatise on Fluxions, art. 416 to 493; where he gives a great variety of expressions for these forces, and several elegant methods of investigating them.


#### Laws of CENTRAL FORCES.

1. The following is a very clear and comprehensive rule, for which we are obliged to the marquis de l'Hôpital: Suppose a body of any determinate weight to revolve uniformly about a centre, with any given velocity; find from what height it must have fallen, by the force of gravity, to acquire that velocity; then, as the radius of the circle it describes is to double that height, so is its weight to its centrifugal force. So that, if  $b$  be the body, or its weight or quantity of matter,  $v$  its velocity, and  $r$  the radius of the circle described, also  $g = 16\frac{1}{2}$  feet; then, first  $4g^2 : v^2 :: g : \frac{v^2}{4g}$  the height due to the velocity  $v$ ; and as

$r : \frac{v^2}{2g} :: b : \frac{v^2 b}{2gr} = f$  the centrifugal force. And hence, if the centrifugal force be equal to the gravity, the velocity is equal to that acquired by falling through half the radius.

2. The central force of a body moving in the periphery of a circle, is as the versed sine AM of the indefinitely small arc AE; or it is as the square of that arc AE directly, and as the diameter AB inversely. For AM is the space through which the body is drawn from the tangent in the given time, and  $2AM$  is the proper measure of the central force. But, AE being very small, and therefore nearly equal to its chord, by the nature of the circle

$$AB : AE :: AE : AM = \frac{AE^2}{AB}$$



3. If two bodies revolve uniformly in different circles; their central forces are in the duplicate ratio of their velocities directly, and the diameters or radii of the circles inversely;

$$\text{that is } F : f :: \frac{V^2}{D} : \frac{v^2}{d} :: \frac{V^2}{R} : \frac{v^2}{r}$$

For the force, by the last article, is as



$\frac{AE^2}{AB}$  or  $\frac{AE^2}{D}$ ; and the velocity  $v$  is as the space  $AE$  uniformly described.

4. And hence, if the radii or diameters be reciprocally in the duplicate ratio of the velocities, the central forces will be reciprocally in the duplicate ratio of the radii, or directly as the 4th power of the velocities; that is, if  $V^2 : v^2 :: r : R$ , then  $F : f :: r^2 : R^2 :: V^4 : v^4$ .

5. The central forces are as the diameters of the circles directly, and squares of the periodic times inversely. For if  $c$  be the circumference described in the time  $t$ , with the velocity  $v$ ; then the space  $c = tv$ , or  $v = \frac{c}{t}$ ; hence, using this value of  $v$  in the 3d rule, it becomes

$F : f :: \frac{C^2}{DF^2} : \frac{c^2}{dt^2} :: \frac{D}{T^2} : \frac{d}{t^2} :: \frac{R}{T^2} : \frac{r}{t^2}$ ; since the diameter is as the circumference.

6. If two bodies, revolving in different circles, be acted on by the same central force; the periodic times are in the subduplicate ratio of the diameters or radii of the circles; for when  $F = f$ , then  $\frac{D}{T^2} = \frac{d}{t^2}$ , and  $D : d :: T^2 : t^2$ , or  $T : t :: \sqrt{D} : \sqrt{d} :: \sqrt{R} : \sqrt{r}$ .

7. If the velocities be reciprocally as the distances from the centre, the central forces will be reciprocally as the cubes of the same distances, or directly as the cubes of the velocities. That is, if  $V : v :: r : R$ , then is  $F : f :: r^3 : R^3 :: V^3 : v^3$ .

8. If the velocities be reciprocally in the subduplicate ratio of the central distances, the squares of the times will be as the cubes of the distances: for if  $V^2 : v^2 :: r : R$ , then is  $T^2 : t^2 :: R^3 : r^3$ .

9. Wherefore, if the forces be reciprocally as the squares of the central distances, the squares of the periodic times will be as the cubes of the distances; or when  $F : f :: r^2 : R^2$ , then is  $T^2 : t^2 :: R^3 : r^3$ .

*Exam.* From this, and some of the foregoing theorems, may be deduced the velocity and periodic time of a body revolving in a circle, at any given distance from the earth's centre, by means of its own gravity. Put  $g = 16\frac{1}{2}$  feet, the space described by gravity, at the surface, in the first second of time, viz =  $AM$  in the foregoing fig. and by rule 2; then, putting  $r =$  the radius  $AC$ ; it is  $AE = \sqrt{AB \times AM} = \sqrt{2gr}$  the velocity in a circle at its surface, in one second of time; and hence, putting  $c = 3.14159$  &c, the circumference of the earth being  $2\pi r = 25,000$  miles, or  $132,000,000$

feet, it will be  $\sqrt{2gr} : 2\pi r :: 1'' : c\sqrt{\frac{2r}{g}} = 5078$  seconds nearly, or  $1^h 24^m 38^s$ , the periodic time at the circumference: Also the velocity there, or  $\sqrt{2gr}$  is  $= 26000$  feet per second nearly. Then, since the force of gravity varies in the inverse duplicate ratio of the distance, by rules 8 and 9, it is  $\sqrt{R} : \sqrt{r} :: v$  or  $26000 : 26000\sqrt{\frac{r}{R}} = V$  the velocity of a body revolving about the earth at the distance  $R$ ; and  $\sqrt{r^3} : \sqrt{R^3} :: t$  or  $5078'' : 5078\sqrt{\frac{R^3}{r^3}} = T$  the time of revolution in the same. So if, for instance, it be the moon

revolving about the earth at the distance of 60 semidiameters; then  $R = 60r$ , and the above expressions become  $V = 26000\sqrt{\frac{r}{60r}} = 3357$  feet per second, or  $38\frac{1}{7}$  miles per minute, for the velocity of the moon in her orbit; and  $T = 5078\sqrt{\frac{R^3}{r^3}} = 2360051$  seconds or  $27\frac{3}{5}$  days nearly, for the periodic time of the moon in her orbit at that distance.

Thus also the ratio of the forces of gravitation of the moon towards the sun and earth may be estimated. For, 1 year or  $365\frac{1}{4}$  days being the periodic time of the earth and moon about the sun, and  $27\frac{3}{5}$  days the periodic time of the moon about the earth, also 60 being the distance of the moon from the earth, and 23920 the distance from the sun, in semidiameters of the earth, by art. 5 it is

$\frac{60}{27.3^2} : \frac{23920}{365.25^2} :: f \text{ or } 1 : \frac{23920}{60} \times \frac{27.3^2}{365.25^2} = 2\frac{2}{9}$  that is, the proportion of the moon's gravitation towards the sun, is to that towards the earth, as  $2\frac{2}{9}$  to 1 nearly.

Again, we may hence compute the centrifugal force of a body at the equator, arising from the earth's rotation. For, the periodic time when the centrifugal force is equal to the force of gravity, it has been shewn above, is 5078 seconds, and 23 hours, 56 minutes, or 86160 seconds, is the period of the earth's rotation on its axis; therefore, by art. 5, as  $86160^2 : 5078^2 :: 1 : \frac{1}{289}$ , the centrifugal force required, which therefore is the 289th part of gravity at the earth's surface. Simpson's Flux. pa. 240, &c.

Also for another example, suppose  $A$  to be a ball of 1 ounce, which is whirled about the centre  $C$ , so as to describe the circle  $ABE$ , each revolution being made in half a second; and the length of the cord  $AC$  equal to 2 feet. Here then  $t = \frac{1}{2}$ ,  $r = 2$ ; and it having been

found above that  $c\sqrt{\frac{2R}{g}} = T$  is the periodic time at the circumference of the earth when the centrifugal force is equal to gravity; hence then, by art. 5, as

$\frac{R}{T^2} : \frac{r}{t^2} :: F \text{ or } 1 : f$ , which proportion becomes


$\frac{g}{2c^2} : \frac{r}{t^2} :: 1 : \frac{2c^2r}{gt^2} = \frac{16c^2}{g} = \frac{16 \times 3.1416^2}{16 \times \frac{1}{2}} = 9.819$  = the centrifugal force, or that by which the string is stretched, viz, nearly 10 ounces, or 10 times the weight of the ball.

Lastly, suppose the string and ball be suspended from a point  $D$ , and describes in its motion a conical surface  $ADB$ ; then putting  $DC = a$ ,  $AC = r$ , and  $AD = b$ ; and putting  $F = 1$  the force of gravity as before; then will the body  $A$  be affected by three forces, viz, gravity acting parallel to  $DC$ , a centrifugal force in the direction  $CA$ , and the tension of the string, or force by which it is stretched, in the direction  $DA$ ; hence these three powers will be as the three sides of the triangle  $ADC$  respectively, and therefore as  $CD$  or  $a : AD$  or  $b :: 1 : \frac{b}{a}$  the tension of the string as compared with the weight

of the body. Also  $CD$  or  $a : AC$  or  $r :: 1 : \frac{2c^2r}{gt^2}$  the general expression for the centrifugal force above found; hence



hence,  $gt^2 = 2ac^2$  and so  $t = c\sqrt{\frac{2a}{g}} = 1.108\sqrt{a} =$  the periodic time. And



10. When the force by which a body is urged towards a point is not always the same, but is either increased or decreased as some power of the distance; several curves will thence arise according to that power. If the force decrease as the squares of the distances increase, the body will describe an ellipse, and the force is directed towards one of its foci; so that in every revolution the body once approaches towards it, and once recedes from it: also the eccentricity of the ellipse is greater or less, according to the projectile force; and the curve may sometimes become a circle, when the eccentricity is nothing; the body may also describe the other two conic sections, the parabola and hyperbola, which do not return into themselves, by supposing the velocity greater in certain proportions. Also if the force increase in the simple ratio as the distance increases, the body will still describe an ellipse; but the force will in this case be directed to the centre of the ellipse; and the body, in each revolution, will twice approach towards it, and again twice recede from that point.

**CENTRAL RULE**, is a rule or method discovered by Mr. Thomas Baker, rector of Nympton in Devonshire, which he published in his *Geometrical Key*, in the year 1684, for determining the centre of a circle which shall cut a given parabola in as many points as a given equation, to be constructed, has real roots; which he has applied with good success in the construction of all equations as far as the 4th power inclusive.

The *Central Rule* is chiefly founded on this property of the parabola; that if a line be inscribed in the curve perpendicular to any diameter, the rectangle of the segments of this line, is equal to the rectangle of the intercepted part of the diameter and the parameter of the axis.

The *Central Rule* has the advantage over the methods of constructing equations by DesCartes and DeLatteres, which are liable to the trouble of preparing the equations by taking away the second term; whereas Baker's method effects the same thing without any previous preparation whatever. See also *Philos. Transf.* N° 157.

**CENTRE**, or **CENTER**, in a general sense, signifies a point equally remote from the extremes of a line, plane, or solid; or a middle point dividing them so that some certain effects are equal on all sides of it.

**CENTRE of Attraction**, or *Gravitation*, is the point to which bodies tend by gravity; or that point to which

a revolving planet or comet is impelled or attracted, by the force or impetus of gravity.

**CENTRE of a Bastion**, is a point in the middle of the gorge, where the capital line commences, and which is usually at the angle of the inner polygon of the figure. Or it is the point where the two adjacent curtains produced intersect each other.

**CENTRE of a Circle**, is the point in the middle of a circle, or circular figure, from which all lines drawn to the circumference are equal.

**CENTRE of a Conic Section**, is the middle point of any diameter, or the point in which all the diameters intersect and bisect one another.

In the ellipse the centre is within the figure; but in the hyperbola it is without, or between the conjugate hyperbolas; and in the parabola it is at an infinite distance from the vertex.

**CENTRE of Conversion**, in Mechanics, a term first used by M. Parent, and may be thus conceived: Suppose a stick laid on stagnant water, and then drawn by a thread fastened to it, so that the thread always makes the same angle with the stick, either a right angle or any other; then it will be found that the stick will turn about one point of it, which will be immovable; and this point is termed the centre of conversion.

This effect arises from the resistance of the fluid to the stick partly immersed in it. And if, instead of the body thus floating on a fluid, the same be conceived to be laid on the surface of another body; then the resistance of this plane to the stick will always have the same effect, and will determine the same centre of conversion. And this resistance is precisely what is called friction, so prejudicial to the effects of machines.

M. Parent has determined this centre in some certain cases, with much laborious calculation. When the thread is fastened to the extremity of the stick, he found that the distance of the centre from this extremity would be nearly  $\frac{1}{2}\frac{2}{3}$  of the whole length. But when it is a surface or a solid, there will be some change in the place of this centre, according to the nature of the figure. See *Mem. of the Acad. of Sciences*, vol. 1, pa. 191.

**CENTRE of a Curve**, of the higher kind, is the point where two diameters meet.—When all the diameters meet in the same point, it is called, by Sir Isaac Newton, the *general centre*.

**CENTRE of a Dial**, is the point where its gnomon or stile, which is placed parallel to the axis of the earth, meets the plane of the dial; and from hence all the hour-lines are drawn, in such dials as have centres, viz, all except that whose plane is parallel to the axis of the world; all the hour-lines of which are parallel to the stile, and to one another, the centre being as it were at an infinite distance.

**CENTRE of an Ellipse**, is the middle of any diameter, or the point where all the diameters intersect.

**CENTRE of the Equant**, in the Old Astronomy, is a point in the line of the aphelion; being as far distant from the centre of the eccentric, towards the aphelion, as the sun is from the same centre of the eccentric towards the perihelion.

**CENTRE of Equilibrium**, is the same with respect to bodies immersed in a fluid, as the centre of gravity is to bodies in free space; being a certain point, upon which



which if the body or bodies be suspended, they will rest in any position. To determine this centre, see Emmer-son's Mechanics, prop. 92, pa. 134.

**CENTRE of Friction**, is that point in the base of a body on which it revolves, into which if the whole surface of the base, and the mass of the body were collected, and made to revolve about the centre of the base of the given body, the angular velocity destroyed by its friction would be equal to the angular velocity destroyed in the given body by its friction in the same time.—See Vince on the Motion of Bodies affected by friction, in the Philos. Transf. 1785.

**CENTRE of Gravity**, is that point about which all the parts of a body do in any situation exactly balance each other. Hence, by means of this property, if the body be supported or suspended by this point, the body will rest in any position into which it is put; as also that if a plane pass through the same point, the segments on each side will equiponderate, neither of them being able to move the other.

The whole gravity, or the whole matter, of a body may be conceived united in the centre of gravity; and in demonstrations it is usual to conceive all the matter as really collected in that point.

Through the centre of gravity passes a right line, called the *diameter of gravity*; and therefore the intersection of two such diameters determines the centre. Also the plane upon which the centre of gravity is placed, is called the *plane of gravity*; so that the common intersection of two such planes determines the diameter of gravity.

In homogeneous bodies, which may be divided lengthways into similar and equal parts, the centre of gravity is the same with the centre of magnitude. Hence therefore the centre of gravity of a line is in the middle point of it, or that point which bisects the line. Also the centre of gravity of a parallelogram, or cylinder, or any prism whatever, is in the middle point of the axis. And the centre of gravity of a circle or any regular figure, is the same as the centre of magnitude.

Also, if a line can be so drawn as to divide a plane into equal and similar parts, that line will be a diameter of gravity, or will pass through the centre of gravity; and it is the same as the axis of the plane. Thus the line drawn from the vertex and perpendicular to the base of the isosceles triangle, is a diameter of gravity; and thus also the axis of an ellipse, or a parabola, &c, is a diameter of gravity. The centre of gravity of a segment or arc of a circle, is in the radius or line perpendicularly bisecting its chord or base.

Likewise, if a plane divide a solid in the same manner, making the parts on both sides of it perfectly equal and similar in all respects, it will be a plane of gravity, or will pass through the centre of gravity. Thus, as the intersection of two such planes determines the diameter of gravity, the centre of gravity of a right cone, or spherical segment, or conoid, &c, will be in the axis of the same.

**Common Centre of Gravity** of two or more bodies, or the different parts of the same body, is such a point as that, if it be suspended or supported, the system of bodies will equiponderate, and rest in any position. Thus, the point of suspension in a common balance beam, or steelyard, is the centre of gravity of the same.

### Laws and Determination of the Centre of Gravity.

1. In two equal bodies, or masses, the centre of gravity is equally distant from their two respective centres. For these are as two equal weights suspended at equal distances from the point of suspension; in which case they will equiponderate, and rest in any position.



2. If the centres of gravity of two bodies A and B be connected by the right line AB, the distances AC and BC from the common centre of gravity C, are reciprocally as the weights or bodies A and B; that is,  $AC : BC :: B : A$ .

See this demonstrated under the article BALANCE.

Hence, if the weights of the bodies A and B be equal, their common centre of gravity C will be in the middle of the right line AB, as in the foregoing article. Also since  $A : B :: BC : AC$ , therefore  $A \times AC = B \times BC$ ; whence it appears that the powers of equiponderating bodies are to be estimated by the product of the mass multiplied by the distance from the centre of gravity; which product is usually called the *momentum* of the weights.

Further, from the foregoing proportion, by composition it will be  $A + B : A :: AB : BC$ , or  $A + B : B :: AB : AC$ . So that the common centre of gravity C of two bodies will be found, if the product of one weight by the whole distance between the two, be divided by the sum of the two weights. Suppose, for example, that  $A = 12$  pounds,  $B = 4$  lb, and  $AB = 36$  inches; then  $16 : 12 :: 36 : 27 = BC$ , and consequently  $AC = 9$ , the two distances from the common centre of gravity.



3. *The Common Centre of Gravity of three or more given bodies or points A, B, C, D, &c, will be thus determined.*—If the given bodies lie all in the same straight line AD; by the last article, find P the centre of gravity of the two A and B, and Q the centre of gravity of C and D; then, considering P as the place of a body equal to the sum of A and B, and Q as the place of another body equal to both C and D, find S the common centre of gravity of these two sums, viz  $A + B$  collected in P, and  $C + D$  united in Q; so shall S be the common centre of gravity of all the four bodies A, B, C, D. And the same for any other number of bodies, always considering the sum of any number of them as united or placed in their common centre of gravity, when found.

*Otherwise, thus.* Take the distances of the given bodies from some fixed point as V, calling the distance  $VA = a$ ,  $VB = b$ ,  $VC = c$ ,  $VD = d$ , and the distance of the centre of gravity  $VS = x$ ; then  $SA = x - a$ ,  $SB = x - b$ ,  $SC = c - x$ ,  $SD = d - x$ , and by the nature of the lever  $A \cdot x - a + B \cdot x - b = C \cdot c - x + D \cdot d - x$ ; hence  $Ax + Bx + Cx + Dx = Aa + Bb + Cc + Dd$ , and  $x = \frac{Aa + Bb + Cc + Dd}{A + B + C + D} = VS$

the distance sought; which therefore is equal to the sum



A geometric diagram illustrating the properties of parallel lines. Two horizontal lines are shown, intersected by three transversal lines labeled AB, SC, and QD. The top horizontal line contains points V, P, S, and Q from left to right. The bottom horizontal line contains points A, B, C, and D from left to right. Transversal AB connects point A on the bottom line to point B on the top line. Transversal SC connects point S on the top line to point C on the bottom line. Transversal QD connects point Q on the top line to point D on the bottom line. Vertical dotted lines connect corresponding points between the two horizontal lines: BB' (labeled B at the bottom), SS' (labeled S at the bottom), and DD' (labeled D at the bottom). Point V is located to the left of point A on the bottom line.


A rectangle is shown with vertices labeled A (top-left), B (top-right), C (bottom-right), and D (bottom-left). A point E is located on the side AB. A horizontal line segment connects A and B, and a vertical line segment connects B and C. The rectangle is shaded with a light gray background.

8. In




8. *In a Trapezium.* Divide the figure into two triangles by the diagonal AC, and find the centres of gravity E and F of these triangles; join EF, and find the common centre G of these two by this proportion,  $ABC : ADC :: FG : EG$ , or  $ABCD : ADC :: EF : EG$ .

In like manner, for any other figure, whatever be the number of sides, divide it into several triangles, and find the centre of gravity of each; then connect two centres together, and find their common centre as above; then connect this and the centre of a third, and find the common centre of these; and so on, always connecting the last found common centre to another centre, till the whole are included in this process; so shall the last common centre be that which is required.



9. *In the Parabola BAC.* Put  $AD = x$ ,  $BD = y$ , and the parameter  $= p$ . Then, by the nature of the figure,  $px = y^2$ , and  $2y = 2\sqrt{px}$ ; hence  $2x\sqrt{px}$  is the fluxion of the weights, and  $2x\dot{x}\sqrt{px}$  is the fluxion of the momenta; then the fluent of the latter divided by that of the former, or  $\frac{4}{3}x^{\frac{5}{2}}\sqrt{p}$  divided by  $\frac{3}{2}x^{\frac{3}{2}}\sqrt{p}$ , gives  $\frac{2}{3}x = \frac{2}{3}AD$ , for AG, the distance of the centre of gravity G from the vertex A of the parabola.





10. *In the Circular Arc ABD,* considered as a physical line having gravity. It is manifest that the centre of gravity G of the arc, will be somewhere in the axis, or middle radius BC, C being the centre of the circle, which is considered as the point of suspension. Suppose F indefinitely near to A, and FH parallel to BC. Put the radius BC or AC  $= r$ , the semicircle  $AB = z$ , and the semichord  $AE = x$ ; then is  $AH = \dot{x}$ , and  $AF = \dot{z}$  the fluxion of the weights, and therefore  $CE \times \dot{z}$  is the fluxion of the momenta. But, by similar triangles,  $AC$  or  $r : CE :: AF$  or  $\dot{z} : AH$  or  $\dot{x}$ , therefore  $r\dot{x} = CE \times \dot{z}$ , and so  $r\dot{x}$  is also the fluxion of the momenta; the fluent of which is  $rx$ , and this divided by  $z$  the weight, gives  $\frac{rx}{z} = \frac{AC \times AE}{AB} = \frac{AC \times AED}{ABD} = CG$  the distance of the centre of gravity from the centre C of the circle; being a 4th proportional to the given arc, its chord, and the radius of the circle.

Vol. I.

Hence, when the arc becomes the semicircle ABK, the above expression becomes  $\frac{IC^2}{IB}$  or  $\frac{r^2}{1.5708r} = \frac{r}{1.5708} = .6366r$ , viz a third proportional to a quadrant and the radius.

11. *In the Circular Sector ABDC.* Here also the centre of gravity will be in the axis or middle radius BC. Now with any smaller radius describe the concentric arc LMN, and put the radius AC or BC  $= r$ , the arc  $ABD = a$ , its chord  $AED = c$ , and the variable radius CL or CM  $= y$ ; then as  $r : y :: a : \frac{ay}{r}$  = the arc LMN, and  $r : y :: c : \frac{cy}{r}$  = the chord LON; also, by the last article, the distance of the centre of gravity of the arc LMN is  $\frac{CM \times LON}{LMN} = \frac{CM \times AED}{ABD} = \frac{cy}{a}$ ; hence the arc LMN or  $\frac{ay}{r}$  multiplied by  $y$  gives  $\frac{ay^2}{r}$  the fluxion of the weights, and this multiplied by  $\frac{cy}{a}$  the distance of the common centre of gravity, gives  $\frac{cy^2y}{r}$  the fluxion of the momenta; the fluent of which, viz  $\frac{cy^3}{3r}$ , divided by  $\frac{ay^2}{2r}$ , the fluent of the weights, gives  $\frac{2cy}{3a}$  for the distance of the centre of gravity of the sector CLEN from the centre C; and when  $y = r$ , it becomes  $\frac{2cr}{3a} = CG$  for that of the sector CABD proposed; being  $\frac{2}{3}$  of a 4th proportional to the arc of the sector, its chord, and the radius of the circle.

Hence, when the sector becomes a semicircle, the last expression becomes  $\frac{4r^2}{3a} = \frac{2IC^2}{3IB}$  or  $\frac{2}{3}$  of a 3d proportional to a quadrantal arc and the radius. Or it is equal to  $\frac{4r}{5p} = .4244r$  from the centre C; where  $p = 3.1416$ .



12. *In the Cone ADB.* Putting  $a = DC$ ,  $b =$  area of the base AEB, and  $x = Dc$  any variable altitude; then as  $a^2 : x^2 :: b : \frac{bx^2}{a^2} =$  area  $aeb$ ; hence the fluxion

M m

ion



ion of the weights is  $\frac{bx^2\dot{x}}{a^2}$ , whose fluent, or the solid, is  $\frac{bx^3}{3a^2}$ ; and the fluxion of the momenta is  $\frac{bx^3\dot{x}}{a^2}$ , whose fluent is  $\frac{bx^4}{4a^2}$ ; then this fluent divided by the former fluent gives  $\frac{3}{4}x$  or  $\frac{3}{4}Dc$  for the distance of the centre of gravity of the cone  $Dab$ , or  $\frac{3}{4}DC$  for that of the cone  $DAB$  below the vertex  $D$ .

And the same is the distance in any other pyramid. So that all pyramids of the same altitude, have the same centre of gravity.

13. In like manner are we to proceed for the centre of gravity in other bodies. Thus, the altitude of the segment of a sphere, or spheroid, or conoid, being  $x$ ,  $a$  being the whole of that axis itself; then the distance of the centre of gravity in each of these bodies, from the vertex, will be as follows, viz,

$\frac{4a-3x}{6a-4x}x$  in the sphere or spheroid,

$\frac{5}{8}x$  in the semisphere or semispheroid,

$\frac{3}{5}x$  in the parabolic conoid,

$\frac{4a+3x}{6a+3x}x$  in the hyperbolic conoid.

14. To determine the Centre of Gravity in any Body Mechanically. Lay the body on the edge of any thing, as a triangular prism, or such like, moving it backward and forward till the parts on both sides are in equilibrio; then is that line just in, or under the centre of gravity. Balance it again in another position, to find another line passing through the centre of gravity; then the intersection of these two lines will give the place of that centre itself.

The same may be done by laying the body on an horizontal table, as near the edge as possible without its falling, and that in two positions, as lengthwise and breadthwise: then the common intersection of the two lines contiguous to the edge, will be its centre of gravity. Or it may be done by placing the body on the point of a style, &c, till it rest in equilibrio. It was by this method that Borelli found that the centre of gravity in a human body, is between the nates and pubis; so that the whole gravity of the body is collected into the place of the genitals; an instance of the wisdom of the creator, in placing the membrum virile in the part, which is the most convenient for copulation.

The same otherwise thus. Hang the body up by any point; then a plumb line hung over the same point, will pass through the centre of gravity; because that centre will always descend to the lowest point when the body comes to rest, which it cannot do except when it falls in the plumb line. Therefore, marking that line upon it, and suspending the body by another point, with the plummet, to find another such line, the intersection of the two will give the centre of gravity.

Or thus. Hang the body by two strings from the same tack, but fixed to different points of the body; then a plummet, hung by the same tack, will fall on the centre of gravity.

In the 4th volume of the New Acts of the Academy of Petersburg, is the demonstration of a very general theorem concerning centres of gravity, by M. Lhuillier;

a particular example only of the general proposition, will be as follows: Let  $A, B, C$ , be the centres of gravity of three bodies;  $a, b, c$  their respective masses, and  $Q$  their common centre of gravity. Let right lines  $QA, QB, QC$ , be drawn from the common centre to that of each body, and the latter be connected by right lines  $AB, AC$ , and  $BC$ ; then

$$QA^2 \times a + QB^2 \times b + QC^2 \times c = AB^2 \times \frac{ab}{a+b+c} + AC^2 \times \frac{ac}{a+b+c} + BC^2 \times \frac{bc}{a+b+c}.$$

Uses of the Centre of Gravity. This point is of the greatest use in mechanics, and many important concerns in life, because the place of that centre is to be considered as the place of the body itself in computing mechanical effects; as in the oblique pressures of bodies, banks of earth, arches of bridges, and such like.

The same centre is even useful in finding the superficial and solid contents of bodies; for it is a general rule, that the superficies or solid generated by the rotation of a line or plane about any axis, is always equal to the product of the said line or plane drawn into the circumference or path described by the centre of gravity. For example, it was found above at art. 11, that in a semicircle, the distance of the centre of gravity from the centre of the circle, is  $\frac{4r}{3p}$ ; and therefore the path of that

centre, or circumference described by it whilst the semicircle revolves about its diameter, is  $\frac{8}{3}r$ ; also the area of the semicircle is  $\frac{1}{2}pr^2$ ; hence the product of the two is  $\frac{4}{3}pr^3$ , which, it is well known, is equal to the solidity of the sphere generated by the revolution of the semicircle.

And hence also is obtained another method of finding mathematically the centre of gravity of a line or plane, from the contents of the superficies or solid generated by it. For if the generated superficies or solid be divided by the generating line or plane, the quotient will be the circumference described by the centre of gravity; and consequently this divided by  $2p$  gives the radius, or distance of that centre from the axis of rotation. So, in the semicircle, whose area is  $\frac{1}{2}pr^2$ , and the content of the sphere generated by it  $\frac{4}{3}pr^3$ ; here the latter divided by the former is  $\frac{8}{3}r$ , and this divided by

$2p$  gives  $\frac{4r}{3p}$  for the distance of the centre of gravity from the axis, or from the centre of the semicircle. The property last mentioned, relative to the relation between the centre of gravity and the figure generated by the revolution of any line or plane, is mentioned by Pappus, in the preface to his 7th book; and father Guldin has more fully demonstrated it in his 2d and 3d books on the Centre of Gravity.

The principal writers on the centre of gravity are Archimedes, Pappus, Guldini, Wallis, Casatus, Carré, Hays, Wolfius, &c.

CENTRE of Gyration, is that point in which if the whole mass be collected, the same angular velocity will be generated in the same time, by a given force acting at any place, as in the body or system itself. This point differs from the centre of oscillation, in as much as in this latter case the motion of the body is produced by the gravity of its own particles, but in the case of the centre



centre of gyration the body is put in motion by some other force acting at one place only.

To determine the Centre of Gyration, in any body, or system of bodies composed of the parts A, B, C, &c, moving about the point S, when urged by a force  $f$  acting at any point P. Let R be that centre: then, by mechanics, the angular velocity generated in the system

by the force  $f$ , is as  $\frac{f \cdot SP}{A \cdot SA^2 + B \cdot SB^2 + C \cdot SC^2 \&c}$ , and, by the same, the angular velocity of the matter placed all in the point R, is  $\frac{f \cdot AP}{A + B + C \&c \times SR^2}$ ;

then since these two are to be equal, their equation will give  $SR = \sqrt{\frac{A \cdot SA^2 + B \cdot SB^2 + C \cdot SC^2 \&c}{A + B + C \&c}}$ , for the distance of the centre of gyration sought, below the axis of motion.

Now because the quantity  $A \cdot SA^2 + B \cdot SB^2 + \&c$  is  $= SG \cdot SO \cdot b$ , where G is the centre of gravity, O the centre of oscillation, and  $b$  the whole body or sum of A, B, C, &c; therefore it follows that  $SR^2 = SG \cdot SO$ ; that is, the distance of the centre of gyration, is a mean proportional between those of gravity and oscillation.

And hence also, if  $p$  denote any particle of a body, placed at the distance  $d$  from the axis of motion; then is  $SR = \frac{\text{sum of all the } pd^2}{\text{body } b}$ ; from whence the point

R may be determined in bodies by means of Fluxions.

CENTRE of an Hyperbola, is the middle of the axis, or of any other diameter, being the point without the figure in which all the diameters intersect one another; and it is common to all the four conjugate hyperbolas.

CENTRE of Magnitude, is the point which is equally distant from all the similar external parts of a body. This is the same as the centre of gravity in homogeneous bodies that can be cut into like and equal parts according to their length, as in a cylinder or any other prism.

CENTRE of Motion, is the point about which any body, or system of bodies, moves, in a revolving motion.

CENTRE of Oscillation, is that point in the axis or line of suspension of a vibrating body, or system of bodies, in which if the whole matter or weight be collected, the vibrations will still be performed in the same time, and with the same angular velocity, as before. Hence, in a compound pendulum, its distance from the point of suspension is equal to the length of a simple pendulum whose oscillations are isochronal with those of the compound one.


Mr. Huygens, in his Horologium Oscillatorium, first shewed how to find the centre of oscillation. At the beginning of his discourse on this subject, he says, that Merfennus first proposed the problem to him while he was yet very young, requiring him to resolve it in the cases of sectors of circles suspended by their angles, and by the middle of their bases, both when they oscillate sideways and flatways; as also for triangles and the segments of circles, either suspended from their vertex or the middle of their bases. But, says he, not

having immediately discovered any thing that would open a passage into this business, I was repulsed at first setting out, and stopped from a further prosecution of the thing; till being farther incited to it by adjusting the motion of the pendulums of my clock, I surmounted all difficulties, going far beyond Descartes, Fabry, and others, who had done the thing in a few of the most easy cases only, without any sufficient demonstration; and solving not only the problems proposed by Merfennus, but many others that were much more difficult, and shewing a general way of determining this centre, in lines, superficies, and solids.

In the Leipzig Acts for 1691 and 1714, this doctrine is handled by the two Bernoullis: and the same is also done by Herman, in his treatise De Motu Corporum Solidorum et Fluidorum.

It may also be seen in treatises on the Inverse Method of Fluxions, where it is introduced as one of the examples of that method. See Hayes, Carré, Wollius, &c.

To determine the Centre of Oscillation, in any Compound Mass or Body MN, or of any System of Bodies A, B, C, &c.



Let MN be the plane of vibration, to which plane conceive all the matter to be reduced by letting fall perpendiculars to this plane from every particle in the body; a supposition which will not alter the vibration of the body, because the particles are still at the same distance from the axis of motion. Let O be the centre of oscillation, and G the centre of gravity; through the axis S draw SGO, and the horizontal line ST; then from every particle A, B, C, &c, let fall perpendiculars Aa and Ap, Bb and Bq, Cc and Cr, &c, to these two lines; and join SA, SB, SC; also draw Gm and On perpendicular to ST.

Now the forces of the weights A, B, C, to turn the body about the axis, are  $A \cdot Sp$ ,  $B \cdot Sq$ ,  $-C \cdot Sr$ ; and, by mechanics, the forces opposing that motion are  $A \cdot SA^2$ ,  $B \cdot SB^2$ ,  $C \cdot SC^2$ ; therefore the angular motion generated in the system is  $\frac{A \cdot Sp + B \cdot Sq - C \cdot Sr}{A \cdot SA^2 + B \cdot SB^2 + C \cdot SC^2}$ ;

In like manner, the angular velocity which any body or particle  $p$ , situated in O, generates in the system, by its weight, is  $\frac{p \cdot Sn}{p \cdot SO^2}$ , or  $\frac{Sn}{SO^2}$ , or  $\frac{Sm}{SG \cdot SO}$  because of the



the similar triangles  $SGm$ ,  $SON$ . But, by the conditions of the problem, the vibrations are performed alike in both these cases; therefore these two expressions must be equal to each other, that is  $\frac{Sm}{SG \cdot SO} =$

$$\frac{A \cdot Sp + B \cdot Sq - C \cdot Sr}{A \cdot SA^2 + B \cdot SB^2 + C \cdot SC^2}, \text{ and consequently } SO = \frac{Sm}{SG} \times \frac{A \cdot SA^2 + B \cdot SB^2 + C \cdot SC^2}{A \cdot Sp + B \cdot Sq - C \cdot Sr}.$$

But, by mechanics again, the sum of the forces  $A \cdot Sp + B \cdot Sq - C \cdot Sr$  is equal  $A + B + C \cdot Sm$  the force of the same matter collected all into its centre of gravity  $G$ ; and therefore

$$SO = \frac{A \cdot SA^2 + B \cdot SB^2 + C \cdot SC^2}{(A + B + C) \cdot SG}, \text{ which is the}$$

distance of the centre of oscillation  $O$  below the axis of suspension.

Farther, because it was found under the article *Centre of Gravity*, that  $(A + B + C) \cdot SG = A \cdot Sa + B \cdot Sb + C \cdot Sc$ , therefore  $SO = \frac{A \cdot SA^2 + B \cdot SB^2 + C \cdot SC^2}{A \cdot Sa + B \cdot Sb + C \cdot Sc}$

is the same distance of the centre of oscillation; where any of the products  $A \cdot Sa$ ,  $B \cdot Sb$ , &c are to be taken negatively when the points  $a$ ,  $b$ , &c lie above the point  $S$ , or where the axis passes through.

Again, because, by Eucl. II 12 and 13,  
it is  $SA^2 = SG^2 + GA^2 - 2SG \cdot Ga$ ,  
and  $SB^2 = SG^2 + GB^2 + 2SG \cdot Gb$ ,  
and  $SC^2 = SG^2 + GC^2 + 2SG \cdot Gc$ , &c;

and because by Mechanics, the sum of the last terms is nothing, namely  $-2SG \cdot Ga + 2SG \cdot Gb + 2SG \cdot Gc$  &c  $= 0$ ; therefore the sum of the others, or  $A \cdot SA^2 + B \cdot SB^2$  &c  $= A + B$  &c  $\cdot SG^2 + A \cdot GA^2 + B \cdot GB^2 + C \cdot GC^2$  &c, or  $= b \cdot SG^2 + A \cdot GA^2 + B \cdot GB^2 + C \cdot GC^2$  &c;

where  $b$  denotes the body, or sum  $A + B + C$  &c of all the parts: this value then being substituted in the numerator of the 2d value of  $SO$  above-found, it becomes

$$SO = \frac{b \cdot SG^2 + A \cdot GA^2 + B \cdot GB^2 \text{ \&c}}{b \cdot SG}, \text{ or}$$

$$SO = SG + \frac{A \cdot GA^2 + B \cdot GB^2 \text{ \&c}}{b \cdot SG}.$$

From which it appears that the centre of oscillation is always below the centre of gravity, and that the difference or distance between them is

$$GO = \frac{A \cdot GA^2 + B \cdot GB^2 \text{ \&c}}{b \cdot SG}.$$

It farther follows from hence, that  $SG \cdot GO = \frac{A \cdot GA^2 + B \cdot GB^2 \text{ \&c}}{\text{the body } b}$ ; that is, the rectangle  $SG \cdot GO$  is

always the same constant quantity, wherever the point of suspension  $S$  is placed, since the point  $G$  and the bodies  $A$ ,  $B$ , &c, are constant. Or  $GO$  is always reciprocally as  $SG$ , that is  $GO$  is less as  $SG$  is greater; and the points  $G$  and  $O$  coincide when  $SG$  is infinite; but when  $S$  coincides with  $G$ , then  $GO$  is infinite, or  $O$  is at an infinite distance.

*To find the Centre of Oscillation by means of Fluxions.* From the premises is derived this general method for the centre of oscillation, viz, let  $x$  be the abscissa of an oscillating body, and  $y$  its corresponding ordinate or section; then will the distance  $SO$  of the centre of oscil-

lation below the axis of suspension  $S$ , be equal to the fluent of  $yx^2x$  divided by the fluent of  $yx$ . So that, if from the nature or equation of any given figure, the value of  $y$  be expressed in terms of  $x$ , or otherwise, and substituted in these two fluxions; then the fluents being duly found, and the one divided by the other, the quotient will be the distance to the centre of oscillation in terms of the absciss  $x$ .

But when the body is suspended by a very fine thread of a given length  $a$ , then the fluent of  $\frac{a+x}{2} \cdot yx$  divided by the fluent of  $a+x \cdot yx$  gives the distance of the same centre of oscillation below the point of suspension.

*Ex.* For example, in a right line, or rectangle or cylinder or any other prism, whose constant section is  $y$ , or the constant quantity  $a$ ; then  $yx^2x$  is  $ax^2x$ , whose fluent is  $\frac{1}{3}ax^3$ ; also  $yx$  is  $ax$ , whose fluent is  $\frac{1}{2}ax^2$ ; and the quotient of the former  $\frac{1}{3}ax^3$  divided by the latter  $\frac{1}{2}ax^2$ , is  $\frac{2}{3}x$  for the distance of the centre of oscillation below the vertex in any such figure, namely having every where the same breadth or section, that is, at two-thirds of its length.

In like manner the centre of oscillation is found for various figures, vibrating flatways, and are as they are expressed below, viz,

Nature of the Figure.	When suspended by Vertex.
Isoceles triangle	$\frac{3}{4}$ of its altitude
Common Parabola	$\frac{5}{7}$ of its altitude
Any Parabola	$\frac{2m+1}{3m+1} \times$ its altitude.
As to figures moved laterally or sideways, or edgewise, that is about an axis perpendicular to the plane of the figure, the finding the centre of oscillation is somewhat difficult; because all the parts of the weight in the same horizontal plane, on account of their unequal distances from the point of suspension, do not move with the same velocity; as is shewn by Huygens, in his Horol. Oscil. He found, in this case, the distance of the centre of oscillation below the axis, viz,	
In a circle,	$\frac{3}{4}$ of the diameter:
In a rectangle, susp. by one angle,	$\frac{2}{3}$ of the diagonal:
In a parabola susp. by its vertex,	$\frac{5}{7}$ axis $+$ $\frac{1}{3}$ param.
The same susp. by mid. of base,	$\frac{4}{7}$ axis $+$ $\frac{1}{2}$ param.
In a sector of a circle	$\frac{3 \text{ arc} \times \text{radius}}{4 \text{ chord}}$ :
In a cone	$\frac{4}{5}$ axis $+$ $\frac{\text{radius base}^2}{5 \text{ axis}}$ :
In a sphere	$g + \frac{2r^2}{5g}$ , where $r$ is the radius, and $g = a + r$ the rad. added to the length of the thread.

See also Simpson's Fluxions, art. 183 &c.

*To find the Centre of Oscillation Mechanically or Experimentally.* Make the body oscillate about its point of suspension; and hang up also a simple pendulum of such a length that it may vibrate or just keep time with the other body: then the length of the simple pendulum is equal to the distance of the centre of oscillation of the body below the point of suspension.

Or it will be still better found thus: Suspend the body very freely by the given point, and make it vibrate



in small arcs, counting the vibrations it makes in any portion of time, as a minute, by a good stop watch; and let that number of oscillations made in a minute be called  $n$ : then shall the distance of the centre of oscillation be  $SO = \frac{140850}{nn}$  inches. For, the length of

the pendulum vibrating seconds, or 60 times in a minute, being  $39\frac{1}{8}$  inches, and the lengths of pendulums being reciprocally as the square of the number of vibrations made in the same time, therefore  $n^2 : 60^2 ::$

$39\frac{1}{8} : \frac{140850}{nn}$  the length of the pendulum which vibrates  $n$  times in a minute, or the distance of the centre of oscillation below the axis of motion.

**CENTRE of Percussion**, in a moving body, is that point where the percussive stroke is the greatest, in which the whole percussive force of the body is supposed to be collected; or about which the impetus of the parts is balanced on every side, so that it may be stopped by an immovable obstacle at this point, and rest on it, without acting on the centre of suspension.

1. When the percussive body revolves about a fixed point, the centre of percussion is the same with the centre of oscillation; and is determined in the same manner, viz, by considering the impetus of the parts as so many weights applied to an inflexible right line void of gravity; namely, by dividing the sum of the products of the forces of the parts multiplied by their distances from the point of suspension, by the sum of the forces. And therefore what has been above shewn of the centre of oscillation, will hold also of the centre of percussion when the body revolves about a fixed point. For instance, that the centre of percussion in a cylinder is at  $\frac{2}{3}$  of its length from the point of suspension, or that a flick of a cylindrical figure, supposing the centre of motion at the hand, will strike the greatest blow at a point about two-thirds of its length from the hand.

2. But when the body moves with a parallel motion, or all its parts with the same celerity, then the centre of percussion is the same as the centre of gravity. For the momenta are the products of the weights and celerities; and to multiply equiponderating bodies by the same velocity, is the same thing as to take equimultiples; but the equimultiples of equiponderating bodies do also equiponderate; therefore equivalent momenta are disposed about the centre of gravity, and consequently in this case the two centres coincide, and what is shewn of the one will hold in the other.

**Centre of Percussion in a fluid**, is the same as out of it.

**CENTRE of a Parallelogram**, the point in which its diagonals intersect.

**CENTRE of Pressure**, of a fluid against a plane, is that point against which a force being applied equal and contrary to the whole pressure, it will just sustain it, so as that the body pressed on will not incline to either side.—This is the same as the centre of percussion, supposing the axis of motion to be at the intersection of this plane with the surface of the fluid; and the centre of pressure upon a plane parallel to the horizon, or upon any plane where the pressure is uniform, is the same as the centre of gravity of that plane. *Emerson's Mechanics*, prop. 91.

**CENTRE of a Regular Polygon**, or *Regular Body*, is the same as that of the inscribed, or circumscribed circle or sphere.

**CENTRE of a Sphere**, is the same as that of its generating semicircle, or the middle point of the sphere, from whence all right lines drawn to the superficies, are equal.

**CENTRING of an Optic Glass**, the grinding it so as that the thickest part be exactly in the middle.

Cassini the younger has a discourse expressly on the necessity of well centring the object glass of a large telescope, that is, of grinding it so as that the centre may fall exactly in the axis of the telescope. *Mem. Acad.* 1710.

Indeed one of the greatest difficulties in grinding large optic glasses is, that in figures so little convex, the least difference will throw the centre two or three inches out of the middle. And yet Dr. Hook remarks, that though it were better the thickest part of a long object glass were exactly in the middle, yet it may be a very good one when it is an inch or two out of it. *Philos. Trans.* No 4.

**CENTRIFUGAL FORCE**, is that by which a body revolving about a centre, or about another body, endeavours to recede from it. And


**CENTRIPETAL FORCE**, is that by which a moving body is perpetually urged towards a centre, and made to revolve in a curve, instead of a right line.

Hence, when a body revolves in a circle, these two forces, viz, the centrifugal and centripetal, are equal and contrary to each other, since neither of them gains upon the other, the body being in a manner equally balanced by them. But when, in revolving, the body recedes farther from the centre, then the centrifugal exceeds the centripetal force; as in a body revolving from the lower to the higher apsis, in an ellipse, and respecting the focus as the centre. And when the revolving body approaches nearer to the centre, the centrifugal is less than the centripetal force; as while the body moves from the farther to the nearer extremity of the transverse axis of the ellipse: the two forces being equal to each other only at the very extremities of that axis.

It is one of the established laws of nature, that all motion is of itself rectilinear, and that the moving body never recedes from its first right line, till some new impulse be superadded in a different direction: after that new impulse the motion becomes compounded, but it is still rectilinear, though not in the same line or direction as before. To move in a curve, it must receive a new impulse in a different direction every moment; a curve not being reducible to any number of finite right lines. If then a body, continually drawn towards a centre, be projected in a line that does not pass through that centre, it will describe a curve; in each point of which, as *A*, it will endeavour to recede from the curve, and proceed in the tangent *AD*; and if nothing hindered, it would actually proceed in it; so as in the same time in which it describes the arch *AE*, it would recede the length of the line *DE*, perpendicular to *AD*, by its centrifugal force: Or being projected in the direction *AD*, but being continually drawn out of its direction into a curve by a centripetal force.



force, so as to fall below the line of direction by the perpendicular space DE: Then the centrifugal or centripetal force is as this line of deviation DE; supposing the arch AE indefinitely small.



The doctrine of centrifugal forces was first mentioned by Huygens, at the end of his *Horologium Oscillatorium*, published in 1673, and demonstrated in the volume of his *Posthumous Works*, as also by Guido Grandi; where he has given a few easy cases in bodies revolving in the circumference of circles. But Newton, in his *Principia*, was the first who fully handled this doctrine; at least as far as regards the conic sections. After him there have been several other writers upon this subject; as Leibnitz, Varignon in the *Mem. de l'Acad.* Keil in the *Philos. Trans.* and in his *Physics*, Bernoulli, Herman, Cotes in his *Harmonia Mensurarum*, Maclaurin in his *Geometrica Organica*, and in his *Fluxions*, and Euler in his book *de Motu*, where he considers the curves described by a body acted on by centripetal forces tending to several fixed points.

See also the art. *Central Forces*, where this doctrine is more fully explained.

**CENTROBARICO**, the same as centre of gravity.

**CENTROBARIC-Method**, is a method of determining the quantity of a surface or solid, by means of the generating line or plane, and its centre of gravity. The doctrine is chiefly comprized in this theorem:

Every figure, whether superficial or solid, generated by the motion of a line or plane, is equal to the product of the generating magnitude and the path of its centre of gravity, or the line which its centre of gravity describes.

See more of this subject under the article *Centre of Gravity*.

**CENTRUM**, in Geometry, Mechanics, &c. See **CENTRE**.

**CENTRUM Phonicum**, in Acoustics, is the place where the speaker stands in polysyllabical and articulate echoes.

**CENTRUM Phonocampticum**, is the place or object that returns the voice in an echo.

**CEPHEUS**, a constellation of the northern hemisphere, being one of the 48 old asterisms. The Greeks fable that Cepheus was a king of Ethiopia, and the father of Andromeda, the princess who was delivered up to be devoured by a sea monster, from which she was rescued by Perseus.

The stars of this constellation, in Ptolemy's catalogue are 13, in Tycho's 11, in Hevelius's 51, and in the *Britannic catalogue* 35.

**CERBERUS**, one of the new constellations formed by Hevelius out of the unformed stars, and added to the 48 old asterisms. It contains only 4 stars, which are enumerated under Hercules in the *Britannic catalogue*.

**CETUS**, the *Whale*, a southern constellation, and one of the 48 old asterisms. The Greeks pretend that it was the sea-monster sent by Neptune to devour Andromeda, but was killed by Perseus.

In the neck of the whale is a remarkable star, *Collo Ceti*, which appears and disappears periodically, or rather grows brighter and fainter by turns, owing it is supposed to the alternate turning of its bright and dark sides towards us, as it revolves upon its axis, or else owing to the star having a flattish form. The period of its changes is about 312 days. Bullialdus in *Phil. Trans.* vol. 2, Hevelius *ibid.* vol. 6, Herschel *ibid.* vol. 70, Marald. in *Mem. Acad.* 1719.

The stars in the constellation Cetus, in Ptolemy's catalogue, are 22, in Tycho's 21, in Hevelius's 45, and in the *Britannic catalogue* 97.

**CHAIN**, in Surveying, is a lineal measure, consisting of a certain number of iron links, usually 100: serving to take the dimensions of fields &c.

At every 10th link is usually fastened a small brass plate, with a figure engraven upon it, or else cut into different shapes, to shew how many links it is from one end of the chain.

Chains are of various kinds and lengths; as

1. A chain of 100 feet long, each link one foot, for measuring of large distances only, when regard is not proposed to be had to acres &c, in the superficial content.

2. A chain of one pole or 16 feet and a half in length; especially useful in measuring and laying out gardens and orchards, or the like, by the pole or rod measure.

3. A chain of 4 poles, or 66 feet, or 22 yards, in length, called Gunter's chain, and is peculiarly adapted to the business of Surveying or Land-measuring, because that 10 square chains just make an English acre of land; so that the dimensions being taken in these chains, and thence the contents computed in square chains, they are readily turned into acres by dividing by 10, or barely cutting off the last figure from the square chains. But it is still better in practice to proceed thus, viz, count the dimensions, not in chains, but all in links; then the contents are in square links; and five figures being cut off for decimals, the rest are acres; that is four figures to bring the square links to square chains, and one more to bring the square chains to acres.

In this chain, the links are each 7 inches and  $\frac{92}{1000}$ , or 7.92 inches in length, which is very nearly  $\frac{2}{3}$  of a foot. And hence any number of chains or links are easily brought to feet or inches, or the contrary: the best way of doing which is this: multiply the number of links by 66, then cut off two figures for decimals, and the rest are feet: or multiply links by 22 for yards, cutting off two figures.

**CHALDRON**, of Coals, an English dry measure of capacity consisting of 36 bushels heaped up.

The chaldron of coals is accounted to weigh about 2000 pounds.—On ship board, 21 chaldrons of coals are allowed to the score.

**CHAMBER** of a *Mortar*, or *some cannon*, is a cell or cavity at the bottom of the bore, to receive the charge of powder.

It is not found by experience that chambers have any



any sensible effect on the velocity of the shot, unless in the largest ordnance, as mortars or very large cannon. Neither is it found that the form of them is very material; a small cylinder is as good as any; though mathematical speculations may shew a preference of one form over another. But in practice, the chief point to be observed, is to have the chamber of a size just to contain the charge of powder, and no more, that the ball may lie close to the charge; and that its entrance may point exactly to the centre of the ball.

CHAMBERS (EHRRAIM), author of the dictionary of sciences called the *Cyclopædia*. He was born at Milton in the county of Westmoreland, where he received the common education for qualifying a youth for trade and commerce. When he became of a proper age, he was put apprentice to Mr. Senex the globe-maker, a business which is connected with literature, especially with geography and astronomy. It was during Mr. Chambers's residence with this skilful artist, that he acquired that taste for literature which accompanied him through life, and directed all his pursuits. It was even at this time that he formed the design of his grand work, the *Cyclopædia*; some of the first articles of which were written behind the counter. To have leisure to pursue this work, he quitted Mr. Senex, and took chambers at Gray's-Inn, where he chiefly resided during the rest of his life. The first edition of the *Cyclopædia*, which was the result of many years intense application, appeared in 1728, in 2 vols. folio. The reputation that Mr. Chambers acquired by the execution of this work, procured him the honour of being elected F. R. S. Nov. 6, 1729. In less than ten years time, a second edition became necessary; which accordingly was printed, with corrections and additions, in 1738; and this was followed by a third edition the very next year.

Although the *Cyclopædia* was the chief business of Mr. Chambers's life, and may be regarded as almost the sole foundation of his fame, his attention was not wholly confined to this undertaking. He was concerned in a periodical publication, called, *The Literary Magazine*, which was begun in 1735. In this work he wrote a variety of articles; particularly a review of Morgan's *Moral Philosopher*. He was also concerned with Mr. John Martyn, professor of botany at Cambridge, in preparing for the press a translation and abridgment of the *Philosophical History and Memoirs of the R. Acad. of Sciences at Paris*; which work was not published till 1742, some time after our author's decease, in 5 volumes 8vo. Mr. Chambers was also author of the translation of the *Jesuit's Perspective*, from the French, in 4to; which has gone through several editions.

Mr. Chambers's close and unremitting attention to his studies at length impaired his health, and obliged him occasionally to take a country lodging, but without much benefit; he afterwards visited the south of France, but still with little effect; he therefore returned to England, where he soon after died, at Islington, May 15, 1740, and was buried at Westminster Abbey.

After the author's death, two more editions of his *Cyclopædia* were published. The proprietors afterwards procured a supplement to be compiled, by Mr.

Scott and Dr. Hill, but chiefly by the latter, which extended to two volumes more; and the whole has since been reduced into one alphabet in 4 volumes, by Dr. Rees, forming a very valuable body of the sciences.

CHAMBRANLE, the border, frame, or ornament of stone or wood, surrounding the three sides of doors, windows, and chimneys. This is different in the different orders: when it is plain, and without mouldings, it is called simply and properly, *band, case, or frame*. In an ordinary door, it is mostly called *door-case*; in a window, the *window-frame*.

The Chambranle consists of three parts; the two sides, called *ascendants*; and the top, called the *traverse* or *supercilium*.

CHAMFER, or CHAMFERET, an ornament, in architecture, consisting of half a scotia; being a kind of small furrow or gutter on a column.

CHAMFERING, is used for cutting the edge or the end of any thing bevel, or aslope.

CHANCE, *the Doctrine and Laws of*, are the same as those of expectation, or probability, &c; which see. Chances, in play, consist of the number of ways by which events may happen. Thus, if a halfpenny, or other piece of money, be tossed up, there are two events, or chances, or sides that may turn up, namely one chance for turning up a head, and one for the contrary; that is, it is an equal chance to throw a head or not. And in throwing a common die, which has 6 faces, there are in all 6 chances, that is one chance for throwing an ace or any other single point, and 5 chances against it; or it is 5 to 1 that such assigned point does not come up.

Upon this subject, see De Moivre, Simpson, &c.

CHANDELIERS, in Fortification, a kind of wooden parapet, consisting of upright timbers supporting others laid across the tops of them, 6 feet high, and fortified with fascines &c. They are used to cover the workmen in approaches, galleries, and mines. And they differ from blinds only in this, that the former serve to cover the men before, and the latter over head.

CHANGES, the permutations or variations of any number of things, with regard to their position, order, &c; as how many changes may be rung on a number of bells, or how many different ways any number of persons may be placed, or how many several variations may be made of any number of letters, or any other things proposed to be varied.

To find out such number of changes, multiply continually together all the terms in a series of arithmetical progression, whose first term and common difference are each unity or 1, and the last term the number of things proposed to be varied, thus  $1 \times 2 \times 3 \times 4 \times 5$  &c. till the last number be the proposed number of things. For,

If there be only two things, as *a* and *b*, they admit of a double order or position only; for they may be placed either thus *ab* or thus *ba*, viz,  $1 \times 2 = 2$  ways.

If there be three things, *a*, *b*, and *c*, they will admit of 6 variations =  $1 \times 2$  since as in the margin, and no more; three each of the three may be combined there different ways with each of the other two.

<i>a</i>	<i>b</i>	<i>c</i>
<i>a</i>	<i>c</i>	<i>b</i>
<i>b</i>	<i>a</i>	<i>c</i>
<i>b</i>	<i>c</i>	<i>a</i>
<i>c</i>	<i>a</i>	<i>b</i>
<i>c</i>	<i>b</i>	<i>a</i>

And



And if there be 4 things, each of them may be combined 4 ways with each order of the other three, that is 4 times 6 ways, or  $1 \times 2 \times 3 \times 4 = 24$  ways.

In like manner, the combinations  
of 5 things are  $1 \times 2 \times 3 \times 4 \times 5 = 120$   
of 6 things are  $1 \times 2 \times 3 \times 4 \times 5 \times 6 = 720$   
&c.

So that if it be proposed to assign how many different ways a company of 6 persons may be placed, at table for instance, the answer will be 720 ways. Also the number of changes that can be rung on 7 bells, are  $1 \times 2 \times 3 \times 4 \times 5 \times 6 \times 7$  or  $720 \times 7 = 5040$  changes.

CHAPTERS, the crowns or upper parts of a pillar or column.

CHAPPE (JEAN D'AUTEROCHÉ), a French astronomer, was born at Mauriac, in Auvergne, March 2, 1728. A taste for drawing and mathematics appeared in him at a very tender age; and he owed to Dom Germain a knowledge of the first elements of mathematics and astronomy. M. Cassini, after assuring himself of the genius of this young man, undertook to improve it. He employed him upon the map of France, and the translation of Halley's tables, to which he made considerable additions. The king charged him in 1753 with drawing the plan of the county of Bitche, in Lorraine, all the elements of which he determined geographically. He occupied himself greatly with the two comets of 1760; and the fruit of his labour was his Elementary treatise on the theory of those comets, enriched with observations on the zodiacal light, and on the aurora borealis. He soon after went to Tobolsk, in Siberia, to observe the transit of Venus over the sun; a journey which greatly impaired his health. After two years absence he returned to France in 1762, where he occupied himself for some time in putting in order the great quantity of observations he had made. M. Chappe also went to observe the next transit of Venus, viz that of 1769, at California, on the west side of North America, where he died of a dangerous epidemic disease, the 1st of August 1769. He had been named Adjunct Astronomer to the Academy the 17th of January 1759.

The published works of M. Chappe, are,

1. The Astronomical Tables of Dr. Halley; with observations and additions: in 8vo, 1754.

2. Voyage to California to observe the transit of Venus over the sun, the 3d of June 1769: in 4to, 1772.

3. He had a considerable number of papers inserted in the Memoirs of the Academy, for the years 1760, 1761, 1764, 1765, 1766, 1767, and 1768; chiefly relating to astronomical matters.

CHAPTREL, the same with Imposst.

CHARACTERISTIC, of a Logarithm, the same as Index, or Exponent. This term was first used by Briggs in the 4th section of his Arithmetica Logarithmica, where he treats particularly of it; meaning by it, the integral or first part of a logarithm towards the left hand, which expresses 1 less than the integer places or figures in the number answering to that logarithm, or how far the first figure of this number is removed from the place of units; namely, that 0 is the characteristic of all numbers from 1 to 10; and 1 the characteristic

of all those from 10 to 100; and 2 the characteristic of all those from 100 to 1000; and so on.

CHARACTERS, are certain marks used by Astronomers, Mathematicians, &c, to denote certain things, whether for the sake of brevity, or perspicuity, in their operations.

#### I. ASTRONOMICAL CHARACTERS.

Planets &c.      The twelve Signs or Constellations of the Zodiac.

☉ The Sun	♈ Aries, the Ram
☾ The Moon	♉ Taurus, the Bull
⊕ The Earth	♊ Gemini, the Twins
☿ Mercury	♋ Cancer, the Crab
♀ Venus	♌ Leo, the Lion
♂ Mars	♍ Virgo, the Maid
♃ Jupiter	♎ Libra, the Balance
♄ Saturn	♏ Scorpio, the Scorpion
♁ Herschel, or the Georgian Planet	♐ Sagittary, the Archer
♊ Ascending Node	♑ Capricorn, the Goat
♋ Descending Node	♒ Aquarius, the Water-bearer
	♓ Pisces, the Fishes

The Aspects, Time, Motion, &c.

♌ Conjunction	° Degrees
♍ Opposition	' Minutes or Primes
* Sextile	" Seconds, &c.
□ Quartile	A. M. Ante merid. or in morn.
△ Trine	P. M. Post merid. or a aftern.
	h, m, s, Hours, min. sec.

#### 2. MATHEMATICAL &c. CHARACTERS.

Numeral Characters used by different Nations.

The most common numeral characters, are those called Arabic or Indian, viz. 1, 2, 3, 4, 5, 6, 7, 8, 9, with 0 or cipher for nothing.

The Roman numeral characters are seven, viz, I one, V five, X ten, L fifty, C a hundred, D or 𐀀 five hundred, M or 𐀁 or 𐀂 a thousand. Other combinations are as in the following synopsis of the Roman Notation.

1 = I	
2 = II	As often as any character is repeated, so many times its value is repeated.
3 = III	
4 = IIII or IV	A less character before a greater diminishes its value.
5 = V	
6 = VI	A less character after a greater increases its value.
7 = VII	
8 = VIII	
9 = IX	
10 = X	
50 = L	
100 = C	
500 = D or 𐀀	For every 𐀀 added, this becomes 10 times as many.
1000 = M or 𐀁	For every C and 𐀀, set one at each end, it becomes 10 times as much.
2000 = MM	
5000 = 𐀂 or 𐀃	A line over any number increases it 1000 fold.
6000 = 𐀄	
10000 = 𐀅 or 𐀆	
50000 = 𐀇	
60000 = 𐀈	
100000 = 𐀉 or 𐀊	
1000000 = 𐀋 or 𐀌	
2000000 = 𐀍, &c.	



*Greek Numerals.*

The Greeks had three ways of expressing numbers. First, The most simple was, for every single letter, according to its place in the alphabet, to denote a number from  $\alpha$  1 to  $\omega$  24; in which manner the books of Homer's Ilias are distinguished. Secondly, Another way was by dividing the alphabet into (*first*) 8 units,  $\alpha$  1,  $\beta$  2, &c; (*2nd*) 8 tens,  $\iota$  10,  $\kappa$  20, &c; (*3d*) 8 hundreds,  $\rho$  100,  $\sigma$  200, &c: And thousands they expressed by a point or accent under a letter, as  $\alpha$  1000,  $\beta$  2000, &c. Thirdly, A third way was by six capital letters, thus,  $\text{I}$  ( $\iota\alpha$  for  $\mu\iota\alpha$ ) 1,  $\text{II}$  ( $\omega\epsilon\upsilon\tau\epsilon$ ) 5,  $\Delta$  ( $\delta\epsilon\kappa\alpha$ ) 10,  $\text{H}$  ( $\text{Hex}\alpha\delta\omicron\nu$ ) 100,  $\text{X}$  ( $\chi\iota\lambda\iota\alpha$ ) 1000,  $\text{M}$  ( $\mu\upsilon\epsilon\iota\alpha$ ) 10000: and when the letter  $\text{II}$  inclosed any of these, except  $\text{I}$ , it shewed that the inclosed letter was five times its own value, as  $\overline{\Delta}$  50,  $\overline{\text{H}}$  500,  $\overline{\text{X}}$  5000,  $\overline{\text{M}}$  50000.

*Hebrew Numerals.*

The Hebrew alphabet was divided into, Nine Units, as  $\aleph$  1,  $\beth$  2, &c; Nine Tens, as  $\aleph$  10,  $\beth$  20, &c; Nine Hundreds, as  $\aleph$  100,  $\beth$  200, &c,  $\aleph$  500,  $\beth$  600,  $\aleph$  700,  $\beth$  800,  $\aleph$  900. Thousands were sometimes expressed by the units prefixed to hundreds, as  $\aleph\aleph$  1534, &c; and even to tens, as  $\aleph$  1070, &c. But more commonly thousands were expressed by the word  $\aleph$  1000,  $\aleph\aleph$  2000; and  $\aleph\aleph$  with the other numerals prefixed to signify the number of thousands, as  $\aleph\aleph$  3000, &c.

*Characters used in Arithmetic and Algebra.*

The first letters of the alphabet,  $a, b, c$ , &c, denote given quantities; and the last letters  $z, y, x$ , &c, denote such as are unknown or sought. Stifelius first used the capitals  $A, B, C$ , &c, for the unknown or required quantities. After that, Vieta used the capital vowels  $A, E, I, O, U, Y$  for the unknown or required quantities, and the consonants  $B, C, D$ , &c, for known or given numbers. Harriot changed Vieta's capitals into the small letters, viz  $a, e, i, o, u$ , for unknown, and  $b, c, d$ , &c, for known quantities. And Descartes changed Harriot's vowels for the latter letters  $z, y, x$ , &c, and the consonants for the leading letters  $a, b, c, d$ , &c.

Newton denotes the several orders of the fluxions of variable quantities by as many points over the latter letters;

as  $\dot{x}, \dot{y}, \dot{z}$  are the 1st fluxions,

$\ddot{x}, \ddot{y}, \ddot{z}$  are the 2d fluxions,

$\dddot{x}, \dddot{y}, \dddot{z}$  are the 3d fluxions,

&c, of  $x, y, z$ . And Leibnitz denotes the differentials of the same quantities by prefixing  $d$  to each of them, thus  $dx, dy, dz$ .

Powers of quantities are denoted by placing the index or exponent after them, towards the upper part; thus  $a^2$  is the 2d power,  $a^3$  the third power, and  $a^n$  the  $n$  power of  $a$ . Diophantus marked the powers by their initials, thus  $\delta', \kappa', \delta\delta', \delta\kappa', \kappa\kappa'$ , &c, for dynamis, cubus, dynamodynamis, &c, or the 2d, 3d, 4th, &c powers; and the same method has been used by several of the early writers, since the introduction of Algebra into Europe: but the first of them, as Paciolus, Cardan, &c, used no mark for powers, but the words

themselves. Stifel, and others about his time, used the initials or abbreviations,  $2, 3, \kappa, 33$ , &c, of res or cofs, zenzus, cubus, zenzizenzus, &c, barbarous corruptions of the Italian cofa, census, cubo, censi-census, &c. But he used also numeral exponents, both positive and negative, to the general characters or roots  $A, B, C$ , &c. Bombelli used a half circle thus  $\cup$  as a general character for the unknown or quantity required to be found in any question, and the several powers of it he denoted by figures set above it; thus  $\overset{1}{\cup}, \overset{2}{\cup}, \overset{3}{\cup}$ , are the 1st, 2d, 3d powers of  $\cup$ ; which powers he called dignities. Stevinus used a whole circle for the same unknown quantity, with the numeral index within it, and that both integral and fractional; thus  $\circ, \textcircled{1}, \textcircled{2}, \textcircled{3}$ , are the 0, 1, 2, 3 powers of the general quantity  $\circ$ ; also  $\textcircled{\frac{1}{2}}, \textcircled{\frac{2}{3}}, \textcircled{\frac{3}{4}}$ , he uses as the square root, cubic root, 4th root of the same; and  $\textcircled{\frac{2}{3}}$ , the cube root of the square,

and  $\textcircled{\frac{3}{4}}$ , the square root of the cube, and so on.

And these fractional exponents were adopted and farther used by his commentator Albert Girard. So that Stevinus ought to be esteemed the first person who rendered general the notation of all powers and roots in the same way, the former by integral, and the latter by fractional exponents. Harriot denoted his powers by a repetition of the letters; thus  $a, aa, aaa$ , &c. And Descartes, instead of this, set the numeral index at the upper part of the letters, as at present thus  $a, a^2, a^3$ , &c. Though, I am informed, by such as have seen Harriot's posthumous papers, that he also there makes use of exponents.

The character  $\sqrt{\phantom{x}}$  is the sign of radicality, or of a root, being derived from the initial  $R$  or  $r$ , which was used at first by Paciolus, Cardan, &c. This character  $\sqrt{\phantom{x}}$  I first find used by Stifel, in 1544, and by Robert Recorde in 1557. The character  $\sqrt{\phantom{x}}$  alone denotes the square root only; but at first they used the initial of the name after it, to denote the several roots: as  $\sqrt{q}$  the quadrature or square root, and  $\sqrt{c}$  the cubic root. But the numeral indices of the root were prefixed by Albert Girard, exactly the same as they are used at present, viz  $\sqrt[2]{\phantom{x}}, \sqrt[3]{\phantom{x}}, \sqrt[4]{\phantom{x}}$ , the 2d, 3d, or 4th root.

The character  $+$  denotes addition, and a positive quantity. At first the word itself was used, plus, piu, or the initial  $p$ . by Paciolus, Cardan, Tartalea, &c. And the character  $+$  for addition occurs in Stifelius.

The character  $-$  denotes subtraction, and a negative quantity; which also first occurs in the same author Stifelius. Before that, the word minus, mene, or the initial  $m$ . was used. Other characters have also been sometimes used by other authors, for addition and subtraction; but they are now no longer in use.

$\times$  denotes multiplication, and was introduced by Oughtred.

$\div$  denoting division, was introduced by Dr. Pell. Division is also denoted like a fraction,

$$\text{thus } \frac{a}{b} \text{ or } \frac{6}{3} = 2.$$

$=$  denotes equality, and was used by Robert Recorde. Descartes uses  $\infty$  for the same purpose.

The character  $::$  for proportionality, or equality of ratios, was introduced by Oughtred; as was also the mark  $\div$  for continued proportion.



> for greater, and < for less, were used by Harriot.

And  $\sqsupset$  and  $\sqsubset$  were used by Oughtred for the same purposes.

Dr. Pell used  $\odot$  for involution, and  $\oslash$  for evolution.

$\infty$  denotes a general difference between any two quantities, and was introduced by Dr. Wallis.

The Parenthesis ( ), as a vinculum, was invented by Albert Girard, and used in such expressions as these,  $\sqrt[3]{(72 + \sqrt{5120})}$ , and  $B(Bq + Cq)$ , both for universal roots, and multiplication, &c.

The straight-lined vinculum,  $\frac{\quad}{\quad}$ , was used by Vieta for the same purpose; thus  $A - B$  in  $B + C$ .

#### *Characters in Geometry and Trigonometry.*

$\square$ A Square	$\angle$ An Angle
$\triangle$ A Triangle	$\perp$ A Rightangle
$\square$ A Rectangle	$\perp$ Perpendicular
$\bigcirc$ or $\odot$ A Circle	$\parallel$ Parallel.

**CHARGE**, in Electricity, in a strict sense, imports the accumulation of the electric matter on one surface of an electric, as the Leyden phial, a pane of glass, &c, whilst an equal quantity passes off from the opposite surface. Or, more generally, electrics are said to be charged, when the equilibrium of the electric matter on the opposite surface is destroyed, by communicating one kind of electricity to one side, and the contrary kind to the opposite side: nor can the equilibrium be restored till a communication be made by means of conducting substances between the two opposite surfaces: and when this is done, the electric is said to be discharged. The charge properly refers to one side, in contradistinction from the other; since the whole quantity in the electric is the same before and after the operation of charging; and the operation cannot succeed, unless what is gained on one side is lost by the other, by means of conductors applied to it, and communicating either with the earth, or with a sufficient number of non-electrics. To facilitate the communication of electricity to an electric plate &c, the opposite surfaces are coated with some conducting substance, usually with tin-foil, within some distance of the edge; in consequence of which the electricity communicated to one part of the coating, is readily diffused through all parts of the surface of the electric in contact with it; and a discharge is easily made by forming a communication with any conductor from one coating to the other. If the opposite coatings approach too near each other, the electric matter forces a passage from one surface to the other before the charge is complete. And some kinds of glass have the property of conducting the electricity over the surface, so that they are altogether unfit for the operation of charging and discharging. If indeed the charge be too high, and the glass plate or phial too thin, the attraction between the two opposite electricities forces a passage through the glass, making a spontaneous discharge, and the glass becomes unfit for farther use. See *Conductors, Electrics, Leyden Phial, &c.*

**CHARGE**, in Gunnery, the load of a piece of ordnance, or the quantity of powder and ball, or shot, with which it is prepared for execution.

The charge of powder, for proving guns, is equal to the weight of the ball; but for service, the charge is  $\frac{1}{2}$  or  $\frac{1}{3}$

the weight of the ball, or still less; and indeed in most cases of service, the quantity of powder used is too great for the intended execution. In the British navy, the allowance for 32 pounders is but  $\frac{7}{12}$  of the weight of the ball. But it is probable that, if the powder in all ship guns was reduced to  $\frac{1}{2}$  the weight of the ball, or even less, it would be a considerable advantage, not only by saving ammunition, but by keeping the guns cooler and quieter, and at the same time more effectually injuring the vessels of the enemy. With the present allowance of powder, the guns are heated, and their tackle and furniture strained, and all this only to render the ball less efficacious: for a ball which can but just pass through a piece of timber, and in the passage loses almost all its motion, is found to rend and fracture it much more, than when it passes through with a much greater velocity. See Robins's Tracts, vol. 1. pa. 290, 291.

Again, the same author observes, that the charge is not to be determined by the greatest velocity that may be produced; but that it should be such a quantity of powder as will produce the least velocity necessary for the purpose in view; and if the windage be moderate, no field-piece should ever be loaded with more than  $\frac{1}{2}$ , or at the utmost  $\frac{1}{3}$  of the weight of its ball in powder; nor should the charge of any battering piece exceed  $\frac{1}{3}$  of the weight of its bullet. Ib. pa. 266.

Different charges of powder, with the same weight of ball, produce different velocities in the ball, which are in the subduplicate ratio of the weights of powder; and when the weight of powder is the same, and the ball varied, the velocity produced is in the reciprocal subduplicate ratio of the weight of the ball: which is agreeable both to theory and practice. See my paper on Gunpowder in the Philos. Transf. 1778, pa. 50; and my Tracts, vol. 1. pa. 266.

But this is on a supposition that the gun is of an indefinite length; whereas, on account of the limited length of guns, there is some variation from this law in practice, as well as in theory; in consequence of which it appears that the velocity of the ball increases with the charge only to a certain point, which is peculiar to each gun, where the velocity is the greatest; and that by farther increasing the charge, the velocity gradually diminishes, till the bore is quite full of powder. By an easy fluxionary process it appears that, calling the length of the bore of the gun  $b$ , the length of the charge pro-

ducing the greatest velocity, ought to be  $\frac{b}{2.718281828}$

or about  $\frac{3}{8}$  of the length of the bore; where 2.718281828 is the number whose hyp. log. is 1. But, for several reasons, in practice the length of the charge producing the greatest velocity, falls short of that above mentioned, and the more so as the gun is longer. From many experiments I have found the length of the charge producing the greatest velocity, in guns of various lengths of bore, from 15 to 40 calibres, as follows.

Length of Bore in Calibres.	Length of Charge for greatest Veloc.
15	$\frac{3}{10}$
20	$\frac{3}{12}$
30	$\frac{4}{16}$
40	$\frac{3}{20}$



CHARLES's WAIN, a name by which some of the astronomical writers, in our own language, have called Urfa Major, or the great bear; though some writers say the lesser bear. Indeed both of the two bears have been called waggons or wains, and by the Latins, who have followed the Arabians, two biers, *Feretrum majus & minus*.

CHART, or SEA-CHART, a hydrographical or sea-map, for the use of navigators; being a projection of some part of the sea in plano, shewing the sea coasts, rocks, sands, bearings, &c. Fournier ascribes the invention of sea-charts to Henry son of John king of Portugal. These charts are of various kinds, the Plain chart, Mercator's or Wright's chart, the Globular chart, &c.

In the construction of charts, great care should be taken that the several parts of them preserve their position to one another, in the same order as on the earth; and it is probable that the finding out of proper methods to do this, gave rise to the various modes of projection.

There are many ways of constructing maps and charts; but they depend chiefly on two principles. First, by considering the earth as a large extended flat surface; and the charts made on this supposition are usually called Plain Charts. Secondly, by considering the earth as a sphere; and the charts made on this principle are sometimes called Globular Charts, or Mercator's Charts, or Reduced Charts, or Projected Charts.

*Plain Charts* have the meridians, as well as the parallels of latitude, drawn parallel to each other, and the degrees of longitude and latitude everywhere equal to those at the equator. And therefore such charts must be deficient in several respects. For, 1st, since in reality all the meridians meet in the poles, it is absurd to represent them, especially in large charts, by parallel right lines. 2dly, As plain charts shew the degrees of the several parallels as equal to those of the equator, therefore the distances of places lying east and west, must be represented much larger than they really are. And 3dly, In a plain chart, while the same rhumb is kept, the vessel appears to sail on a great circle, which is not really the case. Yet plain charts made for a small extent, as a few degrees in length and breadth, may be tolerably exact, especially for any part within the torrid zone; and even a plain chart made for the whole of this zone will differ but little from the truth.

*Mercator's Chart*, like the plain charts, has the meridians represented by parallel right lines, and the degrees of the parallels, or longitude, everywhere equal to those at the equator, so that they are increased more and more, above their natural size, as they approach towards the pole; but then the degrees of the meridians, or of latitude, are increased in the same proportion at the same part; so that the same proportion is preserved between them as on the globe itself. This chart has its name from that of the author, Girard Mercator, who first proposed it for use in the year 1556, and made the first charts of this kind; though they were not altogether on true or exact principles, nor does it appear that he perfectly understood them. Neither indeed was the thought originally his own, viz. of lengthening the degrees of the meridian in some proportion for this was hinted by Ptolemy near two thousand year ago. It was not perfected however till Mr. Wright

first demonstrated it about the year 1590, and shewed a ready way of constructing it, by enlarging the meridian line by the continual addition of the secants. See his *Correction of Errors in Navigation*, published in 1599.

*Globular Chart*, is a projection so called from the conformity it bears to the globe itself; and was proposed by Messrs Senex, Wilson, and Harris. This is a meridional projection, in which the parallels are equidistant circles, having the pole for their common centre, and the meridians curvilinear and inclined, so as all to meet in the pole, or common centre of the parallels. By which means the several parts of the earth have their proper proportion of magnitude, distance, and situation, nearly the same as on the globe itself; which renders it a good method for geographical maps.

*Hydrographical Charts*, are sheets of large paper, on which several parts of the land and sea are described, with their respective coasts, harbours, sounds, flats, rocks, shelves, sands, &c, also the points of the compass, and the latitudes and longitudes of the places.

*Selenographic Charts*, are particular descriptions of the appearances, spots and maculæ of the moon.

*Topographic Charts*, are draughts of some small parts only of the earth, or of some particular place, without regard to its relative situation, as London, York, &c.

For the Construction of Charts, see GEOGRAPHY, MAPS, PROJECTION, &c.

CHASE, of a Gun, is its bore or cylinder.

CHAULNES (THE DUKE DE), a peer of France, but more honourable and remarkable as an astronomer and mathematician. He was born at Paris Dec. 30, 1714. He soon discovered a singular taste and genius for the sciences; and in the tumults of armies and camps, he cultivated mathematics, astronomy, mechanics, &c. He was named honorary-academician the 27th of February 1743, and few members were more punctual in attending the meetings of that body; where he often brought different constructions and corrections of instruments of astronomy, of dioptrics, and achromatic telescopes. These researches were followed with a new parallaxic machine, more solid and convenient than those that were in use; as also with many reflections on the manner of applying the micrometer to those telescopes, and of measuring exactly the value of the parts of that instrument. The duke of Chaulnes proposed many other works of the same kind, when death surprised him the 23d Sept. 1769.

He had several papers published in the volumes of *Memoirs of the Academy of Sciences*, as follow:

1. Observations on some Experiments in the 4th part of the 2d book of Newton's Optics: an. 1755.
2. Observations on the Platform for dividing mathematical instruments: 1765.
3. Determination of the distance of Arcturus from the Sun's limb, at the summer solstice: 1765.
4. On some means of perfecting astronomical instruments: 1765.
5. On some experiments relative to dioptrics: 1767.
6. The art of dividing mathematical instruments: 1768.
7. Observations of the Transit of Venus, June 3, 1769: 1769.
8. New method of dividing mathematical and astronomical instruments.



**CHAUSE TRAPPES**, or *Calltrops*, or *Crowsfeet*, are iron instruments of spikes about 4 inches long, made like a star, in such a manner that whichever way they fall, one point stands always upwards, like a nail. They are usually thrown and scattered into moats and breaches, to gall the horses feet, and stop the hasty approach of the enemy.

**HAZELLES** (JOHN MATTHEW), a French mathematician and engineer, was born at Lyons in 1657, and educated there in the college of Jesuits, from whence he removed to Paris in 1675. He first became acquainted with Du Hamel, secretary to the Academy of Sciences, and through him with Cassini, who employed him with himself at the Observatory, where Chazelles greatly improved himself, and also assisted Cassini in the measurement of the southern part of the meridian of France. Having, in 1684, instructed the duke of Montemar in the mathematical sciences, this nobleman procured him the appointment of hydrography-professor to the galleys of Marseilles. In discharging the duties of this department, he made numerous geometrical and astronomical observations, from which he drew a new map of the coast of Provence.—He also performed many other services in that department, and as an engineer, along with the armies and naval expeditions. To make observations in Geography and Astronomy, he undertook also a voyage to the Levant, and among other things he measured the pyramids of Egypt, and found the four sides of the largest of them exactly to face the four cardinal points of the compass. He made a report of his voyage, on his return, to the Academy of Sciences, upon which he was named a member of their body in 1695, and had many papers inserted in the volumes of their Memoirs, from 1693 to 1708. Chazelles died at Marseilles the 16th of January 1710.

**CHEMIN des Ronds**, in Fortification, the way of the rounds, or a space between the rampart and the low parapet under it, for the rounds to go about it.

**CHEMISE**, a wall that lines a bastion, or ditch, or the like, for its greater support and strength.

**CHERSONESUS**, a peninsula, or part of the land almost encompassed round with the sea, only joining to the main land by a narrow neck or isthmus. Varenus enumerates 14 of these.

**CHEVAL de Frise**, pl. *Chevaux de Frise*, or Frise-land horse, so called because it was first used in that country. It consists of a joist or piece of timber, about a foot in diameter, and 10 or 12 long, pierced and transversed with a great number of wooden spikes of 5 or 6 feet long, and armed or pointed with iron. It is sometimes also called *turnpike*, or *tourniquet*. It is chiefly used to stop a breach, defend a passage, or make a retrenchment to stop the cavalry.


**CHEVRE'TTE**, in Artillery, an engine to raise guns or mortars into their carriage. It is formed of two pieces of wood of about 4 feet long, standing upright upon a third, which is square. The uprights are about a foot asunder, and pierced with holes exactly opposite to one another, to receive a bolt of iron, which is put in either higher or lower at pleasure, to serve as a support to a handspike by which the gun is raised

**CHEYNE** (GEORGE), a British physician, and mathematician, was born in Scotland, 1671, and educated at Edinburgh under Dr. Pitcairn. He passed his youth in close study and great abstemiousness; but coming to London, when about 30 years of age, he suddenly changed his whole manner of living; which had such an effect upon his constitution, that his body grew to a most enormous bulk, weighing it is said about 448 pounds. From this load of oppression however he was afterward in a great measure relieved, by means of a milk and vegetable diet, which reduced his weight to about one-third of what it had been, and restored him to a good state of health; by which his life was prolonged to the 72d year of his age.

Dr. Cheyne was author of various medical and other tracts, and of a treatise on the Inverse method of Fluxions, under the title of *Fluxionum Methodus Inversa; sive quantitatum fluentium leges generaliores*: in 4to, 1703. Upon this book De Moivre wrote some animadversions in an 8vo vol. 1704; which were replied to by Cheyne in 1705.

**CHILIAD**, an assemblage of several things ranged by thousands. It was particularly applied to tables of logarithms, because they were at first divided into thousands. Thus, in the year 1624, Mr. Briggs published a table of logarithms for 20 chiliads of absolute numbers; afterward, he published 10 chiliads more; and lastly, one more; making in all 31 chiliads.

**CHILIAGON**, a regular plane figure of a thousand sides and angles.



**CHORD**, a right line connecting the two extremes of an arch; so called from its resemblance to the chord or string of a bow; as AB, which is common to the two parts or arches ADB, AEB that make up the whole circle. The chords have several properties:

1. The Chord is bisected by a perpendicular CF drawn to it from the centre.
2. Chords of equal arcs, in the same or equal circles, are themselves equal.
3. Unequal Chords have to one another a less ratio than that of their arcs.
4. The chord of an arc, is a mean proportional between the diameter and the versed sine of that arc.

*Scale or Line of Chords.* See *Plane Scale*.

**CHORD**, or **CORD**, in Music, denotes the string or line by whose vibrations the sensation of sound is excited; and by whose divisions the several degrees of tune are determined.

*To divide a Chord AB in the most simple manner, so as to exhibit all the original concords.*



Divide



Divide the given line into two equal parts at C ; then subdivide the part CB equally in two at D, and again the part CD into two equal parts at E. Here AC to AB is an octave ; AC to AD a fifth ; AD to AB a fourth ; AC to AE a greater third, and AE to AD a less third ; AE to EB a greater sixth, and AE to AB a less sixth. Malcolm's Treatise of Music, ch. 6. sec. 3. See MONOCHORD.

To find the number of Vibrations made by a Musical Chord or String in a given time ; having given its weight, length, and tension. Let  $l$  be the length of the chord in feet,  $w$  its weight, or rather a small weight fixed to the middle and equal to that of the whole chord, and  $t$  the tension, or a weight by which the chord is stretched. Then shall the time of one vibration be expressed by  $\frac{1}{2} \sqrt{\frac{l}{32 \frac{1}{2} w}}$ , and consequently the number of vibrations per second is equal to  $\frac{1}{\frac{1}{2} \sqrt{\frac{l}{32 \frac{1}{2} w}}} = \frac{1}{\frac{1}{2}} \sqrt{\frac{32 \frac{1}{2} w}{l}}$ .

For example, suppose  $w = 28800$ , or the tension equal to 28800 times the weight of the chord, and the length of it 3 feet ; then the last theorem gives 354 nearly for the number of vibrations made in each second of time.

But if  $w$  were 14400, there would be made but 250 vibrations per second ; and if  $w$  were only 288, there would be no more than  $35 \frac{1}{2}$  vibrations per second. See my Select Exerc. prob. 21. pag. 200.

CHORD, in Music, is used for the union of two or more sounds uttered at the same time, and forming together a complete harmony.

Chords are divided into perfect and imperfect. The perfect chord is composed of the fundamental sound below, of its third, its fifth, and its octave : they are likewise subdivided into major and minor, according as the thirds which enter into their composition are flat or sharp. Imperfect chords are those in which the sixth, instead of the fifth, prevails ; and in general all those whose lowest are not their fundamental sounds.

Chords are again divided into consonances and dissonances. The consonances are the perfect chord, and its derivatives. Every other chord is a dissonance. A table of both, according to the system of M. Rameau, may be seen in Rousseau's Musical Dictionary, vol. 1, pa. 27.

CHOROGRAPHY, the art of delineating or describing some particular country or province.

This differs from geography as the description of a particular country differs from that of the whole earth : And from topography, as the description of a country differs from that of a town or a district.

CHROMATIC, a species of music which proceeds by semitones and minor thirds. The word is derived from the Greek *χρῶμα*, which signifies colour, and perhaps the shade or intermediate shades of colour, which mingle and connect colours, like as the small intervals in this scale easily slide or run into each other.

Boethius and Zarlino ascribe the invention of the chromatic genus to Timotheus, a Milesian, in the time of Alexander the Great. The Spartans banished it from their city on account of its softness. The character of this genus, according to Aristides Quintilianus, was sweetness and pathos.

CHROMATICS, is that part of optics which explains the several properties of the colours of light, and of natural bodies.

Before the time of Sir I. Newton, the notions concerning colour were very vague and wild. The Pythagoreans called colour the superficies of bodies : Plato said that it was a flame issuing from them : According to Zeno, it is the first configuration of matter : And Aristotle said it was that which made bodies actually transparent. Descartes accounted colour a modification of light, and he imagined that the difference of colour proceeds from the prevalence of the direct or rotatory motion of the particles of light. Grimaldi, Dechales, and many others, imagined that the differences of colour depended upon the quick or slow vibrations of a certain elastic medium with which the universe is filled. Rohault conceived, that the different colours were made by the rays of light entering the eye at different angles with respect to the optic axis. And Dr. Hooke imagined that colour is caused by the sensation of the oblique or uneven pulse of light ; which being capable of no more than two varieties, he concluded there could be no more than two primary colours.

Sir I. Newton, in the year 1666, began to investigate this subject ; when finding that the coloured image of the sun, formed by a glass prism, was of an oblong, and not of a circular form, as, according to the laws of equal refraction, it ought to be, he conjectured that light is not homogeneous ; but that it consists of rays of different colours, and endued with divers degrees of refrangibility. And, from a farther prosecution of his experiments, he concluded that the different colours of bodies arise from their reflecting this or that kind of rays most copiously. This method of accounting for the different colours of bodies soon became generally adopted, and still continues to be the most prevailing opinion. It is hence agreed that the light of the sun, which to us seems white and perfectly homogeneous, is composed of no fewer than seven different colours, viz red, orange, yellow, green, blue, purple, and violet or indigo : that a body which appears of a red colour, has the property of reflecting the red rays more plentifully than the rest ; and so of the other colours, the orange, yellow, green, &c : also that a body which appears black, instead of reflecting, absorbs all or the most part of the rays that fall upon it ; while, on the contrary, a body which appears white, reflects the greatest part of all the rays indiscriminately, without separating them one from another.

The foundation of a rational theory of colours being thus laid, the next inquiry was, by what peculiar mechanism, in the structure of each particular body, it was fitted to reflect one kind of rays more than another ; and this is attributed, by Sir I. Newton, to the density of these bodies. Dr. Hooke had remarked, that thin transparent substances, particularly soap-water blown into bubbles, exhibited various colours, according to their thinness ; and yet, when they have a considerable degree of thickness, they appear colourless. And Sir Isaac himself had observed, that as he was compressing two prisms hard together, in order to make their sides (which happened to be a little convex) to touch one another, in the place of contact they were both



both perfectly transparent, as if they had been but one continued piece of glass: but round the point of contact, where the glasses were a little separated from each other, rings of different colours appeared. And when he afterwards, farther to elucidate this matter, employed two convex glasses of telescopes, pressing their convex sides upon one another, he observed several series of circles or rings of such colours, different, and of various intensities, according to their distance from the common central pellucid point of contact.

As the colours were thus found to vary according to the different distances between the glass plates, Sir Isaac conceived that they proceeded from the different thickness of the plate of air intercepted between the glasses; this plate of air being, by the mere circumstance of thinness or thickness, disposed to reflect or transmit the rays of this or that particular colour. Hence therefore he concluded, that the colours of all natural bodies depend on their density, or the magnitude of their component particles: and hence also he constructed a table, in which the thickness of a plate necessary to reflect any particular colour, was expressed in millionth parts of an inch.

From a great variety of such experiments, and observations upon them, our author deduced his theory of colours. And hence it seems that every substance in nature is transparent, provided it be made sufficiently thin; as gold, the densest substance we know of, when reduced into thin leaves, transmits a bluish-green light through it. If we suppose any body therefore, as gold for instance, to be divided into a vast number of plates, so thin as to be almost perfectly transparent; it is evident that all, or the greatest part of the rays, will pass through the upper plates, and when they lose their force will be reflected from the under ones. They will then have the same number of plates to pass through which they had penetrated before; and thus, according to the number of those plates through which they are obliged to pass, the object appears of this or that colour, just as the rings of colours appeared different in the experiment of the two plates, according to their distance from one another, or the thickness of the plate of air between them.

This theory of the colours has been illustrated and confirmed by various experiments, made by other philosophers. Mr. E. H. Delaval produced similar effects by the infusions of flowers of different colours, and by the intimate mixture of the metals with the substance of glass, when they are reduced to very fine parts; the more dense metals imparting to the glass the less refrangible colours, and the lighter ones those colours that are more easily refrangible. Dr. Priestley and Mr. Canton also, by laying very thin leaves or slips of the metals upon glass, ivory, wood, or metal, and passing an electrical stroke through them, found that the same effect was produced, viz, that the substratum was tinged with different colours, according to the distance from the point of explosion.

However, the Abbe Mazeas and M. du Tour contended, that the colours between the glasses are not to be ascribed to the thin stratum of air, since they equally produced them by rubbing and pressing together two flat glasses, which cohered so closely that it required

the greatest force to move or slide them over one another. See Priestley's History of Vision.

*Of Newton's 8th Exper. in the 2d Book of Optics.*

The event of this experiment, which has been contradicted by repetitions of the same by other philosophers, having been the occasion of much controversy; and relating to a material part of the doctrine of chromatics, it will not be improper here to give an account of what has passed concerning it. Newton found, he says, that when light, by contrary refractions through different mediums, is so corrected, that it emerges in lines parallel to the incident rays, it continues ever after to be white. But that if the emergent rays be inclined to the incident ones, the whiteness of the emerging light will, by degrees, in passing on from the place of emergence, become tinged at its edges with colours. And these laws he inferred from experiments made by refracting light with prisms of glass, placed within a prismatic vessel of water.

By theorems deduced from this experiment he infers, that the refraction of the rays of every sort, made out of any medium into air, are known by having the refraction of the rays of any one sort: and also, that the refraction out of one medium into another is found, whenever we have the refractions out of them both, into any third medium.

Now the same experiment, when since performed by other persons, turning out contrary to what is stated above, some rather free reflections have been thrown upon Newton concerning it; but which however have been very satisfactorily obviated by Mr. Peter Dollond, in a late pamphlet on this subject; as we shall shew below.

In the first place then, M. Klingenshierna, a Swedish philosopher, having in the year 1755 considered the controversy between Euler and Mr. John Dollond, relative to the refraction of light, formed a theorem of his own, from geometrical reasoning, by which he was induced to believe that the result of Newton's experiment could not be as he had related it; except when the angles of the refracting mediums are small. See the paper on this matter by Klingenshierna in the pamphlet above cited by Mr. Peter Dollond.

This paper of Klingenshierna being communicated to Mr. John Dollond by Mr. Mallet, to whom it was sent for that purpose, made Dollond entertain doubts concerning Newton's report of the result of his experiment, and determined him to have recourse to experiments of his own, which he did in the year 1757, as follows.

He cemented two glass planes together by their edges, so as to form a prismatic vessel when closed at the ends or bases; and the edge being turned downward, he placed it in a glass prism with one of its edges upward, filling up the vacancy with clear water; so that the refraction of the prism was contrived to be contrary to that of the water, in order that a ray of light, transmitted through both these refracting mediums, might be affected by the difference only between the two refractions. As he found the water to refract more or less than the glass prism, he diminished or augmented the angle between the glass plates, till the two contrary refractions became equal, which he discovered by view-

ing



ing an object through this double prism. For when it appeared neither raised nor depressed, he was satisfied that the refractions were equal, and that the emergent rays were parallel to the incident ones.

Now, according to the prevailing opinion, he observes, that the object ought to have appeared through this double prism in its natural colour; for if the difference of refrangibility had been in all respects equal, in the two equal refractions, they would have rectified each other. But this experiment fully proved the fallacy of the received opinion, by shewing that the divergency of the light by the glass prism, was almost double of that by the water; for the image of the object, though not at all refracted, was yet as much infected with prismatic colours, as if it had been seen through a glass wedge only, having its angle of near 30 degrees.

This experiment is the very same with that of Sir Isaac Newton above-mentioned, notwithstanding the result was so remarkably different. Mr. Dollond plainly saw however, that if the refracting angle of the water-vessel could have admitted of a sufficient increase, the divergency of the coloured rays would have been greatly diminished, or entirely rectified; and that there would have been a very great refraction without colour, as he had already produced a great discolouring without refraction: but the inconveniency of so large an angle as that of the prismatic vessel must have been, to bring the light to an equal divergency with that of the glass prism, whose angle was about 60 degrees, made it necessary to try some experiments of the same kind with smaller angles.

Accordingly he procured a wedge of plate-glass, whose angle was only 9 degrees; and, using it in the same circumstances, he increased the angle of the water-wedge, in which it was placed, till the divergency of the light by the water was equal to that by the glass; that is, till the image of the object, though considerably refracted by the excess of the refraction of the water, appeared nevertheless quite free from any colours proceeding from the different refrangibility of the light.

Many conjectures were made as to the cause of so striking a difference in the results of the same experiment; but none that gave any great satisfaction, till lately that it has been shewn to be probably owing to the nature of the glass then used by Newton. This conjecture is made by Mr. Peter Dollond, son of John, the inventor of the achromatic telescope, in a pamphlet by him lately published in defence of his father's invention, against the misrepresentations of some persons who have unjustly attempted to give the invention to other philosophers, who themselves never imagined that they had any right to it. After a full and satisfactory vindication of his father, Mr. P. Dollond then adds,

"I now come to a more agreeable part of this paper, which is, to endeavour to reconcile the different results of the 8th experiment of the 2d part of the 1st book of Newton's Optics, as related by himself, and as it was found by Dollond, when he tried the same experiment, in the year 1757. Newton says, that light, as often as by contrary refractions it is so corrected, that it emergeth in lines parallel to the incident, continues ever after to be white. Now Dollond says, when he tried the same

experiment, and made the mean refraction of the water equal to that of the glass prism, so that the light emerged in lines parallel to the incident, he found the divergency of the light by the glass prism to be nearly double to what it was by the water prism. The light appeared to be so evidently coloured, that it was directly said by some persons, that if Newton had actually tried the experiment, he must have perceived it to have been so. Yet who could for a moment doubt the veracity of such a character? Therefore different conjectures were made by different persons. Mr. Murdoch in particular gave a paper to the Royal Society in defence of Newton; but it was such as very little tended to clear up the matter. *Philos. Trans.* vol. 53. pa. 192.—Some have supposed that Newton made use of water strongly impregnated with saccharum saturni, because he mentions sometimes using such water, to increase the refraction, when he used water prisms instead of glass prisms. Newton's Opt. pa. 62.—And others have supposed, that he tried the experiment with so strong a persuasion in his own mind that the divergency of the colours was always in the same proportion to the mean refraction, in all sorts of refracting mediums, that he did not attend so much to that experiment as he ought to have done, or as he usually did. None of these suppositions having appeared at all satisfactory, I have therefore endeavoured to find out the true cause of the difference, and thereby shew, how the experiment may be made to agree with Newton's description of it, and to get rid of those doubts, which have hitherto remained to be cleared up.

"It is well known, that in Newton's time the English were not the most famous for making optical instruments: Telescopes, opera-glasses, &c, were imported from Italy in great numbers, and particularly from Venice; where they manufactured a kind of glass which was much more proper for optical purposes than any made in England at that time. The glass made at Venice was nearly of the same refractive quality as our own crown-glass, but of a much better colour, being sufficiently clear and transparent for the purpose of prisms. It is probable that Newton's prisms were made with this kind of glass; and it appears to be the more so, because he mentions the specific gravity of common glass to be to water as 2.58 to 1, Newton's Opt. pa. 247, which nearly answers to the specific gravity we find the Venetian glass generally to have. Having a very thick plate of this kind of glass, which was presented to me about 25 years ago by the late professor Allemand, of Leyden, and which he then informed me had been made many years; I cut a piece from this plate of glass to form a prism, which I conceived would be similar to those made use of by Newton himself. I have tried the Newtonian experiment with this prism, and find it answers so nearly to what Newton relates, that the difference which remains may very easily be supposed to arise from any little difference which may and does often happen in the same kind of glass made at the same place at different times. Now the glass prism made use of by Dollond to try the same experiment, was made of English flint-glass, the specific gravity of which I have never known to be less than 3.22. This difference in the densities of the prisms, used,



used by Newton and Dollond, was sufficient to cause all the difference which appeared to the two experimenters in trying the same experiment.

“ From this it appears, that Newton was accurate in this experiment as in all others, and that his not having discovered that, which was discovered by Dollond so many years afterwards, was owing entirely to accident; for if his prism had been made of glass of a greater or less density, he would certainly have then made the discovery, and refracting telescopes would not have remained so long in their original imperfect state.” See *Achromatic*, and *Telescope*.

*Mr. Delaval's experiments on the colours of opaque bodies.*—Beside the experiments of this gentleman, before-mentioned, on the colours of transparent bodies, he has lately published an account of some made upon the permanent colours of opaque substances, the discovery of which must be of the utmost consequence in the arts of colour-making and dyeing.

The changes of colour in permanently coloured bodies, our author observes, are produced by the same laws that take place in transparent colourless substances; and the experiments by which they are investigated consist chiefly of various methods of uniting the colouring particles into larger masses, or dividing them into smaller ones. Sir Isaac Newton made his experiments chiefly on transparent substances; and in the few places where he treats of others, he acknowledges his want of experiments. He makes the following remark however on those bodies which reflect one kind of light and transmit another, viz, that if these glasses or liquors were so thick and massy that no light could get through them, he questioned whether they would not, like other opaque bodies, appear of one and the same colour in all positions of the eye; though he could not yet affirm it from experience. Indeed it was the opinion of this great philosopher, that all coloured matter reflects the rays of light, some reflecting the more refrangible rays most copiously, and others those that are less so; and that this is at once the true and only reason of these colours. He was likewise of opinion that opaque bodies reflect the light from their anterior surface, by some power of the body evenly diffused over and external to it. With respect to transparent coloured bodies he thus expresses himself: “ A transparent body which looks of any colour by transmitted light, may also look of the same colour by reflected light; the light of that colour being reflected by the farther surface of that body, or by the air beyond it: and then the reflected colour will be diminished, and perhaps cease, by making the body very thick, and pitching it on the back-side to diminish the reflection of its farther surface, so that the light reflected from the tinging particles may predominate. In such cases the colour of the reflected light will be apt to vary from that of the light transmitted.”

To search out the truth of these opinions, Mr. Delaval entered upon a course of experiments with transparent coloured liquors and glasses, as well as with opaque and semitransparent bodies. And from these experiments he discovered several remarkable properties of the colouring matter; particularly, that in transparent coloured substances it does not reflect any light;

and when, by intercepting the light which was transmitted, it is hindered from passing through such substances, they do not vary from their former colour to any other, but become entirely black.

This incapacity of the colouring particles of transparent bodies to reflect light, being deduced from very numerous experiments, may therefore be taken as a general law. It will appear the more extensive, if it be considered that, for the most part, the tinging particles of liquors, or other transparent substances, are extracted from opaque bodies; that the opaque bodies owe their colours to those particles, in like manner as the transparent substances do; and that by the loss of them they are deprived of their colours.

Notwithstanding these and many other experiments, the theory of colours seems not yet determined with certainty; and it must be acknowledged that very strong objections might be brought against every hypothesis on this subject that has been invented. The discoveries of Sir Isaac Newton however are sufficient to justify the following Aphorisms.

*Aphorism 1.* All the colours in nature arise from the rays of light.

2. There are seven primary colours, namely red, orange, yellow, green, blue, indigo, and violet.

3. Every ray of light may be separated into these seven primary colours.

4. The rays of light, in passing through the same medium, have different degrees of refrangibility.

5. The difference in the colours of light arises from its different refrangibility: that which is the least refrangible producing red; and that which is the most refrangible, violet.

6. By compounding any two of the primary, as red and yellow, or yellow and blue, the intermediate colour, orange or green, may be produced.


7. The colours of bodies arise from their disposition to reflect one sort of rays, and to absorb the others: those that reflect the least refrangible rays appearing red; and those that reflect the most refrangible, violet.

8. Such bodies as reflect two or more sorts of rays, appear of various colours.

9. The whiteness of bodies arises from their disposition to reflect all the rays of light promiscuously.

10. The blackness of bodies proceeds from their incapacity to reflect any of the rays of light.—And from their thus absorbing all the rays of light that are thrown upon them, it arises, that black bodies, when exposed to the sun, become hot sooner than all others.

*Of the Diatonic Scale of Colours.*—Sir Isaac Newton, in the course of his investigations of the properties of light, discovered that the lengths of the spaces occupied in the spectrum by the seven primary colours, exactly correspond to the lengths of chords that sound the seven notes in the diatonic scale of music: which is made evident by the following experiment.





On a paper, in a dark chamber, let a ray of light be largely refracted into the spectrum ABCDEF, marking upon it the precise boundaries of the several colours, as *a*, *b*, *c*, &c; and across the spectrum draw the perpendicular lines *ag*, *bb*, &c. Then it will be found that the spaces, by which the several colours are bounded, viz, *BagF* containing the red, *abbg* containing the orange, *bcib* containing the yellow, &c, will be in exact proportion to the divisions of a musical chord for the notes of an octave; that is, as the intervals of these numbers 1,  $\frac{9}{8}$ ,  $\frac{5}{4}$ ,  $\frac{3}{2}$ ,  $\frac{4}{3}$ ,  $\frac{3}{2}$ ,  $\frac{9}{8}$ ,  $\frac{1}{2}$ .

CHRONOLOGICAL, belonging to chronology.

*Chronological Characters*, are characters by which times are distinguished. Of these, some are natural, or astronomical; others are artificial or historical. The natural characters are such as depend on the motions of the stars or luminaries; as eclipses, solstices, equinoxes, the different aspects of planets, &c. And the artificial characters are those that have been invented and established by men; as the solar cycle, the lunar cycle, &c. Historical Chronological Characters are those supported by the testimonies of historians, when they fix the dates of certain events to certain periods.

CHRONOLOGY, the art of measuring and distinguishing time; with the doctrine of dates, epochs, eras, &c.

The measurement of time in the most early periods, was by means of the seasons, or the revolutions of the sun and moon. The succession of Juno's priestesses at Argos served Hellanicus for the regulation of his narrative; while Ephorus reckoned his matters by generations. Even in the histories of Herodotus and Thucydides, there are no regular dates for the events recorded; nor were there any endeavours to establish a fixed era until the time of Ptolemy Philadelphus, who attempted it by comparing and correcting the dates of the olympiads, the kings of Sparta, and the succession of the priestesses of Juno at Argos. Eratosthenes and Apollodorus digested the events related by them, according to the succession of the olympiads and of the Spartan kings.

The chronology of the Latins is still more uncertain. The records of the Romans were destroyed by the Gauls; and Fabius Pictor, the most ancient of their historians, was obliged to borrow the chief part of his information from the Greeks. In other European nations the chronology is still more imperfect, and of a later date: and even in modern times a considerable degree of confusion and inaccuracy has arisen, from the want of attention in the historians to ascertain the dates and epochs with precision.

Hence it is evident, how necessary a proper system of chronology must be for the right understanding of history, and also how difficult it must be to establish such a system. For this purpose, however, several learned men have spent much time, particularly Julius Africanus, Eusebius of Cæsarea, George Cynceile, John of Antioch, Dennis, Petau, Clavius, Calvisius, Scaliger, Vieta, Newton, Usher, Simson, Marsham, Helvicus, Vossius, Strauchius, Blair, and Playfair.

Such a system is founded, 1st, On astronomical observations, especially of the eclipses of the sun and moon, combined with calculations of the years and eras of different nations. 2d, The testimonies of cre-

dible authors. 3d, Such epochs in history as are so well attested and determined, that they have never been controverted. 4th, Ancient medals, coins, monuments, and inscriptions.

The most obvious division of time, as has been observed, is derived from the apparent or real revolutions of the luminaries, the sun and moon. Thus, the apparent revolution of the sun, or the real rotation of the earth on her axis causing the sun to appear to rise and set, constitutes the vicissitudes of day and night, which must be evident to the most barbarous and ignorant nations. The moon, by her revolution about the earth, and her changes, as naturally and obviously forms months; while the great annual course of the sun through the several constellations of the zodiac, points out the larger division of the year.

The Day is divided into hours, minutes, &c; while the month is divided into weeks, and the year into months, having particular names, and a certain number of days.—See a particular account of each of these under the respective words.

Beside the natural divisions of time arising immediately from the revolutions of the heavenly bodies, there are others which are formed from some of the less obvious consequences of these revolutions, and are called *cycles*, or *circles*. The most remarkable of these are, 1, The *Solar Cycle*, or cycle of the sun, a period or revolution of 28 years, in which time the days of the months return to the same days of the week, the sun's place to the same signs and degrees of the ecliptic on the same months and days, and the leap-years begin the same course over again with respect to the days of the week on which the days of the months fall. 2, The *Lunar Cycle*, or cycle of the moon, commonly called the *Golden Number*, is a revolution of 19 years; in which time the conjunctions, oppositions, and other aspects of the moon, are on the same days of the months as they were 19 years before, and within an hour and a half of the same time of the day.

The *Indiction*, or Roman Indiction, is a period of 15 years, used only by the Romans for indicating the times of certain payments made by the subjects to the republic.

The *Cycle of Easter*, called also the *Dionysian Period*, is a revolution of 532 years, and is produced by multiplying the solar cycle 28, by the lunar cycle 19.

The *Julian Period*, is a revolution of 7980 years, and is produced from the continual multiplication of the three numbers 28, 19, 15, of the three former cycles, viz, the solar, lunar, and indiction.

As there are certain fixed points in the heavens, from which astronomers begin their computations, so there are certain points of time, from which historians begin to reckon; and these points or roots of time are called *eras* or *epochs*. The most remarkable of these are, those of the Creation, the Greek Olympiads, the building of Rome, the era of Nabonnassar, the death of Alexander, the birth of Christ, the Arabian Hegira, or flight of Mahomet, and the Persian Jeshgird. All which, with some others of less note, have their beginnings fixed by chronologers to the years of the Julian period, to the age of the world, and to the years before and after the birth of Christ.

The testimony of authors is the second principal  
O o part



part of historical chronology. Though no man has a right to be considered as infallible, it would however be making a very unfair judgment of mankind, to treat them all as dupes or impostors; and it would be an injury offered to public integrity, to doubt the veracity of authors universally esteemed, and facts that are truly worthy of belief. When the historian is allowed to be completely able to judge of an event, and to have no intent of deceiving by his relation, his testimony cannot be refused.

The *Epochs* form the 3d principal part of chronology; being those fixed points in history that have never been contested, and of which there cannot reasonably be any doubt. Notwithstanding that chronologers fix upon the events which are to serve as epochs, in a manner quite arbitrary; yet this is of little consequence, provided the dates of these epochs agree, and that there is no contradiction in the facts themselves.

Medals, Monuments, and Inscriptions, form the last of the four principal parts of chronology; and this study is but of very modern date, scarce more than 150 years having elapsed since close application has been made to the study of these. To the celebrated Spanheim we owe the greatest obligations, for the progress that is made in this method; and it is by the aid of medals that M. Vaillant has composed his judicious history of the kings of Syria, from the time of Alexander the Great to that of Pompey. Nor have they been of less service in elucidating all ancient history, especially that of the Romans; and even sometimes that of the middle ages.

Besides the foregoing general account, there are some few systems of chronology which may deserve some more particular notice, as follows.

*Sacred Chronology.* There have been various systems relating to sacred chronology; which is not to be wondered at, as the three chief copies of the Bible give a very different account of the first ages of the world. For while the Hebrew text reckons about 4000 years from the creation to the birth of Christ, and to the flood 1656 years; the Samaritan makes the former much longer, though it counts from the creation to the flood only 1307 years; and the Septuagint removes the creation of the world to 6000 years before Christ, and 2250 years before the flood. Many attempts have been made to reconcile these differences; though none of them are quite satisfactory. Walton and Vossius give the preference to the account of the Septuagint; while others have defended the Hebrew text. See an abstract of the different opinions of learned men on this subject, in Strauchius's *Brev. Chron.* translated by Sault, p. 166 and 176.

*The Chinese Chronology.* No nation has boasted more of its antiquity than the Chinese: but though they be allowed to trace their origin as far back as the deluge, they have few or no authentic records of their history for so long a period as 500 years before the Christian era. This indeed may be owing to the general destruction of ancient remains by the tyrant Tsin-chi-hoang, in the year 213, or some say 246, before Christ. From a chronology of the Chinese history (for which we are obliged to an illustrious Tartar who was viceroy of Canton in the year 1724,

and of which a Latin translation was published at Rome in 1730), we learn that the most remote epoch of the Chinese chronology does not surpass the first year of Guai-lie-wang, or 424 years before our vulgar era. And this opinion is confirmed by the practice of two of the most approved historians of China, who admit nothing into their histories previous to this period.

The Chinese, in their computation, make use of a cycle of 60 years, called *kia-tse*, from the name given to the first year of it, which serves as the basis of their whole chronology. Every year of this cycle is marked with two letters which distinguish it from the others; and all the years of the emperors, for upwards of 2000 years, have names in history common to them with the corresponding years of the cycle. *Philos. Trans. Abridg.* vol. 8, part 4, pag. 13.

According to M. Freret, in his *Essays*, the Chinese date the epocha of Yao, one of their first emperors, about the year 2145, or 2057, before Christ; and they reckon that their first astronomical observations, and the composition of their calendar, preceded Yao 150 years: from whence it is inferred that the era of their astronomical observations coincides with that of the Chaldeans. But later authors date the rise and progress of the sciences in China from the grand dynasty of Tcheou, about 1200 years before the Christian era, and shew that all historical relations of events prior to the reign of Yao are fabulous. *Mem. de l'Histoire des Sciences &c. Chinois.* vol. 1, Paris 1776.

*Babylonian, Egyptian, and Chaldean Annals.* These, M. Gibert has attempted to reduce to our chronology, in a letter published at Amsterdam in 1743. He begins with shewing, by the authorities of Macrobius, Eudoxus, Varro, Diodorus Siculus, Pliny, Plutarch, St. Augustin, &c, that by a year, the ancients meant the revolution of any planet in the heavens; so that it might consist sometimes of only one day. Thus, according to him, the solar day was the astronomical year of the Chaldeans; and so the boasted period of 473,000 years, assigned to their observations, is reduced to 1297 years 9 months; the number of years which, according to Eusebius, elapsed from the first discoveries of Atlas in astronomy, in the 384th year of Abraham, to the march of Alexander into Asia in the year 1682 of the same era. And the 17,000 years added by Berofus to the observations of the Chaldeans, reduced in the same manner, will give 46 years and 6 or 7 months; being the exact interval between Alexander's march, and the first year of the 123d Olympiad, or the time to which Berofus carried his history.

Epigenius ascribes 720,000 years to the observations preserved at Babylon; but these, according to M. Gibert's system, amount only to 1971 years 3 months; which differ from Callisthenes's period of 1903 years, allotted to the same observations, only by 68 years, the period elapsed from the taking of Babylon by Alexander, which terminated the latter account, and to the time of Ptolomy Philadelphus, to which Epigenius extended his account.

*The Newtonian Principles of Chronology.*—Sir Isaac Newton has shewn, that the chronology of ancient kingdoms is involved in the greatest uncertainty: that



the Europeans in particular had no chronology before the Persian empire, which commenced 536 years before the birth of Christ, when Cyrus conquered Darius the Mede: that the antiquities of the Greeks are full of fables, because their writings were in verse only, till the conquest of Asia by Cyrus the Persian; about which time prose was introduced by Pherecides Syrius and Cadmus Milefius. After this time several of the Greek historians introduced the computation by generations. The chronology of the Latins was still more uncertain: their old records having been burnt by the Gauls 120 years after the expulsion of their kings, or 64 years before the death of Alexander the Great, answering to 388 before the birth of Christ. The chronologers of Gaul, Spain, Germany, Scythia, Sweden, Britain, and Ireland, are of a still later date. For Scythia, beyond the Danube, had no letters till Ulphilas, their bishop, formed them, about the year 276. Germany had none till it received them from the western empire of the Latins, about the year 400. The Huns had none in the days of Procopius, about the year 526. And Sweden and Norway received them still later.

Sir Isaac Newton, after a general account of the obscurity and defects of the ancient chronology, observes that, though many of the ancients computed by successions and generations, yet the Egyptians, Greeks, and Latins, reckoned the reigns of kings equal to generations of men, and three of them to a hundred, and sometimes to 120 years; and this was the foundation of their technical chronology. He then proceeds, from the ordinary course of nature, and a detail of historical facts, to shew the difference between reigns and generations; and that, though a generation from father to son may at an average be reckoned about 33 years, or three of them equal to 100 years, yet, when they are taken by the eldest sons, three of them cannot be estimated at more than about 75 or 80 years; and the reigns of kings are still shorter; so that 18 or 20 years may be allowed a just medium. Sir Isaac then fixes on four remarkable periods, viz, the return of the Heraclidæ into the Peloponnesus, the taking of Troy, the Argonautic expedition, and the return of Sesostris into Egypt, after his wars in Thrace; and he settles the epoch of each by the true value of a generation. To instance only his estimate of that of the Argonautic expedition: Having fixed the return of the Heraclidæ to about the 159th year after the death of Solomon, and the destruction of Troy to about the 76th year after the same period, he observes, that Hercules the Argonaut was the father of Hyllus, the father of Clerdius, the father of Aristomachus, the father of Aristodemus, who conducted the Heraclidæ into Peloponnesus; so that, reckoning by the chief of the family, their return was four generations later than the Argonautic expedition, which therefore happened about 43 years after the death of Solomon. This is farther confirmed by another argument: Æsculapius and Hercules were Argonauts: Hippocrates was the 18th inclusively from the former by the father's side, and the 19th from the latter by the mother's side: now, allowing 28 or 30 years to each of them, the 17 intervals by the father, and the 18 intervals by the mo-

ther, will on a medium give 507 years; and these, reckoning back from the commencement of the Peloponnesian war, or the 431st year before Christ, when Hippocrates began to flourish, will place the Argonautic expedition in the 43d year after the death of Solomon, or 937 years before Christ.

The other kind of reasoning by which Newton endeavours to establish this epoch, is purely astronomical. The sphere was formed by Chiron and Musæus for the use of the Argonautic expedition, as is plainly shewn by several of the asterisms referring to that event: and at the time of the expedition the cardinal points of the equinoxes and solstices were placed in the middle of the constellations Aries, Cancer, Chelæ, and Capricorn. This point is established by Newton from the consideration of the ancient Greek calendar, which consisted of 12 lunar months, and each month of 30 days, which required an intercalary month. Of course this lunisolar year, with the intercalary month, began sometimes a week or two before or after the equinox or solstice; and hence the first astronomers were led to the before-mentioned disposition of the equinoxes and solstices: and that this was really the case, is confirmed by the testimonies of Eudoxus, Aratus, and Hipparchus. Upon these principles Sir Isaac proceeds to argue in the following manner. The equinoctial colure in the end of the year 1689 cut the ecliptic in  $8^{\circ} 44'$ ; and by this reckoning the equinox had then gone back  $36^{\circ} 44'$  since the time of the Argonautic expedition. But it recedes  $50'$  in a year, or  $1^{\circ}$  in 72 years, and consequently  $36^{\circ} 44'$  in 2645 years; and this, counted backwards from the beginning of 1690, will place this expedition about 25 years after the death of Solomon. But as there is no necessity for allowing that the middle of the constellations, according to the general account of the ancients, should be precisely the middle between the prima Arietis and ultima Caudæ, our author proceeds to "examine what were those stars through which Eudoxus made the colures to pass in the primitive sphere, and in this way to fix the position of the cardinal points." Now from the mean of five places he finds, that the great circle, which in the primitive sphere, described by Eudoxus, or which at the time of the Argonautic expedition was the equinoctial colure, did in the end of 1689 cut the ecliptic in  $8^{\circ} 29' 15''$ . In the same manner our author determines that the mean place of the solstitial colure is  $\Omega 6^{\circ} 28' 46''$ , and as it is at right angles with the other, he concludes that it is rightly drawn. And hence he infers that the cardinal points, in the interval between that expedition and the year 1689, have receded from those colures  $1^{\circ} 6' 29'$ ; which, allowing 72 years to a degree, amounts to 2627 years; and these counted backwards, as above, will place the Argonautic expedition 43 years after the death of Solomon. Our author has, by other methods also of a similar nature, established this epoch, and reduced the age of the world 500 years.

This elaborate system has not escaped censure; Mess. Freret and Souciet having both attacked it, and on much the same ground: but the former has confounded reigns and generations, which are carefully distinguished in this system. The astronomical objections of both



have been answered by Sir Isaac Newton himself, and by Dr. Halley. *Philos. Trans. abr.* vol. 8, part 4, pa. 4. *Newton's Chronol.* ch. 1.

**CHRONOMETER**, is any instrument or machine used in measuring time; such as dials, clocks, watches, &c.

The term is however more particularly used for a kind of clock, so contrived as to measure a small portion of time, even to the 16th, or the 40th part of a second; one of this latter kind I have seen, made by an ingenious artist; but it could not be stopped to the 10th part of the proposed degree of accuracy. There is a description of one also in Desaguliers's *Experimental Philosophy*, invented by the late ingenious Mr. George Graham; which might be of great use for measuring small portions of time in astronomical observations, the time of the fall of bodies, the velocity of running waters, &c. But long intervals of time cannot be measured by it with sufficient exactness, unless its pendulum be made to vibrate in a cycloid; for otherwise it is liable to err considerably, as is the case of all clocks with short pendulums that swing in large arches of the circle.

Various other contrivances, besides clocks, have been used for measuring time for some particular purposes. See a musical chronometer described in Malcolm's *Treatise of Music*, pa. 407.

**CHRONOSCOPE**, a word sometimes used for a pendulum, or machine, to measure time.

**CHRYSTALLINE**. See **CRYSTALLINE**.

**CIMA**, or **SIMA**, in Architecture, a member, or moulding, called also ogee, and cimatum.

**CINCTURE**, in Architecture, is a ring or list around the shaft of a column, at its top and bottom.

**CINTRE**, in Building, the mould on which an arch is turned or built; popularly called centre, and sometimes a cradle.

**CIPHER**, or **Cypher**, one of the numeral characters, or figures, thus formed 0. The word comes from the Hebrew *saphar*, to number.

The cipher of itself signifies nothing, or implies a privation of value; but when combined with other numeral characters, it alters their value in a tenfold proportion, for every cipher so annexed; viz, when set after a figure in common integral arithmetic, it increases its value in that proportion, though it has no effect when set before or to the left hand side of figures; but on the contrary, in decimal arithmetic, it decreases their value in that proportion when set before the figures, but has no effect when set after them.

Thus, 5 is five,

but 50 is fifty,

and 500 is five hundred;

whereas 05, or 005, &c, is still but 5 or five.

Also .5 is five tenths,

but .05 is five hundredths,

and .005 is five thousandths;

whereas .50, or .500, &c, is still but .5 or five tenths.

The invention and use of the cipher, as in the common arithmetic and notation of numbers, is one of the happiest devices that can be imagined; and is ascribed to the Indians, by the Arabians, through whom it came into Europe, along with the revival of literature.

**CIRCLE**, a plane figure, bounded by a curve line

which returns into itself, called its circumference, and which is every where equally distant from a point within, called its centre.

The circumference or periphery itself is called the circle, though improperly, as that name denotes the space contained within the circumference.

A circle is described with a pair of compasses, fixing one foot in the centre, and turning the other round to trace out the circumference.

The circumference of every circle is supposed to be divided into 360 equal parts, called degrees, and marked °; each degree into 60 minutes or primes, marked ′; each minute into 60 seconds, marked ″; and so on. So 24° 12′ 15″ 20″, is 24 degrees 12 minutes 15 seconds and 20 thirds.

Circles have many curious properties, some of the most important of which are these:

1. The circle is the most capacious of all plain figures, or contains the greatest area within the same perimeter, or has the least perimeter about the same area; being the limit and last of all regular polygons, having the number of its sides infinite.

2. The area of a circle is always less than the area of any regular polygon circumscribed about it, and its circumference always less than the perimeter of the polygon. But on the other hand, its area is always greater than that of its inscribed polygon, and its circumference greater than the perimeter of the said inscribed polygon. However, the area and perimeter of the circle approach always nearer and nearer to those of the two polygons, as the number of the sides of these is greater; the circle being always limited between the two polygons.

3. The area of a circle is equal to that of a triangle whose base is equal to the circumference, and perpendicular equal to the radius. And therefore the area of the circle is found by drawing half the circumference into half the diameter, or the whole circumference into the whole diameter, and taking the 4th part of the product. Demonstrated by Euclid.

4. Circles, like other similar plane figures, are to one another, as the squares of their diameters. And the area of the circle is to the square of the diameter, as 11 to 14 nearly, as proved by Archimedes; or as 7854 to 1 more nearly; or still more nearly as

7853981633, 9744830961, 5660845819, 8757210492, 9234984377, 6455243736, 1480769541, 0157155224, 9657008706, 3355292669, 9553702162, 8318076661, 7734611 + to 1;

as it has been found by modern mathematicians.

In Wallis's *Arithmetic of Infinites* are contained the first infinite series for expressing the ratio of a circle to the square of its diameter: viz,

1st, The circle is to the square of its diameter, as 1 to  $\frac{3 \times 3 \times 5 \times 5 \times 7 \times 7 \times \&c}{2 \times 4 \times 4 \times 6 \times 6 \times 8 \times \&c}$ , found out by Wallis himself;

or 1 to  $\frac{9}{8} \times \frac{25}{24} \times \frac{49}{48} \&c$ ,

or as 1 to  $1 + \frac{1}{2 + \frac{9}{2 + \frac{25}{2 + \frac{49}{2 + \&c}}}}$ , by Ld. Brounker;



$$\text{or as } 1 - \frac{1}{2 \times 3} - \frac{1}{2 \times 4 \times 5} - \frac{1 \times 3}{2 \times 4 \times 6 \times 7} - \frac{1 \times 3 \times 5}{2 \times 4 \times 6 \times 8 \times 9} \&c \text{ to } 1, \text{ by Sir I. Newton;}$$

or as  $1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} - \frac{1}{11} \&c$  to 1, by Gregory and Leibnitz; and a great many other forms of series have been invented by different authors, to express the same ratio between the circle and circumscribed square.

5. The circumferences of circles are to one another, as their diameters, or radii. And as the areas of circles are proportional to the rectangles of their radii and circumferences; therefore the quadrature of the circle will be effected by the rectification of its circumference; that is, if the true length of the circumference could be found, the true area could be found also. But whilst several mathematicians have endeavoured to determine the true area and circumference, others have even doubted of the possibility of the same. Of this latter opinion is Dr. Isaac Barrow: towards the end of his 15th Mathematical-Lecture he says, he is of opinion, that the radius and circumference of a circle, are lines of such a nature, as to be not only incommensurable in length and square, but even in length, square, cube, bi-quadrato, and all other powers to infinity: for, continues he, the side of the inscribed square is incommensurable to the radius, and the square of the side of the inscribed octagon is incommensurable to the square of the radius; and consequently the square of the octagonal perimeter is incommensurable to the square of the radius: and thus the ambits of all regular polygons, inscribed in a circle, may have their superior powers incommensurate with the co-ordinate powers of the radius; from whence the last polygon, that is, the circle itself, seems to have its periphery incommensurate with the radius. Which, if true, will put a final stop to the quadrature of the circle, since the ratio of the circumference to the radius is altogether inexplicable from the nature of the thing, and consequently the problem requiring the explication of such a ratio is impossible to be solved, or rather it requires that for its solution which is impossible to be apprehended. But, concludes he, this great mystery cannot be explained in a few words: But if time and opportunity had permitted, I would have endeavoured to produce many things for the explication and confirmation of this conjecture. Sir Isaac Newton too, in book 1 of his Principia, has attempted to demonstrate the impossibility of the general quadrature of oval figures, by the description of a spiral, and the impossibility of determining, by a finite equation, the intersections of that oval and spiral, which must be the case, if the oval be quadrable. And several other authors have attempted to demonstrate the impossibility of the general quadrature of the circle by any means whatever. On the other hand, many authors not only believe in the possibility of the quadrature of the circle, but some have even pretended to have discovered the same, and have published to the world their pretended discoveries: of which no one has rendered himself more remarkable than our countryman Mr. Hobbes, though a great scholar, and of excellent understanding in other matters. See QUADRATURE.

The approximate quadrature of the circle however, or

the determination of the ratio between the diameter and the circumference, is what the mathematicians of all ages have successfully attempted, and with different degrees of accuracy, according to the improved state of the science. Archimedes, in his book *de Dimensione Circuli*, first gave a near value of that ratio in small numbers, being that of 7 to 22, which are still used as very convenient numbers for this purpose in common measurements. Other and nearer ratios have since been successively assigned, but in larger numbers,

as 106 to 333,  
or 113 to 355,  
or 1702 to 5347,  
or 1815 to 5702, &c,

which are each more accurate than the foregoing. Vieta, in his *Universalium Inspectionum ad Canonum Mathematicum*, published 1579, by means of the inscribed and circumscribed polygons of 393216 sides, has carried the ratio to ten places of figures, shewing that if the diameter of a circle be 1000 &c, the circumference will be greater than 314,159,265,35, -

but less than 314,159,265,37.

And Van Colen, in his book *de Circulo & Adscriptis*, has, by the same means, carried that ratio to 36 places of figures; which were also recomputed and confirmed by Willibrord Snell. After these, that indefatigable computer, Mr. Abraham Sharp, extended the ratio to 72 places of figures, in a sheet of paper, published about the year 1706, by means of the series of Dr. Halley, from the tangent of an arc of 30 degrees. And the ingenious Mr. Machin carried the same to a hundred places, by other series, depending on the differences of arcs whose tangents have certain relations to one another. See this method explained in my *Mensuration*, pa. 120 second edit. And, finally, M. De Lagny, in the *Memoirs de l'Acad.* 1719, by means of the tangent of the arc of 30 degrees, has extended the same ratio to the amazing length of 128 places of figures; finding, that, if the diameter be 1000 &c, the circumference will be

31415,92653,58979,32384,62643,38327,95028,  
84197,16939,93751,05820,97494,45923,07816,  
40628,62089,98628,03482,53421,17067,98214,  
80865,13272,30664,70938,446 + or 447 -

From such methods as the foregoing, a variety of series have been discovered for the length of the circumference of a circle, such as the following, viz, If the diameter be 1, the circumference  $c$  will be variously expressed thus,

$$c = 4 \times \left( 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} - \frac{1}{11} + \frac{1}{13} - \frac{1}{15} \&c \right),$$

$$c = \sqrt{8} \times \left( 1 + \frac{1}{3} - \frac{1}{5} - \frac{1}{7} + \frac{1}{9} + \frac{1}{11} - \frac{1}{13} - \frac{1}{15} \&c \right),$$

$$c = \sqrt{12} \times \left( 1 - \frac{1}{3 \cdot 3} + \frac{1}{5 \cdot 3^2} - \frac{1}{7 \cdot 3^3} + \frac{1}{9 \cdot 3^4} \&c \right),$$

$$c = 8 \times \left( \frac{1}{1 \cdot 1 \cdot 3} + \frac{1}{1 \cdot 3 \cdot 5} - \frac{1}{3 \cdot 5 \cdot 7} + \frac{1}{5 \cdot 7 \cdot 9} - \frac{1}{7 \cdot 9 \cdot 11} \&c \right),$$

$$c = 8 \times \left( \frac{2}{3} - \frac{1}{5} - \frac{1}{4 \cdot 7} - \frac{1 \cdot 3}{4 \cdot 6 \cdot 9} - \frac{1 \cdot 3 \cdot 5}{4 \cdot 6 \cdot 8 \cdot 11} \&c \right),$$

$$c = 4\sqrt{2} \times \left( \frac{2}{3} - \frac{1}{5 \cdot 2} - \frac{1}{4 \cdot 7 \cdot 2^2} - \frac{1 \cdot 3}{4 \cdot 6 \cdot 9 \cdot 2^3} - \frac{1 \cdot 3 \cdot 5}{4 \cdot 6 \cdot 8 \cdot 11 \cdot 2^4} \&c \right),$$

$$c = 4 \times \left( 1 - \frac{1}{2 \cdot 3} - \frac{1}{2 \cdot 4 \cdot 5} - \frac{1 \cdot 3}{2 \cdot 4 \cdot 6 \cdot 7} - \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6 \cdot 8 \cdot 9} \&c \right).$$

And

*John King*  
12. 41



And many other series might here be added. See my Menfuration in feveral places; alfo my paper on fuch series in the Philof. Tranf. 1776; Euler's *Introductio in Analyfin Infinitorum*; and many other authors.

6. Some of the more remarkable properties of the circle, are as follow.

If two lines AB, CD cut the circle, and interfect within it, the angle of interfection E is meafured by half the *sum* of the intercepted arcs AC, DB.

But if the lines interfect without the circle, the angle E is meafured by half the *difference* of the intercepted arcs AC, DB.

7. The angle at the centre of a circle is double the angle at the circumference, ftanding on the fame arc; and all angles in the fame fegment are equal. Alfo the angle at the centre is meafured by the arc it ftands upon, and the angle at the circumference by half the fame arc.

8. If the chords FG, HI crofs at right angles, the fums of the oppofite arcs are equal; viz  $FH + GI = FI + GH$ .

9. If one fide NO of a trapezium infcribed in a circle be continued out, the external angle, LOP will be equal to the oppofite internal angle M.

10. An angle, as RQS, formed by a tangent QR and chord QS, is meafured by half the arc of the chord QS, and is equal to any angle T formed in the oppofite arc QTS.

11. If VW be a diameter, and XYZ a chord perpendicular to it; then is XZ or ZY a mean proportional between the fegments YZ, ZW. So that if  $d$  denote the diameter VW,  $x$  the abfcifs VZ, and  $y$  the ordinate ZX; then is  $y^2 = d - x \times x$  or  $= dx - x^2$ ; which is called the equation to the circle.

The chord VX is a mean proportional between the diameter VW and the verfed fine VZ; and the chord WX is a mean proportional between the diameter and the verfed fine WZ; alfo each verfed fine is proportional to the fquare of the corresponding chord; viz  $VZ : WZ :: VX^2 : WX^2$ .

12. When two lines cut the circle, whether they interfect within the circle, or without it, as in the two figures to article 6, the fegments between the common interfection and the two points where each line cuts the curve, are reciprocally proportional, and their rectan-

gles are equal; viz,  $EA : EC :: ED : EB$ ,  
or  $EA \times EB = EC \times ED$ .

13. In a trapezium infcribed in a circle, the rectangle of the two diagonals is equal to the fum of the two rectangles of the two pairs of oppofite fides; viz,  $AC \times BD = AB \times DC + AD \times BC$ .

14. If any chords EF, EG, drawn from the fame point E in the circumference, be cut by any other line HI, the rectangles will be all equal which are made of each chord and the part intercepted by this line, viz,  $EF \times EI = EG \times EH = EK^2$ .













15. In a circle whose centre is N and radius NO, if two points M, P, in the radius produced, be fo placed that the three NM, NO, MP, be in continued proportion; then if from the points M and P lines be drawn to any, or every point in the circumference, as Q; these lines will be always in the given ratio of MO to PO; viz,  $MQ : PQ :: MO : PO$ .

16. If VW, be two points in the diameter, equidiftant from the centre T; and if two lines be drawn from these to any point X in the circumference; the fum of their fquares will be equal to the fum of the fquares of the fegments of the diameter made by either point; viz,  $VX^2 + WX^2 = RV^2 + VS^2$ , or  $= RW^2 + WS^2$ , or  $= 2RT^2 + 2VT^2$ .

17. If a line FE perpendicular to the diameter AB, meet any other chord CD in the point E; then is  $AF \times FB = CE \times ED + EF^2$ .


18. If upon the diameter GH of a circle there be formed a rectangle GHKI, whose breadth GI or HK is equal to GL or HL, the chord of a quadrant, or fide of the infcribed fquare; then if from I and K lines be drawn to any point M in the circle GMH, they will cut the diameter GH in fuch a manner that  $GO^2 + HN^2 = GH^2$ .

19. If the arcs PQ, QR, RS, &c, be equal, and there be drawn the chords PQ, PR, PS, PT, &c, then it will







will be  $PQ : PR :: PR : PQ + PS :: PS : PR + PT :: PT : PS + PV$ , &c.



20. The centre of a circle being  $O$ , and  $P$  a point in the radius, or in the radius produced; if the circumference be divided into as many equal parts  $AB, BC, CD$ , &c, as there are units in  $2n$ , and lines be drawn from  $P$  to all the points of division; then shall the continual product of all the alternate lines viz  $PA \times PC \times PE$  &c be  $= r^n - x^n$  when  $P$  is within the circle, or  $= x^n - r^n$  when  $P$  is without the circle; and the product of the rest of the lines, viz  $PB \times PD + PF$  &c  $= r^n + x^n$ : where  $r = AO$  the radius, and  $x = OP$  the distance of  $P$  from the centre.



21. A circle may thus be divided into any number of parts that shall be equal to one another both in area and perimeter. Divide the diameter  $QR$  into the same number of equal parts at the points  $S, T, V$ , &c; then, on one side of the diameter describe semicircles on the diameters  $QS, QT, QV$ , and on the other side of it describe semicircles on  $RV, RT, RS$ ; so shall the parts 1 7, 3 5, 5 3, 7 1 be all equal, both in area and perimeter. See my Tracts, pa. 93.



22. To describe a Circle either about or within a given Regular Polygon. Bisect two of its angles, or two of its sides, with perpendiculars, and the intersection of the bisecting lines will, in either case, be the centre of the circles.

Parallel, or Concentric CIRCLES, are such as are equally distant from each other in every point of their peripheries; or that have the same centre. As, on the other hand, those are called the *eccentric* circles, that have not the same point for their centres.

The *Quadrature of the CIRCLE*, is the manner of describing, or assigning, a square, whose surface shall be perfectly equal to that of a circle. This problem has exercised the geometricians of all ages, but it is now generally given up as a problem impossible to be effected,


by most persons that have any just claim to that rank. Des Cartes insists on the impossibility of it, for this reason, that a right line and a circle being of different natures, there can be no strict proportion between them. Dr. Barrow shews the strongest probability of the same thing; and not only that the diameter and circumference themselves, but that all powers of them to infinity, are incommensurate.

The Emperor Charles V offered a reward of 100,000 crowns to any person who should resolve this celebrated problem: and the States of Holland also proposed a reward for the same thing. See *Quadrature*.

CIRCLES of the Higher Orders, are curves in which

$$WY^m : YZ^m :: YZ : YX,$$

$$\text{or } WY^m : YZ^m :: YZ^n : YX^n.$$



When  $m$  and  $n$  are each equal to 1, then  $WY : YZ :: YZ : YX$ , which is the property of the common circle.


Put  $WY = x$ ,  $YZ = y$ ,  $WX = a$ ; then is  $YX = a - x$ , and the above proportions become

$$x^m : y^m :: y : a - x, \text{ or } y^{m+1} = x^m \cdot a - x^m, \text{ and}$$

$$x^m : y^m :: y^n : a - x, \text{ or } y^{m+n} = x^m \cdot a - x^m, \text{ the}$$

equations to curves of this kind.

Curves defined by this equation will be ovals when  $m$  is an odd number. Thus suppose  $m = 1$ , then the equation becomes  $y^2 = x \cdot a - x^2$  or  $ax - x^2$ , the equation to the common circle. And if  $m = 3$ , it becomes  $y^4 = x^3 \cdot a - x^3$  or  $ax^3 - x^4$ , which denotes a curve of this form AB.



But when  $m$  denotes an even number, the curve will have two infinite legs. So if  $m = 2$ , the equation becomes  $y^3 = x^2 \cdot a - x^2$  or  $ax^2 - x^3$ , for a circle of the 2d order, and which defines one of Newton's defective hyperbolas, being his 37th species of curves, whose asymptote is the right line EF, making an angle of 40 degrees with the absciss AB.

CIRCLE of Curvature, or circle of equi-curvature, is that circle which has the same curvature with a given curve at a certain point; or that circle whose radius is equal to the radius of curvature of the given curve at that point.

CIRCLES of the Sphere, are such as cut the mundane sphere, and have their circumference in its surface.

These circles are either fixed or moveable.

The latter are those whose peripheries are in the moveable



moveable or revolving surface; and which therefore move or turn with it; as the meridians, &c. The former, having their periphery in the immoveable surface, do not revolve; as the ecliptic, equator, and its parallels.

The circles of the sphere are either great or little.

*A Great Circle* of the Sphere, is that which divides it into two equal parts or hemispheres, having the same centre and diameter with it. As the horizon, meridian, equator, ecliptic, the colures, and the azimuths.

*A Little, or Lesser Circle* of the Sphere, divides the sphere into two unequal parts, having neither the same centre nor diameter with the sphere; its diameter being only some chord of the sphere less than its axis. Such as the parallels of latitude, &c.

*CIRCLES of Altitude, or Almucantars*, are little circles parallel to the horizon, having their common pole in the zenith, and still diminishing as they approach it. They are so called from their use, which is to shew the altitude of a star above the horizon.

*CIRCLES of Declination*, are great circles intersecting each other in the poles of the world.

*CIRCLE of Dissipation*, in Optics. See the article DISSIPATION.

*Diurnal CIRCLES*, are parallels to the equinoctial, supposed to be described by the several stars, and other points of the heavens, in their apparent diurnal rotation about the earth.

*CIRCLE Equant*, in the Ptolomaic Astronomy, is a circle described on the centre of the equant. Its chief use is, to find the variation of the first inequality.

*CIRCLES of Excursion*, are little circles parallel to the ecliptic, and at such a distance from it, as that the excursions of the planets towards the poles of the ecliptic, may be included within them; being usually fixed at about 10 degrees.

It may here be observed, that all the circles of the sphere, described above, are transferred from the heavens to the earth; and so come to have a place in geography as well as in astronomy: all the points of each circle being conceived as let fall perpendicularly on the surface of the terrestrial globe, and thus tracing out circles perfectly similar to them. So, the terrestrial equator is a circle conceived precisely under the equinoctial line, which is in the heavens: and so of the rest.

*Horary CIRCLES*, in Dialling, are the lines which shew the hours on dials. These are straight lines on the dials, but called circles as being the projections of the meridians.

*Horary CIRCLE, or Hour Circle*, on the globe, is a small brazen circle fixed to the north pole, divided into 24 hours, and furnished with an index to point them out, thereby shewing the difference of meridians in time, and serving for the solution of many problems, on the artificial globes.

*CIRCLE of Illumination*, is that imaginary circle on the surface of the earth, which separates the illuminated side or hemisphere of the earth from the dark side: and all lines passing from the sun to the earth, being physically parallel, are perpendicular to the plane of this circle.

*CIRCLES of Latitude, or Secondaries of the Ecliptic*, are great circles perpendicular to the plane of the ecliptic,

intersecting one another in its poles, and passing through every star and planet, &c.—These are so called, because they serve to measure the latitude of the stars, which is an arch of one of these circles, intersected between the star and the ecliptic.

*CIRCLES of Longitude*, are lesser circles parallel to the ecliptic, diminishing more and more as they recede from it, or as they approach the pole of that circle.

They are so called, because the longitudes of the stars are counted upon them.

*CIRCLE of Perpetual Apparition*, one of the lesser circles parallel to the equator, described by the most northern point of the horizon, as the sphere revolves round by its diurnal motion.—All the stars included within this circle, are continually above the horizon, and so never set.

*CIRCLE of Perpetual Occultation*, is another lesser circle at a like distance from the equator, but on the other side of it, being described by the most southern point of the horizon, and contains all those stars which never appear in our hemisphere, or which never rise.

All other stars, being contained between these two circles, do alternately rise and set, at certain moments of the diurnal rotation.

*Polar CIRCLES*, are immoveable circles, parallel to the equator, and at such a distance from the pole as is equal to the greatest declination of the ecliptic, which now is  $23^{\circ} 28'$ . That next the northern pole is called the arctic, and that next the southern one the antarctic.

*CIRCLES of Position*, are circles passing through the common intersections of the horizon and meridian, and through any degree of the ecliptic, or the centre of any star, or other point in the heavens; and are used for finding out the situation or position of any star. These are usually six in number, cutting the equinoctial into 12 equal parts, which the astrologers call the Celestial Houses, and hence they are sometimes called Circles of the Celestial Houses.

*CIRCUIT, Electrical*, denotes the course of the electric fluid from the charged surface of an electric body, to the opposite surface, into which the discharge is made. Some electricians at first apprehended, that the same particles of the electric fluid that were thrown on one side of the charged glass, actually made the whole circuit of the intervening conductors, and arrived at the opposite side: whereas Dr. Franklin's theory only requires, that the redundancy of electric matter on the charged surface should pass into those bodies which form that part of the circuit which is contiguous to it, driving forward that part of the fluid which they naturally possess; and that the deficiency of the exhausted surface should be supplied by the neighbouring conductors, which form the last part of the circuit. On this supposition, a vibrating motion is successively communicated through the whole length of the circuit.

Many attempts were made, both in France and England, at an early period in the practice of electricity, to ascertain the distance to which the electric shock might be carried, and the velocity of its motion. The French philosophers, at different times, caused it to pass through circuits of 900 and even 2000 toises, or about 2 English miles and a half; and they discharged the Leyden phial through a basin of water, whose surface

was



was equal to about one acre. M. Monier found that, in passing through an iron wire of 950 toises in length, it did not spend a quarter of a second; and that its motion was instantaneous through a wire of 1319 feet. In 1747, Dr. Watfon, and other English philosophers, after many experiments of a similar kind, conveyed the electric matter through a circuit of 4 miles; and, from two several trials, they concluded that its passage is instantaneous. By all which doubtless is meant, that its motion is too rapid to be measured. Priestley's Hist. of Elect. vol. 1, pa. 128, 8vo edit.

**CIRCULAR**, appertaining to a circle; as a circular form, circular motion, &c.

**CIRCULAR Lines**, a name given by some authors to such straight lines as are divided by means of the divisions made in the arch of a circle. Such as the Sines, Tangents, Secants, &c.

**CIRCULAR Numbers**, are such as have their powers ending in the roots themselves. As the number 5, whose square is 25, and its cube 125, &c.

Napier's **CIRCULAR Parts**, are five parts of a right-angled or a quadrantal spherical triangle; they are the two legs, the complement of the hypotenuse, and the complements of the two oblique angles.

Concerning these circular parts, Napier gave a general rule in his *Logarithmorum Canonis Descriptio*, which is this; "*The rectangle under the radius and the sine of the middle part, is equal to the rectangle under the tangents of the adjacent parts, and to the rectangle under the cosines of the opposite parts.*" The right angle or quadrantal side being neglected, the two sides and the complements of the other three natural parts are called the circular parts; as they follow each other as it were in a circular order. Of these, any one being fixed upon as the middle part, those next it are the adjacent, and those farthest from it the opposite parts." Lord Buchan's Life of Napier, pa. 98.

This rule contains within itself all the particular rules for the solution of right-angled spherical triangles, and they were thus brought into one general comprehensive theorem, for the sake of the memory; as thus, by charging the memory with this one rule alone: All the cases of right-angled spherical triangles may be resolved, and those of oblique ones also, by letting fall a perpendicular, excepting the two cases in which there are given either the three sides, or the three angles.—And for these a similar expedient has been devised by Lord Buchan and Dr. Minto. "M. Pingre, in the *Memoires de Mathematique et de Physique* for the year 1756, reduces the solution of all the cases of spherical triangles to four analogies. These four analogies are in fact, under another form, Napier's rule of the circular parts, and his second or fundamental theorem, with its application to the supplemental triangle. Although it would be no difficult matter to get by heart the four analogies of M. Pingre, yet there are few persons blessed with a memory capable of retaining them for any considerable time. For this reason, the rule for the circular parts ought to be kept under its present form. If the reader attends to the circumstance of the second letters of the words *tangents* and *cosines* being the same with the first of the words *adjacent* and *opposite*, he will find it almost impossible to forget the rule. And

VOL. I.

the rule for the solution of the two cases of spherical triangles, for which the former of itself is insufficient, may be thus expressed: *Of the circular parts of an oblique spherical triangle, the rectangle under the tangents of half the sum and half the difference of the segments at the middle part* (formed by a perpendicular drawn from an angle to the opposite side), *is equal to the rectangle under the tangents of half the sum and half the difference of the opposite parts.* By the circular parts of an oblique spherical triangle are meant its three sides and the supplements of its three angles. Any of these six being assumed as a middle part, the opposite parts are those two of the same denomination with it, that is, if the middle part is one of the sides, the opposite parts are the other two, and, if the middle part is the supplement of one of the angles, the opposite parts are the supplements of the other two.—Since every plane triangle may be considered as described on the surface of a sphere of an infinite radius, these two rules may be applied to plane triangles, provided the middle part be restricted to a side.

"Thus it appears that two simple rules suffice for the solution of all the possible cases of plane and spherical triangles. These rules, from their neatness and the manner in which they are expressed, cannot fail of engraving themselves deeply on the memory of every one who is a little versed in trigonometry. It is a circumstance worthy of notice, that a person of a very weak memory may carry the whole art of trigonometry in his head." Napier's Life, pa. 102.

**CIRCULAR Sailing**, is that performed in the arch of a great circle.—It is chiefly on account of the shortest distance that this method of sailing has been proposed; and for the most part it is advantageous for a ship to reach her port by the shortest course.

As the solutions of the cases in Mercator's sailing are performed by plane triangles; so the cases in great-circle sailing are resolved by the solution of spherical triangles. But, after all, the several cases in this kind of sailing serve rather for exercises in the solution of spherical triangles, than for any real use towards the navigating of a ship.

**CIRCULAR Spots** are made on pieces of metal by large electrical explosions. See Philos. Trans. vol. 58, pa. 68; also Priestley's Hist. of Electricity, vol. 2, sect. 9, edit. 8vo.

These beautiful spots, produced by the moderate charge of a large battery, discharged between two smooth surfaces of metals, or semi-metals, lying at a small distance from each other, consist of one central spot, and several concentric circles, which are more or less distinct, and more or fewer in number, as the metal upon which they are marked is more easy or difficult of fusion, and as a greater or less force is employed. They are composed of dots or cavities, which indicate a real fusion. If the explosion of a battery, issuing from a pointed body, be repeatedly taken on the plain surface of a piece of metal near the point, or be received from the surface on a point, the metal will be marked with a spot, consisting of all the prismatic colours disposed in circles, and formed by scales of the metal separated by the force of the explosion.

**CIRCULAR Velocity**, a term in astronomy signifying that



that velocity of a planet, or revolving body, which is measured by the arch of a circle.

**CIRCULATING Decimals**, called also recurring or repeating decimals, are those in which a figure or several figures are continually repeated. They are distinguished into *single* and *multiple*, and these again into *pure* and *mixed*.

A *pure single* circulate, is that in which one figure only is repeated; as  $\cdot 222$  &c, and is marked thus  $\cdot 2$ .

A *pure multiple* circulate, is that in which several figures are continually repeated; as  $\cdot 232323$  &c, marked  $\cdot 23$ ; and  $\cdot 524524$  &c, marked  $\cdot 524$ .

A *mixed single* circulate, is that which consists of a terminate part, and a single repeating figure; as  $4\cdot 222$  &c, or  $4\cdot 2$ . And

A *mixed multiple* circulate is that which contains a terminate part with several repeating figures; as  $45\cdot 524$ .

That part of the circulate which repeats, is called the *repetend*: and the whole repetend, supposed infinitely continued, is equal to a vulgar fraction, whose numerator is the repeating number, or figures, and its denominator the same number of nines: so  $\cdot 2$  is  $= \frac{2}{9}$ ; and  $\cdot 23$  is  $= \frac{23}{99}$ ; and  $\cdot 524$  is  $= \frac{524}{999}$ .

It seems it was Dr. Wallis who first distinctly considered, or treated of infinite circulating decimals, as he himself informs us in his *Treatise of Infinites*. Since his time many other authors have treated on this part of arithmetic; the principal of these however, to whom the art is mostly indebted, are Messrs. Brown, Cunn, Martin, Emerson, Malcolm, Donn, and Henry Clarke, in whose writings the nature and practice of this art may be fully seen, especially in the last mentioned ingenious author.


**CIRCUMFERENCE**, in a general sense denotes the line or lines bounding any figure. But it is commonly used in a more limited sense, to denote the curve line which bounds a circle, and which is otherwise called the *periphery*; the boundary of a right-lined figure being expressed by the term *perimeter*.

The circumference of a circle is every where equidistant from the centre. And the circumferences of different circles are to one another as their radii or diameters, or the ratio of the diameter to the circumference is a constant ratio, in every circle, which is nearly as 7 to 22, as it was found by Archimedes, or, more nearly, as 1 to 3.1416. Under the article *Circle* may be seen various other approximations to that ratio, one of which is carried to 128 places of figures, viz by M. De Lagny.

The Circumference of every circle is supposed to be divided into 360 equal parts, called *degrees*.—Any part of a circumference is called an *arc* or *arch*; and a right line drawn from one end of an arc to the other, is called its *chord*.—The *angle* at the circumference is equal to half the angle at the centre, standing on the same arc; and therefore it is measured by the half of that arc.

**CIRCUMFERENTOR**, a particular instrument used by surveyors for taking angles. It consists of a brass circle and index all of a piece; the diameter of the circle is commonly about 7 inches; the index about 14 inches long, and an inch and a half broad. On the circle is a card or compass, divided into 360 degrees; the

meridian line of which answers to the middle of the breadth of the index. On the limb or circumference of the circle is foldered a brass ring; which, with another fitted with a glass, forms a kind of box for the needle, which is suspended on a pivot in the centre of the circle. There are also two sights to screw on, and slide up and down the index, as also a spangle and socket screwed on the under side of the circle, to receive the head of the three-legged staff.



To take, or observe the Quantity of an Angle by the Circumferentor. The angle proposed being EKG; place the instrument at K, with the flower-de-luce of the card towards you; then direct the sights to E, and observe what degrees are cut by the south end of the needle, which let be 295; then, turning the instrument about on its stand, direct the sights to G, noting again what degrees are cut by the south end of the needle, which suppose are 213. This done, subtract the less number from the greater, viz, 213 from 295, and the remainder, or 82 degrees, is the quantity of the angle EKG sought.

**CIRCUMGYRATION**, is the whirling motion of any body about a centre; as of the planets about the sun, &c.

**CIRCUM POLAR Stars**, are those stars which, by reason of their vicinity to the pole, move round it, without setting.

**CIRCUMSCRIBED Figure**, is a figure which is drawn about another, so that all its sides or planes touch the latter or inscribed figure.

The area and perimeter of every polygon that can be circumscribed about a circle, are greater than those of the circle; and the area and perimeter of every inscribed polygon, are less than those of the circle; but they approach always nearer to equality as the number of sides is more. And on these principles Archimedes, and some other authors since his time, attempted the quadrature of the circle; which is nothing else, in effect, but the measuring the area or capacity of a circle.

**CIRCUMSCRIBED Hyperbola**, is one of Newton's hyperbolas of the 2d order, that cuts its asymptotes, and contains the parts cut off within its own space.

**CIRCUMVALLATION**, or *Line of Circumvallation*, in the Art of War, is a trench, bordered with a parapet, thrown up around the besieger's camp, as a security against any army that may attempt to relieve the place, as well as to prevent desertion.


**CIRCUMVOLUTION**, in Architecture, the torus of the spiral line of the Ionic Order.

**CISSOID**, is a curve line of the second order, invented



vented by Diocles for the purpose of finding two continued mean proportionals between two other given lines. The generation or description of this curve is as follows:

On the extremity B of the diameter AB of the circle AOB, erect the indefinite perpendicular CBD, to which from the other extremity A draw several lines, cutting the circle in I, O, N, &c; and upon these lines set off the corresponding equal distances, viz,  $HM = AI$ , and  $FO = AO$ , and  $CL = AN$ , &c; then the curve line drawn through all the points M, O, L, &c, is the cissoid of Diocles, who was an ancient Greek geometer.



This curve is, by Newton, reckoned among the defective hyperbolas, being the 42d species in his *Enumeratio Linearum tertii ordinis*. And in his appendix de *Æquationum Constructione Lineari*, at the end of his *Arithmetica Universalis*, he gives another elegant method of describing this curve by the continual motion of a square ruler. Other methods have also been devised by different authors for the same thing.

*The Properties of the Cissoid* are the following :

1. The curve has two infinite legs AMOL, Amol meeting in a cusp A, and tending continually towards the indefinite line CBD, which is their common asymptote.

2. The curve passes through O and o, points in the circle equally distant from A and B; or it bisects each semicircle.

3. Letting fall perpendiculars MP, IK from any corresponding points I, M; then is  $AP = BK$ , and  $AM = HI$ , because  $AI = MH$ .

4.  $AP : PB :: MP^2 : AP^2$ . So that, if the diameter AB be  $= a$ , the absciss  $AP = x$ , and the ordinate  $PM = y$ ; then is  $x : a - x :: y^2 : x^2$ , or  $x^3 = a - x \times y^2$ ; which is the equation of the curve.

5. Sir Isaac Newton, in his last letter to M. Leibnitz, has shewn how to find a right line equal to one of the legs of this curve, by means of the hyperbola; but he suppressed the investigation, which however may be seen in his Fluxions.

6. The whole infinitely long cissoidal space, contained between the infinite asymptote BCD, and the curves LOAol &c, of the cissoid, is equal to triple the generating circle AOB oA.

See more of this curve in Dr. Wallis, vol. 1, pa. 545.

CIVIL Day. See Day.

CIVIL Month. See Month.

CIVIL Year, is the legal year, or annual account of time, which every government appoints to be used within its own dominions.

It is so called in contradistinction to the natural year,

which is measured exactly by the revolution of the heavenly bodies.

CLAIRAUT (ALEXIS-CLAUDE), a celebrated French mathematician and academician, was born at Paris the 13th of May 1713, and died the 17th of May 1765, at 52 years of age. His father, a teacher of mathematics at Paris, was his sole instructor, teaching him even the letters of the alphabet on the figures of Euclid's Elements, by which he was able to read and write at 4 years of age. By a similar stratagem it was that calculations were rendered familiar to him. At 9 years of age he put into his hands Guisnée's Application of Algebra to Geometry; at 10 he studied l'Hopital's Conic Sections; and between 12 and 13 he read a memoir to the Academy of Sciences concerning four new Geometrical curves of his own invention. About the same time he laid the first foundation of his work upon curves that have a double curvature, which he finished in 1729, at 16 years of age. He was named Adjoint-Mechanician to the Academy in 1731, at the age of 18, Associate in 1733, and Pensioner in 1738; during his connection with the Academy, he had a great multitude of learned and ingenious communications inserted in their Memoirs, beside several other works which he published separately; the list of which is as follows.

1. On Curves of a Double Curvature; in 1730, 4to.
2. Elements of Geometry; 1741, 8vo.
3. Theory of the Figure of the Earth; 1743, 8vo.
4. Elements of Algebra; 1746, 8vo.
5. Tables of the Moon; 1754, 8vo.

His papers inserted in the Memoirs of the Academy are too numerous to be particularised here; but they may be found from the year 1727, for almost every year till 1762; being upon a variety of subjects, astronomical, mathematical, optical, &c.

CLAVIUS (CHRISTOPHER), a German Jesuit, was born at Bamberg in Germany, in 1537. He became a very studious mathematician, and elaborate writer; his works, when collected, and closely printed, making 5 large folio volumes; being a complete body or course of the mathematics. They are mostly elementary, and commentaries on Euclid and others; having very little of invention of his own. His talents and writings have been variously spoken of, and it must be acknowledged that they are heavy and elaborate. He was sent for to Rome, to assist, with other learned men, in the reformation of the calendar, by pope Gregory; which he afterward undertook a defence of, against Scaliger, Vieta, and others, who attacked it. He died at Rome, the 6th of February, 1612, at 75 years of age, after more than 50 years close application to the mathematical sciences.

CLEFF. See Cliff.

CLERC (JOHN LE), a celebrated writer and universal scholar, was born at Geneva in 1657. After passing through the usual course of study at Geneva, he went to France in 1678; but returning the year after, he took holy orders. In 1682 Le Clerc visited England, to learn the language; but the smoky air of London not agreeing with his lungs, he soon returned to Holland, where he settled; and was appointed professor of philosophy, polite literature, and the Hebrew tongue, in the school at Amsterdam. Here he long continued



to read lectures; for which purpose he drew up and published his Logic, Ontology, Pneumatology, and Natural Philosophy. He published also *Ars Critica*; a Commentary on the Old Testament; a Compendium of Universal History; an Ecclesiastical History of the two first centuries; a French translation of the New Testament, and other works. In 1686, he began, jointly with M. de la Crose, his *Bibliothèque Universelle et Historique*, in imitation of other literary journals; which was continued to the year 1693, making 26 volumes. In 1703 he began his *Bibliothèque Choisie*, and continued it to 1714, when he commenced another work on the same plan, called *Bibliothèque Ancienne et Moderne*, which he continued to the year 1728; all of them justly esteemed excellent stores of useful knowledge. He published also, in 1713, a neat little treatise on Practical Geometry, in 2 vols. small 8vo, called, *Pratique de la Geometrie, sur le papier et sur le terrain*. In 1728 he was seized with a palsy and fever; and, after spending the last six years of his life with little or no understanding, he died in 1736, at 79 years of age.

**CLEPSYDRA**, a kind of water clock, or an hour-glass serving to measure time by the fall of a certain quantity, commonly out of one vessel into another.—There have been also clepsydræ made with quicksilver; and the term is also used for hour-glasses of sand.

By this instrument the Egyptians measured their time and the course of the sun. Also Tycho Brahe, in modern times, made use of it to measure the motion of the stars, &c; and Dudley used the same contrivance in making all his maritime observations.

The use of Clepsydræ is very ancient. They were probably invented in Egypt under the Ptolemys; though some authors ascribe the invention of them to the Greeks, and others to the Romans. Pliny informs us, that Scipio Nasica, about 150 years before Christ, gave the first hint for the construction of them: and Pancirollus has particularly described them. According to his account, the clepsydra was a vessel made of glass, with a small hole in the bottom, edged with gold: in the upper part of this vessel a line was drawn, and marked with the 12 hours: the vessel was filled with water, and a cork with a pin fixed in it floated on the surface, pointing to the first hour; and as the water sunk in the vessel by issuing out of the small hole, the pin indicated the other hours as it descended.

Clepsydræ were chiefly used in the winter; as sundials served for the summer. They had however two great defects; the one, that the water ran out more or less easily, as the air was more or less dense; the other, that the water flowed more rapidly at the beginning, than towards the conclusion when its quantity and pressure were much decreased. Amontons has invented a clepsydra which, it is said, is free from both these inconveniences; and the same effect is produced by one described by Mr. Hamilton, in the *Philos. Transf.* vol. 44, pa. 171, or *Abridg.* vol. 10, pa. 248. Varignon too, in the *Memoires de l'Acad.* 1699, delivers a general geometrical method of making clepsydræ, or water-clocks, with any kind of vessels, and with any given orifices for the water to run through.

Vitruvius, in lib. 9 of his *Achitecture*, treats of these instruments; and Pliny in chap. 60, lib. 7, says that Scipio Nasica was the first who measured time at Rome

by clepsydræ, or water-clocks. Gefner, in his *Pandects* pa. 91, gives several contrivances for these instruments. Solomon de Caus also treats on this subject in his *Reasons of Moving Forces* &c. So also does Ozanam, in his *Mathematical Recreations*, in which is contained a Treatise on Elementary Clocks, translated from the Italian of Dominique Martinelli. There is likewise a treatise on Hour-Glasses by Arcangelo Maria Radi, called *Nova Scienza de Horologi Polvere*. See also the *Technica Curiosa* of Gasper Schottus; and Amonton's *Remarques & Experiences Physiques sur la Construction d'une nouvelle Clepsydre, exempte des défauts des autres*.

**CLIFF**, or **CLEFF**, a term in Music, for a certain mark, from the position of which the proper places of all other notes in a piece of music are known.

**CLIMACTERIC**, a critical year in a person's life.

According to some, this is every 7th year: but others allow it only to those years produced by multiplying 7 by the odd numbers 3, 5, 7, 9. These years, say they, bring with them some remarkable change with respect to health, life, or fortune: the grand climacteric is the 63d year; but some add also the 81st to it: the other remarkable climacterics are the 7th, 21st, 35th, 49th, and 56th.

**CLIMATE**, or *Clima*, in Geography, a part of the surface of the earth, bounded by two lesser circles parallel to the equator; and of such a breadth, as that the longest day in the parallel nearer the pole exceeds the longest day in that next the equator, by some certain space, as half an hour, or an hour, or a month.

The beginning of a climate, is a parallel circle in which the day is the shortest; and the end of the climate, is that in which the day is the longest. The climates therefore are reckoned from the equator to the pole; and are so many zones or bands, terminated by lines parallel to the equator: though, in strictness, there are several climates, or different degrees of light or temperature, in the breadth of one zone. Each climate only differs from its contiguous ones, in that the longest day in summer is longer or shorter, by half an hour, for instance, in the one place than in the other.

As the climates commence at the equator, at the beginning of the first climate, that is at the equator, the day is just 12 hours long; but at the end of it, or at the beginning of the 2d climate, the longest day is 12 hours and a half long; and at the end of the 2d, or beginning of the 3d climate, the longest day is 13 hours long; and so of the rest, as far as the polar circles, where the hour climates terminate, and month climates commence. And as an hour climate is a space comprised between two parallels of the equator, in the first of which the longest day exceeds that in the latter by half an hour; so the month climate is a space contained between two circles parallel to the polar circles, and having its longest day longer or shorter than that of its contiguous one, by a month, or 30 days. But some authors, as Ricciolus, make the longest day of the contiguous climates to differ by half hours, to about the latitude of 45 degrees; then to differ by an hour, or sometimes 2 hours, to the polar circle; and after that by a month each. See tables of climates in Varenius, chap. 25, prop. 13.

The ancients, who confined the climates to what they thought the habitable parts of the earth, reckoned only seven,



seven, the middles of which they made to pass through some remarkable places; as the 1st through Meroe, the 2d through Sienna, the 3d through Alexandria, the 4th through Rhodes, the 5th through Rome, the 6th through Pontus, and the 7th through the mouth of the Borysthenes. But the moderns, who have failed farther toward the poles, make 30 climates on each side.

Vulgarly the term Climate is bestowed on any country or region differing from another either in respect of the seasons, the quality of the soil, or even the manners of the inhabitants; without any regard to the length of the longest day. Abulfeda, an Arabic author, distinguishes the first kind of climates by the term *real climates*, and the latter by that of *apparent climates*.

CLOCK, a machine now constructed in such a manner, and so regulated by the uniform motion of a pendulum, as to measure time, and all its subdivisions, with great exactness. Before the invention of the pendulum, a balance, not unlike the fly of a kitchen-jack, was used instead of it.—They were at first called nocturnal dials to distinguish them from sun-dials, which shewed the hour by the shadow of the sun.

The invention of clocks with wheels is ascribed to Pacificus, archdeacon of Verona, in the 9th century, on the credit of an epitaph quoted by Ughelli, and borrowed by him from Panvinus. Others attribute the invention to Boethius, about the year 510.

Mr. Derham, however, makes clock-work of a much older date; ranking Archimedes's sphere, mentioned by Claudian, and that of Posidonius, mentioned by Cicero, among machines of this kind: not that either their form or use were the same with those of ours; but that they had their motion from some hidden weights, or springs, with wheels or pulleys, or some such clock-work principle.

In the *Disquisitiones Monasticæ* of Benedictus Haëften, published in the year 1644, he says, that clocks were invented by Silvester the 4th, a monk of his order, about the year 998, as Dithmarus and Bozius have shewn; for before that time, they had nothing but sun-dials and clepsydræ to shew the hour.—Conrade Gesner, in his *Epitome*, pa. 604, says, that Richard Wallingford, an English abbot of St. Albans, who flourished in the year 1326, made a wonderful clock by a most excellent art, the like of which could not be produced by all Europe.—Moreri, under the word *Horologe* du Palais, says, that Charles the Fifth, called the wise king of France, ordered at Paris the first large clock to be made by Henry de Vie, whom he sent for from Germany, and set it upon the tower of his palace, in the year 1372.—John Froissart, in his *Histoire & Chronique*, vol. 2, ch. 28, says, the duke of Bourgogne had a clock, which sounded the hour, taken away from the city of Courtray, in the year 1382: and the same thing is said by Wm. Paradin, in his *Annales de Bourgogne*.

Clock-makers were first introduced into England in 1368, when Edward the 3d granted a licence for three artists to come over from Delft in Holland, and practise their occupation in this country.

The water-clocks, or clepsydræ, and sun-dials, have both a much better claim to antiquity. The French annals mention one of the former kind, sent by Aaron, king of Persia, to Charlemagne, about the year 807, which it would seem bore some resemblance to the

modern clocks: it was of brass, and shewed the hours by 12 little balls of the same metal, which at the end of each hour fell upon a bell, and made a sound. There were also figures of 12 cavaliers, which at the end of each hour came out through certain apertures, or windows, in the side of the clock, and shut them again, &c.

The invention of pendulum clocks is owing to the happy industry of the last age; and the honour of that discovery is disputed between Galileo and Huygens. The latter, who wrote an excellent volume on the subject, declares it was first put in practice in the year 1657, and the description of it printed in 1658. Becher, *De Nova Temporis dimetiendi Theoria*, anno 1680, contends for Galileo; and relates, though at second-hand, the whole history of the invention; adding that one Tresler, clock-maker to the father of the then grand duke of Tuscany, made the first pendulum clock at Florence, under the direction of Galileo Galilei, a pattern of which was brought to Holland. And the Academy del Cimento say expressly, that the application of the pendulum to the movement of a clock was first proposed by Galileo, and put in practice by his son Vincenzo Galilei, in 1649. But whoever may have been the inventor, it is certain that the invention never flourished till it came into the hands of Huygens, who insists on it, that if ever Galileo thought of such a thing, he never brought it to any degree of perfection. The first pendulum clock made in England was in the year 1662, by one Fromantil, a Dutchman.

Among the modern clocks, those of Strasburg and Lyons are very eminent for the richness of their furniture, and the variety of their motions and figures. In the former, a cock claps his wings, and proclaims the hour: the angel opens a door, and salutes the Virgin; and the holy spirit descends on her, &c. In the latter, two horsemen encounter, and beat the hour upon each other: a door opens, and there appears on the theatre the Virgin, with Jesus Christ in her arms; the Magi, with their retinue, marching in order, and presenting their gifts; two trumpeters sounding all the while to proclaim the procession.

These, however, are far excelled by two that have lately been made by English artists, as a present from the East-India company to the emperor of China. These two clocks are in the form of chariots, in each of which a lady is placed, in a fine attitude, leaning her right hand upon a part of the chariot, under which appears a clock of curious workmanship, little larger than a shilling, that strikes and repeats, and goes for eight days. Upon the lady's finger sits a bird, finely modelled, and set with diamonds and rubies, with its wings expanded in a flying posture, and actually flutters for a considerable time, on touching a diamond button below it; the body of the bird, in which are contained part of the wheels that animate it as it were, is less than the 16th part of an inch. The lady holds in her left hand a golden tube little thicker than a large pin, on the top of which is a small round box, to which is fixed a circular ornament not larger than a sixpence, set with diamonds, which goes round in near three hours in a constant regular motion. Over the lady's head is a double umbrella, supported by a small fluted pillar not thicker than a quill, and under the larger of which a bell is fixed at a considerable distance from the clock,



clock, with which it seems not to have any connection; but from which a communication is secretly conveyed to a hammer, that regularly strikes the hour, and repeats the same at pleasure, by touching a diamond button fixed to the clock below. At the feet of the lady is a golden dog; before which, from the point of the chariot, are two birds fixed on spiral springs, the wings and feathers of which are set with stones of various colours, and they appear as if flying away with the chariot, which, from another secret motion, is contrived to run in any direction, either straight or circular, &c; whilst a boy, that lays hold of the chariot behind, seems also to push it forwards. Above the umbrella are flowers and ornaments of precious stones; and it terminates with a flying dragon set in the same manner. The whole is of gold, most curiously executed, and embellished with rubies and pearls.

The ingenious Dr. Franklin contrived a clock to shew the hours, minutes, and seconds, with only three wheels and two pinions in the whole movement. The dial-plate has the hours engraven upon it in spiral spaces along two diameters of a circle, containing four times 60 minutes. The index goes round in four hours, and counts the minutes from any hour by which it has passed to the next following hour. The small hand, in an arch at top, goes round once in a minute, and shews the seconds. The clock is wound up by a line going over a pulley, on the axis of the great wheel, like a common 30 hour clock. Many of these very simple machines have since been constructed, that measure time exceedingly well. This clock is subject, however, to the inconvenience of requiring frequent winding, by drawing up the weight; as also to some uncertainty as to the particular hour shewn by the index.

Mr. Ferguson has proposed to remedy these inconveniences by another construction, which is described in his *Select Exercises*, pa. 4. This clock will go a week without winding, and always shews the precise hour; but, as Mr. Ferguson acknowledges, it has two disadvantages which do not belong to Dr. Franklin's clock: when the minute hand is adjusted, the hour plate must also be set right, by means of a pin; and the smallness of the teeth in the pendulum wheel will cause the pendulum ball to describe but small arcs in its vibrations; and therefore the momentum of the ball will be less, and the times of the vibrations will be more affected by any unequal impulse of the pendulum wheel on the pallets. Besides, the weight of the flat ring, on which the seconds are engraven, will load the pivots of the axis of the pendulum wheel with a great deal of friction, which ought by all possible means to be avoided. To remedy this inconvenience, the seconds plate might be omitted.

Mr. Ferguson also contrived a clock, shewing the apparent diurnal motions of the sun and moon, the age and phases of the moon, with the time of her coming to the meridian, and the times of high and low water; and all this by having only two wheels and a pinion added to the common movement. See his *Select Exercises* before mentioned. In this clock the figure of the sun serves as an hour index, by going round the dial-plate in 24 hours; and a figure of the moon goes round in 24 h. 50½ min. the time of her going round in the heavens from any meridian to the same meridian again.

A clock of this kind was adapted by Mr. Ferguson to the movement of an old watch. See also a description and drawing of an astronomical clock, shewing the apparent daily motions of the sun, moon, and stars, with the times of their rising, southing, and setting; the places of the sun and moon in the ecliptic, and the age and phases of the moon for every day of the year, in the same book, pa. 19.

There have been several treatises upon clocks; the principal of which are the following. Hieronymus Cardan, de Varietate Rerum libri 17.—Conrade Dasypodius, *Descriptio Horologii Astronomici Argentinenfis in summa Templi erecti*.—Guido Paucirollus, *Antiqua deperdita & nova reperta*.—L'Usage du Cadran, ou de l'Horloge Physique Univerfelle, par Galilée, Mathématicien du Duc de Florence.—Oughtred's *Opuscula Mathematica*.—Huygens's *Horologium Oscillatorium*.—*Pendule perpetuelle*, par l'Abbé de Hautefeuille.—J. J. Becheri *Theoria & Experientia de nova Temporis dimetiendi Ratione & Horologiorum Constructione*.—Clark's *Oughtredus explicatus, ubi de Constructione Horologiorum*.—*Horological Disquisitions*.—Huygens's *Posthumous Works*.—Sully's *Regle Artificielle du Temps, &c.*—Serviere's *Recueil d'Ouvrages Curieux*.—Derham's *Artificial Clock-maker*.—Camus's *Traité des Forces Mouvantes*.—Alexandre's *Traité Général des Horologies*.—Also *Treatises and Principles of Clock-making*, by Hatton, Cuming, &c. &c.

CLOUD, a collection of vapours suspended in the atmosphere, and rendered visible.

Although it be generally allowed that the clouds are formed from the aqueous vapours, which before were so closely united with the atmosphere as to be invisible: it is, however, not easy to account for the long continuance of some very opaque clouds without dissolving; or to assign the reason why the vapours, when they have once begun to condense, do not continue to do so till they at last fall to the ground in the form of rain or snow, &c. It is now known that a separation of the latent heat from the water of which vapour is composed is attended with a condensation of that vapour in some degree; in such case, it will first appear as a smoke, mist, or fog; which, if interposed between the sun and earth, will form a cloud; and the same causes continuing to operate, the cloud will produce rain or snow. It is however abundantly evident that some other cause besides mere heat or cold is concerned in the formation of clouds, and the condensation of atmospherical vapours. This cause is esteemed in a great measure the electrical fluid; indeed electricity is now so generally admitted as an agent in all the great operations of nature, that it is no wonder to find the formation of clouds attributed to it; and this has accordingly been given by Beccaria as the cause of the formation of all clouds whatsoever, whether of thunder, rain, hail, or snow.

But whether the clouds are produced, that is, the atmospheric vapours rendered visible, by means of electricity or not, it is certain that they do often contain the electric fluid in prodigious quantities, and many terrible and destructive accidents have been occasioned by clouds very highly electrified. The most extraordinary instance of this kind perhaps on record happened



in the island of Java, in the East-Indies, in August, 1772. On the 11th of that month, at midnight, a bright cloud was observed covering a mountain in the district called Cheribou, and several reports like those of a gun were heard at the same time. The people who dwelt upon the upper parts of the mountain not being able to fly fast enough, a great part of the cloud, eight or nine miles in circumference, detached itself under them, and was seen at a distance, rising and falling like the waves of the sea, and emitting globes of fire so luminous, that the night became as clear as day. The effects of it were astonishing; every thing was destroyed for 20 miles round; the houses were demolished; plantations were buried in the earth; and 2140 people lost their lives, besides 1500 head of cattle, and a vast number of horses, goats, &c. Another remarkable instance of the dreadful effects of electric clouds, which happened at Malta the 29th of October 1757, is related in Brydone's Tour through Malta.

The height of the clouds is not usually great: the summits of high mountains being commonly quite free from them, as many travellers have experienced in passing these mountains. It is found that the most highly electrified clouds descend lowest, their height being often not more than 7 or 800 yards above the ground; and sometimes thunder-clouds appear actually to touch the ground with one of their edges: but the generality of clouds are suspended at the height of a mile, or little more, above the earth.

The motions of the clouds, though often directed by the wind, are not always so, especially when thunder is about to ensue. In this case they are seen to move very slowly, or even to appear quite stationary for some time. The reason of this probably is, that they are impelled by two opposite streams of air nearly of equal strength; and in such cases it seems that both the aerial currents ascend to a considerable height; for Mess. Charles and Robert, when endeavouring to avoid a thunder cloud, in one of their aerial voyages with a balloon, could find no alteration in the course of the current, though they ascended to the height of 4000 feet above the earth. In some cases the motions of the clouds evidently depend on their electricity, independent of any current of air whatever. Thus, in a calm and warm day, small clouds are often seen meeting each other in opposite directions, and setting out from such short distances, that it cannot be supposed that any opposite winds are the cause. Such clouds, when they meet, instead of forming a larger one, become much smaller, and sometimes quite vanish; a circumstance most probably owing to the discharge of opposite electricities into each other. And this serves also to throw some light on the true cause of the formation of clouds; for if two clouds, the one electrified positively, and the other negatively, destroy each other on contact, it follows that any quantity of vapour suspended in the atmosphere, while it retains its natural quantity of electricity, remains invisible, but becomes a cloud when electrified either plus or minus.

The shapes of the clouds are also probably owing to their electricity; for in those seasons in which a great commotion has been excited in the atmospherical electricity, the clouds are seen assuming strange and whimsical shapes, that are continually varying. This, as

well as the meeting of small clouds in the air, and vanishing upon contact, is a sure sign of thunder.

The uses of the clouds are evident, as from them proceeds the rain that refreshes the earth, and without which, according to the present state of nature, the whole surface of the earth must become a mere desert. They are likewise useful as a screen interposed between the earth and the scorching rays of the sun, which are often so powerful as to destroy the grass and other tender vegetables. In the more secret operations of nature too, where the electric fluid is concerned, the clouds bear a principal share; and chiefly serve as a medium for conveying that fluid from the atmosphere into the earth, and from the earth into the atmosphere: in doing which, when electrified to a great degree, they sometimes produce very terrible effects; an instance of which is related above.

CLOUTS, in Artillery, are thin plates of iron nailed on that part of the axle-tree of a gun-carriage which comes through the nave, and through which the limpin goes.

CLUVIER, or CLUVERIUS, (PHILIP), a celebrated geographer, was born at Dantzic in 1580. After an education at home, he travelled into Poland, Germany, and the Netherlands, to improve himself in the knowledge of the law. But, when at Leyden, Joseph Scaliger persuaded him to give way to his genius for geography. In pursuance of this advice, Cluvier visited the greatest part of the European states. He was well skilled in many languages, speaking half a score with facility, viz, Greek, Latin, German, French, English, Dutch, Italian, Hungarian, Polish, and Bohemian. On his return to Leyden, he taught there with great applause; and died in 1623, being only 43 years of age, justly esteemed the first geographer who had put his researches in order, and reduced them to certain principles. He was author of several ingenious works in geography, viz:

1. *De Tribus Rhæni Alveis.*
2. *Germania Antiqua.*
3. *Italia Antiqua, Sicilia, Sardinia, & Corsica.*
4. *Introductio in Universam Geographiam.*

COASTING, is that part of Navigation in which the places are not far asunder, so that a ship may sail in sight of land, or within soundings between them.

COCHLEA, one of the five Mechanical powers, otherwise called the Screw; being so named from the resemblance a screw bears to the spiral shell of a snail, which the Latins call Cochlea. See SCREW, and MECHANICAL Powers.

COCK of a Dial, the pin, style, or gnomon.

COEFFICIENTS, in Algebra, are numbers, or given quantities, usually prefixed to letters, or unknown quantities, by which it is supposed they are multiplied; and so, with such letters, or quantities, making a product, or *coefficient* production; whence the name.

Thus, in  $3a$  the coefficient is 3, in  $bx$  it is  $b$ , and in  $cx^2$  it is  $c$ . If a quantity have no number prefixed, unity or 1 is understood; as  $a$  is the same as  $1a$ , and  $bc$  the same as  $1bc$ . The name *coefficient* was first given by Vieta.

In any equation so reduced as that its highest power or term has 1 for its coefficient; then the coefficient of the 2d term is equal to the sum of all the roots, both positive



positive and negative ; so that if the 2d term is wanting in an equation, the sum of the positive roots of that equation is equal to the sum of the negative roots, as they mutually balance and cancel each other. Also the coefficient of the 3d term of an equation is equal to the sum of all the rectangles arising by the multiplication of every two of the roots, how many ways soever they can be combined by twos ; as once in the quadratic, 3 times in the cubic, 6 times in the biquadratic equation, &c. And the coefficient of the 4th term of an equation, is the sum of all the solids made by the continual multiplication of every three of the roots, how often soever such a ternary can be had ; as once in a cubic, 4 times in a biquadratic, 10 times in an equation of 5 dimensions, &c. And thus it will go on infinitely.

**COEFFICIENTS of the same Order**, is a term sometimes used for the coefficients prefixed to the same unknown quantities, in different equations.

$$\text{Thus in the equations } \begin{cases} ax + by + cz = m, \\ dx + ey + fz = n, \\ gx + hy + kz = p, \end{cases}$$

the coefficients  $a, d, g$ , are of the same order, being the coefficients of the same letter  $x$  ; also  $b, e, h$  are of the same order, being the coefficients of  $y$  ; and so on.

**Opposite COEFFICIENTS**, such as are taken each from a different equation, and from a different order of coefficients. Thus, in the foregoing equations,  $a, e, k$ , or  $a, b, f$ , or  $d, b, k$ , &c, are opposite coefficients.

**COELESTIAL**. See **CELESTIAL**.

**COFFER**, in Architecture, a square depressure or sinking, in each interval between the modillions of the Corinthian cornice ; usually filled up with a rose ; sometimes with a pomegranate, or other enrichment.

**COFFER**, in Fortification, denotes a hollow lodgment, athwart a dry moat, 6 or 7 feet deep, and 16 or 18 broad. The upper part of it is made of pieces of timber, raised 2 feet above the level of the moat ; the elevation having hurdles laden with earth for its covering, and serving as a parapet with embrasures.

The coffer is nearly the same with the caponiere, excepting that this last is sometimes made beyond the counterscarp on the glacis, and the coffer always in the moat, taking up its whole breadth, which the caponiere does not.

It differs from the traverse and gallery, in that these are made by the besiegers, and the coffer by the besieged.

The besieged commonly make use of coffers to repulse the besiegers, when they endeavour to pass the ditch. And, on the other hand, the besiegers, to save themselves from the fire of these coffers, throw up the earth on that side towards the coffer.

**COFFER-Dams, or Batardeaux**, in Bridge-building, are inclosures formed for laying the foundation of piers, and for other works in water, to exclude the surrounding water, and so prevent it from interrupting the workmen.

These inclosures are sometimes single, and sometimes double, with clay rammed between them ; sometimes they are made with piles driven close by one another, and sometimes the piles are notched or dove-tailed into one another ; but the most usual method is to drive piles with grooves in them, at the distance of five or six feet from each other, and then boards are let down

between them ; after which the water is pumped out.

**COGGESHALL's Sliding-Rule**, an instrument used in Gauging, and so called from its inventor. See the description and use under **SLIDING-Rule**.

**COHESION**, one of the four species of attraction, denoting that force by which the parts of bodies adhere or stick together.

This power was first considered by Newton as one of the properties essential to all matter, and the cause of all that variety observed in the texture of different terrestrial bodies. He did not, however, absolutely determine that the power of cohesion was an immaterial one ; but that it might possibly arise, as well as that of gravitation, from the action of another. His doctrine of cohesion Newton delivers in these words : " The particles of all hard homogeneous bodies, which touch one another, cohere with a great force ; to account for which, some philosophers have recourse to a kind of hooked atoms, which in effect is nothing else but to beg the thing in question. Others imagine that the particles of bodies are connected by rest, i. e. in effect by nothing at all ; and others by conspiring motions, i. e. by a relative rest among themselves. For myself, it rather appears to me, that the particles of bodies cohere by an attractive force, whereby they tend mutually toward each other : which force, in the very point of contact, is very great ; at little distances is less, and at a little farther distance is quite insensible."

It is uncertain in what proportion this force decreases as the distance increases : Desaguliers conjectures, from some phenomena, that it decreases as the biquadratic or 4th power of the distance, so that at twice the distance it acts 16 times more weakly, &c.

" Now if compound bodies be so hard, as by experience we find some of them to be, and yet have a great many hidden pores within them, and consist of parts only laid together ; no doubt those simple particles which have no pores within them, and which were never divided into parts, must be vastly harder. For such hard particles, gathered into a mass, cannot possibly touch in more than a few points : and therefore much less force is required to sever them, than to break a solid particle, whose parts touch throughout all their surfaces, without any intermediate pores or interstices. But how such hard particles, only laid together, and touching only in a few points, should come to cohere so firmly, as in fact we find they do, is inconceivable ; unless there be some cause, whereby they are attracted and pressed together. Now the smallest particles of matter may cohere by the strongest attractions, and constitute larger, whose attracting force is feebler : and again, many of these larger particles cohering, may constitute others still larger, whose attractive force is still weaker ; and so on for several successions, till the progression end in the biggest particle, on which the operations in chemistry, and the colours of natural bodies, do depend ; and which by cohering compose bodies of a sensible magnitude."

Again, the opinion maintained by many is that which is so strongly defended by J. Bernoulli, *De Gravitate Aetheris* ; who attributes the cohesion of the parts of matter to the uniform pressure of the atmosphere ; confirming this opinion by the known experiment of two polished marble planes, which cohere very strongly



in the open air, but easily drop asunder in an exhausted receiver. However, if two plates of this kind be smeared with oil, to fill up the pores in their surfaces, and prevent the lodgment of air, and one of them be gently rubbed upon the other, they will adhere so strongly, even when suspended in an exhausted receiver, that the weight of the lower plate will not be able to separate it from the upper one. But although this theory might serve tolerably well to explain the cohesion of compositions, or greater collections of matter; yet it falls far short of accounting for that first cohesion of the atoms, or primitive corpuscles, of which the particles of hard bodies are composed.

Again, some philosophers have positively asserted, that the powers, or means, are immaterial, by which matter coheres; and, in consequence of this supposition, they have so refined upon attractions and repulsions, that their systems seem but little short of scepticism, or denying the existence of matter altogether. A system of this kind is adopted by Dr. Priestley, from Messrs. Boscovich and Michell, to solve some difficulties concerning the Newtonian doctrine of light. See his *History of Vision*, vol. 1. pa. 392. "The easiest method," says he, "of solving all difficulties, is to adopt the hypothesis of Mr. Boscovich, who supposes that matter is not impenetrable, as has been perhaps universally taken for granted; but that it consists of physical points only, endued with powers of attraction and repulsion in the same manner as solid matter is generally supposed to be: provided therefore that any body move with a sufficient degree of velocity, or have a sufficient momentum to overcome any powers of repulsion that it may meet with, it will find no difficulty in making its way through any body whatever; for nothing else will penetrate one another but powers, such as we know do in fact exist in the same place, and counterbalance or over-rule one another. The most obvious difficulty, and indeed almost the only one, that attends this hypothesis, as it supposes the mutual penetrability of matter, arises from the idea of the nature of matter, and the difficulty we meet with in attempting to force two bodies into the same place. But it is demonstrable, that the first obstruction arises from no actual contact of matter, but from mere powers of repulsion. This difficulty we can overcome; and having got within one sphere of repulsion, we fancy that we are now impeded by the solid matter itself. But the very same is the opinion of the generality of mankind with respect to the first obstruction. Why, therefore, may not the next be only another sphere of repulsion, which may only require a greater force than we can apply to overcome it, without disordering the arrangement of the constituent particles; but which may be overcome by a body moving with the amazing velocity of light?"

Other philosophers have supposed that the powers both of gravitation and cohesion are material; and that they are only different actions of the ethereal fluid, or elementary fire. In proof of this doctrine, they allege the experiment with the Magdeburg hemispheres, as they are called. The pressure of the atmosphere we see is, in this case, capable of producing a very strong cohesion; and if there be in nature any fluid more penetrating, as well as more powerful in its effects, than the air we breath, it is possible that what is called

the attraction of cohesion may in some measure be an effect of the action of that fluid. Such a fluid as this is the element of fire. Its activity is such as to penetrate all bodies whatever; and in the state in which it is commonly called fire, it acts according to the quantity of solid matter contained in the body. In this state, it is capable of dissolving the strongest cohesions observed in nature. Fire, therefore, being able to dissolve cohesions, must also be capable of causing them, provided its power be exerted for that purpose, which possibly it may be, when we consider its various modes or appearances, viz, as fire or heat, in which state it consumes, destroys, and dissolves; or as light, when it seems deprived of that destructive power; and as the electric fluid, when it attracts, repels, and moves bodies, in a great variety of ways. In the *Philos. Trans.* for 1777 this hypothesis is noticed, and in some measure adopted by Mr. Henly. "Some gentlemen (says he) have supposed that the electric matter is the cause of the cohesion of the particles of bodies. If the electric matter be, as I suspect, a real elementary fire inherent in all bodies, that opinion may probably be well founded; and perhaps the folding of metals, and the cementation of iron, by fire, may be considered as strong proofs of the truth of their hypothesis."

But whatever the cause of cohesion may be, its effects are evident and certain. The different degrees of it constitute bodies of different forms and properties. Thus, Newton observes, the particles of fluids, which do not cohere too strongly, and are small enough to render them susceptible of those agitations which keep liquors in a fluid, are most easily separated and rarefied into vapour, and make what the chemists call *volatile bodies*; being rarefied with an easy heat, and again condensed with a moderate cold. Those that have grosser particles, and so are less susceptible of agitation, or cohere by a stronger attraction, are not separable without a greater degree of heat; and some of them not without fermentation: and these make what the chemists call *fixed bodies*.

Air, in its fixed state, possesses the interstices of solid substances, and probably serves as a bond of union to their constituent parts; for when these parts are separated, the air is discharged, and recovers its elasticity. And this kind of attraction is evinced by a variety of familiar experiments; as, by the union of two contiguous drops of mercury; by the mutual approach of two pieces of cork, floating near each other in a basin of water; by the adhesion of two leaden balls, whose surfaces are scraped and joined together with a gentle twist, which is so considerable, that, if the surfaces are about a quarter of an inch in diameter, they will not be separated by a weight of 100 lb; by the ascent of oil or water between two glass planes, so as to form the hyperbolic curve, when they are made to touch on one side, and kept separate at a small distance on the other; by the depression of mercury, and by the rise of water in capillary tubes, and on the sides of glass vessels; also in sugar, sponge, and all porous substances. And where this cohesive attraction ends, a power of repulsion begins.

To determine the force of cohesion, in a variety of different substances, many experiments have been made, and particularly by professor Muschenbroek. The ad-

hesion



hesion of polished planes, about two inches in diameter, heated in boiling water, and smeared with greafe, required the following weights to separate them :

	Cold greafe	Hot greafe
Planes of Glafs	- 150 lb	- 300 lb
Brass	- 150	- 800
Copper	- 200	- 850
Marble	- 225	- 600
Silver	- 150	- 250
Iron	- 300	- 950

But when the Brass planes were made to adhere by other sorts of matter, the results were as in the following table :

With Water	- - -	12 oz
Oil	- - -	18
Venice Turpentine	- - -	24
Tallow Candle	- - -	800
Rosin	- - -	850
Pitch	- - -	1400

In estimating the *Absolute Cohesion* of solid pieces of bodies, he applied weights to separate them according to their length : his pieces of wood were long square parallelipedons, each side of which was  $\frac{1}{26}$  of an inch, and they were drawn asunder by the following weights :

Fir	- - -	600 lb
Elm	- - -	950
Alder	- - -	1000
Linden tree	- - -	1000
Oak	- - -	1150
Beech	- - -	1250
Ash	- - -	1250

He tried also wires of metal,  $\frac{1}{10}$ th of a Rhinland inch in diameter : the metals and weights were as follow :

Of Lead	- - -	29 $\frac{1}{4}$ lb
Tin	- - -	40 $\frac{1}{4}$
Copper	- - -	299 $\frac{1}{4}$
Yellow Brass	- - -	360
Silver	- - -	370
Iron	- - -	450
Gold	- - -	500

He then tried the *Relative Cohesion*, or the force with which bodies resist an action applied to them in a direction perpendicular to their length. For this purpose he fixed pieces of wood by one end into a square hole in a metal plate, and hung weights towards the other end, till they broke at the hole : the weights and distances from the hole are exhibited in the following table.

	Distance	Weight
Pine	- 9 $\frac{1}{2}$ inc	- 36 $\frac{1}{2}$ oz
Fir	- 9	- 40
Beech	- 7	- 56 $\frac{1}{2}$
Elm	- 9	- 44
Oak	- 8 $\frac{1}{2}$	- 48
Alder	- 9 $\frac{1}{4}$	- 48

See his Elem. Nat. Philos. cap. 19.

COLD, the privation of heat, or the opposite to it.

As it is supposed that heat consists in a particular motion of the parts of the hot body, hence the nature of cold, which is its opposite, is deduced ; for it is found that cold extinguishes, or rather abates heat ;

hence it would seem to follow, that those bodies are cold, which check and restrain the motion of the particles in which heat consists.

In general, cold contracts most bodies, and heat expands them : though there are some instances to the contrary, especially in the extreme cases or states of these qualities of bodies. Thus, though iron, in common with other bodies, expand with heat, yet, when melted, it is always found to expand in cooling again. So also, though water always is found to expand gradually as it is heated, and to contract as it cools, yet in the act of freezing, it suddenly expands again, and that with a most enormous force, capable of rending rocks, or bursting the very thick shells of metal, &c. A computation of the force of freezing water has been made by the Florentine Academicians, from the bursting of a very strong brass globe or shell, by freezing water in it ; when, from the known thickness and tenacity of the metal, it was found that the expansive power of a spherule of water only one inch in diameter, was sufficient to overcome a resistance of more than 27,000 pounds, or 13 tons and a half. See also experiments on bursting thick iron bomb-shells by freezing water in them by Major Edward Williams of the Royal Artillery, in the Edinb. Philos. Transf. vol. 2.

Such a prodigious power of expansion, almost double that of the most powerful steam engines, and exerted in so small a mass, seemingly by the force of cold, was thought a very powerful argument in favour of those who supposed that cold, like heat, is a positive substance. Dr. Black's discovery of latent heat, however, has now afforded a very easy and natural explication of this phenomenon. He has shewn, that, in the act of congelation, water is not cooled more than it was before, but rather grows warmer : that as much heat is discharged, and passes from a latent to a sensible state, as, had it been applied to water in its fluid state, would have heated it to  $135^{\circ}$ . In this process, the expansion is occasioned by a great number of minute bubbles suddenly produced. Formerly these were supposed to be cold in the abstract ; and to be so subtle, that, insinuating themselves into the substances of the fluid, they augmented its bulk, at the same time that, by impeding the motion of its particles upon each other, they changed it from a fluid to a solid. But Dr. Black shews that these are only air extricated during the congelation ; and to the extrication of this air he ascribes the prodigious expansive force exerted by freezing water. The only question therefore now remaining, is, By what means this air comes to be extricated, and to take up more room than it naturally does in the fluid. To this it may be answered, that perhaps part of the heat which is discharged from the freezing water, combines with the air in its unelastic state, and, by restoring its elasticity, gives it that extraordinary force, as is seen also in the case of air suddenly extricated in the explosion of gun-powder.

Cold also usually tends to make bodies electric, which are not so naturally, and to increase the electric properties of such as are so. And it is farther found that all substances do not transmit cold equally well ; but that the best conductors of electricity, viz metals, are likewise the best conductors of cold. It may farther be added, that when the cold has been carried to such



such an extremity as to render any body an electric, it then ceases to conduct the cold so well as before. This is exemplified in the practice of the Laplanders and Siberians; where, to exclude the extreme cold of the winters from their habitations the more effectually, and yet to admit a little light, they cut pieces of ice, which in the winter time must always be electric in those countries, and put them into their windows; which they find to be much more effectual in keeping out the cold than any other substance.

Cold is the destroyer of all vegetable life, when increased to an excessive degree. It is found that many garden plants and flowers, which seem to be very stout and hardy, go off at a little increase of cold beyond the ordinary standard. And in severe winters, nature has provided the best natural defence for the corn fields and gardens, namely, a covering of snow, which preserves such parts green and healthy as are under it, while such as are uncovered by it are either killed or greatly injured.

Dr. Clarke is of opinion, that cold is owing to certain nitrous, and other saline particles, endued with particular figures proper to produce such effects. Hence, sal-ammoniac, saltpetre, or salt of urine, and many other volatile and alkalizate salts, mixed with water, very much increase its degree of cold. In the *Philos. Trans.* number 274, M. Geoffroy relates some remarkable experiments with regard to the production of cold. Four ounces of sal-ammoniac dissolved in a pint of water, made his thermometer descend 2 inches and  $\frac{3}{4}$  in less than 15 minutes. An ounce of the same salt put into 4 or 5 ounces of distilled water, made the thermometer descend 2 inches and  $\frac{1}{4}$ . Half an ounce of sal-ammoniac mixed with 3 ounces of spirit of nitre, made the thermometer descend 2 inches and  $\frac{5}{8}$ ; but, on using spirit of vitriol instead of nitre, it sunk 2 inches and  $\frac{1}{2}$ . In this last experiment it was remarked, that the vapours raised from the mixture had a considerable degree of heat, though the liquid itself was so extremely cold. Four ounces of saltpetre mixed with a pint of water, sunk the thermometer an inch and  $\frac{1}{4}$ ; but a like quantity of sea salt sunk it only  $\frac{1}{2}$  of an inch. Acids always produced heat, even common salt with its own spirit. Volatile alkaline salts produced cold in proportion to their purity, but fixed alkalies heat.

But the greatest degree of cold produced by the mixture of salts and aqueous fluids, was that shewn by Homberg; who gives the following receipt for making the experiment: Take a pound of corrosive sublimate, and as much sal-ammoniac; powder them separately, and mix the powders well; put the mixture into a vial, pouring upon it a pint and a half of distilled vinegar, shaking all well together. This composition grows so cold, that it can scarce be held in the hand in summer; and it happened, as M. Homberg was making the experiment, that the matter froze. The same thing once happened to M. Geoffroy, in making an experiment with sal-ammoniac and water, but it never was in his power to make it succeed a second time.

If, instead of making these experiments with fluid water, it be taken in its congealed state of ice, or rather snow, degrees of cold will be produced greatly superior to any that have yet been mentioned. A mixture of snow and common salt sinks Fahrenheit's thermometer to 0; pot ashes and pounded ice sunk it 8 degrees far-

ther; two affusions of spirit of salt on pounded ice sunk it  $14\frac{1}{2}$  below 0; and by repeated affusions of spirit of nitre M. Fahrenheit sunk it to  $40^{\circ}$  below 0. This is the ultimate degree of cold which the mercurial thermometer will measure; for the mercury itself begins then to congeal; and therefore recourse must afterwards be had to spirit of wine, naptha, or some other fluid that will not congeal. The greatest degree of cold hitherto produced by artificial means, has been  $80^{\circ}$  below 0; which was done at Hudson's Bay by means of snow and vitriolic acid, the thermometer standing naturally at  $20^{\circ}$  below 0. Indeed greater degrees of cold than this have been supposed: Mr. Martin, in his *Treatise on Heat*, relates, that at Kirenga in Siberia, the mercurial thermometer sunk to  $118^{\circ}$  below 0; and professor Brown at Petersburg, when he made the first experiment of congealing quicksilver, fixed the point of congelation at  $350^{\circ}$  below 0; but from later experiments it has been more accurately determined, that  $40^{\circ}$  below 0 is the freezing point of quicksilver.

The most remarkable experiment however was made by Mr. Walker of Oxford, with spirit of nitre poured on Glauber's salt, the effect of which was found to be similar to that of the same spirit poured on ice or snow; and the addition of sal-ammoniac rendered the cold still more intense. The proportions of these ingredients recommended by Mr. Walker, are concentrated nitrous acid two parts by weight, water one part; of this mixture, cooled to the temperature of the atmosphere, 18 ounces; of Glauber's salt, a pound and a half avoirdupois; and of sal-ammoniac, 12 ounces. On adding the Glauber's salt to the nitrous acid, the thermometer fell  $52^{\circ}$ , viz from 50 to  $-2$ ; and on the addition of the sal-ammoniac, it fell to  $-9^{\circ}$ . Thus Mr. Walker was able to freeze quicksilver without either ice or snow, when the thermometer stood at  $45^{\circ}$ ; viz, by putting the ingredients in 4 different pans, and inclosing these within each other.

Excessive degrees of cold occur naturally in many parts of the globe in the winter season.

Although the thermometer in this country hardly ever descends so low as 0, yet in the winter of 1780, Mr. Wilson of Glasgow observed, that a thermometer laid on the snow sunk to  $25^{\circ}$  below 0; and Mr. Derham, in the year 1708, observed in England, that the mercury stood within one-tenth of an inch of its station when plunged into a mixture of snow and salt. At Petersburg, in 1732, the thermometer stood at  $28^{\circ}$  below 0; and when the French academicians wintered near the polar circle, the thermometer sunk to  $33^{\circ}$  below 0; and in the Asiatic and American continents, still greater degrees of cold are often observed.

The effects of these extreme degrees of cold are very surprising. Trees are burst, rocks rent, and rivers and lakes frozen several feet deep: metallic substances blister the skin like red-hot iron: the air, when drawn in by breathing, hurts the lungs, and excites a cough: even the effects of fire in a great measure seem to cease; and it is observed, that though metals are kept for a considerable time before a strong fire, they will still freeze water when thrown upon them. When the French mathematicians wintered at Tornea in Lapland, the external air, when suddenly admitted into their rooms, converted the moisture of the air into whirls of snow; their breasts seemed to be rent when they breathed



it, and the contact of it was intolerable to their bodies; and the spirit of wine, which had not been highly rectified, burst some of their thermometers by the congelation of the aqueous part.

Extreme cold too often proves fatal to animals in those countries where the winters are very severe; thus 7000 Swedes perished at once in attempting to pass the mountains which divide Norway from Sweden. But it is not necessary that the cold, in order to prove fatal to human life, should be so very intense as has just been mentioned; it is only requisite to be a little below  $32^{\circ}$  of Fahrenheit, or the freezing point, accompanied with snow or hail, from which shelter cannot be obtained. The snow which falls upon the clothes, or the uncovered parts of the body, then melts, and by a continual evaporation carries off the animal heat to such a degree, that a sufficient quantity is not left for the support of life. In such cases, the person first feels himself extremely chill and uneasy; he turns listless, unwilling to walk or use exercise to keep himself warm, and at last turns drowsy, sits down to refresh himself with sleep, but wakes no more.

**COLLIMATION**, *Line of*, in a telescope, is a line passing through the intersection of those wires that are fixed in the focus of the object-glass, and the centre of the same glass.

**COLLINS** (JOHN), an eminent accountant and mathematician, was born at Wood Eaton near Oxford, March 5, 1624. At 16 years of age he was put apprentice to a bookseller at Oxford; but his genius appeared so remarkable for the study of the mechanical and mathematical sciences, that he was taken under the tuition of Mr. Marr, who drew several curious dials, which were placed in different positions in the king's gardens; under whom Mr. Collins made no small progress in the mathematics. In the course of the civil wars, he travelled abroad, to prosecute his favourite study; and on his return he took upon him the profession of an accountant, and published, in the year 1652, a large work entitled, *An Introduction to Merchants Accounts*; which was followed by several other publications on different branches of accounts. In 1658, he published a treatise called *The Sector on a Quadrant*; containing the description and use of four several quadrants, each accommodated to the making of sun-dials, &c; to which he afterward added an appendix concerning reflected dialling, from a glass placed reclining.—In 1659, he published his *Geometrical Dialling*; and the same year also his *Mariner's Plain Scale new plained*.—Collins now became a fellow of the Royal Society in London, to which he made various communications; particularly some ingenious chronological rules for the calendar, printed in the *Philos. Transf.* number 46, for April 1669: also a curious dissertation concerning the resolution of equations in numbers, in number 69, for March 1671: an elegant construction of the curious problem, having given the mutual distances of three objects in a plane, with the angles made by them at a fourth place in that plane, to find the distance of this place from each of the three former, vol. 6. pa. 2093: and thoughts about some defects in algebra, vol. 14. pa. 375.

Collins wrote also several commercial tracts, highly acceptable to the public; viz, *A Plea for bringing over Irish cattle, and keeping out the fish caught by foreign-*

*ers*:—For the promotion of the English fishery:—For the working the Tin-mines:—*A Discourse of Salt and Fishery*. He was frequently consulted in nice and critical cases of accounts, of commerce, and engineering. On one of these occasions, being appointed to inspect the ground for cutting a canal or river between the Isis and the Avon, he contracted a disorder by drinking cyder when he was too warm, which ended in his death, the 10th of November 1683, at 59 years of age.

Mr. Collins was a very useful man to the sciences, keeping up a constant correspondence with the most learned men, both at home and abroad, and promoting the publication of many valuable works, which, but for him, would never have been seen by the public; particularly Dr. Barrow's optical and geometrical lectures; his abridgment of the works of Archimedes, Apollonius, and Theodosius; Branker's translation of Rhronius's algebra, with Dr. Pell's additions, &c; which were procured by his frequent solicitations.

It was a considerable time after, that his papers were all delivered into the hands of the learned and ingenious Mr. William Jones, F. R. S. among which were found manuscripts, upon mathematical subjects, of Briggs, Oughtred, Barrow, Newton, Pell, and many others. From a variety of letters from these, and many other celebrated mathematicians, it appears that Collins spared neither pains nor cost to procure what tended to promote real science: and even many of the late discoveries in physical knowledge owe their improvement to him; for while he excited some to make known every new and useful invention, he employed others to improve them. Sometimes he was peculiarly useful, by shewing where the defect was in any useful branch of science, pointing out the difficulties attending the enquiry, and at other times setting forth the advantages, and keeping up a spirit and warm desire for improvement. Mr. Collins was also as it were the register of all the new improvements made in the mathematical sciences; the magazine to which the curious had frequent recourse: in so much that he acquired the appellation of the English *Mersennus*. If some of his correspondents had not obliged him to conceal their communications, there could have been no dispute about the priority of the invention of a method of analysis, the honour of which evidently belongs to Newton; as appears undeniably from the papers printed in the *Commercium Epistolicum D. Joannis Collins & aliorum de Analyti promota; jussu Societatis Regiæ in lucem editum*, 1712; a work that was made out from the letters in the possession of our author.

**COLLINS'S Quadrant.** See **QUADRANT**.

**COLLISION**, is the friction, percussion, or striking of bodies against one another.

Striking bodies are considered either as elastic, or non-elastic. They may also be either both in motion, or one of them in motion, and the other at rest.

When non-elastic bodies strike, they unite together as one mass; which, after collision, either remains at rest, or moves forward as one body. But when elastic bodies strike, they always separate after the stroke.

The principal theorems relating to the collision of bodies, are the following:

1. If any body impinge or act obliquely on a plane surface; the force or energy of the stroke, or action, is as the sine of the angle of incidence. Or the force upon



upon the surface, is to the same when acting perpendicularly, as the line of incidence is to radius.

2. If one body act on another, in any direction, and by any kind of force; the action of that force on the second body is made only in a direction perpendicular to the surface on which it acts.

3. If the plane, acted on, be not absolutely fixed, it will move, after the stroke, in the direction perpendicular to its surface.

4. If a body A strike another body B, which is either at rest, or else in motion, either towards A or from it; then the momenta, or quantities of motion, of the two bodies, estimated in any one direction, will be the very same after the stroke that they were before it.




Thus, first, if A with a momentum of 10, strike B at rest, and communicate to it a momentum of 4, in the direction AB. Then there will remain in A only a momentum of 6 in that direction: which together with the momentum of B, viz 4, makes up still the same momentum between them as before.—But if B were in motion before the stroke, with a momentum of 5, in the same direction, and receive from A an additional momentum of 2: then the motion of A after the stroke will be 8, and that of B, 7; which between them make up 15, the same as 10 and 5, the motions before the stroke.—Lastly, if the bodies move in opposite directions, and meet one another, namely A with a motion of 10, and B, of 5; and A communicate to B a motion of 6 in the direction AB of its motion: then, before the stroke, the whole motion from both, in the direction AB, is  $10 - 5$ , or 5: but after the stroke the motion of A is 4 in the direction AB, and the motion of B is  $6 - 5$ , or 1 in the same direction AB; therefore the sum  $4 + 1$ , or 5, is still the same motion from both as it was before.


5. If a hard and fixed plane be struck either by a soft or a hard unelastic body; the body will adhere to it. But if the plane be struck by a perfectly elastic body, it will rebound from it with the same velocity with which it struck the plane.

6. The effect of the blow of the elastic body, upon the plane, is double to that of the non-elastic one; the velocity and mass being the same in both.

7. Hence, non-elastic bodies lose, by their collision, only half the motion that is lost by elastic bodies; the masses and velocities being equal.



8. If an elastic body A impinge upon a firm plane DE at the point B, it will rebound from it in an angle equal to that in which it struck it; or the angle of incidence will be equal to the angle of reflection: namely, the angle  $ABD = CBE$ .



9. If the non-elastic body B, moving with the velocity V in the direction Bb, and the body b with the velocity v, strike each other, the direction of the mo-

tion being in the line BC; then they will move after the stroke with a common velocity, which will be more or less according as, before the stroke, b moved towards B, or from B, or was at rest; and that common velocity, in each of these cases, will be as follows: viz, it will be

$$\frac{BV + bv}{B + b} \text{ when } b \text{ moved from B,}$$

$$\frac{BV - bv}{B + b} \text{ when } b \text{ moved towards B,}$$

$$\frac{BV}{B + b} \text{ when } b \text{ was at rest.}$$

For example, if the bodies or weights, B and b, be 5lb and 3lb; and their velocities V and v, 60 feet and 40 feet per second; then 300 and 120 will be their momenta BV and bv, and  $18 = B + b$  the sum of the weights. Consequently the common velocity after the stroke, in the three cases above mentioned, will be thus, viz,

$$\frac{300 + 120}{18} = \frac{420}{18} \text{ or } 23\frac{1}{3} \text{ in the first case,}$$

$$\frac{300 - 120}{18} = \frac{180}{18} \text{ or } 10 \text{ in the second case,}$$

$$\frac{300}{18} = 16\frac{2}{3} \text{ in the third case.}$$

10. If two perfectly elastic bodies impinge on each other; their relative velocity is the same both before and after the impulse; that is, they will recede from each other with the same velocity with which they approached and met.

It is not meant however by this theorem, that each body will have the very same velocity after the impulse as it had before; but that the velocity of the one, after the stroke, will be as much increased, as that of the other is decreased, in one and the same direction. So, if the elastic body B move with the velocity V, and overtake the elastic body b, moving the same way, with the velocity v; then their relative velocity, or that with which they strike, is only  $V - v$ ; and it is with this same velocity that they separate from each other after the stroke: but if they meet each other, or the body b move contrary to the body B; then they meet and strike with the velocity  $V + v$ , and it is with the same velocity that they separate again, and recede from each other after the stroke: in like manner, they would separate with the velocity V of B, if b were at rest before the stroke. Also the sum of the velocities of the one body, is equal to the sum of the others. But whether they move forwards or backwards after the impulse, and with what particular velocities, are circumstances that depend on the various masses and velocities of the bodies before the stroke, and are as specified in the next theorem.

11. If the two elastic bodies B and b move directly towards each other, or directly from each other, the former with the velocity V, and the latter with the velocity v; then, after their meeting and impulse, the respective velocities of B and b in the direction BC, in the three cases of motion, will be as follow: viz,

$$\frac{2bv + B - b \cdot V}{B + b} \text{ the velocity of B,}$$

$$\frac{2BV - B - b \cdot v}{B + b} \text{ the velocity of b,}$$

when the bodies both moved towards C before the stroke; and

—2bv



$$\frac{-2bv + \overline{B - b} \cdot V}{B + b} \text{ the velocity of } B,$$

$$\frac{2BV + \overline{B - b} \cdot v}{B + b} \text{ the velocity of } b,$$


when  $B$  moved towards  $C$ , and  $b$  towards  $B$  before the stroke;

$$\frac{B - b}{B + b} \times V \text{ the velocity of } B,$$

$$\frac{2B}{B + b} \times V \text{ the velocity of } b,$$

when  $b$  was at rest before the stroke.

12 The motions of bodies after impact, that strike each other obliquely, are thus determined.



Let the two bodies  $B$ ,  $b$ , move in the oblique directions  $BA$ ,  $bA$ , and strike each other at  $A$  with velocities which are in proportion to the lines  $BA$ ,  $bA$ . Let  $CAH$  represent the plane in which the bodies touch in the point of concourse; to which draw the perpendiculars  $BC$ ,  $bD$ , and complete the rectangles  $CE$ ,  $DF$ . Now the motion in  $BA$  is resolved into the two  $BC$ ,  $CA$ ; and the motion in  $bA$  is resolved into the two  $bD$ ,  $DA$ ; of which the antecedents  $BC$ ,  $bD$  are the velocities with which they directly meet, and the consequents  $CA$ ,  $DA$  are parallel, and therefore by these the bodies do not impinge on each other, and consequently the motions according to these directions will not be changed by the impulse; so that the velocities with which the bodies meet, are as  $BC$  or  $EA$ , and  $bD$  or  $FA$ . The motions therefore of the bodies  $B$ ,  $b$ , directly striking each other with the celerities  $EA$ ,  $FA$ , will be determined by art. 11 or 9, according as the bodies are elastic or non-elastic; which being done, let  $AG$  be the velocity, so determined, of one of them, as  $A$ ; and since there remains also in the same body a force of moving in the direction parallel to  $BE$ , with a velocity as  $BE$ , make  $AH$  equal to  $BE$ , and complete the rectangle  $GH$ : then the two motions in  $AH$  and  $AG$ , or  $HI$ , are compounded into the diagonal  $AI$ , which therefore will be the path and celerity of the body  $B$  after the stroke. And after the same manner is the motion of the other body  $b$  determined after the impact.

13. The state of the common centre of gravity of bodies is not affected by the collision or other actions of those bodies on one another. That is, if it were at rest before their collision, so will it be also at rest after collision: and if it were moving in any direction, and with any velocity, before collision; it will do the very same after it.

See more upon this subject under the article *PERCUSSION*.

**COLONNADE**, a Peristyle of a circular figure; or a series of columns disposed in a circle, and insulated within-side.

**COLOUR**, a property inherent in light, by which, according to the various sizes of its parts, or from some other cause, it excites different vibrations in the optic nerve; which, propagated to the sensorium, affect the mind with different sensations. See the doctrine of colours fully explained under *Chromatics*. See also *Optics*, *Achromatic*, and *Telescope*.

**COLUMBA Noachi**, Noah's Dove, a small constellation in the southern hemisphere, consisting of 10 stars.

**COLUMN**, in Architecture, a round pillar, made to support or adorn a building.

The column is the principal part of an architectural order, and is composed of three parts, the *base*, the *shaft*, and the *capital*; each of which is subdivided into a number of lesser parts, called members, or mouldings.

Columns are different according to the different orders they are used in; and also according to their matter, construction, form, disposition, and use. The proportion of the length of each to its diameter, and the diminution of the diameter upwards, are diversely stated by different authors. The medium of them is nearly as follows:

The *Tuscan* is the simplest and shortest of all; its height  $3\frac{1}{2}$  diameters, or 7 modules; and it diminishes  $\frac{1}{4}$  part of its diameter.

The *Doric* is more delicate, and adorned with flutings; its height  $7\frac{1}{2}$  or 8 diameters.

The *Ionic* is more delicate still, being 9 diameters long. It is distinguished from the rest by the volutes, or curled scrolls in its capital, and by its base which is peculiar to it.

The *Corinthian* is the richest and most delicate of all the columns, being 10 diameters in length, and adorned with two rows of leaves, and stalks or stems, from whence spring out small volutes.

The *Composite* Column is also 10 diameters long, its capital adorned with rows of leaves like the Corinthian, and with angular volutes like the Ionic.

**COLURES**, are two great circles imagined to intersect at right angles in the poles of the world, and to pass, the one through the equinoctial points Aries and Libra, and the other through the solstitial points Cancer and Capricorn; from whence they are called the Equinoctial and Solstitial Colures. By thus dividing the ecliptic into four equal parts, they mark the four seasons, or quarters of the year.

It is disputed over what part of the back of Aries the equinoctial colure passed in the time of Hipparchus. Newton, in his Chronology, takes it to have been over the middle of the constellation. Father Souciet insists that it passed over the dodecatemoron of Aries, or midway between the rump and first of the tail. There are some observations in the *Philos. Trans.* number 466, concerning the position of this colure in the ancient sphere, from a draught of the constellation Aries, in the *Aratæa* published at Leyden and Amsterdam in 1652, which seem to confirm Newton's opinion; but the antiquity and authority of the original draught may still remain in question.

**COMA BERENICES**, *Berenice's Hair*, a modern constellation



stellation of the northern hemisphere; composed of unformed stars between the Lion's tail and Bootes.

It is said that this constellation was formed by Conon, an astronomer, to console the queen of Ptolemy Euergetes, for the loss of a lock of her hair, which was stolen out of the temple of Venus, where she had dedicated it on account of a victory obtained by her husband.

The stars in this constellation are, in Tycho's catalogue 14, in Hevelius's 21, and in the Britannic catalogue 43.

COMBINATIONS, denote the alternations or variations of any number of quantities, letters, sounds, or the like, in all possible ways.

Father Merfenne gives the combinations of all the notes and sounds in music, as far as 64; the sum of which amounts to a number expressed by 90 places of figures. And the number of possible combinations of the 24 letters of the alphabet, taken first two by two, then three by three, and so on, according to Pretet's calculation, amounts to

1391724288887252999425128493402200.

Father Truchet, in Mem. de l'Acad. shews, that two square pieces, each divided diagonally into two colours, may be arranged and combined 64 different ways, so as to form so many different kinds of chequer-work: a thing that may be of use to masons, paviours, &c.

*Doctrine of COMBINATIONS.*

I. Having given any number of things, with the number in each combination; to find the number of combinations.

1. When only two are combined together.

One thing admits of no combination.

Two, *a* and *b*, admit of one only, viz *ab*.

Three, *a*, *b*, *c*, admit of three, viz *ab*, *ac*, *bc*.

Four admit of six, viz, *ab*, *ac*, *ad*, *bc*, *bd*, *cd*.

Five admit of 10, viz, *ab*, *ac*, *ad*, *ae*, *bc*, *bd*, *be*, *cd*, *ce*, *de*.

Whence it appears that the numbers of combinations, of two and two only, proceed according to the triangular numbers 1, 3, 6, 10, 15, 21, &c, which are produced by the continual addition of the ordinal series 0, 1, 2, 3, 4, 5, &c. And if *n* be the number of things, then the general formula for expressing the sum of all their combinations by twos, will be  $\frac{n \cdot n - 1}{1 \cdot 2}$ .

Thus, if *n* = 2; this becomes  $\frac{2 \cdot 1}{2} = 1$ .

If *n* = 3; it is  $\frac{3 \cdot 2}{2} = 3$ .

If *n* = 4; it is  $\frac{4 \cdot 3}{2} = 6$ . &c.

2. When three are combined together; then

Three things admit of one order, *abc*.

Four admit of 4; viz *abc*, *abd*, *acd*, *bcd*.

Five admit of 10; viz *abc*, *abd*, *abe*, *acd*, *ace*, *ade*, *bcd*, *bce*, *bde*, *cde*. And so on according to the first pyramidal numbers 1, 4, 10, 20, &c, which are formed by the continual addition of the former, or triangular numbers 1, 3, 6, 10, &c. And the general formula for any number *n* of combinations, taken by threes,

is  $\frac{n \cdot n - 1 \cdot n - 2}{1 \cdot 2 \cdot 3}$ .

So, if *n* = 3; it is  $\frac{3 \cdot 2 \cdot 1}{1 \cdot 2 \cdot 3} = 1$ .

If *n* = 4; it is  $\frac{4 \cdot 3 \cdot 2}{6} = 4$ .

If *n* = 5; it is  $\frac{5 \cdot 4 \cdot 3}{2} = 6$ . &c.

Proceeding thus, it is found that a general formula for any number *n* of things, combined by *m* at each time, is  $s = \frac{n \cdot n - 1 \cdot n - 2 \cdot n - 3 \cdot \&c}{1 \cdot 2 \cdot 3 \cdot 4 \cdot \&c}$ , continued to *m* factors, or terms, or till the last factor in the denominator be *m*.

So, in 6 things, combined by 4's, the number of combinations is  $\frac{6 \cdot 5 \cdot 4 \cdot 3}{1 \cdot 2 \cdot 3 \cdot 4} = 15$ .

3. By adding all these series together, their sum will be the whole number of possible combinations of *n* things combined both by twos, by threes, by fours, &c. And as the said series are evidently the coefficients of the power *n* of a binomial, wanting only the first two 1 and *n*; therefore the said sum, or whole number of all such combinations, will be

$1 + 1^1 = n + 1$ , or  $2^n - n - 1$ . Thus if the number of things be 5; then  $2^5 - 5 - 1 = 32 - 6 = 26$ .

II. To find the number of Changes and Alterations which any number of quantities can undergo, when combined in all possible varieties of ways, with themselves and each other, both as to the things themselves, and the Order or Position of them.

One thing admits but of one order or position.

Two things may be varied four ways; thus, *aa*, *ab*, *ba*, *bb*.

Three quantities, taken by twos, may be varied nine ways; thus *aa*, *ab*, *ac*, *ba*, *ca*, *bb*, *bc*, *cb*, *cc*.

In like manner four things, taken by twos, may be varied  $4^2$  or 16 ways; and 5 things, by twos,  $5^2$  or 25 ways; and, in general, *n* things, taken by twos, may be changed or varied  $n^2$  different ways.

For the same reason, when taken by threes, the changes will be  $n^3$ ; and when taken by fours, they will be  $n^4$ ; and so generally, when taken by *n*'s, the changes will be  $n^n$ .

Hence, then, adding all these together, the whole number of changes, or combinations in *n* things, taken both by 2's, by 3's, by 4's, &c, to *n*'s, will be the sum of the geometrical series  $n + n^2 + n^3 + n^4 + \dots + n^n$ , which sum is  $= \frac{n^n - 1}{n - 1} \times n$ .

For example, if the number of things *n* be 4; this gives  $\frac{4^4 - 1}{4 - 1} \times 4 = \frac{255}{3} \times 4 = 340$ .

And if *n* be 24, the number of letters in the alphabet; the theorem gives

$$\frac{24^{24} - 1}{24 - 1} \times 24 = \frac{24^{24} - 1}{23} \times 24 =$$

1391724288887252999425128493402200. In so many different ways, therefore, may the 24 letters of the alphabet be varied or combined among themselves, or so many different words may be made out of them.

COMBUST, or COMBUSTION, is said of a planet when



when it is in conjunction with the sun, or not distant from it above half their disc.

But according to Argol, a planet is Combust, or in Combustion, when it is within eight degrees and a half of the sun.

COMET, a heavenly body in the planetary region, appearing suddenly, and again disappearing; and during the time of its appearance moving in a proper, though very eccentric orbit, like a planet.

Comets are vulgarly called *Blazing Stars*, and have this to distinguish them from other stars, that they are usually attended with a long train of light, tending always opposite to the sun, and being of a fainter lustre the farther it is from the body of the comet. And hence arises a popular division of comets, into three kinds; viz *bearded*, *tailed*, and *hairy* comets; though in reality, this division rather relates to the several circumstances of the same comet, than to the phenomena of several. Thus, when the comet is eastward of the sun, and moves from him, it is said to be *bearded*, because the light precedes it in the manner of a beard: When the comet is westward of the sun, and sets after him, it is said to be *tailed*, because the train of light follows it in the manner of a tail: And lastly, when the sun and comet are diametrically opposite, the earth being between them, the train is hid behind the body of the comet, excepting the extremities, which, being broader than the body of the comet, appear as it were around it, like a border of hair, or *coma*, from which it is called *hairy*, and a comet.

But there have been comets whose disc was as clear, round, and well defined, as that of Jupiter, without either tail, beard or coma.

*Of the Nature of Comets.*—Philosophers and Astronomers, of all ages, have been much divided in their opinions as to the nature of comets. Their strange appearance has in all ages been matter of terror to the vulgar, who have uniformly considered them as evil omens, and forerunners of war, pestilence, &c. Diodorus Siculus and Appollinus Myndius, in Seneca, inform us, that many of the Chaldeans held them to be lasting bodies, having stated revolutions as well as the planets, but in orbits vastly more extensive; on which account they are only visible while near the earth, but disappear again when they go into the higher regions. Others of them were of opinion, that the comets were only meteors raised very high in the air, which blaze for a while, and disappear when the matter of which they consist is consumed or dispersed.

Some of the Greeks, before Aristotle, supposed that a comet was a vast heap or assemblage of very small stars meeting together, by reason of the inequality of their motions, and so uniting into a visible mass, by the union of all their small lights; which must again disappear, as those stars separated, and each proceeded in its course. Pythagoras, however, accounted them a kind of planets or wandering stars, disappearing in the superior parts of their orbits, and becoming visible only in the lower parts of them.

But Aristotle held, that comets were only a kind of transient fires, or meteors, consisting of exhalations raised to the upper region of the air, and there set on fire; far below the course of the moon.

Seneca, who lived in the first century, and who had seen two or three comets himself, plainly intimates that he thought them above the moon; and argues strongly against those who supposed them to be meteors, or who held other absurd opinions concerning them; declaring his belief that they were not fires suddenly kindled, but the eternal productions of nature. He points out also the only way to come at a certainty on this subject, viz, by collecting a number of observations concerning their appearance, in order to discover whether they return periodically or not. "For this purpose, says he, one age is not sufficient; but the time will come when the nature of comets and their magnitudes will be demonstrated, and the routes they take, so different from the planets, explained. Posterity will then wonder, that the preceding ages should be ignorant of matters so plain and easy to be known."

For a long time this prediction of Seneca seemed not likely to be fulfilled; and Tycho Brahe was the first among the moderns, who restored the comets to their true rank in the creation; for after diligently observing the comet of 1577, and finding that it had no sensible diurnal parallax, he assigned it its true place in the planetary regions. See his book *De Cometa*, anni 1577.

Before this however, there were various opinions concerning them. In the dark and superstitious ages, comets were held to be forerunners of every kind of calamity, and it was supposed they had different degrees of malignity, according to the shape they assumed; from whence also they were differently denominated. Thus, it was said that some were bearded, some hairy; that some represented a beam, sword, or spear; others a target, &c; whereas modern astronomers acknowledge only one species of comets, and account for their different appearances from their different situations with respect to the sun and earth.

Kepler, in other respects a very great genius, indulged the most extravagant conjectures, not only concerning comets, but the whole system of nature in general. The planets he imagined were huge animals swimming round the sun; and the comets monstrous and uncommon animals generated in the celestial spaces.

A still more ridiculous opinion, if possible, was that of John Bodin, a learned Frenchman in the 16th century; who maintained that comets "are spirits, which having lived on the earth innumerable ages, and being at last arrived on the confines of death, celebrate their last triumph, or are recalled to the firmament like shining stars! This is followed by famine, plague, &c, because the cities and people destroy the governors and chiefs who appease the wrath of God."—Others again have denied even the existence of comets, and maintained that they were only false appearances, occasioned by the refraction or reflection of light.

Hevelius, from a great number of observations, proposed it as his opinion, that the comets, like the solar maculæ or spots, are formed or condensed out of the grosser exhalations of his body; in which he differs but little from the opinion of Kepler.

James Bernoulli, in his *Systema Cometarum*, imagined that comets were no other than the satellites of some



some very distant planet, which was itself invisible to us on account of its distance, as were also the satellites unless when in a certain part of their orbits.

Des Cartes advances another opinion: He conjectures that comets are only stars, formerly fixed, like the rest, in the heavens; but which becoming gradually covered with maculæ or spots, and at length wholly deprived of their light, cannot keep their places, but are carried off by the vortices of the circumjacent stars; and in proportion to their magnitude and solidity, moved in such a manner, as to be brought nearer the orb of Saturn; and thus coming within reach of the sun's light, rendered visible.

But the vanity of all these hypotheses now abundantly appears from the observed phenomena of comets, and from the doctrine of Newton, which is as follows:

The comets, he says, are compact, solid, fixed, and durable bodies; in fact a kind of planets, which move in very oblique and eccentric orbits, every way with the greatest freedom; persevering in their motions, even against the course and direction of the planets: and their tail is a very thin and slender vapour, emitted by the head or nucleus of the comet, ignited or heated by the sun. This theory of the comets at once solves their principal phenomena, which are as below.

*The Principal Phenomena of the Comets.*

1. First then, those comets which move according to the order of the signs, do all, a little before they disappear, either advance slower than usual, or else go retrograde, if the earth be between them and the sun; but more swiftly, if the earth be placed in a contrary part. On the other hand, those which proceed contrary to the order of the signs, move more swiftly than usual, if the earth be between them and the sun; and more slowly, or else retrograde, when the earth is in a contrary part.—For since this course is not among the fixed stars, but among the planets; as the motion of the earth either conspires with them, or goes against them; their appearance, with respect to the earth, must be changed; and, like the planets, they must sometimes appear to move swifter, sometimes slower, and sometimes retrograde.

2. So long as their velocity is increased, they nearly move in great circles; but towards the end of their course, they deviate from those circles; and when the earth proceeds one way, they go the contrary way. Because, in the end of their course, when they recede almost directly from the sun, that part of the apparent motion which arises from the parallax, must bear a greater proportion to the whole apparent motion.

3. The comets move in ellipses, having one of their foci in the centre of the sun; and by radii drawn to the sun, describe areas proportional to the times. Because they do not wander precariously from one fictitious vortex to another; but, making a part of the solar system, return perpetually, and run a constant round. Hence, their elliptic orbits being very long and eccentric, they become invisible when in that part which is most remote from the sun. And from the curvity of the paths of comets, Newton concludes, that when they disappear, they are much beyond the orbit of Jupiter; and that in their perihelion they frequently descend within the orbits of Mars and the inferior planets.

VOL. I.

4. The light of their nuclei, or bodies, increases as they recede from the earth toward the sun; and on the contrary, it decreases as they recede from the sun. Because, as they are in the regions of the planets, their access towards the sun bears a considerable proportion to their whole distance.

5. Their tails appear the largest and brightest, immediately after their transit through the region of the sun, or after their perihelion. Because then, their heads being the most heated, will emit the most vapours.—From the light of the nucleus we infer their vicinity to the earth, and that they are by no means in the region of the fixed stars, as some have imagined; since, in that case, their heads would be no more illuminated by the sun, than the planets are by the fixed stars.

6. The tails always incline from a just opposition to the sun towards those parts which the nuclei or bodies pass over, in their progress through their orbits. Because all smoke, or vapour, emitted from a body in motion, tends upwards obliquely, still receding from that part towards which the smoking body proceeds.

7. This declination, *cæteris paribus*, is the smallest when the nuclei approach nearest the sun: and it is also less near the nucleus, or head, than towards the extremity of the tail. Because the vapour ascends more swiftly near the head of the comet, than in the higher extremity of its tail; and also when the comet is nearer the sun, than when it is farther off.

8. The tails are somewhat brighter, and more distinctly defined in their convex, than in their concave part. Because the vapour in the convex part, which goes first, being somewhat nearer and denser, reflects the light more copiously.

9. The tails always appear broader at their upper extremity, than near the centre of the comet. Because the vapour in a free space continually rarefies and dilates.

10. The tails are always transparent, and the smallest stars appear through them. Because they consist of infinitely thin vapour.

*The Phases of Comets.*—The nuclei, which are also called the heads, and bodies, of comets, viewed through a telescope, shew a face very different from those of the fixed stars or planets. They are liable to apparent changes, which Newton ascribes to changes in the atmosphere of comets: and this opinion was confirmed by observations of the comet in 1744. *Hist. Acad. Scienc.* 1744. Sturmius says that, observing the comet of 1680 with a telescope, it appeared like a coal dimly glowing, or a rude mass of matter illuminated with a dusky humid light, less sensible to the extremes than in the middle; whereas a star appears with a round disc, and a vivid light.

Of the comet of 1661, Hevelius observes, that its body was of a yellowish colour, very bright and conspicuous, but without any glittering light: in the middle was a dense ruddy nucleus, almost equal to Jupiter, encompassed by a much fainter, thinner matter. February 5th, its head was somewhat larger and brighter, and of a gold colour; but its light more dusky than the stars: and here the nucleus appeared divided into several parts. Feb. 6th, the disc was lessened; the parts of the nucleus still existed, though less

R r



less than before: one of them, on the lower part of the disc, on the left, much denser and brighter than the rest; its body round, and representing a very lucid little star: the nuclei still encompassed with another kind of matter. Feb. 10th, the head somewhat more obscure, and the nuclei more confused, but brighter at top than bottom. Feb. 13th, the head diminished much both in size and splendor. March 2d, its roundness a little impaired, and its edges lacerated, &c. March 28th, very pale, and exceeding thin; its matter much dispersed; and no distinct nucleus at all appearing.

Weigelius, who saw the comet of 1664, as also the moon, and a small cloud in the horizon illuminated by the sun at the same time, observed, that through the telescope the moon appeared of a continued luminous surface: but the comet very different; being exactly like the little cloud. And from these observations it was that Hevelius formed his opinion, that comets are like maculæ or spots formed out of the solar exhalations.

*Of the Magnitude of Comets.*—The estimates that have been given of the magnitude of comets by Tycho Brahe, Hevelius, and some others, are not very accurate; as it does not appear that they distinguished between the nucleus and the surrounding atmosphere. Thus Tycho computes that the true diameter of the comet in 1577 was in proportion to the diameter of the earth, as 3 is to 14; and Hevelius made the diameter of the comet of 1652 to that of the earth, as 52 to 100. But the diameter of the atmosphere is often 10 or 15 times as great as that of the nucleus: the former, in the comet of 1682, was measured by Flamsteed, and found to be 2', when the diameter of the nucleus alone was only 11 or 12". Though some comets, estimated by a comparison of their distance and apparent magnitude, have been judged much larger than the moon, and even equal to some of the primary planets. The diameter of that of 1744, when at the distance of the sun from us, measured about 1', which makes its diameter about three times that of the earth: at another time the diameter of its nucleus was nearly equal to that of the planet Jupiter.

*Of the Tails of Comets.*—There have been various conjectures about the nature of the tails of comets, the principal of which are those of Newton, and the others that follow. Newton shews that the atmospheres of comets will furnish vapour sufficient to form their tails. This he argues from that wonderful rarefaction in our air at a distance from the earth; which is such, that a cubic inch of common air, expanded to the rarity of that at the distance of half the earth's diameter, or 4000 miles, would fill a space larger than the whole region of the stars. Since then the coma, or atmosphere of a comet, is 10 times higher than the surface of the nucleus, from the centre; the tail, ascending still much higher, must necessarily be immensely rare: so that it is no wonder the stars are visible through it.

Now the ascent of vapours into the tail of the comet, he supposes occasioned by the rarefaction of the matter of the atmosphere at the time of the perihelion. Smoke, it is observed, ascends the chimney by the impulse of the air in which it floats; and air, rarefied by

heat, ascends by the diminution of its specific gravity, carrying up the smoke along with it: in the same manner then it may be supposed that the tail of a comet is raised by the sun.

The tails therefore thus produced in the perihelions of comets, will go off along with their head into remote regions; and either return from thence, together with the comets, after a long series of years; or rather be there lost, and vanish by little and little, and the comet be left bare; till at its return, descending towards the sun, some short tails are again gradually produced from the head; which afterwards, in the perihelion, descending down into the sun's atmosphere, will be immensely increased.

Newton farther observes, that the vapours, when thus dilated, rarefied, and diffused through all the celestial regions, may probably, by means of their own gravity, be gradually attracted down to the planets, and become intermingled with their atmospheres. He adds that this intermixture may be useful and necessary for the conservation of the water and moisture of the planets, dried up or consumed in various ways. And I suspect, adds our author, that the spirit, which makes the finest, subtlest, and best part of our air, and which is absolutely requisite for the life and being of all things, comes principally from the comets.—On this principle there may seem to be some foundation for the popular opinion of presages from comets; since the tail of a comet thus intermingled with our atmosphere, may produce changes very sensible in animal and vegetable bodies.

It may here be added that another use which Newton conjectures comets may be designed to serve, is that of recruiting the sun with fresh fuel, and repairing the consumption of his light by the streams continually sent forth in every direction from that luminary. In support of this conjecture he observes, that comets in their perihelion may suffer a diminution of their projectile force, by the resistance of the solar atmosphere; so that by degrees their gravitation towards the sun may be so far increased, as to precipitate their fall into his body.

Other opinions on the tails of comets, are the following.

Apian, Tycho Brahe, and some others, think they were produced by the sun's rays transmitted through the nucleus of the comet, which they supposed was transparent, and there refracted as in a glass lens, so as to form a beam of light behind the comet. Des Cartes accounted for the phenomenon of the tail by the refraction of light from the head of the comet to the spectator's eye. Mairan supposes that the tails are formed out of the luminous matter composing the sun's atmosphere: and M. De la Lande combines this hypothesis with that of Newton recited above. But Mr. Rowning, not satisfied with Newton's opinion, accounts for the tails of comets in the following manner: It is well known, says he, that when the sun's light passes through the atmosphere of any body, as the earth, that which passes on one side, is by the refraction made to converge towards that which passes on the opposite side; and this convergency is not wholly effected either at the entrance of the light into the atmosphere, or at its exit on going out; but beginning at



its entrance, it increases in every point of its progress: It is also agreed that the atmospheres of the comets are very large and dense: he therefore supposes that by such time as the light of the sun has passed through a considerable part of the atmosphere of the comet, the rays are so far refracted towards each other, that they then begin sensibly to illuminate it, or rather the vapours floating in it; and so render that part they have yet to pass through, visible to us: and that this portion of the atmosphere of a comet thus illuminated, appears to us in form of a beam of the sun's light, and passes under the denomination of a comet's tail, Rowning's Nat. Philos. part 4. chap. 11.

M. Euler, Mem. Berlin tom. 2. pa. 117, thinks there is a great affinity between the tails of comets, the zodiacal light, and the aurora borealis, and that the common cause of all of them, is the action of the sun's light on the atmospheres of the comets, of the sun, and of the earth. He supposes that the impulse of the rays of light on the atmosphere of comets, may drive some of the finer particles of that atmosphere far beyond its limits; and that this force of impulse combined with that of gravity towards the comet, would produce a tail, which would always be in opposition to the sun, if the comet did not move. But the motion of the comet in its orbit, and about an axis, must vary the position and figure of the tail, giving it a curvature, and deviation from a line joining the centres of the sun and comet; and that this deviation will be greater, as the orbit of the comet has the greater curvature, and as the motion of the comet is more rapid. It may even happen, that the velocity of the comet, in its perihelion, may be so great, that the force of the sun's rays may produce a new tail, before the old one can follow; in which case the comet might have two or more tails. The possibility of this is confirmed by the comet of 1744, which was observed to have several tails while it was in its perihelion.

Dr. Hamilton urges several objections against the Newtonian hypothesis; and concludes that the tail of a comet is formed of matter which has not the power of refracting or reflecting the rays of light; but that it is a lucid or self-shining substance: and from its similarity to the Aurora borealis, that it is produced by the same cause, and is properly an electrical phenomenon. Dr. Halley too seemed inclined to this hypothesis, when he said, that the streams of light in an Aurora borealis so much resembled the long tails of comets, that at first sight they might well be taken for such: and that this light seems to have a greater affinity to that which the effluvia of electric bodies emit in the dark. Philos. Trans. N° 347. Hamilton's Philos. Essays, pa. 91.

*The Motion of Comets.*—If it be supposed that the paths of comets are perfectly parabolical, as some have imagined, it will follow that, being impelled towards the sun by a centripetal force, they descend as from spaces infinitely distant; and that by their falls they acquire such a velocity as will carry them off again into the remotest regions, never more to return. But the frequency of their appearance, and their degree of velocity, which does not exceed what they might acquire by their gravity towards the sun, seem to put it past doubt that they move like the planets, in elliptic orbits,

though exceedingly eccentric; and so return again after very long periods.

The apparent velocity of the comet of 1472, as observed by Regiomontanus, was such as to carry it through 40° of a great circle in 24 hours: and it was observed that the comet of 1770 moved through more than 45° in the last 25 hours.

About the return of comets there have been different opinions. Newton, Flamsteed, Halley, and other English astronomers, seem satisfied of the return of comets: Cassini and some of the French think it highly probable; but De la Hire and others oppose it. Those on the affirmative side suppose that the comets describe orbits prodigiously eccentric, inasmuch that we can see them only in a very small part of their revolution: out of this, they are lost in the immensity of space; hid not only from our eyes, but our telescopes: that little part of their orbit next us passing sometimes within those of all the inferior planets.

M. Cassini gives the following reasons in favour of the return of comets. 1. It is found that they move a considerable time in the arch of a great circle, when referred to the fixed stars, that is a circle whose plane passes through the centre of the earth; deviating but a little from it chiefly towards the end of their appearance; a deviation however common to them with the planets.—2. Comets, as well as planets, appear to move so much the faster as they are nearer the earth; and when they are at equal distances from their perigee, their velocities are nearly the same. By subtracting from their motion the apparent inequality of velocity occasioned by their different distance from the earth, their equal motion might be found: but we should not still be certain that this is their true motion; because they might have considerable inequalities, not distinguishable in that small part of their orbit visible to us. It is rather probable that their real motion, as well as that of the planets, is unequal in itself; and hence we have a reason why the observations made during the appearance of a comet, cannot give the just period of their revolution.—3. There are no two different planets whose orbits cut the ecliptic in the same angle, whose nodes are in the same points of the ecliptic, and having the same apparent velocity in their perigee: consequently, two comets seen at different times, yet agreeing in all those three circumstances, can only be one and the same comet. Not that this exact agreement, in these circumstances, is absolutely necessary to determine their identity: for the moon herself is irregular in all of them, so that it seems there may be cases in which the same comet, at different periods of revolution, may disagree in these points.

As to the objections against the return of comets, the principal is that of the rarity of their appearance, with regard to the number of revolutions assigned to them. In 1702 there was a comet, or rather the tail of one, seen at Rome, which M. Cassini takes to be the same with that observed by Aristotle, and again lately in the year 1668; which would imply a period of 34 years: Now, it may seem strange that a star which has so short a revolution, and of consequence such frequent returns, should be so seldom seen. Again, in April of the same year 1702, a comet was observed by Messrs. Bianchini and Maraldi, which the latter sup-



posed was the same with that of 1664, both on account of its motion, velocity, and direction. M. de la Hire thought it had some relation to another he had observed in 1698, which Cassini refers to that of 1652; which would make it a period of 43 months, and the number of revolutions between 1652 and 1692, 14: now, it is hard to suppose, that in this age, when the heavens are so narrowly watched, a star should make 14 revolutions unperceived; especially such a star as this, which might appear above a month together; and consequently be often disengaged from the crepuscula. For this reason M. Cassini is very reserved in maintaining the hypothesis of the return of comets, and only proposes those for planets where the motions are easy and simple, and are solved without straining, or allowing any irregularities.

M. de la Hire proposes one general difficulty against the whole system of the return of comets, which would seem to prevent any comet from returning as a planet: which is this; that by the disposition necessarily given to their courses, they ought to appear as small at first as at last; and always increase till they arrive at their nearest proximity to the earth; or if they should chance not to be observed, as soon as they are capable of being seen, it is yet hardly possible but they must often shew themselves before they have arrived at their full magnitude and brightness: but, adds he, none were ever yet observed till they had arrived at it. However, the appearance of a comet in the month of October 1723, while at a great distance, so as to be too small and dim to be viewed without a telescope, as well as the observations of several others since, may serve to remove this obstacle, and set the comets still on the same footing with the planets.

It is a conjecture of Newton, that as those planets which are nearest to the sun, and revolve in the least orbits, are the smallest; so among the comets, such as in their perihelion come nearest the sun, are the smallest, and revolve in the least orbits.

#### *Of the Writings and Lists of Comets.*

There have been many writings upon the subject of comets, beside the notices of historians as to the appearance of certain particular ones.

Regiomontanus first shewed how to find the magnitudes of comets, their distance from the earth, and their true place in the heavens. His 16 problems De Co-

metæ Magnitudine, Longitudine, ac Loco, are to be found in a book published in the year 1544, with the title of Scripta Joannis Regiomontani.

Peter Apian observed and wrote upon the comets of 1631, 1632, &c. Other writers are Tycho Brahe, in his *Progymnasmata Astronomiæ Instauratæ*.—Kepler, of the comet in the year 1607, and *de Cometis Libellus* tres.—Ricciolus, in his *Almagestum Novum*, published 1651, enumerates 154 comets cited by historians down to the year 1618.—Hevelius's *Prodromus Cometicus*, containing the history of the comet of the year 1664. Also his *Cometographia*.—Lubienietz, in a large folio work expressly on this subject, published 1667, extracts, with immense labour, from the passages of all historians, an account of 415 comets, ending with that of 1665.—Dr. Hook, in his *Posthumous works*.—M. Cassini's little *Tract of Comets*.—Sturmius's *Differtatio de Cometarum Natura*.—Newton, in his *Principia*, lib. 3; who first assigned their proper orbits, and by calculations compared the observations of the great comet of 1680 with his theory.—Dr. Halley, his *Synopsis Cometica*, in the *Philos. Transf.* number 218, &c; who computed the elements and orbits of 24 comets, and who first ventured to predict the return of one in 1759, which happened accordingly.—De la Lande, *Théorie des Comètes*, 1759; also, in his *Astronomie*, vol. 3.—Clairaut, *Théorie du mouvement des Comètes*, 1760.—D'Alembert, *Opusculs Mathématiques*, vol. 2 pa. 97.—M. Albert Euler, 1762.—Séjour, *Essai sur les Comètes*, 1775.—Besides Boscovich, De la Grange, De la Place, Frisi, Lexel, Barker, Hancocks, Cole, with many others.—And M. Pingré's *Cométographie*, in 2 vols. 4to, 1784; in which is contained the most ample list of such comets as have been well observed, and their elements computed, to the number of 67. And accounts of a very few more that have been observed since that time, may be seen in the *Mem. de l'Acad.* and in the *Philos. Transf.*—And while this work is printing, there has just come out a very ingenious and ample work upon comets, by Sir Henry Englefield, entitled, "On the Determination of the Orbits of Comets."

The whole list of comets that have been noticed, on record, amount to upwards of 500, but the following is a complete list of all that have been properly observed, and their elements computed, the mean distance of the earth from the sun being 100000.

TABLE OF THE ELEMENTS OF COMETS.

Year	Ascending Node	Inclin. of Orbit	Perihelion	Perihel. Dist.	Time of Perihelion	Motion	Calculated by
	s o '	o '	s o '		d h		
837	6 26 33	11 abt.	9 19 3	58000	March 1 0	Ret.	Pingré
1231	0 13 30	6 5	4 14 48	94776	Jan. 30 7	Dir.	Pingré
1264	5 19 0	36 30	9 21 0	44500	July 6 8	Dir.	Duntherne
	5 28 45	30 25	9 5 45	41081	17 6	Dir.	Pingré
1299	3 17 8	68 57	0 3 20	31793	March 31 8	Ret.	Pingré
1301	0 15 about	70 abt.	9 about	45700	Oct. 22 about	Ret.	Pingré
1337	2 24 21	32 11	1 7 59	40666	June 2 7	Ret.	Halley
	2 6 22	32 11	0 20 0	64450	1 1	Ret.	Pingré
1456	1 18 30	17 56	10 1 0	58550	June 8 22	Ret.	Pingré
1472	9 11 46	5 20	1 15 34	54273	Feb. 28 23	Ret.	Halley



Year	Ascending Node	Inclin. of Orbit	Perihelion	Perihel. Dist.	Time of Perihelion	Motion	Calculated by
	s o /	o /	s o /		d h		
1531	1 19 25	17 56	10 1 39	56700	Aug. 24 21	Ret.	Halley
1532	2 20 27	32 36	3 21 7	50910	Oct. 19 22	Dir.	Halley
1533	4 5 44	35 49	4 27 16	20280	June 16 20	Ret.	Douwes
1556	5 25 42	32 7	9 8 50	66390	April 21 20	Dir.	Halley
1577	0 25 52	74 33	4 9 22	18342	Oct. 26 19	Ret.	Halley
1580	0 18 57	64 40	3 19 6	59628	Nov. 28 15	Dir.	Halley
	0 19 8	64 52	3 19 12	59553	28 14	Dir.	Pingré
1582	7 21 7	61 28	8 5 23	22570	May 6 16	Ret.	Pingré
	7 4 43	59 29	9 11 27	4006	7 9	Ret.	Pingré
					NEW STYLE		
1585	1 7 43	6 4	0 8 51	109358	Oct. 7 20	Dir.	Halley
1590	5 15 31	29 41	7 6 55	57661	Feb. 8 4	Ret.	Halley
1593	5 14 15	87 58	5 26 19	8911	July 18 14	Dir.	La Caille
1596	10 12 13	55 12	7 18 16	51293	Aug. 10 20	Ret.	Halley
	10 15 37	52 10	7 28 31	54942	8 16	Ret.	Pingré
1607	1 20 21	17 2	10 2 16	58680	Oct. 16 4	Ret.	Halley
1618	9 23 25	21 28	10 18 20	51298	Aug. 17 3	Dir.	Pingré
1618	2 16 1	37 34	0 2 14	37975	Nov. 8 13	Dir.	Halley
1652	2 28 10	79 28	0 28 19	84750	Nov. 12 16	Dir.	Halley
1661	2 22 31	32 36	3 25 59	44851	Jan. 26 24	Dir.	Halley
1664	2 21 14	21 19	4 10 41	102575	Dec. 4 12	Ret.	Halley
1665	7 18 2	76 5	2 11 55	10649	April 24 5	Ret.	Halley
1672	9 27 31	83 22	1 17 0	69739	March 1 9	Dir.	Halley
1677	7 26 49	79 3	4 17 37	28059	May 6 1	Ret.	Halley
1678	5 11 40	3 4	10 27 46	123801	Aug. 26 14	Dir.	Douwes
1680	9 2 2	60 56	8 22 40	612½	Dec. 18 0	Dir.	Halley
	9 2 2	61 7	8 22 44	617	17 23	...	Halley
	9 2 59	58 40	8 23 27	656	17 21	...	Euler
	9 1 53	61 20	8 23 43	592	18 0	...	Newton
	9 1 57	61 23	8 22 40	603	18 0	...	Pingré
1682	1 21 17	17 56	10 2 53	58328	Sept. 14 8	Ret.	Halley
	1 20 48	17 42	10 1 36	58250	14 22	...	Halley
1683	5 23 23	83 11	2 25 30	56020	July 13 3	Ret.	Halley
1684	8 28 15	65 49	7 28 52	96015	June 8 10	Dir.	Halley
1686	11 20 35	31 22	2 17 1	32500	Sept. 16 15	Dir.	Halley
1689	10 23 45	69 17	8 23 45	1689	Dec. 1 15	Ret.	Pingré
1698	8 27 44	11 46	9 0 51	69129	Oct. 18 17	Ret.	Halley
1699	10 21 46	69 20	7 2 31	74400	Jan. 13 9	Ret.	La Caille
1702	6 9 25	4 30	4 18 41	64590	March 13 14	Dir.	La Caille
1706	0 13 12	55 14	2 12 29	42581	Jan. 30 5	Dir.	La Caille
	0 13 11	55 14	2 12 36	42686	30 5	...	Struyck
1707	1 22 8	88 50	2 17 4	86350		Dir.	Houtteryn
	1 22 47	88 36	2 19 55	85974	Dec. 12 0	...	La Caille
	1 22 50	88 38	2 19 58	85904	12 0	...	Struyck
1718	4 8 43	30 20	4 1 30	102655	Jan. 15 0	Ret.	La Caille
	4 7 55	31 13	4 1 27	102565	15 1	...	Douwes
1723	0 14 16	49 59	1 12 52	99865	Sept. 27 16	Ret.	Bradley
1729	10 10 35	77 2	10 22 17	406980	June 23 7	Dir.	Douwes
	10 10 33	76 58	10 22 40	426140	25 11	...	La Caille
	10 10 17	76 53	10 27 22	416927	23 0	...	Maraldi
	10 10 52	77 19	10 16 27	394927	May 22 11	...	Kies
	10 10 33	77 1	10 22 37	408165	June 25 9	...	De l'Isle
1737	7 16 22	18 21	10 25 55	22282	Jan. 30 9	Dir.	Bradley
1739	6 27 18	55 53	3 12 34	67160	June 17 11	Ret.	Zanotti
	6 27 25	55 43	3 12 39	67358	17 10	...	La Caille
1742	6 5 35	67 4	7 7 33	76555	Feb. 8 5	Ret.	Struyck
	6 5 33		7 7 32	76550	8 4	...	Le Monnier
	6 5 38	66 59	7 7 35	76568	8 5	...	La Caille
	6 5 43	66 52	7 7 39	76530	8 8	...	Zanotti
	6 16 9	56 35	7 16 42	73766	1 22	...	Euler



Year	Ascending Node	Inclm. of Orbit	Perihelion	Perihel. Dist.	Time of Perihelion	Motion	Calculated by
	s o "	o "	s o "		d h		
1742	6 9 32	61 44	7 10 49	73668	Feb. 7 4	Ret.	Euler
	6 5 47	68 14	7 7 33	76890	7 22	..	Wright
	6 5 29	67 11	7 7 26	76620	8 5	...	Klinkenberg
	6 5 42	66 51	7 7 38	76545	8 7	...	Houtteryn
1743	2 8 11	2 16	3 2 58	83811	Jan. 10 21	Dir.	Struyck
	2 8 21	2 20	3 2 42	83501	10 21	...	La Caille
1743	0 5 16	45 48	8 6 34	52057	Sept. 20 21	Ret.	Klinkenberg
1744	1 15 45	47 9	6 17 13	22206	March 1 8	Dir.	Bets and Bliss
	1 15 47	47 4	6 17 6	22322	1 8	...	Maraldi
1744	1 15 46	47 5	6 17 10	22250	March 1 8	Dir.	La Caille
	1 15 51	47 18	6 17 18	22156	1 8	...	Zanotti
	1 16 5	47 50	6 17 19	22192	1 9	...	Chéseaux
	1 15 46	47 11	6 17 12	22222	1 8	...	Euler
	1 15 48	47 8	6 17 13	22223	1 8	...	Pingré
	1 15 49	47 18	6 17 15	22200	1 8	...	Klinkenberg
	1 15 49	47 14	6 17 16	22176	1 9	...	Hiorter
1747	4 26 58	77 57	9 10 6	229388	Feb. 28 12	Ret.	Chéseaux
	4 27 19	79 7	9 7 2	219859	March 3 10	...	Maraldi
	4 27 19	79 6	9 7 2	219851	3 7	...	La Caille
1748	7 22 52	85 27	7 5 1	84066	April 28 20	Ret.	Maraldi
	7 22 46	85 35	7 4 39	84150	29 1	...	Le Monnier
	7 22 52	85 28	7 5 23	84040	28 19	...	La Caille
1748	1 4 40	56 59	9 6 9	65525	June 18 2	Dir.	Struyck
1757	7 4 13	12 50	4 2 58	33754	Oct. 21 8	Dir.	Bradley
	7 4 6	12 39	4 2 39	33907	21 10	...	La Caille
	7 4 4	12 48	4 2 49	33797	21 10	...	Pingré
	7 4 7	12 41	4 2 36	33932	21 9	...	De Ratte
1758	7 20 50	68 19	8 27 38	21535	June 11 3	Dir.	Pingré
1759	1 23 48	17 38	10 3 14	58255	March 12 14	Ret.	Mellier
	1 23 46	17 40	10 3 8	58490	12 14	...	De la Lande
	1 23 49	17 35	10 3 16	58360	12 13	...	Maraldi
	1 23 49	17 38	10 3 16	58380	12 13 <sup>1</sup> / <sub>2</sub>	...	La Caille
	1 23 49	17 39	10 3 16	58350	12 13 <sup>2</sup> / <sub>3</sub>	...	La Caille
	1 23 46	17 40	10 3 19	58298	12 13	...	Klinkenberg
	1 24 7	17 29	10 1 0	59708	13 10	...	Klinkenberg
	1 23 45	17 41	10 3 23	58234	12 13	...	Bailly
1759	4 19 40	79 7	1 23 34	80139	Nov. 27 0	Dir.	Pingré
	4 19 39	78 59	1 23 24	79851	27 2	...	La Caille
	4 19 40	79 3	1 23 38	80208	27 1	...	Chappe
1759	2 19 51	4 52	4 18 25	96599	Dec. 16 21	Ret.	La Caille
1759	2 19 20	4 42	4 19 4	96180	Dec. 16 13	Ret.	Chappe
1762	11 18 56	85 22	3 15 22	101415	May 29 0	Dir.	Maraldi
	11 19 20	84 45	3 15 15	101249	28 15	...	De la Lande
	11 18 58	85 12	3 14 24	101065	29 2	...	Bailly
	11 18 35	85 40	3 13 43	100686	28 2	...	Klinkenberg
	11 19 2	85 3	3 14 30	100986	28 7	...	Struyck
1763	11 26 23	72 41	2 24 52	49876	Nov. 1 20	Dir.	Pingré
1764	4 0 5	52 54	0 15 15	55522	Feb. 12 14	Ret.	Pingré
1766	8 4 11	40 50	4 23 15	50532	Feb. 17 9	Ret.	Pingré
1766	2 14 23	11 8	8 2 18	33275	April 22 21	Dir.	Pingré
1769	5 25 1	40 38	4 24 6	12376	Oct. 7 13	Dir.	De la Lande
	5 25 2	40 43	4 24 14	12287	7 12	...	Wallot
	5 25 3	40 47	4 24 11	12258	7 13	...	Cassin, jun.
	5 25 7	40 49	4 24 11	12272	7 14	...	Prosperin
	5 25 3	40 41	4 24 9	12289	7 14	...	Audiffredi
	5 25 11	41 1	4 24 33	12100	7 12	...	Slop
	5 19 41	29 41	4 13 15	15880	16 10	...	Zanotti
	5 25 5	40 41	4 24 7	12308	7 14	...	Asclépi
	5 24 42	41 28	4 25 46	11640	7 11	...	Lambert
	5 25 14	40 43	4 24 22	12280	7 18	...	Widder



Year	Ascending Node	Inclin. of Orbit	Perihelion	Perihel. Dist.	Time of Perihelion	Motion	Calculated by
	s o /	o /	s o /		d h		
1769	5 25 3	40 50	4 24 16	12264	Oct. 7 15	Dir.	Euler
	5 25 5	40 50	4 24 11	12269	7 16	...	Lexell
	5 25 6	40 47	4 24 16	12274	7 16	...	Pingré
1770	4 16 39	1 44	11 26 7	62959	Aug. 9 0	Dir.	Pingré
	4 13 39	1 41	11 25 5	65800	10 22	...	Pingré
	4 15 29	1 47	11 26 7	62955	9 0	...	Prosperin
	4 15 4	1 45	11 22 51	64456	8 9	...	Prosperin
	4 14 30	1 23	0 7 14	71717	25 2	...	Prosperin
	4 12 56	1 46	11 29 45	64946	12 21	...	Widder
1770	4 12 0	1 34	11 26 16	67438	Aug. 13 13	Dir.	Lexell
	4 12 17	1 35	11 26 26	67689	14 0	...	Pingré
	4 16 14	1 45	11 26 13	62872	9 1	...	Slop
	4 12 0	1 55	11 25 57	63100	9 4	...	Lambert
	4 14 22	1 49	11 26 19	62758	8 19	...	Rittenhouse
1770	3 18 42	31 26	6 28 23	52824	Nov. 22 6	Ret.	Pingré
1771	0 27 51	11 15	3 13 28	90576	April 18 22	Dir.	Pingré
	0 27 50	11 17	3 13 48	90188	19 1	...	Prosperin
1772	8 12 43	19 0	3 18 6	101814	Feb. 18 21	Dir.	La Lande
1773	4 1 16	61 25	2 15 36	113390	Sept. 5 11	Dir.	Pingré
	4 3 15	62 33	2 21 40	123800	2 12	...	Lambert
	4 3 35	62 36	2 20 43	121550	2 19	...	Schultz
	4 1 12	61 21	2 15 16	113010	5 6	...	Lexell
	4 1 5	61 13	2 14 58	112530	5 9	...	Pingré
1774	6 0 57	82 48	10 16 28	142525	Aug. 14 4	Dir.	De Saron
	6 0 50	82 49	10 16 48	142525	14 18	...	De Saron
	6 1 22	82 21	10 17 26	142600	15 5	...	Boscovich
	6 0 50	83 0	10 17 22	142860	15 11	...	Mechain
1779	0 25 3	32 26	2 27 14	71322	Jan. 4 3	Dir.	De Saron
	0 25 6	32 24	2 27 13	71313	4 2	...	Mechain
	0 25 4	32 26	2 27 14	71319	4 2	...	D'Angos
1780	4 4 0	53 56	8 6 30	9781	Sept. 30 20	Ret.	Lexell
	4 4 30	53 15	8 6 19	10047	30 16	...	Lexell
	4 4 9	53 48	8 6 21	9926	30 18	...	Mechain
1781	2 25 13	5 16	5 22 17	1027558	March 13 0	Dir.	Boscovich
			5 28 13	944040	Jan. 27 6	...	La Place
1781	2 23 1	81 43	7 29 11	77586	July 7 5	Dir.	Mechain
1781	2 17 23	27 13	0 16 3	96101	Nov. 20 13	Ret.	Mechain

M. Facio has suggested, that some of the comets have their nodes so very near the annual orbit of the earth, that if the earth should happen to be found in that part next the node at the time of a comet's passing by; as the apparent motion of the comet will be immensely swift, so its parallax will become very sensible; and its proportion to that of the sun will be given: whence, such transits of comets will afford the best means of determining the distance between the earth and sun.

The comet of 1472, for instance, had a parallax above 20 times greater than the sun's: and if that of 1618 had come down in the beginning of March to its descending node, it would have been much nearer the earth, and its parallax much more notable. But hitherto none has threatened the earth with a nearer appulse than that of 1680: for, Dr. Halley finds, by calculation, that Nov. 11th, at 1 h. 6 min. afternoon, that comet was not more than one semidiameter of the earth to the northward of the earth's path; at which time had the earth been in that part of its orbit, the comet would have had a parallax equal to that of the moon.—

What might have been the consequence of so near an appulse, a contact, or lastly, a shock of these bodies? Mr. Whiston says, a deluge!

*To determine the Place and Course of a Comet.*—Observe the distance of the comet from two fixed stars, whose longitudes and latitudes are known: then from the distances thus known, calculate the place of the comet by spherical trigonometry.

Longomontanus shews an easy method of finding and tracing out the places of a comet mechanically, which is, to find two stars in the same line with the comet, by stretching a thread before the eye over all the three; then do the same by two other stars and the comet: this done, take a celestial globe, or a planisphere, and draw a line upon it first through the former two stars, and then through the latter two; so shall the intersection of the two lines be the place of the comet at that time. If this be repeated from time to time, and all the points of intersection connected, it will shew the path of the comet in the heavens.

COMETARIUM, a machine adapted to give a representation of the revolution of a comet about the sun.



It is so contrived as, by elliptical wheels, to shew the unequal motion of a comet in every part of its orbit. The comet is represented by a small brass ball, carried by a radius vector, or wire, in an elliptical groove about the sun in one of its foci; and the years of its period are shewn by an index moving with an equable motion over a graduated silvered circle. See a particular description, with a figure of it, in Ferguson's Astron. 8vo. pa. 400.

**COMMANDINE** (FREDERICK), a celebrated mathematician and linguist, was born at Urbino in Italy, in 1509; and died in 1575; consequently at 66 years of age. He was famous for his learning and knowledge in the sciences. To a great depth, and just taste in the mathematics, he joined a critical skill in the Greek language; a happy conjunction which made him very well qualified for translating and expounding the writings of the Greek mathematicians. And accordingly, with a most laudable zeal and industry, he translated and published several of their works, to which no former writer had done that good office. On which account, Francis Moria, duke of Urbino, who was very conversant in those sciences, proved a very affectionate patron to him. He is greatly applauded by Bianchanus, and other writers; and he justly deserved their encomiums.

Of his own works Commandine published the following:

1. Commentarius in Planisphærium Ptolomæi: 1558, in 4to.

2. De Centro Gravitatis Solidorum: Bonon. 1565, in 4to.

3. Horologiorum Descriptio: Rom. 1562, in 4to.

He translated and illustrated with notes the following works, most of them beautifully printed, in 4to. by the celebrated printer Aldus:

1. Archimedis Circuli Dimensio; de Lineis Spiraliibus; Quadratura Parabolæ; de Conoidibus & Sphæroidibus; de Arenæ Numero: 1558.

2. Ptolomæi Planisphærium; & Planisphærium Jordani: 1558.

3. Ptolomæi Analemma: 1562.

4. Archimedis de iis quæ vehuntur in aqua: 1565.

5. Apollonii Pergæi Conicorum libri quatuor, una cum Pappi Alexandrini Lemmatibus, & Commentariis Eutocii Ascalonitæ, &c: 1566.

6. Machometes Bagdadinus de Superficierum Divisionibus: 1570.

7. Elementa Euclidis: 1572.

8. Aristarchus de Magnitudinibus & Distantiis Solis & Lunæ: 1572.

9. Heronis Alexandrini Spiritualium liber: 1583.

10. Pappi Alexandrini Collectiones Mathematicæ: 1588.

**COMMANDING Ground**, in Fortification, an eminence, or rising ground, overlooking any post or strong place. This is of three sorts. 1st, *A Front Commanding Ground*, or a height opposite to the face of the post, which plays upon its front. 2dly, *A Reverse Commanding Ground*, or an eminence that can play upon the rear or back of the post. 3dly, *An Enfilade Commanding Ground*, or an eminence in flank which can, with its shot, scour all the length of a straight line.

**COMMENSURABLE Quantities, or Magnitudes**, are such as have some common aliquot part, or which may be measured or divided, without a remainder, by one and the same measure or divisor, called their common measure. Thus, a foot and a yard are commensurable, because there is a third quantity that can measure each, viz an inch; which is 12 times contained in the foot, and 36 times in the yard.—Commensurables are to each other, as one rational whole number is to another; but incommensurables are not so: And therefore the ratio of commensurables is rational; but that of incommensurables, irrational: hence also the exponent of the ratio of commensurables, is a rational number.

**COMMENSURABLE Numbers**, whether integers, or fractions, or surds, are such as have some other number, which will measure or divide them exactly, or without a remainder. Thus, 6 and 8 are commensurable, because 2 measures or divides them both. And  $\frac{2}{3}$  and  $\frac{3}{4}$ , or  $\frac{8}{12}$  and  $\frac{9}{12}$  are commensurable fractions, because the fraction  $\frac{1}{12}$ , or  $\frac{1}{12}$ , &c, will measure them both: and in this sense, all fractions may be said to be commensurable. Also, the surds  $2\sqrt{2}$  and  $3\sqrt{2}$  are commensurable, being measured by  $\sqrt{2}$ , or being to each other as 2 to 3.

**COMMENSURABLE in Power**. Euclid says, right lines are commensurable in power, when their squares are measured by one and the same space or superficies.

**COMMON**, is applied to an angle, line, measure, or the like, that belongs to two or more figures, or other things. As, a common angle, a common side, a common base, a common measure, &c.

**COMMON MEASURE**, or divisor, is that which measures two or more things without a remainder. So of 8 and 12, a common measure is 2, and so is 4.

*The greatest COMMON Measure*, is the greatest number that can measure two other numbers. So, of 8 and 12, the greatest common measure is 4.

*To find the greatest common measure* of two numbers. Divide the greater term by the less; then divide the divisor by the remainder, if there be any; and so on continually, always dividing the last divisor by the last remainder, till nothing remains; and then is the last divisor the greatest common measure sought.

Thus, to find the greatest common measure of 816 and 1488.

$$\begin{array}{r} 816)1488(1 \\ \underline{816} \end{array}$$

$$\begin{array}{r} 672)816(1 \\ \underline{672} \end{array}$$

$$\begin{array}{r} 144)672(4 \\ \underline{576} \end{array}$$

$$\begin{array}{r} 96)144(1 \\ \underline{96} \end{array}$$

$$\begin{array}{r} \text{the common meas. } 48)96(2 \\ \underline{96} \\ 0 \end{array}$$

Therefore



AIR - BALLOONS .

Montgolfier's.  
*Fauxbourg of St. Antoine .*



Montgolfier's .  
*Fauxbourg of St. Germain .*



Charles & Robert's.  
*Champ de Mars .*



Versailles .



Blanchard's .










Fig. 1. AIR GUN.




Fig. 2. The Lock.



Fig. 3. Improved AIR GUN.




Fig. 4. AIR Cane.




Fig. 5. Magazine AIR GUN.




Fig. 6.





Fig. 8.



ASTROLABE,

Fig. 7.









AIR PUMPS.

Fig. 1






Fig. 2



ANEMOMETERS.

Wolfius's

Fig. 3



Lind's.

Fig. 4





Fig. 5

Martin's.









B A R O M E T E R S .

Fig. 1.



Fig. 2.



Fig. 3.




Fig. 5.




Fig. 4.



Fig. 8.



Fig. 7.




Fig. 13.



Fig. 6.




Fig. 11.



Fig. 12.




Fig. 9.





Fig. 10.









BALLISTA.




Fig. 1.

BATTERING RAM.




Fig. 2

CATAPULTS.




Fig. 3

Fig. 4








COMPASS.

Fig. 1



D<sup>r</sup> Knight's COMPASSES.

Fig. 2






Fig. 3




Azimuth COMPASS.

Fig. 4



COMPASS Card.


Fig. 5



Proportional Compasses.

Elliptic Compasses.

Fig. 6





FOUNTAINS.

Fig. 9



Fig. 8









Therefore 48 is the greatest common measure of 816 and 1488, thus :

$$\begin{array}{r} 48 \overline{) 816} (17 \\ \underline{336} \\ 336 \\ \underline{\phantom{000}} \end{array} \qquad \begin{array}{r} 48 \overline{) 1488} (31 \\ \underline{144} \\ 48 \\ \underline{48} \\ \phantom{000} \end{array}$$

The common measure is useful in fractions, to reduce a fraction to its least terms, by dividing those that are given by their greatest common measure. So  $\frac{816}{1488}$  reduces to  $\frac{17}{31}$ , by dividing 816 and 1488 both by their greatest common measure 48.

**COMMUNICATION of Motion**, that act of a moving body, by which it gives motion, or transfers its motion to another body.

Father Mallebranche considers the communication of motion, as something metaphysical; that is, as not necessarily arising from any physical principles, or any properties of bodies, but flowing from the immediate agency of God.

The communication of motion results from, and is an evidence of the impenetrability and inertia of matter, as such; unless we admit the hypothesis of the penetrability of matter, advanced by Boscovich and Michell, and ascribe to the power of repulsion those effects which have been usually ascribed to its solidity and actual resistance.

Newton shews that action and reaction are equal and opposite; so that one body striking or acting against another, and thence causing a change in its motion, does itself undergo the very same change in its own motion, the contrary way. And hence, a moving body striking directly another at rest, it loses just as much of its motion as it communicates to the other. For the laws and quantity of motion so communicated, either in elastic or nonelastic bodies, see **COLLISION**.

**COMMUTATION**, *Angle of*, is the distance between the sun's true place seen from the earth, and the place of a planet reduced to the ecliptic: which therefore is found by taking the difference between the sun's longitude and the heliocentric longitude of the planet.

**COMPANY**, *Rule of*, or *Rule of Fellowship*, in Arithmetic, is a rule by which are determined the true shares of profit or loss, due to the several partners, or associates, in any enterprize, or trade, in due proportion to the stock contributed by each, and the time it was employed. To do which properly, see the *Rule of Fellowship*.

**COMPARTMENT**, a design composed of several different figures, disposed with symmetry; to adorn a parterre, a cieling, pannel of joinery, or the like.

**COMPARTITION**, the useful and graceful distribution of the whole ground-plot of an edifice, into rooms of office, and of reception, or entertainment.

**COMPASS**, or *Mariner's Compass*, is an instrument used at sea by mariners to direct and ascertain the course of their ships. It consists of a circular brass box, which contains a paper card with the 32 points of the compass, or winds, fixed on a magnetic needle that always turns to the north, excepting a small deviation, which is variable at different places, and at the same

place at different times. See **VARIATION of the Compass**.

The needle with the card turns on an upright pin fixed in the centre of the box. To the middle of the needle is fixed a brass conical socket or cap, by which the card hanging on the pin turns freely round the centre.

The top of the box is covered with a glass, to prevent the wind from disturbing the motion of the card. The whole is inclosed in another box of wood, where it is suspended by brass hoops or gimbals, to keep the card in a horizontal position during the motions of the ship. The whole is to be so placed in the ship, that the middle section of the box, parallel to its sides, may be parallel to the middle section of the ship along its keel.

The invention of the compass is usually ascribed to Flavio Gioia, or Flavio of Malphi, about the year 1302; and hence it is that the territory of Principato, the part of the kingdom of Naples where he was born, has a compass for its arms. He divided his compass only into 8 points. Others ascribe the invention to the Chinese; and Gilbert, in his book *de Magnete*, affirms that Marcus Paulus, a Venetian, making a journey to China, brought back the invention with him in 1260. What strengthens this conjecture is, that at first they used the compass, in the same manner as the Chinese still do, viz, letting it float on a small piece of cork, instead of suspending it on a pivot. It is added, that their emperor Chiningus, a celebrated astrologer, had a knowledge of it 1120 years before Christ. The Chinese divide their compass only into 24 points. But Ludi Vertomanus affirms, that when he was in the East-Indies, about the year 1500, he saw a pilot of a ship direct his course by a compass, fastened and framed as those now commonly used. And Barlow, in his book called *The Navigator's Supply*, anno 1597, says, that in a personal conference with two East-Indians, they affirmed, that instead of our compass, they use a magnetical needle of 6 inches, and longer, upon a pin in a dish of white China earth, filled with water; in the bottom of which they have two cross lines for the 4 principal winds, the rest of the divisions being left to the skill of their pilots. Also in the same book he says that the Portuguese, in their first discovery of the East-Indies, got a pilot of Mahinde, who brought them from thence in 33 days, within sight of Calicut.

But Fauchette relates some verses of Guoyot de Provence, who lived in France about the year 1200, which seem to make mention of the compass under the name of *marinette*, or *mariner's stone*; which shew it was used in France near 100 years before either the Mafite or Venetian one. The French even lay claim to the invention, from the fleur de lys with which most people distinguish the north point of the card. With as much reason Dr. Wallis ascribes it to the English, from its name *compass*, by which name most nations call it, and which he observes is used in many parts of England to signify a circle.

The mariner's compass was long very rude and imperfect, but at length received great improvement from the invention and experiments of Dr. Knight, who discovered the useful practice of making artificial magnets; and the farther emendations of Mr. Smeaton, and



Mr. M'Culloch, by which the needles are larger and stronger than formerly, and instead of swinging in gimbals, the compass is supported in its very centre upon a prop, and the centres of motion, gravity, and magnetism are brought almost all to the same point.

After the discovery of that most useful property of the magnet, or loadstone, viz, its giving a polarity to hardened iron or steel, the compass was many years in use before it was known in anywise to deviate from the poles of the world. About the middle of the 16th century, so confident were some persons that the needle invariably pointed due north, that they treated with contempt the notion of the variation, which about that time began to be suspected. However, careful observations soon discovered that in England, and its neighbourhood, the needle pointed to the eastward of the true north line; and the quantity of this deviation being known, mariners became as well satisfied as if the compass had none; because the true course could be obtained by making allowance for the true variation.

From succeeding observations it was afterwards found, that the deviation of the needle from the north was not a constant quantity, but that it gradually diminished, and at last, namely, about the year 1657, it was found that the needle pointed due north at London, and has ever since been going to the westward, till now the variation is upwards of two points of the compass: indeed it was  $22^{\circ} 41'$  about the middle of the year 1781, as appears by the *Philos. Trans.* pa. 225, for that year, and is probably now somewhat more, which it would be of consequence to know; but why such useful observations and experiments, as those of the variation and dip of the magnetic needle, have been so long discontinued, to the prejudice of science, is best known to the learned President of that Society, and his Council. So that in any one place it may be suspected the variation has a kind of libratory motion, traversing through the north, to unknown limits eastward and westward. But the settling of this point must be left to time and future experiments. See *Variation*, also *Inclination*, and *Dip*. Also for a farther description of different compasses and their uses, see that useful book, Robertson's *Naviga-tion*, vol. 2. p. 231.

The *Azimuth Compass* differs from the common sea compass in this; that the circumference of the card or box is divided into degrees; and there is fitted to the box an index with two sights, which are upright pieces of brass, placed diametrically opposite to each other, having a slit down the middle of them, through which the sun or star is to be viewed at the time of observation.

The *Use of the Azimuth Compass*, is to take the bearing of any celestial object, when it is in, or above the horizon, that from the magnetical azimuth or amplitude, the variation of the needle may be known. See *Azimuth*, and *Amplitude*.

The figure of the compass card, with the names of the 32 points or winds, are as in fig. 5, plate vi; where other compasses are also exhibited.

As there are 32 whole points quite around the circle, which contains 360 degrees, therefore each point of the compass contains the 32d part of 360, that is  $11\frac{1}{4}$  degrees, or  $11^{\circ} 15'$ ; consequently the half point is  $5^{\circ} 37' 30''$ , and the quarter point  $2^{\circ} 48' 45''$ .

The points of the compass are otherwise called Rhumbs; and the numbers of degrees, minutes and seconds made by every quarter point with the meridian, are exhibited in the following table.

A TABLE of Rhumbs, shewing the Degrees, Minutes, and Seconds, that every Point and Quarter-point of the Compass makes with the Meridian.

North		Pts. qr.	°	'	"	Pts. qr.	South	
N b E	N b W	0 1	2	48	45	0 1	S b E	S b W
		0 2	5	37	30	0 2		
		0 3	8	26	15	0 3		
		1 0	11	15	0	1 0		
N N E	N N W	1 1	14	3	45	1 1	S S E	S S W
		1 2	16	51	30	1 2		
		1 3	19	41	15	1 3		
		2 0	22	30	0	2 0		
N E b N	N W b N	2 1	25	18	45	2 1	S E b S	S W b S
		2 2	28	7	30	2 2		
		2 3	30	56	15	2 3		
		3 0	33	45	0	3 0		
N E	N W	3 1	36	33	45	3 1	S E	S W
		3 2	39	22	30	3 2		
		3 3	42	11	15	3 3		
		4 0	45	0	0	4 0		
N E b E	N W b W	4 1	47	48	45	4 1	S E b E	S W b W
		4 2	50	37	30	4 2		
		4 3	53	26	15	4 3		
		5 0	56	15	0	5 0		
N N E	W N W	5 1	59	3	45	5 1	E S E	W S W
		5 2	61	52	30	5 2		
		5 3	64	41	15	5 3		
		6 0	67	30	0	6 0		
E b N	W b N	6 1	70	18	45	6 1	E b S	W b S
		6 2	73	7	30	6 2		
		6 3	75	56	15	6 3		
		7 0	78	45	0	7 0		
East	West	7 1	81	33	45	7 1	East	West
		7 2	84	22	30	7 2		
		7 3	87	11	15	7 3		
		8 0	90	0	0	8 0		

COMPASS Dials, are small dials, fitted in boxes, for the pocket, to shew the hour of the day by the direction of the needle that indicates how to place them right, by turning the dial about till the cock or style stand directly over the needle. But these can never be very exact, because of the variation of the needle itself; unless that variation be allowed for, in making and placing the instrument.

COMPASSES, or *Pair of COMPASSES*, a mathematical instrument for describing circles, measuring and dividing lines, or figures, &c.

The common compasses consist of two sharp-pointed branches or legs of iron, steel, brass, or other metal, joined together at the top by a rivet, about which they move as on a centre. Those compasses are of the best sort in which the pin or axle, on which the joint turns, is made of steel, and also half the joint itself, as the opposite metals wear more equally: the points should also be made of hard steel, well polished; and the joint should open and shut with a smooth, easy, and uniform motion. In some compasses, the points are both fixed; but in others, one is made to take out occasionally, and a drawing-pen, or pencil, put in its place.

There are in use compasses of various kinds and contrivances, adapted to the various purposes they are intended for; as,

COMPASSES of *three Legs*, or *Triangular Compasses*; the construction of which is like that of the common compasses, with the addition of a third leg or point, which



which has a motion every way. Their use is to take three points at once, and so to form triangles, and lay down three positions of a map to be copied at once.

*Beam COMPASSES* consist of a long straight beam or bar, carrying two brass cursors; one of these being fixed at one end, the other sliding along the beam, with a screw to fasten it on occasionally. To the cursors may be screwed points of any kind, whether steel, pencils, or the like. To the fixed cursor is sometimes applied an adjusting or micrometer screw, by which an extent is obtained to very great nicety. The beam compasses are used to draw large circles, to take great extents, or the like.

*Bow COMPASSES*, or *Bows*, are a small sort of compasses, that shut up in a hoop, which serves for a handle. Their use is to describe arcs or circumferences with a very small radius.

*Caliber COMPASSES*. See *CALIBER*.

*Clockmakers COMPASSES* are jointed like the common compasses, with a quadrant or bow, like the spring compasses; only of different use, serving here to keep the instrument firm at any opening. They are made very strong, with the points of their legs of well-tempered steel, as being used to draw or cut lines in pasteboard, or copper, &c.

*Cylindrical and Spherical COMPASSES*, consist of four branches, joined in a centre, two of them being circular and two flat, a little bent at the ends. The use of them is to take the diameter, thickness, or caliber of round or cylindrical bodies; as cannons, balls, pipes, &c.

There are also spherical compasses, differing in nothing from the common ones, but that their legs are arched; serving to take the diameters of round bodies.

There is also another sort of compasses lately invented, for measuring the diameter of round bodies, as balls, &c, which consist of two flat pieces of metal set at right angles on a straight bar or beam of the same; the one piece being fixed, and the other sliding along it, so far as just to receive the round body between them; and then its diameter, or distance between the two pieces, is shewn by the divisions marked on the beam.

*Elliptical COMPASSES*, are used to draw ellipses or ovals of any kind. The instrument consists of a beam AB (Plate vi. fig. 6.) about a foot long, bearing three cursors; to one of which may be screwed points of any kind; and to the bottom of the other two are rivetted two sliding dove-tails, adjusted in grooves made in the cross branches of the beam. The dove-tails having a motion every way, by turning about the long branch, they go backward and forward along the cross; so that when the beam has gone half way round, one of these will have moved the whole length of one of the branches; and when the beam has gone quite round, the same dove-tail has gone back the whole length of the branch. Understand the same of the other dove-tail.

Note, the distance between the two sliding dove-tails, is the distance between the two foci of the ellipse; so that by changing that distance, the ellipse will be rounder or flatter. Under the ends of the branches of the cross, are placed four steel points to keep it fast.

The use of this compass is easy: by turning round

the long branch, the pen, pencil, or other points will draw the ellipse required.

Its figure shews both its use and construction.

*German COMPASSES* have their legs a little bent outwards, near the top; so that when shut, the points only meet.

*Hair COMPASSES* are so contrived within side by a small adjusting screw to one of the legs, as to take an extent to a hair's breadth, or great exactness.

*Proportional COMPASSES* are those whose joint lies, not at the end of the legs, but between the points terminating each leg. These are either simple, or compound. In the former sort the centre, or place of the joint is fixed; so that one pair of these serves only for one proportion.

*Compound Proportional COMPASSES* have the joint or centre moveable. They consist of two parts or sides of brass, which lie upon each other so nicely as to seem but one when they are shut. These sides easily open, and move about the centre, which is itself moveable in a hollow canal cut through the greatest part of their length. To this centre on each side is fixed a sliding piece, of a small length, with a fine line drawn on it serving as an index, to be set against other lines or divisions placed upon the compasses on both sides. These lines are, 1, A line of lines; 2, a line of superficies, areas, or planes, the numbers on which answer to the squares of those on the line of lines; 3, a line of solids, the numbers on which answer to the cubes of those on the line of lines; 4, a line of circles, or rather of polygons to be inscribed in circles. These lines are all unequally divided, the first three from 1 to 20, and the last from 6 to 20. The use of the first is to divide a line into any number of equal parts; by the 2d and 3d are found the sides of like planes or solids in any given proportion; and by the 4th, circles are divided into any number of equal parts, or any polygons inscribed in them. See Plate vi. fig. 7.

*Spring COMPASSES*, or *Dividers*, are made of hardened steel, with an arched head, which by its spring opens the legs; the opening being directed by a circular screw fastened to one of the legs, let through the other, and worked with a nut.

*Trisecting COMPASSES*, for the trisecting of angles geometrically, for which purpose they were invented by M. Tarragon.

The instrument consists of two central rules, and an arch of a circle of 120 degrees, immoveable, with its radius: the radius is fastened with one of the central rules, like the two legs of a sector, that the central rule may be carried through all the points of the circumference of the arch. The radius and rule should be as thin as possible; and the rule fastened to the radius should be hammered cold, to be more elastic; and the breadth of the other central rule must be triple the breadth of the radius: in this rule also is a groove, with a dove-tail fastened on it, for its motion; there must also be a hole in the centre of each rule.

*Turn-up COMPASSES*, a late contrivance to save the trouble of changing the points: the body is like the common compasses; and towards the bottom of the legs without side, are added two other points, besides the usual ones; the one carrying a drawing pen point, and the



the other a port-crayon; both being adjusted to turn up, to be used or not, as occasion may require.

COMPLEMENT in general, is what is wanting, or necessary, to complete some certain quantity or thing. As, the

COMPLEMENT of an arch or angle, as of  $90^\circ$  or a quadrant, is what any given arch or angle wants of it; so the complement of  $50^\circ$  is  $40^\circ$ , and the complement of  $100^\circ$  degrees is  $-10^\circ$ , a negative quantity.—The complement to  $180^\circ$  is usually called the supplement, to distinguish it from the complement to  $90^\circ$ , properly so called.—The sine of the complement of an arc, is contracted into the word cosine; the tangent of the complement, into cotangent; &c.

Arithmetical COMPLEMENT, is what a number or logarithm wants of unity or 1 with some number of ciphers. It is best found, by beginning at the left-hand side, and subtracting every figure from 9, except the last, or right-hand figure, which must be subtracted from 10. So, the arithmetical comp. of the log. 9.5329714, by subtracting from 9's, &c, is 0.4670286.

The arithmetical complements are much used in operations by logarithms, to change subtractions into additions, which are more conveniently performed, especially when there are more than one of them in the operation.


COMPLEMENT, in Astronomy, is used for the distance of a star from the zenith; or the arc contained between the zenith and the place of a star which is above the horizon. It is the same as the complement of the altitude, or co-altitude, or the zenith distance.

COMPLEMENT of the Course, in Navigation, is the quantity which the course wants of  $90^\circ$ , or 8 points, viz, a quarter of the compass.

COMPLEMENT of the Curtain, in Fortification, is that part of the anterior side of the curtain, which makes the demigorge.

COMPLEMENT of the Line of Defence, is the remainder of that line, after the angle of the flank is taken away.

COMPLEMENTS of a Parallelogram, or in a Parallelogram, are the two lesser parallelograms, made by drawing two right lines parallel to each side of the given parallelogram, through the same point in the diagonal. So P and Q are the complements in the parallelogram ABCD.



In every case, these complements are always equal, viz, the parallelogram  $P = Q$ .

COMPLEMENT of Life, a term much used, in the doctrine of Life Annuities, by De Moivre, and, according to him, it denotes the number of years which a given life wants of 86, this being the age which he considered as the utmost probable extent of life. So 56 is the complement of 30, and 30 is the complement of 56.

That author supposed an equal annual decrement of life through all its stages, till the age of 86. Thus, if there be 56 persons living at 30 years of age, it is supposed that one will die every year, till they be all dead in 56 years. This hypothesis in many cases is very near the truth; and it agrees so nearly with Halley's

table, formed from his observations of the mortuary bills of Breslaw, that the value of lives deduced either from the hypothesis, or the table, need not be distinguished; hence it very much eases the labour of calculating them. See Life ANNUITIES, also De Moivre's Treatise on Annuities, pa. 83, and Price on Reverfionary Payments, pa. 2.

COMPOSITE Number, is one that is compounded of, or made up by the multiplication of two other numbers, greater than 1, or which can be measured by some other number greater than 1. As 12, which is composed, or compounded of 2 and 6, or 3 and 4, viz by multiplying together 2 and 6, or 3 and 4, both products making the same number 12; which therefore is a composite number.

Composites are opposed to prime numbers, or primes, which cannot be exactly measured by any other number, and cannot be produced by multiplying together two other factors.

Composite Numbers between themselves, are the same with commensurable numbers, or such as have a common measure or factor; as 15 and 12, which have the common term 3.

The doctrine of Prime and Composite numbers is pretty fully treated in the 7th and 8th books of Euclid's Elements.

COMPOSITE Order, in Architecture, is the last of the five orders of columns; and is so called because its capital is composed out of those of the other orders. Thus, it borrows a quarter-round from the Tuscan and Doric; a double row of leaves from the Corinthian; and volutes from the Ionic. Its cornice has single modillions, or dentils; and its column is 10 diameters in height.

This order is also called the Roman order, and Italic order, as having been invented by the Romans, like as the other orders are denominated from the people among whom they had their rise.

COMPOSITION, is a species of reasoning by which we proceed from things that are known and given, step by step, till we arrive at such as were before unknown or required; viz, proceeding upon principles self-evident, on definitions, postulates, and axioms, with a previously demonstrated series of propositions, step by step, till it gives a clear knowledge of the thing to be known or demonstrated. Composition, otherwise called the synthetical method, is opposed to Resolution, or the analytical method, and is chiefly used by the ancients, Euclid, Apollonius, &c. See Pappus; also the term Analysis.


COMPOSITION of forces, or of motion, is the union or assemblage of several forces or motions that are oblique to one another, into an equivalent one in another direction.

1. When several forces or motions are united, that act in the same line of direction, the combined force or motion will be in the same line of direction still. But when oblique forces are united, the compounded force takes a new direction, different from both, and is either a right line or a curve, according to the nature of the forces compounded.

2. If two compounding motions be both equable ones, whether equal to each other or not, the line of the compound




compound motion will be a straight line. Thus, if the one equable in the direction AB be sufficient to carry a body over the space AB in any time, and the other motion sufficient to pass over AC in the same time; then by the compound motion, or both acting on the body together, it would in the same time pass over the diagonal AD of the parallelogram ABDC. For because the motions are uniform, any spaces  $Ab$ ,  $Ac$  passed over in the same time, are proportional to the velocities, or to AB and AC; and consequently all the points A,  $d$ ,  $d$ , D, of the path are in the same right line.




3. And though the compounding motions be not equable, but variable, either accelerated or retarded, provided they do but vary in a similar manner, the compounded motion will still be in a straight line. Thus, suppose, for instance, that the motions both vary in such a manner, as that the spaces passed over in the same time, whether they be equal to each other or not, are both as the same power  $n$  of the time; then  $AB^n : AC^n :: Ab^n : Ac^n$ , and hence  $AB : AC :: Ab : Ac$ , and therefore, as before,  $Ad$  and  $D$  is still a right line.

4. But if the compounding motions be not similar to each other, as when the one is equable and the other variable, or when they are varied in a dissimilar manner; then the compounded motion is in some curve line. So if the motion in the one direction EF be in a less proportion, with respect to the time, than that in the direction EG is, then the path will be a curve line EIH concave to EF; but if the motion in EF be in a greater proportion than that in EG, then the path of the compound motion will be a curve E $b$ H convex towards EF: that is, in general, the curvilinear path is convex towards that direction in which the motion is in the less proportion to the time. Hence, for a particular instance, if the motion in the direction EF be a motion of projection, which is an equable motion, and the motion in the direction EG that arising from gravity, which is a uniformly accelerated motion, or in proportion to the squares of the times; then is EG as  $GH^2$ , and Eg as  $gb^2$ , that is  $EG : Eg :: GH^2 : gb^2$ , which is the property of the parabola; and therefore the path E $b$ H of any body projected, is the common parabola.



5. If there be three forces united, or acting against the same point A at the same time, viz, the force or weight B in the direction AB, and the forces or tensions in the directions AC, AD; and if these three forces mutually balance each other, so as to keep the common point A in equilibrio; then are these forces directly proportional to the respective sides of a triangle formed by drawing lines parallel to the directions of these forces; or indeed perpendicular to those directions, or making any one and the same angle with them. So that, if BE be drawn parallel, for instance, to AD, and meet CA produced in E, forming the triangle

ABE; then are the three forces in the directions AB, AC, AD, respectively proportional to the sides AB, AE, BE.



And this theorem, with its corollaries, Dr. Kēil observes, is the foundation of all the new mechanics of M. Varignon: by help of which may the force of the muscles be computed, and most of the mechanic theorems in Borelli, *De Motu Animalium*, may immediately be deduced.

See more of the Composition of Forces under the article COLLISION.

COMPOSITION of Numbers and Quantities. See COMBINATION.

COMPOSITION of Proportion, according to the 15th definition of the 5th book of Euclid's Elements, is when, of four proportionals, the sum of the 1st and 2d is to the 2d, as the sum of the 3d and 4th is to the 4th:

as if it be  $a : b :: c : d$ ,

then by composition  $a + b : b :: c + d : d$ .

Or, in numbers, if  $2 : 4 :: 9 : 18$ ,

then by composition  $6 : 4 :: 27 : 18$ .

COMPOSITION of Ratios, is the adding of ratios together: which is performed by multiplying together their corresponding terms, viz, the antecedents together, and the consequents together, for the antecedent and consequent of the compounded ratio; like as the addition of logarithms is the same thing as the multiplication of their corresponding numbers. Or, if the terms of the ratios be placed fraction-wise, then the addition or composition of the ratios, is performed by multiplying the fractions together.

Thus, the ratio of  $a : b$ , or of  $2 : 4$ ,

added to the ratio of  $c : d$ , or of  $6 : 8$ ,

makes the ratio of  $ac : bd$ , or of  $12 : 32$ ; and so the ratio of  $ac$  to  $bd$  is said to be compounded of the ratios of  $a$  to  $b$ , and  $c$  to  $d$ . So likewise, if it were required to compound together the three ratios, viz, of  $a$

to  $b$ ,  $c$  to  $d$ , and  $e$  to  $f$ ; then  $\frac{a}{b} \times \frac{c}{d} \times \frac{e}{f} = \frac{ace}{bdf}$  are the

terms of the compound ratio; or the ratio of  $ace$  to  $bdf$  is compounded, or made up of the ratios of  $a$  to  $b$ ,  $c$  to  $d$ , and  $e$  to  $f$ .

Hence, if the consequent of each ratio be the same as the antecedent of the preceding ratio, then is the ratio of the first term to the last, compounded, or made up of all the other ratios, viz, the ratio of  $a$  to  $e$ , equal to the sum of all the ratios of  $a$  to  $b$ , of  $b$  to  $c$ , of  $c$  to  $d$ , and of  $d$  to  $e$ ; for  $\frac{a}{b} \times \frac{b}{c} \times \frac{c}{d} \times \frac{d}{e} = \frac{a}{e}$  the terms or exponents of the compounded ratio.

Hence



Hence also, in a series of continual proportionals, the ratio of the first term to the third is double of the ratio of the first to the second, and the ratio of the 1st to the 4th is triple of it, and the ratio of the 1st to the 5th is quadruple of it, and so on; that is, the exponents are double, triple, quadruple, &c, of the first exponent: as in the series  $1, a, a^2, a^3, a^4, \&c$ ; where the ratio 1 to  $a^2$  is double, of 1 to  $a^3$  triple, &c, of the ratio of 1 to  $a$ ; or the exponent of  $a^2, a^3, a^4, \&c$ , double, triple, quadruple, &c, of  $a$ .

COMPOUND INTEREST, called also *Interest upon Interest*, is that which is reckoned not only upon the principal, but upon the interest itself forborn, which thus becomes a sort of secondary principal.

If  $r$  be the amount of 1 pound for 1 year, that is the sum of the principal and interest together for one year; then is  $r^2$  the amount for 2 years, and  $r^3$  the amount for 3 years, and in general  $r^t$  the amount for  $t$  years; that is  $r^t$  is the sum or total amount of all the principals and interests together of 1 l. for the whole time or number of years  $t$ ; consequently, if  $p$  be any other principal sum, forborn for  $t$  years, then its amount in that time at compound interest, is  $a = pr^t$ .

The Rule therefore in words is this, to one pound add its interest for one year, or half year, or for the first time at which the interest is reckoned; raise the sum  $r$  to the power denoted by the time or number of terms; then this power multiplied by the principal, or first sum lent, will produce the whole amount.

For example, To find how much 50l. will amount to in 5 years at 5 per cent. per annum, compound interest.—Here the interest of 1 l. for 1 year is .05, and therefore  $r = 1.05$ ; hence the 5th power of it for 5 years, is  $r^5 = 1.27628 \&c$ ; multiply this by  $p$  or 50

gives the amount  $pr^t$  or 63.8141.

or 63 l. 16s. 3 $\frac{1}{4}$ d. for the amount sought.

But Compound Interest is best computed by means of such a table as the following, being the amounts of 1 pound for any number of years, and at several rates of compound interest.

As an example of the use of this table, suppose it were required to find the amount of 250l. for 35 years at 4 per cent. compound interest.

In the column of 4 per cent, and line of 35 years, is 3.94609,

which multiplied by the principal 250 gives 986.52250 or 986 l 10s 5 $\frac{1}{4}$ d, which is the amount sought.

Note, By a bare inspection of this table, it appears how many years are required for any sum of money to double itself, at any rate of compound interest; viz, by looking in the columns when the amount becomes the number 2. So it is found that at the several rates the respective times requisite for doubling any sum, are nearly thus: viz,

Rate 3 3 $\frac{1}{2}$  4 4 $\frac{1}{2}$  5 6  
Years 23 $\frac{1}{2}$  20 $\frac{3}{4}$  17 $\frac{3}{4}$  15 $\frac{3}{4}$  14 $\frac{1}{4}$  12

TABLE of the Amount of 1l. at Compound Interest for many Years and several Rates of Interest.

Yrs.	at 3 per cent	at 3 $\frac{1}{2}$ per cent	at 4 per cent	at 4 $\frac{1}{2}$ per cent	at 5 per cent	at 6 per cent
1	1.03000	1.03500	1.04000	1.04500	1.05000	1.06000
2	1.06090	1.07123	1.08160	1.09203	1.10250	1.12360
3	1.09273	1.10872	1.12486	1.14117	1.15763	1.19102
4	1.12551	1.14752	1.16986	1.19252	1.21551	1.26248
5	1.15927	1.18769	1.21665	1.24618	1.27628	1.33823
6	1.19405	1.22926	1.26532	1.30226	1.34010	1.41852
7	1.22987	1.27228	1.31593	1.36086	1.40710	1.50363
8	1.26677	1.31681	1.36857	1.42210	1.47746	1.59385
9	1.30477	1.36290	1.42331	1.48610	1.55133	1.68948
10	1.34392	1.41060	1.48024	1.55297	1.62890	1.79085
11	1.38423	1.45997	1.53945	1.62285	1.71034	1.89830
12	1.42576	1.51107	1.60103	1.69588	1.79586	2.01220
13	1.46853	1.56396	1.66507	1.77220	1.88565	2.13293
14	1.51259	1.61869	1.73168	1.85194	1.97993	2.26090
15	1.55797	1.67535	1.80094	1.93528	2.07893	2.39656
16	1.60471	1.73399	1.87298	2.02237	2.18287	2.54035
17	1.65285	1.79468	1.94790	2.11338	2.29202	2.69277
18	1.70243	1.85749	2.02582	2.20848	2.40662	2.85434
19	1.75351	1.92250	2.10685	2.30786	2.52695	3.02560
20	1.80611	1.98979	2.19112	2.41171	2.65330	3.20714
21	1.86029	2.05943	2.27877	2.52024	2.78596	3.39956
22	1.91610	2.13151	2.36992	2.63365	2.92526	3.60354
23	1.97359	2.20611	2.46472	2.75217	3.07152	3.81975
24	2.03279	2.28333	2.56330	2.87601	3.22510	4.04893
25	2.09378	2.36324	2.66584	3.00543	3.38635	4.29187
26	2.15659	2.44596	2.77247	3.14068	3.55567	4.54938
27	2.22129	2.53157	2.88337	3.28201	3.73346	4.82235
28	2.28793	2.62017	2.99870	3.42970	3.92013	5.11169
29	2.35657	2.71188	3.11865	3.58404	4.11614	5.41839
30	2.42726	2.80679	3.24340	3.74532	4.32194	5.74349
31	2.50008	2.90503	3.37313	3.91386	4.53804	6.08810
32	2.57508	3.00671	3.50806	4.08998	4.76494	6.45339
33	2.65234	3.11194	3.64838	4.27403	5.00319	6.84059
34	2.73191	3.22086	3.79432	4.46636	5.25335	7.25103
35	2.81386	3.33359	3.94609	4.66735	5.51602	7.68609
36	2.89828	3.45027	4.10393	4.87738	5.79182	8.14725
37	2.98523	3.57103	4.26809	5.09686	6.08141	8.63609
38	3.07478	3.69601	4.43881	5.32622	6.38548	9.15425
39	3.16703	3.82537	4.61637	5.56590	6.70475	9.70351
40	3.26204	3.95926	4.80102	5.81636	7.03999	10.28572
41	3.35990	4.09783	4.99306	6.07810	7.39199	10.90286
42	3.46070	4.24126	5.19278	6.35162	7.76159	11.55703
43	3.56452	4.38970	5.40050	6.63744	8.14967	12.25045
44	3.67145	4.54334	5.61652	6.93612	8.55715	12.98548
45	3.78160	4.70236	5.84118	7.24825	8.98501	13.76461
46	3.89504	4.86694	6.07482	7.57442	9.43426	14.59049
47	4.01190	5.03728	6.31782	7.91527	9.90597	15.46592
48	4.13225	5.21359	6.57053	8.27146	10.40127	16.39387
49	4.25622	5.39606	6.83335	8.64367	10.92133	17.37750
50	4.38391	5.58493	7.10668	9.03264	11.46740	18.42015

COMPOUND Motion, that motion which is the effect of several conspiring powers or forces, viz, such forces as are not directly opposite to each other: as when the radius of a circle is considered as revolving about a centre, and at the same time a point as moving straight along it; which produces a kind of a spiral for the path of the point. And hence it is easily perceived, that all curvilinear motion is compound, or the effect of two or more forces; although every compound motion is not curvilinear.

It is a popular theorem in Mechanics, that in uniform compound motions, the velocity produced by the conspiring powers or forces, is to that of either of the two compounding powers separately, as the diagonal of



of a parallelogram, according to the direction of whose sides they act separately, is to either of the sides. See *COMPOSITION of Motion*, and *COLLISION*.

*COMPOUND Numbers*, those composed of the multiplication of two or more numbers; as 12, composed of 3 times 4. See *COMPOSITE*.

*COMPOUND Pendulum*, that which consists of several weights constantly keeping the same distance, both from each other, and from the centre about which they oscillate.

*COMPOUND Quantities*, are such as are connected together by the signs + or -. Thus,  $a + b$ , or  $a - c + d$ , or  $aa - za$ , are compound quantities.

Compound quantities are distinguished into binomials, trinomials, quadrinomials, &c, according to the number of terms in them; viz, the binomial having two terms; the trinomial, three; the quadrinomial, 4; &c. Also, those that have more than two terms, are called by the general name of multinomials, as also polynomials.

*COMPOUND Ratio*, is that which is made by adding two or more ratios together; viz, by multiplying all their antecedents together for the antecedent, and all the consequents together for the consequent of the compound ratio. So 6 to 72 is a ratio compounded of the ratios of 2 to 6, and 3 to 12; because  $\frac{6}{72} = \frac{2}{6} \times \frac{3}{12}$ : also  $ab$  to  $cd$  is a ratio compounded of the ratio of  $a$  to  $c$ , and  $b$  to  $d$ ; for  $\frac{ab}{cd} = \frac{a}{c} \times \frac{b}{d}$ . See *COMPOSITION of Ratios*.

*COMPRESSION*, the act of pressing, or squeezing something, so as to bring its parts nearer together, and make it occupy less space.

Compression differs from condensation as the cause from the effect, compression being the action of any force on a body, without regarding its effects; whereas condensation denotes the state of a body that is actually reduced into a less bulk, and is an effect of compression, though it may be effected also by other means. Nevertheless, compression and condensation are often confounded.

Pumps, which the ancients imagined to act by suction, do in reality act by compression; the piston, in working in the narrow pipe, compresses the inclosed air, so as to enable it, by the force of its increased elasticity, to raise the valve, and make its escape; upon which, the balance being destroyed, the pressure of the atmosphere on the stagnant surface, forces up the water in the pipe, thus evacuated of its air.

It was long thought that water was not compressible into less bulk: and it was believed, till lately, that after the air had been purged out of it, no art or violence was able to press it into less space. In an experiment made by the Academy del Cimento, water, when violently squeezed, made its way through the fine pores of a globe of gold, rather than yield to the compression.

But the ingenious Mr. Canton, attentively considering this experiment, found that it was not sufficiently accurate to justify the conclusion which had always been drawn from it; since the Florentine philosophers had no method of determining that the alteration of figure in their globe of gold, occasioned

such a diminution of its internal capacity, as was exactly equal to the quantity of water forced into its pores. To bring this matter therefore to a more accurate and decisive trial, he procured a small glass tube of about two feet long, with a ball at one end, of an inch and a quarter in diameter. Having filled the ball and part of the tube with mercury, and brought it exactly to the heat of 50° of Fahrenheit's thermometer, he marked the place where the mercury stood in the tube, which was about six inches and a half above the ball; he then raised the mercury by heat to the top of the tube, and there sealed the tube hermetically; then upon reducing the mercury to the same degree of heat as before, it stood in the tube  $\frac{3}{8}$  of an inch higher than the mark. The same experiment was repeated with water exhausted of air, instead of mercury, and the water stood in the tube  $\frac{4}{8}$  of an inch above the mark. Since the weight of the atmosphere on the outside of the ball, without any counterbalance from within, will compress the ball, and equally raise both the mercury and water, it appears that the water expands  $\frac{1}{8}$  of an inch more than the mercury by removing the weight of the atmosphere. Having thus determined that water is really compressible, he proceeded to estimate the degree of compression corresponding to any given weight. For this purpose he prepared another ball, with a tube joined to it; and finding that the mercury in  $\frac{2}{3}$  of an inch of the tube was the hundred thousandth part of that contained in the ball, he divided the tube accordingly. He then filled the ball and part of the tube with water exhausted of air; and leaving the tube open, placed this apparatus under the receiver of an air-pump, and observed the degree of expansion of the water answering to any degree of rarefaction of the air: and again by putting it into the glass receiver of a condensing engine, he noted the degree of compression of the water corresponding to any degree of condensation of the air. He thus found, by repeated trials, that, in a temperature of 50°, and when the mercury has been at its mean height in the barometer, the water expands one part in 21740; and is as much compressed by the weight of an additional atmosphere; or the compression of water by twice the weight of the atmosphere, is one part in 10870 of its whole bulk. Should it be objected, that the compressibility of the water was owing to any air which it might be supposed to contain, he answers, that more air would make it more compressible; he therefore let into the ball a bubble of air, and found that the water was not more compressed by the same weight than before.

In some farther experiments of the same kind, Mr. Canton found that water is more compressible in winter than in summer; but he observed the contrary in spirit of wine, and oil of olives.

The following table was formed, when the barometer was at 29 inches and a half, and the thermometer at 50 degrees.

Compression of	Millionth parts.	Spec. grav.
Spirit of wine	- 66 -	846
Oil of olives	- 48 -	918
Rain water	- 46 -	1000
Sea water	- 40 -	1028
Mercury	- 3 -	13595



He infers that these fluids are not only compressible, but elastic; and that the compressions of them, by the same weight, are not in the inverse ratio of their densities, or specific gravities, as might be supposed. *Phil. Transf.* vol. lii. 1762. art. 103. and vol. liv. 1764. art. 47.

The compression of the air, by its own weight, is surprisingly great: but the air may be still further compressed by art. See *Elasticity of Air*.

This immense compression and dilatation, Newton observes, cannot be accounted for in any other way, but by a repelling force, with which the particles of air are endued; by virtue of which, when at liberty, they mutually fly each other.

This repelling power, he adds, is stronger and more sensible in air, than in other bodies; because air is generated out of very fixed bodies, but not without great difficulty, and by the help of fermentation: now those particles always recede from each other with the greatest violence, and are compressed with the greatest difficulty, which, when contiguous, cohere the most strongly. See *AIR, ATTRACTION, COHESION, DILATATION, and REPULSION*.

COMPUTATION, the manner of accounting and estimating time, weights, measures, money, &c. See *CALCULATION*, which it is also used for.

CONCAVE, an appellation used in speaking of the inner surface of hollow bodies, more especially of spherical or circular ones.

CONCAVE glasses, lenses, and mirrors, have either one side or both sides concave.

The property of all concave lenses is, that the rays of light, in passing through them, are deflected, or made to recede from one another; as in convex lenses they are inflected towards each other; and that the more as the concavity or convexity has a smaller radius. Hence parallel rays, as those of the sun, by passing through a concave lens, become diverging; diverging rays are made to diverge more; and converging rays are made either to converge less, or to become parallel, or go out diverging. And hence it is, that objects viewed through concave lenses, appear diminished; and the more so, as they are portions of less spheres. See *LENS*.

Concave mirrors have the contrary effect to lenses: they reflect the rays which fall on them, so as to make them approach more to, or recede from each other, than before, according to the situation of the object; and that the more as the concavity is greater, or as the radius of concavity is less. Hence it is that concave mirrors magnifying objects that are presented to them; and that in a greater proportion, as they are portions of greater spheres. And hence also concave mirrors have the effect of burning glasses. See *MIRROR, and BURNING GLASS*.

CONCAVITY, that side of a figure or body which is hollow.

An arch of a curve has its concavity turned all one way, when the right lines that join any two of its points are all on the same side of the arch.

Archimedes, intending to include in his definition such lines as have rectilinear parts, says, a line has its concavity turned one way, when the right lines that join any two of its points, are either all upon one side

of it, or while some fall upon the line itself, none fall upon the opposite side. *Archim. de Sphær. et Cyl. Def. 2,* and *Maclaurin's Fluxions*, art. 180.

When two lines that have their concavity turned the same way, have the same extremes, and the one includes the other, or has its concavity towards it, the perimeter of that which includes, is greater than that which is included. *Archim. ib. ax. 2.*


CONCENTRIC, having the same centre. It is opposed to excentric, or having different centres.

The word is chiefly used in speaking of round bodies and figures, such as circular, and elliptic ones; but it may likewise be used for polygons that are drawn parallel to each other, from the same centre.

Nonnius's method of graduating instruments consists in describing with the same quadrant 45 concentric arches, dividing the outermost into 90 equal parts, the next into 89, and so on.

CONCHOID, or CONCHILES, the name of a curve invented by Nicomedes. It was much used by the ancients in the construction of solid problems, as appears by what Pappus says.

It is thus constructed: AP and BD being two lines intersecting at right angles; from P draw a number of other lines PFDE, &c, on which take always  $DE = DF = AB$  or  $BC$ ; so shall the curve line drawn through all the points E, E, E, be the first conchoid, or that of Nicomedes; and the curve drawn through all the other points F, F, F, is called the second conchoid; though in reality, they are both but parts of the same curve, having the same pole P, and four infinite legs, to which the line DBD is a common asymptote.



The inventor, Nicomedes, contrived an instrument for describing his conchoid by a mechanical motion: thus, in the rule AD is a channel or groove cut, so that a smooth nail, firmly fixed in the moveable rule CB, in the point F, may slide freely within it: into the rule EG is fixed another nail at K, for the moveable rule CB to slide upon. If therefore the rule BC be so moved, as that the nail F passes along the canal AD; the style, or point in C, will describe the first conchoid.

To determine the equation of the curve: put  $AB = BC = DE = DF = a$ ,  $PB = b$ ,  $BG = EH = x$ , and  $GE = BH = y$ ; then the equation to the first conchoid



choid will be  $x^2 \times \overline{b+x}^2 + x^2 y^2 = a^2 \times \overline{b+x}^2$ , or  $x^4 + 2bx^3 + b^2 x^2 + x^2 y^2 = a^2 b^2 + 2a^2 bx + a^2 x^2$ ; and, changing only the sign of  $x$ , as being negative in the other curve, the equation to the 2d conchoid will be  $x^2 \times \overline{b-x}^2 + x^2 y^2 = a^2 \times \overline{b-x}^2$ , or  $x^4 - 2bx^3 + b^2 x^2 + x^2 y^2 = a^2 b^2 - 2a^2 bx + a^2 x^2$ .

Of the whole conchoid, expressed by these two equations, or rather one equation only, with different signs, there are three cases or species; as first, when BC is less than BP, the conchoid will be as in fig. 1; when BC is equal to BP, the conchoid will be as in fig. 2; and when BC is greater than BP, the conchoid will be as in fig. 3.

Newton approves of the use of the conchoid for trisecting angles, or finding two mean proportionals, or for constructing other solid problems. Thus, in the Linear Construction of equations, towards the end of his Universal Arithmetic, he says, "The ancients at first endeavoured in vain at the trisection of an angle, and the finding of two mean proportionals by a right line and a circle. Afterwards they began to consider several other lines, as the conchoid, the cissoid, and the conic sections, and by some of these to solve these problems." Again, "Either therefore the trochoid is not to be admitted at all into geometry, or else, in the construction of problems, it is to be preferred to all lines of a more difficult description: and there is the same reason for other curves; for which reason we approve of the trisections of an angle by a conchoid, which Archimedes in his Lemmas, and Pappus in his Collections, have preferred to the inventions of all others in this case; because we ought either to exclude all lines, besides the circle and right line, out of geometry, or admit them according to the simplicity of their descriptions, in which case the conchoid yields to none, except the circle." Lastly, "That is arithmetically more simple which is determined by the more simple equations, but that is geometrically more simple which is determined by the more simple drawing of lines; and in geometry, that ought to be reckoned best which is geometrically most simple: wherefore I ought not to be blamed, if, with that prince of mathematicians, Archimedes, and other ancients, I make use of the conchoid for the construction of solid problems."

**CONCRETE Numbers** are those that are applied to express or denote any particular subject; as 3 men, 2 pounds, &c. Whereas, if nothing be connected with a number, it is taken abstractedly or universally: thus, 4 signifies only an aggregate of 4 units, without any regard to a particular subject, whether men or pounds, or any thing else.

**CONCURRING, or CONGRUENT Figures**, in Geometry, are such as, being laid upon one another, do exactly correspond to, and cover one another, and consequently must be equal among themselves. Thus, triangles having two sides and the contained angle equal, each to each, are equal to each other in all respects.

**CONDENSATION**, is the compressing or reducing of a body into a less bulk or space; by which means it is rendered more dense and compact.

Wolffius, and some other writers, restrain the use of the word *condensation* to the action of cold: that which is done by external application, they call compression.

Condensation however, in general, consists in bring-

ing the parts closer to each other, and increasing their contact, whatever be the means by which it is effected: in opposition to rarefaction, which renders the body lighter and looser, by setting the parts farther asunder, and diminishing their contact, and of consequence their cohesion.

Air easily condenses, either by cold, or by pressure, but much more by the latter; but most of all by chemical process. Water condenses also both by cold and by pressure; but it suddenly expands by congelation: indeed almost all matter, both solids and fluids, has the same property of condensation by those means. See **COMPRESSION**. So also vapour is condensed, or converted into water, by distillation, or naturally in the clouds. The way in which vapour commonly condenses, is by the application of some cold substance. On touching it, the vapour parts with its heat which it had before absorbed: and on doing so, it immediately loses the proper characteristics of vapour, and becomes water. But though this be the most common and usual way in which we observe vapour to be condensed, nature certainly proceeds after another manner; since we often observe the vapours most plentifully condensed when the weather is really warmer than at other times.

**CONDENSER**, a pneumatic engine, or syringe, by which an extraordinary quantity of air may be crowded or pushed into a given space; so that frequently ten atmospheres, or ten times as much air as the space naturally contains, without the engine, may be thrown in by means of it, and its egress prevented by valves properly disposed.

The condenser is made either of metal, or glass, and either in a cylindrical or globular form, into which the air is thrown with an injecting syringe.

The receiver, or vessel containing the condensed air, should be made very strong, to bear the force of the air's elasticity thus increased; for which reason it is commonly made of brass. When glass is used, it will not bear so great a condensation of air; but then the experiment will be more entertaining, as the effect may be viewed of the condensed air upon any subject put within it.

**CONDUCTOR**, in Electricity, a term first introduced in this science by Dr. Defaguliers, and used to denote those substances which are capable of receiving and transmitting electricity; in opposition to electrics, in which the matter or virtue of electricity may be excited and accumulated, or retained. The former are also called *non-electrics*, and the latter *non-conductors*. And all bodies are ranked under one or other of these two classes, though none of them are *perfect electrics*, nor *perfect conductors*, so as wholly to retain, or freely and without resistance to transmit the electric fluid.

To the class of conductors, belong all metals and semi-metals, ores, and all fluids (except air and oils), together with the substances containing them, the effluvia of flaming bodies, ice (unless very hard frozen), and snow, most saline and stony substances, charcoals, of which the best are those that have been exposed to the greatest heat, smoke, and the vapour of hot water.

It seems probable that the electric fluid passes through the substance, and not merely over the surfaces of metallic conductors; because, if a wire of any kind of metal be covered with some electric substance, as resin, sealing-



Sealing-wax, &c, and a jar be discharged through it, the charge will be conducted as well as without the electric coating.

It has also been alleged, that electricity will pervade a vacuum, and be transmitted through it almost as freely as through the substance of the best conductor: but Mr. Wallh found, that the electric spark or shock would no more pass through a perfect vacuum, than through a stick of solid glass. In other instances however, when the vacuum has been made with all possible care, the experiment has not succeeded.

It may also be observed, that many of the forementioned substances are capable of being electrified, and that their conducting power may be destroyed and recovered by different processes: for example, green wood is a conductor; but baked, it becomes a non-conductor: again its conducting power is restored by charring it; and lastly it is destroyed by reducing this to ashes.

Again, many electric substances, as glass, resin, air, &c, become conductors by being made very hot: however, air heated by glass must be excepted.

See, on this subject, Priestley's *Hist. of Electricity*, vol. 1; Franklin's *Letters &c*, pa. 96 and 262 edit. 1769; Cavallo's *Complete Treat. of Electr.* chap. 2; Henley's *Exper. and Obser. in Electr.* also *Philos. Transf.* vol. 67 pa. 122; and elsewhere in the different volumes of the *Transactions*.

*Prime CONDUCTOR*, is an insulated conductor, so connected with the electrical machine, as to receive the electricity immediately from the excited electric.

Mr. Grey first employed metallic conductors in this way, in 1734; and these were several pieces of metal suspended on filken strings, which he charged with electricity. Mr. Du Fay fastened to the end of an iron bar, which he used as his prime conductor, a bundle of linen threads, to which he applied the excited tube: but these were afterwards changed for small wires suspended from a common gun-barrel, or other metallic rod.

In the present advanced state of the science, this part of the electrical apparatus has been considerably improved. The prime conductor is made of hollow brass, and usually of a cylindrical form. Care should be taken, that it be perfectly smooth and round, without points and sharp edges. The ends of the conductor are spherical; and it is necessary, that the part most remote from the electric should be made round and much larger than the rest, the better to prevent the electric matter from escaping, which it always endeavours most to do at the greatest distance from the electric: and the other end should be furnished with several pointed wires or needles, either suspended from, or fixed to an open metallic ring, and pointing to the globe or cylinder, or plate, to collect the fire. It is best supported by pillars of solid glass, covered with sealing-wax or good varnish. Prime conductors of a large size are usually made of paste-board, covered with tin-foil or gilt paper; these being useful for throwing off a longer and denser spark than those of a smaller size: they should terminate in a smaller knob or obtuse edge, at which the sparks should be solicited. Mr. Nairne prepared a conductor 6 feet in length, and 1 foot in diameter, from which he drew electrical sparks at the distance of 16, 17, or 18 inches; and Dr. Van Marum

still far exceeded this, with a conductor of 8 inches diameter, and upwards of 20 feet long, formed of different pieces, and applied to the large electrical machine in Teyler's Museum at Harlem, the most powerful machine of the kind ever yet constructed. But the size of the conductor is always limited by that of the electric, there being a maximum which the size of the former should not exceed; for it may be so large, that the dissipation of the electricity from its surface may be greater than that which the electric is capable of supplying.

Dr. Priestley recommends a prime conductor of polished copper, in the form of a pear, supported by a pillar and a firm basis of baked wood: this receives its fire by a long arched wire of soft brass, which may be easily bent, and raised or lowered to the globe: it is terminated by an open ring, in which some sharp-pointed wires are hung. In the body of this conductor are holes for the insertion of metalline rods. This, he says, collects the fire perfectly well, and retains it equally everywhere. *Philos. Transf.* vol. 64, art. 7. *Hist. Electr.* vol. 2, § 2.

Mr. Henley has contrived a new kind of prime conductor, which, from its use, is called the *luminous conductor*. It consists of a glass tube 18 inches long, and 2 inches diameter. The tube is furnished at both ends with brass caps and females about 2 inches long, cemented and made air-tight, and terminated by brass balls. In one of these caps is drilled a small hole, which is covered by a strong valve, and serves for exhausting the tube of its air. Within the tube at each end there is a knobbed wire, projecting to the distance of 2 inches and a half from the brass caps. To one of the balls is annexed a fine-pointed wire for receiving and collecting the electricity, and to the other a wire with a knob or ball for discharging it. The conductor, thus prepared, is supported on pillars of sealing-wax or glass. Beside the common purposes of a prime conductor to an electrical machine, this apparatus serves to exhibit and ascertain the direction of the electric matter in its passage through it. See a figure of this conductor in the *Philos. Transf.* with a description of experiments, &c. with it, vol. 64, pa. 403.

*CONDUCTORS of Lightning*, are pointed metallic rods fixed to the upper parts of buildings, to secure them from strokes of lightning. These were invented and proposed by Dr. Franklin for this purpose, soon after the identity of electricity and lightning was ascertained; and they exhibit a very important and useful application of modern discoveries in this science. This ingenious philosopher, having found that pointed bodies are better fitted for receiving and throwing off the electric fire, than such as are terminated by blunt ends or flat surfaces, and that metals are the readiest and best conductors, soon discovered that lightning and electricity resembled each other in this and other distinguishing properties: he therefore recommended a pointed metalline rod, to be raised some feet above the highest part of a building, and to be continued down into the ground, or the nearest water. The lightning, should it ever come within a certain distance of this rod or wire, would be attracted by it, and pass through it preferably to any other part of the building, and be conveyed into the earth or water, and there dissipated, without doing any damage to the building. Many facts have occurred



red to evince the utility of this simple and seemingly trifling apparatus. And yet some electricians, of whom Mr. Wilfon was the chief, have objected to the pointed termination of this conductor; preferring rather a blunt end: because, they pretend, a point invites the electricity from the clouds, and attracts it at a greater distance than a blunt conductor. *Philos. Transf.* vol. 54, pa. 234; vol. 63, pa. 49; and vol. 68, pa. 232.

This subject has indeed been very accurately examined and discussed; and pointed conductors are almost universally, and for the best reasons, recommended as the most proper and eligible. A sharp-pointed conductor, as it attracts the electric fire of a cloud at a greater distance than the other, draws it off gradually: and by conveying it away gently, and in a continued stream, prevents an accumulation and a stroke; whereas a conductor with a blunt termination receives the whole discharge of a cloud at once, and is much more likely to be exploded, whenever a cloud comes within a striking distance. To this may be added experience; for buildings guarded by either natural or artificial conductors terminating in a point, have very seldom been struck by lightning; but others, having flat or blunt terminations, have often been struck and damaged by it.

The best conductor for this purpose, is a rod of iron, or rather of copper, as being a better conductor of electricity, and less liable to rust, about 3 quarters of an inch thick, which is either to be fastened to the walls of a building by wooden cramps, or supported by wooden posts, at the distance of a foot or two from the wall; though less may do: the upper end of it should terminate in a pyramidal form, with a sharp point and edges; and, when made of iron, gilt or painted near the top, or else pointed with copper; and be elevated 5 or 6 feet above the highest part of the building, or chimneys, to which it may be fastened. The lower end should be driven 5 or 6 feet into the ground, and directed away from the foundations of the building, or continued till it communicates with the nearest water: and if this part be made of lead, it will be less apt to decay. When the conductor is formed of different pieces of metal, care should be taken that they are well joined: and it is farther recommended, that a communication be made from the conductor by plates of lead, 8 or 10 inches broad, with the lead on the ridges and gutters, and with the pipes that carry down the rain water, which should be continued to the bottom of the building, and be made to communicate either with water or moist earth, or with the main pipe which serves the house with water. If the building be large, two, three, or more conductors should be applied to different parts of it, in proportion to its extent. *Philos. Transf.* vol. 64, pa. 403.

Chains have been used as conductors for preserving ships; but as the electric matter does not pass readily through the links of it, copper wires, a little thicker than a goose quill, have been preferred, and are now generally used. They should reach 2 or 3 feet above the highest mast, and be continued down in any convenient direction, so as always to touch the sea water. *Philos. Transf.* vol. 52, pa. 633. See also Franklin's *Letters &c* 1769, pa. 65, 124, 479, &c; and Cavallo's *Electr.* chap. 9.


For the *Construction* and management of *Electrical Kites*, and *Conduitors* or *Machines* for drawing electricity from the clouds, see Priestley's *Hist. of Electr.* vol. 2, pa. 103 edit. 1775.

CONE, a kind of round pyramid, or a solid body having a circle for its base, and its sides formed by right lines drawn from the circumference of the base to a point at top, being the vertex or apex of the cone.

Euclid defines a cone to be a solid figure, whose base is a circle, and is produced by the entire revolution of a right-angled triangle about its perpendicular leg, called the axis of the cone. If this leg, or axis, be greater than the base of the triangle, or radius of the circular base of the cone, then the cone is *acute-angled*, that is, the angle at its vertex is an acute angle; but if the axis be less than the radius of the base, it is an *obtuse-angled* cone; and if they are equal, it is a *right-angled* cone.

But Euclid's definition only extends to a *right cone*, that is, to a cone whose axis is perpendicular or at right-angles to its base; and not to oblique ones, in which the axis is oblique to the base, the general definition, or description of which may be this:

If a line VA continually pass through the point V, turning upon that point as a joint, and the lower part of it be carried round the circumference ABC of a circle; then the space inclosed between that circle and the path of the line, is a cone. The circle ABC is the base of the cone; V is its vertex; and the line VD, from the vertex to the centre of the



base, is the axis of the cone. Also the other part of the revolving line, produced above V, will describe another cone *Vacb*, called the opposite cone, and having the same common axis produced DVd, and vertex V.

*Properties of the CONE.*—1. The area or surface of every right cone, exclusive of its base, is equal to a triangle whose base is the periphery, and its height the slant side of the cone. Or, the curve superficies of a right cone, is to the area of its circular base, as the slant side is to the radius of the base. And therefore, the same curve surface of the cone is equal to the sector of a circle whose radius is the slant side, and its arch equal to the circumference of the base of the cone.

2. Every cone, whether right or oblique, is equal to one-third part of a cylinder of equal base and altitude; and therefore the solid content is found by multiplying the base by the altitude, and taking  $\frac{1}{3}$  of the product; and hence also all cones of the same or equal base and altitude, are equal.

3. Although the solidity of an oblique cone be obtained in the same manner with that of a right one, it is otherwise with regard to the surface, since this cannot be reduced to the measure of a sector of a circle, because all the lines drawn from the vertex to the base are not equal. See a *Memoir* on this subject, by M. Euler, in the *Nouv. Mem. de Petersburg* vol. 1. Dr. Barrow has demonstrated, in his *Lectiões Geometricæ*, that the solidity of a cone with an elliptic base, forming part of a right cone, is equal to the product of its surface by a third part of one of the perpendiculars



drawn from the point in which the axis of the right cone intersects the ellipse; and that it is also equal to  $\frac{1}{3}$  of the height of the cone multiplied by the elliptic base: consequently that the perpendicular is to the height of the cone, as the elliptic base is to the curve surface. For the curve surface of all the oblique parts of a cone, see my Mensur. pa. 234 &c.

4. To find the Curve Surface of the Frustum of a Cone. Multiply half the sum of the circumferences of the two ends, by the slant side, or distance between these circumferences.

5. For the Solidity of the Frustum of a Cone, add into one sum the areas of the two ends and the mean proportional between them, multiply that sum by the perpendicular height, and  $\frac{1}{3}$  of the product will be the solidity. See also my Mensuration, pa. 189.

6. The Centre of Gravity of a cone is  $\frac{3}{4}$  of the axis distant from the vertex.

CONES of the Higher Kinds, are those whose bases are circles of the higher kinds; and are generated, like the common cone, by conceiving a line turning on a point or vertex on high, and revolving round the circle of the higher kind.

CONE of Rays, in Optics, includes all the several rays which fall from any point of a radiant object, on the surface of a glass.

DOUBLE CONE, or Spindle, in Mechanics, is a solid formed of two equal cones joined at their bases. If this be laid on the lower part of two rulers, making an angle with each other, and elevated in a certain degree above the horizontal plane, the cones will roll up towards the raised ends, and seem to ascend, though in reality its centre of gravity descends perpendicularly lower.

CONFIGURATION, the exterior surface or shape that bounds bodies, and gives them their particular figure.

CONFIGURATION of the planets, in Astrology, is a certain distance or situation of the planets in the zodiac, by which it is supposed that they assist or oppose each other.

CONFUSED Vision. See VISION.


CONGELATION, or FREEZING, the act of fixing the fluidity of any liquid, by cold, or the application of cold bodies: in which it differs from coagulation, which is produced by other causes. See FREEZING, FROST, and ICE.

CONGRUITY, in Geometry, is applied to lines and figures, which exactly correspond when laid over one another; as having the same terms, or bounds. It is assumed, as an axiom, that those things are equal and similar, between which there is a congruity. Euclid, and most geometers after him, demonstrate great part of their elements from the principle of congruity: though Leibnitz and Wolfius substitute the notion of Similitude instead of that of congruity.


CONIC SECTIONS, are the figures made by cutting a cone by a plane.

2. According to the different positions of the cutting plane, there arise five different figures or sections, viz, a triangle, a circle, an ellipse, a parabola, and an hyperbola: the last three of which only are peculiarly called conic sections.


3. If the cutting plane pass through the vertex of the cone, and any part of the base, the section will evidently be a triangle; as VAB.




4. If the plane cut the cone parallel to the base, or make no angle with it, the section will be a circle, as ABD.




5. The section DAB is an ellipse, when the cone is cut obliquely through both sides, or when the plane is inclined to the base in a less angle than the side of the cone is.



6. The section is a parabola, when the cone is cut by a plane parallel to the side, or when the cutting plane and the side of the cone make equal angles with the base.



7. The section is an hyperbola, when the cutting plane makes a greater angle with the base than the side of the cone makes. And if the plane be continued to cut the opposite cone, this latter section is called the opposite hyperbola to the former; as dBe.




8. The vertices of any section, are the points where the cutting plane meets the opposite sides of the cone, or the sides of the vertical triangular section; as A and B. —Hence, the ellipse and the opposite hyperbolas have each two vertices; but the parabola only one; unless we consider the other as at an infinite distance.

9. The axis, or transverse diameter of a conic section, is the line or distance AB between the vertices. —Hence the axis of a parabola is infinite in length.

Ellipse.

Oppos. Hyperb.

Parabola.



10. The centre C is the middle of the axis. —Hence the centre of a parabola is infinitely distant from the vertex. And of an ellipse, the axis and centre lie within the curve; but of an hyperbola, without.



11. A *Diameter* is any right line, as  $AB$  or  $DE$ , drawn through the centre, and terminated on each side by the curve: and the extremities of the diameter, or its intersections with the curve, are its *vertices*.—Hence all the diameters of a parabola are parallel to the axis, and infinite in length; because drawn through the centre, a point at an infinite distance. And hence also every diameter of the ellipse and hyperbola have two vertices; but of the parabola, only one; unless we consider the other as at an infinite distance.


12. The *conjugate* to any diameter, is the line drawn through the centre, and parallel to the tangent of the curve at the vertex of the diameter. So  $FG$ , parallel to the tangent at  $D$ , is the conjugate to  $DE$ ; and  $HI$ , parallel to the tangent at  $A$ , is the conjugate to  $AB$ .—Hence the conjugate  $HI$ , of the axis  $AB$ , is perpendicular to it; but the conjugates of other diameters are oblique to them.

13. An *ordinate* to any diameter, is a line parallel to its conjugate, or to the tangent at its vertex, and terminated by the diameter and curve. So  $DK$  and  $EL$  are ordinates to the axis  $AB$ ; and  $MN$  and  $NO$  ordinates to the diameter  $DE$ .—Hence the ordinates of the axis are perpendicular to it; but of other diameters, the ordinates are oblique to them.

14. An *absciss* is a part of any diameter, contained between its vertex and an ordinate to it; as  $AK$  or  $BK$ , and  $DN$  or  $EN$ .—Hence, in the ellipse and hyperbola, every ordinate has two abscisses; but in the parabola, only one; the other vertex of the diameter being infinitely distant.

15. The *parameter* of any diameter, is a third proportional to that diameter and its conjugate.

16. The *focus* is the point in the axis where the ordinate is equal to half the parameter: as  $K$  and  $L$ , where  $DK$  or  $EL$  is equal to the femiparameter.—Hence, the ellipse and hyperbola have each two foci; but the parabola only one.—The foci, or burning points, were so called, because all rays are united or reflected into one of them, which proceed from the other focus, and are reflected from the curve.



17. If  $DAE$ ,  $FBG$  be two opposite hyperbolas, having  $AB$  for their first or transverse axis, and  $ab$  for their second or conjugate axis; and if  $dae$ ,  $fbg$  be two other opposite hyperbolas, having the same axis, but in the contrary order, viz,  $ab$  their first axis, and  $AB$  their second; then these two latter curves  $dae$ ,  $fbg$ , are called the *conjugate hyperbolas* to the two former  $DAE$ ,  $FBG$ ; and each pair of opposite curves mutually conjugate to the other.

18. And if tangents be drawn to the four vertices of the curves, or extremities of the axis, forming the inscribed rectangle  $HIKL$ ; the diagonals  $HCK$  and  $ICL$ , of this rectangle, are called the *asymptotes* of the curves.

19. *Scholium*. The rectangle inscribed between the


four conjugate hyperbolas, is similar to a rectangle circumscribed about an ellipse, by drawing tangents, in like manner, to the four extremities of the two axes; also the asymptotes or diagonals in the hyperbola, are analogous to those in the ellipse, cutting this curve in similar points, and making the pair of equal conjugate diameters. Moreover, the whole figure, formed by the four hyperbolas, is, as it were, an ellipse turned inside out, cut open at the extremities  $D$ ,  $E$ ,  $F$ ,  $G$ , of the said equal conjugate diameters, and those four points drawn out to an infinite distance, the curvature being turned the contrary way, but the axes, and the rectangle passing through their extremities, remaining fixed, or unaltered.

From the foregoing definitions are easily derived the following general corollaries to the sections.

Ellipse.

Hyperbola.

Parabola.



20. *Corol. 1*. In the ellipse, the semiconjugate axis,  $CD$  or  $CE$ , is a mean proportional between  $CO$  and  $CP$ , the parts of the diameter  $OP$  of a circular section of the cone, drawn through the centre  $C$  of the ellipse, and parallel to the base of the cone. For  $DE$  is a double ordinate in this circle, being perpendicular to  $OP$  as well as to  $AB$ .

21. In like manner, in the hyperbola, the length of the semiconjugate axis,  $CD$  or  $CE$ , is a mean proportional between  $CO$  and  $CP$ , drawn parallel to the base, and meeting the sides of the cone in  $O$  and  $P$ . Or, if  $AO'$  be drawn parallel to the side  $VB$ , and meet  $PC$  produced in  $O'$ , making  $CO' = CO$ ; and on this diameter  $O'P$  a circle be drawn parallel to the base: then the semiconjugate  $CD$  or  $CE$  will be an ordinate of this circle, being perpendicular to  $O'P$  as well as to  $AB$ .

Or, in both figures, the whole conjugate axis  $DE$  is a mean proportional between  $QA$  and  $BR$ , parallel to the base of the cone. See my Conic Sections, pa. 6.

In the parabola, both the transverse and conjugate are infinite; for  $AB$  and  $BR$  are both infinite.

22. *Corol. 2*. In all the sections,  $AG$  will be equal to the parameter of the axis, if  $QG$  be drawn making the angle  $AQG$  equal to the angle  $BAR$ . In like manner  $Bg$  will be equal to the same parameter, if  $Rg$  be drawn to make the angle  $BRg =$  the angle  $ABQ$ .

23. *Corol. 3*. Hence the upper hyperbolic section, or section of the opposite cone, is equal and similar to the lower one. For the two sections have the same transverse or first axis  $AB$ , and the same conjugate or second axis  $DE$ , which is the mean proportional between  $AQ$  and  $RB$ ; and they have also equal parameters.



ters AG, Bg. So that the two opposite sections make, as it were, but the two opposite ends of one entire section or hyperbola, the two being every where mutually equal and similar. Like the two halves of an ellipse, with their ends turned the contrary way:

24. *Corol. 4.* And hence, although both the transverse and conjugate axis in the parabola be infinite, yet the former is infinitely greater than the latter, or has an infinite ratio to it. For the transverse has the same ratio to the conjugate, as the conjugate has to the parameter, that is, as an infinite to a finite quantity, which is an infinite ratio.

The peculiar properties of each particular curve, will be best referred to the particular words ELLIPSE, HYPERBOLA, PARABOLA; and therefore it will only be proper here to lay down a few of the properties that are common to all the conic sections.

*Some other General Properties.*


25. From the foregoing definitions, &c, it appears, that the conic sections are in themselves a system of regular curves, naturally allied to each other; and that one is changed into another perpetually, when it is either increased, or diminished, in infinitum. Thus, the curvature of a circle being ever so little increased or diminished, passes into an ellipse; and again, the centre of the ellipse going off infinitely, and the curvature being thereby diminished, is changed into a parabola; and lastly, the curvature of a parabola being ever so little changed, there ariseth the first of the hyperbolas; the innumerable species of which will all of them arise orderly by a gradual diminution of the curvature; till this quite vanishing, the last hyperbola ends in a right line. From whence it is manifest, that every regular curvature, like that of a circle, from the circle itself to a right line, is a conical curvature, and is distinguished with its peculiar name, according to the divers degrees of that curvature.

26. That all diameters in a circle and ellipse intersect one another in the centre of the figure within the section: that in the parabola they are all parallel among themselves, and to the axis: but in the hyperbola, they intersect one another, without the figure, in the common centre of the opposite and conjugate sections.

27. In the circle, the *latus rectum*, or parameter, is double the distance from the vertex to the focus, which is also the centre. But in ellipses, the parameters are in all proportions to that distance, between the double and quadruple, according to their different species. While, in the parabola, the parameter is just quadruple that distance. And, lastly in hyperbolas, the parameters are in all proportions beyond the quadruple, according to their various kinds.

28. The first general property of the conic sections, with regard to the abscisses and ordinates of any diameter, is, that the rectangles of the abscisses are to each other, as the squares of their corresponding ordinates. Or, which is the same thing, that the square of any diameter is to the square of its conjugate, as the rectangle of two abscisses of that diameter, to the square of the ordinate which divides them. That is, in all the figures,

the rect. AC . CB : rect. AE . EB :: CD<sup>2</sup> : EF<sup>2</sup> :



But as, in the parabola the infinites CB and EB are in a ratio of equality, for this curve the same property becomes  $AC : AE :: CD^2 : EF^2$ , that is, in the parabola, the abscisses are as the squares of their ordinates.

Or, when one of the ordinates is the semiconjugate GH, dividing the diameter equally in the centre, the same general property becomes,

$AG . GB$  or  $AG^2 : AC . CB :: GH^2 : CD^2$ ,  
or  $AB^2 : HI^2 :: AC . CB : CD^2$ .

29. From hence is derived the equation of the curves of the conic sections; thus, putting the diameter  $AB = d$ , its conjugate  $HI = c$ , absciss  $AC = x$ , and its ordinate  $CD = y$ ; then is the other absciss  $CB = d - x$  in the ellipse, or  $d + x$  in the hyperbola, or  $d$  in the parabola; and hence the last analogy above, becomes  $d^2 : c^2 :: x . d \mp x$  or  $dx \mp x^2 : y^2$ ,

or  $dy^2 y^2 = c^2 . dx \mp x^2$  is the general equation for all the conic sections; and, in particular, it is

$d^2 y^2 = c^2 . dx - x^2$  in the ellipse,

$d^2 y^2 = c^2 . dx + x^2$  in the hyperbola, and

$d^2 y^2 = c^2 dx$ , or  $dy^2 = c^2 x$  in the parabola: Or the three equations may be otherwise expressed thus:

$y^2 = \frac{c^2}{d^2} . dx - x^2$  in the ellipse,

$y^2 = \frac{c^2}{d^2} . dx + x^2$  in the hyperbola, and

$y^2 = \frac{c^2}{d} x$  or  $= px$  in the parabola, where the para-

meter  $p = \frac{c^2}{d}$  the third proportional to the diameter and its conjugate, by the definition of it.

And from this one general proposition alone, which is easily derived from the section in the solid cone itself, together with the definitions only, as laid down above, all the other properties of all the sections may easily be derived, without any farther reference to the cone, and without mechanical descriptions of the curves in plano; as is done in my Treatise on Conic Sections, for the use of the Royal Mil. Acad.; in which also all the similar propositions in the ellipse and hyperbola are carried on word for word in them both.

The more ancient mathematicians, before the time of Apollonius Pergæus, admitted only the right cone into their geometry, and they supposed the section of it to be made by a plane perpendicular to one of its sides; and as the vertical angle of a right cone may be either right, acute, or obtuse, the same method of cutting these several cones, viz, by a plane perpendicular to one side, produced all the three conic sections. The parabola was called the section of a right-angled cone; the ellipse, the section of the acute-angled cone; and the hyperbola, the section of the obtuse-angled cone. But Apollonius, who, on account of his writings on this subject,



subject, obtained the appellation of *Magnus Geometra*, the Great Geometrician, observed, that these three sections might be obtained in every cone, both oblique and right, and that they depended on the different inclinations of the plane of the section to the cone itself. Apollon. Con. Halley's edit. lib. 1, p. 9.

Instead of considering these curves as sections cut from the solid cone, which is the true genuine way of all the ancients, and of the most elegant writers among the moderns, Descartes, and some others of the moderns, have given arbitrary constructions of curves on a plane, from which constructions they have demonstrated the properties of these, and have afterwards proved that some principal property of them belongs to such curves or sections as are cut from a cone; and hence it is inferred by them that those curves, so described on a plane, are the same with the conic sections.

The doctrine of the conic sections is of great use in physical and geometrical astronomy, as well as in the physico-mathematical sciences. The doctrine has been much cultivated by both ancient and modern geometricians, who have left many good treatises on the subject. The most ancient of these is that of Apollonius Pergæus, containing 8 books, the first 4 of which have often been published; but Dr. Halley's edition has all the eight. Pappus, in his Collect. Mathem. lib. 7, says that the first four of these were written by Euclid, though perfected by Apollonius, who added the other 4 to them. Among the moderns, the chief writers are Mydorgius de Sectionibus Conicis; Gregory St. Vincent's Quadratura Circuli & Sectionum Coni; De la Hire de Sectionibus Conicis; Trevigar Elem. Section. Con.; De Witt's Elementa Curvarum; Dr. Wallis's Conic Sections; De l'Hospital's Anal. Treat. of Conic Sections; Dr. Simson's Section. Con.; Milne's Elementa Section. Conicarum; Muller's Conic Sections; Steel's Conic Sections; Dr. Hamilton's elegant treatise; my own treatise, above cited; and at the writing of this, my friend Mr. Abram Robertson of Oxford is preparing a curious work on this subject, containing at the same time a treatise on the science, and a history of the writings relating to it.

CONICS, that part of the higher geometry, or geometry of curves, which considers the cone, and the several curve lines arising from the sections of it.

CONJUGATE *Axis*, or *Diameter*, in the Conic Sections, is the axis, or a diameter parallel to a tangent to the curve at the vertex of another axis, or diameter, to which that is a conjugate. Indeed the two are mutually conjugates to each other, and each is parallel to the tangent at the vertex of the other.

CONJUGATE HYPERBOLAS, also called *Adjacent Hyperbolas*, are such as have the same axes, but in the contrary order, the first or principal axis of the one being the 2d axis of the other, and the 2d axis of the former, the 1st axis of the latter. See art. 17 of CONIC SECTIONS.

CONJUNCTION, in Astronomy, is the meeting of the stars and planets in the same point or place in the heavens; and is either true or apparent.

True CONJUNCTION is when the line drawn through the centres of the two stars passes also through the centre of the earth. And Apparent CONJUNCTION is when that line does not pass through the earth's centre.

CONOID, is a figure resembling a cone, except that the slant sides from the base to the vertex are not straight lines as in the cone, but curved. It is generated by the revolution of a conic section about its axis; and it is therefore threefold, answering to the three sections of the cone, viz, the *Elliptical Conoid*, or spheroid, the *Hyperbolic Conoid*, and the *Parabolic Conoid*.

If a conoid be cut by a plane in any position, the section will be of the figure of some one of the conic sections; and all parallel sections, of the same conoid, are like and similar figures. When the section of the solid returns into itself, it is an ellipse; which is always the case in the sections of the spheroid, except when it is perpendicular to the axis; which position is also to be excepted in the other solids, the section being always a circle in that position. In the parabolic conoid, the section is always an ellipse, except when it is parallel to the axis. And in the hyperbolic conoid, the section is an ellipse, when its axis makes with the axis of the solid, an angle greater than that made by the said axis of the solid and the asymptote of the generating hyperbola; the section being an hyperbola in all other cases, but when those angles are equal, and then it is a parabola.

But when the section is parallel to the fixed axis, it is of the same kind with, and similar to the generating plane itself; that is, the section parallel to the axis, in the spheroid, is an ellipse similar to the generating ellipse; in the parabolic conoid it is a parabola, similar to the generating one; and in the hyperbolic conoid, it is an hyperbola similar to the generating one.

The section through the axis, which is the generating plane, is, in the spheroid the greatest of the parallel sections, but in the hyperboloid it is the least, and in the paraboloid those parallel sections are all equal.

The analogy of the sections of the hyperboloid to those of the cone, are very remarkable, all the three conic sections being formed by cutting an hyperboloid in the same positions as the cone is cut. Thus, let an hyperbola and its asymptote be revolved together about the transverse axis, the former describing an hyperboloid, and the latter a cone circumscribing it: then let it be supposed that they are both cut by one plane in any position; so shall the two sections be like, similar, and concentric figures: that is, if the plane cut both the sides of each, the sections will be concentric and similar ellipses; but if the cutting plane be parallel to the asymptote, or to the side of the cone, the sections will be parabolas; and in all other positions, the sections will be similar and concentric hyperbolas.

And this analogy of the sections will not seem strange, when it is considered that a cone is a species of the hyperboloid; or a triangle a species of the hyperbola, the axes being infinitely small. See my Mensuration, prop. 1, part 3, sect. 4, pag. 265 edit. 2vo.

CONON (of Samos), a respectable mathematician and philosopher, who flourished about the 130th olympiad, being a contemporary and friend of Archimedes, to whom Conon communicated his writings, and sent him some problems, which Archimedes received with approbation, saying they ought to be published while Conon was living, for he comprehends them with ease, and can give a proper demonstration of them.

At another time he laments the loss of Conon, thus  
admir



admiring his genius. "How many theorems in geometry, says he, which at first seemed impossible, would in time have been brought to perfection! Alas! Conon, though he invented many, with which he enriched geometry, had not time to perfect them, but left many in the dark, being prevented by death." He had an uncommon skill in mathematics, joined to an extraordinary patience and application. This is farther confirmed by a letter sent to Archimedes by a friend of Conon's. "Having heard of Conon's death, with whose friendship I was honoured, and with whom you kept an intimate correspondence; as he was thoroughly versed in geometry, I greatly lament the loss of a sincere friend, and a person of surprising knowledge in mathematics. I then determined to send to you, as I had before done to him, a theorem in geometry, hitherto observed by no one."

Conon had some disputes with Nicoteles, who wrote against him, and treated him with too much contempt. Apollonius confesses it; though he acknowledges that Conon was not fortunate in his demonstrations.

Conon invented a kind of volute, or spiral, different from that of Dynostriatus; but because Archimedes explained the properties of it more clearly, the name of the inventor was forgotten, and it was hence called Archimedes's volute or spiral.

As to Conon's astrological or astronomical knowledge, it may in some measure be gathered from the poem of Catullus, who describes it in the beginning of his verses on the hair of Berenice, the sister and wife of Ptolomy Euergetes, upon the occasion of Conon having given out that it was changed into a constellation among the stars, to console the queen for the loss, when it was stolen out of the temple, where she had consecrated it to the gods.

CONSECTARY, or *Corollary*, a consequence deduced from some foregoing principles.

CONSEQUENT, is the latter of the two terms of a ratio; or that to which the antecedent is referred and compared. Thus, in the ratio  $a : b$ , or  $a$  to  $b$ , the latter term  $b$  is the consequent, and  $a$  is the antecedent.

CONSISTENT *Bodies*, is a term much used by Mr. Boyle, for such as are usually called *firm*, or *fixed bodies*; in opposition to *fluid* ones.

CONSOLE, in Architecture, is an ornament cut upon the key of an arch, having a projecture or jetting, and occasionally serving to support small cornices, busts, and bases.

CONSONANCE, in Music, is commonly used in the same sense with *concord*, viz, for the union or agreement of two sounds produced at the same time, the one grave, the other acute, which is compounded together by such a proportion of each, as proves agreeable to the ear.

An unison is the first consonance, an eighth is the 2d, a fifth is the 3d; and then follow the fourth, with the third and sixths, major and minor.

CONSTANT *Quantities* are such as remain invariably the same, while others increase or decrease. Thus, the diameter of a circle is a constant quantity; for it remains the same while the abscisses and ordinates, or the sines, tangents, &c, are variable.

These are sometimes called *given*, or *invariable* or *per-*

*manent* quantities; and in algebra it is now usual to represent them by the leading letters of the alphabet,  $a, b, c$ , &c; while the variable ones are denoted by the last letters,  $z, y, x$ , &c.

CONSTELLATIONS, certain imaginary figures of birds, beasts, fishes, and other things in the heavens, within which are arranged certain stars. These assemblages are also sometimes called asterisms.

The ancients portioned out the firmament into several parts, or constellations; reducing a certain number of stars under the representation of certain images, to assist the imagination and memory, to conceive or retain their number, order, and disposition, or even to distinguish the virtues they attributed to them.

The division of the heavens into constellations is very ancient; being known to the most early authors, whether sacred or profane. In the book of Job the names of some of them are mentioned; witness that sublime expostulation, *Canst thou restrain the sweet influence of the Pleiades, or loosen the bands of Orion?* And the same may be observed of the oldest among the heathen writers, Hesiod and Homer.

The division of the ancients took in only the visible firmament, or so much as came under their notice, as visible to the naked eye. The first or earliest of these, is contained in the catalogue of Ptolomy, given in the 7th book of his *Almagest*, prepared, as he assures us, from his own observations, compared with those of Hipparchus, and the other ancient astronomers. In this catalogue Ptolomy has formed 48 constellations. Of these, 12 are about the ecliptic, commonly called the 12 signs; 21 to the north of it; and 15 to the south. The northern constellations are, the Little Bear, the Great Bear, the Dragon, Cepheus, Bootes, the Northern Crown, Hercules, the Harp, the Swan, Cassiopeia, Perseus, Auriga, Ophiucus or Serpentary, the Serpent, the Arrow, the Eagle, the Dolphin, the Horse, Pegasus, Andromeda, and the Triangle.

The constellations about the ecliptic are Aries, Taurus, Gemini, Cancer, Leo, Virgo, Libra, Scorpio, Sagittarius, Capricorn, Aquarius, and Pisces: or according to the English names, the Ram, the Bull, the Twins, the Crab, the Lion, the Virgin, the Balance, the Scorpion, the Archer, the Goat, the Water bearer, and the Fishes.

The Southern constellations are, the Whale, Orion, the Eridanus, the Hare, the Great Dog, the Little Dog, the Ship, the Hydra, the Cup, the Raven, the Centaur, the Wolf, the Altar, the Southern Crown, and the Southern Fish.

The other stars not comprehended under these constellations, yet visible to the naked eye, the ancients called *informes*, or *sporades*, some of which the modern astronomers have since reduced into new figures, or constellations. Ptolomy has set down the longitude and latitude of all these stars to about the year of Christ 137, amounting to the number of 1022, viz,

in the northern constellations	-	-	360
in the zodiacal constellations	-	-	346
in the southern constellations	-	-	316
<hr/>			
in all of Ptolomy's catalogue	-	-	1022

Among the modern astronomers, Tycho Brahe is the first



first who determined, with exactness, and in consequence of his own observations, the long. and lat. of the fixed stars, out of which he formed 45 constellations; of these, 43 were of the old ones described by Ptolomy, to which Tycho added the Coma Berenices, and Antinous; but he omits 5 of the old southern constellations, viz, the Centaur, the Wolf, the Altar, the Southern Crown, and Southern Fish; which he could not observe, because of the high northern latitude of Uranibourg.

After Tycho, Bayer gave the figures of 60 constellations, very exactly represented, and with tables annexed, having added, to the 48 old ones of Ptolomy, the following 12 about the south pole, viz, the Peacock, the Toucan, the Crane, the Phoenix, the Dorado, the Flying Fish, the Hydra, the Chameleon, the Bee, the Bird of Paradise, the Triangle, and the Indian. Besides accurately distinguishing the relative size and the situation of every star, Bayer marks the stars in each constellation with the letters of the Greek and Roman alphabets, setting the first letter  $\alpha$  to the first or principal star in each constellation,  $\beta$  to the 2d in order,  $\gamma$  to the 3d, and so on; a very useful method of noting and describing the stars, which has been used by all astronomers since, and who have farther enlarged this method, by adding the ordinal numbers 1, 2, 3, &c, to the other stars discovered since his time, when any constellation contains more than can be marked by the two alphabets. The number and order of the stars, as mentioned by Bayer, are,

of the 1st magnitude	-	-	17
of the 2d magnitude	-	-	63
of the 3d magnitude	.	.	195
of the 4th magnitude	-	-	415
of the 5th magnitude	-	-	348
of the 6th magnitude	-	-	341
of the unformed stars	-	-	326
			<hr/>
in all	-	-	1706
			<hr/>

After Bayer, a catalogue, with new constellations, was published by Schiller, in 1627, in a work called *Coelum Stellatum Christianum*, the Christian Starry Heaven, in which he substitutes, very improperly, other figures of the constellations, and names, taken from the sacred scriptures, instead of the old ones.

In the year 1665, Riccioli published his *Astronomy Reformed*, containing a catalogue of the stars in 62 constellations, viz, the 60 of Bayer, with the Coma Berenices and Antinous of Tycho. He distributes the stars in all the constellations into four classes. In the first of these classes are contained those stars determined by his own observations, and those of Grimaldi. In the second are those stars which had been ascertained by Tycho Brahe and Kepler. In the 3d are the stars determined by Hipparchus and Ptolomy. And the 4th class consists of those of the southern hemisphere discovered by Navigators, who have ascertained their places in a more or less accurate manner; in which he has marked the longitudes and latitudes for the year 1700, the period to which he has reduced all his observations. This catalogue was followed by a number of celestial schemes and maps of the heavens, published in 1673 by Pardies, who has represented very carefully all the constellations, with the stars they contain. After this, Vi-

talis published a catalogue of the fixed stars in his *Tables of the Primum Mobile*, in which their longitudes and latitudes, with the right-ascensions and declinations are set down for the year 1675.

Some time after this, Royer published maps of the heavens, reduced into 4 tables, with a catalogue of the fixed stars for the year 1700. To the stars marked by Bayer, he adds a number of stars not before seen, with others taken from the tables of Riccioli, and not mentioned by Bayer: he also forms, out of the unformed stars, eleven other constellations. Five of these are to the north, and are called the Giraffe, the River Jordan, the River Tigris, the Sceptre, and the Flower-de-luce; with 6 on the south part, which are the Dove, the Unicorn, the Cross, the Great Cloud, the Little Cloud, and the Rhomboide. To this work Royer has joined the catalogue of the southern stars observed by Dr. Halley at the island of St. Helena.

Hevelius has also improved upon the labours of those who went before him, and collected together several stars of the before unformed class into some new constellations. These are, the Unicorn, the Camelopardalis, described by Bartschius, the Sextant of Urania, the Dogs, the Little Lion, the Lynx, the Fox and Goose, the Sobieski's Crown, the Lizard, the Little Triangle, and the Cerberus; to which Gregory has added the Ring and the Armilla. Some of these new constellations however answer to those of Royer, as the Camelopardal to the Giraffe, the Dogs to the River Jordan, and the Fox to the River Tigris. The latitudes and longitudes are added for the year 1700.

Finally, Flamsteed has given a catalogue of the fixed stars, not only much more correct, but much larger than those of all that went before him. He has set down the longitude, latitude, right ascension, and polar distance of 2934 stars, as they were at the beginning of 1690, all determined from his own observations. He distinguishes all the stars into seven classes, or orders of magnitude, distinguishing those of Bayer by his letters, and marking their variation in right ascension, for shewing their situation in the succeeding years. See the term *CATALOGUE*.

This catalogue was followed by an *Atlas Cœlestis*, published at London in the year 1729, describing, in several schemes, the figures of the constellations seen in our hemisphere, with the exact position of the fixed stars, with respect to the circles of the sphere, as resulting from the last catalogue corrected by Flamsteed. And still later observations, made with farther improved telescopes, have greatly enlarged the number and accuracy of the stars; but the number of the constellations remains the same as above described, except that an attempt has lately been made by Dr. Hill to add to the list 14 new ones, formed out of more of the clusters of unformed stars.

Beside the literal marks of the stars introduced by Bayer, it is usual also to distinguish them by that part of the constellation in which they are placed; and many of them again have their peculiar names; as Arcturus, between the knees of Bootes; Gemina, or Lucida, in the Corona Septentrionalis, or Northern Crown; Palilium, or Aldebaran, in the Bull's eye, Pleiades in his neck, and Hyades in his forehead; Castor and Pollux in the heads of Gemini; Capella, with the Hædi in the



shoulder of Auriga; Regulus, or Cor Leonis, the Lion's Heart; Spica Virginis in the hand, and Vindemiatrix in the shoulder of Virgo; Antares, or Cor Scorpionis, the Scorpion's Heart; Fomalhaut, in the mouth of Piscis Australis, or Southern Fish; Regel, in the foot of Orion; Sirius, in the mouth of Canis Major, the Great Dog; Procyon, in the back of Canis Minor, the Little Dog; and the Pole Star, the last in the tail of Urfa Minor, the Little Bear.

The Greek and Roman poets, from the ancient theology, give wild and romantic fables about the origin of the constellations, probably derived from the hieroglyphics of the Egyptians, and transmitted, with some alterations, from them to the Greeks, who probably obscured them greatly with their own fables. See Hyginus's *Poeticon Astron.*; Riccioli *Almagest*. lib. 6. cap. 3, 4, 5; Shelburne's *Notes upon Manilius*; Bailly's *Antient Astronomy*; and Gebelin's *Monde Primitif*, vol. 4: from the whole of which it is made probable, that the invention of the signs of the zodiac, and probably of most of the other constellations of the sphere, is to be ascribed to some very ancient nation, inhabiting the northern temperate zone, probably what is now called Tartary, or the parts to the northward of Persia and China; and from thence transmitted through China, India, Babylon, Arabia, Egypt, Greece, &c.

It is a very probable conjecture, that the figures of the signs in the zodiac, are descriptive of the seasons of the year, or months, in the sun's path: thus, the first sign Aries, denotes, that about the time when the sun enters that part of the ecliptic, the lambs begin to follow the sheep; that on the sun's approach to the 2d constellation, Taurus, the Bull, is about the time of the cows bringing forth their young. The third sign, now Gemini, was originally two kids, and signified the time of the goats bringing forth their young, which are usually two at a birth, while the former, the sheep and cow, commonly produce only one. The 4th sign, Cancer, the Crab, an animal that goes side-ways and backwards, was placed at the northern solstice, the point where the sun begins to return back again from the north to the southward. The 5th sign, Leo, the Lion, as being a very furious animal, was thought to denote the heat and fury of the burning sun, when he has left Cancer, and entered the next sign Leo. The succeeding constellation, the 6th in order, received the sun at the time of ripening corn and approaching harvest; which was aptly expressed by one of the female reapers, with an ear of corn in her hand; viz, Virgo the maid. The ancients gave to the next sign Scorpio, two of the 12 divisions of the zodiac: Autumn, which affords fruits in great abundance, affords the means and causes of diseases, and the succeeding time is the most unhealthy of the year; expressed by this venomous animal, here spreading out his long claws into the one sign, as threatening mischief, and in the other brandishing his tail to denote the completion of it. The fall of the leaf was the season of the ancient hunting; for which reason the stars which marked the sun's place at this season, into the constellation Sagittary, a huntsman with his arrows and his club, the weapons of destruction for the large creatures he pursued. The reason of the Wild Goat's being chosen to mark the southern sol-

stice, when the sun has attained his extreme limit that way, and begins to return and mount again to the northward, is obvious enough; the character of that animal being, that it is mostly climbing, and ascending some mountain as it browses. There yet remain two of the signs of the zodiac to be considered with regard to their origin, viz, Aquarius and Pisces. As to the former, it is to be considered that the winter is a wet and uncomfortable season; this therefore was expressed by Aquarius, the figure of a man pouring out water from an urn. The last of the zodiacal constellations was Pisces, a couple of fishes, tied together, that had been caught: The lesson was, the severe season is over, your flocks do not yet yield their store; but the seas and rivers are open, and there you may take fish in abundance.

Through a vain and blind zeal, rather than through any love for the science, some persons have been moved to alter either the figures of the constellations, or their names. Thus, venerable Bede, instead of the profane names and figures of the twelve zodiacal constellations, substituted those of the 12 apostles; which example was followed by Schiller, who completed the reformation, and gave scripture names to all the constellations in the heavens. Thus, Aries, or the Ram, was changed into Peter; Taurus, or the Bull, into St. Andrew; Andromeda, into the Sepulchre of Christ; Lyra, into the Manger of Christ; Hercules, into the Magi coming from the East; the Great Dog, into David; and so on. And Weigelius, professor of Mathematics in the university of Jena, made a new order of constellations; changing the firmament into a *Cælum Heraldicum*; and introducing the arms of all the princes in Europe, by way of constellations. Thus Urfa major, the Great Bear, he transformed into the elephant of the kingdom of Denmark; the Swan, into the Ruta with swords of the House of Saxony; Ophiuchus, into the Cross of Cologne; the Triangle, into Compasses, which he calls the Symbol of Artificers; and the Pleiades into the Abacus Pythagoricus, which he calls that of merchants; &c.

But the more judicious among astronomers never approved of such innovations; as they only tend to introduce confusion into astronomy. The old constellations are therefore still retained; both because better could not be substituted, and likewise to keep up the greater correspondence and uniformity between the old astronomy and the new. See CATALOGUE.

**CONSTRUCTION**, in Geometry, the art or manner of drawing or describing a figure, scheme, the lines of a problem, or such like.

**CONSTRUCTION of Equations**, in Algebra, is the finding the roots or unknown quantities of an equation, by geometrical construction of right lines or curves; or the reducing given equations into geometrical figures. And this is effected by lines or curves according to the order or rank of the equation.

The roots of any equation may be determined, that is, the equation may be constructed, by the intersections of a straight line with another line or curve of the same dimensions as the equation to be constructed: for the roots of the equation are the ordinates of the curve at the points of intersection with the right line; and it is well known that a curve may be cut by a right line in as many points as its dimensions amount to. Thus, then,



then, a simple equation will be constructed by the intersection of one right line with another: a quadratic equation, or an affected equation of the 2d rank, by the intersections of a right line with a circle, or any of the conic sections, which are all lines of the 2d order; and which may be cut, by the right line, in two points, thereby giving the two roots of the quadratic equation. A cubic equation may be constructed by the intersection of the right line with a line of the 3d order: and so on.


But if, instead of the right line, some other line of a higher order be used; then the 2d line, whose intersections with the former are to determine the roots of the equation, may be taken as many dimensions lower, as the former is taken higher. And, in general, an equation of any height will be constructed by the intersections of two lines whose dimensions, multiplied together, produce the dimension of the given equation. Thus, the intersections of a circle with the conic sections, or of these with each other, will construct the biquadratic equations, or those of the 4th power, because  $2 \times 2 = 4$ ; and the intersections of the circle or conic sections with a line of the 3d order, will construct the equations of the 5th and 6th power; and so on.—For example,

*To construct a Simple Equation.* This is done by resolving the given simple equation into a proportion, or finding a third or 4th proportional, &c. Thus, 1. If the equation be  $ax = bc$ ; then  $a : b :: c : x = \frac{bc}{a}$ , the fourth proportional to  $a, b, c$ .

2. If  $ax = b^2$ ; then  $a : b :: b : x = \frac{b^2}{a}$ , a third proportional to  $a$  and  $b$ .


3. If  $ax = b^2 - c^2$ ; then, since  $b^2 - c^2 = \overline{b+c} \times \overline{b-c}$ , it will be  $a : b+c :: b-c : x = \frac{b+c \times b-c}{a}$ , a fourth proportional to  $a, b+c$  and  $b-c$ .

4. If  $ax = b^2 + c^2$ ; then construct the right-angled triangle ABC, whose base is  $b$ , and perpendicular is  $c$ , so shall the square of the hypotenuse be  $b^2 + c^2$ , which call  $b^2$ ; then the equation is  $ax = b^2$ , and  $x = \frac{b^2}{a}$  a third proportional to  $a$  and  $b$ .



*To construct a Quadratic Equation.*


1. If it be a simple quadratic, it may be reduced to this form  $x^2 = ab$ ; and hence  $a : x :: x : b$ , or  $x = \sqrt{ab}$  a mean proportional between  $a$  and  $b$ . Therefore upon a straight line take  $AB = a$ , and  $BC = b$ ; then upon the diameter AC describe a semi-circle, and raise the perpendicular BD to meet it in D; so shall BD be  $= x$  the mean proportional sought between AB and BC, or between  $a$  and  $b$ .



2. If the quadratic be affected, let it first be  $x^2 + 2ax = b^2$ ; then form the right-angled triangle whose base AB is  $a$ , and perpendicular BC is  $b$ ; and with the centre A and radius AC describe the semi-

circle DCE; so shall DB and BE be the two roots of the given quadratic equation  $x^2 + 2ax = b^2$ .

3. If the quadratic be  $x^2 - 2ax = b^2$ , then the construction will be the very same as of the preceding one  $x^2 + 2ax = b^2$ .



4. But if the form be  $2ax - x^2 = b^2$ : form a right-angled triangle whose hypotenuse FG is  $a$ , and perpendicular GH is  $b$ ; then with the radius FG and centre F describe a semi-circle IGK; so shall IH and HK be the two roots of the given equation  $2ax - x^2 = b^2$ , or  $x^2 - 2ax = -b^2$ . See Maclaurin's Algebra, part 3, cap. 2, and Simpson's Algebra, pa. 267.

*To construct Cubic and Biquadratic Equations.*—These are constructed by the intersections of two conic sections; for the equation will rise to 4 dimensions, by which are determined the ordinates from the 4 points in which these conic sections may cut one another; and the conic sections may be assumed in such a manner, as to make this equation coincide with any proposed biquadratic: so that the ordinates from these 4 intersections will be equal to the roots of the proposed biquadratic. When one of the intersections of the conic section falls upon the axis, then one of the ordinates vanishes, and the equation, by which these ordinates are determined, will then be of 3 dimensions only, or a cubic; to which any proposed cubic equation may be accommodated. So that the three remaining ordinates will be the roots of that proposed cubic. The conic sections for this purpose should be such as are most easily described; the circle may be one, and the parabola is usually assumed for the other.

Vieta, in his *Canonica Recensione Effectuum Geometricarum*, and Ghetaldus, in his *Opus Posthumum de Resolutione & Compositione Mathematica*, as also Des Cartes, in his *Geometria*, have shewn how to construct simple and quadratic Equations. Des Cartes has also shewn how to construct cubic and biquadratic equations, by the intersection of a circle and a parabola: And the same has been done more generally by Baker in his *Clavis Geometrica*, or *Geometrical Key*. But the genuine foundation of all these constructions was first laid and explained by Slusius in his *Mesolabium*, part 2. This doctrine is also pretty well handled by De la Hire, in a small treatise, called *La Construction des Equations Analytiques*, annexed to his *Conic Sections*. Newton, at the end of his *Algebra*, has given the construction of cubic and biquadratic equations mechanically; as also by the conchoid and cissoid, as well as the conic sections. See also Dr. Halley's *Construction of Cubic and Biquadratic Equations*; Colson's, in the *Philos. Trans.*; the Marquis de l'Hospital's *Traité Analytique des Sections Coniques*; Maclaurin's *Algebra*, part 3, c. 3 &c.


CONTACT, the relative state of two things that touch each other, but without cutting or entering; or whose surfaces join to each other without any interstice.



The contact of curve lines or surfaces, with either straight or curved ones, is only in points; and yet these points have different proportions to one another, as is shewn by Mr. Robartes, in the *Philos. Trans.* vol. 27 pa. 470; or *Abr.* vol. 4. pa. 1.

Because few or no surfaces are capable of touching in all points, and the cohesion of bodies is in proportion to their contact, those bodies will adhere fastest together, that are capable of the greatest contact.

*Angle of CONTACT* is the opening between a curve line and a tangent to it, particularly the circle and its tangent; as the angle formed at A between BA and AC, at the point of contact A.



It is demonstrated by Euclid, that the line CA standing perpendicular to the radius DA, touches the circle only in one point: and that no other right line can be drawn between the tangent and the circle.

Hence, the angle of contact is less than any rectilinear angle; and the angle of the semi-circle between the radius DA and the arch AB, is greater than any rectilinear acute angle.

This seeming paradox of Euclid has exercised the wits of mathematicians: it was the subject of a long controversy between Peletarius and Clavius; the former of whom maintained that the angle of contact is heterogeneous to a rectilinear one; as a line is to a surface; the latter maintained the contrary.

Dr. Wallis has a formal treatise on the angle of contact, and of the semi-circle; where, with other great mathematicians, he approves of the opinion of Peletarius.

**CONTENT**, a term often used for the measurement of bodies and surfaces, whether solid or superficial; or the capacity of a vessel and the area of a space; being the quantity either of matter or space included within certain bounds or limits.

**CONTIGUITY**, the relation of bodies touching one another.

**CONTIGUOUS**, a relative term, understood of things disposed so near each other, that they join their surfaces, or touch.

**CONTIGUOUS Angles**, are such as have one leg or side common to each angle; and are otherwise called *adjoining angles*; in contradistinction to those made by continuing their legs through the point of contact, which are called *opposite* or *vertical angles*.

**CONTINENT**, a terra firma, main-land, or a large extent of country, not interrupted by seas: so called, in opposition to island, peninsula, &c.

The world is usually divided into two grand continents, the old and the new: the old continent comprehends Europe, Asia, and Africa; the new, North and South America. Since the discovery of New Holland and New South Wales, it is a doubt with many whether to call that vast country an island or a continent.

**CONTINGENT Line**, the same with tangent line in Dialling, being the intersection of the planes of the dial and equinoctial, and at right angles to the substyle line.

**CONTINUAL PROPORTIONALS**, are a series of three or more quantities compared together, so that the ratio is the same between every two adjacent terms, viz between the 1st and 2d, the 2d and 3d, the 3d and 4th, &c. As 1, 2, 4, 8, 16, &c, where the terms continually increase in a double ratio; or 12, 4,  $\frac{2}{3}$ ,  $\frac{1}{3}$ , where the terms decrease in a triple ratio.

A series of continual or continued proportionals, is otherwise called a *progression*.

**CONTINUED Quantity**, or *Body*, is that whose parts are joined and united together.

**CONTINUED Proportion**, is that in which the consequent of the first ratio is the same with the antecedent of the second; as in these,  $3 : 6 :: 6 : 12$ . See *CONTINUAL Proportion*.

On the contrary, if the consequent of the first ratio be different from the antecedent of the second, the proportion is called *discrete*: as  $3 : 6 :: 4 : 8$ .

**CONTRA-HARMONICAL Proportion**, that relation of three terms, in which the difference of the first and second is to the difference of the 2d and 3d, as the 3d is to the first. Thus, for instance, 3, 5, and 6, are numbers contra-harmonically proportional; for  $2 : 1 :: 6 : 3$ .

**CONTRA-MURE**, in Fortification, is a little wall built before another partition wall, to strengthen it, so that it may receive no damage from the adjacent buildings.

**CONTRATE-WHEEL**, is that wheel in watches which is next to the crown, whose teeth and hoop lie contrary to those of the other wheels; from whence comes its name.

**CONTRAVALLATION**, *Line of*, in Fortification, is a trench, guarded with a parapet; being made by the besiegers, between them and the place besieged, to secure themselves on that side, and stop the sallies of the garrison. It is made without musket-shot of the town; sometimes going quite around it, and sometimes not, as occasion may require. The besiegers lie between the lines of circumvallation and contravallation: but it is now seldom used.

**CONVERGING Curves**. See *CURVE*.

**CONVERGING**, or **CONVERGENT Lines**, in Geometry, are those that continually approximate, or whose distance becomes continually less and less the farther they are continued, till they meet: in opposition to *divergent* lines, whose distance becomes continually greater.

Lines that converge the one way, diverge the other.

**CONVERGING Rays**, in Optics, are such as incline towards one another in their passage, and in Dioptrics, are those rays which, in their passage out of one medium into another of a different density, are refracted towards one another; so that, if far enough continued, they will meet in a point or focus.

**CONVERGING Series**, a series of terms or quantities, that always decrease the farther they proceed, or which tend to a certain magnitude or limit: in opposition to *diverging* series, or such as become larger and larger continually. See *SERIES*.

**CONVERSE**. A proposition is said to be the converse of another, when, after drawing a conclusion from something first supposed, we return again, and, making a supposition of what had before been concluded, draw from thence as a conclusion what before was made the supposition.



supposition. Thus, when it is supposed that the two sides of a triangle are equal, and thence demonstrate or conclude that the two angles opposite to those sides are equal also; then the converse is to suppose that the two angles of a triangle are equal, and thence to prove or conclude that the sides opposite to those angles are also equal.

CONVERSE *Direction*, in Astrology, is used in opposition to *direct* direction; that is, by the latter the promoter is carried to the significator, according to the order of the signs: whereas by the other it is carried from east to west, contrary to the order of the signs.

CONVERSION, or CONVERTENDO, is when there are four proportionals, and it is inferred, that the first is to its excess above the 2d, as the third to its excess above the 4th: according to Euclid, lib. 5, def. 17. Thus, if it be  $8 : 6 :: 4 : 3$ , then convertendo, or by conversion,  $8 : 2 :: 4 : 1$ . Or if there be  $a : b :: c : d$ , then convertendo, or by conversion,  $a : a - b :: c : c - d$ .

CONVEX, round or curved and protuberant outwards, as the outside of a globular body.

CONVEX *Lens*, *Mirror*, &c. See LENS, MIRROR, &c.

CONVEXITY, the exterior or outward surface of a convex or round body.

COPERNICAN, something relating to Copernicus. As, the

COPERNICAN *Sphere*. See SPHERE.

COPERNICAN *System*, is that system of the world, in which it is supposed that the sun is at rest in the centre, and the earth and planets all moving around him in their own orbits.

Here it is supposed, that the heavens and stars are at rest; and the diurnal motion which they appear to have, from east to west, is imputed to the earth's diurnal motion from west to east.

This system was maintained by many of the ancients; particularly Ecphantus, Seleucus, Aristarchus, Philolaus, Cleanthes Samius, Nicetas, Heraclides Ponticus, Plato, and Pythagoras; from the last of whom it was anciently called the Pythagoric, or Pythagorean System.

This system was also held by Archimedes, in his book of the number of the Grains of Sand; but after him it became neglected, and even forgotten, for many ages; till about 300 years since, when Copernicus revived it; from whom it took the new name of the Copernican System. See the next article.

COPERNICUS (NICHOLAS), an eminent astronomer, was born at Thorn in Prussia, January 19, 1473. He was instructed in the Latin and Greek languages at home; and afterward sent to Cracow, where he studied philosophy, mathematics, and medicine: though his genius was naturally turned to mathematics, which he chiefly studied, and pursued through all its various branches.

He set out for Italy at 23 years of age; stopping at Bologna, that he might converse with the celebrated astronomer of that place, Dominic Maria, whom he assisted for some time in making his observations. From hence he passed to Rome, where he was presently considered as not inferior to the famous Regiomontanus. Here he soon acquired so great a reputation, that he

was chosen professor of mathematics, which he taught there for a long time with the greatest applause; and here also he made some astronomical observations about the year 1500.

Afterward, returning to his own country, he began to apply his fund of observations and mathematical knowledge, to correcting the system of astronomy which then prevailed. He set about collecting all the books that had been written by philosophers and astronomers, and to examine all the various hypotheses they had invented for the solution of the celestial phenomena; to try if a more symmetrical order and constitution of the parts of the world could not be discovered, and a more just and exquisite harmony in its motions established, than what the astronomers of those times so easily admitted. But of all their hypotheses, none pleased him so well as the Pythagorean, which made the sun to be the centre of the system, and supposed the earth to move both round the sun, and also round its own axis. He thought he discerned much beautiful order and proportion in this; and that all the embarrassment and perplexity, from epicycles and excentrics, which attended the Ptolemaic hypotheses, would here be entirely removed.

This system he began to consider, and to write upon, when he was about 35 years of age. He carefully contemplated the phenomena; made mathematical calculations; examined the observations of the ancients, and made new ones of his own; till, after more than 20 years chiefly spent in this manner, he brought his scheme to perfection, establishing that system of the world which goes by his name, and is now universally received by all philosophers.

This system however was at first looked upon as a most dangerous heresy, and his work had long been finished and perfected, before he could be prevailed upon to give it to the world, being strongly urged to it by his friends. At length yielding to their intreaties, it was printed, and he had but just received a perfect copy, when he died the 24th of May 1543, at 70 years of age; by which it is probable he was happily relieved from the violent fanatical persecutions of the church, which were but too likely to follow the publication of his astronomical opinions; and which indeed was afterward the fate of Galileo, for adopting and defending them.

The above work of Copernicus, first printed at Norimberg in folio, 1543, and of which there have been other editions since, is intitled *De Revolutionibus Orbium Caelestium*, being a large body of astronomy, in 6 books.

When Rheticus, the disciple of our author, returned out of Prussia, he brought with him a tract of Copernicus, on plane and spherical trigonometry, which he had printed at Norimberg, and which contained a table of sines. It was afterward printed at the end of the first book of the *Revolutions*. An edition of our author's great work was also published in 4to at Amsterdam in 1617, under the title of *Astronomia Instaurata*, illustrated with notes by Nicolas Muler of Groningen.

COPERNICUS, the name of an astronomical instrument, invented by Whiston, to shew the motion and phenomena of the planets, both primary and secondary. It is founded upon the Copernican system, and therefore called by his name.



**COR CAROLI**, Charles's Heart, an extra-constellated star of the 2d magnitude in the northern hemisphere, between the Coma Berenices and Urfa Major; so called by Sir Charles Scarborough, in honour of king Charles I.

**COR HYDRÆ**, the Hydra's Heart, a star of the 2d magnitude, in the Heart of the constellation Hydra.

**COR LEONIS**, Lion's Heart, or Regulus, a star of the first magnitude in the constellation Leo.

**COR SCORPII**. See **ANTARES**.

**CORBEILS**, in Fortification, are little baskets about a foot and a half high, 8 inches broad at the bottom, and 12 at the top; which being filled with earth, are set against one another on the parapet, or elsewhere, leaving certain port-holes, from whence to fire under cover upon the enemy.

**CORBEL**, in Architecture, the representation of a basket, sometimes seen on the heads of caryatides.

**CORBEL**, or **CORBIL**, is also used, in Building, for a short piece of timber placed in a wall, with its end projecting out 6 or 8 inches, as occasion serves, in the manner of a shouldering-piece.

**CORBET**, the same as **CORBEL**.

**CORDON**, in Fortification, a row of stones jutting out between the rampart and the basis of the parapet, like the tore of a column. The cordon ranges round the whole fortrefs, and serves to join the rampart, which is aslope, and the parapet, which is perpendicular, more agreeably together.

In fortifications raised of earth, this space is filled up with pointed stakes, instead of a cordon.

**CORDS**, in Music, are the sounds produced by an instrument or the voice.

**CORIDOR**, or **CORRIDOR**, in Fortification, is the covert-way lying round about the whole compass of the works of a place, between the outside of the moat and the pallisadoes, being about 20 yards broad.

**CORIDOR** is also used, in Architecture, for a gallery, or long aisle, around a building, leading to several chambers at a distance from each other, sometimes wholly inclosed, and sometimes open on one side.

**CORINTHIAN Order**, of Architecture, is the 4th in order, or the 5th and last according to Scamozzi and Le Clerc.

This order was invented by an Athenian Architect, and is the richest and most delicate of them all; its capital being adorned with rows of leaves, and of 8 volutas, which support the abacus. The height of its column is 10 diameters, and its cornice is supported by modillions.

**CORNEA Tunica**, the second coat of the eye; so called from its substance resembling the horn of a lantern. This is situated in the fore-part; and is surrounded by the sclerotica. It has a greater convexity than the rest of the eye, and is a portion of a small sphere, or rather spheroid, and consolidates the whole eye.

**CORNICE**, **CORNICHE**, or **CORNISH**, the third and uppermost part of the entablature of a column, or the uppermost ornament of any wainscotting, &c.

**COROLLARY**, or **CONSECTARY**, a consequence drawn from some proposition or principles already advanced or demonstrated, and without the aid of any other proposition: as if from this theorem, *That a triangle which has two equal sides, has also two equal*

*angles, this consequence should be drawn, that a triangle which hath the three sides equal, has also its three angles equal.*

**CORONA**, *Crown* or *Crowning*, in Architecture, the flat and most advanced part of the cornice; so called, because it crowns the cornice and entablature: by the workmen it is called the *drip*, as serving by its projection to screen the rest of the building from the rain.

**CORONA**, in Optics, a luminous circle, usually coloured, round the sun, moon, or largest planets. See **HALO**.

**CORONA Borealis**, or *Septentrionalis*, the *Northern Crown* or *Garland*, a constellation of the northern hemisphere, being one of the 48 old ones. It contains 8 stars according to the catalogue of Ptolemy, Tycho, and Hevelius; but according to the Britannic Catalogue, 21.

**CORONA Australis**, or *Meridionalis*, the *Southern Crown*, a constellation of the southern hemisphere, whose stars in Ptolemy's catalogue are 13; in the British Catalogue, 12.

**CORPUSCLE** the diminutive of corpus, used to express the minute parts, or particles, that constitute natural bodies; meaning much the same as *atoms*.

Newton shews a method of determining the sizes of the corpuscles of bodies, from their colours.

**CORPUSCULAR Philosophy**, that scheme or system of physics, in which the phenomena of bodies are accounted for, from the motion, rest, position, &c, of the corpuscles or atoms of which bodies consist.

The Corpuscular philosophy, which now flourishes under the name of the mechanical philosophy, is very ancient. Leucippus and Democritus taught it in Greece; from them Epicurus received it, and improved it; and from him it was called the *Epicurean Philosophy*.

Leucippus, it is said, received it from one Mochus, a Phenician philosopher, before the time of the Trojan war, and the first who philosophized about atoms: which Mochus is, according to the opinion of some, the Moses of the Scriptures.

After Epicurus, the corpuscular philosophy gave way to the peripatetic, which became the popular system. Thus, instead of atoms, were introduced specific and substantial forms, qualities, sympathies, &c, which amused the world, till Gassendus, Charleton, Descartes, Boyle, Newton, and others, retrieved the corpuscularian hypotheses; which is now become the basis of the mechanical and experimental philosophy.

Boyle reduces the principles of the corpuscular philosophy to the 4 following heads.

1. That there is but one universal kind of matter, which is an extended, impenetrable, and divisible substance, common to all bodies, and capable of all forms.—On this head, Newton finely remarks thus: "All things considered, it appears probable to me, that God in the beginning created matter in solid, hard, impenetrable, moveable particles; of such sizes and figures, and with such other properties, as most conduced to the end for which he formed them: and that these primitive particles, being solids, are incomparably harder than any of the sensible porous bodies compounded of them; even so hard as never to wear, or break in pieces: no other power being able to divide what God made one in the first creation. While these corpuscles remain entire, they may compose bodies of one



and the same nature and texture in all ages: but should they wear away, or break in pieces, the nature of things depending on them would be changed: water and earth, composed of old worn particles, of fragments of particles, would not be of the same nature and texture now, with water and earth composed of entire particles at the beginning. And therefore, that nature may be lasting, the changes of corporeal things are to be placed only in the various separations, and new associations, of these permanent corpuscles."

2. That this matter, in order to form the vast variety of natural bodies, must have motion in some, or all its assignable parts; and that this motion was given to matter by God, the creator of all things; and has all manner of directions and tendencies.—“These corpuscles, says Newton, have not only a vis inertiae, accompanied with such passive laws of motion as naturally result from that force; but also are moved by certain active principles; such as that of gravity, and that which causes fermentation, and the cohesion of bodies."

3. That matter must also be actually divided into parts; and each of these primitive particles, fragments, or atoms of matter, must have its proper magnitude, figure, and shape.

4. That these differently sized and shaped particles, have different orders, positions, situations, and postures, from whence all the variety of compound bodies arises.

CORRIDOR. See CORIDOR.

CORVUS, the *Raven*, a southern constellation, fabled by the Greeks, as taken up to heaven by Apollo, to whom it tattled that the beautiful maid Coronis, the daughter of Phlegos, and mother of Esculapius by Apollo, played the deity false with Ischys, under a tree upon which the animal happened to be perched.

The stars in this constellation, in Ptolemy's and Tycho's catalogues are 7; but in the Britannic catalogue, 9.

COSECANT, COSINE, COTANGENT, COVERSED SINE, are the secant, sine, tangent, and versed sine of the complement of an arch or angle; *Co* being, in this case, a contraction of the word complement, and was first introduced by Gunter.

COSMICAL ASPECT, among astrologers, is the aspect of a planet with respect to the earth.

COSMICAL RISING, or *Setting*, is said of a star when it rises or sets at the same time when the sun rises.

But, according to Kepler, to rise or set cosmically, is only simply to rise or set, that is, to ascend above, or descend below, the horizon; as much as to say, to rise or set to the world.

COSMOGONY, the science of the formation of the universe; as distinguished from cosmography, which is the science of the parts of the universe, supposing it formed, and in the state as we behold it; and from cosmology, which reasons on the actual and permanent state of the world as it now is; whereas cosmogony reasons on the variable state of the world at the time of its formation.

COSMOGRAPHY, the description of the world; or the art that teaches the construction, figure, disposition, and relation of all the parts of the world, with

the manner of representing them on a plane. It consists chiefly of two parts; viz, *Astronomy*, which shews the structure of the heavens, with the disposition of the stars; and *Geography*, which shews those of the earth.

COSMOLOGY, the science of the world in general.

COSS, *Rule of*, meant the same as Algebra, by which name it was for some time called, when first introduced into Europe through the Italians, who named it *Regola de Cosa*, the *Rule of the thing*; the unknown quantity, or that which was required in any question, being called *cosa*, the *thing*; from whence we have Coss, and Cossic numbers, &c.

COTES (ROGER), a very eminent mathematician, philosopher, and astronomer, was born July 10, 1682, at Burbach in Leicestershire, where his father Robert was rector. He was first placed at Leicester school; where, at 12 years of age, he discovered a strong inclination to the mathematics. This being observed by his uncle, the Rev. Mr. John Smith, he gave him all the encouragement he could; and prevailed on his father to send him for some time to his house in Lincolnshire, that he might assist him in those studies: and here he laid the foundation of that deep and extensive knowledge in that science, for which he was afterward so deservedly famous. He was hence removed to St. Paul's school, London, where he made a great progress in classical learning; and yet he found so much leisure as to support a constant correspondence with his uncle, not only in mathematics, but also in metaphysics, philosophy, and divinity. His next remove was to Trinity College Cambridge, where he took his degrees, and became fellow.

Jan. 1706, he was appointed professor of astronomy and experimental philosophy, upon the foundation of Dr Thomas Plume, archdeacon of Rochester; being the first that enjoyed that office, to which he was unanimously chosen, on account of his high reputation and merits. He entered into orders in 1713; and the same year, at the desire of Dr. Bentley, he published at Cambridge the second edition of Newton's *Mathematica Principia*; inserting all the improvements which the author had made to that time. To this edition he prefixed a most admirable preface, in which he pointed out the true method of philosophizing, shewing the foundation on which the Newtonian philosophy was raised, and refuting the objections of the Cartesians and all other philosophers against it.

The publication of this edition of Newton's *Principia* added greatly to his reputation; nor was the high opinion the public now conceived of him in the least diminished, but rather much increased, by several productions of his own, which afterward appeared. He gave in the *Philos. Transactions*, two papers, viz, 1, *Logometria*, in vol. 29; and a Description of the great fiery meteor that was seen March 6, 1716, in vol. 31.

This extraordinary genius in the mathematics died, to the great regret of the university, and all the lovers of the sciences, June 5, 1716, in the very prime of his life, being not quite 34 years of age.

Mr. Cotes left behind him some very ingenious, and indeed admirable tracts, part of which, with the *Logometria*:




gometria above mentioned, were published, in 1722, by Dr. Robert Smith, his cousin and successor in his professorship, afterward master of Trinity College, under the title of *Harmonia Mensurarum*, which contains a number of very ingenious and learned works: see the Introduction to my Logarithms. He wrote also a *Compendium of Arithmetic*; of the *Resolution of Equations*; of *Dioptrics*; and of the *Nature of Curves*. Beside these pieces, he drew up, in the time of his lectures, a course of *Hydrostatical and Pneumatical Lectures*, in English, which were published also by Dr. Smith in 8vo, 1737, and are held in great estimation.

So high an opinion had Sir Isaac Newton of our author's genius, that he used to say, "If Cotes had lived, we had known something."

**COTESIAN theorem**, in Geometry, an appellation used for an elegant property of the circle discovered by Mr. Cotes. The theorem is this:

If the factors of the binomial  $a^c + x^c$  be required, the index  $c$  being an integer number. With the centre  $O$ , and radius  $AO = a$ , describe a circle, and di-



vide its circumference into as many equal parts as there are units in  $2c$ , at the points  $A, B, C, D, \&c$ ; then in the radius, produced if necessary, take  $OP = x$ , and from the point  $P$ , to all the points of division in the circumference, draw the lines  $PA, PB, PC, \&c$ ; so shall these lines taken alternately be the factors sought; viz,

$PB \times PD \times PF \&c = a^c + x^c$ ,  
and  $PA \times PC \times PE \&c = a^c - x^c$ , viz,  $a^c - x^c$  or  $x^c - a^c$ , according as the point  $P$  is within or without the circle.

For instance, if  $c = 5$ , divide the circumference into 10 equal parts, and the point  $P$  being within the circle, then will  $OA^5 + OP^5 = BP \times DP \times FP \times HP \times KP$ , and  $OA^5 - OP^5 = AP \times CP \times EP \times GP \times IP$ .

In like manner, if  $c = 6$ , having divided the circumference into 12 equal parts, then will  $OA^6 + OP^6 = BP \times DP \times FP \times HP \times KP \times MP$ ,  $OA^6 - OP^6 = AP \times CP \times EP \times GP \times IP \times LP$ .

The demonstration of this theorem may be seen in Dr. Pemberton's *Epist. de Cotesii inventis*. See also Dr. Smith's *Theoremata Logometrica* and *Trigonometrica*, added to Cotes's *Harm. Mens.* pa. 114; De Moivre *Miscel. Analyt.* pa. 17; and Waring's *Letter to Dr. Powell*, pa. 39.

By means of this theorem, the acute and elegant author was enabled to make a farther progress in the inverse method of Fluxions, than had been done before. But in the application of his discovery there

still remained a limitation, which was removed by Mr. De Moivre. Vide ut supra.

**COVERT-WAY**, in fortification, a space of ground level with the adjoining country, on the outer edge of the ditch, ranging quite round all the works. This is otherwise called the *corridor*, and has a parapet with its banquette and glacis, which form the height of the parapet. It is sometimes also called the counter-scarp, because it is on the edge of the scarp.

One of the greatest difficulties in a siege, is to make a lodgment on the covert-way; because it is usual for the besieged to palisade it along the middle, and undermine it on all sides.

**COVING**, in Building, is when houses are built projecting over the ground plot, and the turned projection formed into an arch.

**COVING Cornice**, is one that has a large casemate or hollow in it.

**COUNT-WHEEL**, is a wheel in the striking part of a clock, moving round once in 12 or 24 hours. It is sometimes called the *locking-wheel*, because it has usually 11 notches in it at unequal distances from one another, to make the clock strike.

**COUNTER-APPROACHES**, in Fortification, lines or trenches made by the besieged, where they come out to attack the lines of the besiegers in form.

**COUNTER-BATTERY**, a battery raised to play on another, to dismount the guns, &c.

**COUNTER-BREAST-WORK**, the same as *Fausse-Braye*.

**COUNTER-FORTS**, *Buttresses*, or *Spurs*, are pillars of masonry serving to prop or sustain walls, or terraces, subject to bulge, or be thrown down.

**COUNTER-FUGUE**, in Music, is when fugues proceed contrary to one another.

**COUNTER-GUARD**, in Fortification, a work commonly serving to cover a bastion. It is composed of two faces, forming a salient angle before the flanked angle of a bastion.

**COUNTER-HARMONICAL**. See **CONTRA-HARMONICAL**.

**COUNTER-MINE**, a subterraneous passage, made by the besieged, in search of the enemy's mine, to give air to it, to take away the powder; or by any other means to frustrate the effect of it.

**COUNTER-PART**, a term in Music, only denoting that one part is opposite to another, so, the bass and treble are counterparts to each other.

**COUNTER-POINT**, in Music, the art of composing harmony; or disposing and concerting several parts so together, as that they may make an agreeable whole.

**COUNTER-POISE**, any thing serving to weigh against another; particularly a piece of metal, usually of brass or iron, making an appendage to the Roman *statera*, or steel-yard. It is contrived to slide along the beam; and from the division at which it keeps the balance in equilibrio, the weight of the body is determined. It is sometimes called the *pear*, on account of its figure; and *mass*, by reason of its weight.

Rope-dancers make use of a pole by way of counterpoise, to keep their bodies in equilibrio.

**COUNTERSCARP**, is that side of the ditch that is next the country; or properly the talus that supports the earth of the covert-way: though by this word is often understood the whole covert-way, with its



its parapet and glacis. And so it must be understood when it is said, The enemy lodged themselves on the counterfearp.

COUNTER-SWALLOWS-TAIL, is an outwork in Fortification, in form of a single tenaille, wider towards the place, or at the gorge, than at the head, or next the country.

COUNTER-TENOR, one of the mean or middle parts of music: so called, as being opposite to the tenor.

COURSE, in Navigation, the point of the compass, or horizon, which a ship steers on; or the angle which the rhumb line on which it sails makes with the meridian; being sometimes reckoned in degrees, and sometimes in points of the compass.

When a ship sails either due north or south, she sails on a meridian, makes no departure, and her distance and difference of latitude are the same.

When she sails due east or west, her course makes right-angles with the meridian, and she sails either upon the equator, or a parallel to it; in which case she makes no difference of latitude, but her distance and departure are the same.

But when the ship sails between the cardinal points, on a course making always the same oblique angle with the meridians, her path is then the loxodromic curve, being a spiral cutting all the meridians in the same angle, and terminating in the pole.

COURTAIN. See CURTIN.

CRAB, in Mechanics, an engine used for mounting guns on their carriages. See GIN.

CRAB, on ship-board, is a wooden pillar, whose lower end is let down through the ship's decks, and rests upon a socket like the capstan: in its upper end are three or four holes, at different heights, through the middle of it, above one another, to receive long bars, against which men act by pushing or thrusting.—It is employed to wind in the cable, and for other purposes requiring a great mechanical power.

The Crab with three claws is used to launch ships, and to heave them into the dock, or off the key.

CRANE, a machine used in building, and in commerce, for raising large stones and other weights.

M. Perrault, in his notes on Vitruvius, makes the crane the same with the corvus, or raven, of the ancients.

The modern crane consists of several members or pieces, the principal of which is a strong upright beam, or arbor, firmly fixed in the ground, and sustained by eight arms, coming from the extremities of four pieces of wood laid across, through the middle of which the foot of the beam passes. About the middle of the arbor the arms meet, and are mortised into it: its top ends in an iron pivot, on which is borne a transverse piece, advancing out to a good distance like a crane's neck; whence the name. The middle and extremities of this are again sustained by arms from the middle of the arbor: and over it comes a rope, or cable, to one end of which the weight is fixed; the other is wound about the spindle of a wheel, which, turned, draws the rope, and that heaves up the weight; to be afterwards applied to any side or quarter, by the mobility of the transverse piece on the pivot.

Several improvements of this useful machine are mentioned in Defaguliers's Exper. Philos. pa. 178 & VOL. I.

seq. particularly how to prevent the inconveniences arising from sudden jerks, as well as to increase its force by using a double axis in peritrochio, and two handles.

The Crane is of two kinds; in the first kind, called the rat-tailed crane, the whole machine, with the load, turns upon a strong axis: in the second kind, the gibbet alone moves on its axis. See Defaguliers, as above, for a particular account of the different cranes, and of the gradual improvements they have received. See also the Supplement to Ferguson's Lectures, pa. 3, &c., or Philos. Transf. vol. 54, pa. 24, for a description of a new and safe crane, with four different powers adapted to different weights.

CRANE is the name of a southern constellation. See GRUS.

CRANE is also a popular name for a syphon.


CRANK, a contrivance in machines, in manner of an elbow, only of a square form; projecting out from an axis, or spindle; and serving, by its rotation, to raise and fall the pistons of engines for raising water, or the like.

CRATER, the Cup, a constellation in the southern hemisphere; whose stars, in Ptolemy's catalogue, are 7; in Tycho's, 8; in Hevelius's, 10; and in the Britannic catalogue, 31.

CREEK, a part of a haven, where any thing is landed from the sea. It is also said to be a shore or bank on which the water beats, running in a small channel from any part of the sea.

CREPUSCULUM, *Twilight*; the time from the first dawn or appearance of the morning, to the rising of the sun; and again, between the setting of the sun, and the last remains of day.

The Crepusculum, or twilight, it is supposed, usually begins and ends when the sun is about 18 degrees below the horizon; for then the stars of the 6th magnitude disappear in the morning, and appear in the evening. It is of longer duration in the solstices than in the equinoxes, and longer in an oblique sphere, than in a right one; because, in those cases the sun, by the obliquity of his path, is longer in ascending through 18 degrees of altitude.



Twilight is occasioned by the sun's rays refracted in our atmosphere, and reflected from the particles of it to the eye. For let A be the place of an observer on the earth ADL, AB the sensible horizon, meeting in B the circle CBM bounding that part of the atmosphere which is capable of refracting and reflecting light to the eye. It is plain that when the sun is under this horizon,



horizon, no direct rays can come to the eye at A: but the sun being in the refracted line CG, the particle C will be illuminated by the direct rays of the sun; and that particle may reflect those rays to A, where they enter the eye of the spectator. And thus the sun's light illuminating an innumerable multitude of particles, may be all reflected to the spectator at A.—From B draw BD touching the circle ADL in D; and let the sun be in the line BD at S: Then the ray SB will be reflected into BA, and will enter the eye, because the angle of incidence DBE is equal to the angle of reflection ABE: And that will be the first ray that reaches the eye in the morning, when the dawning begins; or the last that falls upon the eye at night, when the twilight ends: for when the sun goes lower down, the particles at B can be no longer illuminated.

Kepler indeed assigns another cause of the crepusculum, viz, the luminous matter or atmosphere about the sun; which, arising near the horizon, in a circular figure, exhibits the crepusculum; in no wise, he thinks, owing to the refraction of the atmosphere.—The sun's luminous atmosphere indeed, though neither the sole nor principal cause of twilight, may lengthen its duration, by illuminating our air, when the sun is too low to reach it with his own light. Gregor. Astr. lib. 2, prop. 8.

*The depth of the sun below the horizon, at the beginning of the morning, or end of the evening twilight, is determined in the same manner as the arch of vision; viz, by observing the moment when the air first begins to shine in the morning, or ceases to shine in the evening; then finding the sun's place for that moment, and thence the time till his rising in the horizon, or from his setting in it in the evening. It is now generally agreed that this depth is about 18 degrees upon an average.—Alhazen found it to be 19°; Tycho, 17°; Rothmann, 24°; Stevenius, 18°; Cassini, 15°; Riccioli, in the equinox in the morning 16°, in the evening 20° 30'; in the summer solstice in the morning 21° 25', in the winter solstice in the morning 17° 25'.*

Nor is this difference among the determinations of astronomers to be wondered at; the cause of the crepusculum being inconstant: for, if the exhalations in the atmosphere be either more copious, or higher, than ordinary; the morning twilight will begin sooner, and the evening hold longer than ordinary: for the more copious the exhalations are, the more rays will they reflect, consequently the more will they shine; and the higher they are, the sooner will they be illuminated by the sun. On this account too, the evening twilight is longer than the morning, at the same time of the year in the same place. To this it may be added, that in a denser air, the refraction is greater; and that not only the brightness of the atmosphere is variable, but also its height from the earth: and therefore the twilight is longer in hot weather than in cold, in summer than in winter, and also in hot countries than in cold, other circumstances being the same. But the chief differences are owing to the different situations of places upon the earth, or to the difference of the sun's place in the heavens. Thus, the twilight is longest in a parallel sphere, and shortest in a right sphere, and longest to places in an oblique sphere in proportion as they

are nearer to one of the poles; a circumstance which affords relief to the inhabitants of the more northern countries, in their long winter nights. And the twilights are longest in all places of north latitude, when the sun is in the tropic of cancer; and to those in south latitude, when he is in the tropic of capricorn. The time of the shortest twilight is also different in different latitudes; in England, it is about the beginning of October and of March, when the sun is in the signs ♋ and ♏. For the method of determining it by trigonometry, see Gregor. Astron. lib. 2, prob. 41. See also Robertson's Navigation, book 5, prob. 12.—Hence, when the difference between the sun's declination and the depth of the equator is less than 18°, so that the sun does not descend more than 18° below the horizon; the crepusculum will continue the whole night, as is the case in England from about the 22d of May to the 22d of July.

*Given the latitude of the place, and the sun's declination; to find the beginning of the morning, and end of the evening twilight.*—In the oblique-angled spherical triangle ZPS, are given ZP the colatitude, PS the codeclination, and  $ZS = 108^\circ$ , being the sum of  $90^\circ$  the quadrant and  $18^\circ$  the depression at the extremity of the twilight. Then, by spherical trigonometry, calculate the angle ZPS the hour-angle from noon; which changed into time, at the rate of  $15^\circ$  to the hour, gives the time from noon at the beginning or end of twilight. See Robertson, ubi supra.

Of the Height of the sensible Atmosphere, as determined from the duration of twilight, see Keil's Astron. Lect. lect. 20, pa. 235, ed. 1721; or Long's Astron. vol. 1, pa. 260; where it is determined that the height where the atmosphere is dense enough to reflect the rays of light, is about 42 miles.

CRESCENT, the new moon, which, as it begins to recede from the sun, shews a small rim of light, terminating in horns or points, which are still increasing, till it becomes full, and round in the opposition.

The term is sometimes also used for the same figure of the moon in her wane, or decrease, but improperly; both because the horns are then turned towards the west, and because the figure is on the decrease; the *crescent* properly signifying increase, from *cresco*, I grow.

CRONICAL. See ACRONICAL.

CRONOS, a name given to Saturn by some of the old astronomical writers.

CROSIER, four stars, in form of a cross; by the help of which, those that sail in the southern hemisphere find the antarctic pole.


CROSS, in Surveying, is an instrument consisting of a brass circle, divided into 4 equal parts, by two lines crossing each other in the centre. At each extremity of these lines is fixed a perpendicular sight, with small holes below each slit, for the better discovering of distant objects. The cross is mounted on a staff, or stand, to fix it in the ground, and is very useful for measuring small pieces of land, and taking offsets, &c.

*Ex.* Suppose it be required to survey the field ABCDE with the Cross. Measure along the diagonal line AC, and observe, with the Cross, when you are perpendicularly opposite to the corners, as at F, G, H, and from thence measure the perpendiculars FE, GB,

HD.



**HD.** When you think you are nearly opposite a corner, set up the cross, with one of the bars or cross lines in the direction AC; then look through the sights of the other cross bar for the corner, as B; if it be seen through them, the cross is fixed in the right place; if not, take it up and move it backward or forward in the line AC, till the point B be seen through those sights; and then you have the true place of the perpendicular.




*Invention of the Cross, Inventio Crucis*, an ancient feast, which is still retained in our calendar, and solemnized on the 3d of May, in memory of the finding of the true Cross of Christ, deep in the ground, on Mount Calvary, by St. Helena, the mother of Constantine; where she erected a church for the preservation of part of it: the rest being brought to Rome, and deposited in the church of the Holy Cross of Jerusalem.

*Exaltation of the Cross*, an ancient feast, held on the 14th of September, in memory of this, that Heraclitus restored to Mount Calvary the true cross, in 642, which had been carried off, 14 years before, by Cosroes king of Persia, upon his taking Jerusalem from the emperor Phocas. This feast is still retained in our calendar, on Sept. 14, under the denomination of *Holy Rood*, or *Holy Cross*.

**CROSS-MULTIPLICATION**, a method used chiefly by artificers in multiplying feet and inches by feet and inches, or the like; so called, because the factors are multiplied cross-wise, thus:

f.	i.	
9	10	
6	8	
<hr/>		
59	0	
6	6	8
<hr/>		
65	6	8

**CROSS-STAFF**, or *Fore-Staff*, is a mathematical instrument of box, or pear tree, consisting of a square staff, of about 3 feet long, having each of its faces divided like a line of tangents, and having 4 cross pieces of unequal lengths to fit on to the staff, the halves of these being as the radii to the tangent lines on the faces of the staff.—The instrument was used in taking the altitudes of the celestial bodies at sea.



**CROUSAZ** (JOHN PETER DE), a learned philosopher and mathematician, was born at Lausanne in Switzerland, April 13, 1663; where he died in 1748, at 85 years of age. Having made great progress in mathematics and the philosophy of Des Cartes, he travelled into Geneva, Holland, and France. He was successively professor in several universities; and at length was chosen governor to Prince Frederick of Hesse-Cassel, nephew to the king of Sweden.

Croufaz was author of many works, in various branches; belles-lettres, logic, philosophy, divinity, &c. &c; but the most esteemed of them are, 1. His *Logic*; the best edition of which is that of 1741, in 6 vols. 8vo.—2. *A Treatise on Beauty*.—3. *A Treatise on Education*, 2 vols, 12mo.—4. *A Treatise on the Human Understanding*.—5. *Several Treatises on Philosophy and Mathematics*; as a *Treatise on Motion*, &c. with several papers inserted in the *Memoirs of the French Academy of Sciences*.

**CROW**, in Mechanics, an iron lever, made with a sharp point at one end, and two claws at the other; being used in heaving and purchasing great weights, &c.

**CROWN**, in Astronomy, a name given to two constellations, the southern and the northern.

**CROWN**, in Geometry, a plane ring included between two parallel or concentric peripheries, of unequal circles.

The area of this is had, by multiplying its breadth by the length of a middle periphery, which is an arithmetical mean between the two peripheries that bound it; or by multiplying half the sum of the circumferences by half the difference of the diameters; or lastly by multiplying the sum of the diameters by the difference of the diameters, and this last product by .7854. See my *Mensuration*, pa. 148, 2d ed.

**CROWN-Post**, is a post in some buildings standing upright in the middle, between two principal rafters; and from which proceed struts or braces to the middle of each rafter. It is otherwise called a *king-post*, or *king's-piece*, or *joggle-piece*.

**CROWN-Wheel**, of a Watch, is the upper wheel next the balance, or that which drives the balance.

**CROWN-Work**, in Fortification, is an out-work running into the field; designed to keep off the enemy, gain some hill, or advantageous post, and cover the other works of the place. It consists of two demi-bastions at the extremities, and an entire bastion in the middle, with curtains.

**CROWNED Horn-work**, is a Horn-work with a crown-work before it.

**CRYSTALLINE Humour**, is a thick compact humour of the eye, in form of a flattish convex lens, placed in the middle of the eye, and serving to make that refraction of the rays of light which is necessary to have them meet in the retina, and form an image there, by which vision may be performed.

**CRYSTALLINE Heavens**, in the Old Astronomy, two orbs imagined between the primum mobile and the firmament, in the Ptolomaic system, which supposed the heavens solid, and only susceptible of a single motion.

King Alphonfus of Arragon, it is said, introduced the Crystallines, to explain what they called the *motion of trepidation*, or *titubation*.

The first Crystalline, according to Regiomontanus,  
X x 2 &c,



&c, serves to account for the slow motion of the fixed stars; by which they advance a degree in about 70 years, according to the order of the signs, or from west to east; which occasions a precession of the equinox. The 2d serves to account for the motion of libration, or trepidation; by which the celestial sphere librates from one pole towards the other, causing a difference in the sun's greatest declination.

**CUBATURE**, or **CUBATION**, of a solid, is the measuring the space contained in it, or finding the solid content of it, or finding a cube equal to it.



The cubature regards the content of a body, as the quadrature does the superficies or area of a figure.

**CUBE**, a regular or solid body, consisting of six equal sides or faces, which are squares.—A die is a small cube.

It is also called a *hexaedron*, because of its six sides, and is the 2d of the five Platonic or Regular bodies.

The cube is supposed to be generated by the motion of a square plane, along a line equal and perpendicular to one of its sides.

*To describe a Rete, or Net, for forming a cube, or with which it may be covered.*—Describe six squares as in the annexed figure, upon card paper, paste-board, or the like, of the size of the faces of the proposed cube; and cut it half through by the lines AB, CD, EF, AC, BD; then fold up the several squares till their edges meet, and so form the cube, or a covering over one, as in the figure annexed.



*To determine the Surface and Solidity of a Cube.*—Multiply one side by itself, which will give one square or face; then this multiplied by 6, the number of faces, will give the whole surface. Also multiply one side twice by itself, that is, cube it, and that will be the solid content.

*Duplication of a CUBE.* See **DUPLICATION**.

**CUBES**, or **CUBIC Numbers**, are formed by multiplying any numbers twice by themselves. So the cubes of 1, 2, 3, 4, 5, 6, &c, are 1, 8, 27, 64, 125, 216, &c.

The third differences of the cubes of the natural numbers are all equal to each other, being the constant number 6. For, let  $m^3$ ,  $n^3$ ,  $p^3$  be any three adjacent cubes in the natural series as above, that is, whose roots  $m$ ,  $n$ ,  $p$  have the common difference 1; then because

$n = m + 1$ , theref.  $n^3 = m^3 + 3m^2 + 3m + 1$ ,

$p = n + 1$ , theref.  $p^3 = n^3 + 3n^2 + 3n + 1$ ; so that

the difference between the 1st and 2d, and between the 2d and 3d cubes, are

$n^3 - m^3 = 3m^2 + 3m + 1$ ,  
 $p^3 - n^3 = 3n^2 + 3n + 1$ , } the 1st differences;

and the dif. of these differences, is

$3 \cdot n^2 - m^2 + 3 \cdot n - m = 3 \cdot n + m + 1 = 6 \cdot m + 1$ ,  
 the 2d difference.

In like manner the next 2d dif. is  $6 \cdot n + 1$ : hence the dif. of these two 2d difs. is  $6 \cdot n - m = 6$ , which is therefore the constant 3d difference of all the series of cubes. And hence that series of cubes will be formed by addition only, viz, adding always the 3d dif. 6 to find the column or series of 2d difs, then these added always for the 1st difs, and lastly these always added for the cubes themselves, as below:

3d Difs.	2d Difs.	1st Difs.	Cubes.
6	6	1	0
6	12	7	1
6	18	19	8
6	24	37	27
6	30	61	64
6	36	91	125
6	42	127	216
6	48	169	343

Peletarius, among various speculations concerning square and cubic numbers, shews that the continual sums of the cubic numbers, whose roots are 1, 2, 3, &c, form the series of squares whose roots are 1, 3, 6, 10, 15, 21, &c.

Thus,

$$1 = 1^2,$$

$$1 + 8 = 9 = 3^2,$$

$$1 + 8 + 27 = 36 = 6^2,$$

$$1 + 8 + 27 + 64 = 100 = 10^2, \text{ \&c.}$$

Or, in general,

$$1^3 + 2^3 + 3^3 + 4^3 \text{ \&c. to } n^3 = 1 + 2 + 3 + 4 + \dots + n = \frac{1}{2} n \cdot n + 1.$$

It is also a pretty property, that any number, and the cube of it, being divided by 6, leave the same remainder; the series of remainders being 0, 1, 2, 3, 4, 5, continually repeated. Or that the differences between the numbers and their cubes, divided by 6, leave always 0 remaining; and the quotients, with their successive differences, form the several orders of figurate numbers. Thus,

Num.	Cubes.	Difs.	Quot.	1 Dif.	2 Dif.
1	1	0	0	0	0
2	8	6	1	1	1
3	27	24	4	3	2
4	64	60	10	6	3
5	125	120	20	10	4
6	216	210	35	15	5
7	343	336	56	21	6

The following is a Table of the first 1000 cubic numbers.



TABLE OF CUBES.

Num.	Cubes	Num.	Cubes	Num.	Cubes	Num.	Cubes	Num.	Cubes	Num.	Cubes
1	1	60	216000	119	1685159	178	5639752	237	13312053	296	25934336
2	8	61	226981	120	1728000	179	5735339	238	13481272	297	26198073
3	27	62	238328	121	1771561	180	5832000	239	13651919	298	26463592
4	64	63	250047	122	1815848	181	5929741	240	13824000	299	26730899
5	125	64	262144	123	1860867	182	6028568	241	13997521	300	27000000
6	216	65	274625	124	1906624	183	6128487	242	14172488	301	27270901
7	343	66	287496	125	1953125	184	6229504	243	14348907	302	27543608
8	512	67	300763	126	2000376	185	6331625	244	14526784	303	27818127
9	729	68	314432	127	2048383	186	6434856	245	14706125	304	28094464
10	1000	69	328509	128	2097152	187	6539203	246	14886936	305	28372625
11	1331	70	343000	129	2146689	188	6644672	247	15069223	306	28652616
12	1728	71	357911	130	2197000	189	6751269	248	15252992	307	28934443
13	2197	72	373248	131	2248091	190	6859000	249	15438249	308	29218112
14	2744	73	389017	132	2299968	191	6967871	250	15625000	309	29503629
15	3375	74	405224	133	2352637	192	7077888	251	15813251	310	29791000
16	4096	75	421875	134	2406104	193	7189057	252	16003008	311	30080231
17	4913	76	438976	135	2460375	194	7301384	253	16194277	312	30371328
18	5832	77	456533	136	2515456	195	7414875	254	16387064	313	30664297
19	6859	78	474552	137	2571353	196	7529536	255	16581375	314	30959144
20	8000	79	493039	138	2628072	197	7645373	256	16777216	315	31255875
21	9261	80	512000	139	2685619	198	7762392	257	16974593	316	31554496
22	10648	81	531441	140	2744000	199	7880599	258	17173512	317	31855013
23	12167	82	551368	141	2803221	200	8000000	259	17373979	318	32157432
24	13824	83	571787	142	2863288	201	8120601	260	17576000	319	32461759
25	15625	84	592704	143	2924207	202	8242408	261	17779581	320	32768000
26	17576	85	614125	144	2985984	203	8365427	262	17984728	321	33076161
27	19683	86	636056	145	3048625	204	8489664	263	18191447	322	33386248
28	21952	87	658503	146	3112136	205	8615125	264	18399744	323	33698267
29	24389	88	681472	147	3176523	206	8741816	265	18609625	324	34012224
30	27000	89	704969	148	3241792	207	8869743	266	18821096	325	34328125
31	29791	90	729000	149	3307949	208	8998912	267	19034163	326	34645976
32	32768	91	753571	150	3375000	209	9123329	268	19248832	327	34965783
33	35937	92	778688	151	3442951	210	9261000	269	19465109	328	35287552
34	39304	93	804357	152	3511808	211	9393931	270	19683000	329	35611289
35	42875	94	830584	153	3581577	212	9528128	271	19902511	330	35937000
36	46656	95	857375	154	3652264	213	9663597	272	20123648	331	36264691
37	50653	96	884736	155	3723875	214	9800344	273	20346417	332	36594368
38	54872	97	912673	156	3796416	215	9938375	274	20570824	333	36926037
39	59319	98	941192	157	3869893	216	10077696	275	20796875	334	37259704
40	64000	99	970299	158	3944312	217	10218313	276	21024576	335	37595375
41	68921	100	1000000	159	4019679	218	10360232	277	21253933	336	37933056
42	74088	101	1030301	160	4096000	219	10503459	278	21484952	337	38272753
43	79507	102	1061208	161	4173281	220	10648000	279	21717639	338	38614472
44	85184	103	1092727	162	4251528	221	10793861	280	21952000	339	38958219
45	91125	104	1124864	163	4330747	222	10941048	281	22188041	340	39304000
46	97336	105	1157625	164	4410944	223	11089567	282	22425768	341	39651821
47	103823	106	1191016	165	4492125	224	11239424	283	22665187	342	40001688
48	110592	107	1225043	166	4574296	225	11390625	284	22906304	343	40353607
49	117649	108	1259712	167	4657463	226	11543176	285	23149125	344	40707584
50	125000	109	1295029	168	4741632	227	11697083	286	23393656	345	41063625
51	132651	110	1331000	169	4826809	228	11852352	287	23639903	346	41421736
52	140608	111	1367631	170	4913000	229	12008989	288	23887872	347	41781923
53	148877	112	1404928	171	5000211	230	12167000	289	24137569	348	42144192
54	157464	113	1442897	172	5088448	231	12326391	290	24389000	349	42508549
55	166375	114	1481544	173	5177717	232	12487168	291	24642171	350	42875000
56	175616	115	1520875	174	5268024	233	12649337	292	24897088	351	43243551
57	185193	116	1560896	175	5359375	234	12812904	293	25153757	352	43614208
58	195112	117	1601613	176	5451776	235	12977875	294	25412184	353	43986977
59	205379	118	1643032	177	5545233	236	13144256	295	25672375	354	44361864



Num.	Cubes	Num.	Cubes	Num.	Cubes	Num.	Cubes	Num.	Cubes	Num.	Cubes
355	44738875	417	72511713	479	109902239	541	158340421	603	219256227	665	294079625
356	45118016	418	73034632	480	110592000	542	159220088	604	220348864	666	295408296
357	45499293	419	73560059	481	111284641	543	160103007	605	221445125	667	296740963
358	45882712	420	74088000	482	111980168	544	160989184	606	222545016	668	298077632
359	46268279	421	74618461	483	112678587	545	161878625	607	223648543	669	299418309
360	46656000	422	75151448	484	113379904	546	162771336	608	224755712	670	300763000
361	47045881	423	75686967	485	114084125	547	163667323	609	225866529	671	302111711
362	47437923	424	76225024	486	114791256	548	164566592	610	226981000	672	303464448
363	47832147	425	76765625	487	115501303	549	165469149	611	228099131	673	304821217
364	48228544	426	77308776	488	116214272	550	166375000	612	229220928	674	306182024
365	48627125	427	77854483	489	116930169	551	167284151	613	230346397	675	307546875
366	49027896	428	78402752	490	117649000	552	168196608	614	231475544	676	308915776
367	49430863	429	78953589	491	118370771	553	169112377	615	232608375	677	310288733
368	49836032	430	79507000	492	119095488	554	170031464	616	233744896	678	311665752
369	50243409	431	80062991	493	119823157	555	170953875	617	234885113	679	313046839
370	50653000	432	80621568	494	120553784	556	171879616	618	236029032	680	314432000
371	51064811	433	81182737	495	121287375	557	172808693	619	237176659	681	315821241
372	51478848	434	81746504	496	122023936	558	173741112	620	238328000	682	317214568
373	51895117	435	82312875	497	122763473	559	174676879	621	239483061	683	318611987
374	52315624	436	82881856	498	123505992	560	175616000	622	240641848	684	320013504
375	52734375	437	83453453	499	124251499	561	176558481	623	241804367	685	321419125
376	53157376	438	84027672	500	125000000	562	177504328	624	242970624	686	322828856
377	53582633	439	84604519	501	125751501	563	178453547	625	244140625	687	324242703
378	54010152	440	85184000	502	126506008	564	179406144	626	245314376	688	325660672
379	54439939	441	85766121	503	127263527	565	180362125	627	246491883	689	327082769
380	54872000	442	86350888	504	128024064	566	181321496	628	247673152	690	328509000
381	55306341	443	86938307	505	128787625	567	182284263	629	248858189	691	329939371
382	55742968	444	87528384	506	129554216	568	183250432	630	250047000	692	331373888
383	56181887	445	88121125	507	130323843	569	184220009	631	251239591	693	332812557
384	56623104	446	88716536	508	131096512	570	185193000	632	252435968	694	334255384
385	57066625	447	89314623	509	131872229	571	186169411	633	253636137	695	335702375
386	57512456	448	89915392	510	132651000	572	187149248	634	254840104	696	337153536
387	57960603	449	90518849	511	133432831	573	188132517	635	256047875	697	338608873
388	58411072	450	91125000	512	134217728	574	189119224	636	257259456	698	340068392
389	58863869	451	91733851	513	135005697	575	190109375	637	258474853	699	341532099
390	59319000	452	92345408	514	135796744	576	191102976	638	259694072	700	343000000
391	59776471	453	92959677	515	136590875	577	192100033	639	260917119	701	344472101
392	60236288	454	93576664	516	137388096	578	193100552	640	262144000	702	345948008
393	60698457	455	94196375	517	138188413	579	194104539	641	263374721	703	347428927
394	61162984	456	94818816	518	138991832	580	195112000	642	264609288	704	348913664
395	61629875	457	95443993	519	139798359	581	196122941	643	265847707	705	350402625
396	62099136	458	96071912	520	140608000	582	197137368	644	267089984	706	351895816
397	62570773	459	96702579	521	141420761	583	198155287	645	268336125	707	353393243
398	63044792	460	97336000	522	142236648	584	199176704	646	269586136	708	354894912
399	63521199	461	97972181	523	143055667	585	200201625	647	270840023	709	356400829
400	64000000	462	98611128	524	143877824	586	201230056	648	272097792	710	357911000
401	64481201	463	99252847	525	144703125	587	202262003	649	273359449	711	359425431
402	64964808	464	99897344	526	145531576	588	203297472	650	274625000	712	360944128
403	65450827	465	100544625	527	146363183	589	204336469	651	275894451	713	362467097
404	65939264	466	101194696	528	147197952	590	205379000	652	277167808	714	363994344
405	66430125	467	101847563	529	148035889	591	206425071	653	278445077	715	365525875
406	66923416	468	102503232	530	148877000	592	207474688	654	279726264	716	367061696
407	67419143	469	103161709	531	149721291	593	208527857	655	281011375	717	368601813
408	67911312	470	103823000	532	150568768	594	209584584	656	282300416	718	370146232
409	68417929	471	104487111	533	151419437	595	210644875	657	283593393	719	371694959
410	68921000	472	105154048	534	152273304	596	211708736	658	284890312	720	373248000
411	69426531	473	105823817	535	153130375	597	212776173	659	286191179	721	374805361
412	69934528	474	106496424	536	153990656	598	213847192	660	287496000	722	376367048
413	70444997	475	107171875	537	154854153	599	214921799	661	288804781	723	377933067
414	70957944	476	107850176	538	155720872	600	216000000	662	290117528	724	379503424
415	71473375	477	108531333	539	156590819	601	217081801	663	291434247	725	381078125
416	71991296	478	109215352	540	157464000	602	218167208	664	292754944	726	382657176



Num.	Cubes.	Num.	Cubes.	Num.	Cubes.	Num.	Cubes.	Num.	Cubes.
727	384240583	782	478211768	837	586376253	892	709732288	947	849278123
728	385828352	783	480048687	838	588480472	893	712121957	948	851971392
729	387420489	784	481890304	839	590589719	894	714516984	949	854670349
730	389017080	785	483736625	840	592704000	895	716917375	950	857375000
731	390617891	786	485587656	841	594823321	896	719323136	951	860085351
732	392223168	787	487443403	842	596947688	897	721734273	952	862801408
733	393832837	788	489303872	843	599077107	898	724150792	953	865523177
734	395446904	789	491169069	844	601211584	899	726572699	954	868250664
735	397065375	790	493039000	845	603351125	900	729000000	955	870933875
736	398688256	791	494913671	846	605495736	901	731432701	956	873722816
737	400315553	792	496793088	847	607645423	902	733870808	957	876467493
738	401947272	793	498677257	848	609800192	903	736314327	958	879217912
739	403583419	794	500566184	849	611960049	904	738763264	959	881974079
740	405224000	795	502459875	850	614125000	905	741217625	960	884736000
741	406869021	796	504358336	851	616295051	906	743677416	961	887503681
742	408518488	797	506261573	852	618470208	907	746142643	962	890277128
743	410172407	798	508169592	853	620650477	908	748613312	963	893056347
744	411830784	799	510082399	854	622835864	909	751089129	964	895841344
745	413493625	800	512000000	855	625026375	910	753571000	965	898632125
746	415160936	801	513922401	856	627222016	911	756058031	966	901428696
747	416832723	802	515849608	857	629422793	912	758550528	967	904231063
748	418508992	803	517781627	858	631628712	913	761048497	968	907039232
749	420189749	804	519718464	859	633839779	914	763551944	969	909853209
750	421875000	805	521660125	860	636056000	915	766060875	970	912673000
751	423564751	806	523606616	861	638277381	916	768575296	971	915498611
752	425259008	807	525557943	862	640503928	917	771095213	972	918330048
753	426957777	808	527514112	863	642735647	918	773620632	973	921167317
754	428661064	809	529475129	864	644972544	919	776151559	974	924010424
755	430368875	810	531441000	865	647214625	920	778688000	975	926859375
756	432081216	811	533411731	866	649461896	921	781229961	976	929714176
757	433798093	812	535387328	867	651714363	922	783777448	977	932574833
758	435519512	813	537366797	868	653972032	923	786330467	978	935441352
759	437245479	814	539353144	869	656234909	924	788889024	979	938313739
760	438976000	815	541343375	870	658503000	925	791453125	980	941192000
761	440711081	816	543338496	871	660776311	926	794022776	981	944076141
762	442450728	817	545338513	872	663054848	927	796597983	982	946966168
763	444194947	818	547343432	873	665338617	928	799178752	983	949862087
764	445943744	819	549353259	874	667627624	929	801765089	984	952763904
765	447697125	820	551368000	875	669921875	930	804357000	985	955671625
766	449455096	821	553387661	876	672221376	931	806954491	986	958585256
767	451217663	822	555412248	877	674526133	932	809557568	987	961504803
768	452984832	823	557441767	878	676836152	933	812166237	988	964430272
769	454756609	824	559476224	879	679151439	934	814780504	989	967361669
770	456533000	825	561515625	880	681472000	935	817400375	990	970297000
771	458314011	826	563559976	881	683797841	936	820025856	991	973242271
772	460099648	827	565609283	882	686128968	937	822656953	992	976191488
773	461889917	828	567663552	883	688465387	938	825293672	993	979146657
774	463684824	829	569722789	884	690807104	939	827936019	994	982107784
775	465484375	830	571787000	885	693154125	940	830584000	995	985074375
776	467288576	831	573856191	886	695506456	941	833237621	996	988047936
777	469097433	832	575930368	887	697864103	942	835896888	997	991026973
778	470910955	833	578009537	888	700227072	943	838561807	998	994011992
779	472729130	834	580093704	889	702595369	944	841232384	999	997002999
780	474552000	835	582182875	890	704969000	945	843908625	1000	1000000000
781	476379541	836	584277056	891	707347971	946	846590536		



*The Cube of a Binomial*, is equal to the cubes of the two parts or members, together with triple of the two parallelopipedons under each part and the square of the other; viz,  $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$ . And hence the common method of extracting the cube root.

*Cubic Equations*, are those in which the unknown quantities rise to three dimensions; as  $x^3 = a$ , or  $x^3 + ax^2 = b$ , or  $x^3 + ax^2 + bx = c$ , &c.

All cubic equations may be reduced to this form,  $x^3 + px = q$ ; viz, by taking away the 2d term.

All cubic equations have three roots; which are either all real, or else one only is real, and the other two imaginary; for all roots become imaginary by pairs.

But the nature of the roots as to real and imaginary, is known partly from the sign of the co-efficient  $p$ , and partly from the relation between  $p$  and  $q$ : for the equation has always two imaginary roots when  $p$  is positive; it has also two imaginary roots when  $p$  is negative, provided  $\frac{1}{3}p^3$  is less  $\frac{1}{2}q^2$ , or  $4p^3$  less than  $27q^2$ ; otherwise the roots are all real, namely, whenever  $p$  is negative, and  $4p^3$  either equal to, or greater than  $27q^2$ .

Every cubic equation of the above form, viz, wanting the 2d term, has both positive and negative roots, and the greatest root is always equal to the sum of the two less roots; viz, either one positive root equal to the sum of the two negative ones, or else one negative root equal to the sum of two smaller and positive ones. And the sign of the greatest, or single root, is positive or negative, according as  $q$  is positive or negative when it stands on the right-hand side of the equation, thus  $x^3 + px = q$ ; and the two smaller roots have always the contrary sign to  $q$ .

So that, in general, the sign of  $p$  determines the nature of the roots, as to real and imaginary; and the sign of  $q$  determines the affection of the roots, as to positive and negative. See my Tract on Cubic Equations in the Philos. Transf. for 1780.

*To find the Values of the Roots of Cubic Equations.* Having reduced the equation to this form  $x^3 + px = q$ , its root may be found in various ways; the first of these, is that which is called Cardan's Rule, by whom it was first published, but invented by Ferreus and Tartalea. See ALGEBRA. The rule is this: Put  $a = \frac{1}{3}p$ , and  $b = \frac{1}{2}q$ ; then is Cardan's root

$x = \sqrt[3]{b + \sqrt{b^2 + a^3}} + \sqrt[3]{b - \sqrt{b^2 + a^3}}$ ; or if there be put  $s = \sqrt[3]{b + \sqrt{b^2 + a^3}}$ , and  $d = \sqrt[3]{b - \sqrt{b^2 + a^3}}$ ; then  $s + d = x$ , the 1st or Cardan's root,

also  $-\frac{s+d}{2} + \frac{s-d}{2}\sqrt{-3}$  is the 2d root,

and  $-\frac{s+d}{2} - \frac{s-d}{2}\sqrt{-3}$  is the 3d root.

Now the first of these, or Cardan's root, is always a real root, though it is not always the greatest root, as it has been commonly mistaken for. And yet this rule always exhibits the root in the form of an imaginary quantity when the equation has no imaginary roots at all; but in the form of a real quantity when the equation has two imaginary roots. See the reason of this explained in my Tract above cited, pa. 407. As to

the other two roots, viz  $-\frac{s+d}{2} \pm \frac{s-d}{2}\sqrt{-3}$ , though, in their general form, they have an imaginary appearance; yet, by substituting certain particular numbers, they come out in a real form in all such cases as they ought to be so.

But, after the first root is found, by Cardan's rule, the other two roots may be found, or exhibited, in several other different ways; some of which are as follow:

Let  $r$  denote the 1st, or Cardan's root,

and  $v$  and  $w$  the other two roots:

then is  $v + w = -r$ , and  $vw = -q$ ;

and the resolution of these two equations will give the other two roots  $v$  and  $w$ .

Or resolve the quadratic equation  $x^2 + rx + r^2 + p = 0$ , and its two roots will be those sought.

Or the same two roots will be

either  $-\frac{1}{2}r + \frac{1}{2}\sqrt{-p - \frac{3q}{r}}$  and  $-\frac{1}{2}r - \frac{1}{2}\sqrt{-p - \frac{3q}{r}}$ ,

or  $-\frac{1}{2}r + \frac{1}{2}\sqrt{r^2 - \frac{4q}{r}}$  and  $-\frac{1}{2}r - \frac{1}{2}\sqrt{r^2 - \frac{4q}{r}}$ ,

or  $-\frac{1}{2}r + \frac{1}{2}\sqrt{1 - \frac{4q}{q - pr}}$  and  $-\frac{1}{2}r - \frac{1}{2}\sqrt{1 - \frac{4q}{q - pr}}$ .

Ex. 1. If the equation be  $x^3 - 15x = 4$ : here  $p = -15$ ,  $q = 4$ ,  $a = -5$ ,  $b = 2$ ; hence  $\sqrt{b^2 + a^3} = \sqrt{4 - 125} = \sqrt{-121} = 11\sqrt{-1}$ ,  $s = \sqrt[3]{2 + \sqrt{-121}} = 2 + \sqrt{-1}$ , and  $d = \sqrt[3]{2 - \sqrt{-121}} = 2 - \sqrt{-1}$ : therefore  $r = s + d = 4$ , the 1st root; and

$-\frac{s+d}{2} \pm \frac{s-d}{2}\sqrt{-3} = -2 \mp \sqrt{3}$ , the other two roots.

Ex. 2. If  $x^3 - 6x = 4$ : here  $a = -2$ , and  $b = 2$ ; therefore  $\sqrt{b^2 + a^3} = \sqrt{4 - 8} = \sqrt{-4} = 2\sqrt{-1}$ ,  $s = \sqrt[3]{2 + 2\sqrt{-1}} = -1 + \sqrt{-1}$ , &  $d = \sqrt[3]{2 - 2\sqrt{-1}} = -1 - \sqrt{-1}$ : hence then  $r = s + d = -2$ , the first root; and  $1 \pm \sqrt{3}$  the other two roots.

Ex. 3. If  $x^3 + 18x = 6$ : here  $a = 6$ , and  $b = 3$ ; then  $\sqrt{b^2 + a^3} = \sqrt{9 + 216} = \sqrt{225} = 15$ ,  $s = \sqrt[3]{3 + 15} = \sqrt[3]{18}$ , and  $d = \sqrt[3]{3 - 15} = \sqrt[3]{-12} = -\sqrt[3]{12}$ : therefore  $r = s + d = \sqrt[3]{18} - \sqrt[3]{12} = .331313$ , the 1st root, and  $-\frac{\sqrt[3]{18} - \sqrt[3]{12}}{2} \pm \frac{\sqrt[3]{18} + \sqrt[3]{12}}{2}\sqrt{-3}$  are the two other roots.

2. Another method for the roots of the equation  $x^3 + px = q$ , is by means of infinite series, as shewn at pa. 415 and seq. of my Tract above cited; whence it appears that the roots are exhibited in various forms of series as follow: viz,

$$2\sqrt[3]{b} \times : 1 - \frac{2c^2}{3 \cdot 6b^2} - \frac{2 \cdot 5 \cdot 8c^4}{3 \cdot 6 \cdot 9 \cdot 12b^4} \&c$$

for the 1st root, and

$$-\sqrt[3]{b} \times : 1 - \frac{2c^2}{3 \cdot 6b^2} - \frac{2 \cdot 5 \cdot 8c^4}{3 \cdot 6 \cdot 9 \cdot 12b^4} \&c$$

$$\pm \frac{c\sqrt{-3}}{\sqrt[3]{b^2}} \times : \frac{1}{3} + \frac{2 \cdot 5c^2}{3 \cdot 6 \cdot 9b^2} + \frac{2 \cdot 5 \cdot 8 \cdot 11c^4}{3 \cdot 6 \cdot 9 \cdot 12 \cdot 15b^4} \&c$$

for the two other roots: where  $a = \frac{1}{3}p$ ,  $b = \frac{1}{2}q$ , and  $c = \sqrt{b^2 + a^3}$ .

And



And various other series for the same purpose may also be seen in my Tract, so often before cited.

3. A third method for the roots of cubic equations, is by angular sections, and the table of sines. It was first hinted by Bombelli, in his Algebra, that angles are trisected by the resolution of cubic equations. Afterwards, Vieta gave the resolution of cubics, and the higher equations, by angular sections. Next, Albert Girard, in his *Invention Nouvelle en l'Algebre*, shews how to resolve the irreducible case in cubics by a table of sines: and he also constructs the same, or finds the roots, by the intersection of the hyperbola and circle. Halley and De Moivre also gave rules and examples of the same sort of resolutions by a table of sines. And, lastly, Mr. Anthony Thacker invented, and Mr. William Brown computed, a large set of similar tables, for resolving affected quadratic and cubic equations, with their application to the resolution of biquadratic ones.

4. Lastly, the several methods of approach, or approximation, for the roots of all affected equations, which have been used in various ways by Stevin, Vieta, Newton, Halley, Raphson, and others.

To these may be added the method of Trial-and-error, or of Double Position, one of the easiest and best of any. Of this method, let there be taken the last example, viz,  $x^3 + 18x = 6$ , in which it is evident that  $x$  is very nearly equal to  $\frac{1}{3}$ , but a little less; take it therefore  $x = .33$ ; then  $x^3 + 18x = 5.975937$ , but should be 6, and therefore the error is .024063 in defect.

Again suppose  $x = .34$ ; then  $x^3 + 18x = 6.159304$ , which is .159304 in excess.


Therefore 
$$\frac{.34 \times .024063 + .33 \times .159304}{.024063 + .159304} = \frac{.60752}{.183367} = .3313,$$
 the root as before very nearly.

For the construction of cubic equations, see CONSTRUCTION.

CUBIC Foot, of any thing, is so much of it as is contained in a cube whose side is one foot.

CUBIC Hyperbola, is a figure expressed by the equation  $xy^2 = a$ , having two asymptotes, and consisting of two hyperbolas, lying in the adjoining angles of the asymptotes, and not in the opposite angles, like the Apollonian hyperbola; being otherwise called by Newton, in his *Enumeratio Linearum Tertii Ordinis*, an hyperbolismus of a parabola; and is the 65th species of those lines according to him.

CUBIC Numbers. See CUBES.




CUBIC Parabola, a curve, as BCD, of the 2d order, having two infinite legs CB, CD, tending contrary

ways. And if the absciss, AP or  $x$ , touch the curve in C, the relation between the absciss and ordinate, viz,  $AP = x$ , and  $PM = y$ , is expressed by the equation  $y = ax^3 + bx^2 + cx + d$ ; or when A coincides with C, by the equation  $y = ax^3$ , which is the simplest form of the equation of this curve.

If the right line AP (fig. 2) cut the cubical parabola in three points A, B, C; and from any point P there be drawn the right line or ordinate PM, cutting the curve in one point M only: then will PM be always as the solid  $AP \times BP \times CP$ ; which is an essential property of this curve.

And hence it is easy to construct a cubic equation, as  $x^3 + a^2x = b^3$ , by the intersection of this curve and a right line. See the Construction of a cubic equation by means of the cubic parabola and a right line, by Dr. Wallis, in his Algebra: As also the Construction of equations of 6 dimensions, by means of the cubic parabola and a circle, by Dr. Halley, in a lecture formerly read at Oxford.

The curve of this parabola cannot be rectified, not even by means of the conic sections. But a circle may be found equal to the Curve Surface, generated by the rotation of the curve AM about the tangent AP to the principal vertex A. Let MN be an ordinate, and MT a tangent at the point M; and let PM be parallel to AN. Divide MN in the point O, so that MO be to ON as TM to MN. Then a mean proportional between TM + ON and  $\frac{1}{3}$  of AN will be the radius of a circle, whose area is equal to the superficies described by that rotation, viz, of AM about AP.



The Area of a Cubic Parabola is  $\frac{3}{4}$  of its circumscribing parallelogram.

CUBIC Root, of any number, or quantity, is such a quantity as being cubed, or twice multiplied by itself, shall produce that which was given. So, the cubic root of 8 is 2, because  $2^3$  or  $2 \times 2 \times 2$  is equal to 8.

The common method of extracting the cube root, founded on the property given above, viz,  $(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$ , is found in every book of common arithmetic, and is as old at least as Lucas de Burgo, where it is first met with in print. Other methods for the cube root may be seen under the article EXTRACTION of Roots, particularly this one, viz, the cube

root of  $n$ , or  $\sqrt[3]{n} = \frac{2n + a^3}{n + 2a^3} \times a$ , very nearly, or  $2a^3 + n$ :  
 $2n + a^3 :: a : \sqrt[3]{n}$  the cube root of  $n$  nearly; where  $n$  is

Y y

is



is any number given whose cube root is sought, and  $a^3$  is the nearest complete cube to  $n$ , whether greater or less.

For example, suppose it were proposed to double the cube, or, which comes to the same thing, to extract the cube root of the number 2. Here the nearest cube is 1, whose cube root is 1 also, that is,  $a^3 = 1$ , and  $a = 1$ , also  $n = 2$ ; therefore

$$2a^3 + n \quad 2n + a^3 \quad a$$

$$4 : 5 :: 1 : \frac{5}{4} = 1.25 = \sqrt[3]{2} \text{ nearly.}$$

But, for a nearer value, assume now  $a = \frac{5}{4}$ , or  $a^3 = \frac{125}{64}$ ; then is  $2a^3 + n = \frac{250}{64} + 2 = \frac{378}{32}$ , and  $2n + a^3 = 4 + \frac{125}{64} = \frac{381}{32}$ ; hence  $378 : 381 :: \frac{5}{4} : \frac{635}{256}$  or  $1.259921 = \sqrt[3]{2}$ , or the cube root of 2, which is true in the last place of decimals.

And this is the simplest and easiest method for the cube root of any number. See its investigation in my *Traacts*, vol. 1, pa. 49.

Every number or quantity has three cubic roots, one that is real, and two imaginary: So, the cube root of 1 is either 1, or  $-\frac{1 + \sqrt{-3}}{2}$ , or  $-\frac{1 - \sqrt{-3}}{2}$ ; and if

$r$  be the real root of any cube  $r^3$ , the two imaginary cubic roots of it will be

$$-\frac{1 + \sqrt{-3}}{2}r, \text{ and } -\frac{1 - \sqrt{-3}}{2}r : \text{ for any one of}$$

these being cubed, gives the same cube  $r^3$ .

**CUBING of a Solid.** See **CUBATURE**.

**CUBO-CUBE**, the 6th power.

**CUBO-CUBO-CUBE**, the 9th power.

**CULMINATION**, the passage of a star or planet over the meridian, or that point of its orbit which it is in at its greatest altitude. Hence a star is said to culminate, when it passes the meridian.

*To find the time of a Star's culminating.* Estimate the time nearly; and find the right ascension both of the sun and star corrected for this estimated time; then the difference between these right ascensions, converted into solar time, at the rate of 15 degrees to the hour, gives the time of southing. See an example of this calculation every year in *White's Ephemeris*, pa. 45.

**CULVERIN**, was the name of a piece of ordnance; but is now disused.

**CUNETTE.** See **CUVETTE**.

**CUNEUS.** See **WEDGE**.

**CUNITIA (MARIA)**, a lady of Silesia, who was famous for her extensive knowledge in many branches of learning, but more particularly in mathematics and astronomy, upon which she wrote several ingenious treatises; particularly one, entitled *Urania Propitia*, printed in 1650, in Latin and German, and dedicated to Ferdinand the third, emperor of Germany. In this work are contained astronomical tables of great ease and accuracy, founded upon Kepler's hypothesis. But notwithstanding her merit shines with such peculiar lustre as to reflect honour on her sex, history does not inform us of the time of her birth or death.

**CURRENT**, a stream or flux of water in any direction. The *setting* of the current, is that point of the

compass towards which the waters run; and the *drift* of a current is the rate it runs an hour.


Currents in the sea, are either natural and general, as arising from the diurnal rotation of the earth on its axis, or the tides, &c; or accidental and particular, caused by the waters being driven against promontories, or into gulphs and streights; from whence they are forced back, and thus disturb the natural flux of the sea.

The currents are so violent under the equator, where the motion of the earth is greatest, that they hurry vessels very speedily from Africa to America; but absolutely prevent their return the same way: so that ships are forced to run as far as the 40th or 45th degree of north latitude, to fall into the return of the current again, to bring them home to Europe. It is shewn by Governor Pownal, that this current performs a continual circulation, setting out from the Guinea coast in Africa, for example, from thence crossing straight over the Atlantic ocean, and so setting into the Gulph of Mexico by the south side of it; then sweeping round by the bottom of the Gulph, it issues out by the north side of it, and thence takes a direction north-easterly along the coast of North America, till it arrives near Newfoundland, where it is turned by a rounding motion backward across the Atlantic again, upon the coasts of Europe, and from thence southward again to the coast of Africa, from whence it set out.

In the streights of Gibraltar, the currents set in by the south side, sweep along the coast of Africa to Egypt, by Palestine, and so return by the northern side, or European coasts, and issue out again by the northern side of the streights. In St. George's Channel too they usually set eastward. The great violence and danger of the sea in the Streights of Magellan, is attributed to two contrary currents setting in, one from the south, and the other from the north sea.

Currents are of some consideration in the art of navigation, as a ship is by them either accelerated or retarded in her course, according as the set is with or against the ship's motion. If a ship sail along the direction of a current, it is evident that the velocity of the current must be added to that of the vessel: but if her course be directly against the current, it must be subtracted: and if she sail athwart it, her motion will be compounded with that of the current; and her velocity augmented or retarded, according to the angle of her direction with that of the current; that is, she will proceed in the diagonal of the parallelogram formed according to the two lines of direction, and will describe or pass over that diagonal in the same time in which she would have described either of the sides, by the separate velocities.

For suppose ABDC be a parallelogram, the diagonal of which is AD. Now if the wind alone would drive the ship from A to B, in the same time as the current alone would drive it from A to C: Then, as the wind neither helps nor hinders the ship from coming towards the line CD, the current will bring it there in the same time as if the wind did not act: And as the current neither helps nor hinders the ship from coming





coming towards the line BD, the wind will bring it there in the same time as if the current did not act: Therefore the ship must, at the end of that time, be found in both those lines, that is, in their meeting D: Consequently the ship must have passed from A to D in the diagonal AD. See COMPOSITION of Forces.

See the Sailing in Currents largely exemplified in Robertson's Navigation, vol. 2, book 7, sect. 8.

CURSOR, a small piece of brass &c that slides; as, the piece in an equinoctial ring dial that slides to the day of the month; or the little ruler or label of brass sliding in a groove along the middle of another label, representing the horizon in the analemma; or the point that slides along the beam compass; &c.

CURTATE Distance, is the distance of a planet's place from the sun, reduced to the ecliptic; or, the interval between the sun and that point where a perpendicular, let fall from the planet, meets the ecliptic.

CURTATION, the interval between a planet's distance from the sun, and the curtate distance.


From the foregoing article it is easy to find the curtate distance; whence the manner of constructing tables of curtation is obvious; the quantity of inclination, reduction, and curtation of a planet, depending on the argument of latitude. Kepler, in his Rodolphine Tables, reduces the tables of them all into one, under the title of *Tabula Latitudinaria*.

CURTIN, CURTAIN, or COURTINE, in Fortification, that part of a wall or rampart that joins two bastions, or lying between the flank of one and that of another.—The curtain is usually bordered with a parapet 5 feet high; behind which the soldiers stand to fire upon the covert-way, and into the moat.

CURVATURE of a Line, is its bending, or flexure; by which it becomes a curve, of any peculiar form and properties. Thus, the nature of the curvature of a circle is such, as that every point in the periphery is equally distant from a point within, called the centre; and so the curvature of the same circle is every where the same; but the curvature in all other curves is continually varying.—The curvature of a circle is so much the more, as its radius is less, being always reciprocally as the radius; and the curvature of other curves is measured by the reciprocal of the radius of a circle having the same degree of curvature as any curve has, at some certain point.

Every curve is bent from its tangent by its curvature, the measure of which is the same as that of the angle of contact formed by the curve and tangent. Now the same tangent AB is common to an infinite number of circles, or other curves, all touching it and each other in the same point of contact C. So that any curve DCE may be touched by an infinite number of different circles at the same point C; and some of these circles fall wholly within it, being more curved, or having a greater curvature than that curve; while others fall without it near the point of contact, or between the curve and tangent at that point, and so, being less deflected from the tangent than the curve is, they have a less degree of curvature there. Consequently there is one, of this infinite number of circles, which neither falls below it nor above it, but, being equally deflected from the tangent, coincides with it most intimately of all the circles; and the radius of this circle is called the

radius of curvature of the curve; also the circle itself is called the circle of curvature, or the osculatory circle of that curve, because it touches it so closely that no other circle can be drawn between it and the curve.



As a curve is separated from its tangent by its flexure or curvature, so it is separated from the osculatory circle by the increase or decrease of its curvature; and as its curvature is greater or less, according as it is more or less deflected from the tangent, so the variation of curvature is greater or less, according as it is more or less separated from the circle of curvature.

It appears, however, from the demonstration of geometers, that circles may touch curve lines in such a manner, that there may be indefinite degrees of more or less intimate contact between the curve and its osculatory circle; and that a conic section may be described that shall have the same curvature with a given line at a given point, and the same variation of curvature, or a contact of the same kind with the circle of curvature.

If we conceive the tangent of any proposed curve to be a base, and that a new line or curve be described, whose ordinate, upon the same base or absciss, is a 3d proportional to the ordinate and base of the first; this new curve will determine the chord of the circle of curvature, by its intersection with the ordinate at the point of contact; and it will also measure the variation of curvature, by means of the tangent of the angle in which it cuts that circle: the less this angle is, the closer is the contact of the curve and circle of curvature; and of this contact there may be indefinite degrees.

For example, let EMH be any curve, to which ET is a tangent at the point E; then let there be always taken  $MT:ET::ET:TK$ , and through all the points K draw the curve BKF; then from the point of contact E draw EB parallel to the ordinate TK, meeting the last curve in B; and finally, describe a circle ERQB through the point B and touching ET in E; and it shall be the osculatory circle to the given curve EMH. And the contact of EM and ER is always the closer, the less the angle KBQ is. See Maclaurin's Fluxions, art. 366.

Hence it follows, that the contact of the curve EMH and its osculatory circle is closest, when the curve BK touches the arch BQ in B, the angle BET being given; and it is farthest from this, or most open, when BK touches the right line BE in B.

Hence also there may be indefinite degrees of more and more intimate contact between a circle and a curve. The first degree is when the same right line touches



touches them both in the same point; and a contact of this sort may take place between any circle and any arch of a curve. The 2d is when the curve EMH and circle ERB have the same curvature, and the tangents of the curve BKF and circle BQE intersect each other at B in any assignable angle. The contact of the curve EM and circle of curvature ER at E, is of the 3d degree, or order, and their osculation is of the 2d, when the curve BKF touches the circle BQE at B, but so as not to have the same curvature with it. The contact is of the 4th degree, or order, and their osculation of the 3d, when the curve BKF has the same curvature with the circle BQE at B, but so as that their contact is only of the 2d degree. And this gradation of more and more intimate contact, or of approximation towards coincidence, may be continued indefinitely, the contact of EM and ER at E being always of an order two degrees closer than that of BK and BQ at B. There is also an indefinite variety comprehended under each order: thus, when EM and ER have the same curvature, the angle formed by the tangents of BK and BQ admits of indefinite variety, and the contact of EM and ER is the closer the less that angle is. And when that angle is of the same magnitude, the contact of EM and ER is the closer, the greater the circle of curvature is. When BK and BQ touch at B, they may touch on the same or on different sides of their common tangent; and the angle of contact KBQ may admit of the same variety with the angle of contact MER; but as there is seldom occasion for considering those higher degrees of more intimate contact of the curve EMH and circle of curvature ERB, Mr. Maclaurin calls the contact or osculation of the same kind, when, the chord EB and angle BET being given, the angle contained by the tangents of BK and BQ is of the same magnitude.

When the curvature of EMH increases from E towards H, and consequently corresponds to that of a circle gradually less and less, the arch EM falls within ER, the arch of the osculatory circle, and BK is within BQ. The contrary happens when the curvature of EM decreases from E towards H, and consequently corresponds to that of a circle which is gradually greater and greater, the arch EM falls without ER, and BK is without BQ. And according as the curvature of EM varies more or less, it is more or less unlike to the uniform curvature of a circle; the arch of the curve EMH separates more or less from the arch of the osculatory circle ERB, and the angle contained by the tangents of BKF and BQE at B, is greater or less. Thus the quality of curvature, as it is called by Newton, depends on the angle contained by the tangents of BK and BQ at B; and the measure of the inequability or variation of curvature, is as the tangent of this angle, the radius being given, and the angle BET being a right one.

The radii of curvature of similar arcs in similar figures, are in the same ratio as any homologous lines of these figures; and the variation of curvature is the same. See Maclaurin, art. 370.

When the proposed curve EMH is a conic section, the new line BKF is also a conic section; and it is a right line when EMH is a parabola, to the axis of which the ordinates TK are parallel. BKF is also a

right line when EMH is an hyperbola, to one asymptote of which the ordinate TK is parallel.

When the ordinate EB, at the point of contact E, instead of meeting the new curve BK, is an asymptote to it, the curvature of EM will be less than in any circle; and this is the case in which it is said to be infinitely little, or that the radius of curvature is infinitely great. And of this kind is the curvature at the points of contrary flexure in lines of the 3d order.

When the curve BK passes through the point of contact E, the curvature is greater than in any circle, or the radius of curvature vanishes; and in this case the curvature is said to be infinitely great. Of this kind is the curvature at the cusps of the lines of the 3d order.

As to the degree of curvature in lines of the 3d and higher orders, see Maclaurin, art. 379; also art. 380, when the proposed curve is mechanical.

As curves which pass through the same point have the same tangent when the first fluxions of the ordinates are equal, so they have the same curvature when the 2d fluxions of the ordinate are likewise equal; and half the chord of the osculatory circle that is intercepted between the points where it intersects the ordinate, is a 3d proportional to the right lines that measure the 2d fluxion of the ordinate and first fluxion of the curve, the base being supposed to flow uniformly. When a ray revolving about a given point, and terminated by the curve, becomes perpendicular to it, the first fluxion of the radius vanishes; and if its 2d fluxion vanish at the same time, that point must be the centre of curvature. The same may be said, when the angular motion of the ray about that point is equal to the angular motion of the tangent of the curve; as the angular motion of the radius of a circle about its centre is always equal to the angular motion of the tangent of the circle. Hence the various properties of the circle may suggest several theorems for determining the centre of curvature.

See art. 396 of the said book, also the following, concerning the curvature of lines that are described by means of right lines revolving about given poles, or of angles that either revolve about such poles, or are carried along fixed lines.


It is to be observed that, as when a right line intersects an arc of a curve in two points, if by varying the position of that line the two intersections unite in one point, it then becomes the tangent of the arc; so when a circle touches a curve in one point, and intersects it in another, if, by varying the centre, this intersection joins the point of contact, the circle has then the closest contact with the arc, and becomes the circle of curvature; but it still continues to intersect the curve at the same point where it touches it, that is, where the same right line is their common tangent, unless another intersection join that point at the same time. In general, the circle of curvature intersects the curve at the point of osculation, only when the number of the successive orders of fluxions of the radius of curvature, that vanishes when this radius comes to the point of osculation, is an even number.

It has been supposed by some, that two points of contact, or four intersections of the curve and circle of curvature, must join to form an osculation. But Mr.

James



James Bernoulli justly insisted, that the coalition of one point of contact and one intersection, or of three intersections, was sufficient. In which case, and in general, when an odd number of intersections only join each other, the point where they coincide continues to be an intersection of the curve and circle of curvature, as well as a point of their mutual contact and osculation. See Maclaurin's Flux. art. 493.



From these principles may be determined the circle of curvature at any point of a conic section. Thus, suppose AEMHG be any conic section, to the point E of which the circle or radius of curvature is to be found. Let ET be a tangent at E, and draw EGB and the tangent HI parallel to the chords of the circle of curvature; then take EB to EG as EI<sup>2</sup> to HI<sup>2</sup>; or, when the section has a centre O, as in the ellipse and hyperbola, as the square of the semi-diameter Oa parallel to ET, is to the square of the semi-diameter OA parallel to EB; and a circle EB described upon the chord EB that touches ET, will be the circle of curvature sought.

When BET is a right angle, or EB is the diameter of the circle of curvature, EG will be the axis of the conic section, and EB will be the parameter of this axis; also when the point G, where the conic section cuts EB, and the point B, are on the same side of E, then EMG will be an ellipsis, and EG the greater or less axis, according as EG is greater or less than EB.

The propositions relating to the curvature of the conic sections, commonly given by authors, follow without much difficulty from this construction.

1. When the chord of curvature, thus found, passes through the centre of the conic section, it will then be equal to the parameter of the diameter that passes through the point of contact.

2. The square of the semi-diameter Oa, is to the rectangle of half the transverse and half the conjugate axis, as the radius of curvature CE is to Oa. And therefore the cube of the semi-diameter Oa, parallel to the tangent ET, is equal to the solid contained by the radius of curvature CE, and the rectangle of the two axes.

3. The perpendicular to either axis bisects the angle made by the chord of curvature, and the common tangent of the conic section and circle of curvature.

4. The chord of the osculatory circle that passes through the focus, the diameter conjugate to that which passes through the point of contact, and the transverse axis of the figure, are in continued proportion.

5. When the section is an ellipse, if the circle of curvature at E meet Oa in d, the square of Ed will be equal to twice the square of Oa. Hence Ed : Oa ::  $\sqrt{2} : 1$ . Which gives an easy method of determining the circle of curvature to any point E, when the semi-diameter Oa is given in magnitude and position.

Several other properties of the circle of curvature, and methods of determining it when the section is given; or vice versa, of determining the section when the circle of curvature is given, may be seen in Maclaurin's Flux. art. 375. See also the Appendix to Maclaurin's Algebra, sect. 1.

To determine the Radius and Circle of Curvature by the Method of Fluxions. Let AEE be any curve, concave towards its axis AD; draw an ordinate DE to the point E where the curve is required to be found; and suppose EC perpendicular to the curve, and equal to the radius of the circle BEE of curvature sought; lastly, draw Ed parallel to AD, and de parallel and indefinitely near to DE; thereby making Ed the fluxion or increment of the absciss AD, also de the fluxion of the ordinate DE, and Ee that of the curve AE. Now put  $x = AD$ ,  $y = DE$ ,  $z = AE$ ; and  $r = CE$  the radius of curvature; then is  $ED = \dot{x}$ ,  $de = \dot{y}$ , and  $Ee = \dot{z}$ .

Now, by sim. tri. the 3 lines  $Ed$ ,  $de$ ,  $Ee$ , or  $\dot{x}$ ,  $\dot{y}$ ,  $\dot{z}$ , are respectively as the three  $GE$ ,  $GC$ ,  $CE$ ; therefore  $GC \cdot \dot{x} = GE \cdot \dot{y}$ ; and the flux. of this equa. is

$$GC \cdot \ddot{x} + \dot{GC} \cdot \dot{x} = GE \cdot \ddot{y} + \dot{GE} \cdot \dot{y},$$

or because  $GC = -BG$ , it is

$$GC \cdot \dot{x} - BG \cdot \dot{x} = GE \cdot \ddot{y} + \dot{GE} \cdot \dot{y}.$$

But, since the two curves AE and BE have the same curvature at the point E, their abscissas and ordinates have the same fluxions at that point, that is Ed or  $\dot{x}$  is the fluxion both of AD and BG, and de or  $\dot{y}$  is the fluxion both of DE and GE. In the above equation

therefore substitute  $\dot{x}$  for BG, and  $\dot{y}$  for GE, and it becomes  $GC \cdot \dot{x} - \dot{x} \dot{x} = GE \cdot \ddot{y} + \dot{y} \dot{y}$ , or  $GC \cdot \dot{x} - GE \cdot \ddot{y} = \dot{x}^2 + \dot{y}^2$  or  $\dot{z}^2$ .

Now mult. the three terms of this equation respectively by these three quantities,  $\frac{\dot{y}}{GC}$ ,  $\frac{\dot{x}}{GE}$ ,  $\frac{\dot{z}}{CE}$ , which are equal, and it becomes  $\dot{y} \dot{x} - \dot{x} \dot{y} = \frac{\dot{z}^3}{CE}$ , or  $\frac{\dot{z}^3}{CE}$ ;

and hence  $r = \frac{\dot{z}^3}{\dot{y} \dot{x} - \dot{x} \dot{y}}$ , which is the general value of the radius of curvature for all curves whatever, in terms of the fluxions of the absciss and ordinate.

Farther, as in any case either  $x$  or  $y$  may be supposed to flow equably, that is, either  $\dot{x}$  or  $\dot{y}$  constant quantities, or  $\dot{x}$  or  $\dot{y} = 0$ , by this supposition either of the terms in the denominator of the value of  $r$  may be made to vanish. So that when  $\dot{x}$  is constant, the value of  $r$  is  $\frac{\dot{z}^3}{-\dot{x} \dot{y}}$ , but  $r$  is  $\frac{\dot{z}^3}{\dot{y} \dot{x}}$  when  $\dot{y}$  is constant.

For example, suppose it were required to find the radius or circle of curvature to any point of a parabola, its vertex being A, and axis AD.—Now the equation



tion of the curve is  $ax = y^2$ ; hence  $ax = 2yy$ , and  $ax = 2y^2$ , supposing  $y$  constant, also  $a^2x^2 = 4y^2y^2$ ; hence  $r$  or

$$\frac{z^3}{y \ddot{x}} \text{ or } \frac{\dot{x}^2 + y^2}{y \ddot{x}}^{\frac{3}{2}} \text{ is } \frac{a^2 + 4y^2}{2a^2}^{\frac{3}{2}} \text{ or } \frac{a + 4x}{2\sqrt{a}}^{\frac{3}{2}},$$

the general value of the radius of curvature for any point E, the ordinate to which cuts off the absciss AD =  $x$ .



Hence, when  $x$  or the absciss is nothing, the last expression becomes barely  $\frac{a^{\frac{3}{2}}}{2\sqrt{a}} = \frac{1}{2}a = r$  for the radius of curvature at the vertex of the parabola; that is, the diameter of the circle of curvature at the vertex of a parabola, is equal to  $a$  the parameter of axis.

*Variation of CURVATURE.* See VARIATION.

*Double CURVATURE*, is used for the curvature of a line, which twists so that all the parts of it do not lie in the same plane.

*CURVE*, a line whose several parts proceed bowing, or tend different ways; in opposition to a straight line, all whose parts have the same course or direction.

The doctrine of curves, and of the figures and solids generated from them, constitute what is called the higher geometry.



In a curve, the line AD, which bisects all the parallel lines MN, is called a *diameter*; and the point A, where the diameter meets the curve, is called the *vertex*: if AD bisect all the parallels at right angles, it is called the *axis*. The parallel lines MN are called *ordinates*, or *applicates*; and their halves, PM, or PN, *semi-ordinates*. The portion of the diameter AP, between the vertex, or any other fixed point, and an ordinate, is called the *absciss*; also the concurrence of all the diameters, if they meet all in one point, is the *centre*. This definition of the diameter, as bisecting the parallel ordinates, respects only the conic sections, or such curves as are cut only in two points by the ordinates; but in the lines of the 3d order, which may be cut in three points by the ordinates, then the diameter is that line which cuts the ordinates so, that the sum of the two parts that lie on one side of it, shall be equal to the part on the other side: and so on for curves of higher orders, the sum of the parts of the ordinates on one side of the diameter, being always equal to the sum of the parts on the other side of it.

Curve lines are distinguished into *algebraical* or *geometrical*, and *transcendental* or *mechanical*.

*Algebraical* or *Geometrical CURVES*, are those in which the relation of the absciss AP, to the ordinates PM, can be expressed by a common algebraic equation

And *Transcendental* or *Mechanical Curves*, are such as cannot be so defined or expressed by an algebraical equation. See *TRANSCENDENTAL Curve*.

Thus, suppose, for instance, the curve be the circle; and that the radius AC =  $r$ , the absciss AP =  $x$ , and the ordinate PM =  $y$ ; then, because the nature of the circle is such, that the rectangle AP  $\times$  PB is always = PM<sup>2</sup>, therefore the equation is  $x \cdot 2r - x = y^2$ , or  $2rx - x^2 = y^2$ , defining this curve, which is therefore an algebraical or geometrical line. Or, suppose CP =  $x$ ; then is CM<sup>2</sup> - CP<sup>2</sup> = PM<sup>2</sup>, that is  $r^2 - x^2 = y^2$ ; which is another form of the equation of the curve.

The doctrine of curve lines in general, as expressed by algebraical equations, was first introduced by Des Cartes, who called algebraical curves geometrical ones; as admitting none else into the construction of problems, nor consequently into geometry. But Newton, and after him Leibnitz and Wolfius, are of another opinion; and think, that in the construction of a problem, one curve is not to be preferred to another for its being defined by a more simple equation, but for its being more easily described.

Algebraical or geometrical lines are best distinguished into orders according to the number of dimensions of the equation expressing the relation between its ordinates and absciss, or, which is the same thing, according to the number of points in which they may be cut by a right line. And curves of the same kind or order, are those whose equations rise to the same dimension. Hence, of the first order, there is the right line only; of the 2d order of lines, or the first order of curves, are the circle and conic sections, being 4 species only, viz,  $dx - x^2 = y^2$  the circle,  $\frac{c^2}{a^2} \cdot dx - x^2 = y^2$  the

ellipse,  $\frac{c^2}{a^2} \cdot dx + x^2 = y^2$  the hyperbola, and  $dx = y^2$

the parabola: the lines of the 3d order, or curves of the 2d order, are expressed by an equation of the 3d degree, having three roots; and so on. Of these lines of the 3d order, Newton wrote an express treatise, under the title of *Enumeratio Linearum Tertii Ordinis*, shewing their distinctive characters and properties, to the number of 72 different species of curves: but Mr. Stirling afterwards added four more to that number; and Mr. Nic. Bernoulli and Mr. Stone added two more.

*Curves of the 2d and other higher kinds*, Newton observes, have parts and properties similar to those of the 1st kind: Thus, as the conic sections have diameters and axes; the lines bisected by these are ordinates; and the intersection of the curve and diameter, the vertex: so, in curves of the 2d kind, any two parallel right lines being drawn to meet the curve in 3 points; a right line cutting these parallels so, as that the sum of the two parts between the secant and the curve on one side, is equal to the 3d part terminated by the curve on the other side, will cut, in the same manner, all other right lines parallel to these, and that meet the curve in three points, that is, so as that the sum of the two parts on one side, will still be equal to the 3d part on the other side. These three parts therefore thus equal, may be called *ordinates*, or *applicates*; the cutting line, the *diameter*;

meter;



*meter*; and where it cuts the ordinates at right angles, the *axis*; the intersection of the diameter and the curve, the *vertex*; and the concurrence of two diameters, the *centre*; also the concurrence of all the diameters, the *common* or *general centre*.

Again, as an hyperbola of the first kind has two asymptotes; that of the 2d has 3; that of the 3d has 4; &c: and as the parts of any right line between the conic hyperbola and its two asymptotes, are equal on either side; so, in hyperbolas of the 2d kind, any right line cutting the curve and its three asymptotes in three points; the sum of the two parts of that right line, extended from any two asymptotes, the same way, to two points of the curve, will be equal to the 3d part extended from the 3d asymptote, the contrary way, to the 3d point of the curve.

Again, as in the conic sections that are not parabolical, the square of an ordinate, i. e. the rectangle of the ordinates drawn on the contrary sides of the diameter, is to the rectangle of the parts of the diameter terminated at the vertices of an ellipse or hyperbola; in the same proportion as a given line called the *latus rectum*, is to that part of the diameter which lies between the vertices, and called the *latus transversum*: so, in curves of the 2d kind, not parabolical, the parallelopiped under three ordinates, is to the parallelopiped under the parts of the diameter cut off at the ordinates and the three vertices of the figure, in a given ratio: in which, if there be taken three right lines situate at the three parts of the diameter between the vertices of the figure, each to each; then these three right lines may be called the *latera recta* of the figure; and the parts of the diameter between the vertices, the *latera transversa*.

And, as in a conic parabola, which has only one vertex to one and the same diameter, the rectangle under the ordinates is equal to the rectangle under the part of the diameter cut off at the ordinates and vertex, and a given right line called the *latus rectum*: so, in curves of the 2d kind, which have only two vertices to the same diameter, the parallelopiped under three ordinates, is equal to the parallelopiped under two parts of the diameter cut off at the ordinates and the two vertices, and a given right line, which may therefore be called the *latus transversum*.

Further, as in the conic sections, where two parallels, terminated on each side by a curve, are cut by two other parallels terminated on each side by a curve, the 1st by the 3d, and the 2d by the 4th; the rectangle of the parts of the 1st is to the rectangle of the parts of the 3d, as that of the 2d is to that of the 4th: so, when four such right lines occur in a curve of the 2d kind, each in three points; the parallelopiped of the parts of the 1st, will be to that of the parts of the 3d, as that of the 2d to that of the 4th.

Lastly, the legs of curves, both of the 1st, 2d, and higher kinds, are either of the parabolic or hyperbolic kind: an hyperbolic leg being that which approaches infinitely towards some asymptote; and a parabolic one, that which has no asymptote.

These legs are best distinguished by their tangents; for, if the point of contact go off to an infinite distance, the tangent of the hyperbolic leg will coincide with the asymptote; and that of the parabolic leg, recede

infinitely, and vanish. Therefore the asymptote of any leg is found, by seeking the tangent of that leg to a point infinitely distant; and the direction of an infinite leg is found, by seeking the position of a right line parallel to the tangent, where the point of contact is infinitely remote, for this line tends that way towards which the infinite leg is directed.

#### *Reduction of CURVES of the 2d kind.*

Newton reduces all curves of the 2d kind to four cases of equations, expressing the relation between the ordinate and absciss, viz,

in the 1st case,  $xy^2 + cy = ax^3 + bx^2 + cx + d$ ;

in the 2d,  $- \quad xy = ax^3 + bx^2 + cx + d$ ;

in the 3d,  $\cdot \quad y^2 = ax^3 + bx^2 + cx + d$ ;

in the 4th,  $- \quad y = ax^3 + bx^2 + cx + d$ .

See Newton's *Enumeratio*, sect. 3; and Stirling's *Linæ*, &c, pa. 83.

#### *Enumeration of the CURVES of the 2d kind.*

Under these four cases, the author brings a great number of different forms of curves, to which he gives different names. An hyperbola lying wholly within the angle of the asymptotes, like a conic hyperbola, he calls an *inscribed hyperbola*; that which cuts the asymptotes, and contains the parts cut off within its own periphery, a *circumscribed hyperbola*; that which has one of its infinite legs inscribed and the other circumscribed, he calls *ambigenal*; that whose legs look towards each other, and are directed the same way, *converging*; that where they look contrary ways, *diverging*; that where they are convex different ways, *cross-legged*; that applied to its asymptote with a concave vertex, and diverging legs, *conchoidal*; that which cuts its asymptote with contrary flexures, and is produced each way into contrary legs, *anguineous*, or *snake-like*; that which cuts its conjugate across, *cruciform*; that which returning around cuts itself, *nodated*; that whose parts concur in the angle of contact, and there terminate, *cuspidated*; that whose conjugate is oval, and infinitely small, i. e. a point, *pointed*; that which, from the impossibility of two roots, is without either oval, node, cusp, or point, *pure*. And in the same manner he denominates a parabola *converging*, *diverging*, *cruciform*, &c. Also when the number of hyperbolic legs exceeds that of the conic hyperbola, that is more than two, he calls the hyperbola *redundant*.

Under those 4 cases the author enumerates 72 different curves: of these, 9 are redundant hyperbolas, without diameters, having three asymptotes including a triangle; the first consisting of three hyperbolas, one inscribed, another circumscribed, and the third ambigenal, with an oval; the 2d, nodated; the 3d, cuspidated; the 4th, pointed; the 5th and 6th, pure; the 7th and 8th, cruciform; the 9th or last, anguineal. There are 12 redundant hyperbolas, having only one diameter: the 1st, oval; the 2d, nodated; the 3d, cuspidated; the 4th, pointed; 5th, 6th, 7th, and 8th, pure; the 9th and 10th, cruciform; the 11th and 12th, conchoidal. And to this class Stirling adds 2 more. There are 2 redundant hyperbolas, with three diameters. There are 9 redundant hyperbolas, with three asymptotes converging to a common point; the 1st being formed of the 5th and 6th redundant parabolas, whose



whose asymptotes include a triangle; the 2d formed of the 7th and 8th; the 3d and 4th, of the 9th; the 5th is formed of the 5th and 7th of the redundant hyperbolas, with one diameter; the 6th, of the 6th and 7th; the 7th, of the 8th and 9th; the 8th, of the 10th and 11th; the 9th, of the 12th and 13th: all which conversions are effected, by diminishing the triangle comprehended between the asymptotes, till it vanish into a point.

Six are defective parabolas, having no diameters: the 1st, oval; the 2d, nodated; the 3d, cuspidated; the 4th, pointed; the 5th, pure; &c.

Seven are defective hyperbolas, having diameters: the 1st and 2d, conchoidal, with an oval; the 3d, nodated; the 4th, cuspidated, which is the cissoid of the ancients; the 5th and 6th, pointed; the 7th, pure.

Seven are parabolic hyperbolas, having diameters: the 1st, oval; the 2d, nodated; the 3d, cuspidated; the 4th, pointed; the 5th, pure; the 6th, cruciform; the 7th, anguineous.

Four are parabolic hyperbolas: four are hyperbolisms of the hyperbola: three, hyperbolas of the ellipsis: two, hyperbolisms of the parabola.

Six are diverging parabolas; one, a trident; the 2d, oval; the 3d, nodated; the 4th, pointed; the 5th, cuspidated (which is Neil's parabola, usually called the semi-cubical parabola); the 6th, pure.

Lastly, one, commonly called the cubical parabola.

Mr. Stirling and Mr. Stone have shewn that this enumeration is imperfect, the former having added four new species of curves to the number, and the latter two, or rather these two were first noticed by Mr. Nic. Bernoulli. Also Mr. Murdoch and Mr. Geo. Sanderfon have found some new species; though some persons dispute the reality of them. See the *Genesis Curvarum per umbras*, and the *Ladies' Diary* 1788 and 1789, the prize question.

*Organical Description of CURVES.*—Sir Isaac Newton shews that curves may be generated by shadows. He says, if upon an infinite plane, illuminated from a lucid point, the shadows of figures be projected; the shadows of the conic sections will always be conic sections; those of the curves of the 2d kind, will always be curves of the 2d kind; those of the curves of the 3d kind, will always be curves of the 3d kind; and so on ad infinitum.

And, like as the projected shadow of a circle generates all the conic sections, so the 5 diverging parabolas, by their shadows, will generate and exhibit all the rest of the curves of the 2d kind: and thus some of the most simple curves of the other kinds may be found, which will form, by their shadows upon a plane, projected from a lucid point, all the other curves of that same kind. And in the French Memoirs may be seen a demonstration of this projection, with a specimen of a few of the curves of the 2d order, which may be generated by a plane cutting a solid formed from the motion of an infinite right line along a diverging parabola, having an oval, always passing through a given or fixed point above the plane of that parabola. The above method of Newton has also been pursued and illustrated with great elegance by Mr. Murdoch, in his treatise entitled *Newtoni Genesis Curvarum per umbras, seu Perspectiva Universalis Elementa*.

Mr. Maclaurin, in his *Geometria Organica*, shews how to describe several of the species of curves of the 2d order, especially those having a double point, by the motion of right lines and angles: but a good commodious description by a continued motion of those curves which have no double point, is ranked by Newton among the most difficult problems. Newton gives also other methods of description, by lines or angles revolving about given poles; and Mr. Brackenridge has given a general method of describing curves, by the intersection of right lines moving about points in a given plane. See *Philos. Transf.* No. 437, or *Abr.* vol. 8, pa. 58; and some particular cases are demonstrated in his *Exerc. Geometrica de Curvarum Descriptione*.

*CURVES above the 2d Order.* The number of species in the higher orders of curves increase amazingly, those of the 3d order only it is thought amounting to some thousands, all comprehended under the following ten particular equations,

$$\text{viz, 1. } y^4 + fx^2y^2 + gxy^3 + hx^2y + iy^2 + hxy + ly,$$

$$\text{or 2. } y^4 + fxy^3 + gx^2y + hxy^2 + ixy + ky,$$

$$\text{or 3. } x^2y^2 + fy^3 + gx^2y + by^3 + ky,$$

$$\text{or 4. } x^2y^2 + fy^3 + gy^2 + hxy + iy,$$

$$\text{or 5. } y^3 + fxy^2 + gx^2y + hy,$$

$$\text{or 6. } y^3 + fxy^2 + gxy + hy$$

$$= ax^4 + bx^3 + cx^2 + dx + e;$$

$$\text{or 7. } y^4 + ex^3y + fxy^3 + gxy^2 + hy^2 + ixy + ky,$$

$$\text{or 8. } x^3y + exy^3 + fx^2y + gy^2 + hxy + iy,$$

$$\text{or 9. } x^3y + ey^3 + fxy^2 + gxy + hy,$$

$$\text{or 10. } x^3y + cy^3 + fy^2 + gxy + hy,$$

$$= ax^3 + bx^2 + cx + d.$$

Those who wish to see how far this doctrine has been advanced, with regard to curves of the higher orders, as well as those of the 1st and 2d orders, may consult Mr. Maclaurin's *Geometria Organica*, and Brackenridge's *Exerc. Geom.*

All geometrical lines of the odd orders, viz, the 3d, 5th, 7th, &c, have at least one leg running on infinitely; because all equations of the odd dimensions have at least one real root. But vast numbers of the lines of the even orders are only ovals; among which there are several having very pretty figures, some being like single hearts, some double ones, some resembling fiddles, and others again single knots, double knots, &c.

Two geometrical lines of any order will cut one another in as many points, as are denoted by the product of the two numbers expressing those orders.

The theory of curves forms a considerable branch of the mathematical sciences. Those who are curious of advancing beyond the knowledge of the circle and the conic sections, and to consider geometrical curves of a higher nature, and in a general view, will do well to study Cramer's *Introduction à l'Analyse des Lignes Courbes Algebraïques*, which the learned and ingenious author composed for the use of beginners. There is an excellent posthumous piece too of Maclaurin's, printed as an Appendix to his *Algebra*, and entitled *De Linearum Geometricarum Proprietatibus Generalibus*. The same author, at a very early age, gave a remarkable specimen of his genius and knowledge in his *Geometria Organica*; and he carried these speculations farther afterwards, as may be seen in the theorems he



has given in the *Philos. Trans.* See *Abr.* vol. 8, pa. 62. Other writings on this subject, beside the *Treatises* on the Conic Sections, are *Archimedes de Spiralibus*; *Des Cartes Geometria*; *Dr. Barrow's Lectiones Geometricæ*; *Newton's Enumeratio Linearum Tertii Ordinis*; *Stirling's Illustratio Tractatus Newtoni de Lineis Tertii Ordinis*; *Maclaurin's Geometria Organica*; *Brackenridge's Descriptio Linearum Curvarum*; *M. De Gua's Usages de l'Analyse de Des Cartes*; beside many other Tracts on Curves in the *Memoirs* of several Academies &c.

*Use of CURVES in the Construction of Equations.* One great use of curves in Geometry is, by means of their interfections, to give the solution of problems. See CONSTRUCTION.

Suppose, *ex. gr.* it were required to construct the following equation of 9 dimensions,

$x^9 + bx^7 + cx^6 + dx^5 + ex^4 + m + f.x^3 + gx^2 + hx + k = 0$ : assume the equation to a cubic parabola  $x^3 = y$ ; then, by writing  $y$  for  $x^3$ , the given equation will become  $y^3 + bxy^2 + cy^2 + dx^2y + exy + my + fx^3 + gx^2 + hx + k = 0$ ; an equation to another curve of the 2d kind, where  $m$  or  $f$  may be assumed  $= 0$  or any thing else: and by the descriptions and interfections of these curves will be given the roots of the equation to be constructed. It is sufficient to describe the cubic parabola once. When the equation to be constructed, by omitting the two last terms  $bx$  and  $k$ , is reduced to 7 dimensions; the other curve, by expunging  $m$ , will have the double point in the beginning of the absciss, and may be easily described as above: If it be reduced to 6 dimensions, by omitting the last three terms,  $gx^2 + bx + k$ ; the other curve, by expunging  $f$ , will become a conic section. And if, by omitting the last three terms, the equation be reduced to three dimensions, we shall fall upon Wallis's construction by the cubic parabola and right line.

*Rectification, Inflection, Quadrature, &c of CURVES.* See the respective terms.

*CURVE of a Double Curvature*, is such a curve as has not all its parts in the same plane.

M. Clairaut has published an ingenious treatise on curves of a double curvature. See his *Recherches sur les Courbes à Double Courbure*. Mr. Euler has also treated this subject in the Appendix to his *Analysis Infinitorum*, vol. 2, pa. 323.

*Family of CURVES*, is an assemblage of several curves of different kinds, all defined by the same equation of an indeterminate degree; but differently, according to the diversity of their kind. For example, Suppose an equation of an indeterminate degree,  $a^{m-1}x = y^m$ : if  $m = 2$ , then will  $ax = y^2$ ; if  $m = 3$ , then will  $a^2x = y^3$ ; if  $m = 4$ , then is  $a^3x = y^4$ ; &c: all which curves are said to be of the same family or tribe.

The equations by which the families of curves are defined, are not to be confounded with the transcendental ones: for though with regard to the whole family, they be of an indeterminate degree; yet with respect to each several curve of the family, they are determinate; whereas transcendental equations are of an indefinite degree with respect to the same curve.

All Algebraical curves therefore compose a certain family, consisting of innumerable others, each of which comprehends infinite kinds. For the equations by

which curves are defined involve only products, either of powers of the absciss and ordinates by constant coefficients; or of powers of the absciss by powers of the ordinates; or of constant, pure, and simple quantities by one another. Moreover, every equation to a curve may have 0 for one member or side of it; for example,  $ax = y^2$ , by transposition becomes  $ax - y^2 = 0$ . Therefore the equation for all algebraic curves will be

$$\left. \begin{aligned} ay^m + bxy^{m-1} + nx^2y^{m-2} &\&c - - - fy^m \\ + fy^{m-1} + kxy^{m-2} \\ + qy^{m-2} \end{aligned} \right\} = 0.$$

*Catacaustic, and Diacaustic CURVES.* See CATACAUSTIC, and DIACAUSTIC.

*Exponential CURVE*, is that which is defined by an exponential equation, as  $ax^x = y$ , &c.

*CURVES by the Light*, or *COURBES à la Lumière*, a name given to certain curves by M. Kurdwanowski, a Polish gentleman. He observed that any line, straight or curved, exposed to the action of a luminous point, received the light differently in its different parts, according to their distance from the light. These different effects of the light upon each point of the line, may be represented by the ordinates of some curve, which will vary precisely with these effects. Priestley's *Hist. of Vision*, pa. 752.


*Logarithmic CURVE.* See LOGARITHMIC Curve.

*CURVE Reflectoire*, so called because it is the appearance of the plane bottom of a basin covered with water, to an eye perpendicularly over it. In this position, the bottom of the basin will appear to rise upwards, from the centre outwards; but the curvature will be less and less, and at last the surface of the water will be an asymptote to it. M. Mairan, who first conceived this idea from the phenomena of light, found also several kinds of these curves; and he gives a geometrical deduction of their properties, shewing their analogy to caustics by refraction. *Mem. Ac.* 1740; Priestley's *Hist. of Vision*, pa. 752.

*Radical CURVES*, a name given by some authors to curves of the spiral kind, whose ordinates, if they may be so called, do all terminate in the centre of the including circle; and appear like so many radii of that circle: whence the name.

*Regular CURVES*, are such as have their curvature turning regularly and continually the same way; in opposition to such as bend contrary ways, by having points of contrary flexure, which are called irregular curves.

*Characteristic Triangle of a CURVE*, is the differential or elementary right-angled triangle whose three sides are, the fluxions of the absciss, ordinate, and curve; the fluxion of the curve being the hypotenuse. So, if  $pq$  be parallel to, and indefinitely near to the ordinate  $PQ$ , and  $Qr$  parallel to the absciss  $AP$ ; then  $Qr$  is the fluxion of the absciss  $AP$ , and  $qr$  the fluxion of the ordinate  $PQ$ , and  $Qq$  the fluxion of the curve  $AQ$ ; hence the elementary triangle  $Qqr$  is the characteristic triangle of the curve  $AQ$ ; and the three sides are  $\dot{x}$ ,  $\dot{y}$ ,  $\dot{z}$ ; in which  $\dot{x}^2 + \dot{y}^2 = \dot{z}^2$ .



*CURVILINEAR Angle, Figure, Superficies, &c,*  
Z z are



are such as are formed or bounded by curves; in opposition to rectilinear ones, which are formed by straight lines or planes.

CUSP, in Astronomy, is used to express the points or horns of the moon, or other luminary.

CUSP, in Astrology, is used for the 1st point of each of the twelve houses, in a figure or scheme of the heavens. See HOUSE.

CUSP, in the Higher Geometry, is used for the point or corner formed by two parts of a curve meeting and terminating there. See CURVE.

CUSPIDATED *Hyperbola*, &c. See CURVE.

CUT-BASTION. See BASTION.

CUVETTE, or CUNETTE, in Fortification, is a kind of ditch within a ditch, being a pretty deep trench, about four fathoms broad, sunk and running along the middle of the great dry ditch, to hold water; serving both to keep off the enemy, and prevent him from mining.

CYCLE, a certain period or series of numbers proceeding orderly from first to last, then returning again to the first, and so circulating perpetually.

Cycles have chiefly arisen from the incommensurability of the revolutions of the earth and celestial bodies to one another. The apparent revolution of the sun about the earth, has been arbitrarily divided into 24 hours, which is the basis or foundation of all our mensuration of time, whether days, years, &c. But neither the annual motion of the sun, nor that of the other heavenly bodies, can be measured exactly, and without any remainder, by hours, or their multiples. That of the sun, for example, is 365 days 5 hours 49 minutes nearly; that of the moon, 29 days 12 hours 44 minutes nearly.

Hence, to swallow up these fractions in whole numbers, and yet in numbers which only express days and years, cycles have been invented; which, comprehending several revolutions of the same body, replace it, after a certain number of years, in the same points of the heaven whence it first departed; or, which is the same thing, in the same place of the civil calendar.

There are various cycles; as, the cycle of Indiction, the cycle of the moon, the cycle of the sun, &c.

CYCLE of *Indiction*, is a series of 15 years, returning constantly around like the other cycles; and commenced from the third year before Christ; whence it happens that if 3 be added to any given year of Christ, and the sum be divided by 15, what remains is the year of the indiction. See INDICATION.

CYCLE of the *Moon*, or the *Lunar Cycle*, is a period of 19 years; in which time the new and full moons return to the same day of the Julian year. See CALIPPIC.

This cycle is also called the *Metonic period* or *cycle*, from its inventor Meton, the Athenian; and also the *Golden Number*, from its excellent use in the calendar: though, properly speaking, the golden number is rather the particular number which shews the year of the lunar cycle, which any given year is in. This cycle of the moon only holds true for  $310\frac{7}{10}$  years: for, though the new moons do return to the same day after 19 years; yet not to the same time of the day, but near an hour and a half sooner; an error which in  $310\frac{7}{10}$

years amounts to an entire day. Yet those employed in reforming the calendar went on a supposition that the lunations return precisely from 19 years to 19 years for ever.

The use of this cycle, in the ancient calendar, is to shew the new moon of each year, and the time of Easter. In the new one, it only serves to find the Epacts; which shew, in either calendar, that the new moon falls 11 days too late.

As the Orientals began the use of this cycle at the time of the Council of Nice in 325, they assumed, that the first year of the cycle the paschal new moon fell on the 13th of March: on which account the lunar cycle 3 fell on the 1st of January in the third year.

The Occidentals, on the contrary, placed the number 1 to the 1st of January, which occasioned a considerable difference in the time of Easter. Hence, Dionysius Exiguus, on framing a new calendar, persuaded the Christians of the west to save the difference, and come into the practice of the church of Alexandria.

To find the Year of the Lunar Cycle, is to find the golden number. See GOLDEN-Number.

CYCLE of the *Sun*, or *Solar Cycle*, is a period or revolution of 28 years; beginning with 1, and ending with 28; which elapsed, the Dominical or Sunday-letters, and those that express the other feasts, &c, return into their former place, and proceed in the same order as before. The days of the month return again to the same days of the week; the sun's place to the same signs and degrees of the ecliptic on the same months and days, so as not to differ one degree in a hundred years; and the leap years begin the same course with respect to the days of the week on which the days of the month fall.

This is called the cycle of the *sun*, or the *solar cycle*, not from any regard to the sun's course, which has no concern in it; but from *Sunday*, anciently called *dies solis*, the *sun's day*; as the dominical or sunday letter is chiefly sought for from this revolution.

The reformation of the calendar under pope Gregory the 13th, occasioned a considerable alteration of this cycle: In the Gregorian calendar, the solar cycle is not constant and perpetual; because every 4th secular year is common; whereas, in the Julian, it is bissextile. The epoch, or beginning of the solar cycle, both Julian and Gregorian, is the 9th year before Christ. And therefore,

To find the Cycle of the Sun for any given year: add 9 to the number given, and divide the sum by 28; the remainder will be the number of the cycle, and the quotient the number of revolutions since Christ. If there be no remainder, it will be the 28th or last year of the cycle.

CYCLE of the Sun, with the correspondent Sunday letters, in Julian Years.

1	GF	5	BA	9	DC	13	FE	17	AG	21	CB	25	ED
2	E	6	G	10	B	14	D	18	F	22	A	26	C
3	D	7	F	11	A	15	C	19	E	23	G	27	B
4	C	8	E	12	G	16	B	20	D	24	F	28	A




CYCLE of the Sun, and Sunday Letters, from the Gregorian year 1700 to the year 1800.

1 DC	5 FE	9 AG	13 CB	17 ED	21 GF	25 BA
2 B	6 D	10 F	14 A	18 C	22 E	26 G
3 A	7 C	11 E	15 G	19 B	23 D	27 F
4 G	8 B	12 D	16 F	20 A	24 C	28 E

*Great Pascal CYCLE*, is another name for the Victorian or Dionysian Period. Which see.

**CYCLOID**, or **TROCHOID**, a mechanical or transcendental curve, which is thus generated: Suppose a wheel, or a circle, *AE*, to roll along a straight line *AB*, beginning at the point *A*, and ending at *B*, where it has completed just one revolution, thereby measuring out a right line *AB* exactly equal to the circumference of the generating circle *AE*, whilst a nail or point *A* in the circumference of the wheel, or circle, traces out or describes a curvilinear path *ADB*; then this curve *ADB* is the cycloid, or trochoid.



Schooten, in his *Commentary on Des Cartes*, says that Des Cartes first conceived the notion of this elegant curve, and after him it was first published by Father Merfenne, in the year 1615. But Torricelli, in the *Appendix de Dimensione Cycloidis*, at the end of his treatise *De Dimensione Parabolæ*, published 1644, says that this curve was considered and named a cycloid, by his predecessors, and particularly by Galileo about 45 years before, i. e. about 1599. And Dr. Wallis shews that it is of a much older standing, having been known to Bovilli about the year 1500, and even considered by cardinal Cusanus much earlier, viz, before the year 1451. *Philos. Transf. Abr.* vol. 1, pa. 116. It would seem however that Torricelli's was the first regular treatise on the Cycloid; though several particular properties of it might be known prior to his work. He first shewed, that the cycloidal space is equal to triple the generating circle, (though Pascal contends that Roberval shewed this): also that the solid generated by the rotation of that space about its base, is to the circumscribing cylinder, as 5 to 8: about the tangent parallel to the base, as 7 to 8: about the tangent parallel to the axis, as 3 to 4: &c.

Honoratus Fabri, in his *Synopsis Geom.* has a short treatise on the cycloid, containing demonstrations of the above, and many other theorems concerning the centres of gravity of the cycloidal space, &c; which he says he found out before the year 1658.


From the preface to Dr. Wallis's treatise on the cycloid we learn, that, in the year 1658, M. Pascal publicly proposed at Paris, under the name of D'Ettonville, the two following problems as a challenge, to

be solved by the mathematicians of Europe, with a reward of 20 pistoles for the solution: viz, to find the area of any segment of the cycloid, cut off by a right line parallel to the base; also the content of the solid generated by the rotation of the same about the axis, and about the base of that segment. This challenge set the Doctor upon writing that treatise upon the cycloid, which is a much better and compleater piece than had been given before upon this curve. He here gives the curve surfaces of the solids generated by the rotation of the cycloidal space about its axis, and about its base, with determinations of the centres of gravity, &c. He here asserts too, that Sir Christopher Wren, in 1658, was the first who found out a right line equal to the curve of the cycloid; and Mr. Huygens, in his *Horolog. Oscillat.* says that he himself was the first inventor of the segment of a cycloidal space, cut off by a right line parallel to the base at the distance of  $\frac{1}{4}$  the axis of the curve from the centre, being equal to a rectilinear space, viz, to a regular hexagon inscribed in the generating circle; the demonstration of which may be seen in Wallis's treatise.

Several other authors have spoken or treated of the cycloid: as Pascal, in his treatise, under the name of D'Ettonville: Schooten in his *Commentary on Des Cartes's Geometry*, near the end of the 2d book; M. Reinau, in his *Analyse Démontrée*, tom. 2, pa. 595: also Newton, Leibnitz, de la Loubere, Roberval, Des Cartes, Wren, Fabri, the Bernoulli's, De la Hire, Cotes, &c, &c.

*Properties of the CYCLOID.*—The circle *AE*, by whose revolution the cycloid is traced out, is called the *generating circle*; the line *AB*, which is equal to the circumference of the circle, is the *base* of the cycloid; and the perpendicular *DC* on the middle of the base, is its *axis*. The properties of the cycloid are among the most beautiful and useful of all curve lines: some of the most remarkable of which are as follow:

1. The circular arc *DG* = the line *GH* parallel to *AB*.
2. The semicircumf. *DGC* = the semibase *AC*.
3. The arc *DH* = double the chord *DG*.
4. The arc *DA* = double the diam. *DC*.
5. The tang. *HI* is parallel to the chord *DG*.
6. The space *ADBA* = triple the circle *AE* or *CGD* &c.
7. The space *ADGCA* = the same circle *AE*, &c.



8. A body falls through any arc *KL* of a cycloid reversed, in the same time, whether that arc be great or small; that is, from any point *L*, to the lowest point *K*, which is the vertex reversed: and that time




is to the time of falling perpendicularly through the axis MK, as the semicircumference of a circle is to its diameter, or as 3.1416 to 2. And hence it follows that, if a pendulum be made to vibrate in the arc LKN of a cycloid, all the vibrations will be performed in the same time.


9. The evolute of a cycloid, is another equal cycloid. So that if two equal femicycloids OP, OQ, be joined at O, so that OM be = MK the diameter of the generating circle, and the string of a pendulum hung up at O, having its length = OK or = the curve OP; then, by plying the string round the curve OP, to which it is equal, and then the ball let go, it will describe, and vibrate in the other cycloid PKQ.

10. The cycloid is the curve of swiftest descent: or a heavy body will fall from one given point to another, by the way of the arc of a cycloid passing through those two points, in a less time, than by any other rout. See the Works of James and John Bernoulli for many other curious properties concerning the descents in cycloids, &c.

CYCLOIDS are also either *curtate* or *prolate*.



CYCLOID, *Curtate*, or *contracted*, is the path described by some point without the circle, while the circumference rolls along a straight line; and a



CYCLOID, *Prolate*, or *Inflected*, is in like manner the path of some point taken within the generating circle.

Thus, if, while the circle rolls along the line AB, the point R be taken without the circle, it will describe or trace out the curtate or contracted cycloid RST; but the point being taken within the circle, it will describe the prolate or inflected cycloid RVW.

These two curves were both noticed by Torricelli and Schooten, and more fully treated of by Wallis, in his 'Treatise on the Cycloid, printed at Oxford in 1659; where he shews that these have properties similar to the first or primary cycloid; only the last of these is a curve having a point of inflection, and the other crossing itself, and forming a node.

By continuing the motion of the wheel, or circle, so as to describe a right line equal to the generating circumference several times repeated, there will be produced as many repetitions of the cycloids, which so united together will appear as in these figures following:

Curtate Cycloid.



Common Cycloid.



Inflected Cycloid.




CYGNUS, the *Swan*, a constellation of the northern hemisphere, being one of the 48 old ones, and fabled by the Greeks to be the swan, under the form of which Jupiter deceived Leda or Nemesis, from which embrace sprung the beauteous Helen.

The stars in the constellation Cygnus, in Ptolemy's catalogue are 19, in Tycho's 18, in Hevelius's 47, and in the Britannic catalogue 81.

CYLINDER, a solid having two equal circular ends, and every plane section parallel to the ends a circle equal to them also.

The cylinder may be conceived to be thus generated:



Suppose two parallel circles AB and CD, and a right line carried continually round them, always parallel to itself; this line will describe the curve surface of a cylinder, ABDC, of which the two parallel circles AB and CD form the two ends. When the line, or sides is perpendicular to the ends, the cylinder is a right or perpendicular one; otherwise it is oblique.

Or the right cylinder may be conceived to be generated by the rotation of a rectangle about one of its sides. The *axis* of the cylinder is the line connecting the centres of its two parallel circular ends; and is equal to the altitude of the cylinder when this is a right one, but exceeds the altitude in the oblique cylinder, in the proportion of radius to the sine of the angle of its inclination to the base.

The convex surface of a cylinder is equal to the product of the axis multiplied by the circumference of its base.

The solidity of a cylinder is equal to the area of its base multiplied by its perpendicular altitude.

Cylinders of equal bases and altitudes, are equal.

Cylinders are to each other, as the product of their bases and altitudes. And equal cylinders have their bases reciprocally as their altitudes.

A cylinder is to its inscribed sphere, or spheroid, as 3 to 2; and to its inscribed cone as 3 to 1.

The



The oblique plane sections of a cylinder, are ellipses; but all the sections parallel to the ends, are circles.

For the surfaces and solidities of the ungulas, or oblique slices, of a cylinder, see my *Mensuration*, pa. 218, 2d edition.

CYLINDRICAL, pertaining to a cylinder.

CYLINDROID, a solid resembling the figure of a cylinder; but differing from it as having ellipses for its ends or bases, instead of circles, in the cylinder.

In the cylindroid, the solidity and curve superficies are found the same way as those of the cylinder; viz,

by multiplying the circumference of the base by the length or axis, for the surface; and the area of the base by the altitude, for the solidity.

CYMATIUM, CIMATIUM, or CIMA, in Architecture, a member, or moulding of the cornice, whose profile is waved; i. e. concave at top, and convex at bottom.

CYNOSURA, a name given by the Greeks, to urfa minor, or little bear, otherwise called Charles's wain; the star in the extremity of the tail being called the pole star.

CYPHER, or nought. See CIPHER.

## D.

### D A R

**D**ACTYLOLOGY, the art of counting or numbering by the fingers.—The rule is this: the left thumb is reckoned 1, the index or fore-finger 2, and so on to the right thumb, which is the tenth or last, and consequently is denoted by the cipher 0.

DADO, that part in the middle of the pedestal of a column &c, between its base and cornice.

DAILY, in Astronomy. See DIURNAL.

DARCY (COUNT), an ingenious philosopher and mathematician, was born in Ireland in 1725; but his friends being attached to the Stuart family, he was sent to France, at 14 years of age, where he spent the rest of his life. Being put under the care of the celebrated Clairaut, he improved so rapidly in the mathematics, that at 17 years of age he gave a new solution of the problem concerning the curve of equal pressure in a resisting medium. This was followed the year after by a determination of the curve described by a heavy body, sliding by its own weight along a moveable plane, at the same time that the pressure of the body causes a horizontal motion in the plane. Darcy served in the war of 1744, and was taken prisoner by the English: and yet, during the course of the war he gave two memoirs to the academy; the first of these contained a general principle in mechanics, that of *the preservation of the rotatory motion*; a principle which he again brought forward in 1750, by the name of *the principle of the preservation of action*.

In 1760, Darcy published *An Essay on Artillery*, containing some curious experiments on the charges of gunpowder, &c, &c, and improvements on those of the ingenious Robins; a kind of experiments which our author carried on occasionally to the end of his life.

In 1765, he published his *Memoir on the Duration of the Sensation of Sight*, the most ingenious of his works: the result of these researches was, that a body may sometimes pass by our eyes without being seen, or marking its presence, otherwise than by weakening the brightness of the object it covers.

All Darcy's works bear the character which results

### D A T

from the union of genius and philosophy; but as he measured every thing upon the largest scale, and required extreme accuracy in experiment, neither his time, fortune, nor avocations, allowed him to execute more than a very small part of what he projected. In his disposition, he was amiable, spirited, lively, and a lover of independence, a passion to which he nobly sacrificed, even in the midst of literary society.—He died of a cholera morbus in 1779, at 54 years of age.

Darcy was admitted of the French academy in 1749, and was made pensioner-geometrician in 1770.—His essays, printed in the *Memoirs of the Academy of Sciences*, are various and very ingenious, and are contained in the volumes for the years 1742, 1747, 1749, 1750, 1751, 1752, 1753, 1754, 1758, 1759, 1760, 1765, and in tom. I, of the *Savans Etrangers*.

DARK Chamber. See CAMERA Obscura.

DARK Tent, a portable camera obscura, made somewhat like a desk, and fitted with optic glasses, to take prospects of landscapes, buildings, &c.

DATA, in General Mathematics, denote certain things or quantities, supposed given or known, from which other quantities are discovered that were unknown, or sought. A problem or question usually consists of two parts, *data* and *quesita*.

Euclid has an express and excellent treatise of *Data*; in which he uses the word for such spaces, lines, angles, &c, as are given; or to which others can be found equal.

Euclid's *Data* is the first in order of the books that have been written by the ancient geometricians, to facilitate and promote the method of resolution or analysis. In general a thing is said to be given which is either actually exhibited, or can be found out, that is, which is either known by hypothesis, or that can be demonstrated to be known: and the propositions in the book of Euclid's *Data* shew what things can be found out or known, from those that by hypothesis are already known: so that in the analysis or investigation of a problem, from the things that are laid down as given or



or known, by the help of these propositions, it is demonstrated that other things are given, and from these last that others again are given, and so on, till it is demonstrated that that which was proposed to be found out in the problem is given; and when this is done, the problem is solved, and its composition is made and derived from the compositions of the Data which were employed in the analysis. And thus the Data of Euclid are of the most general and necessary use in the solution of problems of every kind.

Marinus, at the end of his Preface to the Data, is mistaken in asserting that Euclid has not used the synthetical, but the analytical method in delivering them: for, though in the analysis of a theorem, the thing to be demonstrated is assumed in the analysis; yet, in the demonstrations of the Data, the thing to be demonstrated, which is, that something is given, is never once assumed in the demonstration; from which it is manifest that every one of them is demonstrated synthetically: though indeed if a proposition of the Data be turned into a problem, the demonstration of the proposition becomes the analysis of the problem. See Simson's edition of Euclid's Data, which is esteemed the best.

DAVIS's *Quadrant*, the common sea quadrant, or back-staff. See BACK-STAFF. See also Robertson's Navigation, book 9, sect. 7.

DAY, a division of time arising from the appearance and disappearance of the sun.

DAY is either *natural* or *artificial*.

*Artificial DAY* is that which is primarily meant by the word Day, and is the time of its being light, or the time while the sun is above the horizon. Though sometimes the twilight is included in the term daylight; in opposition to night or darkness, being the time from the end of twilight to the beginning of daylight.

*Natural DAY* is the portion of time in which the sun performs one revolution round the earth; or rather the time in which the earth makes a rotation on its axis. And this is either astronomical or civil.

*Astronomical DAY* begins at noon, or when the sun's centre is on the meridian, and is counted 24 hours to the following noon.

*Civil DAY* is the time allotted for day in civil purposes, and begins differently in different nations, but still including one whole rotation of the earth on its axis; beginning either at sun-rise, sun-set, noon, or midnight.

1st, At sun-rising, among the ancient Babylonians, Persians, Syrians, and most other eastern nations, with the present inhabitants of the Balearic islands, the Greeks, &c. 2dly, At sun-setting, among the ancient Athenians and Jews, with the Austrians, Bohemians, Marcomanni, Silesians, modern Italians, and Chinese. 3dly, At noon, with astronomers, and the ancient Umbri and Arabians. And 4thly, at midnight, among the ancient Egyptians and Romans, with the modern English, French, Dutch, Germans, Spaniards, and Portuguese.

The day is divided into hours; and a certain number of days makes a week, a month, or a year.

The different length of the natural day in different climates, has been matter of controversy, viz, whether the natural days be all equally long throughout the year; and if not, what their difference is? A professor

of mathematics at Seville, in the *Philos. Transf.* vol. 10, pa. 425, asserts, from a continued series of observations for three years, that they are all equal. But Mr. Flamsteed, in the same *Transf.* pa. 429, refutes the opinion; and shews that one day, when the sun is in the equinoctial, is shorter than when he is in the tropics, by 40 seconds; and that 14 tropical days are longer than 50 many equinoctial ones, by 10 minutes. This inequality of the days flows from two several principles: the one, the eccentricity of the earth's orbit; the other, the obliquity of the ecliptic with regard to the equator, which is the measure of time. As these two causes happen to be differently combined, the length of the day is varied. See EQUATION of time.

DAY's-Work, in Navigation, denotes the reckoning or account of the ship's course, during 24 hours, or between noon and noon.

DECAGON, a plane geometrical figure of ten sides and ten angles. When all the sides and angles are equal, it is a *regular* decagon, and may be inscribed in a circle; otherwise, not.

If the radius of a circle, or the side of the inscribed hexagon, be divided in extreme and mean proportion, the greater segment will be the side of a decagon inscribed in the same circle. And therefore, as the side of the decagon is to the radius, so is the radius to the sum of the two. Whence, if the radius of the circle be  $r$ , the side of the inscribed decagon will be  $\frac{\sqrt{5}-1}{2} \times r$ .

If the side of a regular decagon be 1, its area will be  $\frac{5}{2}\sqrt{5} + 2\sqrt{5} = 7.6942088$ ; therefore as 1 is to 7.6942088, so is the square of the side of any regular decagon, to the area of the same: so that, if  $s$  be the side of such a decagon, its area will be equal to  $7.6942088s^2$ . See REGULAR Figure.

To inscribe a decagon in a circle geometrically. See my Mensuration, prob. 35, pa. 25, 2d edit.

DECEMBER, the last month of the year; in which the sun enters the tropic of Capricorn, making the winter solstice.

In the time of Romulus, December was the 10th month; whence the name, viz, from decem, ten; for the Romans began their year in March, from which December is the 10th month.

The month of December was under the protection of Vesta. Romulus assigned it 30 days; Numa reduced it to 29; which Julius Cæsar increased to 31.

At the latter part of this month they had the *Juveniles Ludi*, and the country people kept the feast of the goddess Vacuna in the fields, having then gathered in their fruits, and sown their corn; whence it seems is derived our popular festival called Harvest-home.

DECHALES (CLAUD-FRANCIS-MILLIET), an excellent mathematician, mechanist, and astronomer, was born at Chambery, the capital of Savoy, in 1611. He chiefly excelled in a just knowledge of the mathematical and mechanical sciences: not that he was bent upon new discoveries, or happy in making them; as his talent rather lay in explaining those sciences with ease and accuracy; which perhaps rendered him equally useful and deserving of esteem. Indeed it was generally allowed that he made the best use of the productions of other men, and that he drew the several parts of the mathematical sciences together with great judgment and



and perspicuity. It is also said of him, that his probity was not inferior to his learning; and that both these qualities made him generally admired and beloved at Paris, where for four years together he read public mathematical lectures in the college of Clermont.—From hence he removed to Marseilles, where he taught the art of navigation and the practical mathematical sciences.—He afterward became professor of mathematics in the university of Turin, where he died March 28, 1678, at 67 years of age.

Among other works which do honour to his memory, are,

1. An edition of Euclid's Elements; in which he has omitted the less important propositions, and explained the uses of those he has retained.

2. A Discourse on Fortification; and another on Navigation.

3. These performances, with some others, were collected in 3 volumes folio, under the title of *Mundus Mathematicus*, being indeed a complete course of mathematics. And the same was afterward much enlarged, and published at Lyons, 1690, in 4 large volumes, folio.

DECIL, *Decilis*, an aspect or position of two planets, when they are distant from each other a 10th part of the zodiac, or 36 degrees; and is one of the new aspects invented by Kepler.

DECIMALS, any thing proceeding by tens; as Decimal arithmetic, Decimal fractions, Decimal scales, &c.

DECIMAL Arithmetic, in a general sense, may be considered as the common arithmetical computation in use, in which the decimal scale of numbers is used, or in which the places of the figures change their value in a tenfold proportion, being 10 times as much for every place more towards the left hand, or 10 times less for every place more towards the right hand; the places being supposed indefinitely continued, both to the right and left. In this sense, the word includes both the arithmetic of integers, and decimal fractions. In a more restrained sense however, it means only.

DECIMAL Fractions, which are fractions whose denominator is always a 1 with some number of ciphers annexed, more or fewer according to the value of the fraction, the numerator of which may be any number whatever; as  $\frac{1}{10}$ ,  $\frac{3}{100}$ ,  $\frac{7}{1000}$ , &c.

As the denominator of a decimal is always one of the numbers 10, 100, 1000, &c, the inconvenience of writing these denominators down may be saved, by placing a proper distinction before the figures of the numerator only, to distinguish them from integers, for the value of each place of figures will be known in decimals, as well as in integers, by their distance from the 1st or unit's place of integers, having similar names at equal distances, as appears by the following scale of places, both in decimals and integers:

&c	8	8	8	8	8	8	8	.	8	8	8	8	8	8	&c
	millions	hundred	tens of thousands	thousands	hundreds	tens	units		tenths	hundredths	thousandths	ten thousandths	hundred thousandths	millionths	

The mark of distinction for decimals, called the *separatrix*, has been various at different times, according to the fancy of different authors; sometimes a semi-parenthesis, or a semicrochet, or a perpendicular bar, or the same with a line drawn under the figures, or simply this line itself, &c; but it is usual now to write either a comma or a full point near the bottom of the figures; I place the point near the upper part of the figures, as was done also by Newton; a method which prevents the separatrix from being confounded with mere marks of punctuation.

In setting down a decimal fraction without its denominator, the numerator must consist of as many places as there are ciphers in the denominator; and if it has not so many figures, the defect must be supplied by setting before them as many ciphers as will make them up so many: thus  $\frac{3}{10}$  is .3; and  $\frac{14}{100}$  is .14; and  $\frac{14}{1000}$  is .014; and  $\frac{3}{10000}$  is .003; &c.

So that, as ciphers on the right-hand side of integers *increase* their value decimally, or in a tenfold proportion, as 2, 20, 200, &c; so, when set on the left-hand of decimal fractions, they *decrease* the value decimally, or in a tenfold proportion, as .2, .02, .002, &c. But ciphers set on the other sides of these numbers, make no alteration in their value, neither of increase nor decrease, viz, on the left-hand of integers, or on the right-hand of decimals; so 2, or 02, or 002, &c, are all the same; as are also .2, or .20, or .200, &c.

Decimal fractions may be considered as having been introduced by Regiomontanus, about the year 1464, viz, when he transformed the tables of sines from a sexagesimal to a decimal scale. They were also used by Ramus, in his Arithmetic, written in 1550; and before his time by our countrymen Buckley and Recorde. But it was Stevinus who first wrote an express treatise on decimals, viz, about the year 1582, in *La Pratique d'Arithmetique*; since which time, this has commonly made a part in most treatises on arithmetic.

To reduce any Vulgar fraction, or parts of any thing, as suppose  $\frac{3}{8}$ , to a decimal fraction of the same value; add ciphers at pleasure to the numerator, and divide by the denominator: thus,

$$8 \overline{) 3.000}$$

$$.375 = \frac{3}{8};$$

and therefore .375 or  $\frac{375}{1000}$  is a decimal of the same value with the proposed vulgar fraction  $\frac{3}{8}$ .

Some vulgar fractions can never be reduced into decimals without defect; as  $\frac{1}{3}$ , which by division is .33333 &c infinitely.

Such numbers are very properly called circulating decimals, and repetends, because of the continual return of the same figures. See REPETENDS and CIRCULATES.

The common arithmetical operations are performed the same way in decimals, as they are in integers; regard being had only to the particular notation, to distinguish the fractional from the integral part of a sum. Thus,

In Addition and Subtraction, all figures of the same place or denomination are set straight under each other, the separatrix, or decimal points, forming a straight column.

In Multiplication, set down the numbers, and multiply



multiply them as integers; and point off from the product as many places of decimals as there are in both factors; prefixing ciphers if there be any defect of figures.

In Division, set down the numbers and divide also as in integers; making as many decimals in the quotient, as those in the dividend are more than those in the divisor.


Examples are numerous and common in most books of arithmetic.

DECIMAL Scales, are any scales divided decimally, or by tens.

DECLINATION, in Astronomy, is the distance of the sun, star, planet, &c, from the equinoctial, either northward or southward; being the same with latitude in geography, or distance from the equator.

Declination is either *real* or *apparent*, according as the real or apparent place of the point or object is considered.

The declination of any point S is an arch of the meridian SE, contained between the given point and the equinoctical EQ. The declination of a star &c, is found by knowing or observing the latitude of the place, i. e. the height of the pole, and then the meridian altitude of the star, &c; hence the difference between the co-latitude and the altitude of the star &c, is the declination, viz, the difference between EH the co-latitude, and SH the altitude, is ES the declination. For ex. Tycho found at Uranibourg the meridian altitude of Cauda Leonis, viz,



the co-latitude is

HS =	50°	59′	00″
HE =	34	5	45
<hr/>			
ES =	16	53	15

rem. declin. north      -      -      ES = 16 53 15

To find the Sun's Declination at any time; having given his place in the ecliptic; the rule is, as radius is to the sine of the sun's longitude, so is the sine of the greatest declination, or obliquity of the ecliptic, to his present declination sought.

In constructing tables of declination of the sun, planets, and stars, regard should be had to refraction, aberration, nutation, and parallax.

By comparing ancient observations with the modern, it appears that the declination of the fixed stars is variable; and that differently in different stars; for in some it increases, and in others decreases, and that in different quantities.

Circles of DECLINATION, are great circles of the sphere passing through the poles of the world, on which the declination is measured; and consequently are the same as meridians in geography.

Parallax of DECLINATION, is an arch of the circle of declination, by which the parallax in altitude increases or diminishes the declination of a star.

Parallels of DECLINATION, are lesser circles parallel to the equinoctial. The tropic of Cancer is a parallel of declination at 23° 28′ distance from the equinoctial northward; and the tropic of Capricorn is the parallel of declination as far distant southward.

Refraction of the DECLINATION, an arch of the circle of declination, by which the declination of a star is increased or diminished by means of the refraction.

DECLINATION of the Compass, or Needle, is its deviation from the true meridian. See VARIATION.

DECLINATION of a Vertical Plane, or Wall, in Dialling, is an arch of the horizon, comprehended either between the plane and the prime vertical circle, when it is counted from the east or west; or between the plane and the meridian, if it be accounted from the north or south.

DECLINATOR, or DECLINATORY, an instrument in dialling, by which the declination, inclination, and reclinacion of planes are determined.

DECLINERS, or DECLINING Dials, are those which cut obliquely, either the plane of the prime vertical circle, or the plane of the horizon.

The use of declining vertical dials is very frequent; because the erect walls of houses, on which dials are commonly drawn, mostly decline from the cardinal points. But incliners and recliners are very rare.

DECLIVITY, a sloping or oblique descent.

DECREMENT, *Equal, of Life*. See COMPLEMENT of life.

DECREMENTS are the small parts by which a variable and decreasing quantity becomes less and less. The indefinitely small decrements are proportional to the fluxions, which in this case are negative. See FLUXIONS, also INCREMENTS.

DECUPLE, a term of relation or proportion in arithmetic, implying a tenfold change or scale of variation, or one thing 10 times as much as another.

DECUSSATION, a term in geometry and optics, signifying the crossing of any two lines or rays &c: or the action itself of crossing.

The rays of light decussate in the chryselline, before they reach the retina.

Many of the lines of the 3d order decussate themselves. See Newton's Enumeratio, &c.

DEE (JOHN), a famous mathematician and astrologer, was born at London 1527. In 1542 he was sent to St. John's College, Cambridge. After five years close application to study, chiefly in the mathematical and astronomical sciences, he went over to Holland, to visit some mathematicians on the Continent; whence, after a year's absence, he returned to Cambridge, and was there elected one of the Fellows of Trinity College, then first erected by King Henry the 8th. In 1548 he left England a second time, his stay at home being rendered uneasy to him, by the suspicions that were entertained of his being a conjurer, arising chiefly from his application to astronomy, and from some mechanical inventions of his.

He now visited the university of Louvain; where he was much careffed, and visited by several persons of high rank. After two years he went into France, and read lectures, in the college of Rheims, upon Euclid's Elements. In 1551, he returned to England, and was introduced to King Edward, who assigned him a pension of 100 crowns, which he afterward relinquished for the rectory of Upton upon Severn. But soon after the accession of Queen Mary, having some correspondence with her sister Elizabeth, he was accused of practising against the queen's life by enchantment: on which account he suffered a tedious confinement, and was several times examined; till, in the year 1555, he obtained his liberty by an order of council.

When



When Queen Elizabeth ascended the throne, Dee was consulted concerning a propitious day for the coronation: on which occasion he was introduced to the queen, who made him great promises, which were but ill performed. In 1564, he made another voyage to the continent, to present a book which he had dedicated to the Emperor Maximilian. He returned to England the same year; but in 1571 we find him in Lorrain; where, being dangerously ill, the queen sent over two physicians to his relief. Having once more returned to his native country, he settled at Mortlake in Surry, where he continued his studies with much ardour, and collected a great library of printed books and manuscripts, with a number of instruments; most of which were afterward destroyed by the mob, as belonging to one who dealt with the devil.

In 1578, the queen being much indisposed, Mr. Dee was sent abroad to consult with German physicians and philosophers (astrologers no doubt) on the occasion; though some have said she employed him as a spy; probably he acted in a double capacity. We next find him again in England, where he was soon after employed in a more rational service. The queen, desirous to be informed concerning her title to those countries which had been discovered by Englishmen, ordered Dee to consult the ancient records, and to furnish her with proper geographical descriptions. Accordingly, in a short time, he presented to the queen, at Richmond, two large rolls, in which the discovered countries were geographically described and historically illustrated. His next employment was the reformation of the calendar, on which subject he wrote a rational and learned treatise, preserved in the Ashmolean library at Oxford.

Hitherto the extravagancies of our eccentric philosopher seem to have been tempered with a tolerable proportion of reason and science; but henceforward he is to be considered as a mere necromancer and credulous alchymist. In the year 1581 he became acquainted with one Edward Kelly, by whose assistance he performed divers incantations, and maintained a frequent imaginary intercourse with spirits and angels; one of whom made him a present of a black speculum (a polished piece of cannel-coal), in which these appeared to him as often as he had occasion for them, answering his questions, &c. Hence Butler says,

Kelly did all his feats upon  
The devil's looking-glass, a stone.

HUDIBRAS.

In 1583 they became acquainted with a certain Polish nobleman, then in England, named Albert Laski, a person equally addicted to the same ridiculous pursuits: he was so charmed with Dee and Kelly, that he persuaded them to accompany him to his native country; by whose means they were introduced to Rodolph king of Bohemia; who, though a credulous man, was soon disgusted with their nonsense. They were afterward introduced to the king of Poland, but with no better success. Soon after this they were entertained at the castle of a rich Bohemian nobleman, where they lived for some time in great affluence; owing, as they asserted, to their art of transmutation by means of a certain powder in the possession of Kelly.

VOL. I.

Dee, now quarrelling with his companion, quitted Bohemia, and returned to England, where he was once more graciously received by the queen; who, in 1595, made him warden of Manchester college, in which town he resided several years. In 1604 he returned to his house at Mortlake, where he died in 1608, at 81 years of age; leaving a large family and many works behind him.

The books that were printed and published by Dee, are, 1. *Propædumata Aphoristica*, &c. in 1558, in 12mo. — 2. *Monas Hieroglyphica ad Regem Romanorum Maximilianum*; 1564. — 3. *Epistola ad eximium ducis Urbini mathematicum, Fredericum Commandinum, prefixa libello Machometi Bagdadini de Superficierum Divisionibus* &c; 1570. — 4. *The British Monarchy, otherwise called, The Petty Navy Royal*; 1576. — 5. *Preface Mathematical to the English Euclid*, published by Henry Billingsley, 1570: certainly a very curious and elaborate composition, and where he says, many more arts are wholly invented by name, definition, property, and use, than either the Grecian or Roman mathematicians have left to our knowledge. — 6. *Divers and many annotations and inventions dispersed and added after the 10th book of English Euclid*; 1570. — 7. *Epistola prefixa Ephemeridibus Joannis Feldi à 1557, cui rationem declaraverat Ephemerides conscribendi*. — 8. *Parallaticæ Commentationis Paxcosque Nucleus quidam*; 1573.

This catalogue of Dee's printed and published works is to be found in his *Compendious Rehearsal* &c, as well as in his letter to Abp. Whitgift: and from the same places might be transcribed more than 40 titles of books unpublished, that were written by him.

DEFENCE, in Sieges, is used for any thing that serves to preserve or screen the soldiers, or the place. So the parapets, flanks, casemates, ravelins, and out-works, that cover the place, are called the defences, or covers of the place: and when the cannon have beaten down or ruined these works, so that the men cannot fight under cover, the defences of the place are said to be demolished.

*Line of DEFENCE*, is that which flanks a bastion, being drawn from the flank opposite to it.

The line of defence should not exceed a musket shot, i. e. 120 fathoms: indeed Melder allows 130, Scheiter 140, Vauban and Pagan 150.

*Line of DEFENCE, greater, or fichant*, is a line drawn from the point of the bastion to the concurrence of the opposite flank and curtain.

*Line of DEFENCE, lesser, or rasant, or flaquant*, is the face of the bastion continued to the curtain.

DEFERENT, or DEFERENS, in the ancient astronomy, an imaginary circle, which, as it were, carries about the body of a planet, and is the same with the eccentric; being invented to account for the eccentricity, perigee, and apogee of the planets.

DEFICIENT *Hyperbola*, is a curve having only one asymptote, though two hyperbolic legs running out infinitely by the side of the asymptote, but contrary ways. See CURVE.

This name was given to the curves by Newton, in his *Enumeratio Linearum tertii Ordinis*. There are 6 different species of them, which have no diameters, expressed by the equation  $xyy + cy = -ax^3 + bx^2 + cx + d$ , the term  $ax^3$  being negative. When the equation  $ax^4$

3 A

=  $bx^3$



$=bx^3 + cx^2 + dx + \frac{1}{4}ee$  has all its roots real and unequal, the curve has an oval joined to it. When the two middle roots are equal, the oval joins to the legs, which then cut one another in shape of a noose. When three roots are equal, the nodus is changed into a very acute cusp or point. When, of three roots with the same sign, the two greatest are equal, the oval vanishes into a point. When any two roots are imaginary, there is only a pure serpentine hyperbola, without any oval, decussation, cusp, or conjugate point; and when the terms  $b$  and  $d$  are wanting, it is of the 6th species.

There are also 7 different species of these curves, having each one diameter, expressed by the above equation when the term  $ey$  is wanting: according to the various conditions of the roots of the equation  $ax^3 = bx^2 + cx + d$ , as to their reality, equality, their having the same signs, or two of them being imaginary.

**DEFICIENT Numbers**, are those whose aliquot parts added together, make a sum less than the whole number: as 8, whose parts 1, 2, 4, make only 7; or the number 16, whose parts 1, 2, 4, 8 make only 15.

**DEFILE**, in *Fortification*, a narrow line or passage through which troops can pass only in file, making a small front, so that the enemy may easily stop their march, and charge them with the more advantage, as the front and rear cannot come to the relief of one another.

**DEFINITION**, an enumeration, or specification of the chief simple ideas of which a compound idea consists, in order to ascertain or explain its nature and character.

Definitions are of two kinds; the one *nominal*, or of the *name*; the other *real*, or of the *thing*.

**Nominal DEFINITION**, is an enumeration of such known characters as are sufficient for distinguishing any proposed thing from others; as is that of a square, when it is said that it is a quadrilateral, equilateral, rectangular figure.

**Real DEFINITION**, a distinct notion, explaining the genesis of a thing; that is, how the thing is made or done: as is this definition of a circle, viz, that it is a figure described by the motion of a right line about a fixed point.

**DEFLECTION**, the turning any thing aside from its former course, by some adventitious or external cause.

The word is often applied to the tendency of a ship from her true course, by reason of currents, &c, which turn her out of her right way.

**DEFLECTION of the Rays of Light**, is a property which Dr. Hook observed in 1675. He found it different both from reflection and refraction; and that it was made perpendicularly towards the surface of the opacous body.

This is the same property which Newton calls *inflection*. And by others it is called *diffraction*.

**DEGREE**, in *Algebra*, is used in speaking of equations, when they are said to be of such a degree according to the highest power of the unknown quantity. If the index of that power be 2; the equation is of the 2d degree; if 3, it is of the 3d degree, and so on.

**DEGREE**, in *Geometry* or *Trigonometry*, is the 360th part of the circumference of any circle; for every

circle is considered as divided into 360 parts, called degrees; which are marked by a small ° near the top of the figure; thus 45° is 45 degrees.

The degree is subdivided into 60 smaller parts, called minutes, meaning first minutes; the minute into 60 others, called seconds; the second into 60 thirds; &c. Thus 45° 12' 20" are 45 degrees, 12 minutes, 20 seconds.

The magnitude or quantity of angles is accounted in degrees; for because of the uniform curvature of a circle in all its parts, equal angles at the centre are subtended by equal arcs, and by similar arcs in peripheries of different diameters; and an angle is said to be of so many degrees, as are contained in the arc of any circle comprehended between the legs of the angle, and having the angular point for its centre. Thus we say an angle of 90°, or of 45° 24', or of 12° 20' 30". It is also usual to say, such a star is mounted so many degrees above the horizon, or declines so many degrees from the equator; or such a town is situate in so many degrees of latitude or longitude.—A sign of the ecliptic, or zodiac, contains 30 degrees.

The division of the circle into 360 degrees is usually ascribed to the Egyptians, probably from the circle of the sun's annual course, or according to their number of days in the year, allotting a degree to each day. It is a convenient number too, as admitting of a great many aliquot parts, as 2, 3, 4, 5, 6, 8, 9, &c. The sexagesimal subdivision, however, has often been condemned as improper, by many eminent mathematicians, as Stevinus, Oughtred, Wallis, Briggs, Gellibrand, Newton, &c; who advise a decimal division instead of it, or else that of centesms; as the degree into 100 parts, and each of these into 100 parts again, and so on. Stevinus even holds, that this division of the circle which he contends for, obtained in the wise age, *in seculo sapienti*. Stev. Cosmog. lib. 1, def. 6. And several large tables of sines &c have been constructed according to that plan, and published, by Briggs, Newton, and others. And I myself have carried the idea still much farther, in a memoir published in the Philos. Trans. of 1783, containing a proposal for a new division of the quadrant, viz, into equal decimal parts of the radius; by which means the degrees or divisions of the arch would be the real lengths of the arcs, in terms of the radius: and I have since computed those lengths, of the arcs, with their sines, &c, to a great extent and accuracy.

**DEGREE of Latitude**, is the space or distance on the meridian through which an observer must move, to vary his latitude by one degree, or to increase or diminish the distance of a star from the zenith by one degree; and which, on the supposition of the perfect sphericity of the earth, is the 360th part of the meridian.

The quantity of a degree of a meridian, or other great circle, on the surface of the earth, is variously determined by different observers: and the methods made use of are also various.

Eratosthenes, 250 years before Christ, first determined the magnitude of a degree of the meridian, between Alexandria and Syene on the borders of Ethiopia, by measuring the distance between those places, and comparing it with the difference of a star's zenith distances at those places; and found it 694½ stadia.

Pofidonius,



Pofidonius, in the time of Pompey the Great, by means of the different altitudes of a ftar near the horizon, taken at different places under the fame meridian, compared in like manner with the diftance between thofe places, determined the length of a degree only 600 ftadia.

Ptolomy fixes the degree at  $68\frac{2}{3}$  Arabic miles, counting  $7\frac{1}{2}$  ftadia to a mile. The Arabs themfelves, who made a computation of the diameter of the earth, by meafuring the diftance of two places under the fame meridian, in the plains of Sennar, by order of Almanon, make it only 56 miles. Kepler, determining the diameter of the earth by the diftance of two mountains, makes a degree 13 German miles; but his method is far from being accurate. Snell, seeking the diameter of the earth from the diftance between two parallels of the equator, finds the quantity of a degree, by one method 57064 Paris toifes, or 342384 feet; by another meth. 57057 - - toifes, or 342342 feet. The mean between which two numbers, M. Picard found by menfuration, in 1669, from Amiens to Malvoisin, the moft certain, and he makes the quantity of a degree 57060 toifes, or 342360 feet. However, M. Caffini, at the king's command, in the year 1700, repeated the fame labour, and meafuring the fpace of  $6^{\circ} 18'$ , from the obfervatory at Paris, along the meridian, to the city of Collioure in Rouffillon, that the greatnefs of the interval might diminifh the error, found the length of the degree equal to 57292 toifes, or 343742 Paris feet, amounting to 365184 Englifh feet.

And with this account nearly agrees that of our countryman Norwood, who, about the year 1635, meafured the diftance between London and York, and found that diftance 905751 Englifh feet; the difference of latitude being  $2^{\circ} 28'$ , hence, he determined the quantity of one degree at 367196 Englifh feet, or 57300 Paris toifes, or 69 miles, 288 yards. See Newt. Princ. Phil. prop. 19; and Hift. Acad. Scienc. anno 1700, pa. 153.

M. Caffini, the fon, completed the work of meafuring the whole arc of the meridian through France, in 1718. For this purpofe he divided the meridian of France into two arcs, which he meafured feparately. The one from Paris to Collioure gave him 57097 toifes; the other from Paris to Dunkirk - - 56960; and the whole arc from Dunkirk to Collioure 57060, the fame as M. Picard's.

M. Mufchenbroek, in 1700, refolving to correct the errors of Snell, found by particular obfervations, that the degree between Alemaer and Bergen-op-zoom contained 57033 toifes.

Meffieurs Maupertuis, Clairaut, Camus, Monnier, and Outhier of France, were fent on a northern expedition, and began their operations, affifted by M. Cellus, an eminent aftronomer of Sweden, in Swedifh Lapland, in July 1736, and finifhed them by the end of May following. They obtained the meafure of that degree, whole middle point was in lat.  $66^{\circ} 20'$  north, and found it 57439 toifes, when reduced to the level of the fea. About the fame time another company of philofophers was fent to South America, viz, Meflieurs Godin, Bouguer, and Condamine of France, to whom were joined Don Jorge Juan, and Don An-

tonio de Ulloa of Spain. They left Europe in 1735, and began their operations in the province of Quito in Peru, about October 1736, and finifhed them, after many interruptions, about 8 years after. The Spanifh gentlemen published a feparate account, and affign for the meafure of a degree of the meridian at the equator 56768 toifes. M. Bouguer makes it 56753 toifes, when reduced to the level of the fea; and M. Condamine ftates it at 56749 toifes.

M. Caille, being at the Cape of Good Hope in 1752, found the length of a degree of the meridian in lat.  $33^{\circ} 18' 30''$  fouth, to be 57037 toifes. In 1755, father Bofcovich found the length of a degree in lat.  $43^{\circ}$  north to be 56972 toifes, as meafured between Rome and Rimini in Italy. In the year 1740, Mefirs Caffini and La Caille again examined the former meafures in France, and, after making all the neceffary corrections, found the meafure of a degree, whole middle point is in lat.  $49^{\circ} 22'$  north, to be 57074 toifes; and in the lat. of  $45^{\circ}$ , it was 57050 toifes.

In 1764, F. Beccaria completed the meafurement of a portion of the meridian near Turin; from which it is deduced that the length of a degree, whole middle lat. is  $44^{\circ} 44'$  north, is 57024 Paris toifes.

At Vienna, 3 degrees of the meridian were meafured; and the medium, for the latitude of  $47^{\circ} 40'$  north may be taken at 57091 Paris toifes. See an account of this meafurement, by father Jofeph Liefganig, in the Philof. Tranf. 1768, pa. 15.

Finally, in the fame vol. too is an account of the meafurement of a part of the meridian in Maryland and Penfylvania, North America, 1766, by Mefirs Mafon and Dixon; from which it follows that the length of a degree whole middle point is  $39^{\circ} 12'$  north, was 363763 Englifh feet, or  $56904\frac{1}{2}$  Paris toifes.

Hence, from the whole we may collect the following table of the principal meafures of a degree in different parts of the earth, as meafured by different perfons, viz,

Mean Latitude.	Length of a Degree in Paris toifes.	Names of the Meafurers.	Years of Meafurement.
$66^{\circ} 20' N$	57422	Maupertuis &c	1736 & 1737
$49^{\circ} 23' N$	57074	{ Maupertuis &c and Caffini	1739 & 1740
$47^{\circ} 40' N$	57091	Liefganig	1766
$45^{\circ} 0' N$	57028	Caffini	1739 & 1740
$44^{\circ} 44' N$	57069	Beccaria	1760 to 1764
$43^{\circ} 0' N$	56979	{ Bofcovich and Le Maire	1752
$39^{\circ} 12' N$	56888	Mafon & Dixon	1764 to 1768
$0^{\circ} 0'$	56750	{ Bouguer and Condamine	1736 to 1744
$33^{\circ} 18' S$	57037	La Caille	1752

The method of obtaining the length of a degree of the terreftrial meridian, is to meafure a certain diftance upon it by a feries of triangles, whole angles may be found by actual obfervation, connected with a bafe, whole length may be taken by an actual furvey, or otherwife; and then to obferve the different altitudes of



of some star at the two extremities of that distance, which gives the difference of latitude between them: then, by proportion, as this difference of latitude is to one degree, so is the measured length to the length of one degree of the meridian sought. This method was first practised by Eratosthenes, in Egypt. See GEOGRAPHY, and the beginning of this article.

DEGREE of *longitude*, is the space between two meridians that make an angle of  $1^{\circ}$  with each other at the poles; the quantity or length of which is variable, according to the latitude, being every where as the cosine of the latitude; viz, as the cosine of one lat. is to the cosine of another, so is the length of a degree in the former lat. to that in the latter; and from this theorem is computed the following Table of the length of a degree of long. indifferent latitudes, supposing the earth to be a globe.

Degr. lat.	English miles.	Degr. lat.	English miles.	Degr. lat.	English miles.
0	69.07	31	59.13	61	33.45
1	69.06	32	58.51	62	32.40
2	69.03	33	57.87	63	31.33
3	68.97	34	57.20	64	30.24
4	68.90	35	56.51	65	29.15
5	68.81	36	55.81	66	28.06
6	68.62	37	55.10	67	26.96
7	68.48	38	54.37	68	25.85
8	68.31	39	53.62	69	24.73
9	68.15	40	52.85	70	23.60
10	67.95	41	52.07	71	22.47
11	67.73	42	51.27	72	21.32
12	67.48	43	50.46	73	20.17
13	67.21	44	49.63	74	19.02
14	66.95	45	48.78	75	17.86
15	66.65	46	47.93	76	16.70
16	66.31	47	47.06	77	15.52
17	65.98	48	46.16	78	14.35
18	65.62	49	45.26	79	13.17
19	65.24	50	44.35	80	11.98
20	64.84	51	43.42	81	10.79
21	64.42	52	42.48	82	9.59
22	63.97	53	41.53	83	8.41
23	63.51	54	40.56	84	7.21
24	63.03	55	39.58	85	6.00
25	62.53	56	38.58	86	4.81
26	62.02	57	37.58	87	3.61
27	61.48	58	36.57	88	2.41
28	60.93	59	35.54	89	1.21
29	60.35	60	34.50	90	0.00
30	59.75				

Note, This table is computed on the supposition that the length of the degrees of the equator are equal to those of the meridian at the medium latitude of  $45^{\circ}$ , which length is  $69\frac{1}{5}$  English miles.

The expressions *Latitude* and *Longitude*, are borrowed from the ancients, who happened to be acquainted with a much larger extent of the earth in the direction east and west, than in that of north and south; the former of which therefore passed, with them, for the length of the earth, or longitude, and the latter for the breadth or shorter dimension, viz, the latitude.

DEJECTION, in Astrology, is applied to the planets when in their detriment, as astrologers speak, i. e. when they have lost their force, or influence, as is pretended, by reason of their being in opposition to some others, which check and counteract them.

Or, it is used when a planet is in a sign opposite to that in which it has its greatest effect, or influence, which is called its exaltation. Thus, the sign Aries being the exaltation of the Sun, the opposite sign Libra is its dejection.

DEINCLINERS, or DEINCLINING *Dials*, are such as both decline and incline, or recline, at the same time. Suppose, for instance, a plane cutting the prime vertical circle at an angle of 30 degrees, and the horizontal plane at an angle of 24 degrees, the latitude of the place being 52 degrees; a dial drawn on this plane, is called a deincliner.

DELIACAL *Problem*, a celebrated problem among the ancients, concerning the duplication of the cube.

DELPHINUS, the *Dolphin*, a constellation of the northern hemisphere; whose stars, in Ptolomy's catalogue, are 10; in Tycho's the same; in Hevelius's 14; and in Flamsteed's 18.

DEMETRIUS, a Cynic philosopher, and disciple of Apollonius Thyaneus, in the age of Caligula. That emperor wishing to gain the philosopher to his interest by a large present, he refused it with indignation, saying, If Caligula wishes to bribe me, let him send me his crown. Vespasian was displeased with his insolence, and banished him to an island. The cynic derided the punishment, and bitterly inveighed against the emperor.

Demetrius lived to a very great age. And Seneca observes, that "nature had brought him forth to shew mankind that an exalted genius can live securely without being corrupted by the vice of the surrounding world."

DEMI-*Bastion*, in Fortification, one that has only one face and one flank.

DEMI-*Cannon*, and DEMI-*Culverin*, names of certain species of cannon, now no longer used.

DEMI-*Cross*, an instrument used by the Dutch to take the altitude of the sun or a star at sea; instead of which we use the cross staff, or forestaff.

DEMI-*Gorge*, is half the gorge or entrance into the bastion; not taken directly from angle to angle, where the bastion joins to the curtain, but from the angle of the flank to the centre of the bastion; or the angle the two curtains would make, were they thus protracted to meet in the bastion.

DEMI-*Lune*, or *Half-moon*, an outwork consisting of two faces, and two little flanks. It is often built before the angle of a bastion, and sometimes also before the curtain; though now it is very seldom used.

DEMOCRITUS, one of the greatest philosophers of antiquity, was born at Abdera, a town of Thrace, about the 80th olympiad, or about 400 years before Christ. His father, says Valerius Maximus, was able to entertain the army of Xerxes; and Diogenes Laertius adds, upon the testimony of Herodotus, that the king, in requital, presented him with some Magi and Chaldeans. From these, it seems, Democritus received the first part of his education; and from them, whilst yet a boy, he learned theology and astronomy. He next applied to Leucippus, from whom he learned the system.



system of atoms and a vacuum. His father dying, he and his two brothers divided his effects. Democritus made choice of that part which consisted in money, as being, though the least share, the most convenient for travelling; and it is said that his portion amounted to more than 100 talents, which is near 20 thousand pounds sterling. His extraordinary inclination for knowledge and the sciences, induced him to travel into all parts of the world where he might find learned men. He went to visit the priests of Egypt, from whom he learned geometry: He consulted the Chaldean and Persian philosophers: and it is said that he penetrated even into India and Ethiopia, to confer with the Gymnosophists. In these travels he wasted his substance; after which, at his return he was obliged for some time to be maintained by his brother. Settling himself at Abdera, he there governed in the most absolute manner, by virtue of his consummate wisdom. The magistrates of that city made him a present of 500 talents, and erected statues to him, even in his lifetime: but being naturally more inclined to contemplation than delighted with public honours and employments, he withdrew into solitude and retirement.

Democritus always laughed at human life, as a continued farce, which made the people think he was mad; on which they sent for Hippocrates to cure him: but that celebrated physician having discoursed with the philosopher, told the people that he had a great veneration for Democritus; and that, in his opinion, those who esteemed themselves the most healthy, were the most distempered.

It is said, though with little probability, that Democritus put out his own eyes, that he might meditate more profoundly on philosophical subjects. He died, according to Diogenes Laertius, in the 361st year before the Christian era, at 109 years of age. He was the author of many books, which are lost; from which Epicurus borrowed his philosophy.

DEMOIVRE (ABRAHAM), a celebrated mathematician, of French original, but who spent most of his life in England. He was born at Vitri in Champagne 1667. The revocation of the edict of Nantz, in 1685, determined him, with many others, to take shelter in England; where he perfected his mathematical studies, the foundation of which he had laid in his own country. A mediocrity of fortune obliged him to employ his talent in this way in giving lessons, and reading public lectures, for his better support: in the latter part of his life too, he chiefly subsisted by giving answers to questions in chances, play, annuities, &c, and it is said most of these responses were delivered at a Coffee-house in St. Martin's-lane, where he spent most of his time. The Principia Mathematica of Newton, which chance is said to have thrown in his way, soon convinced Demoivre how little he had advanced in the science he professed. This induced him to redouble his application; which was attended by a considerable degree of success; and he soon became connected with, and celebrated among, the first rate mathematicians. His eminence and abilities in this line, opened him an entrance into the Royal Society of London, and into the academies of Berlin and Paris. By the former his merit was so well known and esteemed, that they judged

him a fit person to decide the famous contest between Newton and Leibnitz, concerning the invention of Fluxions.

The collection of the Academy of Paris contains no memoir of this author, who died at London Nov. 1754, at 87 years of age, soon after his admission into it. But the Philosophical Transactions of London have several, and all of them interesting, viz, in the volumes 19, 20, 22, 23, 25, 27, 29, 30, 32, 40, 41, 43.

He published also some very respectable works, viz,

1. *Miscellanea Analytica, de Seriebus & Quadraturis* &c; 1730, in 4to. But perhaps he has been more generally known by his

2. *Doctrine of Chances; or, Method of Calculating the Probabilities of Events at Play*. This work was first printed, 1718, in 4to, and dedicated to Sir Isaac Newton: it was reprinted in 1738, with great alterations and improvements; and a third edition was afterwards printed.

3. *Annuities on Lives*; first printed 1724, in 8vo.—In 1742 the ingenious Thomas Simpson (then only 33 years of age) published his *Doctrine of Annuities and Reversions*; in which he paid some handsome compliments to our author. Notwithstanding which, Demoivre presently brought out a second edition of his *Annuities*, in the preface to which he passed some harsh reflections upon Simpson. To these the latter gave a handsome and effectual answer, 1743, in *An Appendix, containing some Remarks on a late book on the same subject, with answers to some personal and malignant misrepresentations, in the preface thereof*. At the end of this answer, Mr. Simpson concludes, “Lastly, I appeal to all mankind, whether, in his treatment of me, he has not discovered an air of self-sufficiency, ill-nature and inveteracy, unbecoming a gentleman.” Here it would seem the controversy dropped: Mr. Demoivre published the 3d edition of his book in 1750, without any farther notice of Simpson, but omitted the offensive reflections that had been in the preface.

DEMONSTRATION, a certain or convincing proof of some proposition: such as the demonstrations of the propositions in Euclid's Elements.

The method of demonstrating in mathematics, is the same with that of drawing conclusions from principles in logic. Indeed, the demonstrations of mathematicians are no other than series of enthymemes; every thing is concluded by force of syllogism, only omitting the premises, which either occur of their own accord, or are recollected by means of quotations.

DENDROMETER, an instrument lately invented by Messrs Duncombe and Whittel; so called, from its use in measuring trees.

DENEB, an Arabic term, signifying tail; used by astronomers as a name to some of the fixed stars, but especially for the bright star in the Lion's tail.

DENOMINATOR, of a fraction, is the number or quantity placed below the line, which shews the whole integer, or into how many parts the integer is supposed to be divided by the fraction; as that which gives denomination or name to the parts of the fraction. Thus, in the fraction  $\frac{5}{12}$ , five-twelfths, the number 12 is the denominator, and shews that the integer is here divided into 12 parts, or that it consists of 12 of those parts.



parts of which the numerator contains 5. Also  $b$  is the denominator of the fraction  $\frac{a}{b}$ .

**DENOMINATOR** of a ratio, is the quotient arising from the division of the antecedent by the consequent. Thus, 6 is the denominator of the ratio 30 to 5, because 30 divided by 5 gives 6. It is otherwise called the *exponent* of the ratio.

**DENSITY**, that property of bodies, by which they contain a certain quantity of matter, under a certain bulk or magnitude. Accordingly a body that contains more matter than another, under the same bulk, is said to be denser than the other, and that in proportion to the quantity of matter; or if the quantity of matter be the same, but under a less bulk, it is said to be denser, and so much the more so as the bulk is less. So that, in general, the density is directly proportional to the mass or quantity of matter, and reciprocally or inversely proportional to the bulk or magnitude under which it is contained.

The quantities of matter in bodies, or at least the proportions of them, are known by their gravity or weight; every equal particle of matter being endowed with an equal gravity, it is inferred that equal masses or quantities of matter have an equal weight or gravity; and unequal masses have proportionally unequal weights. So that, when body, or mass, or quantity of matter is spoken of, we are to understand their weight or gravity.

From the foregoing general proportion of the density of bodies, viz, that it is as the mass directly, and as the bulk inversely, may be inferred the proportion of the masses, or of the magnitudes; viz, that the mass or quantity of matter, is in the compound ratio of the bulk and density; and that the bulk or magnitude, is as the mass directly, and the density inversely. Hence, if  $B, b$  be two bodies, or masses, or weights;

and  $D, d$  their respective densities;

also  $M, m$  their magnitudes, or bulks:

Then the theorems above are thus expressed,

$$\text{viz, } D \propto \frac{B}{M}, \text{ and } B \propto DM, \text{ and } M \propto \frac{B}{D};$$

$$\text{or } D : d :: \frac{B}{M} : \frac{b}{m}, \text{ and } B : b :: DM : dm, \text{ \&c;}$$

$$\text{or } \frac{D}{d} = \frac{Bm}{bM}, \text{ and } \frac{B}{b} = \frac{DM}{dm}, \text{ and } \frac{M}{m} = \frac{Bd}{bD}.$$

No body is absolutely or perfectly dense; or no space is perfectly full of matter, so as to have no vacuity or interstices; on the contrary, it is the opinion of Newton, that even the densest bodies, as gold &c, contain but a small portion of matter, and a very great portion of vacuity; or that it contains a great deal more of pores or empty space, than of real substance.

It has been observed above, that the relative density of bodies may be known by their weight or gravity; and hence the most general way of knowing those densities, is by actually weighing an equal bulk or magnitude of the bodies, whether solid or fluid; if solid, by shaping them to the same figure and dimensions; if fluid, by filling the same vessel with them, and weighing it.

For fluids, there are also other methods of finding

their density: as 1st, by making an equilibrium between them in tubes that communicate; for, the diameters of the tubes being equal, and the weights or quantities of matter also equal, the densities will be inversely as the altitudes of the liquids in them, that is inversely as the bulk.

2dly, The densities of fluids are also compared together by immersing a solid in them; for if the solid be lighter than the liquids, the part immersed by its own weight, will be inversely as the density of the fluid; or if it be heavier, and sink in the liquids, by weighing it in them; then the weights lost by the body will be directly proportional to the densities of the fluids.

**DENSITY of the Air**, is a property that has much employed the later philosophers, since the discovery of the Torricellian experiment, and the air-pump. By means of the barometer it is demonstrated that the air is of the same density at all places at the same distance from the level of the sea; provided the temperature, or degree of heat, be the same. Also the density of the air always increases in proportion to the compression, or the compressing forces. And hence the lower parts of the atmosphere are always denser than the upper: yet the density of the lower air is not exactly proportional to the weight of the atmosphere, by reason of heat and cold, which make considerable alterations as to rarity and density; so that the barometer measures the elasticity of the air, rather than its density. If the height of the barometer be considered as the measure both of the density and elasticity of the air, when the thermometer is at  $31^\circ$ , and  $b$  be any other height of the barometer, when the thermometer is at  $t$  degrees; then in this case,  $b$  is the measure of the elasticity,

and  $\frac{466 - t}{435}b$  is the measure of the density of the air.

**DENSITY of the Planets**. In homogeneous, unequal, spherical bodies, the gravities on their surfaces, are as their diameters when the densities are equal, or the gravities are as the densities when the bulks are equal; therefore, in spheres of unequal magnitude and density, the gravity is in the compound ratio of the diameters and densities, or the densities are as the gravities divided by the diameters. Knowing therefore the diameters of the planets by observation and comparison, and the gravities at their surface by means of the revolution of the satellites, the relation of their densities becomes known. And as I have found the mean density of the earth to be about  $4\frac{1}{2}$  times that of water, *Philos. Trans.* 1778; hence the densities of the planets, with respect to water, become known, and are as below:

	Densities.
Water	1
The Sun	$1\frac{2}{5}$
Mercury	$9\frac{1}{2}$
Venus	$5\frac{1}{5}$
The Earth	$4\frac{1}{2}$
Mars	$3\frac{2}{7}$
The Moon	$3\frac{1}{11}$
Jupiter	$1\frac{1}{24}$
Saturn	$0\frac{1}{32}$
Georgian Planet	$0\frac{2}{160}$

As it is not likely that any of these bodies are homogeneous,



geneal, the densities here determined are supposed to be the mean densities, or such as the bodies would have if they were homogeneous, and of the same mass of matter and magnitude.

**DENTICLES**, or **DENTILS**, are ornaments in a cornice, cut after the manner of teeth. These are mostly affected in the Ionic and Corinthian orders; and of late also in the Doric. The square member on which they are cut, is called the Denticule.

**DEPARTURE**, in Navigation, is the easting or westing of a ship, with regard to the meridian she *departed* or failed from. Or, it is the difference in longitude, either east or west, between the present meridian the ship is under, and that where the last reckoning or observation was made. This departure, any where but under the equator, must be accounted according to the number of miles in a degree proper to the parallel the ship is in.


The Departure, in Plane and Mercator's Sailing, is always represented by the base of a right-angled plane triangle, where the course is the angle opposite to it, and the distance sailed is the hypotenuse, the perpendicular or other leg being the difference of latitude. And then the theorem for finding it, is always this: As radius is to the sine of the course, so is the distance sailed, to the departure sought.

**DEPRESSION of the Pole.** So many degrees &c. as you sail or travel from the pole towards the equator, so many it is said you depress the pole, because it becomes so much lower, or nearer the horizon.

**DEPRESSION of a Star, or of the Sun,** is its distance below the horizon; and is measured by an arc of a vertical circle, intercepted between the horizon and the place of the star.

**DEPRESSION of the Visible Horizon, or Dip of the Horizon,** denotes its sinking or dipping below the true horizontal plane, by the observer's eye being raised above the surface of the sea; in consequence of which, the observed altitude of an object is by so much too great.

Thus, the eye being at E, the height AE above the surface of the earth, whose centre is C; then EH is the real horizon, and Eb the visible one, below the former by the angle HEB, by reason of the elevation AE of the eye.



*To compute the Depression or Dip of the Horizon.*

In the right-angled triangle CEb, are given Cb the earth's radius = 21000000 feet, and the hypotenuse CE = the radius increased by the height AE of the eye; to find the angle C which is = the angle HEB, or depression sought;

viz, as Cb : CE :: radius : sec.  $\angle C$ ,  
or as CE : Cb :: radius : cofin.  $\angle C$ .

By either of these theorems are computed the numbers in the following table, which shews the depression or dip of the horizon of the sea for different heights of the eye, from 1 foot to 100 feet.

Height of the eye	Dip of the horizon	Height of the eye	Dip of the horizon	Height of the eye	Dip of the horizon
feet	' "	feet	' "	feet	' "
1	0 57	13	3 26	26	4 52
2	1 21	14	3 34	28	5 3
3	1 39	15	3 42	30	5 14
4	1 55	16	3 49	35	5 39
5	2 8	17	3 56	40	6 2
6	2 20	18	4 3	45	6 24
7	2 31	19	4 10	50	6 44
8	2 42	20	4 16	60	7 23
9	2 52	21	4 22	70	7 59
10	3 1	22	4 28	80	8 32
11	3 10	23	4 34	90	9 3
12	3 18	24	4 40	100	9 33

See Robertson's Navigation, book 9 appendix; and Tables requisite to be used with the Nautical Ephemeris, pa. i. See also LEVELLING.

**DEPTH**, the opposite of Height, and one of the dimensions of bodies, or of space. See HEIGHT, ALTITUDE, ELEVATION, &c.

**DERHAM** (Doctor WILLIAM), an eminent English philosopher and divine, was born at Stowton, near Worcester, 1657, and educated in Trinity College, Oxford. In 1682, he was presented to the vicarage of Wargrave in Berkshire; and, in 1689, to the valuable rectory of Upminster in Essex; which, lying at a convenient distance from London, afforded him an opportunity of conversing and corresponding with the principal literary geniuses of the nation. Applying himself there with great eagerness to the pursuit of his studies in natural and experimental philosophy, he soon became a distinguished and useful member of the Royal Society, whose Philosophical Transactions contain a great variety of curious and valuable pieces, the fruits of his laudable industry, in all or most of the volumes, from the 20th to the 39th, both inclusive; the principal of which are:

1. Experiments on the Motion of Pendulums in vacuo.
2. A Description of an instrument for finding the Meridian.
3. Experiments and Observations on the Motion of Sound.
4. On the Migration of Birds.
5. History of the Spots in the Sun, from 1703 to 1711.
6. Observations on the Northern Lights, Oct. 8, 1726, and Oct. 13, 1728.
7. Tables of the Eclipses of Jupiter's Satellites.
8. The difference of Time in the meridian of different places.
9. Of the meteor called Ignis Fatuus.
10. The History of the Death Watch.
11. Meteorological Diaries for several years.

In his younger days he published his Artificial Clock-maker, a very useful little work, that has gone through several editions. In 1711, 1712, 1714, he preached those sermons at Boyle's lecture, which he afterward digested under the well-known titles of *Physico-Theology* and *Astro-Theology*, or Demonstrations of the being and attributes.



attributes of God, from his works of creation, and a survey of the heavens.

In 1716 he was made a canon of Windsor, being at that time chaplain to the Prince of Wales; and in 1730 received, from the university of Oxford, the degree of Doctor of Divinity. He revised the *Miscellanea Curiosa*, in 3 vols 8vo, containing many curious papers of Dr. Halley and several other ingenious philosophers. To him also the world is indebted for the publication of the Philosophical Experiments of the late eminent Dr. Hooke, and other ingenious men of his time; as well as notes and illustrations of several other works.

Dr. Derham was very well skilled in medical as well as in physical knowledge; and was constantly a physician to the bodies as well as the souls of his parishioners. This great and good man, after spending his life in the most agreeable and improving study of nature, and the diligent and pious discharge of his duty, died at Upminster in 1735, at 78 years of age.

DESAGULIERS (JOHN THEOPHILUS), an eminent experimental philosopher, was the son of the Rev. John Desaguliers, a French Protestant refugee, and born at Rochelle in 1683. His father brought him to England an infant; and having taught him the classics himself, he sent him at a proper age to Christ-church College, Oxford; where in 1702 he succeeded Dr. Keil in reading lectures on experimental philosophy at Hart Hall. In 1712 he married, and settled in London, when he first of any introduced the reading of lectures in experimental philosophy in the metropolis, which he continued during the rest of his life with the greatest applause, having several times the honour of reading his lectures before the king and royal family. In 1714 he was elected F. R. S. and proved a very useful member, as appears from the great number of his papers that are printed in their Philos. Transf. on the subjects of optics, mechanics, and meteorology. The magnificent duke of Chandos made Dr. Desaguliers his chaplain, and presented him to the living of Edgware, near his seat at Cannons; and he became afterward chaplain to Frederick prince of Wales. In the latter part of his life, he removed to lodgings over the Great Piazza in Covent Garden, where he carried on his lectures with great success till the time of his death in 1749, at 66 years of age.

He was a member of several foreign academies, and corresponding member of the Royal Academy of Sciences at Paris; from which academy he obtained the prize, proposed by them for the best account of electricity. He communicated a multitude of curious and valuable papers to the Royal Society, for the year 1714 to 1743, or from vol. 29 to vol. 42.

Beside those numerous communications, he published a valuable *Course of Experimental Philosophy*, 1734, in 2 large vols. 4to; and gave an edition of *Gregory's Elements of Catoptrics and Dioptrics*, with an Appendix on Reflecting Telescopes, 8vo, 1735. This appendix contains some Original Letters that passed between Sir Isaac Newton and Mr. James Gregory, relating to those telescopes.

DESCENDING, a going or moving from above, downwards.

There are ascending and descending stars, and ascending and descending degrees, &c.

DESCENDING Latitude, is the latitude of a planet in its return from the nodes to the equator.

DESCENSION, in Astronomy, is either *right*, or *oblique*.

*Right* DESCENSION is a point, or arch, of the equator, which descends with a star, or sign, below the horizon, in a *right* sphere, and

*Oblique* DESCENSION is a point, or arch, of the equator, which descends at the same time with a star, or sign, below the horizon, in an oblique sphere.

Descensions, both right and oblique, are counted from the first point of Aries, or the vernal intersection, according to the order of the signs, i. e. from west to east. And, as they are unequal, when it happens that they answer to equal arcs of the ecliptic, as for example to the 12 signs of the zodiac, it follows, that sometimes a greater part of the equator rises or descends with a sign, in which case the sign is said to ascend or descend rightly: and sometimes again, a less part of the equator rises or sets with the same sign, in which case it is said to ascend or descend obliquely. See ASCENSION.

*Refraction of the* DESCENSION. See REFRACTION.

DESCENSIONAL Difference, is the difference between the right and oblique descension of the same star, or point of the heavens.

DESCENT, or *Fall*, in mechanics, &c, is the motion, or tendency, of a body towards the centre of the earth, either directly or obliquely.

The descent of bodies may be considered either as freely, like as in a vacuum, or as clogged or resisted by some external force, as an opposing body, or a fluid medium, &c.

1st, If the body  $b$  descend freely, and perpendicularly, by the force of gravity; then the motive force urging it downwards, is equal to its whole weight  $b$ ; and the quantity of matter being  $b$  also, the accelerative force will be  $\frac{b}{b}$  or 1.

2dly, If the body  $b$  descending, be opposed by some mechanical power, suppose a wedge or inclined plane, that is, instead of pursuing the perpendicular line of gravity, it is made to descend in a sloping direction down the inclined plane: then if the sine of the angle the plane makes with the horizon be  $s$ , to the radius 1, the motive force urging the body down the plane will be  $bs$ ; and therefore the accelerative force is  $\frac{bs}{b}$  or  $s$ ; which is less than in the former case in the proportion of  $s$  to 1.

3dly, In a medium, a body suspended loses as much of its weight, as is the weight of a like bulk of the medium; and when descending, it loses the same, beside the obstruction arising from the cohesion of the parts of the medium, and the opposing force of the particles struck, which last produces a greater or less resistance, according to the velocity of the motion. But, the weight of the body being  $b$ , and that of a like bulk of the fluid medium  $m$ , the motive force urging the body to descend, is only  $b - m$ ; that is, the body only falls by the excess of its weight above that of an equal bulk of the medium.

Hence, the power that sustains a body in a medium,



is equal to the excess of the absolute weight of the body above an equal bulk of the medium. Thus, a piece of copper weighing  $47\frac{1}{3}$  lb, loses  $5\frac{1}{3}$  lb of its weight in water: and therefore a power of 42 lb will sustain it in the water.

4thly, If two bodies have the same specific gravity, the less the bulk of the descending body is, the more of its gravity does it lose, and the slower does it descend, in the same medium. For, though the proportion of the specific gravity of the body to that of the fluid be still the same, whether the bulk be greater or less, yet the smaller the body, the more the surface is, in proportion to the mass; and the more the surface, the more the resistance of the parts of the fluid, in proportion.

5thly, If the specific gravities of two bodies be different; that which has the greatest specific gravity will descend with greater velocity in the air, or resisting medium, than the other body. Thus, a ball of lead descends swifter than wood or cork, because it loses less of its weight, though in a vacuum they both fall equally swift.

The cause of this descent, or tendency downwards, has been greatly controverted. Two opposite hypotheses have been advanced; the one, that it proceeds from an internal principle, and the other from an external one: the first is maintained by the Peripatetics, Epicureans, and the Newtonians; and the latter by the Cartesians and Gassendists. See also ACCELERATION.

#### *Laws of DESCENT of Bodies.*

1st, Heavy bodies, in an unresisting medium, fall with an uniformly accelerated motion. For, it is the nature of all constant and uniform forces, such as that of gravity at the same distance from the centre of the earth, to generate or produce equal additions of velocity in equal times. So that, if in one second of time there be produced 1 degree of velocity, in 2 seconds there will be 2 degrees of velocity, in 3 seconds 3 degrees, and so on, the degree or quantity of velocity being always proportional to the length of the time.

2nd, The space descended by an uniform gravity, in any time, is just the half of the space that might be uniformly described in the same time by the last velocity acquired at the end of that time, if uniformly continued. For, as the velocity increases uniformly in an arithmetical progression, the whole space descended by the variable velocity, will be equal to the space that would be described with the middle velocity uniformly continued for the same time; and this again will be only half the space that would be described with the last velocity, also uniformly continued for the same time, because the last velocity is double of the middle velocity, being produced in a double time.

3d, The spaces descended by an uniform gravity, in different times, are proportional to the squares of the times, or to the squares of the velocities. For the whole space descended in any number of particles of time, consists of the sums of all the particular spaces, or velocities, which are in arithmetical progression; but the sum of such an arithmetical progression, beginning at 0, and having the last term and the number of terms the same quantity, is equal to half the square of the last term, or of the number of terms; therefore

VOL. I.

the whole sums are as the squares of the times, or of the velocities.

This theory of the descents by gravity was first discovered and taught by Galileo, who afterwards confirmed the same by experiments; which have often been repeated in various ways by many other persons since his time, as Grimaldi, Riccioli, Huygens, Newton, and many others, all confirming the same laws.

The experiments of Grimaldi and Riccioli were made by dropping a number of balls, of half a pound weight, from the tops of several towers, and measuring the times of falling by a pendulum. Ricciol. Almag. Nov. tom. 1 lib. 2, cap. 21, prop. 4. An abstract of their experiments is exhibited here below:

Vibrations of the pendulum	The time		Space at the end of the time	Space descended each time
	"	'''	Rom. feet	Rom. feet
5	0	50	10	10
10	1	40	40	30
15	2	30	90	50
20	3	20	160	70
25	4	10	250	90
6	1	0	15	15
12	2	0	60	45
18	3	0	135	75
24	4	0	240	105

The space descended by a heavy body in any given time, being determined by experiment, is sufficient, in connection with the preceding theorems, for determining every inquiry concerning the times, velocities, and spaces descended, depending on an uniform force of gravity. From many accurate experiments made in England, it has been found that a heavy body descends freely through 16 feet 1 inch, or  $16\frac{1}{2}$  feet, in the first second of time; and consequently, by theorem 2, the velocity gained at the end of 1 second, is  $32\frac{1}{2}$  feet per second. Hence, by the same, and theorem 3, the velocity gained in any other time  $t$  is  $32\frac{1}{2}t$ , and the space descended is  $16\frac{1}{2}t^2$ . So that, if  $v$  denote the velocity, and  $s$  the space due to the time  $t$ , and there be put  $g = 16\frac{1}{2}$ ; then

$$\text{is } v = 2gt = 2\sqrt{gs} = \frac{2s}{t}.$$

$$s = gt^2 = \frac{v^2}{4g} = \frac{1}{2}tv.$$

$$t = \frac{v}{2g} = \sqrt{\frac{s}{g}} = \frac{2s}{v}.$$

The experiments with pendulums give also the same space for the descent of a heavy body in a second of time. Thus, in the latitude of London, it is found by experiment, that the length of a pendulum vibrating seconds is just  $39\frac{1}{8}$  inches; and it being known that the circumference of a circle is to its diameter, as the time of one vibration of any pendulum, is to the time in which a heavy body will fall through half the length of the pendulum; therefore as  $3.1416 : 1 :: 1 : \frac{1}{3.1416}$  which is the time of descending through  $19\frac{1}{8}$  inches, or half

3 B



half the length of the pendulum; then, spaces being as the squares of the times, as  $\frac{1}{3 \cdot 1416^2} : 1^2 :: 19\frac{2}{3} : 193$  inches, or 16 feet 1 inch, which therefore is the space a heavy body will descend through in one second; the very same as before.

4th, For any other constant force, instead of the perpendicular free descent by gravity, find by experiment, or otherwise, the space descended in one second by that force, and substitute that instead of  $16\frac{1}{2}$  for the value of  $g$  in these formulæ: or, if the proportion of the force to the force of gravity be known, let the value of  $g$  be altered in the same proportion, and the same formulæ will still hold good. So, if the descent be on an inclined plane, making, for instance, an angle of  $30^\circ$  with the horizon; then, the force of descent upon the plane being always as the sine of the angle it makes with the horizon, in the present case it will be as the sine of  $30^\circ$ , that is, as  $\frac{1}{2}$  the radius; therefore in this case the value of  $g$  will be but half the former,  $8\frac{1}{4}$ , in all the foregoing formulæ.

Or, if one body descending perpendicularly draw another after it, by means of a cord sliding over a pulley; then it will be, as the sum of the two bodies is to the descending body, so is  $16\frac{1}{2}$  to the value of  $g$  in this case; which value of it being used in the said formulæ, they will still hold good. And in like manner for any other constant forces whatever.

5th, The time of the oblique descent down any chord of a circle, drawn either from the uppermost point or lowermost point of the circle, is equal to the perpendicular descent through the diameter of the circle.

6th. The descent, or vibration, through all arcs of the same cycloid are equal, whether great or small.

7th. But the descent, or vibration, through unequal arcs of a circle, are unequal; the times being greater in the greater arcs, and less in the less.

8th, For Descents by Forces that are variable, see FORCES, &c. See also INCLINED PLANE, CYCLOID, PENDULUM, &c.

*Line of Swiftest DESCENT*, is that which a body, falling by the action of gravity, describes in the shortest time possible, from one given point to another. And this line, it is proved by philosophers, is the arc of a cycloid, when the one point is not perpendicularly over the other. See CYCLOID.

DESCRIBENT, a term in Geometry, signifying a line or superficies, by the motion of which a superficies or solid is described.

DETENTS, in a clock, are those stops which, by being lifted up or let down, lock and unlock the clock in striking.

DETENT-Wheel, or HOOP-Wheel, that wheel in a clock which has a hoop almost round it, in which there is a vacancy, where the clock locks.

DETERMINATE Number. See NUMBER.

DETERMINATE Problem, is that which has but one solution, or a certain limited number of solutions; in contradistinction to an indeterminate problem, which admits of infinite solutions.

Such, for instance, is the problem, To form an isosceles triangle on a given line, so that each of the angles at the base shall be double of that at the ver-

tex; which has only one solution: or this, To find an isosceles triangle whose area and perimeter are given; which admits of two solutions.

DETERMINATE Section, the name of a Tract, or General Problem, written by the ancient geometrician Apollonius. None of this work has come down to us, excepting some extracts and an account of it by Pappus, in the preface to the 7th book of his Mathematical Collections. He there says that the general problem was, "To cut an infinite right line in one point so, that, of the segments contained between the point of section sought, and given points in the said line, either the square on one of them, or the rectangle contained by two of them, may have a given ratio, either to the rectangle contained by one of them and a given line, or to the rectangle contained by two of them."

Pappus farther informs us, that this Tract of Apollonius was divided into two books; that the first book contained 6 problems, and the second 3; that the 6 problems of the first book contained 16 epitagmas, or cases, respecting the dispositions of the points; and the second book 9. Farther, that of the epitagmas of the 6 problems of the first book, 4 were maxima, and one a minimum: that the maxima are at the 2d epitagma of the 2d problem, at the 3d of the 4th, the 3d of the 5th, and the 3d of the 6th; but that the minimum was at the 3d epitagma of the 3d problem. Also, that the second book contained three determinations; of which the 3d epitagma of the 1st problem, and the 3d of the 2d were minima, and the 3d of the 3d a maximum. Moreover, that the first book had 27 lemmas, and the second book 24; and lastly, that both books contained 83 theorems.

From such account of the contents of this Tract, and the lemmas also given by Pappus, several persons have attempted to restore, or recompose what they thought might be nearly the form of Apollonius's tract, or the subject of each problem, case, determination, &c; among whom are, Snellius, an eminent Dutch mathematician of the last century; a translation of whose work was published in English by Mr. John Lawton, in 1772, together with a new restoration of the whole work by his friend Mr. William Wales.

DEW, a thin light insensible mist, or rain, ascending with a slow motion, and falling while the sun is below the horizon.

To us it appears to differ from rain, as less from more. Its origin and matter are doubtless from the vapours and exhalations that rise from the earth and water. See EXHALATION. Some define it a vapour liquefied, and let fall in drops. M. Huet, in one of his letters, shews that dew does not fall, but rises; and others have adopted the same opinion.

M. du Fay made several experiments, first with glasses, then with pieces of cloth stretched horizontally at different heights; and he found that the lower bodies, with their under surfaces, were wetted before those that were placed higher, or their upper surfaces. And Du Fay and Muschenbroek both found, that different substances, and even different colours, receive the dew differently, and some little or not at all.

From the principles laid down under the article EVAPORATION, the several phenomena of dews are easily accounted for. Such as, for instance, that dews are more



more copious in the spring, than in the other seasons of the year; there being then a greater stock of vapour in readiness, than at other times, by reason of the small expence of it in the winter's cold and frost. Hence it is too, that Egypt, and some other hot countries, abound with dews throughout all the heats of summer; for the air there being too hot to conflate the vapours in the day-time, they never gather into clouds; and hence they have no rain: but in climates that are excessively hot, the nights are remarkably cold; so that the vapours raised after sun-set, are readily condensed into dews.

It is natural to conclude, from the different substances which are combined with dew, that it must be either salutary or injurious, both to plants and animals.

It is not easy to ascertain the quantity of dew that rises every night, or in the whole year, because of the winds which disperse it, the rains which carry it down, and other inconveniences: but it is known that it rises in greater abundance after rain than after dry weather, and in warm countries than in cold ones. There are some places in which dew is observed only to ascend, and not to fall: and others again in which it is carried upwards in greater plenty than downwards, being dispersed by the winds.


Dr. Hales made some experiments, to determine the quantity of dew that falls in the night. For this purpose, on the 15th of August, at 7 in the evening, he filled two glazed earthen pans with moist earth; the dimensions of the pans being, 3 inches deep, and 12 inches diameter: and he observes, that the moister the earth, the more dew falls on it in a night; and that more than a double quantity of dew falls on a surface of water, than on an equal surface of moist earth. These pans increased in weight by the night's dew, 180 grains; and decreased in weight by the evaporation of the day, 1 oz 282 grs: so that 540 grains more are evaporated from the earth every 24 hours in summer, than the dew that falls in the night: i. e. in 21 days near 26 ounces from a circular area of a foot diameter. Now if 180 grains of dew, falling in one night on such an area, which is equal to 113 square inches, be equally spread on the surface, its depth will be the 150th part of an inch. He likewise found that the depth of dew in a winter's night was the 90th part of an inch. If therefore we allow 159 nights for the extent of the summer's dew, it will in that time amount to one inch in depth; and reckoning the remaining 206 nights for the extent of the winter's dew, it will produce 2.28 inches depth; and the dew of the whole year will amount to 3.28 inches depth. But the quantity which evaporated in a fair summer's day from the same surface, being 1 oz and 282 grs, gives the 40th part of an inch deep for evaporation, which is 4 times as much as fell at night. Dr. Hales observes that the evaporation of a winter's day is nearly the same as in a summer's day; the earth's greater moisture in winter compensating for the sun's greater heat in summer. Hales's Vegetable Statics, vol. 1, pa. 52 of 4th edit. See EVAPORATION.

Signor Beccaria made several experiments to demonstrate the existence of the electricity that is produced by dew. He observes in general, that such electricity took place in clear and dry weather, during which no strong wind prevailed; and that it depends on the

quantity of the dew, as the electricity of the rain depends on the quantity of the rain. He sometimes found that it began before sun-set; at other times not till 11 o'clock at night. *Artificial Electricity*, Appendix, letter 3.

DE WIT (JOHN), the famous Dutch pensionary, was born at Dort, in 1625; where he prosecuted his studies so diligently, that at 23 years of age, he published *Elementa Curvarum Linearum*, one of the deepest books in mathematics at that time. After taking his degrees, and travelling, he, in 1650, became pensionary of Dort, and distinguished himself very early in the management of public affairs, which soon after raised him to the rank of pensionary of Holland. After rendering the greatest benefits to his country in many important instances, and serving it in several high capacities, with the greatest ability, diligence, and integrity, by some intrigues of the court, it is said, he and his brother were thrown into prison, from whence they were dragged by the mob, and butchered with the most cruel and savage barbarity.

DIACAUSTIC *Curve*, or the *Cauptic by Refraction*, is a species of caustic curves, the genesis of which is in the following manner. Imagine an infinite number of rays BA, BM, BD, &c, issuing from the same luminous point B, refracted to or from the perpendicular MC, by the given curve AMD; and so, that CE the sines of the angles of incidence CME, be always to CG the sines of the refracted angles CMG, in a given ratio: then the curve HFN that touches all the refracted rays AH, MF, DN, &c, is called the **Diacoustic**, or *Cauptic by Refraction*.



DIACOUSTICS, or DIAPHONICS, the consideration of the properties of sound refracted in passing through different mediums; that is, out of a denser into a more subtile, or out of a more subtile into a denser medium. See SOUND.

DIADROME, a term sometimes used for the vibration, motion, or swing of a pendulum.

DIAGONAL, is a right line drawn across a figure, from one angle to another; and is sometimes called a diameter. It is used chiefly in quadrilateral figures, viz, in parallelograms and trapeziums.

1. Every diagonal, as AC, divides a parallelogram into two equal parts or triangles ABC, ADC.

2. Two diagonals, AC, BD, drawn in a parallelogram, do mutually bisect each other; as in the point E.



3. B 2

3. Any



3. Any line, as FG, drawn through the middle of the diagonal of a parallelogram, is bisected by it at the point E; and it divides the parallelogram into two equal parts, BFGC and AFGD.

4. The diagonal of a square is incommensurable with its side.

5. In any parallelogram, the sum of the squares of the four sides is equal to the sum of the squares of the two diagonals.

6. In any trapezium, the sum of the squares of the four sides is equal to the sum of the squares of the two diagonals together with 4 times the square of the distance between the middle points of the diagonals.

7. In any trapezium, the sum of the squares of the two diagonals is double the sum of the squares of two lines bisecting the two pairs of opposite sides.

8. In any quadrilateral inscribed in a circle, the rectangle of the two diagonals is equal to the sum of the two rectangles under the two pairs of opposite sides.

DIAGONAL Scale. See SCALES.

DIAGRAM, is a scheme for the explanation or demonstration of any figure, or of its properties.

DIAL, or SUN-DIAL, an instrument for measuring time by means of the sun's shadow. Or, it is a draught or description of certain lines on the surface of a body, so that the shadow of a style, or ray of the sun through a hole, should touch certain marks at certain hours.

Sun-Dials are doubtless of great antiquity. But the first upon record is, it seems, the dial of Ahaz, who began to reign 400 years before Alexander, and within 12 years of the Building of Rome: it is mentioned in Isaiah, chap. 38, ver. 8.

Several of the antients are spoken of, as makers of dials; as Anaximenes Milesius, Thales. Vitruvius mentions one made by the ancient Chaldee historian Berofus, on a reclining plane, almost parallel to the equator. Aristarchus Samius invented the hemispherical dial. And there were at the same time some spherical ones, with a needle for a gnomon. The discus of Aristarchus was an horizontal dial, with its rim raised up all around, to prevent the shadow from stretching too far.

It was late before the Romans became acquainted with dials. The first sun-dial at Rome was set up by Papyrius Cursor, about the 460th year of the city; before which time, Pliny says there is no mention of any account of time but by the sun's rising and setting: the first dial was set up near the temple of Quirinus; but being inaccurate, about 30 years after, another was brought out of Sicily by the consul M. Valerius Messala, which he placed on a pillar near the Rostrum; but neither did this shew time truly, because not made for that latitude; and, after using it 99 years, Martius Philippus set up another more exact.

The diversity of sun-dials arises from the different situation of the planes, and from the different figure of the surfaces upon which they are described; whence they become denominated *equinoctial*, *horizontal*, *vertical*, *polar*, *direct*, *erect*, *declining*, *inclining*, *reclining*, *cylindrical*, &c. For the general principles of their construction, see DIALLING.

Dials are sometimes distinguished into *primary* and *secondary*.

*Primary* DIALS are such as are drawn either on the

plane of the horizon, and thence called *horizontal* dials; or perpendicular to it, and called *vertical* dials; or else drawn on the polar and equinoctial planes, though neither horizontal nor vertical. And

*Secondary* DIALS are all those that are drawn on the planes of other circles, beside those last mentioned; or those which either decline, incline, recline, or deince.

Each of these again is divided into several others, as follow:

*Equinoctial* DIAL, is that which is described on an equinoctial plane, or one parallel to it.

*Horizontal* DIAL, is described on an horizontal plane, or a plane parallel to the horizon.—This dial shews the hours from sun-rise to sun-set.

*South* DIAL, or an *Erect, direct South Dial*, is that described on the surface of the prime vertical circle looking towards the south.—This dial shews the time from 6 in the morning till 6 at night.

*North* DIAL, or an *Erect, direct North Dial*, is that which is described on the surface of the prime vertical looking northward. This dial only shews the hours before 6 in the morning, and after 6 in the evening.

*East* DIAL, or *Erect, direct East Dial*, is that drawn on the plane of the meridian, looking to the east.—This can only shew the hours till 12 o'clock.

*West* Dial, or *Erect, direct West Dial*, is that described on the western side of the meridian.—This can only shew the hours after noon. Consequently this, and the last preceding one, will shew all the hours of the day between them.

*Polar* DIAL, is that which is described on a plane, passing through the poles of the world, and the east and west points of the horizon. It is of two kinds; the first looking up towards the zenith, and called the *upper*; the latter, down towards the nadir, called the *lower*. The polar dial therefore is inclined to the horizon in an angle equal to the elevation of the pole.—The upper polar dial shews the hours from 6 in the morning till 6 at night, and the lower one shews the hours before 6 in the morning, and after 6 in the evening, viz, from sun-rise and till sun-set.

*Declining* DIALS, are erect or vertical dials which decline from any of the cardinal points; or they are such as cut either the plane of the prime vertical, or of the horizon, at oblique angles.

Declining dials are of very frequent use; as the walls of houses, on which dials are mostly drawn, commonly deviate from the cardinal points.

Of declining dials there are several kinds, which are denominated from the cardinal points which they are nearest to; as decliners from the south, and from the north, and even from the zenith.

*Inclined* DIALS, are such as are drawn on planes not erect, but inclining, or leaning forward towards the south, or southern side of the horizon, in an angle, either greater or less than the equinoctial plane.

*Reclining* DIALS, are those drawn on planes not erect, but reclined, or leaning backwards from the zenith towards the north, in an angle greater or less than the polar plane.

*Deinclined* DIALS, are such as both decline and incline, or recline.—These last three sorts of dials are very rare.



*DIALS without Centres*, are those whose hour lines converge so slowly, that the centre, or point of their concurrence, cannot be expressed on the given plane.

*Quadrantal DIAL.* See *Horodistical QUADRANT*.

*Reflecting DIAL.* See *REFLECTING Dial*.

*Cylindrical DIAL*, is one drawn on the curve surface of a cylinder. This may first be drawn on a paper plane, and then pasted round a cylinder of wood, &c. It will shew the time of the day, the sun's place in the ecliptic, and his altitude at any time of observation.


There are also *Portable DIALS*, or on a *Card*, and *Universal DIALS* on a *Plain Cross*, &c.

*Refracted DIALS*, are such as shew the hour by means of some refracting transparent fluid.

*Ring DIAL*, is a small portable dial, consisting of a brass ring or rim, about 2 inches in diameter, and one-third of an inch in breadth. In a point of this rim there is a hole, through which the sun beams pass, and form a bright speck in the concavity of the opposite semi-circle, which gives the hour of the day in the divisions marked within it.

When the hole is fixed, the dial only shews true about the time of the equinox. But to have it perform throughout the whole year, the hole is made moveable, the signs of the zodiac, or the days of the month, being marked on the convex side of the ring; hence, in using it, the moveable hole is set to the day of the month, or the degree of the zodiac the sun is in; then suspending the dial by the little ring, turn it towards the sun, and his rays through the hole will shew the hour on the divisions within side.

*Universal, or Astronomical Ring DIAL*, is a ring dial which shews the hour of the day in any part of the earth; whereas the former is confined to a certain latitude. Its figure see represented below.



It consists of two rings or flat circles, from 2 to 6 inches in diameter, and of a proportionable breadth &c. The outward ring A represents the meridian of any place you are at, and contains two divisions of 90 degrees each, diametrically opposite to one another, the one serving from the equator to the north pole, the

other to the south pole. The inner ring represents the equator, and turns exactly within the outer, by means of two pivots in each ring at the hour of 12.

Across the two circles goes a thin reglet or bridge, with a cursor C, sliding along the middle of the bridge, and having a small hole for the sun to shine through. The middle of this bridge is conceived as the axis of the world, and the extremities as the poles: on the one side are drawn the signs of the zodiac, and on the other the days of the month. On the edge of the meridian slides a piece, to which is fitted a small ring to suspend the instrument by.

In this dial, the divisions on the axis are the tangents of the angles of the sun's declination, adapted to the semi-diameter of the equator as radius, and placed on either side of the centre: but instead of laying them down from a line of tangents, a scale of equal parts may be made, of which 1000 shall answer exactly to the length of the semi-axis, from the centre to the inside of the equinoctial ring; and then 434 of these parts may be laid down from the centre towards each end, which will limit all divisions on the axis, because 434 is the natural tangent of  $23^{\circ} 28'$ . And thus, by a nonius fixed to the sliding piece, and taking the sun's declination from an ephemeris, and the tangent of that declination from the table of natural tangents, the slider might be always set true within 2 minutes of a degree. This scale of 434 equal parts might be placed right against the  $23^{\circ} 28'$  of the sun's declination, on the axis, instead of the sun's place, which is there of little use. For then the slider might be set in the usual way, to the day of the month, for common use; or to the natural tangent of the declination, when great accuracy is required.

*To use this Dial:* Place the line *a* (on the middle of the sliding piece) over the degree of latitude of the place, as for instance  $51\frac{1}{2}$  degrees for London: put the line which crosses the hole of the cursor to the degree of the sign, or day of the month. Open the instrument so as that the two rings be at right angles to each other, and suspend it by the ring H, that the axis of the dial, represented by the middle of the bridge, may be parallel to the axis of the world. Then turn the flat side of the bridge towards the sun, so that his rays, striking through the small hole in the middle of the cursor, may fall exactly on a line drawn round the middle of the concave surface of the inner ring; in which case the bright spot shews the hour of the day in the said concave surface of the ring.

*Nocturnal or Night-DIAL*, is that which shews the hour of the night, by the light, or shadow projected from the moon or stars.

Lunar or Moon Dials may be either purposely described and adapted to the moon's motion; or the hour may be found on a sun-dial by the moon shining upon it, thus: Observe the hour which the shadow of the index points at by moon light; find the days of the moon's age in the calendar, and take  $\frac{3}{4}$ ths of that number, for the hours to be added to the hour shewn by the shadow, to give the hour of the night. The reason of which is, that the moon comes to the same horary circle later than the sun by about three quarters of an hour every day; and at the time of new moon the solar and lunar hour coincide.




**DIAL Planes**, are the plane superficies upon which the hour lines of dials are drawn.

*Tide DIAL.* See *TIDE Dial*.

**DIALLING**, the art of drawing sun, moon, and star-dials on any sort of surface, whether plane or curved.


Dialling is wholly founded on the first motion of the heavenly bodies, and chiefly the sun; or rather on the diurnal rotation of the earth: so that the elements of spherics, and spherical trigonometry, should be understood, before a person advances to the doctrine of dialling.

The principles of dialling may be aptly deduced from, and illustrated by, the phenomena of a hollow or transparent sphere, as of glass. Thus, suppose  $aPcp$  to re-



present the earth as transparent; and its equator as divided into 24 equal parts by so many meridian semi-circles  $a, b, c, d, e$ , &c, one of which is the geographical meridian of any given place, as London, which it is supposed is at the point  $a$ ; and if the hour of 12 were marked at the equator, both upon that meridian and the opposite one, and all the rest of the hours in order on the other meridians, those meridians would be the hour circles of London: because, as the sun appears to move round the earth, which is in the centre of the visible heavens, in 24 hours, he will pass from one meridian to another in an hour. Then, if the sphere had an opaque axis, as  $PEp$ , terminating in the poles  $P$  and  $p$ , the shadow of the axis, which is in the same plane with the sun and with each meridian, would fall upon every particular meridian and hour, when the sun came to the plane of the opposite meridian, and would consequently shew the time at London, and at all other places on the same meridian. If this sphere were cut through the middle by a solid plane  $ABCD$  in the rational horizon of London, one half of the axis  $EP$  would be above the plane, and the other half below it; and if straight lines were drawn from the centre of the plane to those points where its circumference is cut by the hour circles of the sphere, those lines would be the hour lines of an horizontal dial for London; for the shadow of the axis would fall upon each particular hour line of the dial, when it fell upon the like hour circle of the sphere.

If the plane which cuts the sphere be upright, as  $AFCG$ , touching the given place, for ex. London, at  $F$ ,



and directly facing the meridian of London, it will then become the plane of an erect direct south dial; and if right lines be drawn from its centre  $E$ , to those points of its circumference where the hour circles of the sphere cut it, these will be the hour lines of a vertical or direct south dial for London, to which the hours are to be set in the figure, contrary to those on an horizontal dial; and the lower half  $Ep$  of the axis will cast a shadow on the hour of the day in this dial, at the same time that it would fall upon the like hour circle of the sphere, if the dial plane was not in the way.

If the plane, still facing the meridian, be made to incline, or recline, any number of degrees, the hour circles of the sphere will still cut the edge of the plane in those points to which the hour lines must be drawn straight from the centre; and the axis of the sphere will cast a shadow on these lines at the respective hours. The like will still hold, if the plane be made to decline by any number of degrees from the meridian towards the east or west; provided the declination be less than 90 degrees, or the reclamation be less than the co-latitude of the place; and the axis of the sphere will be the gnomon: otherwise, the axis will have no elevation above the plane of the dial, and cannot be a gnomon.

Thus it appears that the plane of every dial represents the plane of some great circle on the earth, and the gnomon the earth's axis; the vertex of a right gnomon the centre of the earth or visible heavens; and the plane of the dial is just as far from this centre as from the vertex of this stile. The earth itself, compared with its distance from the sun, is considered as a point; and therefore, if a small sphere of glass be placed upon any part of the earth's surface, so that its axis be parallel to the axis of the earth, and the sphere have such lines upon it, and such planes within it, as above described; it will shew the hours of the day as truly as if it were placed at the earth's centre, and the shell of the earth were as transparent as glass. Fergusson, lect. 10.

The principal writers on Dials, and Dialling, are the following: Vitruvius, in his *Architecture*, cap. 4 and



and 7, lib. 9 : Sebastian Munster, his *Horographia* : John Dryander de *Horologiorum varia Compositione* : Conrad Gefner's *Pandectæ* : Andrew Schoner's *Gnomonica* : Fred. Commandine de *Horologiorum Descriptione* : Joan. Bapt. Benedictus de *Gnomonum Umbrarumque Solarium Ufu* : Joannes Georgius Schomberg, *Exegesis Fundamentorum Gnomonicorum* : Solomon de Caus, *Traité des Horologes Solaires* : Joan. Bapt. Trolta, *Praxis Horologiorum* : Defargues, *Maniere Universelle pour poser l'Effieu & placer les Heures & autres Choses aux Cadrans Solaires* : Ath. Kircher, *Ars magna Lucis & Umbræ* : Hallum, *Explicatio Horologii in Horto Regio Londini* : Tractatus *Horologiorum* Joannis Mark : Clavius, *Gnomonices de Horologiis* ; in which he demonstrates both the theory and the operations after the rigid manner of the ancient mathematicians : Dechales, Ozanam, and Schottus, gave much easier treatises on this subject ; as did also Wolfius in his *Elementa* : M. Picard gave a new method of making large dials, by calculating the hour lines ; and M. De la Hire, in his *Dialling*, printed in 1683, gave a geometrical method of drawing hour lines from certain points, determined by observation. Everhard Walper, in 1625, published his *Dialling*, in which he lays down a method of drawing the primary dials on a very easy foundation ; and the same foundation is also described at length by Sebastian Munster, in his *Rudimenta Mathematica*, published in 1651. In 1672, Sturmius published a new edition of Walper's *Dialling*, with the addition of a whole second part, concerning inclining and declining dials, &c. In 1708, the same work, with Sturmius's additions, was re-published, with the addition of a 4th part, containing Picard's and De la Hire's methods of drawing large dials, which makes much the best and fullest book on the subject. Peterfon, Michael, and Muller, have each written on *Dialling*, in the German language : Coetfius, in his *Horologiographia Plana*, printed in 1689 : Gauppen, in his *Gnomonica Mechanica* : Leybourn, in his *Dialling* : Bion, in his *Use of Mathematical Instruments* : Wells, in his *Art of Shadows*. There is also a treatise by M. Deparceux, 1740. Mr. Ferguson has also written on the same subject in his *Lectures on Mechanics* ; besides Emerson, in his *Dialling* ; and Mr. W. Jones, in his *Instrumental Dialling*.

*Universal Dialling Cylinder*, is represented in fig. 2, plate vii, where ABCD is a glass cylindrical tube, closed at both ends with brass plates, in the centres of which a wire or axis EFG is fixed. The tube is either fixed to an horizontal board H, so that its axis may make an angle with the board equal to that which the earth's axis makes with the horizon of any given place, and be parallel to the axis of the world ; or it may be made to move on a joint, and elevated for any particular latitude. The twenty-four hour lines are drawn with a diamond on the outside of the glass, equidistant from each other, and parallel to the axis. The XII next B stands for midnight, and the XII next the board H for noon. When the axis of this instrument is elevated according to the latitude, and the board set level, with the line HN in the plane of the meridian, and the end towards the north ; the axis EFG will serve as a gnomon or stile, and cast a shadow on the hour of the day among the pa-

rallel hour lines, when the sun shines on the instrument. As the plate AD at the top is parallel to the equator, and the axis EFG perpendicular to it, right lines drawn from the centre to the extremities of the parallels will be the hour lines of an equinoctial dial, and the axis will be the stile. An horizontal plate *ef* put down into the tube, with lines drawn from the centre to the several parallels, cutting its edge, will be an horizontal dial for the given latitude ; and a vertical plate *gc*, fronting the meridian, and touching the tube with its edge, with lines drawn from its centre to the parallels, will be a vertical south dial : the axis of the instrument serving in both cases for the stile of the dial : and if a plate be placed within the tube, so as to decline, incline, or recline, by any given number of degrees, and lines be drawn, as above, a declining, inclining, or reclining dial will be formed for the given latitude. If the axis with the several plates fixed to it be drawn out of the tube, and set up in sunshine in the same position as they were in the tube, AD will be an equinoctial dial, *ef* an horizontal dial, and *gc* a vertical south dial ; and the time of the day will be shewn by the axis EFG. If the cylinder were wood, instead of glass, and the parallel lines drawn upon it in the same manner, it would serve to facilitate the operation of making these several dials. The upper plate with lines drawn to the several intersections of the parallels, which appears obliquely in fig. 2, would be an equinoctial dial as in fig. 3, and the axis perpendicular to it be its stile. An horizontal dial for the latitude of the elevation of the axis might be made, by drawing out the axis and cutting the cylinder, as at *efgh*, parallel to the horizontal board H ; the section would be elliptic as in fig. 4. A circle might be described on the centre, and lines drawn to the divisions of the ellipse would be the hour lines ; and the wire put in its place again, as E, would be the stile. If this cylinder were cut by a plane perpendicular to the horizontal board H, or to the line SHN, beginning at g, the plane of the section would be elliptical as in fig. 5, and lines drawn to the points of intersection of the parallels on its edge would be the hour lines of a vertical direct south dial, which might be made of any shape, either circular or square, and F the axis of the cylinder would be its stile. Thus also inclining, declining, or reclining dials might be easily constructed, for any given latitude. Ferguson, *ubi supra*.

*Dialling Globe*, is an instrument made of brass, or wood, with a plane fitted to the horizon, and an index ; particularly contrived to draw all sorts of dials, and to give a clear exhibition of the principles of that art.

*Dialling Lines*, or *Scales*, are graduated lines, placed on rules or the edges of quadrants, and other instruments, to expedite the construction of dials. The principal of these lines are, 1. A scale of six hours, which is only a double tangent, or two lines of tangents each of 45 degrees, joined together in the middle, and equal to the whole line of lines, with the declination set against the meridian altitudes in the latitude of London, suppose, or any place for which it is made : the radius of which line of lines is equal to the dialling scale of six hours. 2. A line of latitudes, which is fitted to the hour scale, and is made by this canon :



canon: as the radius is to the chord of 90 degrees; so are the tangents of each respective degree of the line of latitudes, to the tangents of other arches: and then the natural sines of those arches are the numbers, which, taken from a diagonal scale of equal parts, will graduate the divisions of the line of latitude to any radius. The line of hours and latitudes is generally for pricking down all dials with centres. For the method of constructing these scales, see SCALE.

**DIALLING Sphere**, is an instrument made of brass, with several semicircles sliding over one another, on a moving horizon, to demonstrate the nature of the doctrine of spherical triangles, and to give a true idea of the drawing of dials on all manner of planes.

**DIAMETER**, of a circle, is a right line passing through the centre, and terminated at the circumference on both sides.

The diameter divides the circumference, and the area of the circle, into two equal parts. And half the diameter, or the semi-diameter, is called the radius.

For the proportion between the diameter and the circumference of a circle, see CIRCLE and CIRCUMFERENCE.

**DIAMETER of a Conic Section**, or *Transverse Diameter*, is a right-line passing through the centre of the section, or the middle of the axis.—The diameter bisects all ordinates, or lines drawn parallel to the tangent at its vertex. See CONIC Sections.

**Conjugate DIAMETER**, is a diameter, in Conic Sections, parallel to the ordinates of another diameter, called the transverse; or parallel to the tangent at the vertex of this other.

**DIAMETER, of any Curve**, is a right line which divides two other parallel right lines, in such manner that, in each of them, all the segments or ordinates on one side, between the diameter and different points of the curve, are equal to all those on the other side. This is Newton's sense of a Diameter.

But, according to some, a diameter is that line, whether right or curved, which bisects all the parallels drawn from one point to another of a curve. So that in this way every curve will have a diameter; and hence the curves of the 2d order, have, all of them, either a right-lined diameter, or else the curves of some one of the conic sections for diameters. And many geometrical curves of the higher orders, may also have for diameters, curves of more inferior orders.

**DIAMETER of Gravity**, is a right line passing through the centre of gravity.

**DIAMETER in Astronomy**. The diameters of the heavenly bodies are either *apparent*, i. e. such as they appear to the eye; or *real*, i. e. such as they are in themselves.

The apparent diameters are best measured with a micrometer, and are estimated by the measure of the angle they subtend at the eye. These are different in different circumstances and parts of the orbits, or according to the various distances of the luminary; being in the inverse ratio of the distance.

The sun's vertical diameter is found by taking the height of the upper and lower edge of his disk, when he is in the meridian, or near it; correcting the altitude of each edge on account of refraction and parallax; then the difference between the true altitudes of

the two, is the true apparent diameter sought. Or the apparent diameter may be determined by observing, with a good clock, the time which the sun's disc takes in passing over the meridian: and here, when the sun is in or near the equator, the following proportion may be used; viz, as the time between the sun's leaving the meridian and returning to it again, is to 360 degrees, so is the time of the sun's passing over the meridian, to the number of minutes and seconds of a degree contained in his apparent diameter: but when the sun is in a parallel at some distance from the equator, his diameter measures a greater number of minutes and seconds in that parallel than it would do in a great circle, and takes up proportionally more time in passing over the meridian; in which case say, as radius is to the cosine of the sun's declination, so is the time of the sun's passing the meridian reduced to minutes and seconds of a degree, to the arc of a great circle which measures the sun's apparent horizontal diameter. See TRANSIT.

The sun's apparent diameter may also be taken by the projection of his image in a dark room.

There are several ways of finding the apparent diameters of the planets: but the most certain method is that with the micrometer.

The following is a table of the apparent diameters of the sun and planets, in different circumstances, and as determined by different astronomers.

TABLE OF APPARENT DIAMETERS.											
1. Of the Sun, according to			Greatest			Mean			Least		
Aristarchus and Archimedes	30	0	0	30	0	0	30	0	0	0	0
Ptolomy	33	20	0	32	18	0	31	20	0	0	0
Albategnius	33	40	0	32	28	0	31	20	0	0	0
Regiomontanus	34	0	0	32	27	0	31	0	0	0	0
Copernicus	33	54	0	32	44	0	31	40	0	0	0
Tycho	32	0	0	31	0	0	30	0	0	0	0
Kepler	31	4	0	30	30	0	30	0	0	0	0
Riccioli	32	8	0	31	40	0	31	0	0	0	0
J. D. Cassini	32	46	0	32	13	0	31	40	0	0	0
Gascoigne	32	50	0	-	-	-	31	40	0	0	0
Flamsteed	32	48	0	-	-	-	31	30	0	0	0
Mouton	32	32	0	-	-	-	30	29	0	0	0
De la Hire	32	44	0	32	11	0	31	38	0	0	0
Louville	32	37	7	32	4	36	31	32	50	0	0
M. Cassini, jun.	32	37	30	32	5	0	31	32	30	0	0
Monnier	-	-	-	32	5	0	-	-	-	-	-
Short	32	33	0	-	-	-	31	28	0	0	0
2. Of the Moon.											
Ptolomy	35	30	0	-	-	-	31	20	0	0	0
Tycho, { in Conjunction	28	48	0	-	-	-	25	36	0	0	0
{ in Opposition	36	0	0	-	-	-	32	0	0	0	0
Kepler	32	44	0	-	-	-	30	0	0	0	0
De La Hire	33	30	0	-	-	-	29	30	0	0	0
Newton, { in Syzygy	-	-	-	31	30	0	-	-	-	-	-
{ in Quadrature	-	-	-	31	3	0	-	-	-	-	-
Mouton, Full, in Perigee	33	29	0	-	-	-	-	-	-	-	-
Monnier, { in Syzygy	-	-	-	31	30	0	-	-	-	-	-
{ in Quadrature	-	-	-	31	0	0	-	-	-	-	-



	Greatest	Mean	Least
<i>3. Of Mercury.</i>	<i>" " "</i>	<i>" " "</i>	<i>" " "</i>
Albategnius	- - -	2 5 10	- - -
Alfraganus	- - -	1 15 12	- - -
Tycho	3 57 0	2 10 0	1 29 0
Hevelius	0 11 0	0 6 0	0 4 0
Hortensius	0 28 0	0 19 0	0 10 0
Riccioli	0 25 12	0 13 48	0 9 20
Bradley	0 10 45	- - -	- - -
Monnier	- - -	- - -	0 10 0
<i>4. Of Venus.</i>			
Albategnius	3 8 0	- - -	- - -
Alfraganus	1 34 0	- - -	- - -
Tycho	4 40 0	3 15 0	1 52 0
Hevelius	1 5 0	0 16 0	0 9 0
Hortensius	1 40 0	0 53 0	0 15 20
Kepler	7 6 0	- - -	- - -
Riccioli	4 8 0	1 4 12	0 33 30
Huygens	- - -	- - -	1 25 0
Flamsteed	1 12 0	- - -	- - -
Horrox	1 18 30	- - -	- - -
Crabtree	1 9 0	- - -	- - -
Monnier	- - -	1 17 0	- - -
Transit of 1761	- - -	0 58 0	- - -
Transit of 1769	- - -	0 59 0	- - -
<i>5. Of Mars.</i>			
Albateg. and Alfrag.	- - -	1 34 0	- - -
Tycho	2 46 0	1 40 0	0 57 0
Hevelius	0 20 0	0 5 0	0 2 0
Hortensius	1 4 0	0 36 0	0 9 0
Kepler	6 30 0	- - -	- - -
Riccioli	1 32 0	0 22 0	0 10 6
Huygens	- - -	- - -	0 30 0
Flamsteed	0 33 0	- - -	- - -
Monnier	- - -	- - -	0 26 0
Herschel, } Polar Diam.	- - -	- - -	0 21 29
} Equat. Diam.	- - -	- - -	0 22 25
<i>6. Of Jupiter.</i>			
Albateg. and Alfrag.	- - -	2 36 40	- - -
Tycho	3 59 0	2 45 0	2 14 0
Hevelius	0 24 0	0 18 0	0 14 0
Hortensius	1 1 40	0 50 0	0 38 30
Kepler	0 50 0	- - -	- - -
Riccioli	1 8 46	0 49 46	0 38 18
Huygens	- - -	- - -	1 4 0
Flamsteed	0 54 0	- - -	- - -
Newton, from } Pound's Obf. }	- - -	0 37 15	- - -
Monnier	- - -	0 37 0	- - -
<i>7. Of Saturn.</i>			
Albateg. and Alfrag.	- - -	1 44 28	- - -
Tycho	2 12 0	1 50 0	1 34 0
Hevelius	0 19 0	0 16 0	0 14 0
Hortensius	0 42 40	0 37 0	0 31 0
Kepler	0 30 0	- - -	- - -

	Greatest	Mean	Least
<i>Of Saturn</i>	<i>" " "</i>	<i>" " "</i>	<i>" " "</i>
Riccioli	1 12 0	0 57 0	0 46 0
Huygens	- - -	- - -	0 30 0
Flamsteed	0 25 0	- - -	- - -
Newton, from } Pound's Obferv. }	- - -	0 16 0	- - -
Monnier	- - -	0 16 0	- - -
Huygens, 1/2's Ring	- - -	1 4 0	1 8 0
Newton, from } Pound's Obf. }	- - -	0 40 0	- - -
Monnier	- - -	0 42 0	- - -
<i>8. New Planet.</i>			
Herschel	- - -	0 3 54	- - -

The Mean Apparent diameters of the planets, as seen from the sun, are as follow :

Mercury, Venus, Earth, Moon, Mars, Jup. Sat. Hersch.  
20'' . 30'' 17'' 6'' 11'' 37'' 16'' 4''

For the true diameters of the sun and planets, and their proportions to each other, see PLANETS, SEMI-DIAMETER, and SOLAR SYSTEM.

DIAMETER of a Column, is its thickness just above the base. From this the module is taken, which measures all the other parts of the column.

DIAMETER of the Diminution, is that taken at the top of the shaft.

DIAMETER of the Swelling, is that taken at the height of one third from the base.

DIAPASON, a musical interval, otherwise called an octave, or eighth: so called because it contains all the possible diversities of sound.

If the lengths of two strings be to each other as 1 to 2, the tensions being equal, their tones will produce an octave.

DIAPENTE, is the perfect fifth or second of the concords, making an octave with the Diatessaron. The lengths of the chords are as 3 to 2.

DIAPHONOUS Body or Medium, one that is translucent, or through which the rays of light easily pass; as water, air, glass, talc, fine porcelain, &c. See TRANSPARENCY.

DIAPHONICS, is sometimes used for the science of refracted sound, as it passes through different mediums.

DIASTYLE, a sort of edifice, in which the pillars stand at such a distance from one another, that three diameters of their thickness are allowed for the intercolumnation.

DIATESSARON, is the perfect fourth; or a musical interval, consisting of one greater tone, one lesser, and one greater semitone. The lengths of strings to sound the diatessaron, are as 3 to 4.

DIATONIC, a term signifying the ordinary sort of music, which proceeds by tones or degrees, both ascending and descending. It contains or admits only the greater and lesser tone, and the greater semitone.

DIESIS, a division of a tone, less than a semitone; or an interval consisting of a lesser, or imperfect semitone. The Diesis is the smallest and softest change, or inflexion,



inflexion, of the voice imaginable. It is also called a *feint*, and is expressed by a St. Andrew's Cross, or *faltier*,

**DIFFERENCE**, is the excess by which one magnitude or quantity exceeds another. When a less quantity is subtracted from a greater, the remainder is otherwise called difference.

*Ascensional* DIFFERENCE. See ASCENSIONAL.

DIFFERENCE of *Longitude*, of two places, is an arch of the equator contained between the meridians of those two places, or the measure of the angle formed by their meridians.

**DIFFERENTIAL**, an indefinitely small quantity, part, or difference. By some, the Differential is considered as infinitely small, or less than any assignable quantity; and also as of the same import as fluxion.

It is called a Differential, or Differential Quantity, because often considered as the difference between two quantities; and as such it is the foundation of the Differential Calculus. Newton used the term *moment* in a like sense, as being the momentary increase or decrease of a variable quantity. M. Leibnitz and others call it also an infinitesimal.

DIFFERENTIAL of the 1st, 2d, 3d, &c degree. See DIFFERENTIO-DIFFERENTIAL.

DIFFERENTIAL Calculus, or *Method*, is a method of differencing quantities. See DIFFERENTIAL Method, CALCULUS, and FLUXIONS.

DIFFERENTIO-DIFFERENTIAL Calculus, is a method of differencing differential quantities.

As the sign of a differential is the letter *d* prefixed to the quantity, as *dx* the differential of *x*; so that of a differential of *dx* is *ddx*, and the differential of *ddx* is *ddd*, &c; similar to the fluxions  $\dot{x}$ ,  $\ddot{x}$ ,  $\ddot{\dot{x}}$ , &c.

Thus we have degrees of differentials. The differential of an ordinary quantity, is a differential of the first order or degree, as *dx*; that of the 2d degree is *ddx*; that of the 3d degree, *ddd*, &c. The rules for differentials, are the very same as those for fluxions. See FLUXIONS.

DIFFERENTIAL, in Logarithms. Kepler calls the logarithms of tangents, *differentiales*; which we usually call *artificial tangents*.

DIFFERENTIAL Equation, is an equation involving or containing differential quantities; as the equation  $3a^2dx - 2axdx + aydx + axdy = 0$ . Some mathematicians, as Stirling, &c, have also applied the term differential equation in another sense, to certain equations defining the nature of series.

DIFFERENTIAL Method, a method of finding quantities by means of their successive differences.

This method is of very general use and application, but especially in the construction of tables, and the summation of series, &c. This method was first used, and the rules of it laid down, by Briggs, in his Construction of Logarithms and other Numbers, much the same as they were afterwards taught by Cotes, in his Constructio Tabularum per Differentias; as I have shewn in the Introduction to my Logs. pa. 69 & seq. See Briggs's Arithmetica Logarithmica, cap. 12 and 13, and his Trigonometria Britannica.

The method was next treated in another form by Newton in the 5th Lemma of the 3d book of his Principia, and in his Methodus Differentialis, published

by Jones in 1711, with the other tracts of Newton. This author here treats it as a method of describing a curve of the parabolic kind, through any given number of points. He distinguishes two cases of this problem; the first, when the ordinates drawn from the given points to any line given in position, are at equal distances from one another; and the 2d, when these ordinates are not at equal distances. He has given a solution of both cases, at first without demonstration, which was afterwards supplied by himself and others: see his Methodus Differentialis above mentioned; and Stirling's Explanation of the Newtonian Differential Method, in the Philos. Trans. N<sup>o</sup> 362; Cotes, De Methodo Differentiali Newtoniana, published with his Harmonia Mensurarum; Herman's Phoronomia; and Le Seur & Jacquier, in their Commentary on Newton's Principia. It may be observed, that the methods there demonstrated by some of these authors extend to the description of any algebraic curve through a given number of points, which Newton, writing to Leibnitz, mentions as a problem of the greatest use.

By this method, some terms of a series being given, and conceived as placed at given intervals, any intermediate term may be found nearly; which therefore gives a method for interpolations. Briggs's Arith. Log. ubi supra; Newton Meth. Differ. prop. 5; Stirling, Methodus Differentialis.

Thus also may any curvilinear figure be squared nearly, having some few of its ordinates. Newton, ibid. prop 6; Cotes De Method. Differ.; Simpson's Mathematical Differt. pa. 115. And thus may mathematical tables be constructed by interpolation: Briggs, ibid. Cotes Canonotechnia.

The successive differences of the ordinates of parabolic curves, becoming ultimately equal, and the intermediate ordinate required, being determined by these differences of the ordinates, is the reason for the name Differential Method.

To be a little more particular.—The first case of Newton's problem amounts to this: A series of numbers, placed at equal intervals, being given, to find any intermediate number of that series, when its interval or distance from the first term of the series is given.—

Subtract each term of the series from the next following term, and call the remainders first differences; then subtract in like manner each of these differences from the next following one, calling these remainders 2d differences; again, subtract each 2d difference from the next following, for the 3d differences; and so on: then if A be the 1st term of the series,

$d'$  the first of the 1st differences,  
 $d''$  the first of the 2d differences,  
 $d'''$  the first of the 3d differences, &c;

and if  $x$  be the interval or distance between the first term of the series and any term sought, T, that is, let the number of terms from A to T, both included, be  $= x + 1$ ; then will the term sought, T, be =

$$A + \frac{x}{1}d' + \frac{x}{1} \cdot \frac{x-1}{2}d'' + \frac{x}{1} \cdot \frac{x-1}{2} \cdot \frac{x-2}{3}d''' \text{ \&c.}$$

Hence, if the differences of any order become equal, that is, if any of the diffs.  $d''$ ,  $d'''$ , &c, become  $= 0$ , the above series will give a finite expression for T the term sought; it being evident, that the series must terminate when any of the diffs.  $d''$ ,  $d'''$ , &c, become  $= 0$ .



It is also evident that the co-efficients  $\frac{x}{1}, \frac{x}{1}, \frac{x-1}{2}$ , &c, of the differences, are the same as to the terms of the binomial theorem.

For ex. Suppose it were required to find the log. tangent of  $5^{\circ} 1'' 12''' 24''''$ , or  $5^{\circ} 1'' \frac{2}{3}'''$ , or  $5^{\circ} 1'' \cdot 2066$  &c.

Take out the log. tangents to several minutes and seconds, and take their first and second differences, as below :

	Tang.	d'	d''
5' 0'' - - -	7.1626964		
5' 1 - - -	7.1641417	14453	
5' 2 - - -	7.1655821	14404	-49
5' 3 - - -	7.1670178	14357	-47

Here  $A = 7.1641417$ ;  $x = \frac{2}{3}$ ;  $d' = 14404$ ; and the mean 2d difference  $d'' = -48$ . Hence

$$\begin{array}{r} A - - - - - 7.1641417 \\ xd' - - - - - 2977 \\ \frac{x}{1} \cdot \frac{x-1}{1} d'' - - - - - 4 \end{array}$$

Theref. the tang. of  $5^{\circ} 1'' 12''' 24''''$  is 7.1644398

Hence may be deduced a method of finding the sums of the terms of such a series, calling its terms A, B, C, D, &c. For, conceive a new series having its 1st term = 0, its 2d = A, its 3d = A + B, its 4th = A + B + C, its 5th = A + B + C + D, and so on; then it is plain that assigning one term of this series, is finding the sum of all the terms A, B, C, D, &c. Now since these terms are the differences of the sums, 0, A, A + B, A + B + C, &c; and as some of the differences of A, B, C, &c, are = 0 by supposition; it follows that some of the differences of the sums will be = 0; and since in the series  $A + \frac{x}{1}d' + \frac{x}{1} \cdot \frac{x-1}{2}d''$

&c, by which a term was assigned, A represented the 1st term;  $d'$  the 1st of the 1st differences, and  $x$  the interval between the first term and the last; we are to write 0 instead of A, A instead of  $d'$ ,  $d'$  instead of  $d''$ ,  $d''$  instead of  $d'''$ , &c, also  $x+1$  instead of  $x$ ; which being done, the series expressing the sums will be

$$0 + \frac{x+1}{1}A + \frac{x+1}{1} \cdot \frac{x}{2}d' + \frac{x+1}{1} \cdot \frac{x}{2} \cdot \frac{x-1}{3}d'', \&c.$$

Or, if the real number of terms of the lines be called  $z$ , that is, if  $z = x+1$ , or  $x = z-1$ , the sum of the series will be  $Az + \frac{z}{1} \cdot \frac{z-1}{2}d' + \frac{z}{1} \cdot \frac{z-1}{2} \cdot \frac{z-2}{3}d''$  &c. See De Moivre's Doct. of Chances, pa. 59, 60; or his Miscel. Analyt. pa. 153; or Simpson's Essays, pa. 95.

For ex. to find the sum of six terms of the series of squares  $1 + 4 + 9 + 16 + 25 + 36$ , of the natural numbers.

Terms	d'	d''	d'''
1			
4	3		
9	5	2	0
16	7	2	0
25	9		

Here  $A = 1$ ,  $d' = 3$ ,  $d'' = 2$ ,  $d''' = 0$ , and  $z = 6$ ; therefore the sum is  $6 + \frac{6}{1} \cdot \frac{5}{2} \cdot 3 + \frac{6}{1} \cdot \frac{5}{2} \cdot \frac{4}{3} \cdot 2 = 6 + 45 + 40 = 91$  the sum required, viz. of  $1 + 4 + 9 + 16 + 25 + 36$ .

A variety of examples may be seen in the places above cited, or in Stirling's Methodus Differentialis, &c.

As to the Differential method, it may be observed, that though Newton and some others have treated it as a method of describing an algebraic curve, at least of the parabolic kind, through any number of given points; yet the consideration of curves is not at all essential to it, though it may help the imagination. The description of a parabolic curve through given points, is the same problem as the finding of quantities from their given differences, which may always be done by Algebra, by the resolution of simple equations. See Stirling's Method. Differ. pa. 97. This ingenious author has treated very fully of the differential method, and shewn its use in the solution of some very difficult problems. See also SERIES.

DIFFERENTIAL Scale, in Algebra, is used for the scale of relation subtracted from unity. See Recurring SERIES.

DIFFRACTION, a term first used by Grimaldi, to denote that property of the rays of light, which others have called Inflection; the discovery of which is attributed by some to Grimaldi, and by others to Dr. Hook.

DIGBY (Sir KENELM), a famous English philosopher, was born at Gotherst in Buckinghamshire, 1603. He was descended of an ancient family: his great grandfather, with six of his brothers, fought valiantly at Bosworth-field on the side of Henry the 7th, against Richard the 3d. His father, Everard, engaged in the gunpowder plot against James the 1st, for which he was beheaded. His son, however, was restored to his estate; and had afterwards several appointments under king Charles the 1st. He granted him letters of reprisal against the Venetians, from whom he took several prizes with a small fleet which he commanded. He fought the Venetians near the port of Scanderoon, and bravely made his way through them with his booty.

In the beginning of the civil wars, he exerted himself greatly in the king's cause. He was afterwards imprisoned, by order of the parliament; but was set at liberty in 1643. He afterward compounded for his estate; but being banished from England, he retired to France, and was sent on two embassies to Pope Innocent the 10th, from the queen, widow to Charles the 1st, whose chancellor he then was. On the restoration of Charles the 2d, he returned to London; where he died in 1665, at 62 years of age.

Digby was a great lover of learning, and translated several authors into English, as well as published several works of his own; as, 1. *Observations upon Dr. Brown's Religio Medici*, 1643.—2. *Observations on part of Spenser's Fairy Queen*, 1644.—3. *A Treatise of the Nature of Bodies*, 1644.—4. *A Treatise declaring the Operations and Nature of Man's Soul, out of which the Immortality of reasonable souls is evinced*: works that discover great penetration and extensive knowledge.

He applied much to chemistry; and found out several



ral useful medicines, which he distributed with a liberal hand. He particularly distinguished himself by his sympathetic powder for the cure of wounds at a distance; his discourse concerning which made great noise for a while. He held several conferences with Des Cartes, about the nature of the soul, and the principles of things. At the beginning of the Royal Society, he became a distinguished member, being one of the first council. And he had at his own house regular levees or meetings of learned men, to improve themselves in knowledge, by conversing with one another.

This eminent person was, for the early pregnancy of his talents, and his great proficiency in learning, compared to the celebrated Picus de Mirandola, who was one of the wonders of human nature. Yet his knowledge, though various and extensive, probably appeared greater than it really was; as he had all the powers of elocution and address to recommend it. He knew how to shine in a circle, either of ladies or philosophers; and was as much attended to when he spoke on the most trivial subjects, as when he spoke on the most important. It has been said that one of the princes of Italy, who had no child, was desirous that his princess should bring him a son by Sir Kenelm, whom he esteemed a just model of perfection.

DIGGES (LEONARD), a considerable mathematician in the 16th century, was descended from an ancient family, and born at Digges-court in the parish of Barham in Kent; but in what year is not known; and died about the year 1574. He was educated for some time at Oxford, where he laid a good foundation of learning. Retiring from thence, he prosecuted his studies, and became an excellent mathematician, a skilful architect, and an expert surveyor of land, &c. He composed several books: as, 1. *Tectonicum: briefly shewing the exact Measuring, and speedy Reckoning of all manner of Lands, Squares, Timber, Stones, Steeples, &c.* 1556, 4to. Augmented and published again by his son Thomas Digges, in 1592; and also reprinted in 1647.—2. *A Geometrical Practical Treatise, named Pantometria, in three books.* This he left in manuscript; but after his death, his son supplied such parts of it as were obscure and imperfect, and published it in 1591, folio; subjoining, “A Discourse Geometrical of the five regular and Platonic bodies, containing sundry theoretical and practical propositions, arising by mutual conference of these solids, Inscription, Circumscription, and Transformation.”—3. *Prognostication Everlasting of right good effect: or Choice Rules to judge the Weather by the Sun, Moon, and Stars, &c.* in 4to. 1555, 1556, and 1564: corrected and augmented by his son, with divers general tables, and many compendious rules, in 4to, 1592.

DIGGES (THOMAS), only son of Leonard Digges, after a liberal education from his tenderest years, went and studied for some time at Oxford; and by the improvements he made there, and the subsequent instructions of his learned father, became one of the best mathematicians of his age. When queen Elizabeth sent some forces to assist the oppressed inhabitants of the Netherlands, Mr. Digges was appointed muster-master general of them; by which he became well skilled in military affairs; as his writings afterward shewed. He died in 1595.

Mr. Digges, beside revising, correcting, and enlarg-

ing some pieces of his father's, already mentioned, wrote and published the following learned works himself: viz, 1. *Alae five Scale Mathematicæ: or Mathematical Wings or Ladders*, 1573, 4to. A book which contains several demonstrations for finding the parallaxes of any comet, or other celestial body, with a correction of the errors in the use of the radius astronomicus.—2. *An Arithmetical Military Treatise, containing so much of Arithmetic as is necessary towards military discipline*, 1579, 4to.—3. *A Geometrical Treatise, named Stratioticos, requisite for the perfection of Soldiers*, 1579, 4to. This was begun by his father, but finished by himself. They were both reprinted together in 1590, with several additions and amendments, under this title: “An Arithmetical Warlike Treatise, named Stratioticos, compendiously teaching the science of Numbers, as well in Fractions as Integers, and so much of the Rules and Equations Algebraical, and art of Numbers Cosical, as are requisite for the profession of a souldier. Together with the Moderne militaire discipline, offices, lawes, and orders in every well-governed campe and armie, inviolably to be observed.” At the end of this work there are two pieces; the first, “A briefe and true report of the proceedings of the Earle of Leycester, for the reliefe of the towne of Sluce, from his arrival at Vlissing, about the end of June 1587, untill the surrendrie thereof 26 Julii next ensuing. Whereby it shall plainlie appear, his excellencie was not in anie fault for the losse of that towne:” the second, “A briefe discourse what orders were best for repulsing of foraine forces, if at any time they should invade us by sea in Kent, or elsewhere.”—4. *A perfect Description of the Celestial Orbs, according to the most ancient doctrine of the Pythagoreans, &c.* This was placed at the end of his father's “Prognostication Everlasting, &c.” printed in 1592, 4to.—5. *A humble Motive for Association to maintain the religion established*, 1601, 8vo. To which is added, his *Letter to the same purpose to the archbishops and bishops of England.*—6. *England's Defence: or, A Treatise concerning Invasion.* This is a tract of the same nature with that printed at the end of his Stratioticos, and called, A briefe Discourse, &c. It was written in 1599, but not published till 1686.—7. *A Letter printed before Dr. John Dee's Parallatice Commentationis praxeosque nucleus quidam*, 1573, 4to.—Beside these, and his *Nova Corpora*, he left several mathematical treatises ready for the press; which, by reason of lawfuits and other avocations, he was hindered from publishing.

If our author was great in himself, he was not less so in his son, Sir Dudley Digges, so celebrated as a politician and elegant writer.

DIGIT, in Arithmetic, one of the ten characters, 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, by means of which all numbers are expressed.

DIGIT, in Astronomy, is the measure by which the part of the luminaries in eclipses is estimated, being the 12th part of the diameter of the luminary. Thus, an eclipse is said to be of 10 digits, when 10 parts out of 12 of the diameter are in the eclipsed part; when the whole of the luminary is just all covered, the digits eclipsed are just 12; and when the luminary is more than covered, as often happens in lunar eclipses, then more than 12 digits are said to be eclipsed: Thus, if the diameter or breadth of the earth's shadow, where the



the moon passes through, be equal to one diameter and a half of the moon, then  $18^\circ$  or digits are said to be eclipsed.

These digits are by Wolfius, and some others, called *digiti ecliptici*.

DIGIT is also a measure taken from the breadth of the finger; being estimated at  $\frac{3}{4}$ ths of an inch, and equal to 4 grains of barley, laid breadth-ways, so as to touch each other.

DILATATION, a motion of the parts of a body by which it expands, or opens itself, so as to occupy a greater space.

Many authors confound dilatation with rarefaction; but the more accurate writers distinguish between them; defining dilatation as the expansion of a body into a greater bulk, by its own elastic power; and rarefaction, the like expansion produced by means of heat.

The moderns have observed, that bodies which, after being compressed, and again left at liberty, restore themselves perfectly, do endeavour to dilate themselves with the same force by which they are compressed; and accordingly they sustain a force, and raise a weight equal to that with which they are compressed.

Again, bodies, in dilating by their elastic power, exert a greater force at the beginning of their dilatation, than towards the end; as being at first more compressed; and the greater the compression, the greater the elastic power and endeavour to dilate. So that these three, the compressing power, the compression, and the elastic power, are always equal.

Finally, the motion by which compressed bodies restore themselves, is usually accelerated: thus, when compressed air begins to restore itself, and dilate into a greater space, it is still compressed; and consequently a new impetus is still impressed upon it, from the dilatative cause; and the former remaining, with the increase of the cause, the effect, that is the motion and velocity, must be increased likewise. Indeed it may happen, that where the compression is only partial, the motion of dilatation shall not be accelerated, but retarded; as is evident in the compression of a sponge, soft bread, gauze, &c.

DILUTE. To dilute a body, is to render it liquid; or, if it were liquid before, to render it more so, by the addition of a thinner to it.

DIMENSION, the extension of a body, considered as measurable. Hence, as we conceive a body extended, and measurable in length, breadth, and depth, dimension is considered as threefold, viz, length, breadth, and thickness. So a line has one dimension only, viz length; a superficies two, length and breadth; and a body or solid has three, viz, length, breadth, and thickness.

DIMENSION is also particularly used with regard to the powers of quantities in equations. Thus, in a simple equation,  $x = a + b$ , the unknown quantity is only of one dimension; in a quadratic equation,  $x^2 = a^2 + b^2$ , it is of two dimensions; in a cubic,  $x^3 = a^3 + b^3$ , it is of three dimensions; and so on.

DIMETIENT, has sometimes been used for diameter.

DIMINISHED *Angle*, a term in Fortification. See *ANGLE*.

DIMINUTION, in Music, is the abating something of the full value or quantity of any note.

DIMINUTION, in Architecture, is a contraction of the upper part of a column, by which its diameter is made less than that of the lower part.

DINOCRATES, a celebrated ancient architect of Macedonia, of whom several extraordinary things are related. He was taken, by Alexander the Great, into Egypt, where he employed him in marking out and building the city of Alexandria. He formed a design, in which Mount Athos was to be laid out into the form of a man, in whose left hand were designed the walls of a great city, and all the rivers of the mount flowing into his right, and from thence into the sea. Another memorable instance of Dinocrates's architectonic skill, is his restoring and building, in a more august and magnificent manner than before, the celebrated temple of Diana at Ephesus, after Herostratus, for the sake of immortalizing his name, had destroyed it by fire. A third instance, more extraordinary and wonderful than either of the former, is related by Pliny in his Natural History; who says he had formed a scheme, by building the dome of the temple of Arsinoe at Alexandria of loadstone, to make her image all of iron to hang in the middle of it, as if it were in the air; but the king's death, and his own, prevented the execution or attempt of this project.

DIONYSIUS, the *Periegetic*, an ancient geographer and poet. Pliny says, he was a native of the Persian Alexandria, afterwards called Antioch, and at last Charax; that he was sent by Augustus, to survey the eastern part of the world. Dionysius wrote a great number of pieces, enumerated by Suidas and his commentator Eustathius: but his *Periegesis*, or *Survey of the World*, is the only one now extant; which may be well esteemed one of the most exact systems of ancient geography, since Pliny himself proposed it as his pattern.

DIONYSIAN, or *Victorian Period*. See *PERIOD*.

DIOPHANTUS, a celebrated mathematician of Alexandria, who has been reputed to be the inventor of Algebra; at least his is the earliest work extant on that science. It is not certain when Diophantus lived. Some have placed him before Christ, and some after, in the reigns of Nero and the Antonines; but all with equal uncertainty. It seems he is the same Diophantus who wrote the *Canon Arithmeticus*, which Suidas says was commented on by the celebrated Hypatia, daughter of Theon of Alexandria. His reputation must have been very high among the ancients, since they ranked him with Pythagoras and Euclid in mathematical learning. Bachet, in his notes upon the 5th book *De Arithmetica*, has collected, from Diophantus's epitaph in the Anthologia, the following circumstances of his life; namely, that he was married when he was 33 years old, and had a son born 5 years after; that this son died when he was 42 years of age, and that his father did not survive him above 4 years; from which it appears, that Diophantus was 84 years old when he died.

DIOPHANTUS wrote 13 books of Arithmetic, or Algebra, which Regiomontanus in his preface to Alfraganus, tells us, are still preserved in manuscript in the Vatican library. Indeed Diophantus himself tells us that



that his work consisted of 13 books, viz, at the end of his address to Dionysius, placed at the beginning of the work; and from hence Regiomontanus might be led to say the 13 books were in that library. No more than 6 whole books, with part of a seventh, have ever been published; and I am of opinion there are no more in being; indeed Bombelli, in the preface to his Algebra, written 1572, says there were but 6 of the books then in the library, and that he and another were about a translation of them.

Those 6 books, with the imperfect 7th, were first published at Basil by Xylander in 1575, but in a Latin version only, with the Greek scholia of Maximus Planudes upon the two first books, and observations of his own. The same books were afterwards published in Greek and Latin at Paris in 1621, by Bachet, an ingenious and learned Frenchman, who made a new Latin version of the work, and enriched it with very learned commentaries. Bachet did not entirely neglect the notes of Xylander in his edition, but he treated the scholiast Planudes with the utmost contempt. He seems to intimate, in what he says upon the 28th question of the 2d book, that the 6 books which we have of Diophantus, may be nothing more than a collection made by some novice, of such propositions as he judged proper, out of the whole 13: but Fabricius thinks there is no just ground for such a supposition.

**DIOPHANTINE Problems**, are certain questions relating to square and cubic numbers, and to right-angled triangles, &c; the nature of which were first and chiefly treated of by Diophantes, in his Arithmetic, or rather Algebra.

In these questions, it is chiefly intended to find commensurable numbers to answer indeterminate problems; which often bring out an infinite number of incommensurable quantities. For example, let it be proposed to find a right-angled triangle, whose three sides  $x$ ,  $y$ ,  $z$  are expressed by rational numbers; from the nature of the figure it is known that  $x^2 + y^2 = z^2$ , where  $z$  denotes the hypotenuse. Now it is plain that  $x$  and  $y$  may also be so taken, that  $z$  shall be irrational; for if  $x = 1$ , and  $y = 2$ , then is  $z = \sqrt{5}$ .

Now the art of resolving such problems, consists in ordering the unknown quantity or quantities, in such a manner, that the square or higher power may vanish out of the equation, and then by means of the unknown quantity in its first dimension, the equation may be resolved without having recourse to incommensurables. For ex. in the equation above,  $x^2 + y^2 = z^2$ , suppose  $z = x + u$ , then is  $x^2 + y^2 = x^2 + 2xu + u^2$ , out of which equation  $x^2$  vanishes, and then it is  $y^2 = 2xu + u^2$ , which gives  $x = \frac{y^2 - u^2}{2u}$ . Hence, assuming  $y$  and  $u$  equal to any numbers at pleasure, the three sides of the triangle will be  $y$ ,  $\frac{y^2 - u^2}{2u}$ , and  $\frac{y^2 + u^2}{2u}$ , which are all rational whenever  $y$  and  $u$  are rational. For ex. if  $y = 3$ , and  $u = 1$ , then  $\frac{y^2 - u^2}{2u} = 4$ , and  $x + u$  or  $\frac{y^2 + u^2}{2u} = 5$ . It is evident that this problem admits of infinite numbers of solutions, as  $y$  or  $u$  may be assumed infinitely various. See ALGEBRA, and DIOPHANTUS.

Abundant information on this sort of problems may

be found in the writings of a great many authors, particularly Fermat, Bachet, Ozanam, Kersey, Saunderson, Euler, &c.

**DIOPTER**, or **DIOPTRA**, the same with the index or alidade of an astrolabe, or other such instrument.

**DIOPTRA** was an instrument invented by Hipparchus, which served for several uses; as, to level water-courses; to take the height of towers, or places at a distance; to determine the places, magnitudes, and distances of the planets, &c.

**DIOPTRICS**, called also *anaclystics*, is the doctrine of refracted vision; or that part of Optics which explains the effects of light as refracted by passing through different mediums, as air, water, glass, &c, and especially lenses.

Dioptrics is one of the most useful and pleasant of all the human sciences; bringing the remotest objects near hand, enlarging the smallest objects so as to shew their minute parts, and even giving sight to the blind; and all this by the simple means of the attractive power in glass and water, causing the rays of light in their passage through them to alter their course according to the different substances of the medium; whence it happens, that the object seen through them, do, in appearance, alter their magnitude, distance, and situation.

The ancients have treated of direct and reflected vision; but what we have of refracted vision, is very imperfect. J. Baptista Porta wrote a treatise on refraction, in 9 books, but without any great improvement. Kepler was the first who succeeded in any great degree, on this subject; having demonstrated the properties of spherical lenses very accurately, in a treatise first published anno 1611. After Kepler, Galileo gave somewhat of this doctrine in his Letters; as also an Examination of the Preface of Johannes Pena upon Euclid's Optics, concerning the use of optics in astronomy. Des Cartes also wrote a treatise on Dioptrics, commonly annexed to his Principles of Philosophy, which is one of his best works; in which the true manner of vision is more distinctly explained than by any former writer, and in which is contained the true law of refraction, which was found out by Snell, though the name of the inventor is suppressed: here are also laid down the properties of elliptical and hyperbolical lenses, with the practice of grinding glasses. Dr. Barrow has treated on Dioptrics in a very elegant manner, though rather too briefly, in his Optical Lectures, read at Cambridge. There are also Huygens's Dioptrics, an excellent work of its kind. Molyneux's Dioptrics, a work rather heavy and dull. Hartsoeker's Essai de Dioptrique. Cherubin's Dioptrique Oculaire, et La Vision Parfaite. David Gregory's Elements of Dioptrics. Traber's Nervus Opticus. Zahn's Oculus Artificialis Teledioptricus. Dr. Smith's Optics, a complete work of its kind. Wolfius's Dioptrics, contained in his Elementa Mathematicos Universalis. But over all, the Treatise on Optics, and the Optical Lectures of Newton, in whose experiments are contained far more discoveries than in all the former writers. Lastly, this science was perfected by Dollond's discovery of the acromatic glasses, by which the colours are obviated in refracting telescopes.

The laws of Dioptrics see delivered under the article REFRACTION, LENS, &c; and the application of it in the construction of telescopes, microscopes, and other dioptrical



dioptrical instruments, under the articles TELESCOPE, and MICROSCOPE.

DIP of the Horizon. See DEPRESSION.

DIPPING-Needle, or *Inclinator* Needle, a magnetical needle, so hung, as that, instead of playing horizontally, and pointing out north and south, one end dips, or inclines to the horizon, and the other points to a certain degree of elevation above it.

The inventor of the Dipping-needle was one Robert Norman, a compass-maker at Ratcliffe, about the year 1580; this is not only testified by his own account, in his *New Attractive*, but also by Dr. Gilbert, Mr. William Burrowes, Mr. Henry Bond, and other writers of that time, or soon after it. The occasion of the discovery he himself relates, viz, that it being his custom to finish, and hang the needles of his compasses, before he touched them, he always found that, immediately after the touch, the north point would dip or decline downward, pointing in a direction under the horizon; so that, to balance the needle again, he was always forced to put a piece of wax on the south end, as a counterpoise. The constancy of this effect led him, at length, to observe the precise quantity of the dip, or to measure the greatest angle which the needle would make with the horizon. This, in the year 1576, he found at London was  $71^{\circ} 50'$ . It is not quite certain whether the dip varies, as well as the horizontal direction, in the same place. Mr. Graham made a great many experiments with the dipping-needle in 1723, and found the dip between 74 and 75 degrees. Mr. Nairne, in 1772, found it somewhat above  $72^{\circ}$ . And by many observations made since that time at the Royal Society, the medium quantity is  $72^{\circ} \frac{1}{2}$ . The trifling difference between the first observations of Norman, and the last of Mr. Nairne and the Royal Society, lead to the opinion that the dip is unalterable; and yet it may be difficult to account for the great difference between these and Mr. Graham's numbers, considering the well-known accuracy of that ingenious gentleman. *Philos. Transf.* vol. 45, pa. 279, vol. 62, pa. 476. vol. 69, 70, 71.

It is certain however, from many experiments and observations, that the dip is different in different latitudes, and that it increases in going northward. It appears from a table of observations made with a marine dipping needle of Mr. Nairne's, in a voyage towards the north pole, in 1773, that

in latitude $60^{\circ} 18'$	the Dip was $75^{\circ} 0'$ ,
in latitude $70 45$	the Dip was $77 52$ ,
in latitude $80 12$	the Dip was $81 52$ , and
in latitude $80 27$	the Dip was $82 2 \frac{1}{2}$ .

See Phipps's Voyage, pa. 122. See also the Observations of Mr. Hutchins, made in Hudson's Bay and Straits, *Philos. Transf.* vol. 65, pa. 129.

Burrowes, Gilbert, Ridley, Bond, &c, endeavoured to apply this discovery of the dip to the finding of the latitude; and Bond, going still farther, first of any proposed finding the longitude by it; but for want of observations and experiments, he could not go any length. Mr. Whiston, being furnished with the farther observations of colonel Windham, Dr. Halley, Mr Pound, Mr. Cunningham, M. Noel, M. Feuille, and his own, made great improvements in the doctrine and use of the dipping-needle, brought it to more certain rules, and endeavoured in good earnest to find the longitude by it. For

this purpose, he observes, 1st, That the true tendency of the north or south end of every magnetic needle, is not to that point of the horizon, to which the horizontal needle points, but towards another, directly under it, in the same vertical, and in different degrees under it, in different ages, and at different places. 2dly, That the power by which the horizontal needle is governed, and all our navigation usually directed, it is proved is only one quarter of the power by which the dipping-needle is moved; which should render the latter far the more effectual and accurate instrument. 3dly, That a dipping-needle of a foot long will plainly shew an alteration of the angle of inclination, in these parts of the world, in half a quarter of a degree, or  $7 \frac{1}{2}$  geographical miles; and a needle of 4 feet, in 2 or 3 miles; i. e. supposing these distances taken along, or near a meridian. 4thly, A dipping-needle, 4 feet long, in these parts of the world, will shew an equal alteration along a parallel, as another of a foot long will shew along a meridian; i. e. that will, with equal exactness, shew the longitude, as this the latitude.

This depends on the position of the lines of equal dip, in these parts of the world, which it is found do lie about 14 or 15 degrees from the parallels. Hence he argues, that as we can have needles of 5, 6, 7, 8, or more feet long, which will move with strength sufficient for exact observation; and since microscopes may be applied for viewing the smallest divisions of degrees on the limb of the instrument, it is evident that the longitude at land may thus be found to less than 4 miles.

And as there have been many observations made at sea with the same instrument by Noel, Feuille, &c, which have determined the dip usually within a degree, sometimes within  $\frac{1}{2}$  or  $\frac{1}{3}$  of a degree, and this with small needles, of 5 or 6, or at the most 9 inches long; it is inferred, that the longitude may be found, even at sea, to less than half a quarter of a degree. This premised, the observation itself follows.

*To find the Longitude or Latitude by the Dipping-Needle.*—If the lines of equal dip, below the horizon, be drawn on maps, or sea-charts, from good observations, it will be easy, from the longitude known, to find the latitude; and from the latitude known, to find the longitude either at sea or land.

Suppose, for example, a person travelling or sailing along the meridian of London, should find that the angle of dip, with a needle of one foot, was  $75^{\circ}$ ; the chart will shew that this meridian, and the line of dip, meet in the latitude of  $53^{\circ} 11'$ ; which is therefore the latitude sought.

Or if he be travelling or sailing along the parallel of London, i. e. in  $51^{\circ} 31'$  north latitude, and find the angle of dip  $74^{\circ}$ ; then this parallel, and the line of this dip, will meet on the map in  $1^{\circ} 46'$  of east longitude from London; which therefore is the longitude sought.

DIRECT, in Arithmetic, is when the proportion of any terms, or quantities, is in the natural or direct order in which they stand; being the opposite to *inverse*, which considers the proportion in the inverted order of the terms. So,  $3 : 4 :: 6 : 8$  directly; or  $3 : 4 :: 8 : 6$  inversely.

*Rule of Three* DIRECT, is when both pairs of terms are in direct proportion.

DIRECT, in Astronomy. A planet is said to be direct.



rect, or its motion direct, when it goes forward by its proper motion in the zodiac, according to the succession or order of the signs; or when it appears so to do, to an observer standing upon the earth. Whereas it is said to be, or to move *retrograde*, when it appears to go the contrary way, or backward; and to be *stationary*, when it seems not to move either way.

**DIRECT Dials.** See DIAL and DIALLING.

**DIRECT**, in Optics. *Direct* vision is that performed by direct rays; in contradistinction to vision by refracted, or reflected rays. Direct vision is the subject of Optics, which prescribes the laws and rules of it.

**DIRECT Rays**, are those which pass on in right lines from the object to the eye, without being turned out of their rectilinear direction by any intervening body, either opaque or pellucid; or without being either reflected or refracted.

**DIRECT Sphere.** See RIGHT Sphere.

**DIRECTION**, in Astronomy, the motion and other phenomena of a planet, when direct.

**DIRECTION**, in Astrology, is a kind of calculus, by which they pretend to find the time in which any notable accident shall befall the person whose horoscope is drawn.

For instance, having established the sun, moon, or ascendant, as masters or significators of life; and Mars or Saturn as promisers or portenders of death; the direction is a calculation of the time in which the significator shall meet the portender.

The significator they likewise call *apheta*, or giver of life; and the promiser, *anereta*, *promissor*, or giver of death.

They work the directions of all the principal points of the heavens and stars, as the ascendant, mid-heaven, sun, moon, and part of fortune. The like is done for the planets and fixed stars; but all differently, according to the different authors.

**Line of DIRECTION**, in Gunnery, is the direct line in which a piece is pointed. Sometimes a line of direction is marked on the upper side of the gun, by a small notch or slit, or knob, in the base and muzzle rings; but, unless the two wheels of the carriage stand equally high, this line will be fallacious; for which reason the gunners commonly find a new line of direction every time, by means of a plummet.

**Line of DIRECTION**, in Mechanics, denotes the line in which a body moves, or endeavours to proceed.

**Angle of DIRECTION**, is that comprehended between the lines of direction of two conspiring powers.

**Quantity of DIRECTION**, is used for the product arising from multiplying the velocity of the common centre of gravity, in a system of bodies, by the sum of their masses. In the collision of bodies the quantity of direction is the same both before and after the impulse.

**DIRECTION of the Load Stone**, that property by which the magnet, or a needle touched by it, always presents one of its ends toward one of the poles of the world, and the opposite end to the other pole. This is also called the *polarity* of the magnet or needle.

The attractive property of the magnet was known long before its directive; and the directive long before the inclinatory.

**Number of DIRECTION**, is the number of days that Septuagesima Sunday falls after the 17th of January. See NUMBER.

**DIRECTLY**, in Geometry: we say two lines lie directly against each other, when they are parts of the same right line. Also quantities are said to be directly proportional, when the proportion is according to the order of the terms; in contradistinction to *inversely*, or *reciprocally* proportional, which is taking the proportion contrary to the order of the terms.

In Mechanics, a body is said to strike or impinge *directly* against another body, when the stroke is in a direction perpendicular to the surface at the point of impact.


And a sphere in particular strikes directly against another, when the line of direction passes through both their centres.

**DIRECTRIX.** See DIRIGENT.

**DIRECTRIX**, in a Parabola, a line perpendicular to the axis produced, at the distance of the focus without the vertex.

**DIRIGENT**, a term expressing the line of motion, along which a describent line, or surface, is carried in the genesis of any plane or solid figure.

Thus, if the line AB move parallel to itself, and along the line AD, so that the point A always keeps in the line AD, and the point B in the line BC; a parallelogram ABCD will be formed; of which the line AB is the describent, and the line AD the dirigent. So also, if the surface ABEG be supposed carried along the line AD, in a position always parallel to itself at its first situation, the solid AF will be formed; where the surface AE is the describent, and the line AD is the dirigent.




**DISC**, or **DISK**, the body or face of the sun or moon; such as it appears to us; for though they be really spherical bodies, they are apparently circular planes.

The diameter of the disc is considered as divided into 12 equal parts, called digits; by means of which it is, that the magnitude of an eclipse is measured, or estimated.—In a total eclipse of either of those luminaries, the whole disc is obscured, or darkened; in a partial eclipse, only part of them.

**Illuminated Disc of the Earth.** See CIRCLE of Illumination.

**DISC**, in Optics, the magnitude of a telescope glass, or the width of its aperture, whatever its figure be, whether a plane, convex, meniscus, or the like.

**DISCHARGER**, or **DISCHARGING Rod**, in Electricity, consists of a handle of glass or baked wood, A,



and two bent metal rods BB, terminating in points, and capable of being screwed into the knobs DD, which move



move by a joint C, fixed to the handle A. Thus it may be used either with the points or balls, as occasion requires; and by being made moveable on a joint, it may be applied to larger or smaller jars at pleasure. By bringing one of these knobs or points to one coated side of a charged electric, and the other to the other side, or to any conductor connected with it, the communication is completed between the two sides, and the electric is discharged.

For the description and use of an Universal Discharger, by Mr. Henly, with which many curious experiments may be performed, see Cavallo's Electricity, pa. 164.

DISCORD, the relation of two sounds which are always, and of themselves, disagreeable, whether applied in succession or consonance.

DISCOUNT, or *Rebate*, is used for an allowance made on a bill, or any other debt not yet become due, in consideration of making present payment of the bill or debt.

Among merchants and traders, it is usual to allow a sum for discount that is equal to the interest of the debt, calculated for the time till it becomes due: but this is not just; for as the true value of the discount is equal to the difference between the debt and its present worth, it is equal only to the interest of that present worth, instead of the interest on the whole debt. And therefore the rule for finding the true discount is this:

As the amount of 100l, for the given rate and time:

Is to the given sum or debt ::

So is the interest of 100l, for the given rate and time:

To the discount of the debt.

So that, if  $p$  be the principal or debt,  $r$  the rate of interest per cent. and  $t$  the time;

then as  $100 + rt : p :: rt : \frac{prt}{100 + rt}$ , which is the true

discount. Hence also  $p - \frac{prt}{100 + rt} = \frac{100p}{100 + rt}$  is the present worth, or sum to be received.

For ex. Suppose it be required to find the discount of 250l, for five months, at the rate of 5 per cent. per annum interest. Here  $p = 250$ ,  $r = 5$ , and  $t = \frac{5}{12}$  or 5 months; then

$$\frac{prt}{100 + rt} = \frac{250 \times 5 \times \frac{5}{12}}{100 + 5 \times \frac{5}{12}} = \frac{250 \times 25}{1200 + 25} = \frac{250}{49} = 5\text{l. } 2\text{s. } \frac{2}{7}\text{d. the discount sought.}$$

A Table of Discounts may be seen in Smart's Tables of Interest, the use of which makes calculations of discount very easy.

Discount is also the name of a rule in books of Arithmetic, by which calculations of Discount are made.

DISCRETE, or DISJUNCT, *Proportion*, is that in which the ratio between two or more pairs of numbers is the same, and yet the proportion is not continued, so as that the ratio may be the same between the consequent of one pair and the antecedent of the next pair.

Thus, if the numbers or proportion  $6 : 8 :: 3 : 4$  be considered: the ratio of 6 to 8 is the same as that of 3 to 4, and therefore these four numbers are proportional: but it is only *discretely* or *disjunctly*, and not continued; for 8 to 3 is not the same ratio as the former; that is, the proportion is broken off between 8 and 3, and not continued all along, as it is in these fol-

lowing four numbers, which are called *continual proportionals*, viz,  $3 : 6 :: 12 : 24$ .

DISCRETE *Quantity*, is such as is not continued and joined together. Such for instance is any number; for its parts, being distinct units, cannot be united into one *continuum*; for in a *continuum* there are no actual determinate parts before division, but they are potentially infinite: so that it is usually and truly said that continued quantity is divisible *in infinitum*.

DISDIAPASON, in Music, a compound concord, in the quadruple ratio of 4 to 1, being a fifteenth or double eighth, and is produced when the voice goes from the first tone to the fifteenth.

DISJUNCT *Proportion*. See DISCRETE *Proportion*.

DISK. See DISC.

DISPART, a term in Gunnery, used for a mark set upon the muzzle-ring of a piece of ordnance, of such height, that a sight-line taken from the top of the base ring near the vent or touch-hole to the top of the dispart near the muzzle, may be parallel to the axis of the concave cylinder; for which reason it is evident that the height of the dispart is equal to the difference between the radii of the piece at the base and muzzle-rings, or to half the difference of the diameters there. Hence comes the common method of disparting the gun, which is this: Take, with the calipers, the two diameters, viz, of the base ring and the place where the dispart is to stand, subtract the less from the greater, and take half the difference, which will be the length of the dispart, which is commonly cut to that length from a small bit of wood, and so fixed upright in its place with a bit of wax or pitch.

DISPERSION, in Dioptrics, is the divergency of refracted rays of light.

*Point of DISPERSION*, is a point from which refracted rays begin to diverge, when their refraction renders them divergent.—It is called *point of dispersion* in opposition to the *point of concurrence*, or point in which converging rays concur after refraction. But it is more usual to call the latter the *focus*, and the former the *virtual focus*.

DISPERSION of Light, occasioned by the refrangibility of the rays, or the nature of the refracting medium. See ABERRATION, and INFLECTION.

DISSIPATION, in Physics, a gradual, slow, insensible loss or consumption of the minute parts of a body; or, more properly, the flux by which they fly off and are lost. See EFFLUVIA.

*Circle of DISSIPATION*, or ABERRATION, in Optics, denotes that circular space upon the retina of the eye, which is occupied by the rays of each pencil in indistinct vision: thus, if the distance of the object, or the constitution of the eye, be such, that the image falls beyond the retina, as when objects are too near; or before the retina, when the rays have not a sufficient divergency, the rays of a pencil, instead of being collected into a central point, will be dissipated over this circular space: and, all other circumstances being alike, this circle will be greater or less, according to the distances from the retina of the foci of refracted rays. But this circle causes no perceptible difference in the distinctness of vision, unless it exceed a certain magnitude; as soon as that is the case, we begin to perceive an indistinct-



ness, which increases as that circle increases, till at length the object is lost in confusion. This circle is also greater or less, according to the greater or less magnitude of the visible object: and though it be not easy to assign the diameter of the said circle, it seems very probable that vision continues distinct for all such distances, or so long as these circles, or the pencils of light from them, do not touch one another upon the retina; and the indistinctness begins when the said circles begin to interfere. It has been often observed, that a precise union of the respective rays upon the retina, is not necessary to distinct vision; but the first author who ascertained the fact beyond all doubt, was Dr. Jurin. See a variety of observations and experiments on this subject, in his Essay on Distinct and Indistinct Vision, in Smith's Optics, Appendix. In the Philos. Trans. for 1789, pa. 256, is an excellent paper on this subject by Dr. Maskelyne; in which he computes the diameter of the circle of dissipation at  $\frac{1}{1002667}$  of an inch, making it answer to an external angle of  $15^\circ$ , which he shews is very compatible with distinct vision. See also MOON, and VISION.

*Radius of DISSIPATION*, is the radius of the circle of dissipation.

**DISSOLVENT**, something that dissolves; i. e. divides, and reduces a body into its smallest parts.

**DISSOLUTION**, is a separation of the structure of a body, into small or minute parts.

According to Newton, and others, this is effected by certain powerful attractions.

**DISSONANCE**, or **DISCORD**, is a false concordance or concord; being produced by the mixture or meeting of two sounds which are disagreeable to the ear.

**DISTANCE**, properly speaking, denotes the shortest line between two points, objects, &c.

**DISTANCE**, in Astronomy, as of the sun, planets, comets, &c.

The Real Distances are found from the parallaxes of the planets, &c. See PARALLAX, and PLANET, and TRANSIT. The distance of the earth from the sun has been determined at 95 millions of miles, by the late transits of Venus; and from this one real distance, and the several relative distances, by analogy are found all the other real distances, as in the table below.

The Proportional or Relative Distances of the planets are very well deduced from the theory of gravity: for Kepler has long since discovered, and Newton has demonstrated, that the squares of their periodical times are proportional to the cubes of their distances. Kepler's Epit. Astron. lib. 4; Newton's Principia, lib. 3, phæn. 4; and Gregory's Astron. book 1, prop. 40. If therefore the mean distance of the earth from the sun be assumed, or supposed 10000, we shall then, from the foregoing analogy, and the known periodical times, obtain the relative distances of the other planets: thus,

The Periodical Revolutions in Days and Parts.

Mercury. Venus. Earth. Mars. Jupiter. Saturn. Herschel.

$87\frac{2}{3}$   $224\frac{1}{4}$   $365\frac{1}{4}$   $686\frac{3}{4}$   $4332\frac{1}{2}$   $10759\frac{2}{7}$   $30445$

Relative Mean Distances from the Sun.

3871 7233 10000 15237 52010 95401 190818

Real Distance in Millions of Miles.

37 66 95 145  $493\frac{1}{2}$   $903\frac{1}{2}$  1813

For the distances of the secondary planets from the centres of their respective primaries, see SATELLITES.

As to that of the fixed stars, as having no sensible parallax, we can do little more than guess at.

**DISTANCE of the Sun from the Moon's node, or apogee**, is an arch of the ecliptic, intercepted between the sun's true place and the moon's node, or apogee. See NODE.

*Curtate DISTANCE*. See CURTATE.


**DISTANCE of the Bastions**, in Fortification, is the side of the exterior polygon.

**Accessible DISTANCES**, in Geometry, are measured with the chain, decempeda or ten-foot rod, or the like.

**Inaccessible DISTANCES**, are found by taking bearings to them, from the two extremities of a line whose length is given. Various ways of performing this may be seen in my Treatise on Mensuration, sect. 3, on Heights and Distances.

**DISTANCE**, in Geography, is the arch of a great circle intercepted between two places.



To find the distance of two places, A and B, far remote from each other. Assume two stations, C and D, from which both the places A and B may be seen; and there, with a theodolite, observe the quantity of the angles ACD, BCD, ADC, BDC, and measure any distance as AC.



Then, in the triangle ACD, there are given the angles ACD, ADC, and the side AC; to find the side CD.—Next, in the triangle BCD, there are given the angles BCD, BDC, and the side CD; to find the side BC.—Lastly, in the triangle ABC, there are given the angle ACB, and the sides AC, CB; to find the side AB, which is the distance sought.

And in these operations, the triangles may be computed either as plane triangles, or as spherical ones, as the case may require, or according to the magnitude of the distances.

*The DISTANCE of a remote object may also be found from its height.* This admits of several cases, according as the distances are large or small, &c. 1st, Suppose that from the top of a tower at A, whose height AB is 120 feet, there be taken the angle  $BAC = 33^\circ$ , and the angle  $BAD = 64^\circ\frac{1}{2}$ , to two trees, or other objects, C, D; to find the distance between them CD, and the distance of each from the bottom of the tower at B.



First, rad. : tang.  $\angle BAD :: AB : BD = 251.585$ ,  
next, rad. : tang.  $\angle BAC :: AB : BC = 77.929$ ,

their difference is the dist.

$CD = 173.656$

2d, Suppose



2d, Suppose it be required to find the distance to which an object can be seen, by knowing its altitude; *ex. gr.* the Pike of Teneriffe, whose height is accounted 3 miles above the level of the sea, supposing the circumference of the earth 25,000 miles, or the diameter 7958 miles. Let  $FG$  be the radius = 3979,  $EF = 3$  the height of the mountain, and  $EI$  a tangent to the earth at the point  $H$ , which is the farthest distant point to which the top of the mountain  $E$  can be seen. Here in the right angled triangle  $EGH$ , are given the hypotenuse  $EG = 3982$ , and the leg  $FG = 3979$ ; to find the other leg  $HE = 154\frac{1}{2}$  miles = the distance sought nearly. Or, rather, as  $EG : GH :: \text{rad.} : \text{cofin. } \angle G = 2^\circ 13'\frac{1}{2}$ ; then as  $360^\circ : 2^\circ 13'\frac{1}{2} :: 25,000 : 154\frac{1}{2}$  miles = the arch of dist.  $HF$  sought, the same as before.

3d, If the eye, instead of being in the horizon at  $H$ , were elevated above it at  $I$ , any known height, as suppose 264 feet, or  $\frac{1}{20}$ th of a mile, as on the top of a ship's mast, &c; then the mountain can be seen much farther off along the line  $IE$ , and the distance will be the two tangents  $IH$  and  $HE$ , or rather the two arcs  $KH$  and  $HF$ . Hence, as above, as  $IG : GH :: \text{rad.} : \text{cofin. } \angle IGH = 17'\frac{1}{2}$ ; then as  $360^\circ : 17'\frac{1}{2} :: 25,000 : 20$  miles = the arc  $KH$ ; this added to the former arc  $HF = 154\frac{1}{2}$ , makes the whole arc  $KF = 174\frac{1}{2}$  miles, for the whole distance to which the top of the mountain can be seen in this case.

*Apparent DISTANCE*, in Optics, that distance which we judge an object is placed at when seen afar off, being usually very different from the true distance; because we are apt to think that all very remote objects, whose parts cannot well be distinguished, and which have no other object in view near them, are at the same distance from us, though perhaps the one is thousands of miles nearer than the other, as is the case with regard to the sun and moon.

*M. De la Hire* enumerates five circumstances, which assist us in judging of the distance of objects; viz, their apparent magnitude, the strength of the colouring, the direction of the two eyes, the parallax of the objects, and the distinctness of their small parts. On the contrary, *Dr. Smith* maintains, that we judge of distance principally, or solely, by the apparent magnitude of objects; and concludes universally, that the apparent distance of an object seen in a glass, is to its apparent distance seen by the naked eye, as the apparent magnitude to the naked eye is to its apparent magnitude in the glass: But it was long since observed by *Alhazen*, that we do not judge of distance merely by the angle under which objects are seen; and *Mr. Robins* clearly shews that *Dr. Smith's* hypothesis is contrary to fact, in the most common and simple cases. Thus, if a double convex glass be held upright before some luminous object, as a candle, there will be seen two images, one erect, and the other inverted; the first is made simply by reflexion from the nearest surface; the second by reflexion from the farther surface, the rays undergoing a refraction from the first surface both before and after the reflexion. If this glass has not too short a focal distance, when it is held near the object, the inverted image will appear larger than the other, and also nearer; but if the glass be carried off from the object, though the eye remain as near to it as before,

the inverted image will be diminished so much faster than the other, that at length it will appear much less than it, but still nearer. Here, says *Mr. Robins*, two images of the same object are seen under one view, and their apparent distances immediately compared; and it is evident that those distances have no necessary connexion with the apparent magnitude. This experiment may be made still more convincing, by sticking a piece of paper on the middle of the lens, and viewing it through a short tube. He observes farther, that the apparent magnitude of very distant objects is neither determined by the magnitude of the angle only under which they are seen, nor is the exact proportion of that angle compared with their true distance, but is compounded also with a deception concerning that distance; so that if we had no idea of difference in the distance of objects, each would appear in magnitude proportional to the angle under which it was seen; and if our apprehension of the distance were always just, our idea of their magnitude would be unvaried, in all distances; but in proportion as we err in our conception of their distance, the greater angle suggests a greater magnitude. By not attending to this compound effect, *Mr. Robins* apprehends that *Dr. Smith* was led into his mistake.

*Dr. Porterfield* has made several remarks on the five methods of judging concerning the distance of objects above recited from *M. De la Hire*; and he has also added to them one more, viz, the conformation of each eye. See *Circle of DISSIPATION*. This, he says, can be of no use to us, with respect to objects that are placed without the limits of distinct vision. But the greater or less confusion with which the object appears, as it is more or less removed from those limits, will assist the mind in judging of its distance: the more confused it appears, the farther will it be thought distant. However, this confusion has its limits; for when an object is placed at a certain distance from the eye, to which the breadth of the pupil bears no sensible proportion, the rays proceeding from a point in the object may be considered as parallel; in which case, the picture on the retina will not be sensibly more confused, though the object be removed to a much greater distance. The most universal, and often the most sure means of judging of the distance of objects, he says, is the angle made by the optic axes: our two eyes are like two different stations, by the assistance of which, distances are taken; and this is the reason why those persons who have lost the sight of one eye, so frequently miss their mark in pouring liquor into a glass, snuffing a candle, and such other actions as require that the distance be exactly distinguished. With respect to the method of judging by the apparent magnitude of objects, he observes that this can only serve when we are otherwise acquainted with their real magnitude. Thus he accounts for the deception to which we are liable in estimating distances, by any extraordinary magnitudes that terminate them; as, in travelling towards a large city, castle, or cathedral, we fancy they are nearer than they really are. Hence also, animals and small objects seen in a valley contiguous to large mountains, or on the top of a mountain or high building, appear exceedingly small. *Dr. Jurin* accounts for the last recited phenomenon, by observing that we have no distinct idea of distance




distance in that oblique direction, and therefore judge of them merely by their pictures on the eye.

Dr. Porterfield observes, with respect to the strength of colouring, that if we are assured they are of a similar colour, and one appears more bright and lively than the other, we judge that the brighter object is the nearer. When the small parts of objects appear confused, or do not appear at all, we judge that they are at a great distance, and *vice versa*; because the image of any object, or part of an object, diminishes as the distance of it increases. Finally, we judge of the distance of objects by the number of intervening bodies, by which it is divided into separate and distinct parts; and the more this is the case, the greater will the distance appear. Thus distances upon uneven surfaces appear less than upon a plane, because the inequalities do not appear, and the whole apparent distance is diminished by the parts that do not appear in it: and thus the banks of a river appear contiguous to a distant eye, when the river is low and not seen. *Accidens de la Vue*, pa. 358. *Smith's Optics*, vol. 1, pa. 52, and *Rem.* pa. 51. *Robins's Tracts*, vol. 2, pa. 230, 247, 251. *Porterfield on the Eye*, vol. 1, pa. 105, vol. 2, pa. 387. See *Priestley's Hist. of Vision*, pa. 205, and pa. 689.

**DISTANCE**, in Navigation, is the number of miles or leagues that a ship has sailed from any point or place. See **SAILING**.

**Line of DISTANCE**, in Perspective, is a right line drawn from the eye to the principal point: as the line OF, drawn between the eye at O, and the principal point F. As this is perpendicular to the plane, or table, it is therefore the distance of the eye from the table.



**Point of DISTANCE**, in Perspective, is a point in the horizontal line at the same distance from the principal point, as the eye is from the same. Such are the points P and Q, in the horizontal line PQ, whose distance from the principal point F, is equal to that of the eye from the same F.

**DISTINCT Base**, in Optics, is that distance from the pole of a convex glass, at which objects, beheld through it, appear distinct and well defined: so that the Distinct base is the same with what is otherwise called the focus.

The Distinct base is caused by the collection of the rays proceeding from a single point in the object, into a single point in the representation: and therefore concave glasses, which do not unite, but scatter and dissipate the rays, can have no real Distinct base.

**DISTINCT Vision**. See **VISION**.

**DITCH**, in Fortification, called also *Foss*, and *Moat*, is a trench dug round the rampart, or wall of a fortified place, between the scarp and counterscarp.

Ditches are either *dry*, or *wet*, that is having water in them; both of which have their particular advantages. The earth dug out of the ditch serves to raise the rampart.

The ditch in front should be of such breadth as that tall trees may not reach over it, being from 12 to 24 fathoms wide, and 7 or 8 feet deep. The ditches on

the sides are made smaller. But the most general rule is perhaps, that the dimensions of the ditch be such as that the earth dug out may be sufficient to build the rampart of a proper magnitude. The space sometimes left between the rampart and ditch, being about 6 or 8 feet, is called the *berm*, or *lift*, serving to pass and repass, and to prevent the earth from rolling into the ditch.

**DITONE**, in Music, an interval comprehending two tones, a greater and a less. The ratio of the sounds that form the Ditone, is of 4 to 5; and that of the semi-ditone, of 5 to 6.

**DITTON** (HUMPHREY), an eminent mathematician, was born at Salisbury, May 29, 1675. Being an only son, and his father observing in him an extraordinary good capacity, determined to cultivate it with a good education. For this purpose he placed him in a reputable private academy; upon quitting of which, he, at the desire of his father, though against his own inclination, engaged in the profession of divinity, and began to exercise his function at Tunbridge in the county of Kent, where he continued to preach some years; during which time he married a lady of that place.

But a weak constitution, and the death of his father, induced Mr. Ditton to quit that profession. And at the persuasion of Dr. Harris and Mr. Whiston, both eminent mathematicians, he engaged in the study of mathematics, a science to which he had always a strong inclination. In the prosecution of this science, he was much encouraged by the success and applause he received: being greatly esteemed by the chief professors of it, and particularly by Sir Isaac Newton, by whose interest and recommendation he was elected master of the new Mathematical School in Christ's Hospital; where he continued till his death, which happened in 1715, in the 40th year of his age, much regretted by the philosophical world, who expected many useful and ingenious discoveries from his assiduity, learning, and penetrating genius.

Mr. Ditton published several mathematical and other tracts, as below.—1. *Of the Tangents of Curves*, &c. *Philos. Trans.* vol. 23.

2. *A Treatise on Spherical Catoptrics*, published in the *Philos. Trans.* for 1705; from whence it was copied and reprinted in the *Acta Eruditorum* 1707, and also in the *Memoirs of the Academy of Sciences* at Paris.

3. *General Laws of Nature and Motion*; 8vo, 1705. Wolfius mentions this work, and says, that it illustrates and renders easy the writings of Galileo, Huygens, and the *Principia* of Newton. It is also noticed by La Roche, in the *Memoires de Literature*, vol. 8, pa. 46.

4. *An Institution of Fluxions, containing the first Principles, Operations, and Applications, of that admirable Method, as invented by Sir Isaac Newton*; 8vo, 1706. This work, with additions and alterations, was again published by Mr. John Clarke, in the year 1726.

5. In 1709 he published the *Synopsis Algebraica* of John Alexander, with many additions and corrections.

6. His *Treatise on Perspective* was published in 1712. In this work he explained the principles of that art mathematically; and besides teaching the methods then generally practised, gave the first hints of the new method



method afterward enlarged upon and improved by Dr. Brook Taylor; and which was published in the year 1715.

7. In 1714, Mr. Ditton published several pieces, both theological and mathematical; particularly his *Discourse on the Resurrection of Jesus Christ; and The New Law of Fluids, or a Discourse concerning the Ascent of Liquids, in exact Geometrical Figures, between two nearly contiguous Surfaces*. To this was annexed a tract, to demonstrate the impossibility of thinking or perception being the result of any combination of the parts of matter and motion: a subject much agitated about that time. To this work also was added an advertisement from him and Mr. Whiston, concerning a method for discovering the longitude, which it seems they had published about half a year before. This attempt probably cost our author his life; for although it was approved and countenanced by Sir Isaac Newton, before it was presented to the Board of Longitude, and the method has been successfully put in practice, in finding the longitude between Paris and Vienna, yet that Board then determined against it: so that the disappointment, together with some public ridicule (particularly in a poem written by Dean Swift), affected his health, so that he died the ensuing year, 1715.

In an account of Mr. Ditton, prefixed to the German translation of his *Discourse on the Resurrection*, it is said that he had published, in his own name only, another method for finding the longitude; but which Mr. Whiston denied. However, Raphael Levi, a learned Jew, who had studied under Leibnitz, informed the German editor, that he well knew that Ditton and Leibnitz had corresponded upon the subject; and that Ditton had sent to Leibnitz a delineation of a machine he had invented for that purpose; which was a piece of mechanism constructed with many wheels like a clock, and which Leibnitz highly approved of for land use; but doubted whether it would answer on ship-board, on account of the motion of the ship.

**DIVERGENT Point.** See *Virtual Focus*.

**DIVERGENT, or DIVERGING Lines,** in Geometry, are those whose distance is continually increasing.—Lines which diverge one way, converge the other way.

**DIVERGENT, or DIVERGING,** in Optics, is particularly applied to rays which, issuing from a radiant point, or having, in their passage, undergone a refraction, or reflexion, do continually recede farther from each other.

In this sense the word is opposed to convergent, which implies that the rays approach each other, or that they tend to a centre, where they intersect, and, being continued, go on diverging. Indeed all intersecting rays, or lines, diverge both ways from the centre, or point of intersection.

Concave glasses render the rays diverging; and convex ones, converging.—Concave mirrors make the rays converge; and convex ones, diverge.—It is demonstrated in Optics, that as the diameter of a pretty large pupil does not exceed  $\frac{1}{2}$  of a digit; diverging rays, flowing from a radiant point, will enter the pupil as parallel, to all intents and purposes, if the distance of the radiant from the eye amount to 40,000 feet. See *Focus, Light, and Vision*.

**DIVERGING HYPERBOLA,** is one whose legs turn

their convexities toward each other, and run out quite contrary ways. See *HYPERBOLA*.

**DIVERGING Parabola.** See *Diverging PARABOLA*.

**DIVIDEND,** in Arithmetic, is the number given to be divided by some other number, called the *divisor*. Or it is the number given to be divided, or separated, into a certain number of equal parts, viz. as many as the divisor contains units; and the number of such equal parts is called the *quotient*. Or, more generally, the dividend contains the divisor, as many times as the quotient contains unity.

The Dividend is the numerator of a fraction, whose denominator is the divisor, and the quotient is the value of the fraction. Thus,  $\frac{8}{2} = 4$ , and  $\frac{3}{4} = .75$ .

**DIVIDUAL.** By this name some authors distinguish the several parts of a dividend, from which each separate figure of the quotient is found.

**DIVING,** the art, or act of descending under water, to considerable depths, and remaining there a competent time.

The uses of Diving are very considerable, particularly in the fishing for pearls, corals, sponges, &c.

Various methods have been proposed, and engines contrived, to render the business of diving more safe and easy. The great point in all these, is to furnish the diver with fresh air, without which he must either make but a short stay, or perish.

Those who dive for sponges in the Mediterranean, help themselves by carrying down sponges dipt in oil in their mouths. But considering the small quantity of air that can be contained in the pores of a sponge, and how much that little will be contracted by the pressure of the incumbent water, such a supply cannot long subsist the diver. For it is found by experiment, that a gallon of air included in a bladder, and by a pipe reciprocally inspired and expired by the lungs, becomes unfit for respiration in little more than one minute of time. For though its elasticity be but little altered in passing the lungs, yet it loses its vivifying spirit, and is rendered effete. In effect, a naked diver, Dr. Halley assures us, without a sponge, cannot remain above two minutes inclosed in water; nor much longer with one, without suffocating; nor without long practice, near so long: ordinary persons beginning to be suffocated in about half a minute. Besides, if the depth be considerable, the pressure of the water on the vessels makes the eyes blood thotten, and frequently occasions a spitting of blood. Hence, where there has been occasion to continue long at the bottom, some have contrived double flexible pipes, to circulate air down into a cavity inclosing the diver, as with armour, both to furnish air, and to bear off the pressure of the water, and give leave to his breast to dilate upon inspiration; the fresh air being forced down one of the pipes with bellows, and returning by the other, not unlike to an artery, and vein.

But this method is impracticable when the depth exceeds three fathoms; the water embracing the bare limbs so closely, as to obstruct the circulation of the blood in them; and withal pressing so strongly on all the junctures where the armour is made tight with leather; that if there be the least defect in any of them, the water rushes in, and instantly fills the whole engine, to the great danger of the diver's life.

DIVINE-



*Diving-Bell*, is a machine contrived to remedy all these inconveniencies. In this the diver is safely conveyed to any reasonable depth, and may stay more or less time under the water, as the bell is greater or less. It is most conveniently made in form of a truncated cone, the smallest base being closed, and the larger open. It is to be poised with lead, and so suspended, that it may sink full of air, with its open basis downward, and as near as may be in a situation parallel to the horizon, so as to close with the surface of the water all at once.

Under this covercle the diver sitting, sinks down with the included air to the depth desired; and if the cavity of the vessel can contain a ton of water, a single man may remain a full hour, without much inconvenience, at five or six fathoms deep. But the lower he goes, still the more the included air contracts itself, according to the weight of the water that compresses it; so that at thirty-three feet deep, the bell becomes half full of water; the pressure of the incumbent water being then equal to that of the atmosphere; and at all other depths, the space occupied by the compressed air in the upper part of its capacity, is to the space filled with water, as thirty-three feet to the depth of the surface of the water in the bell below the common surface of it. And this condensed air, being taken in with the breath, soon insinuates itself into all the cavities of the body, and has no ill effect, provided the bell be permitted to descend so slowly as to allow time for that purpose.

One inconvenience that attends it, is found in the ears, within which there are cavities which open only outwards, and that by pores so small, as not to give admission even to the air itself, unless they be dilated and distended by a considerable force. Hence, on the first descent of the bell, a pressure begins to be felt on each ear, which, by degrees, grows painful, till the force overcoming the obstacle, what constricts these pores, yields to the pressure, and letting some condensed air slip in, presently ease ensues. The bell descending lower, the pain is renewed, and afterwards it is again eased in the same manner. But the greatest inconvenience of this engine is, that the water entering it, contracts the bulk of air into so small a compass, that it soon heats, and becomes unfit for respiration: so that there is a necessity for its being drawn up to recruit it; besides the uncomfortable abiding of the diver, who is almost covered with water.

To obviate the difficulties of the diving-bell, Dr. Halley, to whom we owe the preceding account, contrived some further apparatus, by which not only to recruit and refresh the air from time to time, but also to keep the water wholly out of it at any depth; which he effected after the following manner:

His diving-bell (plate vii, fig. 6.) was of wood, three feet wide at top, five feet at bottom, and eight feet high, containing about sixty-three cubic feet in its concavity, coated externally with lead so heavy, that it would sink empty; a particular weight being distributed about its bottom R, to make it descend perpendicularly, and no otherwise. In the top was fixed a meniscus glass D, concave downwards, like a window, to let in light from above; with a cock, as at B, to let out the hot air; and a circular seat, as at LM, for the divers to sit on: and, below, about a yard under the bell, was a stage suspended from it by three ropes, each charged

with a hundred weight, to keep it steady, and for the divers to stand upon to do their business. The machine was suspended from the mast of a ship by a sprit, which was secured by stays to the mast-head, and was directed by braces to carry it overboard clear of the side of the ship, and to bring it in again.

To supply air to this bell when under water, he had a couple of barrels, as C, holding thirty-six gallons each, cased with lead, so as to sink empty, each having a bung-hole at bottom, to let in the water as they descended, and let it out again as they were drawn up. In the top of the barrels was another hole, to which was fixed a leathern pipe, or hose, well prepared with bees wax and oil, long enough to hang below the bung-hole; being kept down by a weight appended. So that the air driven to the upper part of the barrel by the encroachment of the water, in the descent, could not escape up this pipe, unless the lower end were lifted up.

These air-barrels were fitted with tackle, to make them rise and fall alternately, like two buckets; being directed in their descent by lines fastened to the under edge of the bell: so that they came readily to the hand of a man placed on the stage, to receive them; and who taking up the ends of the pipes, as soon as they came above the surface of the water in the barrels, all the air included in the upper part of it was blown forcibly into the bell; the water taking its place.

One barrel thus received, and emptied; upon a signal given, it was drawn up, and at the same time the other let down; by which alternate succession, fresh air was furnished so plentifully, that the learned Doctor himself was one of five, who were all together in nine or ten fathoms deep of water for above an hour and a half, without the least inconvenience; the whole cavity of the bell being perfectly dry.

All the precaution he observed, was, to be let down gradually about twelve feet at a time, and then to stop, and drive out the water that had entered, by taking in three or four barrels of fresh air, before he descended farther. And, being arrived at the depth intended, he let out as much of the hot air that had been breathed, as each barrel would replace with cold, by means of the cock B, at the top of the bell, through whose aperture, though very small, the air would rush with so much violence, as to make the surface of the sea boil.

Thus, he found, any thing could be done that was required to be done underneath. And by taking off the stage, he could, for a space as wide as the circuit of the bell, lay the bottom of the sea so far dry as not to be over shoes in water. Besides, by the glass window so much light was transmitted, that, when the sea was clear, and especially when the sun shone, he could see perfectly well to write or read, much more to fasten, or lay hold of any thing under him that was to be taken up. And by the return of the air barrel he often sent up orders written with an iron pen on a plate of lead, directing how he would be moved from place to place.

At other times, when the water was troubled and thick, it would be as dark as night below; but in such cases he was able to keep a candle burning in the bell.

Dr. Halley observes, that they were subject to one inconvenience in this bell; they felt at first a small pain in



in their ears, as if the end of a tobacco pipe were thrust into them; but after a little while there was a small puff of air, with a little noise, and they were easy.

This he supposes to be occasioned by the condensed air shutting up a valve leading from some cavity in the ear, full of common air; but when the condensed air pressed harder, it forced the valve to yield, and filled every cavity. One of the divers, in order to prevent this pressure, stopped his ear with a pledget of paper; which was pushed in so far, that a surgeon could not extract it without great difficulty.

The same author intimates, that by an additional contrivance he has found it practicable for a diver to go out of the bell to a good distance from it; the air being conveyed to him in a continued stream by small flexible pipes, which serve him as a clue to direct him back again to the bell. For this purpose, one end of these pipes, kept open against the pressure of the sea, by a small spiral wire, and made tight without by painted leather, and sheep's guts drawn over it, being open, was fastened in the bell, as at P, to receive air, and the other end was fixed to a leaden cap on the man's head, reaching down below his shoulders, open at bottom, to serve him as a little bell, full of air, for him to breathe at his work, which would keep out the water from him, when at the level of the great bell, because of the same density as the air in the great bell. But when he stooped down lower than the level of the great bell, he shut the cock F, to cut off the communication between the two bells. Phil. Transf. abr. vol. iv. part ii. p. 188, &c. vol. vi. p. 550, &c.

The air in this bell would serve him for a minute or two; and he might instantly change it, by raising himself above the great bell, and opening the cock F. The diver was furnished with a girdle of large leaden weights, and clogs of lead for the feet, which, with the weight of the leaden cap, kept him firm on the ground; he was also well clothed with thick flannels, which being first made wet, and then warmed in the bell by the heat of his body, kept off the chill of the cold water for a considerable time, when he was out of the bell.

Mr. Martin Triewald, F. R. S. and military architect to the king of Sweden, contrived to construct a diving-bell on a smaller scale, and less expence, than that of Dr. Halley, and yet capable of answering the same intents and purposes. This bell, AB (fig. 7.) sinks with leaden weights DD, suspended from the bottom of it. It is made of copper, and tinned all over on the inside; three strong convex lenses GGG, defended by the copper lids HHH, illuminate this bell. The iron plate E serves the diver to stand upon, when he is at work; this is suspended by chains FFF, at such a distance from the bottom of the bell, that when he stands upright, his head is just above the water in the bell, where he has the advantage of air fitter for respiration, than when he is much higher up; but as there is occasion for the diver to be wholly in the bell, and consequently his head in the upper part of it, Mr. Triewald has contrived, that, even there, after he has breathed the hot air as long as he well can, by means of a spiral copper tube placed close to the inside of the bell, he may draw the cooler and fresher air from the lowermost parts; for which purpose a flexible leather

pipe, about two feet long, is fixed to the upper end of the tube at *b*; and to the other end of the pipe is fastened an ivory mouth-piece, for the diver to hold in his mouth, by which to respire the air from below. We shall only remark, that as air rendered effete by respiration is somewhat heavier than common air, it must naturally subside in the bell; but it may probably be restored by the agitation of the sea-water, and thus become fitter for respiration. See Fixed Air. Phil. Transf. abr. vol. viii. p. 634. Or Defaguliers's Exper. Phil. vol. ii. p. 220, &c.

The famous Corn. Drebell had an expedient in some respects superior even to the diving bell, if what is related of it be true. He contrived not only a vessel to be rowed under water, but also a liquor to be carried in the vessel, which supplied the place of fresh air.

The vessel was made for king James I. carrying twelve rowers, besides the passengers. It was tried in the river Thames; and one of the persons in that submarine navigation, then living, told it one, from whom Mr. Boyle had the relation.

As to the liquor, Mr. Boyle assures us, he discovered by a physician, who married Drebell's daughter, that it was used from time to time, when the air in that submarine boat was clogged by the breath of the company, and rendered unfit for respiration: at which time, by unstopping the vessel full of this liquor, he could speedily restore to the troubled air such a proportion of vital parts, as would make it serve again a good while. The secret of this liquor Drebell would never disclose to above one person, who himself assured Mr. Boyle what it was. Boyle's Exp. Phys. Mech. of the Spring of the Air.

We have had many projects of diving machines, and diving ships of various kinds, which have proved abortive.

*DIVING-Bladder*, a term used by Borelli for a machine which he contrived for diving under the water to great depths, with great facility, which he prefers to the common diving-bell. The vesica, or bladder, as it is usually called, is to be of brass or copper, and about two feet in diameter. This is to contain the diver's head, and is to be fixed to a goat's skin habit, exactly fitted to the shape of the body of the person. Within this vesica there are pipes, by means of which a circulation of air is contrived; and the person carries an air pump by his side, by means of which he may make himself heavier or lighter, as the fishes do, by contracting or dilating their air bladder: by this means, the objections all other diving machines are liable to are obviated, and particularly that of the air; the moisture by which it is clogged in respiration, and by which it is rendered unfit for the same use again, being here taken from it by its circulation through the pipes, to the sides of which it adheres, and leaves the air as free as before. Borelli Opera Posthuma.

*DIVISIBILITY*, a property in quantity, body, or extension, by which it becomes separable into parts; either actually, or at least mentally.

Such divisibility is infinite, if not actually, at least potentially; for no part can be conceived so small, but another may be conceived still smaller; for every part of matter must have some finite extension, and that extension may be bisected, or otherwise divided; for the  
same







the 3's in 8 are 2 times, and 2 remaining. All the numbers expressing how often 3 is contained in each of those parts, are to be written down according to the order of the parts of the dividend, or from left to right, and separated from the dividend itself by a crooked line, thus :

Divisor Dividend Quotient  
3 ) 6758 ( 2252  $\frac{2}{3}$

It appears therefore, that 3 is contained 2252 times in 6758, with 2 remaining over; or that 6758 being divided into 3 parts, each part will be 2252  $\frac{2}{3}$ , viz, the figures of the quotient before found, together with the fraction  $\frac{2}{3}$  formed of the remainder and the divisor.

When the divisor is a single digit, or even as large as the number 12, the division is easily performed by setting down only the quotient as above. But when the divisor is a larger number, it is necessary to set down the several remainders and products &c. This process may be seen at large in most books of arithmetic, as well as various contractions adapted to particular cases: such as, 1st, when the divisor has any number of ciphers at the end of it, they are cut off, as well as the same number of figures from the end of the dividend, and then the work is performed without them both, annexing only the figures last cut off, to the last remainder; 2d, when the divisor is equal to the product of several single digits, it is easier to divide successively by those digits, instead of the divisor at once; 3d, when it is required to continue a quotient to a great many places of figures, as in decimals, a very expeditious method of performing it, is as follows: Suppose it were required to divide 1 by 29, to a great many places of decimals. Adding ciphers to the 1, first divide 10000 by 29 in the common way, till the remainder become a single figure, and annex the fractional supplement to complete the quotient, which gives  $\frac{1}{29} = 0.03448\frac{6}{29}$ : next multiply each of these by the numerator 8, so shall  $\frac{8}{29} = 0.27584\frac{6}{29}$  or rather  $0.27586\frac{6}{29}$ ; which figures substituted instead of the fraction  $\frac{6}{29}$  in the first value of  $\frac{1}{29}$ , it becomes  $\frac{1}{29} = 0.0344827586\frac{6}{29}$ : again, multiply both of these by the last numerator 6, and it will be  $\frac{6}{29} = 0.2068965517\frac{6}{29}$ ; which figures substituted for  $\frac{6}{29}$  in the last-found value of  $\frac{1}{29}$ , it becomes  $\frac{1}{29} = 0.03448275862068965517\frac{6}{29}$ : and again, multiplying these by the numerator 7, gives  $\frac{7}{29} = 0.24137931034482758620\frac{6}{29}$ ; which figures substituted instead of  $\frac{6}{29}$  in the last-found value of  $\frac{1}{29}$ , it becomes  $\frac{1}{29} = 0.0344827586206896551724137931034482758620\frac{6}{29}$  and so on; where every operation will at least double the number of figures before found by the last one.

*Proof of Division.* In every example of division, unity is always in the same proportion to the divisor, as the quotient is to the dividend; and therefore the product of the divisor and quotient is equal to the product of 1 and the dividend, that is, the dividend itself. Hence, to prove division, multiply the divisor by the quotient, to the product add the remainder, and the sum will be equal to the dividend when the work is right; if not, there is a mistake.

*Division, in Decimal Fractions,* is performed the same way as in integers, regard being had to the number of decimals, viz, making as many in the quotient as those of the dividend exceed those in the divisor.

*Division, in Vulgar Fractions,* is performed by dividing the numerators by each other, and the denominators by each other, if they will exactly divide; but if not, then the dividend is multiplied by the reciprocal of the divisor, that is, having its terms inverted; for, taking the reciprocal of any quantity, converts it from a divisor to a multiplier, and from a multiplier to a divisor. For ex.  $\frac{1}{4} \div \frac{5}{6}$  by  $\frac{6}{5}$  gives  $\frac{3}{2}$ , by dividing the numerators and denominators; but  $\frac{1}{4} \div \frac{5}{6}$  by  $\frac{4}{3}$  is the same as  $\frac{1}{4} \times \frac{4}{3} = \frac{1}{3}$ , which is  $= \frac{2}{6}$ . Where  $\times$  is the sign of multiplication, and the character  $\div$  is the mark of division. Or division is also denoted like a vulgar fraction; so 3 divided by 2, is  $\frac{3}{2}$ .

*Division, in Algebra, or Species,* is performed like that of common numbers, either making a fraction of the dividend and divisor, and cancelling or dividing by the terms or parts that are common to both; or else dividing after the manner of long division, when the quantities are compound ones. Thus,

$ab$  divided by  $a$ , gives  $b$  for the quotient:

and  $12ab$  divided by  $4b$ , gives  $3a$  for the quotient:

and  $16abc^2$  divided by  $8ac$ , gives  $2bc$ :

and  $a$  divided by  $3b$ , gives  $\frac{a}{3b}$ :

and  $15abc^3$  divided by  $12bc^2$ , gives  $\frac{15abc^3}{12bc^2} = \frac{5ac}{4}$

and  $a^2 - b^2$  by  $a + b$  gives  $a - b$ ; thus

$$\begin{array}{r} a + b \ ) \ a^2 - b^2 \ ( \ a - b \\ \underline{a^2 + ab} \end{array}$$

$$\underline{-ab - b^2}$$

$$\underline{-ab - b^2}$$

again, to divide  $x^6 - 8x^4 - 124x^2 - 64$  by  $x^2 - 16$ ;

$$\begin{array}{r} x^2 - 16 \ ) \ x^6 - 8x^4 - 124x^2 - 64 \ ( \ x^4 + 8x^2 + 4 \\ \underline{x^6 - 16x^4} \end{array}$$

$$\underline{8x^4 - 124x^2}$$

$$\underline{8x^4 - 128x^2}$$

$$\underline{4x^2 - 64}$$

$$\underline{4x^2 - 64}$$

In some cases, the quotient will run out to an infinite series; and then, after continuing it to any certain number of terms, it is usual to annex, by way of a fraction, the remainder with the divisor set under it.

It is to be noted that, in dividing any terms by one another, if the signs be both alike, either both plus, or both minus, the sign of the quotient will be plus; but when the signs are different, the one plus and the other minus, the sign of the quotient will be minus.

*Division by Logarithms.* See LOGARITHMS.

*Division of Mathematical Instruments.* See GRADUATION, and MURAL Arc or Quadrant.

*Division in Music,* is the dividing the interval of an octave into a number of lesser intervals.

*Division by Napier's Bones.* See NAPIER'S Bones.

*Division of Powers,* is performed by subtracting their exponents. Thus,  $a^6 \div a^4$  is  $= a^2$ ; and

$$4a^{\frac{3}{4}}b^2 \div 2a^{\frac{1}{2}}b^{\frac{4}{3}} \text{ is } = 2a^{\frac{1}{4}}b^{\frac{2}{3}}.$$



*DIVISION of Proportion*, is comparing the difference between the antecedent and consequent, with either of them. Thus,

$$\begin{array}{l} \text{if} \quad \left\{ \begin{array}{l} a : b :: c : d, \\ 12 : 4 :: 6 : 2, \end{array} \right. \\ \text{then by division} \left\{ \begin{array}{l} a - b : a :: c - d : c, \\ 8 : 12 :: 4 : 6, \end{array} \right. \\ \text{or} \quad \left\{ \begin{array}{l} a - b : b :: c - d : d, \\ 8 : 4 :: 4 : 2. \end{array} \right. \end{array}$$

*DIVISOR*, is the dividing number; or that which shews how many parts the dividend is to be divided into.

*Common DIVISOR*. See *COMMON Divisor*.

*DIURNAL*, something relating to the day; in opposition to nocturnal, relating to the night.

*DIURNAL Arch*, is the arch described by the sun, moon, or stars, between their rising and setting.

*DIURNAL Circle*, is the apparent circle described by the sun, moon, or stars, in consequence of the rotation of the earth.

*DIURNAL Motion of a Planet*, is so many degrees and minutes &c as any planet moves in 24 hours.

*The DIURNAL Motion of the Earth*, is its rotation round its axis, the duration of which constitutes the natural day.—The reality of the diurnal rotation of the earth is now past all dispute.

*DIURNAL Parallax*. See *PARALLAX*.

*DIURNAL* is also used in speaking of what belongs to the nycthemeron, or natural day of 24 hours: in which sense it is opposed to annual, menstrual, &c.

The diurnal phenomena of the heavenly bodies are solved from the diurnal revolution of the earth; that is, from the rotation of the earth round its own axis in 24 hours. This rotation is equable, and from west to east, about an axis whose inclination to the ecliptic is now  $66^{\circ} 32'$ . Since the earth is an opaque body, that small part of its surface which comes at the same time under the confined view of the spectator, though really spherical, will seem to be extended like a plane: and the eye, taking a view of the heavens all around, defines a concave spherical superficies, concentric with the earth, or rather with the eye, which the said plane of the earth's superficies will divide into two equal parts, the one of which is visible, but the other, because of the earth's opacity, hid from the view.

And as the earth revolves about its axis, the spectator, standing upon it, together with the said plane he stands upon, called his horizon, dividing the visible from the invisible hemisphere of the heavens, is carried round the same way, viz, towards the east. From hence it is, that the sun and stars, placed towards the east, being before hid, now become visible, the horizon as it were sinking below them; and the stars &c towards the west are covered or hid, and become invisible, the horizon being elevated above them. So that the former stars, to the spectator, who reckons the place he stands on as immoveable, will appear to ascend above the horizon, or rise; and the latter to descend below the horizon, or set.

Since the earth, with the horizon of the spectator fixed to it, continues to move always towards the same parts, and about the same axis equally; all bodies, and

all phenomena, that do not partake of the said motion, (that is, all such things as are entirely separate from the earth) will seem to move in the same time uniformly, but towards the opposite parts, or from east to west: and every one of these objects, according to sense, will describe the circumference of a circle, whose plane is perpendicular to the axis of the earth. And because all these circles, together with the visible objects describing them, appear to be in the concave spherical superficies of the heavens, every visible object will seem to describe a greater or less circle, according to its greater or less distance from the poles, or extremities of the earth's axis produced; the middle circle between these poles, called the equator, is consequently the greatest.

It may farther be observed, that whereas, by the diurnal revolution of the earth, all the several luminaries seem to move in the heavens from east to west, hence this seeming diurnal motion of the celestial lights is called their common motion, as being common to all of them. Besides which, all the luminaries, except the sun, have a proper motion; from which arise their proper phenomena: as for the proper phenomena of the sun, they likewise seem to arise from the proper motion of the sun; though they are really produced by another motion, which the earth has, and by which it moves round the sun once every year, and thence called the annual motion of the earth.

*DODECAGON*, a regular polygon of 12 equal sides and angles.

If the side of a Dodecagon be 1, its area will be equal to 3 times the tang. of  $75^{\circ} = 3 \times 2 + \sqrt{3} = 11.1961524$  nearly; and, the areas of plane figures being as the squares of their sides, therefore 11.1961524 multiplied by the square of the side of any Dodecagon, will give its area. See my *Mensuration*, pa. 114, 2d ed.

*To inscribe a Dodecagon in a given Circle*. Carry the radius 6 times round the circumference, which will divide it into 6 equal parts, or will make a hexagon; then bisect each of those parts, which will divide the whole into 12 parts, for the Dodecagon. See also other methods of describing the same figure in my *Mensur.* pa. 26, &c. See *POLYGON*.

*DODECAHEDRON*, one of the Platonic bodies, or five regular solids, being contained under a surface composed of twelve equal and regular pentagons.

*To form a Dodecahedron*. See *Regular Body*.

If the side, or linear edge, of a Dodecahedron be  $s$ , its surface will be

$$15 s^2 \sqrt{1 + \frac{2}{5} \sqrt{5}} = 20.6457788 s^2$$

$$\text{and its solidity } 5 s^3 \sqrt{\frac{47 + 21 \sqrt{5}}{40}} = 7.66311896 s^3$$

If the radius of the sphere that circumscribes a Dodecahedron be  $r$ , then is

$$\begin{aligned} \text{its side or linear edge} &= \frac{\sqrt{15 - \sqrt{3}}}{3} r, \\ \text{its superficies} &= 10 r^2 \sqrt{2 - \frac{2}{5} \sqrt{5}}, \\ \text{and its solidity} &= \frac{20 r^3}{3} \sqrt{\frac{3 + \sqrt{5}}{30}}. \end{aligned}$$

The side of a Dodecahedron inscribed in a sphere, is equal



equal to the greater part of the side of a cube inscribed in the same sphere, and cut according to extreme and mean proportion.

If a line be cut according to extreme and mean proportion, and the lesser segment be taken for the side of a Dodecahedron, the greater segment will be the side of a cube inscribed in the same sphere.

The side of the cube is equal to the right line which subtends the angle of a pentagon of the Dodecahedron, inscribed in the same sphere. See POLYHEDRON; also my Mensur. pa. 253, &c.

DODECATEMORY, the 12 houses or parts of the zodiac of the primum mobile. Also the 12 signs of the zodiac are sometimes so called, because they contain each the 12th part of the zodiac.

DOG, a name common to two constellations, called the Great and Little Dog; but more usually Canis Major, and Canis Minor.

DOVE, is a round, vaulted, or arched roof, of a church, hall, pavilion, vestibule, stair-case, &c, by way of crowning, or acroter.

DOMINICAL Letter, otherwise called the SUNDAY Letter, is one of these first seven letters of the alphabet ABCDEFG, used in almanacs &c, to mark or denote the Sundays throughout the year.

The reason for using seven letters, is because that is the number of days in a week; and the method of using them is this: the first letter A is set opposite the 1st day of the year, the 2d letter B opposite the 2d day of the year, the 3d letter C opposite the 3d day of the year, and so on through the seven letters; after which they are repeated over and over again, all the way to the end of the year, the letter A denoting the 8th day, the letter B the 9th, &c. Then whichever of the letters so placed, falls opposite the first Sunday in the year, the same letter, it is evident, will fall opposite every future Sunday throughout the year, because the number of the letters is the same as the number of days in the week, being both 7 in number; that is, in common years; for as to leap years, an interruption of the order takes place in them. For, on account of the intercalary day, either the letters must be thrust out of their places for the whole year afterwards, so as that the letter, for ex. which answers to the 1st of March, shall likewise answer to the 2d &c; or else the intercalary day must be denoted by the same letter as the preceding one. This latter expedient was judged the better, and accordingly all the Sundays in the year after the intercalary day have another Dominical letter.

The Dominical letters were introduced into the calendar by the primitive christians, instead of the nundinal letters in the Roman calendar; and in this manner were those seven letters set opposite the days of the year, to denote the days in the week, in most of our common almanacs, till the year 1771, when the initial letters of the days of the week were generally introduced instead of them, excepting the Sunday letter itself, which is still retained.

From the foregoing account it follows that,

1st, As the common year consists of 365 days, or 52 weeks, and one day over; the letters go one day backwards every common year: so that in such a year, if the beginning or first day fall on a Sunday, the next year it will fall on Saturday, the next on Friday, and

so on. Consequently, if G be the Dominical letter for the present year, F will be that for the next year; and so on, in a retrograde order.

2d, As the bissextile or leap year consists of 366 days, or 52 weeks, and 2 days over, the beginning of the year next after bissextile goes back 2 days. Whence, if in the beginning of the bissextile year, the Dominical letter were G, that of the following year will be E.

3d, Since in leap-years the intercalary day falls on the 24th of February, in which case the 24th and 25th days are considered as one day, and denoted by the same letter; after the 24th day of February the Dominical letter goes back one place: thus, if in the beginning of the year the Dominical letter be G, it will afterwards change to the letter F for the remaining Sundays of the year. With us however, this day is now added, at the end of February, and from thence it is that the change takes place.

4th, As every fourth year is bissextile, or leap-year, and as the number of letters is 7; the same order of Dominical letters only returns in 4 times 7, or 28 years; which without the interruption of bissextiles, would return in 7 years.

5th, Hence the invention of the solar cycle of 28 years; upon the expiration of which the Dominical letters are restored successively to the same days of the month, or the same order of the letters returns.

To find the DOMINICAL Letter of any given year. Find the cycle of the sun for that year, as directed under cycle; and the Dominical letter is found corresponding to it. When there are two letters, the proposed year is bissextile; the former of them serving till the end of February, and the latter for the rest of the year.

The Dominical letter for any year of the present century may be found by this canon:

*Divide the odd years, their fourth and 4, by 7,  
What is left take from 7, the letter is given.*

Thus, for the year 1794, the odd years 94  
their 4th 23  
and - 4

divided by 7) 121 (17

remains 2  
from 7

leaves 5

which answers to E the 5th letter in the alphabet.

The Dominical letter may be found universally, for any year of any century, thus:

*Divide the centuries by 4; and twice what does remain  
Take from 6; and then add to the number you gain  
Their odd years and their 4th; which dividing by 7,  
What is left take from 7, the letter is given.*

Thus, for the year 1878 the letter is F.

For the centuries 18 divided by 4, leave 2; the double of which taken from 6 leaves 2 again; to which add the odd years 78, and their 4th part 19, the sum 99 divided by 7 leaves 1; which taken from 7, leaves 6 answering to F the 6th letter in the alphabet.

By the reformation of the calendar under pope Gregory the 13th, the order of the Dominical letters was again disturbed in the Gregorian year; for the year



1582, which had G for its Dominical letter at the beginning; by retrenching 10 days after the 4th of October, came to have C for its Dominical letter: by which means the Dominical letter of the ancient Julian calendar is 4 places before that of the Gregorian, the letter A in the former answering to D in the latter.

DONJON, in Fortification, usually denotes a large strong tower, or redoubt, of a fortress, where the garrison may retreat in case of necessity, and capitulate with greater advantage. See DUNGEON.

DORADO, a southern constellation, not visible in our latitude; called also *Xiphias*, or the *Sword-fish*. The stars of this constellation, in Sharp's catalogue, are six.

DORIC Order, of Architecture, is the second of the five orders, being placed between the Tuscan and the Ionic. The Doric seems the most natural, and best proportioned, of all the orders; all its parts being founded on the natural position of solid bodies; for which reason it is most proper to be used in great and massy buildings, as the outside of churches and public places.

The Doric order has no ornaments on its base, nor its capital. Its column is 8 diameters high, and its freeze is divided between triglyphs and metopes.

DORMER, or DORMANT, in Architecture, denotes a window made in the roof of a building, or above the entablature; being raised upon the rafters.

DOUBLE Aspect, Bastion, Concave, Convex, Cone, Descant, Eccentricity, Position, Ratio, Tenaille, &c. See the respective words.

DOUBLE Horizontal Dial, one with a double gnomon, the one pointing out the hour on the outer circle, the other the hour on the stereographic projection drawn upon it. This dial finds the meridian, the hour, the sun's place, rising, setting, &c, and many other problems of the globe.

DOUBLE Point, in the Higher Geometry, is a point which is common to two parts or legs or branches of some curve of the 2d or higher order: such as, an infinitely small oval, or a cusp, or the cruciform intersection, &c, of such curves. See Newton's *Enumeratio Linearum &c, de Curvarum Punctis Duplicibus*.

DOUBLING a Cape, or Point of Land, in Navigation, signifies the coming up with it, passing by it, and leaving it behind the ship.

The Portuguese pretend that they first doubled the Cape of Good Hope, under their admiral Vasquez de Gama; but there are accounts in history, particularly in Herodotus, that the Egyptians, Carthaginians, &c, had done the same long before them.

DOUCINE, in Architecture, is an ornament on the highest part of the cornice, or a moulding cut in the figure of a wave, half convex, and half concave.

DOVE-tail, in Carpentry, is a method of fastening boards or timbers together, by letting or indenting one piece into another, with a Dove-tail joint, or a joint in form of a Dove's tail.

DRACHM, or DRAM, is the name of a small weight used with us, and is of two kinds, viz, the 8th part of an ounce in Apothecaries weight, and the 16th part of an ounce in Avoirdupois weight.

DRACO, the DRAGON, a constellation of the northern hemisphere, whose origin is variously fabled by the Greeks; some of them representing it as the Dragon

which guarded the Hesperian fruit, or golden apples, but being killed by Hercules, Juno, as a reward for its faithful services, took it up to heaven, and so formed this constellation; while others say, that in the war of the giants, this Dragon was brought into the combat, and opposed to Minerva, when the goddess taking the Dragon in her hand, threw him, twisted as he was, up to the skies, and fixed him to the axis of the heavens, before he had time to unwind his contortions.

The stars in this constellation are, according to Ptolemy, 31; according to Tycho, 32; according to Hevelius, 40; according to Bayer, 33; and according to Flamsteed, 80.

DRAGON, in Astronomy. See DRACO.

DRAGON's Head, and Tail, are the nodes of the planets, but more particularly of the moon, being the points in which the ecliptic is intersected by her orbit, in an angle of about  $5^{\circ} 18'$ .

One of these points looks northward, the moon beginning then to have north latitude; and the other southward, where she commences south latitude; the former point being represented by the knot  $\Omega$  for the head, and the other by the same reversed, or  $\varpi$  for the tail. And near these points it is that all eclipses of the sun and moon happen.

This deviation of the path of the moon from the ecliptic seems, according to the fancy of some, to make a figure like that of a dragon, whose belly is at the part where she has the greatest latitude; the intersections representing the head and tail, from which resemblance the denomination arises.

These intersections are not always in the same two points of the ecliptic, but shift by a retrograde motion, at the rate of  $3' 11''$  per day, and completing their circle in 18 years 225 days.

DRAGON-Beams, in Architecture, are two strong braces or struts, standing under a breast-summer, and meeting in an angle on the shoulder of the king-piece.

DRAM. See DRACHM.

DRAUGHT-Compasses, those provided with several moveable points, to draw fine draughts in architecture &c. See COMPASS.

DRAUGHT-Hooks, are large hooks of iron, fixed on the cheeks of a gun-carriage, two on each side, one near the trunnion hole, and the other at the train.

DRAW-Bridge, a bridge made after the manner of a floor, to be drawn up, or let down, as occasion requires, before the gate of a town or castle.

DRIFT, in Navigation, denotes the angle which the line of a ship's motion makes with the nearest meridian, when she drives with her side to the wind and waves, and is not governed by the power of the helm; and also the distance which the ship drives on that line, so called only in a storm.

DRIP, in Architecture. See LARMIER.

DRY-Moat. See MOAT.

DUCTILITY, a property of certain bodies, by which they are capable of being beaten, pressed, drawn, or stretched forth, without breaking; or by which they are capable of great alterations in their figure and dimensions, and of gaining in one way as they lose in another.

Such are metals, which, being urged by the hammer, gain in length and breadth what they lose in thickness; or, being drawn into wire through holes in iron, grow longer



longer as they become more slender. Such also are gums, glues, resins, and some other bodies; which, though not malleable, may yet be denominated ductile, in as much as, when softened by water, fire, or some other menstruum, they may be drawn into threads.

Some bodies are ductile both when they are hot and cold, and in all circumstances: such are metals, and especially gold and silver; other bodies are ductile only when they have a certain degree of heat; such as glass, and wax, and such like substances: others again are ductile only when cold, and brittle when hot; as some kinds of iron, viz, those called by workmen redshort, as also brass, and some metallic alloys.

The cause of ductility is very obscure; as depending much on hardness, a quality whose nature we know very little about. It is true, it is usual to account for hardness from the force of attraction between the particles of the hard body; and for ductility, from the particles of the ductile body being, as it were jointed, and entangled with each other. But without dwelling on any fanciful hypotheses about ductility, we may amuse ourselves with some truly amazing circumstances and phenomena of it, in the instances of gold, glass, and spider's-webs. Observing however that the ductility of metals decreases in the following order: gold, silver, copper, iron, tin, lead.

*DUCTILITY of Gold.* One of the properties of gold is, to be the most ductile of all bodies; of which the gold beaters and gold wire-drawers, furnish us with abundant proof.

Fa. Merfenne, M. Rohault, Dr. Halley, &c, have made computations of it: but they trusted to the reports of the workmen. M. Reaumur, in the *Memoires de l'Academie Royale des Sciences*, an. 1713, took a surer way; he made the experiment himself. A single grain of gold, he found, even in the common gold leaf, used in most of our gildings, is extended into 36 and a half square inches; and an ounce of gold, which, in form of a cube, is not half an inch either high, broad, or long, is beat under the hammer into a surface of 146 and a half square feet; an extent almost double to what could be done in former times. In Fa. Merfenne's time, it was deemed prodigious, that an ounce of gold should form 1600 leaves; which, together, only made a surface of 105 square feet.

But the distension of gold under the hammer (how considerable soever) is nothing to that which it undergoes in the drawing-iron. There are gold leaves, in some parts scarce the  $\frac{1}{370000}$  part of an inch thick; but  $\frac{1}{370000}$  part of an inch is a considerable thickness, in comparison of that of the gold spun on silk in our gold thread.

To conceive this prodigious ductility, it is necessary to have some idea of the manner in which the wire drawers proceed. The wire, and thread we commonly call gold thread, &c, (which is only silver wire gilt, or covered over with gold), is drawn from a large ingot of silver, usually about thirty pounds weight. This they round into a cylinder, or roll, about an inch and a half in diameter, and twenty-two inches long, and cover it with the leaves prepared by the gold beater, laying one over another, till the cover is a good deal thicker than that in our ordinary gilding; and yet, even then, it is

very thin; as will be easily conceived from the quantity of gold that goes to gild the thirty pounds of silver: two ounces ordinarily do the business; and, frequently, little more than one.

In effect, the full thickness of the gold on the ingot rarely exceeds  $\frac{1}{100}$  or  $\frac{1}{500}$  part; and, sometimes not  $\frac{1}{1000}$  part of an inch.

But this thin coat of gold must be yet vastly thinner: the ingot is successively drawn through the holes of several irons, each smaller than the other, till it be as fine as, or finer than a hair. Every new hole lessens its diameter; but it gains in length what it loses in thickness; and, of consequence, increases in surface: yet the gold still covers it; it follows the silver in all its extension, and never leaves the minutest part bare, not even to the microscope. Yet, how inconceivably must it be attenuated while the ingot is drawn into a thread, whose diameter is 9000 times less than that of the ingot.

M. Reaumur, by exact weighing, and rigorous calculation, found, that one ounce of the thread was 3232 feet long; and the whole ingot 1163520 feet, Paris measure, or 96 French leagues; equal to 1264400 English feet, or 240 miles English; an extent which far surpasses what Fa. Merfenne, Furetiere, Dr. Halley, &c, ever dreamt of.

Merfenne says, that half an ounce of the thread is 100 toises, or fathoms long; on which footing, an ounce would only be 1200 feet: whereas, M. Reaumur finds it 3232. Dr. Halley makes 6 feet of the wire one grain in weight, and one grain of the gold 98 yards; and, consequently, the ten thousandth part of a grain, above one third of an inch. The diameter of the wire he found one-186th part of an inch; and the thickness of the gold one-154500th part of an inch. But this, too, comes short of M. Reaumur; for, on this principle, the ounce of wire would only be 2680 feet.

But the ingot is not yet extended to its full length. The greatest part of our gold thread is spun, or wound on silk; and, before it is spun, they flat it, by passing it between two rolls, or wheels of exceedingly well polished steel; which wheels, in flattening it, lengthen it by above one seventh. So that our 240 miles are now got to 274.

The breadth, now, of these laminæ, or plates, M. Reaumur finds, is only one-8th of a line, or one-96th of an inch; and their thickness one-3072d. The ounce of gold, then, is here extended to a surface of 1190 square feet; whereas, the utmost the gold beaters can do, we have observed, is to extend it to 146 square feet. But the gold, thus exceedingly extended, how thin must it be! From M. Reaumur's calculus, it is found to be one-175000th of a line, or one-2100000th of an inch; which is scarce one-13th of the thickness of Dr. Halley's gold.

But he adds, that this supposes the thickness of the gold every where equal, which is no ways probable; for in beating the gold leaves, whatever care they can bestow, it is impossible to extend them equally. This we easily find, by the greater opacity of some parts than others; for, where the leaf is thickest, it will gild the wire the thickest.

M. Reaumur,



M. Reaumur, computing what the thickness of the gold must be where thinnest, finds it only one-3150000th part of an inch. But what is the one-3150000th part of an inch? Yet this is not the utmost ductility of gold: for, instead of two ounces of gold to the ingot, which we have here computed upon, a single one might have been used; and, then, the thickness of the gold, in the thinnest places, would only be the 6300000th part of an inch.

And yet, as thin as the plates are, they might be made twice as thin, yet still be gilt; by only pressing them more between the flatter's wheels, they are extended to double the breadth and proportionably in length. So that their thickness, at last, will be reduced to one thirteen or fourteen millionth part of an inch.

Yet, with this amazing thinness of the gold, it is still a perfect cover for the silver: the best eye, or even the best microscope, cannot discover the least chasm, or discontinuity. There is not an aperture to admit alcohol of wine, the subtlest fluid in nature, or even light itself, unless it be owing to cracks occasioned by repeated strokes of the hammer. Add, that if a piece of this gold thread, or gold plate be laid to dissolve in aquafortis, the silver will be all excavated, or eat out, and the gold left entire, in little tubules.

It should be observed, that gold, when it has been struck for some time by a hammer, or violently compressed, as by gold wire drawers, becomes more hard, elastic and stiff, and less ductile, so that it is apt to be cracked or torn: the same thing happens to the other metals by percussion and compression. But ductility and tractability may be restored to metals in that state, by annealing them, or making them red hot. Gold seems to be more affected by percussion and annealing, than any other metals.

As to the *DUCTILITY of soft bodies*, it is not yet carried to that pitch. The reader, however, must not be surprised that, among the ductile bodies of this class, we give the first place to the most brittle of all others, glass.

*DUCTILITY of Glass.* We all know, that, when well penetrated with the heat of the fire, the workmen can figure and manage glass like soft wax; but what is most remarkable, it may be drawn, or spun out into threads exceedingly fine and long.

Our ordinary spinners do not form their threads of silk, flax, or the like, with half the ease, and expedition, as the glass spinners do threads of this brittle matter. We have some of them used in plumes for children's heads, and divers other works, much finer than any hair, and which bend and wave like it with every wind.

Nothing is more simple and easy than the method of making them: there are two workmen employed; the first holds one end of a piece of glass over the flame of a lamp; and, when the heat has softened it, a second operator applies a glass hook to the metal thus in fusion; and, withdrawing the hook again, it brings with it a thread of glass, which still adheres to the mass: then, fitting his hook on the circumference of a wheel about two feet and a half in diameter, he turns the wheel as fast as he pleases; which, drawing out the thread winds it on its rim; till, after a certain number of revolutions, it is covered with a skain of glass thread.

The mass in fusion over the lamp diminishes insensibly: being wound out, as it were, like a pelatoon, or clue of silk, upon the wheel; and the parts, as they recede from the flame, cooling, become more coherent to those next to them; and this by degrees: the parts nearest the fire are always the least coherent, and, of consequence, must give way to the effort the rest make to draw them towards the wheel.

The circumference of these threads is usually a flat oval, being three or four times as broad as thick: some of them seem scarce bigger than the thread of a silk worm, and are surprisingly flexible. If the two ends of such threads be knotted together, they may be drawn and bent, till the aperture, or space in the middle of the knot, doth not exceed one-4th of a line, or one-48th of an inch diameter.

Hence M. Reaumur advances, that the flexibility of glass increases in proportion to the fineness of the threads; and that, probably, had we but the art of drawing threads as fine as a spider's web, we might weave stuffs and cloths of them for wear. Accordingly, he made some experiments this way: and found he could make threads fine enough, viz, as fine, in his judgment, as spider's thread, but he could never make them long enough to do any thing with them.

*DUCTILITY of Spider's-webs.* See WEB.

**DUNGEON, DONJON**, in Fortification, the highest part of a castle built after the ancient mode; serving as a watch-tower, or place of observation; and also for the retreat of a garrison, in case of necessity, so that they may capitulate with greater advantage.

**DUPLE**, or *DOUBLE Ratio*, is that in which the antecedent term is double the consequent; or, where the exponent of the ratio is 2. Thus, 6 to 3 is in a duple ratio.

*Sub-DUPLE Ratio*, is that in which the consequent is double the antecedent; or, in which the exponent of the ratio is  $\frac{1}{2}$ . As in 3 to 6, which is in subduple ratio.

**DUPLICATE Ratio**, is the square of a ratio, or the ratio of the squares of two quantities. Thus, the duplicate ratio of  $a$  to  $b$ , is the ratio of  $a^2$  to  $b^2$ , or of the square of  $a$  to the square of  $b$ .—In a series of geometrical proportionals, the 1st term is to the 3d, in a duplicate ratio of the 1st to the 2d, or as the square of the first to the square of the 2d: Thus, in the geometricals 2, 4, 8, 16, &c, the ratio of 2 to 8, is the duplicate of that of 2 to 4, or as the square of 2 to the square of 4, that is, as 4 to 16. So that duplicate ratio is the ratio of the squares, as triplicate ratio is the ratio of the cubes, &c.

**DUPLICATION**, is the doubling of a quantity, or multiplying it by 2, or adding it to itself.

*DUPLICATION of a Cube*, is finding out the side of a cube that shall be double in solidity to a given cube: which is a celebrated problem, much cultivated by the ancient geometricians, about 2000 years ago.

It was first proposed by the oracle of Apollo at Delphos; which, being consulted about the manner of stopping a plague then raging at Athens, returned for answer, that the plague should cease when Apollo's altar, which was cubical, should be doubled. Upon this, they applied themselves in good earnest, to seek the



the duplicature of the cube, which from thence was called the *Delian problem*.

This problem cannot be effected geometrically, as it requires the solution of a cubic equation, or requires the finding of two mean proportionals, viz, between the side of the given cube and the double of the same, the first of which two mean proportionals is the side of the double cube, as was first observed by Hippocrates of Chios. For, let  $a$  be the side of the given cube, and  $z$  the side of the double cube sought; then it is  $z^3 = 2a^3$ ; or  $a^2 : z^2 :: z : 2a$ ; so that, if  $a$  and  $z$  be the first and 2d terms of a set of continued proportionals, then  $a^2 : z^2$  is the ratio of the square of the 1st to the square of the 2d, which, it is known, is the same as the ratio of the 1st term to the 3d, or of the 2d to the 4th, that is of  $z$  to  $2a$ ; therefore  $z$  being the 2d term,  $2a$  will be the 4th. So that  $z$ , the side of the cube sought, is the 2d of four terms in continued proportion, the 1st and 4th being  $a$  and  $2a$ , that is, the side of the double cube is the first of two mean proportionals between  $a$  and  $2a$ .

Eutocius, in his Commentaries on Archimedes, gives several ways of performing this by the mesolabe. In Pappus too are found three different ways; the first according to Archimedes, the second according to Hero, and the 3d by an instrument invented by Pappus, which gives all the proportions required. The fleur de Comiers has likewise published a demonstration of the same problem, by means of a compass with three legs. But all these methods are only mechanical. See Valerius Maximus, lib. 8; also Eutocius's Com. on lib. 2. Archimedes de Sphæra & Cylindro; and Pappus, lib. 3, prop. 5; & lib. 4, prop. 22.

DURER. (ALBERT), descended of an Hungarian family, but born at Nuremberg, in 1471, was one of the best engravers, painters, and practical geometricians of his age. He was at the same time a man of letters and a philosopher; and he was an intimate friend of Erasmus, who revised some of the pieces which he published. He was also a man of business, and for many years the leading magistrate of Nuremberg.

Though not the inventor, he was one of the first and greatest improvers of the art of engraving. He was however the inventor of cutting in wood, which he devised and practised in great perfection, using this way for expedition, as he had a multitude of designs to execute; and as his work was usually done in the most exquisite manner, his pieces took him up much time. For in many of those prints which he executed on copper, the engraving is elegant to a great degree. His Hell-Scene in particular, engraven in the year 1513, is as highly finished a print as ever was engraved, and as hap-

pily executed. In his wooden prints too it is surprising to see so much meaning in so early a master. In fact, Durer was a man of a very extensive genius. His pictures were excellent; as well as his prints, which were very numerous. They were much admired, from the first, and eagerly bought up; which put his wife, who was another Xantippe, upon urging him to spend more time upon engraving than he was inclined to do: for he was rich; and chose rather to practise his art as an amusement, than as a business. He died at Nuremberg, in 1528, at 57 years of age.

Albert Durer wrote several books, in the German language, which were translated into Latin by other persons, and published after his death. viz,

1. His book upon the rules of painting, intitled, *De Symmetria Partium in rectis formis Humanorum Corporum*, is one of them: printed in folio, at Nuremberg, in 1532, and at Paris in 1557. An Italian version also was published at Venice, in 1591.

2. *Institutiones Geometricæ*; Paris 1532.

3. *De Urbibus, Arcibus, Castellisque condendis & muniendis*; Paris 1531.

4. *De Varietate Figurarum, et Flexuris Partium, et Gestibus Imaginum*; Nuremberg 1534.

The figures in these books, which are from wooden plates, are very numerous, and most admirably well executed, indeed far beyond any thing of the kind done in our own days. Some of them also are of a very large size, as much as 16 inches in length, and of a proportional breadth, which being exquisitely worked, must have cost great labour. His geometry is chiefly of the practical kind, consisting of the most curious descriptions, inscriptions, and circumscriptions of geometrical lines, planes, and solids. We here meet, for the first time, with the plane figures, which folded up make the five regular or platonic bodies, as well as that curious construction of a pentagon, being the last method in prob. 23 of my Mensuration.

DYE, in Architecture, the trunk of the pedestal, or that part between the base and the cornice, being so called, because it is often made in the form of a dye or cube.

DYMANICS, is the science of moving powers; more particularly of the motion of bodies that mutually act on one another. See MECHANICS, MOTION, COMMUNICATION of Motion, OSCILLATION, PERCUSSION, &c.

DYPTERE, or DIPTERE, was a kind of temple, encompassed round with a double row of columns; and the pseudo-diptere, or false diptere, was the same, only this was encompassed with a single row of columns, instead of a double row.



## E.

## E A G

**E**AGLE, *Aquila*, is a constellation of the northern hemisphere, having its right wing contiguous to the equinoctial. For the stars in this constellation, see *AQUILA*.

**EARTH**, *Terra*, in Natural Philosophy, one of the four vulgar, or Peripatetical elements; defined a simple, dry, and cold substance; and, as such, an ingredient in the composition of all natural bodies.

*The EARTH*, in Geography, this terraqueous globe or ball, which we inhabit, consisting of land and sea.

*Figure of the EARTH.* The ancients had various opinions as to the figure of the earth: some, as Anaximander and Leucippus, held it cylindrical, or in form of a drum: but the principal opinion was, that it was flat; that the visible horizon was the bounds of the earth, and the ocean the bounds of the horizon; that the heavens and earth above this ocean were the whole visible universe; and that all beneath the ocean was Hades: and of this same opinion were also some of the Christian fathers, as Lactantius, St. Augustine, &c. See Lactan. lib. 3, cap. 24; St. Aug. lib. 16, de Civitate Dei; Aristotle de Cælo, lib. 2, cap. 13.

Such of the ancients however as understood any thing of astronomy, and especially the doctrine of eclipses, must have been acquainted with the round figure of the earth; as the ancient Babylonian astronomers, who had calculated eclipses long before the time of Alexander, and Thales the Grecian, who predicted an eclipse of the sun. It is now indeed agreed on all hands, unless perhaps by the most vulgar and ignorant, that the form of the terraqueous globe is globular, or very nearly so.

That the exterior of the earth is round, or rotund, is manifest to the most common perception, in the case of a ship sailing either from the land, or towards it; for when a person stands upon the shore, and sees a ship sail from the land, out to sea; at first he loses sight of the hull and lower parts of the ship, next the rigging and middle parts, and lastly of the tops of the masts themselves, in every case the rotundity of the sea between the ship and the eye being very visible: the contrary happens when a ship sails towards us; we first see the tops of the masts appear just over the rotundity of the sea; next we perceive the rigging, and lastly the hull of the ship itself: all which is well illustrated by the following figure.

## E A R



The round figure of the earth is also evident from the eclipses of the sun and moon; for in all eclipses of the moon, which are caused by the moon passing through the earth's shadow, that shadow always appears circular upon the face of the moon, what way soever it be projected, whether east, west, north, or south, and howsoever its diameter vary, according to the greater or less distance from the earth. Hence it follows, that the shadow of the earth, in all situations, is really conical; and consequently the body that projects it, i. e. the earth, is at least nearly spherical.

The spherical figure of the earth is also evinced from the rising and setting of the sun, moon, and stars; all which happen sooner to those who live to the east, and later to those living westwardly; and that more or less so, according to the distance and roundness of the earth.

So also, going or sailing to the northward, the north pole and northern stars become more elevated, and the south pole and southern stars more depressed; the elevation northerly increasing equally with the depression southerly; and either of them proportionably to the distance gone. The same thing happens in going to the southward. Besides, the oblique ascensions, descensions,



scensions, emersions, and amplitudes of the rising and setting of the sun and stars in every latitude, are agreeable to the supposition of the earth's being of a spherical form: all which could not happen if it was of any other figure.

Moreover, the roundness of the earth is farther confirmed by its having been often sailed round: the first time was in the year 1519, when Ferd. Magellan made the tour of the whole globe in 1124 days. In the year 1557 Francis Drake performed the same in 1056 days: in the year 1586, Sir Tho. Cavendish made the same voyage in 777 days; Simon Cordes, of Rotterdam, in the year 1590; in the year 1598, Oliver Noort, a Hollander, in 1077 days; Van Schouten, in the year 1615, in 749 days; Jac. Heremites and Joh. Huygens, in the year 1623, in 802 days: and many others have since performed the same navigation, particularly Anson, Bougainville, and Cook. Sometimes sailing round by the eastward, sometimes to the westward; till at length they arrived again in Europe, from whence they set out; and in the course of their voyage, observed that all the phenomena, both of the heavens and earth, correspond to, and evince this spherical figure.

The same globular figure is likewise inferred from the operation of Levelling, in which it is found necessary to make an allowance for the difference between the apparent and the true level.

The natural cause of this sphericity of the globe is, according to Sir Isaac Newton, the great principle of attraction, which the Creator has stamped on all the matter in the universe; and by which all bodies, and all the parts of bodies, mutually attract one another.—And the same is the cause of the sphericity of the drops of rain, quicksilver, &c.

What the earth loses of its sphericity by mountains and valleys, is nothing considerable; the highest eminence being scarce equivalent to the minutest protuberance on the surface of an orange. Its difference from a perfect sphere however is more considerable in another respect, by which it approaches nearly to the shape of an orange, or to an oblate spheroid, being a little flatted at the poles, and raised about the equatorial parts, so that the axis from pole to pole is less than the equatorial diameter. What gave the first occasion to the discovery of this figure of the earth, was the observations of some French and English philosophers in the East-Indies, and other parts, who found that pendulums, the nearer they came to the equator, performed their vibrations slower: from whence it follows, that the velocity of the descent of bodies by gravity, is less in countries nearer to the equator; and consequently that those parts are farther removed from the centre of the earth, or from the common centre of gravity. See the History of the Royal Academy of Sciences, by Du Hamel, p. 110, 156, 206; and l'Hist. de l'Acad. Roy. 1700 and 1701.—This circumstance put Huygens and Newton upon finding out the cause, which they attributed to the revolution of the earth about its axis. If the earth were in a fluid state, its rotation round its axis would necessarily make it put on such a figure, because the centrifugal force being greatest towards the equator, the fluid would there rise and swell most; and that its figure really should be so now, seems necessary, to keep the sea in the equinoctial re-

gions from overflowing the earth about those parts. See this curious subject well handled by Huygens, in his discourse De Causa Gravitatis, pa. 154, where he states the ratio of the polar diameter to that of the equator, as 577 to 578. And Newton, in his Principia, first published in 1686, demonstrates, from the theory of gravity, that the figure of the earth must be that of an oblate spheroid generated by the rotation of an ellipse about its shortest diameter, provided all the parts of the earth were of an uniform density throughout, and that the proportion of the polar to the equatorial diameter of the earth, would be that of 689 to 692, or nearly that of 229 to 230, or as .9956522 to 1.

This proportion of the two diameters was calculated by Newton in the following manner. Having found that the centrifugal force at the equator is  $\frac{1}{289}$ th of gravity, he assumes, as an hypothesis, that the axis of the earth is to the diameter of the equator as 100 to 101, and thence determines what must be the centrifugal force at the equator to give the earth such a form, and finds it to be  $\frac{4}{585}$ ths of gravity: then, by the rule of proportion, if a centrifugal force equal to  $\frac{4}{585}$ th of gravity would make the earth higher at the equator than at the poles by  $\frac{1}{289}$ th of the whole height at the poles, a centrifugal force that is the  $\frac{1}{289}$ th of gravity will make it higher by a proportional excess, which by calculation is  $\frac{1}{289}$ th of the height at the poles; and thus he discovered that the diameter at the equator is to the diameter at the poles, or the axis, as 230 to 229. But this computation supposes the earth to be every where of an uniform density; whereas if the earth is more dense near the centre, then bodies at the poles will be more attracted by this additional matter being nearer; and therefore the excess of the semi-diameter of the equator above the semi-axis, will be different. According to this proportion between the two diameters, Newton farther computes, from the different measures of a degree, that the equatorial diameter will exceed the polar, by 34 miles and  $\frac{1}{2}$ .

Nevertheless, Messrs Cassini, both father and son, the one in 1701, and the other in 1713, attempted to prove, in the Memoirs of the Royal Academy of Sciences, that the earth was an oblong spheroid; and in 1718, M. Cassini again undertook, from observations, to shew that, on the contrary, the longest diameter passes through the poles; which gave occasion for Mr. John Bernoulli, in his Essai d'une Nouvelle Physique Celeste, printed at Paris in 1735, to triumph over the British philosopher, apprehending that these observations would invalidate what Newton had demonstrated. And in 1720, M. De Mairan advanced arguments, supposed to be strengthened by geometrical demonstrations, farther to confirm the assertions of Cassini. But in 1735 two companies of mathematicians were employed, one for a northern, and another for a southern expedition, the result of whose observations and measurement plainly proved that the earth was flatted at the poles. See DEGREE.

The proportion of the equatorial diameter to the polar, as stated by the gentlemen employed on the northern expedition for measuring a degree of the meridian, is as 1 to 0.9891; by the Spanish mathematicians as 266 to 265, or as 1 to 0.99624; by M. Bouguer as 179 to 178, or as 1 to 0.99441.

As to all conclusions however deduced from the



length of pendulums in different places, it is to be observed that they proceed upon the supposition of the uniform density of the earth, which is a very improbable circumstance; as justly observed by Dr. Horsley in his letter to Capt. Phipps. "You finish your article, he concludes, relating to the pendulum with saying, 'that these observations give a figure of the earth nearer to Sir Isaac Newton's computation, than any others that have hitherto been made;' and then you state the several figures given, as you imagine, by former observations, and by your own. Now it is very true, that *if* the meridians be ellipses, or *if* the figure of the earth be that of a spheroid generated by the revolution of an ellipsis, turning on its shorter axis, the particular figure, or the ellipticity of the generating ellipsis, which your observations give, is nearer to what Sir Isaac Newton saith it should be, if the globe were homogeneous, than any that can be derived from former observations. But yet it is not what you imagine. Taking the gain of the pendulum in latitude  $79^{\circ} 50'$  exactly as you state it, the difference between the equatorial and the polar diameter, is about as much less than the Newtonian computation makes it, and the hypothesis of homogeneity would require, as you reckon it to be greater. The proportion of 212 to 211 should indeed, according to your observations, be the proportion of the force that acts upon the pendulum at the poles, to the force acting upon it at the equator. But this is by no means the same with the proportion of the equatorial diameter to the polar. If the globe were homogeneous, the equatorial diameter would exceed the polar by  $\frac{1}{230}$  of the length of the latter: and the polar force would also exceed the equatorial by the like part. But if the difference between the polar and equatorial force be greater than  $\frac{1}{230}$ , (which may be the case in an heterogeneous globe, and seems to be the case in ours,) then the difference of the diameters should, according to theory, be less than  $\frac{1}{230}$ , and vice versa.

"I confess this is by no means obvious at first sight; so far otherwise, that the mistake, which you have fallen into, was once very general. Many of the best mathematicians were misled by too implicit a reliance upon the authority of Newton, who had certainly confined his investigations to the homogeneous spheroid, and had thought about the heterogeneous only in a loose and general way. The late Mr. Clairault was the first who set the matter right, in his elegant and subtle treatise on the figure of the earth. That work hath now been many years in the hands of mathematicians, among whom I imagine there are none, who have considered the subject attentively, that do not acquiesce in the author's conclusions.

"In the 2d part of that treatise, it is proved, that putting  $P$  for the polar force,  $\Pi$  for the equatorial,  $\delta$  for the true ellipticity of the earth's figure, and  $\epsilon$  for the ellipticity of the homogeneous spheroid,

$$\frac{P - \Pi}{\Pi} = 2\epsilon - \delta : \text{therefore } \delta = 2\epsilon - \frac{P - \Pi}{\Pi};$$

and therefore, according to your observation,  $\delta = \frac{1}{257}$ . This is the just conclusion from your observations of the pendulum, taking it for granted, that the meridians are ellipses: which is an hypothesis, upon which all the reasonings of theory have hitherto proceeded. But plausible as it may seem, I must say, that there is much

reason from experiment to call it in question. If it were true, the increment of the force which actuates the pendulum, as we approach the poles, should be as the square of the sine of the latitude: or, which is the same thing, the decrement, as we approach the equator, should be as the square of the cosine of the latitude. But whoever takes the pains to compare together such of the observations of the pendulum in different latitudes, as seem to have been made with the greatest care, will find that the increments and decrements do by no means follow these proportions; and in those which I have examined, I find a regularity in the deviation which little resembles the mere error of observation. The unavoidable conclusion is, that the true figure of the meridians is not elliptical. If the meridians are not ellipses, the difference of the diameters may indeed, or it may not, be proportional to the difference between the polar and the equatorial force; but it is quite an uncertainty, what relation subsists between the one quantity and the other; our whole theory, except so far as it relates to the homogeneous spheroid, is built upon false assumptions, and there is no saying what figure of the earth any observations of the pendulum give."


He then lays down the following table, which shews the different results of observations made in different latitudes; in which the first three columns contain the names of the several observers, the places of observation, and the latitude of each; the 4th column shews the quantity of  $P - \Pi$  in such parts as  $\Pi$  is 100000, as deduced from comparing the length of the pendulum at each place of observation, with the length of the equatorial pendulum as determined by M. Bouguer, upon the supposition that the increments and decrements of force, as the latitude is increased or lowered, observe the proportion which theory assigns. Only the 2d and the last value of  $P - \Pi$  are concluded from comparisons with the pendulum at Greenwich and at London, not at the equator. The 5th column shews the value of  $\delta$  corresponding to every value of  $P - \Pi$ , according to Clairault's theorem:

Observers.	Places.	Latitudes.	$P - \Pi$	$\delta$
Bouguer	Equator	$0^{\circ} \quad 0'$		
Bouguer	Porto Bello.	9 34	741.8	$\frac{1}{84}$
Green	Otaheitee	17 29	563.2	$\frac{1}{27}$
Bouguer	San Domingo	18 27	591.0	$\frac{1}{38}$
Abbé de la } Caille }	Cape of } Good Hope }	33 55	731.5	
- - - -	Paris	48 50	585.1	$\frac{1}{57}$
The Acade- } micians }	Pello	66 48	565.9	$\frac{1}{29}$
Capt. Phipps	- - - -	79 50	471.2	$\frac{1}{71}$

"By this table it appears, that the observations in the middle parts of the globe, setting aside the single one at the Cape, are as consistent as could reasonably be expected; and they represent the ellipticity of the earth as about  $\frac{1}{38}$ . But when we come within 10 degrees of the equator, it should seem that the force of gravity suddenly becomes much less, and within the like distance



The following problem, communicated by Dr Leatherland to Dr. Pemberton, and published by Mr. Robertson, serves for finding the proportion between the axis and the equatorial diameter, from measures taken of a degree of the meridian in two different latitudes, supposing the earth an oblate spheroid.



in each; and let  $P$  denote  $\sqrt{\frac{M}{N}}$ : then

$$\sqrt{\frac{\text{cof.}^2 F - P^2 \times \text{cof.}^2 E}{P^2 \times \text{fin.}^2 E - \text{fin.}^2 F}} \text{ is } = \frac{\text{lesser axis}}{\text{greater axis}}.$$

But M, or the length of a degree, obtained by actual mensuration in different latitudes, is known from the following table :

<i>Names.</i>	<i>Latit.</i> ° /	<i>Value of M.</i> toises
Maupertuis and Assoc.	66 20	M = 57438
Cassini and {	49 22	M = 57074
La Caille {	45 00	M = 57050
Boscovich	43 00	M = 56972
De la Caille	33 18	M = 57037
Juan and Ulloa	00 00	M = 56768
Bouguer	00 00	M = 56753
Condamine	00 00	M = 56749

} at the equator.

Now, the magnitude as well as the figure of the earth, that is the polar and equatorial diameters, may be deduced from the foregoing problem. For, as half the latus rectum of the greater axis  $Aa$  is the radius of curvature at  $A$ , it is given in magnitude from the degree measured there, and thence the axes themselves are given. Thus, the circular arc whose length is equal to the radius being  $57.29578$  degrees, if this number be multiplied by  $56750$  toises, the measure of a degree at the



the equator, as Bouguer has stated it, the product will be the radius of curvature there, or half the latus rectum of the greater axis; and this is to half the lesser axis in the ratio of the less axis to the greater, that is, as 0.9953467 to 1; whence the two axes are 6533820 and 6564366 toises, or 7913 and 7950 English miles; and the difference between the two axes about 37 miles. See Robertson's Navigation, vol. 2, pa. 206 &c. See also Suite des Mem. de l'Acad. 1718, pa. 247, and Maclaurin's Fluxions, vol. 2, book 1, ch. 14.

And very nearly the same ratio is deduced from the lengths of pendulums vibrating in the same time, in different latitudes; provided it be again allowed that the meridians are real ellipses, or the earth a true spheroid, which however can only take place in the case of an uniform gravity in all parts of the earth.

Thus, in the new Petersburg Acts, for the years 1788 and 1789, are accounts and calculations of experiments relative to this subject, by M. Krafft. These experiments were made at different times and in various parts of the Russian empire. This gentleman has collected and compared them, and drawn the proper conclusions from them: thus he infers that the length  $x$  of a pendulum that swings seconds in any given latitude  $\lambda$ , and in a temperature of 10 degrees of Reaumur's thermometer, may be determined by this equation:

$x = 439.178 + 2.321 \sin^2 \lambda$ , lines of a French foot, or  $x = 39.0045 + 0.206 \sin^2 \lambda$ , in English inches, in the temperature of 53 of Fahrenheit's thermometer.

This expression nearly agrees, not only with all the experiments made on the pendulum in Russia, but also with those of Mr. Graham in England, and those of Mr. Lyons in  $79^\circ 50'$  north latitude, where he found its length to be 431.38 lines. It also shews the augmentation of gravity from the equator to the parallel of a given latitude  $\lambda$ : for, putting  $g$  for the gravity under the equator,  $G$  for that under the pole, and  $y$  for that under the latitude  $\lambda$ , M. Krafft finds

$y = (1 + 0.0052848 \sin^2 \lambda) g$ ; and theref.  $G = 1.0052848 g$ .

From this proportion of gravity under different latitudes, the same author infers that, in case the earth is a homogeneous ellipsoid, its oblateness must be  $\frac{1}{231}$ ; instead of  $\frac{1}{230}$ , which ought to be the result of this hypothesis: but on the supposition that the earth is a heterogeneous ellipsoid, he finds its oblateness, as deduced from these experiments, to be  $\frac{1}{237}$ ; which agrees with that resulting from the measurement of some of the degrees of the meridian. This confirms an observation of M. De la Place, that, if the hypothesis of the earth's homogeneity be given up, then theory, the measurement of degrees of latitude, and experiments with the pendulum, all agree in their result with respect to the oblateness of the earth. See Memoires de l'Acad. 1783, pa. 17.

In the Philos. Transf. for 1791, pa. 236, Mr. Dalby has given some calculations on measured degrees of the meridian, from whence he infers, that those degrees measured in middle latitudes, will answer nearly to an ellipsoid whose axes are in the ratio assigned by Newton, viz, that of 230 to 229. And as to the deviations of some of the others, viz, towards the poles

and equator, he thinks they are caused by the errors in the observed celestial arcs.

Tacquet draws some pretty little inferences, in the form of paradoxes, from the round figure of the earth; as, 1st, That if any part of the surface of the earth were quite plane, a man could no more walk upright upon it, than on the side of a mountain. 2d, That the traveller's head goes a greater space than his feet; and a horseman than a footman; as moving in a greater circle. 3d, That a vessel, full of water, being raised perpendicularly, some of the water will be continually flowing out, yet the vessel still remain full; and on the contrary, if a vessel of water be let perpendicularly down, though nothing flow out, yet it will cease to be full: consequently there is more water contained in the same vessel at the foot of a mountain, than on the top; because the surface of the water is compressed into a segment of a smaller sphere below than above. Tacq. Astron. lib. 1, cap. 2.

*Changes of the EARTH.* Mr. Boyle suspects that there are great, though slow internal changes, in the mass of the earth. He argues from the varieties observed in the change of the magnetic needle, and from the observed changes in the temperature of climates. But as to the latter, there is reason to doubt that he could not have diaries of the weather sufficient to direct his judgment. Boyle's Works abr. vol. 1, pa. 292, &c.

*Magnetism of the EARTH.* The notion of the magnetism of the earth was started by Gilbert; and Boyle supposes magnetic effluvia moving from one pole to the other. See his Works abr. vol. 1, pa. 285, 290.

Dr. Knight also thinks that the earth may be considered as a great loadstone, whose magnetical parts are disposed in a very irregular manner; and that the south pole of the earth is analogous to the north pole in magnets, that is, the pole by which the magnetical stream enters. See MAGNET.

He observes, that all the phenomena attending the direction of the needle, in different parts of the earth, in great measure correspond with what happens to a needle, when placed upon a large terrella; if we make allowances for the different dispositions of the magnetical parts, with respect to each other, and consider the south pole of the earth as a north pole with regard to magnetism. The earth might become magnetical by the iron ores it contains, for all iron ores are capable of magnetism. It is true, the globe might notwithstanding have remained unmagnetical, unless some cause had existed capable of making that repellent matter producing magnetism move in a stream through the earth.

Now the doctor thinks that such a cause does exist. For if the earth revolves round the sun in an ellipsis, and the south pole of the earth is directed towards the sun, at the time of its descent towards it, a stream of repellent matter will thence be made to enter at the south pole, and issue out at the north. And he suggests, that the earth's being in its perihelion in winter may be one reason why magnetism is stronger in this season than in summer.

This cause here assigned for the earth's magnetism must continue, and perhaps improve it, from year to year. Hence the doctor thinks it probable, that the earth's magnetism has been improving ever since the creation, and that this may be one reason why the use



of the compass was not discovered sooner. See Dr. Knight's Attempt to demonstrate, that all the phenomena in nature may be explained by Attraction and Repulsion, prop. 87.

*Magnitude and Constitution of the EARTH.* This has been variously determined by different authors, both ancient and modern. The usual way has been, to measure the length of one degree of the meridian, and multiply it by 360, for the whole circumference. See DEGREE. Diogenes Laertius informs us, that Anaximander, a scholar of Thales, who lived about 550 years before the birth of Christ, was the first who gave an account of the circumference of the sea and land; and it seems his measure was used by the succeeding mathematicians, till the time of Eratosthenes. Aristotle, at the end of lib. 2 De Cælo, says, the mathematicians who have attempted to measure the circuit of the earth, make it 40000 stadia: which it is thought is the number determined by Anaximander.

Eratosthenes, who lived about 200 years before Christ, was the next who undertook this business; which, as Cleomedes relates, he performed by taking the sun's zenith distances, and measuring the distance between two places under the same meridian; by which he deduced for the whole circuit about 250000 stadia, which Pliny states at 31500 Roman miles, reckoning each at 1000 paces. But this measure was accounted false by many of the ancient mathematicians, and particularly by Hipparchus, who lived 100 years afterwards, and who added 25000 stadia to the circuit of Eratosthenes.

Possidonius, in the time of Cicero and Pompey the Great, next measured the earth, viz, by means of the altitudes of a star, and measuring a part of a meridian; and he concluded the circumference at 240000 stadia, according to Cleomedes, but only at 180000 according to Strabo.

Ptolomy, in his Geography, says that Marinus, a celebrated geographer, attempted something of the same kind; and, in lib. 1, cap. 3, he mentions that he himself had tried to perform the business in a way different from any other before him, which was by means of places under different meridians: but he does not say how much he made the number; for he still made use of the 180000, which had been found out before him.

Snellius relates, from the Arabian Geographer Abulfeda, who lived about the 1300th year of Christ, that about the 800th year of Christ, Almainon, an Arabian king, having collected together some skilful mathematicians, commanded them to find out the circumference of the earth. Accordingly these made choice of the fields of Mesopotamia, where they measured under the same meridian from north to south, till the pole was depressed one degree lower: which measure they found equal to 56 miles, or  $56\frac{1}{2}$ : so that according to them the circuit of the earth is 20160 or 20340 miles.

It was a long time after this before any more attempts were made in this business. At length however, the same Snell, above mentioned, professor of mathematics at Leyden, about the year 1620, with great skill and labour, by measuring large distances between two parallels, found one degree equal to 28500 perches, each of which is 12 Rhinland feet, amounting to 19

Dutch miles, and so the whole periphery 6840 miles; a mile being, according to him, 1500 perches, or 18000 Rhinland feet. See his treatise called *Eratosthenes Batarus*.

The next that undertook this measurement, was Richard Norwood, who in the year 1635, by measuring the distance from London to York with a chain, and taking the sun's meridian altitude, June 11th old style, with a sextant of about 5 feet radius, found a degree contained 367200 feet, or 69 miles and a half and 14 poles; and thence the circumference of a great circle of the earth is a little more than 25036 miles, and the diameter a little more than 7966 miles. See the particulars of this measurement in his *Seaman's Practice*.

The measurement of the earth by Snell, though very ingenious and troublesome, and much more accurate than any of the ancients, being still thought by some French mathematicians, as liable to certain small errors, the business was renewed, after Snell's manner, by Picard and other mathematicians, by the king's command; using a quadrant of  $3\frac{1}{2}$  French feet radius; by which they found a degree contained 342360 French feet. See Picard's treatise, *La Mesure de la Terre*.

M. Cassini the younger, in the year 1700, by the king's command also, renewed the business with a quadrant of 10 feet radius, for taking the latitude, and another of  $3\frac{1}{8}$  feet for taking the angles of the triangles; and found a degree, from his calculation, contained 57292 toises, or almost  $69\frac{1}{2}$  English miles.

See the results of many other measurements under the article DEGREE. From the mean of all which, the following dimensions may be taken as near the truth:

the circumference	25000 miles,
the diameter	7957 $\frac{3}{4}$ miles,
the superficies	198944206 square miles,
the solidity	263930000000 cubic miles.

Also the seas and unknown parts of the earth, by a measurement of the best maps, contain 160522026 square miles, the inhabited parts 38922180; of which Europe contains 4456065; Asia, 10768823; Africa, 9654807; and America, 14110874.

It is now generally granted that the terraqueous globe has two motions, besides that on which the precession of the equinoxes depends; the one diurnal around its own axis in the space of 24 hours, which constitutes the natural day or nycthemeron; the other annual, about the sun, in an elliptical orbit or track, in 365 days 6 hours, constituting the year. From the former arise the diversities of night and day; and from the latter, the vicissitudes of seasons, spring, summer, autumn, winter.

The terraqueous globe is distinguished into three parts or regions, viz, 1st, The external part or crust, being that from which vegetables spring and animals are nursed. 2d, The middle, or intermediate part, which is possessed by fossils, extending farther than human labour ever yet penetrated. 3d, The internal or central part, which is utterly unknown to us, though by many authors supposed of a magnetic nature; by others, a mass or sphere of fire; by others, an abyss or collection of waters, surrounded by the strata of earth; and by others, a hollow, empty space, inhabited by animals, who have their sun, moon, planets, and other



other conveniences within the same. But others divide the body of the globe into two parts, viz, the external part, called the cortex, including the internal, which they call the nucleus, being of a different nature from the former, and possessed by fire, water, or more probably by a considerable portion of metals, as it has been found, by calculation, that the mean density of the whole earth is near double the density of common stone. See my determination of it, *Philos. Trans.* 1778, pa. 781.

The external part of the globe either exhibits inequalities, as mountains and valleys; or it is plane and level; or dug in channels, fissures, beds, &c, for rivers, lakes, seas, &c. These inequalities in the face of the earth most naturalists suppose have arisen from a rupture or subversion of the earth, by the force either of the subterraneous fires or waters. The earth, in its natural and original state, it has been supposed by Des Cartes, and after him, Burnet, Steno, Woodward, Whiston, and others, was perfectly round, smooth, and equable; and they account for its present rude and irregular form, principally from the great deluge. See *DELUGE*.

In the external, or cortical part of the earth, there appear various strata, supposed the sediments of several floods; the waters of which, being replete with matters of divers kinds, as they dried up, or oozed through, deposited these different matters, which in time hardened into strata of stone, sand, coal, clay, &c.

Dr. Woodward has considered the circumstances of these strata with great attention, viz, their order, number, situation with respect to the horizon, depth, intersections, fissures, colour, consistence, &c. He ascribes the origin and formation of them all, to the great flood or cataclysmus. At that terrible revolution he supposed that all sorts of terrestrial bodies had been dissolved and mixed with the waters, forming all together a chaos or confused mass. This mass of terrestrial particles, intermixed with water, he supposes was at length precipitated to the bottom; and that generally according to the order of gravity, the heaviest sinking first, and the lightest afterwards. By such means were the strata formed of which the earth consists; which, attaining their solidity and hardness by degrees, have continued so ever since. These sediments, he farther concludes, were at first all parallel and concentric; and the surface of the earth formed of them, perfectly smooth and regular; but that in course of time, divers changes happening, from earthquakes, volcanos, &c, the order and regularity of the strata was disturbed and broken, and the surface of the earth by such means brought to the irregular form in which it now appears.

M. De Buffon surmises that the earth, as well as the other planets, are parts struck off from the body of the sun by the collision of comets; and that when the earth assumed its form, it was in a state of liquefaction by fire. But that could not be the method of producing the planets; for if they were struck off from the body of the sun, they would move in orbits that would always pass through the sun, instead of having the sun for their focus, or centre, as they are now

found; so that having been struck off they would fall down into the sun again, terminating their career as it were after one revolution only.

*EARTH, in Astronomy*, is one of the primary planets, according to the system of Copernicus, or Pythagoras; its astronomical character or mark being  $\ominus$ : but according to the Ptolomaic hypothesis, the earth is the centre of the system. For, whether the earth move or remain at rest, that is, whether it be fixed in the centre, having the sun, the heavens and stars moving round it from east to west; or, these being at rest, whether the earth only moves from west to east, is the great article that distinguishes the Ptolomaic system from the Copernican.

*Motion of the EARTH.* It is now universally agreed that, besides the small motion of the earth which causes the precession of the equinoxes, the earth has two great and independent motions; viz, the one by which it turns round its own axis, in the space of 24 hours nearly, and causing the continual succession of day and night; and the other an absolute motion of its whole mass in a large orbit about the sun, having that luminary for its centre, in such manner that its axis keeps always parallel to itself, inclined in the same angle to its path, and by that means causing the vicissitudes of seasons, spring, summer, autumn, winter.

It is indeed true that, as to sense, the earth appears to be fixed in the centre, with the sun, stars and heavens moving round it every day; and such doubtless would be considered as the true nature of the motions in the rude ages of mankind, as they are still by the rude and unlearned. But to a thinking and learned mind, the contrary will soon appear.

Indeed there are traces of the knowledge of these motions in the earliest age of the sciences. Cicero, in his *Tusc. Quæst.* says that Nicetas of Syracuse first discovered that the earth had a diurnal motion, by which it revolved round its axis every 24 hours; and Plutarch, de *Placit. Philosoph.* informs, that Philolaus discovered its annual motion round the sun; and Aristarchus, about 100 years after Philolaus, proposed the motion of the earth in stronger and clearer terms, as we are assured by Archimedes, in his *Arenarius*. And the same, we are farther assured, was the opinion and doctrine of Pythagoras.

But the religious opinions of the heathen world prevented this doctrine from being more cultivated. For, Aristarchus being accused of sacrilege by Cleanthes for moving Vesta and the tutelar deities of the universe out of their places, the philosophers were obliged to dissemble, and seem to relinquish so perilous a position.

Many ages afterwards, Nic. Cusanus revived the ancient system, in his *Doct. de Pignorant.* and asserted the motion of the earth: but the doctrine gained very little ground till the time of Copernicus, who shewed its great use and advantages in astronomy; and who had immediately all the philosophers and astronomers on his side, who dared to differ from the crowd, and were not afraid of ecclesiastical censure, which was not less dangerous under the christian dispensation, than it had been under that of the heathen. For, because certain parts of scripture make mention of the stability of



of the earth, and of the motion of the sun, as the rising and setting, &c, the fathers of the church thought their religion required that they should defend, with all its power, what they conceived to be its doctrines, and to censure and punish every attempt at innovation on such points. They have now however been pretty generally convinced that in such instances the expressions are only to be considered as accommodated to appearances, and the vulgar notions of things.

By the diurnal rotation of the earth on its axis, the same phenomena will take place as if it had no such motion, and as if the sun and stars moved round it. For, turning round from west to east, causes the sun and all the visible heavens to seem to move the contrary way, or from east to west, as we daily see them do. So, when in its rotation it has brought the sun or a star to appear just in the horizon in the east, they are then said to be rising; and as the earth continues to revolve more and more towards the east, other stars seem to rise and advance westwards, passing the meridian of the observer, when they are due south from him, and at their greatest altitude above his horizon; after which, by a continuance of the same motions, viz, of the earth's rotation eastwards, and the luminaries apparent counter motion westwards, these decline from the meridian, or south point, towards the west, where being arrived, they are said to set and descend below it; and so on continually from day to day; thus making it day while the sun is above the horizon, and night while he is below it.

While the earth is thus turning on its axis, it is at the same time carried by its proper motion in its orbit round the sun, as one of the planets, namely, between the orbits of Venus and Mars, having the orbits of Venus and Mercury within its own, or between it and the sun, in the centre, and those of Mars, Jupiter, Saturn, &c, without or above it; which are therefore called superior planets, and the others the inferior ones. This is called the annual motion of the earth, because it is performed in a year, or 365 days 6 hours nearly; or rather 365 days 5<sup>h</sup> 49<sup>m</sup>; from any equinox or solstice to the same again, making the tropical year; but from any fixed star to the same again, as seen from the sun, in 365 days 6 hours 9 minutes, which is called the sidereal year. The figure of this orbit is elliptical, having the sun in one focus, the mean distance being about 95 millions of miles, which is upon the supposition that the sun's parallax is about  $8''\frac{1}{3}$ , or the angle under which the earth's semi-diameter would appear to an observer placed in the sun: and the eccentricity of the orbit, or distance of the sun, in the focus, from the centre of this elliptic orbit, is about  $\frac{1}{80}$ th of the mean distance.

Now this annual motion is performed in such a manner, that the earth's axis is every where parallel, or in the same direction in every part of the orbit; by which means it happens, that at one time of the year the sun enlightens more of the north polar parts, and at the opposite season of the year more of the southern parts, thus shewing all the varieties of seasons, spring, summer, autumn and winter; which may be illustrated in the following manner: Let the candle I (fig. 1, plate viii) represent the sun, about which the

earth E, or F, &c, is moved in its elliptical orbit ABCD, or ecliptic, and cutting the equator *abcd* in the nodes E and G: then, suspending the terrella by its north pole, and moving it, so suspended, round the ecliptic, its axis will always be parallel to its first position, and the various seasons will be represented at the different parts of the path. Thus, when the earth is at  $\odot$  or F, the enlightened half of it includes the south pole, and leaves the north pole in darkness, making our winter; at G it is spring, and the two poles are equally illuminated, and the days are every where of the same length; at H or  $\varpi$  it is our summer, the north polar parts being in the illuminated hemisphere, and the southern in the dark one; lastly at E it is autumn, the poles being equally illuminated again, and the days of equal length every where.

EARTH, its *Quantity of Matter, Density, and Attractive Power*. Although the relative densities of the earth and most of the other planets have been known a considerable time, it is but very lately that we have come to the knowledge of the absolute gravity or density of the whole mass of the earth. This I have calculated and deduced from the observations made by Dr. Maskelyne, Astronomer Royal, at the mountain Schiehallien, in the years 1774, 5, and 6. The attraction of that mountain on a plummet, being observed on both sides of it, and its mass being computed from a number of sections in all directions, and consisting of stone; these data being then compared with the known attraction and magnitude of the earth, gave by proportion its mean density, which is to that of water as 9 to 2, and to common stone as 9 to 5: from which very considerable mean density, it may be presumed that the internal parts contain some great quantities of metals.

From the density, now found, its quantity of matter becomes known, being equal to the product of its density by its magnitude. From various experiments too, we know that its attractive force, at the surface, is such, that bodies fall there through a space of  $16\frac{1}{2}$  feet in the first second of time: from whence the force at any other place, either within or without it, becomes known; for the force at any part within it, is directly as its distance from the centre; but the force of any part without it, reciprocally as the square of its distance from the centre.

EAST, one of the cardinal points of the horizon, or of the compass, being the middle point of it between north and south, on that side where the sun rises, or the point in which it is intersected on that side by the prime vertical.

EASTER, a feast of the church, held in memory of our Saviour's resurrection. This feast has been annually celebrated ever since the time of the apostles, and is one of the most considerable festivals in the christian calendar; being that which regulates and determines the times of all the other moveable feasts.

The rule for the celebration of Easter, fixed by the council of Nice, in the year 325, is, that it be held on the Sunday which falls upon, or next after, the full moon which happens next after the 21st of March; that is, the Sunday which falls upon, or next after the first full moon after the vernal equinox. The reason of which



which decree was, that the christians might avoid celebrating their Easter at the same time with the Jewish Passover, which, according to the institution of Moses, was held the very day of the full moon.

*To find EASTER according to the Old, or Julian Style.*

In the annexed table, find the golden number, with the day of the paschal full moon, and the Sunday letter annexed; compare this letter with the dominical letter of the given year, that it may appear how many days are to be added to the day of the paschal full moon, to give Easter-day.

*For ex.* In the year 1715, the dominical letter is B, and the golden number is 6, opposite to which stands April 10 for the day of the paschal full moon; opposite to which is the Sunday letter B, which happening to be the same with that of the year given, that day is a Sunday; and therefore Easter will fall 7 days after, viz, on the 17th of April.

But in this computation, the vernal equinox is supposed fixed to the 21st of March; and the cycle of 19 years, or golden numbers, is supposed to point out the places of the new and full moons exactly; both which suppositions are

erroneous: so that the Julian Easter never happens at its due time, unless by accident. For instance, in the above example the vernal equinox falls on the 10th of March, eleven days before the rule supposes it; and the paschal full moon on the 7th of April, or 3 days earlier than was supposed: and therefore Easter-day should be held on the 10th of April, instead of the 17th.

This error had grown to such a height, that pope Gregory the 13th thought it necessary to correct it; and accordingly, in the year 1582, by the advice of Aloysius Lilius and others, he ordered 10 days to be thrown out of October, to bring the vernal equinox back again to the 21st of March: and hence arise the terms Gregorian calendar, Gregorian year, &c.

This correction however did not entirely remove the error; for the equinoxes and solstices still anticipate 28' 20" in every 100 Gregorian years; but the difference is so inconsiderable as not to amount to a whole day, or 24 hours, in less than 5082 Gregorian years.

The Gregorian, or New Style, was not introduced into England till the year 1752, when eleven days were thrown out, viz, between the 3d and 14th of September, the error amounting then to that quantity.

Gold. Num.	Paschal full moons.	Dom. letter.
16	March 21	C
5	22	D
	23	E
13	24	F
2	25	G
	26	A
10	27	B
	28	C
18	29	D
7	30	E
	31	F
15	April 1	G
4	2	A
	3	B
12	4	C
1	5	D
	6	E
9	7	F
	8	G
17	9	A
6	10	B
	11	C
14	12	D
3	13	E
	14	F
11	15	G
	16	A
19	17	B
8	18	C

*To find EASTER according to the New or Gregorian Style, till the year 1900 exclusive.*

Look for the golden number of the year in the first column of the table, against which stands the day of the paschal full moon; then look in the 3d column for the Sunday letter, next after the day of the full moon, and the day of the month standing against that Sunday letter is Easter-day. When the full moon happens on a Sunday, then the next Sunday after is Easter-day.

*For Ex.* For the year 1790, the golden number is 5; against which stands March the 30th, and the next Sunday letter, which is C, below that, stands opposite April 4, which is therefore the Easter-day for the year 1790.

Though the Gregorian calendar be much preferable to the Julian, it is yet not without its defects. It cannot, for instance, keep the equinox fixed on the 21st of March, but it will sometimes fall on the 19th, and sometimes on the 23d.

Add, that the full moon happening on the 20th of March, might sometimes be paschal; yet it is not allowed as such in the Gregorian computation; as on the contrary, the full moon of the 22d of March may be allowed for paschal, which it is not. Scaliger and Calvisius have also pointed out other inaccuracies in this calendar. An excellent paper on this subject by the earl of Macclesfield, may be seen in the Philos. Transf. vol. 40, pa. 417.

EASTER Term. See TERM.

EAVES, the margin or lower edge of the roof of a house, being the lowest course of tiles, slates, or the like, which hang over the walls to throw off the water to a distance from the wall.

EAVES-Board, or EAVES-Lath, a thick feather-edged board, usually nailed round the eaves of a house for the lowermost tiles, slate, or shingles, to rest upon.

EBBING and FLOWING of the Sea. See TIDES.

ECCENTRIC. See EXCENTRIC.

ECCENTRICITY. See EXCENTRICITY.

ECHO, or Eccho, a sound reflected, or reverberated from some body, and thence returned or repeated to the ear.

For an echo to be heard, the ear must be in the line of reflection; that the person who made the sound, may hear the echo, it is necessary he should be in a perpendicular to the place which reflects it; and for a multiple

Gold. Num.	Paschal full moon.	Sund. letter.
14	March 21	C
3	22	D
	23	E
11	24	F
	25	G
19	26	A
8	27	B
	28	C
16	29	D
5	30	E
	31	F
13	April 1	G
2	2	A
	3	B
10	4	C
	5	D
18	6	E
7	7	F
	8	G
15	9	A
4	10	B
	11	C
12	12	D
1	13	E
	14	F
9	15	G
	16	A
17	17	B
6	18	C



multiple or tautological echo, it is necessary there be a number of walls and vaults, rocks, and cavities, either placed behind each other, or fronting each other. Those murmurs in the air, that are occasioned by the discharge of great guns, &c, are a kind of indefinite echoes, and are produced from the vaporous particles suspended in the atmosphere, which resist the undulations of sound, and reverberate them to the ear.

There can be no echo, unless the direct and reflex sounds follow one another at a sufficient distance of time; for if the reflex sound arrive at the ear before the impression of the direct sound ceases, the sound will not be doubled, but only rendered more intense. Now if we allow that 9 or 10 syllables can be pronounced in a second, in order to preserve the sounds articulate and distinct, there should be about the 9th part of a second between the times of their appulse to the ear; or, as sound flies about 1142 feet in a second, the said difference should be  $\frac{1}{9}$  of 1142, or 127 feet; and therefore every syllable will be reflected to the ear at the distance of about 70 feet from the reflecting body; but as, in the ordinary way of speaking, 3 or 4 syllables only are uttered in a second, the speaker, that he may have the echo returned as soon as they are expressed, should stand about 500 feet from the reflecting body; and so in proportion for any other number of syllables. Merfenne allows for a monosyllable the distance of 69 feet; Morton, 90 feet; for a dissyllable 105 feet, a trisyllable 160 feet, a tetrasyllable 182 feet, and a pentasyllable 204 feet. Nat. Hist. Northampton, cap. 5, pa. 358.

From what has been said, it follows that echoes may be applied for measuring inaccessible distances. Thus, Mr. Derham, standing upon the banks of the Thames, opposite to Woolwich, observed that the echo of a single sound was reflected back from the houses in 3 seconds; consequently the sum of the direct and reflex rays must have been  $1142 \times 3 = 3426$  feet, and the half of it, 1713 feet, the breadth of the river in that place.

It also follows that the echoing body being removed farther off, it reflects more of the sound than when nearer; which is the reason why some echoes repeat but one syllable, or one word, and some many. Of these, some are tonical, which only return a voice when modulated into some particular musical tone; and others polysyllabical. That fine echo in Woodstock park, Dr. Plot assures us, in the day-time will return very distinctly 17 syllables, and in the night 20. Nat. Hist. Oxf. cap. 1, pa. 7.

Echoing bodies may be so contrived, and placed, as that reflecting the sound from one to the other, a multiple echo, or many echoes, shall arise.—At Rosneath, near Glasgow, in Scotland, there is an echo that repeats a tune played with a trumpet three times completely and distinctly.—At the sepulchre of Metella, wife of Crassus, there was an echo, which repeated what a man said five times.—Authors mention a tower at Cyzicus, where the echo repeated seven times.—There is an echo at Brussels, that answers 15 times.

One of the finest echoes we read of, is that mentioned by Barthius, in his notes on Statius's Thebais, lib. 6, ver. 30, which repeated the words a man uttered 17 times. This was on the banks of the Naha, between Coblentz and Bingen. And whereas, in common echoes, the repetition is not heard till some time after hearing the words

spoken, or the notes sung; in this, the person who speaks, or sings, is scarce heard at all; but the repetition very clearly, and always in surprising varieties; the echo seeming sometimes to approach nearer, and sometimes farther off; sometimes the voice is heard very distinctly, and sometimes scarce at all: one person hears only one voice, and another several; one hears the echo on the right, and the other on the left, &c.

Addison, and other travellers in Italy, mention an echo at Simonetta palace, near Milan, still more extraordinary, returning the sound of a pistol 56 times. The echo is heard behind the house, which has two wings; the pistol is discharged from a window in one of these wings, the sound is returned from a dead wall in the other wing, and heard from a window in the back-front. See Addison's Travels, pa. 32; Mission's Voyag. d'Ital. tom. 2, pa. 196; Philos. Transf. N° 480, pa. 220.

Farther, a multiple echo may be made, by so placing the echoing bodies, at unequal distances, as that they may reflect all one way, and not one on the other; by which means, a manifold successive sound will be heard; one clap of the hands like many; one *ha* like a laughter; one single word like many of the same tone and accent; and so one musical instrument like many of the same kind, imitating each other.

Lastly, echoing bodies may be so ordered, that from any one sound given, they shall produce many echoes, different both as to tone and intension. By which means a musical room may be so contrived, that not only one instrument playing in it shall seem many of the same sort and size, but even a concert of different ones; this may be contrived by placing certain echoing bodies so, as that any note played, shall be returned by them in 3ds, 5ths, and 8ths.

ECHO is also used for the place where the repetition of the sound is produced, or heard. This is either natural or artificial.

In echoes, the place where the speaker stands, is called the *centrum phonicum*; and the object or place that returns the voice, the *centrum phonocampticum*.

ECHO, in Architecture, is applied to certain vaults and arches, mostly of elliptical or parabolical figures; used to redouble sounds, and produce artificial echoes.—The method of making them is taught by F. Biancani, in his Echometria, at the end of his book on the Sphere.

Vitruvius tells us, that in divers parts of Greece and Italy there were brazen vessels, artfully ranged under the seats of the theatres, to render the sound of the actors' voices more clear, and make a kind of echo; by which means, every one of the prodigious multitude of persons, present at those spectacles, might hear with ease and pleasure.

ECLIPSAREON, an instrument invented by Mr. Ferguson for shewing the phenomena of eclipses; as their time, quantity, duration, progress, &c. Ferguson's Astron., or Philos. Transf. vol. 48, pa. 520.

ECLIPSE, a privation of the light of one of the luminaries, by the interposition of some opaque body, either between it and the eye, or between it and the sun.

The ancients had terrible ideas of eclipses; supposing them prefaces of some dreadful events. Plutarch assures



us, that at Rome it was not allowed to talk publicly of any natural causes of eclipses; the popular opinion running so strongly in favour of their supernatural production, at least those of the moon; for as to those of the sun, they had some idea that they were caused by the interposition of the moon between us and the sun; but were at a loss for a body to interpose between us and the moon, which they thought must be the way, if the eclipses of the moon were produced by natural causes. They therefore made a great noise with brazen instruments, and set up loud shouts, during eclipses of the moon; thinking by that means to ease her in labour: whence Juvenal, speaking of a talkative woman, says, *Una laboranti poterit succurrere lunæ*. Others attributed the eclipse of the moon to the arts of magicians, who, by their enchantments, plucked her out of heaven, and made her skim over the grass.

The natives of Mexico keep fast during eclipses; and particularly their women, who beat and abuse themselves, drawing blood from their arms, &c; imagining the moon has been wounded by the sun, in some quarrel between them.

The Chinese fancy that eclipses are occasioned by great dragons, who are ready to devour the sun and moon; and therefore when they perceive an eclipse, they rattle drums and brass kettles, till they think the monster, terrified by the noise, lets go his prey.

The superstitious notions entertained of eclipses, were once of considerable advantage to Christopher Columbus, the discoverer of America, who being driven on the island of Jamaica in the year 1493, and distressed for want of provisions, was refused relief by the natives; but having threatened them with a plague, and foretelling an eclipse as a token of it, which happened according to his prediction, the barbarians were so terrified, that they strove who should be the first in bringing him supplies, throwing them at his feet, and imploring forgiveness.

*Duration of an ECLIPSE*, is the time of its continuance, or between the immersion and emersion.

*Immersion, or Incidence, of an ECLIPSE*, is the moment when the eclipse begins, or when part of the sun, moon, or planet first begins to be obscured.

*Emergence, or Expurgation, of an ECLIPSE*, is the time when the eclipsed luminary begins to re-appear, or emerge out of the shadow.

*Quantity of an ECLIPSE*, is the part of the luminary eclipsed. To determine the quantity eclipsed, it is usual to divide the diameter of the luminary into 12 equal parts, called digits; whence the eclipse is said to be of so many digits according to the number of them contained in that part of the diameter which is eclipsed or obscured.

Eclipses are divided, with respect to the luminary eclipsed, into *Eclipses of the sun, of the moon, and of the satellites*; and with respect to the circumstances, into *total, partial, annular, central, &c.*

*Annular ECLIPSE*, is when the whole is eclipsed, except a ring, or annulus, which appears round the border or edge.


*Central ECLIPSE*, is one in which the centres of the two luminaries and the earth come into the same straight line.

*Partial ECLIPSE*, is when only a part of the luminary is eclipsed. And a

*Total ECLIPSE*, is that in which the whole luminary is darkened.

*ECLIPSE of the Moon*, is a privation of the light of the moon, occasioned by an interposition of the body of the earth directly between the sun and moon, and so intercepting the sun's rays that they cannot arrive at the moon, to illuminate her. Or, the obscuration of the moon may be considered as a section of the earth's conical shadow, by the moon passing through some part of it.

The manner of this eclipse is represented in this figure, where S is the sun, E the earth, and M or M the moon.



Lunar Eclipses only happen at the time of full moon; because it is only then the earth is between the sun and moon: nor do they happen every full moon, because of the obliquity of the moon's path with respect to the sun's; but only in such full moons as happen either at the intersection of those two paths, called the moon's nodes, or very near them; viz, when the moon's latitude, or distance between the centres of the earth and moon, is less than the sum of the apparent semi-diameters of the moon and the earth's shadow.

*The chief Circumstances in Lunar Eclipses*, are the following:—1. All lunar Eclipses are universal, or visible in all parts of the earth which have the moon above their horizon; and are every where of the same magnitude, with the same beginning and end.—2. In all lunar eclipses, the eastern side is what first immerses and emerges again, i. e. the left-hand side of the moon as we look towards her, from the north; for the proper motion of the moon being swifter than that of the earth's shadow, the moon approaches it from the west, overtakes and passes through it with the moon's east side foremost, leaving the shadow behind, or to the westward.—3. Total eclipses, and those of the longest duration, happen in the very nodes of the ecliptic; because the section of the earth's shadow, then falling on the moon, is considerably larger than her disc. There may however be total eclipses within a small distance of the nodes; but their duration is the less as they are farther from it; till they become only partial ones, and at last, none at all.—4. The moon, even in the middle of an eclipse, has usually a faint appearance of light, resembling tarnished copper; which Gassendus, Ricciolus, Kepler, &c, attribute to the light of the sun, refracted by the earth's atmosphere, and so transmitted thither.—Lastly, she grows sensibly paler and dimmer, before entering into the real shadow; owing to a penumbra which surrounds that shadow to some distance.

*Astronomy of Lunar ECLIPSES, or the method of calculating their times, places, magnitudes, and other phenomena.*

The




The 1st preliminary is to find the length of the earth's conical shadow. This may be found either from the distance between the earth and sun, and the proportion of their diameters, or from the angle of the sun's apparent magnitude at the time. Thus, suppose the semiaxis of the earth's orbit 95,000,000 miles, and the eccentricity of the orbit 1,377,000 miles, making the greatest distance 96,377,000 miles, or 24194 semidiameters of the earth; and the sun's semidiameter being to the earth's, as 112 to 1; then as  $AD : BE :: DB : EC$ , that is,  $111 : 1 :: 24194 : 218$  semidiameters of the earth =  $EC$  the length of the earth's shadow. Otherwise, suppose the angle  $AES$ , or the sun's apparent semidiameter be  $15' 56''$ , and the angle  $BAE$ , or the sun's parallax  $8.6''$ , then is their difference, or the angle  $ACE = 15' 47.4''$ ; hence, as  $\text{tang. } 15' 47.4'' : \text{radius} :: BE \text{ or } 1 : 218 \text{ nearly} = CE$ , the same distance as before. Hence, as the moon's least distance from the earth is scarce 56 semidiameters, and the greatest not more than 64, the moon, when in opposition to the sun, in or near the nodes, will fall into the earth's shadow, and will be eclipsed, as the length of the shadow is almost 4 times the moon's distance.

2. To find the apparent semidiameter of the earth's shadow, in the place where the moon passes through it, at any given time. Add together the sun and moon's parallaxes, and from the sum subtract the apparent semidiameter of the sun; so shall the remainder be the apparent semidiameter of the shadow at the place of the moon's passage. For ex. the 28th of April 1790, at midnight, the moon's parallax is  $61' 9''$ , to which add  $8.6''$ , or  $9''$ , for the sun's parallax, from the sum  $61' 18''$  take  $15' 56''$ , the sun's apparent semidiameter, and the remainder  $45' 22''$  is the semidiameter of the shadow at the place where the moon passes through at that time. N. B. Some omit the sun's parallax, as of no consequence; but increase the apparent semidiameter of the shadow by one whole minute, for the shadow of the atmosphere; which would give the semidiameter of the shadow, in the case above,  $46' 13''$ .

3. There must also be had, the true distance of the moon from the node, at the mean opposition; also the true time of the opposition, with the true place of the sun and moon, reduced to the ecliptic; likewise the moon's true latitude at the time of the true opposition; the angle of the moon's way with the ecliptic, and the true horary motions of the sun and moon: from which all the circumstances of her eclipse may be computed by common arithmetic and trigonometry.

#### To Construct an ECLIPSE of the Moon.

Let  $EW$  be a part of the ecliptic, and  $C$  the centre of the earth's shadow, through which draw perpendicular to  $EW$ , the line  $CN$  towards the north, if the moon have north latitude at the time of the eclipse, or  $CS$  southward, if she have south latitude. Make the angle  $NCD$  equal to the angle of the moon's way with the ecliptic, which may be always taken at  $5^\circ 35'$ , on an average, without any sensible error; and bisect this angle by the right line  $CF$ ; in which line it is that the true equal time of opposition of the sun and moon falls, as given by the tables.



From a convenient scale of equal parts, representing minutes of a degree, take the moon's latitude at the true time of full moon, and set it from  $C$  to  $G$ , on the line  $CF$ ; and through the point  $G$ , at right-angles to  $CD$ , draw the right line  $HKGLI$  for the path of the moon's centre. Then is  $L$  the point in the earth's shadow, where the moon's centre is at the middle of the eclipse;  $G$  the point where her centre is at the tabular time of her being full; and  $K$  the point where her centre is at the instant of her ecliptic opposition; also  $I$  the moon's centre at the moment of immersion, and  $H$  her centre at the end of the eclipse.

With the moon's semidiameter as a radius, and the points  $I, L, H$ , as centres, describe circles for the moon at the beginning, middle, and end of the eclipse.

Finally, the length of the line of path  $IH$ , measured on the same scale, will serve to determine the duration of the eclipse, viz, by saying, As the moon's horary motion from the sun is to  $IH :: 1 \text{ hour or } 60 \text{ min. to the whole duration of the eclipse.}$

To Compute a Lunar ECLIPSE. This will be very easy from the foregoing construction. For, 1st, in the triangle  $CGL$ , right-angled at  $L$ , there are given the hypotenuse  $CG =$  the moon's latitude at the time of full moon, and the angle  $GCL =$  the half of  $5^\circ 35'$ ; to find the legs  $CL$  and  $LG$ .—2d, In the right-angled triangle  $CHL$  or  $CIL$ , are given the leg  $CL$ , and  $CH$  or  $CI$ , the sum of the semidiameters of the moon and the earth's shadow; to find  $LH$  or  $LI$ , half the difference of the sun's and moon's motions during the time of the eclipse.—3d, As the difference of the horary motions of the luminaries is to one hour, or 60 min. ::  $HL$  to the semiduration of the eclipse, and ::  $GL$  to the difference between the opposition and middle of the eclipse; this last therefore taken from the time of full moon, gives the time of the middle of the eclipse; from which subtracting the time in  $LI$ , or semiduration before found, gives the beginning of the eclipse; or add the same, and it gives the end of it.—Lastly, from  $CO$  the semidiameter of the shadow, take  $CL$ , leaves  $LO$ ; to which add  $LP$ , the moon's semidiameter, when necessary, gives  $OP$  the quantity eclipsed.


Note, When the moon's distance from the node exceeds  $12^\circ$ , there can be no eclipse of the moon; or, more accurately, the limit is from  $10\frac{1}{2}$  and  $12\frac{1}{8}$  degrees, according to the distances of the sun, earth, and moon.

ECLIPSE of the Sun, is an occultation of the sun's body.



body, occasioned by an interposition of the moon between the earth and sun. On which account it is by some considered as an eclipse of the earth, since the light of the sun is hid from the earth by the moon, whose shadow involves a part of the earth.

The manner of a solar eclipse is represented in this figure; where S is the sun, *m* the moon, and CD the



earth, *rmso* the moon's conical shadow, travelling over a part of the earth CoD, and making a complete eclipse to all the inhabitants residing in that track, but no where else; excepting that for a large space around it there is a fainter shade, included within all the space rCDs, which is called the *Penumbra*.

Hence, Solar Eclipses happen when the moon is in conjunction with the sun, or at the new moon, and also in the nodes or near them, the limit being about 17 degrees on each side of it; and such eclipses only happen when the latitude of the moon, viewed from the earth, is less than the sum of the apparent semidiameters of the sun and moon; because the moon's way is oblique to the ecliptic, or sun's path, making an angle of nearly  $5^{\circ} 35'$  with it.

In the nodes, when the moon has no visible latitude, the occultation is total; and with some continuance, when the disc of the moon in perigee appears greater than that of the sun in apogee, and its shadow is extended beyond the surface of the earth; and without continuance at moderate distances when the cusp, or point of the moon's shadow, barely touches the earth. Lastly, out of the nodes, but near them, the eclipses are partial.

The other circumstances of solar eclipses are, 1. That none of them are universal; that is, none of them are seen throughout the whole hemisphere which the sun is then above; the moon's disc being much too little, and much too near the earth, to hide the sun from the whole disc of the earth. Commonly the moon's dark shadow covers only a spot on the earth's surface, about 180 miles broad when the sun's distance is greatest, and the moon's least. But her partial shadow, or penumbra, may then cover a circular space of 4900 miles in diameter, within which the sun is more or less eclipsed, as the places are nearer to or farther from the centre of the penumbra. In this case the axis of the shade passes through the centre of the earth, or the new moon happens exactly in the node, and then it is evident that the section of the shadow is circular; but in every other case the conical shadow is cut obliquely by the surface of the earth, and the section will be an oval, and very nearly a true ellipsis.

2. Nor does the Eclipse appear the same in all parts

of the earth, where it is seen; but when in one place it is total, in another it is only partial. Farther, when the moon appears much less than the sun, as is chiefly the case when she is in apogee and he in perigee, the vertex of the lunar shadow is then too short to reach the earth, and though she be in a central conjunction with the sun, is yet not large enough to cover his whole disc, but lets his limb appear like a lucid ring or bracelet, and so causes an *Annular Eclipse*.

3. A Solar Eclipse does not happen at the same time, in all places where it is seen; but appears more early to the western parts, and later to the eastern; as the motion of the moon, and consequently of her shadow, is from west to east.

4. In most Solar Eclipses the moon's disc is covered with a faint light; which is attributed to the reflexion of the light from the illuminated part of the earth.

Lastly, in total Eclipses of the sun, the moon's limb is seen surrounded by a pale circle of light; which some astronomers consider as an indication of a lunar atmosphere; but others as the atmosphere of the sun, because it has been observed to move equally with the sun, and not with the moon; and besides, it is generally believed that the moon is without any atmosphere, unless it be one that is very small, and very rare.

#### *To determine the Bounds of a Solar ECLIPSE.*

If the moon's parallax were insensible, the bounds of a solar eclipse would be determined after the same manner as those of a lunar; but because here is a sensible parallax, the method is a little altered, viz.

1. Add together the apparent semidiameters of the luminaries, both in apogee and perigee; which gives  $33' 6''$  for the greatest sum of them, and  $30' 31''$  for the least sum.

2. Since the parallax diminishes the northern latitude and augments the southern, therefore let the greatest parallax in latitude be added to the former sums, and also subtracted from them: Thus in each case there will be had the true latitude, beyond which there can be no eclipse. This latitude being given, the moon's distance from the nodes, beyond which eclipses cannot happen, is found as for a lunar eclipse. This limit is nearly between  $16\frac{1}{2}$  and  $18\frac{1}{3}$  degrees distance from the nodes.

*To find the Digits eclipsed.* Add the apparent semidiameters of the luminaries into one sum; from which subtract the moon's apparent latitude; the remainder is the scruples, or parts of the diameter eclipsed. Then say, As the semidiameter of the sun is to the scruples eclipsed; so are 6 digits reduced into scruples, viz 360 scruples or minutes, to the digits &c eclipsed.

*To determine the Duration of a Solar ECLIPSE.* Find the horary motion of the moon from the sun, for one hour before the conjunction, and another hour after: then say, As the former horary motion is to the seconds in an hour, so are the scruples of half duration (found as in a lunar eclipse) to the time of immersion; and as the latter horary motion is to the same seconds, so are the same scruples of half duration to the time of emersion. Lastly, adding the times of immersion and emersion together, the aggregate is the total duration.

The moon's apparent diameter when largest, exceeds



needs the sun's when least, only  $2'$  of a degree; and at the greatest solar eclipse that can happen at any time and place, the total darkness cannot continue any longer than whilst the moon is moving through this  $2'$  from the sun in her orbit, which is about 4 minutes of time: for the motion of the shadow on the earth's disc is equal to the moon's motion from the sun, which on account of the earth's rotation on its axis towards the same way, or eastward, is about  $30\frac{1}{2}$  minutes of a degree every hour, at a mean rate; but so much of the moon's orbit is equal to  $30\frac{1}{2}$  degrees of a great circle on the earth, because the circumference of the moon's orbit is about 60 times that of the earth; and therefore the moon's shadow goes  $30\frac{1}{2}$  degrees, or 1830 geographical miles in an hour, or  $30\frac{1}{2}$  miles in a minute.

*To determine the Beginning, Middle, and End, of a Solar Eclipse.* From the moon's latitude, for the time of conjunction, find the arch GL (last fig. but one), or the distance of the greatest obscurity. Then say, as the horary motion of the moon from the sun, before the conjunction, is to 1 hour; so is the distance of the greatest darkness, to the interval of time between the greatest darkness and the conjunction. Subtract this interval, in the 1st and 3d quarter of the anomaly, from the time of the conjunction; and in the other quarters, add it to the same; the result is the time of the greatest darkness. Lastly, from the time of the greatest darkness subtract the time of incidence, and add it to the time of emergence; the difference in the first case will be the beginning; and the sum, in the latter case, the end of the eclipse.

*To Calculate ECLIPSES of the Sun.* First, find the mean new moon, and thence the true one; with the place of the luminaries for the apparent time of the true one.—2. For the apparent time of the true new moon, compute the apparent time of the new moon observed.—3. For the apparent time of the new moon seen, compute the latitude seen.—4. Thence determine the digits eclipsed.—5. Find the times of the greatest darkness, immersion, and emergence.—6. Thence determine the beginning, and ending of the eclipse.

From the foregoing problems, it is evident that all the trouble and fatigue of the calculus arises from the parallaxes of longitude and latitude, without which, the calculation of solar eclipses would be the same with that of lunar ones.

See the Construction and Calculation of Eclipses by Flamsteed in Sir Jonas Moor's System of Mathematics, and in Ferguson's Astronomy, &c.

In the Philos. Trans. N<sup>o</sup> 461 is a contrivance to represent solar eclipses, by means of the terrestrial globe, by M. Seguer, professor of Mathematics at Gottingen. And Mr. Ferguson has fitted a terrestrial globe, so as to shew the time, quantity, duration, and progress of solar eclipses, at any place of the earth where they are visible; which he calls the Eclipsareon. He has also given a large catalogue of ancient and modern eclipses, including those recorded in history, from 721 years before Christ, to A. D. 1485; also computed eclipses from 1485 to 1700, and all the eclipses visible in Europe from 1700 to 1800. See his Astron.

*The Number of ECLIPSES*, of both luminaries, in any year, cannot be less than two, nor more than seven; the most usual number is 4, and it is rare to have more

than 6. The reason is obvious; because the sun passes by both the nodes but once in a year, unless he pass by one of them in the beginning of the year; in which case he will pass by the same again a little before the end of the year; because the nodes move backwards  $19\frac{1}{2}$  degrees every year, and therefore the sun will come to either of them 173 days after the other. And if either node be within  $17^\circ$  of the sun at the time of new moon, the sun will be eclipsed; and at the subsequent opposition, the moon will be eclipsed in the other node, and come round to the next conjunction before the former node be  $17^\circ$  beyond the sun, and eclipse him again. When three eclipses happen about either node, the like number commonly happens about the opposite one; as the sun comes to it in 173 days afterward, and 6 lunations contain only 4 days more. Thus there may be two eclipses of the sun, and one of the moon, about each of the nodes. But when the moon changes in either of the nodes, she cannot be near enough the other node at the next full, to be eclipsed; and in 6 lunar months afterward she will change near the other node; in which case there cannot be more than two eclipses in a year, both of the sun.

*Period of ECLIPSES*, is the period of time in which the same eclipses return again; and as the nodes move backwards  $19\frac{1}{2}$  degrees every year, they would shift through every point of the ecliptic in 18 years and 225 days; and this would be the regular period of their return, if any complete number of lunations were finished without a fraction; but this is not the case. However, in 223 mean lunations, after the sun, moon, and nodes have been once in a line of conjunction, they return so nearly to the same state again, as that the same node which was in conjunction with the sun and moon at the beginning of the first of these lunations, will be within  $28' 12''$  of the line of conjunction with the sun and moon again, when the last of these lunations is completed; and in this period there will be a regular return of eclipses for many ages. To the mean time of any solar or lunar eclipse, by adding this period, or 18 Julian years 11 days 7 hours 43 minutes 20 seconds, when the last day of February in leap years is 4 times included, or a day less when it occurs 5 times, we shall have the mean time of the return of the same eclipse. In an interval of 6890 mean lunations, containing 557 years 21 days 18 hours 30 minutes 11 seconds, the sun and node meet so nearly, as to be distant only 11 seconds.

*The Use of ECLIPSES.* In Astronomy, eclipses of the moon determine the spherical figure of the earth; they also shew that the sun is larger than the earth, and the earth than the moon. Eclipses also, that are similar in all circumstances, and that happen at considerable intervals of time, serve to ascertain the period of the moon's motion. In Geography, eclipses discover the longitude of different places; for which purpose those of the moon are the more useful, because they are more often visible, and the same lunar eclipse is of equal magnitude and duration at all places where it is seen. In Chronology, both solar and lunar eclipses serve to determine exactly the time of any past event.

*ECLIPSES of the Satellites.* See SATELLITES of Jupiter.

The chief circumstances here observed, are, 1. That the satellites of Jupiter undergo two or three kinds of eclipses;



eclipses; the first of which are proper, being such as happen when Jupiter's body is directly interposed between them and the sun: and these happen almost every day. Various authors have given tables for computing eclipses of the satellites of Jupiter; as Flamsteed, Cassini, &c, but the latest and best of all, are those of professor Wargentin of Upsal.

The second sort are occultations, rather than observations; when the satellites, coming too near the body of Jupiter, are lost in his light; which Riccioli calls *occidere zeusice, setting jovially*. In which case, the nearest or first satellite exhibits a third kind of eclipse; being observed like a round macula, or dark spot, transiting the disc of Jupiter, with a motion contrary to that of the satellite; like as the moon's shadow projected on the earth, will appear to do, to the lunar inhabitants.

The eclipses of Jupiter's satellites furnish very good means of finding the longitude at sea. Those especially of the first satellite are much surer than the eclipses of the moon, and they also happen much oftener: the manner of applying them is also very easy. See LONGITUDE.

**ECLIPTIC**, in Astronomy, a great circle of the sphere conceived to pass through the middle of the zodiac. It is sometimes called the *via solis*, or *sun's path*, being the track which he appears to describe among the fixed stars; though more properly it is the apparent path of the earth, as viewed from the sun, and thence called the heliocentric circle of the earth. It is called the ecliptic, because all the eclipses of the sun or moon happen when the moon crosses it, or is nearly in one of those two parts of her orbit where it crosses the ecliptic, which points are called the moon's nodes.

Upon the ecliptic are marked and counted the 12 celestial signs, Aries, Taurus, Gemini, &c; and upon it is counted the longitude of the planets and stars. It is placed obliquely with respect to the equator, which it cuts in two opposite points, viz, the beginning of Aries and Libra, which are directly opposite to each other, and called the equinoxes, making the one half of the ecliptic to the north, and the other half on the south side of the equator; the two extreme points of it, to the north and south, which are opposite to each other, and at a quadrant distance from the equinoctial points both ways, are called the solstices, or solstitial points, or also the two tropics, which are at the beginning of Cancer and Capricorn, and which are at the farthest distance of any points of it from the equator, which distance is the measure of the sun's greatest declination, which is the same with the obliquity of the ecliptic, or the angle it makes with the equator.

*This obliquity of the ecliptic* is not permanent, but is continually diminishing, by the ecliptic approaching nearer and nearer to a parallelism with the equator, at the rate of half a second in a year nearly, or from 50'' to 55'' in 100 years, as is deduced from ancient and modern observations compared together; and as the mean obliquity of the ecliptic was  $23^{\circ} 28'$  about the end of the year 1788, or beginning of 1789, by adding half a second for each preceding year, or subtracting the same for each following year, the mean obliquity will be found nearly for any year either before or since that period. The quantity however of this change is variously stated by different authors, from 50'' to 60'' or

70'' for each century or 100 years. Hipparchus, almost two thousand years since, observed the obliquity of the ecliptic, and found it about  $23^{\circ} 51'$ ; and all succeeding astronomers, to the present time, having observed the same, have found it always less and less; being now rather under  $23^{\circ} 28'$ ; a difference of about 23' in 1950 years; which gives a medium of 70'' in 100 years. There is great reason however to think that the diminution is variable.

This diminution of the obliquity of the ecliptic to the equator, according to Mr. Long and some others, is chiefly owing to the unequal attraction of the sun and moon on the protuberant matter about the earth's equator. For if it be considered, say they, that the earth is not a perfect sphere, but an oblate spheroid, having its axis shorter than its equatorial diameter; and that the sun and moon are constantly acting obliquely upon the greater quantity of matter about the equator, drawing it, as it were, towards a nearer and nearer coincidence with the ecliptic; it will not appear strange that these actions should gradually diminish the angle between the planes of those two circles. Nor is it less probable that the mutual attractions of all the planets should have a tendency to bring their orbits to a coincidence: though this change is too small to become sensible in many ages.

It is now however well known that this change in the obliquity of the ecliptic, is wholly owing to the actions of the planets upon the earth, and especially the planets Venus and Jupiter, but chiefly the former. See La Grange's excellent paper upon this subject in the Memoirs of the French Academy for 1774; Cassini's in 1778; and La Lande's Astron. vol. 3, art. 2737. According to La Grange, who proceeds upon theory, the annual change of obliquity is variable, and has its limits: about 2000 years ago, he thinks it was after the rate of about 38'' in 100 years; that it is now, and will be for 400 years to come, 56'' per century; but 2000 years hence, 49'' per century. According to Cassini, who computes from observations of the obliquity between the years 1739 and 1778, the annual change at present is 60'' or 1' in 100 years. But according to La Lande, the diminution is at the rate of 88'' per century; while Dr. Maskelyne makes it only 50'' in the same time.

Beside the regular diminution of the obliquity of the ecliptic, at the rate of near 50 seconds in a century, or half a second a year, which arises from a change of the ecliptic itself, it is subject to two periodical inequalities, the one produced by the unequal force of the sun in causing the precession of the equinoxes, and the other depending on the nutation of the earth's axis. See the Explanation and Use of Dr. Maskelyne's Tables and Observations, pa. vi, where we are shewn how to calculate those inequalities, and where he shews that, from his own observations, the mean obliquity of the ecliptic to the beginning of the year 1769, was  $23^{\circ} 28' 9''.7$ .

*To find the Obliquity of the Ecliptic*, or the greatest declination of the sun: about the time of the summer solstice observe very carefully the sun's zenith distance for several days together; then the difference between this distance and the latitude of the place, will be the obliquity sought, when the sun and equator are both on one side of the place of observation; but their sum will



will be the obliquity when they are on different sides of it. Or, it may be found by observing the meridian altitude, or zenith distance, of the sun's centre, on the days of the summer and winter solstice; then the difference of the two will be the distance between the tropics, the half of which will be the obliquity sought.

By the same method too, the declination of the sun from the equator for any other day may be found; and thus a table of his declination for every day in the year might be constructed. Thus also the declination of the stars might be found.

Authors' Names	Years before Christ	Obliquity		
		°	'	"
Pytheas - - -	324	23	49	23
Eratosthenes and Hipparchus	230 & 140 after Christ	23	51	20
Ptolomy - - -	140	23	48	45
Almahmon - - -	832	23	35	
Albategnius - - -	880	23	35	
Thebat - - -	911	23	33	30
Abul Wafi and Hamed	999	23	35	
Persian Tables in Chrysoceca	1004	23	35	
Albatrunius - - -	1007	23	35	
Arzachel - - -	1104	23	33	30
Almaeon - - -	1140	23	33	30
Choja Nassir Oddin - -	1290	23	30	
Prophatius the Jew - -	1300	23	32	
Ebn Shattir - - -	1363	23	31	
Purbach and Regiomontanus	1460	23	30	
Ulugh Beigh - - -	1463	23	30	17
Walther - - -	1476	23	30	
Do. corrected by refraction &c		23	29	8
Werner - - -	1510	23	28	30
Copernicus - - -	1525	23	28	24
Egnatio Danti - - -	1570	23	29	
Prince of Hesse - - -	1570	23	31	
Rothmann and Byrge - -	1570	23	30	20
Tycho Brahe - - -	1584	23	31	30
Ditto corrected - - -		23	29	
Wright - - -	1594	23	30	
Kepler - - -	1627	23	30	30
Gassendus - - -	1630	23	31	
Ricciolus - - -	1646	23	30	20
Ditto corrected - - -	1655	23	29	
Hevelius - - -	1653	23	30	20
Ditto corrected - - -	1661	23	28	52
Cassini - - -	1655	23	29	15
Montons, corrected, &c	1660	23	29	3
Richer corrected - - -	1672	23	28	52
De la Hire - - -	1686	23	29	
Ditto corrected - - -		23	29	28
Flamsteed - - -	1690	23	29	
Bianchini - - -	1703	23	28	25
Roemer - - -	1706	23	28	41
Louville - - -	1715	23	28	24
Godin - - -	1730	23	28	20
Bradley - - -	1750	23	28	18
Mayer - - -	1756	23	28	16
Maskelyne - - -	1769	23	28	10
Hornsbey - - -	1772	23	28	8

The observations of astronomers of all ages, on the obliquity of the ecliptic, have been collected together; and although some of them may not be quite accurate, yet they sufficiently shew the gradual and continual decrease of the obliquity from the times of the earliest observations down to the present time. The chief of those observations may be seen in the foregoing table; where the first column contains the name of the observer, or author, the 2d the year before or after Christ, and the 3d the obliquity of the ecliptic for that time.

See Ptolom. Alm. lib. 1, cap. 10; Riccioli Alm. vol. 1, lib. 3, cap. 27; Flamsteed Proleg. Hist. Cœl. vol. 3; Philos. Transf. number 163; ib. vol. 63, pt. 1; Long's Astron. vol. 1, cap. 16; Memoirs of the Acad. an. 1716, 1734, 1762, 1767, 1774, 1778; Acta Erud. Lipsiæ 1719; Naut. Alm. 1779; Maskelyne's Observ. Explan. pa. vi; &c.

According to an ancient tradition of the Egyptians, mentioned by Herodotus, the ecliptic had formerly been perpendicular to the equator: they were led into this notion by observing, for a long series of years, that the obliquity was continually diminishing; or, which amounts to the same thing, that the ecliptic was continually approaching to the equator. From thence they took occasion to suspect that those two circles, in the beginning, had been as far off each other as possible, that is, perpendicular to each other. Diodorus Siculus relates, that the Chaldeans reckoned 403,000 years from their first observations to the time of Alexander's entering Babylon. This enormous account may have some foundation, on the supposition that the Chaldeans built on the diminution of the obliquity of the ecliptic at the rate of a minute in 100 years. M. de Louville, taking the obliquity such as it must have been at the time of Alexander's entrance into Babylon, and going back to the time when the ecliptic, at that rate, must have been perpendicular to the equator, actually finds 402,942 Egyptian, or Chaldean years; which is only 58 years short of that epocha. Indeed there is no way of accounting for the fabulous antiquity of the Egyptians, Chaldeans, &c, so probable, as from the supposition of long periods of very slow celestial motions, a small part of which they had observed, and from which they calculated the beginning of the period, making the world and their own nation to commence together. Or perhaps they sometimes counted months or days for years.

Should the diminution always continue at the rate it has lately done, viz at 50'' or 56'' a century, it would take 96,960 years, from the year 1788, to bring the ecliptic exactly to coincide with the equator.

**ECLIPTIC**, in *Geography*, a great circle on the terrestrial globe, in the plane of, or directly under, the celestial ecliptic.

**ECLIPTIC**, *Eclipticus*, something belonging to the ecliptic, or to eclipses; as ecliptic conjunction, opposition, &c.

**ECLIPTIC Bounds**, or *Limits*, are the greatest distances from the nodes at which the sun or moon can be eclipsed; namely, near 18 degrees for the sun, and 12 degrees for the moon.

**ECLIPTIC Digits**, *digiti ecliptici*. See **DIGITS**.

**Poles of the ECLIPTIC**, are the two opposite points of



of the sphere which are each everywhere equally distant from the ecliptic quite around, or  $90^\circ$  distant from it. The distance of the poles of the ecliptic from the poles of the equator, or of the world, is always equal to the varying distance of the obliquity of the ecliptic, and at the beginning of the year 1789 it was just  $23^\circ 28'$ .

*Reduction to the ECLIPTIC.* See REDUCTION.

EFFECT, the result or consequence of the application of a cause, or agent, on some subject. It is one of the great axioms in philosophy, that full effects are always proportional to the powers of their adequate causes.

EFFECTION, denotes the geometrical construction of a proposition. The term is also used in reference to problems and practices; which, when they are deducible from, or founded upon some general propositions, are called the *geometrical effection* of them.

EFFERVESCENCE, is popularly used for a light ebullition, or a brisk intestine motion, produced in a liquor by the first action of heat, with any remarkable separation of its parts.

EFFICIENT Cause, is that which produces an effect. See CAUSE and EFFECT.

EFFICIENTS, in Arithmetic, are the numbers given for an operation of multiplication, and are otherwise called the factors. Hence the term coefficients in Algebra, which are the numbers prefixed to, or that multiply the letters or algebraic quantities.

EFFLUVIUM, a flux or exhalation of minute particles from any body; or an emanation of subtile corpuscles from a mixed, sensible body, by a kind of motion of transpiration.

ELASTIC, an appellation given to all bodies endowed with the property of elasticity or springiness.

ELASTIC Body, is that which changes its figure, and yields to any impulse or pressure, but endeavours by its own nature and force to restore the same again; or, it is a springy body, which, when compressed or condensed, or the like, makes an effort to set itself at liberty, and to repel the body that constrained it. Such, for instance, as a bow, or a sword blade, &c, which are easily bent, but presently return to their former figure and extension. All bodies partake of this property in some degree, though perhaps none are perfectly elastic, as none are found to restore themselves with a force equal to that with which they are compressed.

The principal phenomena observable in Elastic bodies, are 1, That an elastic body (i. e. a body perfectly elastic, if any such there be) endeavours to restore itself with the same force with which it is pressed or bent. 2, An elastic body exerts its force equally towards all sides; though the effect is chiefly found on that side where the resistance is weakest; as is evident in the case of a gun exploding a ball, a bow shooting out an arrow, &c.—3, Elastic bodies, in what manner soever struck, or impelled, are inflected and rebound after the same manner: thus a bell yields the same musical sound, in what manner, or on what side soever it be struck; the same of a tense or musical chord; and a body rebounds from a plane in the same angle in which it meets or strikes it, making the angle of incidence equal to the angle of reflection, whether the intensity of the stroke be greater or less.—4, A body perfectly fluid, if any such

there be, cannot be elastic, if it be allowed that its parts cannot be compressed.—5, A body perfectly solid, if any such there be, cannot be elastic; because, having no pores, it is incapable of being compressed.—6, The elastic properties of bodies seem to differ, according to their greater or less density or compactness, though not in an equal degree: thus, metals are rendered more compact and elastic by being hammered: tempered steel is much more elastic than soft steel; and the density of the former is to that of the latter as 7809 to 7738: cold condenses solid bodies, and renders them more elastic; whilst heat, that relaxes them, has the opposite effect: but, on the contrary, air, and other elastic fluids, are expanded by heat, and rendered more elastic.—For the laws of Motion and Percussion in Elastic bodies, see MOTION, and PERCUSSION.

ELASTIC Curve. See CATENARIA.

ELASTIC Fluids. See AIR, ELECTRICITY, GAS, ELASTIC VAPOURS, &c.

ELASTIC Gum. The same as CAOUTCHOUC, or Indian Rubber.

ELASTIC Vapours, or Fluids, are such as may be compressed mechanically into a less space, and which resume their former state when the compressing force is withdrawn. Such as atmospherical air, and all the aerial fluids, with all kinds of fumes raised by means of heat, whether from solid or fluid bodies.

Of these, some remain elastic only while a considerable degree of heat is applied to them, or to the substance which produces them; while others continue elastic in every degree of cold that has yet been observed. Of the former kind, are the vapours of water, spirit of wine, mercury, sal-ammoniac, and all kinds of sublimable salts: of the latter, those of spirit of salt, mixtures of vitriolic acid and iron, nitrous acid, and various other metals, and in short the several species of aerial fluids indiscriminately.

The elastic force with which any one of these fluids is endowed, has not yet been calculated, as being ultimately greater than any obstacle we can put in its way. Thus, on compressing the atmospherical air, we find that for some little time at first it easily yields to any force applied; but at every succeeding moment the resistance becomes always the stronger, and a greater and greater force must be applied, to compress it farther. As the compression goes on, the vessel containing the air becomes hot; but no power whatever has yet been able in any degree to destroy the elasticity of the contained fluid; for, upon removing the pressure, it is always found to occupy the very same space that it did before. The case is the same with the steam of water, to which a sufficient heat is applied to keep it from condensing into water.

ELASTICITY, or ELASTIC Force, that property of bodies by which they restore themselves to their former figure, after any external pressure.

The cause or principle of this important property, elasticity, is variously accounted for. The Cartesians ascribe it to their subtile matter making an effort to pass through pores that are too narrow for it. Thus, say they, in bending or compressing a hard elastic body, as a bow, for instance, its parts recede from each other on the convex side, and approach on the concave one: conse-



consequently the pores are contracted or straitened on the concave side; and, if they were before round, are now perhaps oval: so that the *materia subtilis*, or matter of the second element, endeavouring to pass out of the pores thus straitened, must make an effort, at the same time, to restore the body to the state it was in when the pores were rounder, i. e. before the bow was bent: and in this consists its Elasticity.

Other later philosophers account for Elasticity much after the same manner as the Cartesians; with this only difference, that instead of the subtle matter of the Cartesians, these substitute *Ether*, or a fine ethereal medium that pervades all bodies.

Others, setting aside the precarious notion of a *materia subtilis*, account for Elasticity from the great law of nature, Attraction, or the cause of the cohesion of the parts of solid and firm bodies. Thus, say they, when a hard body is struck or bent, so that the component parts are moved a little from each other, but not quite disjointed or broken off, or separated so far as to be out of the power of that attracting force by which they cohere; they must, on removing the external violence, spring back to their former natural state.

Others again resolve Elasticity into the pressure of the atmosphere: for a violent tension, or compression, though not so great as to separate the constituent particles of bodies far enough to let in any foreign matter, must yet occasion many little vacuola between the separated surfaces; so that on the removal of the force they will close again by the pressure of the aerial fluid upon the external parts.

Lastly, others attribute the Elasticity of all hard bodies to the power of resiliion in the air included within them; and so make the elastic force of the air the principle of Elasticity in all other bodies. See Desaguliers's Exper. Philos. vol. 2, pa. 38, &c.

The ELASTICITY of *Fluids* is accounted for from their particles being all endowed with a centrifugal force; whence Sir Isaac Newton demonstrates, prop. 23, lib. 2, that particles, which naturally avoid or fly off from one another by such forces as are reciprocally proportional to the distances of their centres, will compose an elastic fluid, whose density shall be proportional to its compression; and vice versa, if any fluid be composed of particles that fly off or avoid one another, and have its density proportional to its compression, then the centrifugal forces of those particles will be reciprocally proportional to the distances of their centres.

ELASTICITY of the *Air* is the force with which that element endeavours to expand, and with which it does actually dilate itself, on removing the force that compressed it. See AIR, and ATMOSPHERE.

The Elasticity or spring of the air was first discovered by lord Bacon, and farther established by Galileo. Its existence is proved by this experiment of that philosopher: An extraordinary quantity of air being intruded, by means of a syringe, into a hollow ball or shell of glass or metal, till such time as the ball, with this accession of air, weigh considerably more in the balance than it did before; then, opening the mouth of the ball, the air rushes out, till the ball sink to its former weight. From hence we infer, that there is just as much air gone out, as compressed air had been crowded

in. Air therefore returns to its former degree of expansion, upon removing the force that compressed or resisted its expansion; and consequently it is endowed with an elastic force. It may be added, that as the air is found to rush out in every situation or direction of the orifice, the elastic force acts every way, or in every direction alike.

The cause of Elasticity in air hath been usually ascribed to a repulsion between its particles; but what is the cause of that repulsion? The term repulsion, like that of attraction, requires to be defined; and probably it will be found in most cases to be the effect of the action of some other fluid. Thus, it is found that the Elasticity of the atmosphere is very considerably affected by heat. Supposing a quantity of air heated to such a degree as to raise Fahrenheit's thermometer to 212, it will then occupy a considerable space; but if it be cooled again to such a degree, as to sink the thermometer to 0, it will shrink up to less than half the former bulk. The quantity of repulsive power therefore acquired by the air, while passing from one of these states to the other, is evidently owing to the heat added to it, or taken away from it. Nor does there seem to be any reason to suppose, that the quantity of Elasticity or repulsive power it still possesses, is owing to any other cause than the fire contained in it. The supposition that repulsion is a primary cause, independent of all others, has given rise to many erroneous theories, and very much embarrassed philosophers in accounting for the phenomena of Elasticity.

The Elasticity of the air is not only proportional to its density, but is always equal to the force which compresses it, because these two exactly balance each other. This Elasticity, in the atmospheric air, is measured by the height of the barometer at any time, allowing for its heat or temperature, after this rate, viz., the 434th part for each degree of Fahrenheit's thermometer, above or below some mean temperature, as 55°; for by that part of the whole it is that air expands or contracts, or else increases or decreases in its Elasticity, for each degree of the thermometer. Sir Geo. Shuckburgh, in the Philos. Transf. for 1777, pa. 561.

ELECTIONS, or *Choice*, signify the several different ways of taking any number of things proposed, either separately, or as combined in pairs, in threes, in fours, &c; not as to the order, but only as to the number and variety of them. Thus, of the things *a, b, c, d, e*, &c, the elections of

one thing are (*a*,)  $1 = 2^1 - 1$ ,  
two things are (*a, b, ab*)  $3 = 2^2 - 1$ ,  
three things are (*a, b, c, ab, ac, bc, abc*)  $7 = 2^3 - 1$ ,  
&c; and of any number, *n*, all the elections are  $2^n - 1$ ; that is, one less than the power of 2 whose exponent is *n*, the number of single things to be chosen, either separately or in combination.

ELECTRIC, in Physics, is a term applied to those substances, in which the electric fluid can be excited, and accumulated, without transmitting it; and which are therefore called *non-conductors*. They are also called *original Electrics*, and *Electrics per se*.

The word is derived from *ηλεκτρον*, *amber*, one of the most observable non-conductors. To this class also belong glass, and all vitrifications, even of metals; all precious stones, of which the most transparent are the best;



best; all resins, and resinous compositions; also sulphur, baked wood, all bituminous substances, wax, silk, cotton, all dry animal substances, as feathers, wool, hair, &c; also paper, white sugar, and sugar-candy; likewise air, oils, chocolate, calces of metals and semi-metals, the ashes of animal and vegetable substances, the rust of metals, all dry vegetable substances, and stones, of which the hardest are the best.

Substances of this kind may be excited, so as to exhibit the Electric appearances of attracting and repelling light bodies, emitting a spark of light, attended with a snapping noise, and yielding a current of air, the sensation of which resembles that of a spider's web drawn over the face, &c, and a smell like that of phosphorus; and this exciting may be either by friction, or by heating and cooling, or by melting, and pouring one melted substance into another.

The term is peculiarly applied to the electric, viz. the globe, or cylinder, &c, used in electrical machines, to collect the electrical matter by rubbing it.

*ELECTRICAL Air Thermometer*, an instrument contrived by Mr. Kinnerley of Philadelphia, and used in determining the effects of the electrical explosion upon air. The description may be seen in Franklin's Letters, &c, pa. 389, 4to, 1769.

*ELECTRICAL Apparatus*, consists of glass tubes, about 3 feet long, and an inch and a half in diameter, one of which should be closed at one end, and furnished at the other end with a brass cap and stop-cock, to rarefy or condense the inclosed air; sticks of sealing wax, or tubes of rough glass, or glass tubes covered with sealing-wax, or cylinders of baked wood for producing the negative electricity; with proper rubbers, as black oiled silk, with amalgam upon it for the former, and soft new flannel, or hare skins, or cat skins, tanned with the hair on, for the latter; coated jars, or plates of glass, either single, or combined in a battery, for accumulating electricity; metal rods, as dischargers; an electrical machine; electrometers, and insulated stools, supported by pillars of glass, covered with sealing-wax, or baked wood, varnished or boiled in linseed oil.

*ELECTRICAL Atmosphere*, is a stream or mass of the Electrical fluid which surrounds an excited or electrified body, to some distance.

*ELECTRICAL Balls*. See BALLS and ELECTROMETER.

*ELECTRICAL Battery*, consists of a large quantity of coated jars, placed near each other in a convenient manner. These being charged, or electrified, and connected with each other, are then suddenly exploded or discharged, with a prodigious effect.

*ELECTRICAL Fluid*, is a fine rare fluid which issues from, and surrounds electrified bodies.

*ELECTRICAL Kite*, was contrived by Dr. Franklin, to verify his hypothesis of the identity of electricity and lightning. It consisted of a large thin silk handkerchief, extended and fastened at the four corners to two slender strips of cedar, and accommodated with a tail, loop, and string, so as to rise in the air like a paper kite. To the top of the upright stick of the cross was fixed a very sharp-pointed wire, rising a foot or more above the wood; and to the end of the twine, next the hand, a silk ribband was tied. From a key

suspended at the junction of the twine and silk, when the kite is raised during a thunder-storm, a phial may be charged, and electric fire collected, as is usually done by means of a rubbed glass tube or globe. Philos. Transf. vol. 47, pa. 565, or Franklin's Letters, pa. 111 and 112.

Kites made of paper, covered with varnish, or with well boiled linseed oil, to preserve them from the rain, with a stick and cane bow, like the common ones used by boys, will answer the purpose extremely well, and are very useful in determining the electricity of the atmosphere. See CONDUCTOR.

*ELECTRICAL Machine*, is a part of the Electrical apparatus, contrived for collecting a great quantity of electricity, and exhibiting its effects in a very sensible manner. It consists of the electric, the moving engine, the rubber, and the prime conductor. In the early state of this science, for the electric, was used sealing-wax, sulphur, or rough glass; but, since the method of insulating the rubber, and so producing negative electricity, was introduced, smooth glass has been used. The form is commonly either that of a globe, or of a cylinder. Each figure has its advantages, and its inconveniences. Dr. Van Marum, a late German writer, has constructed a machine, in which gumlac, in the form of a disc, is used as an electric instead of glass; which has the effect of depending very little on the temperature of the air; described in his *Verhandelinge over het Electrizeeren*, &c, or a *Treatise concerning the method of electrifying*. Groningen, 1776. But he has since procured some others to be made by Mr. Cuthbertson, a very ingenious artist, of large discs, or round plates of glass: one of these is now placed in Teyler's Museum at Harlem, having two of these glass plates, of 65 inches diameter, excited, on both sides of them, by rubbers of waxed taffaty; with which, effects are produced that are truly astonishing and tremendous. See his *Description of this machine*, and its effects, published in 4to, at Harlem, 1785, &c.

There have been various contrivances for giving motion to the electric of a machine. The common method is by a wheel turned by a winch or handle; a cord going round a groove in the periphery of the wheel, and over a pulley in the neck of the globe or cylinder. Others have used multiplying wheels, which are easily turned by a winch; and others again make use of a wheel and pinion, or a wheel and endless screw. But Van Marum's machine it seems has the completest movement, its operation being very uniform, and easily worked; it is kept in motion by a weight, which, after being wound up to the height of 12 feet, will continue the motion uniformly for 6 hours; yielding also a negative power, as well as the positive; and the conductors annexed to it serving easily to convey the electrical power wherever it is required, without the addition of any chain, or wires, &c.

The Rubber is the next material part of a machine. These were formerly made of red-basil skins, stuffed with hair, wool, flax, or bran: Dr. Nooth introduced silk cushions stuffed with hair, over which is laid a piece of leather, rubbed with amalgam, which are better than the others. The rubber may be insulated in any way that best suits the construction of the machine:



chine: and a chain or wire may easily be suspended from it, to communicate with the floor, whenever the insulation is not necessary; and thus positive and negative electricity may be produced at pleasure. Van Marum uses mercury to his rubbers.

The Prime Conductor is another necessary appendage to the Electrical Machine: its use is to receive the electricity from the electric, as it is produced, and accumulate it as in a magazine, ready to be drawn off and employed on all occasions. See *PRIME CONDUCTOR*.

*Description of the most useful Electrical Machines.*

Fig. 1, plate ix, represents Dr. Priestley's Machine, a very extensively useful one, described in his History of Electricity; in which *g* is the globe, or electric; *f* the rubber; in the two pillars *d, d*, of baked wood, are several holes to receive the spindles of different globes or cylinders, several of which may be put on together, to increase the electricity: *klm* is the prime conductor, being a copper tube, supported on a stand of glass or baked wood.

Fig. 2 is Dr. Watson's Machine, for using several globes at once, to accumulate a great quantity of electricity.

Fig. 3 represents a very portable Electrical Machine invented by Mr. Read, and improved by Mr. Lane. *A* is the glass cylinder, moved vertically by means of the pulley at the lower end of the axis, the pulley being turned by the large wheel *B* parallel to the table; there are several pulleys, of different sizes, either of which may be used, according as the motion is required to be quicker or slower. The conductor *C* is furnished with points to collect the fluid, and is screwed to the wire of a coated jar *D*. The figure shews also the manner of applying Mr. Lane's electrometer to this machine.

Electrical Machines have of late years undergone some very essential alterations and improvements; both from the suggestions of private electricians, and the inventions of Messrs. Adams, Nairne, and Jones, instrument makers in London; some of which are as follow:

Fig. 4 represents a very convenient machine for practice. The frame of this machine consists of the bottom board *ABCD*; which, when the machine must be used, is fastened to the table by two metal cramps. *EF* are two round pillars, of baked wood, which support the cylinder *G* by the axles of the brass or wooden caps *H*, turned sometimes by a simple winch *I*, and sometimes by a pulley and wheel, as in the next fig. The rubber is fixed to a glass pillar *K*, which is fastened to a wooden basis *L* at the bottom. The conductor *N* is usually made of brass or tin japanned, and is insulated by a glass pillar, screwed into a wooden basis or foot, which is most conveniently placed parallel to the cylinder.

Fig. 5 represents an Electrical Machine, with a conductor in the shape of a *T*; and an improved medical apparatus, where it is necessary to give the shock in the arms.

Fig. 6 shews Mr. Nairne's patent machine for medical purposes. Its glass cylinder is about 7 inches in diameter, and 12 long, with two conductors parallel to it. The rubber is fastened to the conductor *R*; and consists of a cushion of leather stuffed, having a piece

of silk glewed to its under part. The conductors are of tin covered with black lacker, each of them containing a large coated glass jar, and likewise a smaller one, or a coated tube, which are visible when the caps *NN* are removed. To each conductor is fixed a knob *O*, for the occasional suspension of a chain to produce positive or negative electricity. That part of the winch *C* which acts as a lever in turning the cylinder, is of glass. Thus every part of the machine is insulated, the cylinder itself and its brass caps not excepted; by which means very little of the electricity is dissipated, and hence of course the effects are likely to be the more powerful. And to this the inventor has adapted some flexible conducting joints, a discharging electrometer, &c, for the practice of medical electricity.

The large Electrical Machine placed in Teyler's Museum at Harlem, has been partly described above. It was constructed by Mr. John Cuthbertson, an English instrument maker; and it has, for the electric, two glass plates of 65 inches diameter, made of French glass, as this is found to produce the most electricity next to English flint glass, which could not be made of a sufficient size: these plates are set on the same horizontal axis, at the distance of  $7\frac{1}{2}$  inches, and are excited by 8 rubbers, each  $15\frac{1}{2}$  inches long; and both sides of the plates are covered with a resinous substance to the distance of  $16\frac{1}{2}$  inches from the centre, both to strengthen the plates, and to prevent any electricity from being carried off by the axis. Its battery of jars contains 225 square feet of coated surface, and its effects are astonishingly great.

*ELECTRICAL Phial.* See *LEYDEN Phial*.

*ELECTRICAL Rubber.* See *ELECTRICAL Apparatus*, and *ELECTRICAL Machine*.

*ELECTRICAL Shock*, is the sudden explosion between the opposite sides of a charged electric; so called because if the discharge be made through the body of an animal, it occasions a sudden motion by the contraction of the muscles through which it passes, accompanied with a disagreeable sensation. The force of this shock is proportioned to the quantity of coated surface, the thinness of the glass, and the power of the machine by which it is charged. Its velocity is almost instantaneous, and it has not been found to take up the least sensible time in passing to the greatest distances.

It has been observed that the Electrical Shock is weakened by being communicated through several persons in contact with one another. Indeed it is obstructed in its passage, even through the best conductors, as it will prefer a short passage through the air to a long one through the most perfect conductors; and if the circuit be interrupted, either by electrics, or very imperfect conductors of a moderate thickness, the shock will rend them in its passage, disperse them in every direction, and exhibit the appearance of a sudden expansion of the air about the centre of the shock. A strong shock made to pass through or over the belly of a muscle, forces it to contract; and sent through a small animal body, deprives it instantly of life, and hastens putrefaction. It gives polarity to magnetic needles, reverses their poles, and produces effects precisely similar, though inferior in degree, to those of lightning.

*ELECTRICAL Star.* See *STAR*.



**ELECTRICITY**, or **ELECTRICAL Force**, is that power or property, which was first observed in amber, the lyncurium, or tourmalin, and which sealing-wax, glass, and a variety of other substances, called electrics, are now known to possess, of attracting light bodies, when excited by heat or friction; and which is also capable of being communicated in particular circumstances to other bodies.

**ELECTRICITY** also denotes the science, or that part of natural philosophy, which proposes to investigate the nature and effects of this power. From *ηλεκτρον*, the Greek name for amber, is derived the term Electricity, which is now very extensively applied, not only to the power of attracting light bodies inherent in amber, but to other similar powers, and their various effects, in whatever bodies they reside, or to whatever bodies they may be communicated.

Muschenbroek and Æpinus have observed a considerable analogy, in a variety of particulars, between the powers of Electricity and Magnetism; and they have also pointed out many instances in which they differ.

*History of ELECTRICITY.*—The property which amber possesses of attracting light bodies, was very anciently observed. Thales of Miletus, 600 years before Christ, concluded from hence that it was animated. But the first person who expressly mentioned this substance, was Theophrastus, about 300 years before Christ. The attractive property of amber is also occasionally noticed by Pliny, and other later naturalists, particularly Gassendus, Kenelm Digby, and Sir Thomas Brown. But it was generally apprehended that this quality was peculiar to amber and jet, and perhaps agate, till W. Gilbert, a native of Colchester, and a physician in London, published his treatise *De Magnete*, in the year 1600. Dr. Gilbert made many considerable experiments and discoveries, considering the then infant state of the science. He enlarged the list both of electrics, and of the bodies on which they act: he remarked, that a dry air was most favourable to electrical appearances, whilst a moist air almost annihilates the electric virtue: he also observed the conical figure assumed by electrified drops of water: he considered electrical attraction separately from repulsion, which he thought had no place in Electricity, as a phenomenon similar to the attraction of cohesion, and he imagined, that electrics were brought into contact with the bodies on which they act by their effluvia, excited by friction.

The ingenious Mr. Boyle added to the catalogue of electric substances; but he thought that glass possessed this power in a very low degree: he found, that the Electricity of all bodies, in which it might be excited, was increased by wiping and warming them before they were rubbed; that an excited electric was acted upon by other bodies as strongly as it acted upon them; that diamonds rubbed against any kind of stuff, emitted light in the dark; and that feathers would cling to the fingers, and to other substances, after they had been attracted by electrics. He accounted for electrical attraction, by supposing a glutinous effluvia emitted from electrics, which laid hold of small bodies, in its way, and carried them back to the body from which it proceeded.

Otto Guericke, the celebrated inventor of the air-pump, lived about the same time. This ingenious philosopher discovered, by means of a globe of sulphur, that a body once attracted by an electric, was next repelled, and continued in this state of repulsion till it should be touched by some other body: he also observed the sound and light produced by the excitation of his globe; and that bodies immersed in electrical atmospheres are themselves electrified with an electricity opposite to that of the atmosphere.

The light emitted by electrical bodies was, not long after, observed to much greater advantage by Dr. Wall, who ascribes to light the electrical property which they possess; and he suggests a similarity between the effects of electricity and lightning.

Sir Isaac Newton was not inattentive to this subject: he observed that excited glass attracts light bodies on the side opposite to that on which it is rubbed; and he ascribes the action of electric bodies to an elastic fluid, which freely penetrates glass, and the emission of it to the vibratory motions of the parts of excited bodies.

Mr. Hawksbee wrote on this subject in the year 1709, when a new æra commenced in the history of this science. He first took notice of the great electrical power of glass, and the light proceeding from it; though others had before observed the light proceeding from other electrified substances: he also noted the noise occasioned by it, with a variety of phenomena relating to electrical attraction and repulsion. He first introduced a glass globe into the electrical apparatus, to which circumstance it was that many of his important discoveries were owing.

After his time there was an interval of near 20 years in the progress of this science, till Mr. Stephen Grey established a new æra in the history of Electricity. To him we owe the capital discovery of communicating the power of native electrics to other bodies, in which it cannot be excited, by supporting them on silken lines, hair lines, cakes of resin or glass; and a more accurate distinction than had hitherto obtained between electrics and non-electrics: he also shewed the effect of electricity on water much more obviously than Gilbert had done in the infancy of this science.

The experiments of Mr. Grey were repeated by M. du Fay, member of the Academy of Sciences at Paris, to which he added many new experiments and discoveries of his own. He observed, that electrical operations are obstructed by great heat, as well as by a moist air; that all bodies, both solid and fluid, would receive electricity, when placed on warm or dry glass, or sealing-wax; that those bodies which are naturally the least electric, have the greatest degree of electricity communicated to them by the approach of the excited tube. He transmitted the electric virtue through a distance of 1256 feet; and first observed the electric spark from a living body, suspended on silken lines, and noted several circumstances attending it. M. du Fay also established a principle, first suggested by Otto Guericke, that electric bodies attract all those that are not so, and repel them as soon as they are become electric, by the vicinity or contact of the electric body. He likewise inferred from other experiments, that



that there were two kinds of electricity; one of which he called the *vitreous*, belonging to glass, rock crystal, &c; and the other *resinous*, as that of amber, gumlac, &c, distinguished by their repelling those of the same kind, and attracting each other. He farther observed, that communicated electricity had the same property as the excited; and that electric substances attract the dew more than conductors.

Mr. Grey, resuming his experiments in 1734, suspended several pieces of metal on silken lines, and found that by electrifying them they gave sparks; which was the origin of metallic conductors: and on this occasion he discovered a cone or pencil of electric light, such as is now known to issue from an electrified point. From other experiments he concludes, that the electric power seems to be of the same nature with that of thunder and lightning.

Dr. Desaguliers succeeded Mr. Grey in the prosecution of this science. The account of his first experiments is dated in 1739. To him we owe those technical terms of *conductors* or *non-electrics*, and *electrics per se*; and he first ranked pure air among the electrics per se, and supposed its Electricity to be of the vitreous kind.

After the year 1742, in which Dr. Desaguliers concluded his experiments, the subject was taken up and pursued in Germany: the globe was substituted for the tube, which had been used ever since the time of Hawksbee, and a cushion was soon after used as a rubber, instead of the hand. About this time too, some used cylinders instead of the globes; and some of the German electricians made use of more globes than one at the same time. By thus increasing the electrical power, they were the first who succeeded in setting fire to inflammable substances: this was first done by Dr. Ludolf, in the beginning of the year 1744, who, with sparks excited by the friction of a glass tube, kindled the ethereal spirit of Frobenius. Winkler did the same by a spark from his own finger, by which he kindled French brandy, and other spirits, after previously heating them. Mr. Gralath fired the smoke of a candle just blown out, and so lighted it again; and Mr. Boze fired gun-powder, by means of its inflammable vapour. About this time Ludolf the younger demonstrated, that the luminous barometer was made perfectly electrical by the motion of the quicksilver. The electrical star and electrical bells were also of German invention.

In England Dr. Watson made a distinguished figure from this period in the history of Electricity: he fired a variety of substances by the electrical spark, and first discovered that they are capable of being fired by the repulsive power of Electricity. In the year 1745, the accumulation of the electrical power in glass, by means of the Leyden phial, was first discovered. See *LEYDEN phial*: and for the method practised about this time, of measuring the distance to which the electrical shock may be conveyed, see *Electrical Circuit*. Dr. Watson discovered that the glass tubes and globes do not contain the electric matter in themselves, but only serve as *first-movers* or *determiners*, as he expresses it, of that power; which was also confirmed towards the end of 1746, by Mr. Benjamin Wilson, who made the same discovery, that the electric fluid does not come

from the globe, but from the earth, and other non-electric bodies about the apparatus. Dr. Watson also discovered what Dr. Franklin observed about the same time in America, and called the *plus* and *minus* in Electricity. He likewise shewed that the electric matter passed through the substance of the metal of communication, and not merely over the surface. The history of medical Electricity commenced in the year 1747. We must omit other experiments, and conclusions drawn from them, by Mr. Wilson, Mr. Smeaton, and Dr. Miles in England, and by the Abbé Nollet, with regard to the effect of Electricity on the evaporation of fluids, on solids, and on animal and other organized bodies, in France.

Whilst the philosophers of Europe were busily employed in electrical experiments and pursuits, those of America, and Dr. Franklin in particular, were equally industrious, and no less successful. His discoveries and observations in Electricity were communicated in several letters to a friend; the first of which is dated in 1747, and the last in 1754; and the particulars of his system may be seen under the articles, *Theory of ELECTRICITY, LEYDEN Phial, POINTS, CHARGING, CONDUCTORS, ELECTRICS, &c.*

The similarity between Electricity and Lightning had been suggested by several writers: Dr. Franklin first proposed a method of bringing the matter to the test of experiment, by raising an electrical kite; and he succeeded in collecting electrical fire by this means from the clouds, in 1752, one month after the same theory had been verified in France, and without knowing what had been done there: and to him we owe the practical application of this discovery, in securing buildings from the damage of lightning, by erecting metallic conductors. See *CONDUCTORS, and LIGHTNING*.

In the subsequent period of the history of this science, Mr. Canton in England, and Signior Beccaria in Italy, acquired distinguished reputation. They both discovered, independently of each other, that air is capable of receiving Electricity by communication, and of retaining it when received. Mr. Canton also, towards the latter end of the year 1753, pursued a series of experiments, which prove that the appearances of positive and negative Electricity, which had hitherto been deemed essential and unchangeable properties of different substances, as of glass and sealing-wax for instance, depend upon the surface of the electrics, and that of the rubber.

This hypothesis, verified by numerous experiments, occasioned a controversy between Mr. Canton, and Mr. Delaval, who still maintained that these different powers depended entirely on the substances themselves. About this time too, some curious experiments were performed by four of the principal electricians of that period, viz. Dr. Franklin, and Messrs Canton, Wilcke, and Æpinus, to ascertain the nature of electric atmospheres; the result of which see under that article.

The theory of two electric fluids, always co-existent and counteracting each other, though not absolutely independent, was maintained by a course of experiments on silk stockings of different colours, communicated to the Royal Society by Mr. Symmer, in the year 1759, which were farther pursued by Mr. Cigna  
of.



of Turin, who published an account of them in the Memoirs of the Academy at Turin for the year 1765.

Many instances occur in the history of the science about this period, of the astonishing force of the electric shock, in melting wires, and producing other similar effects: but the most remarkable is an experiment of S. Beccaria, in which he thus revived metals. Several experiments were also made by Dr. Watson, Mr. Smeaton, Mr. Canton, and others, on the passage of the electric fluid through a vacuum, and its luminous appearance, and on the power possessed by certain substances of retaining the light communicated to them by an electric explosion. Mr. Canton, S. Beccaria, and others, made many experiments to identify Electricity and lightning, to ascertain the state of the atmosphere at different times, and to explain the various phenomena of the Aurora Borealis, Water-Spouts, Hurricanes, &c, on the principles of this science.

Those who are desirous of farther information with respect to the history of electrical experiments and discoveries, may consult Dr. Priestley's History and Present State of Electricity. This author however is not merely an historian: his work contains many original experiments and discoveries made by himself. He ascertained the conducting power of charcoal, and of hot glass; the Electricity of fixed and inflammable air, and of oil; the difference between new and old glass, with respect to the diffusion of Electricity over its surface; the lateral explosion in electrical discharges; a new method of fixing circular-coloured spots on the surfaces of metals, and the most probable difference between electrics and conductors, &c. The science is also greatly indebted to many other persons, either for their experiments and improvements of it, or for treatises and other writings upon it; as Mr. Henley, to whom we owe several curious experiments and observations on the electrical and conducting quality of different substances, as chocolate, vapour, &c, with the reason of the difference between them; the fusion of platina; the nature of the electric fluid, and its course in a discharge; the method of estimating the quantity of it in electrical bodies by an electrometer; the influence of points; &c, &c. Also Messrs Van Marum, Van Swinden, Ferguson, Cavallo, Lord Mahon, Nairne, &c, &c, for their several treatises on the subject of Electricity, any of which may be consulted with advantage for the experiments and principles of the science.

*Medical ELECTRICITY.* It is natural to imagine that a power of such efficacy as that of electricity would be applied to medical purposes; especially, since it has been found invariably to increase the sensible perspiration, to quicken the circulation of the blood, and to promote the glandular secretion: accordingly, many instances occur in the latter period of the history of this science, in which it has been applied with considerable advantage and success. And among the variety of cases in which it has been tried, there are none in which it has been found prejudicial except those of pregnancy and the venereal disease. In most disorders, in which it has been used with perseverance, it has given at least a temporary and partial relief, and in many it has effected a total cure. Of which numerous instances may be seen in the Philos. Transf. and the writ-

ings on this science by Messrs Lovet, Westley, Ferguson, Cavallo, &c. &c.

*Theory of ELECTRICITY.* It is hardly necessary to recite the ancient hypotheses on this subject; such as that of the sympathetic powder of the Peripatetics; that of unctuous effluvia emitted by excited bodies, and returning to them again, adopted by Gilbert, Gassendus, Sir Kenelm Digby, &c; or that of the Cartesians, who ascribed electricity to the globules of the first elements, discharged through the pores of the rubbed substance, and in their return carrying with them those light bodies, in whose pores they were entangled: these hypotheses were framed in the infancy of the science, and of philosophy in general, and have long since been exploded. In the more advanced state of electricity there have been two principal theories, each of which has had its advocates. The one, is that of two distinct electric fluids, repulsive with respect to themselves, and attractive of one another, adopted by M. du Fay, on discovering the two opposite species of electricity, viz, the vitreous and resinous, and since new-modelled by Mr. Symmer. It is supposed that these two fluids are equally attracted by all bodies, and exist in intimate union in their pores; and that in this state they exhibit no mark of their existence. But that the friction of an electric by a rubber separates these fluids, and causes the vitreous electricity of the rubber to pass to the electric, and then to the prime conductor of a machine, while the resinous electricity of the conductor and electric is conveyed to the rubber: and thus the quality of the electric fluid, possessed by the conductor and the rubber, is changed, while the quantity remains the same in each. In this state of separation, the two electric fluids will exert their respective powers; and any number of bodies charged with either of them will repel each other, attract those bodies that have less of each particular fluid than themselves, and be still more attracted by bodies that are wholly destitute of it, or that are loaded with the contrary. According to this theory, the electric spark makes a double current; one fluid passing to an electrified conductor from any substance presented to it, whilst the same quantity of the other fluid passes from it; and when each body receives its natural quantity of both fluids, the balance of the two powers is restored, and both bodies are unelectrified. For a further account of the explication of some of the principal phenomena of electricity by this theory, see Dr. Priestley's History, vol. 2, § 3.

The other theory is commonly distinguished under the denomination of *positive and negative electricity*, being first suggested by Dr. Watson, but digested, illustrated, and confirmed by Dr. Franklin; and since that it has been known by the appellation of the Franklinian hypothesis. It is here supposed that all the phenomena of electricity depend on one fluid, *sui generis*, extremely subtle and elastic, dispersed through the pores of all bodies, by which the particles of it are as strongly attracted as they are repelled by one another. When bodies possess their natural share of this fluid, or such a quantity as they can retain by their non-attraction, it is then said they are in an unelectrified state; but when the equilibrium is disturbed, and they either acquire an additional quantity *from* other bodies, or lose part of their own natural share by communication *to* other bodies, they



they exhibit electrical appearances. In the former case it is said they are electrified positively, or *plus*; and in the other negatively, or *minus*. This electric fluid, it is supposed, moves with great ease in those bodies that are called conductors, but with extreme difficulty and slowness in the pores of electrics; whence it comes to pass, that all electrics are impermeable to it. It is farther supposed that electrics contain always an equal quantity of this fluid, so that there can be no surcharge or increase on one side without a proportionable decrease or loss on the other, and vice versa; and as the electric does not admit the passage of the fluid through its pores, there will be an accumulation on one side, and a corresponding deficiency on the other. Then when both sides are connected together by proper conductors, the equilibrium will be restored by the rushing of the redundant fluid from the overcharged surface to the exhausted one. Thus also, if an electric be rubbed by a conducting substance, the electricity is only conveyed from one to the other, the one giving what the other receives; and if one be electrified positively, the other will be electrified negatively, unless the loss be supplied by other bodies connected with it, as in the case of the electric and insulated rubber of a machine. This theory serves likewise to illustrate the other phenomena and operations in the science of electricity. Thus, bodies differently electrified will naturally attract each other, till they mutually give and receive an equal quantity of the electric fluid, and the equilibrium is restored between them. Beccaria supposes, that this effect is produced by the electric matter making a vacuum in its passage, and the contiguous air afterwards collapsing, and so pushing the bodies together.

The influence of points, in drawing or throwing off the electric fluid, depends on the less resistance it finds to enter or pass off through fewer particles than through a greater number, whose resistance is united in flat or round surfaces. The electric light is supposed to be part of the electric fluid, which appears when it is properly agitated; and the sound of an explosion is produced by vibrations, occasioned by the air's being displaced by the electric fluid, and again suddenly collapsing.

As to the nature of the electric fluid, philosophers have entertained very different sentiments: some, and among them Mr. Wilson, have supposed that it is the same with the ether of Sir Isaac Newton, to which the phenomena of attraction and repulsion are ascribed; whilst the light, smell, and other sensible qualities of the electric fluid, are referred to the grosser particles of bodies, driven from them by the forcible action of this ether; and other appearances are explained by means of a subtile medium diffused over the surfaces of all bodies, and resisting the entrance and exit of the ether; which medium, it is supposed, is the same with the electric fluid, and is more rare on the surfaces of conductors, and more dense and resisting on those of electrics: but Dr. Priestley remarks that, though they may possess some common properties, they have others essentially distinct; the ether is repelled by all other matter, whereas the electric fluid is strongly attracted by it. Others have had recourse to the element of fire; and from the supposed identity of fire and the electric fluid, as well as from the similarity of some of their effects, the

latter has been usually called the electric fire: but most electricians have supposed that it is a fluid *sui generis*. Mr. Cavendish has published an attempt to deduce and explain some of the principal phenomena of electricity in a mathematical and systematic manner, from the nature of this fluid, considered as composed of particles that repel each other, and attract the particles of all other matter, with a force inversely as some less power of the distance than the cube, whilst the particles of all other matter repel each other, and attract those of the electric fluid, according to the same law. *Philos. Trans.* vol. 61, pa. 584—677. And a similar hypothesis and method of reasoning was also proposed by M. *Æpinus*, in his *Tentamen Theoriæ Electricitatis & Magnetismi*.

Dr. Priestley concludes, from experiments, that the electric matter either is phlogiston, or contains it, since he found that both produced similar effects. Mr. Henley also apprehends, that the electric fluid is a modification of that element, which, in its quiescent state, is called phlogiston; in its first active state, electricity; and when violently agitated, fire. Perhaps we may be allowed to enlarge our views, and consider the sun as the fountain of the electric fluid, and the zodiacal light, the tails of comets, the aurora borealis, lightning, and artificial electricity, as its various and not very dissimilar modifications. On this subject, see Priestley's *Hist. of Electr.* vol. 2, part 3, § 1, 2, 3; Wilson's *Essay towards an Explication of the Phenomena of Electricity*, &c; Wilson and Hoadley's *Obs.* &c, pa. 55, 1759; Freke's *Essay on the Cause of Electricity*, 1746; Priestley on *Air*, vol. 1, pa. 186, 274, &c; *Philos. Trans.* vol. 67, pa. 129; and Mr. *Eeles's* *Letters*, on the same subject.

ELECTROMETER, is an instrument that measures the quantity, and determines the quality of electricity, in any electrified body. Previous to the invention of instruments of this kind, Mr. Canton estimated the quantity of electricity in a charged phial, by presenting the phial with one hand to an insulated conductor, and giving it a spark, which he took off with the other; proceeding in this manner till the phial was discharged, when he determined the height of the charge by the number of sparks. Electrometers are of 4 kinds: 1, the single thread; 2, the cork or pith balls; 3, the quadrant; and 4, the discharging Electrometer.

The 1st, or most simple Electrometer, is a linen thread, called by Dr. Desaguliers, the *thread of trial*; which, being brought near an electrified body, is attracted by it: but this does little more than determine whether the body is in any degree electrified or not; without determining with any precision its quantity, much less the quality of it. The Abbé Nollet used two threads, shewing the degree of electricity by the angle of their divergency exhibited in their shadow on a board placed behind them.

Mr. Canton's Electrometer consisted of two balls of cork, or pith of elder, about the size of a small pea, suspended by fine linen threads, about 6 inches long, which may be wetted in a weak solution of salt. See fig. 7. If the box containing these balls be insulated, by placing it on a drinking glass, &c, and an excited smooth glass tube be brought near them, they will first be attracted by it, and then be repelled both from the glass,



glass, and from each other; but on the approach of excited wax, they will gradually approach and come together; and vice versa. This apparatus will also serve to determine the electricity of the clouds and air, by holding them at a sufficient distance from buildings, trees, &c; for if the electricity of the clouds or air be positive, their mutual repulsion will increase by the approach of excited glass, or decrease by the approach of amber or sealing-wax; on the contrary, if it be negative, their repulsion will be diminished by the former, and increased by the latter. See *Philos. Trans.* vol. 48, part 1 and 2, for an account of Mr. Canton's curious experiments with this apparatus.

If two balls of this kind be annexed to a prime conductor, they will serve to determine both the degree and quality of its electrification, by their mutual repulsion and divergency.

The Discharging Electrometer, fig. 3, plate ix, was invented by Mr. Lane. It consists of brass work G, the lower part of which is inclosed in the pillar F, made of baked wood, and boiled in linseed oil, and bored cylindrically about two-thirds of its length; the brass work is fixed to the pillar by the screw H, moveable in the groove I; and through the same is made to pass a steel screw L, to the end of which, and opposite to K, a polished hemispherical piece of brass, attached to the prime conductor, is fixed a ball of brass M well polished. To this screw is annexed a circular plate O, divided into 12 equal parts. The use of this Electrometer is to discharge a jar D, or any battery connected with the conductor, without a discharging rod, and to give shocks successively of the same degree of strength; on which account it is very fit for medical purposes. Then, if a person holds a wire fastened to the screw H in one hand, and another wire fixed to E, a loop of brass wire passing from the frame of the machine to a tin plate, on which the phial D stands, he will perceive no shock, when K and M are in contact; and the degree of the explosion, as well as the quantity of electricity accumulated in the phial, will be regulated by the distance between K and M. *Philos. Trans.* vol. 57, pa. 451.—Mr. Henley much improved Mr. Lane's Electrometer, by taking away the screw, the double milled nut, and the sharp-edged graduated plate, and adding other contrivances in their stead. Mr. Henley's discharger of this kind has two tubes, one sliding within the other, to lengthen and accommodate it to larger apparatus.

The Quadrant Electrometer of Mr. Henley, consists of a stem, terminating at its lower end with a brass ferrule and screw, for fastening it upon any occasion; and its upper part ends in a ball. Near the top is fixed a graduated semicircle of ivory, on the centre of which the index, being a very light rod with a cork ball at its extremity, reaching to the brass ferrule of the stem, is made to turn on a pin in the brass piece, so as to keep near the graduated limb of the semicircle. When the Electrometer is not electrified, the index hangs parallel to the stem; but as soon as it begins to be electrified, the index, repelled by the stem, will begin to move along the graduated edge of the semicircle, and so mark the degree to which the conductor is electrified, or the height to which the charge of any jar or battery is advanced.

Mr. Cavallo has also contrived several ingenious Elec-

trometers, for different uses; as may be seen in his *Treatise on Electricity*, pa. 370, &c, and in the *Philos. Trans.* vol. 67, pa. 48 and 399.

ELECTROPHOR, or ELECTROPHORUS, an instrument for shewing perpetual electricity; which was invented by Mr. Volta, of Como, near Milan, in Italy. The machine consists of two plates, fig. 8, one of which B is a circular plate of glass, covered on one side with some resinous electric, and the other A is a plate of brass, or a circular board, coated with tinfoil, and furnished with a glass handle I, which may be screwed into its centre by means of a socket. If the plate B be excited by rubbing it with new white flannel, and the plate A be applied to its coated side, a finger, or any other conductor, will receive a spark on touching this plate; and if the plate A be then separated, by means of the handle I, it will be found strongly electrified, with an electricity contrary to that of the plate B. By replacing the plate A, touching it with the finger, and separating it again, it will be found electrified as before, and give a spark to any conductor, attended with a snapping noise; and by this means a coated phial may be charged. The same phenomena may repeatedly be exhibited, without any renewed excitation of the electric plate B; the electric power of B having continued for several days, and even weeks, after excitation; though there is no reason to imagine that it is perpetual.

Mr. Cavallo prepares this machine by coating the glass plate with sealing-wax; and Mr. Adams, philosophical instrument maker, prepares them with plates formed from a composition of two parts of shell-lac, and one of Venice turpentine, without any glass plate.

The action of this plate depends on a principle discovered and illustrated by the experiments of Franklin, Canton, Wilcke, and Æpinus, viz, that an excited electric repels the electricity of another body, brought within its sphere of action, and gives it a contrary electricity. Thus the plate A, touched by a conductor, whilst in contact with the plate B, electrified negatively, will acquire an additional quantity of the electric fluid from the conductor; but if it were in contact with a plate electrified positively, it would part with its electricity to the conductor connected with it. See an account of several curious experiments with this machine, by Mr. Henley, Mr. Cavallo, and Dr. Ingenhousz, in the *Philos. Trans.* vol. 66, pa. 513; vol. 67, pa. 116 and 389; and vol. 68, pa. 1027 and 1049.

ELEMENTARY, something that relates to the principles or elements of bodies, or sciences; as Elementary Air, Fire, Geometry, Music, &c.

ELEMENTS, the first principles, of which all bodies and things are composed. These are supposed few in number, unchangeable, and by their combinations producing that extensive variety of objects to be met with in the works of nature.

Democritus stands at the head of the Elementary Philosophers, in which he is followed by Epicurus, and many others after them, of the Epicurean and corpuscular philosophers.

Among those who hold the Elements corruptible, some will have only one, and some several. Of the former, the principal are Heraclitus, who held fire; Anaximenes, air; Thales Milesius, water; and Hesiod, earth;



earth; as the only Element. Hesiod is followed by Bernardin, Telefius; and Thales by many of the chemists.

Among those who admit several corruptible Elements, the principal are the Peripatetics; who, after their leader Aristotle, contend for four Elements, viz, fire, air, water, and earth. Aristotle took the notion from Hippocrates; Hippocrates from Pythagoras; and Pythagoras from Ocellus Lucanus, who it seems was the first author of it.

The Cartesians admit only three Elements, fire, air, and earth. See *CARTESIAN Philosophy*.

Newton observes, that it seems probable that God, in the beginning, formed matter in solid, massive, hard, impenetrable, moveable particles, of such sizes and figures, &c, as most conduced to the end for which he formed them; and that these primitive particles, being solids, are incomparably harder than any porous body compounded of them; even so hard as never to wear out; no ordinary power being able to divide what God made one in the first creation. While the particles remain entire, they may compose bodies of one and the same nature and texture in all ages; but should they wear away, or break in pieces, the nature of things, depending on them, would be changed; water and earth, composed of old worn particles, and fragments of particles, would not be of the same nature and texture now, with water and earth composed of entire particles in the beginning. And therefore, that things may be lasting, the changes of corporeal things are to be placed only in the various separations, and new associations and motions of those permanent particles; compound bodies being apt to break, not in the midst of solid particles, but where those particles are laid together, and only touch in a few points. It seems to him likewise, that these particles have not only a vis inertie, with the passive laws of motion thence resulting, but are also moved by certain active principles; such as gravity, and the cause of fermentation, and the cohesion of bodies.

ELEMENTS, a term also used for the first grounds and principles of arts and sciences; as the Elements of geometry, Elements of mathematics, &c. So Euclid's Elements, or simply the Elements, as they were anciently and peculiarly named, denotes the treatise on the chief properties of geometrical figures by that author.

The ELEMENTS of Mathematics have been delivered by several authors in their courses, systems, &c. The first work of this kind is that of Herigon, in Latin and French, and published in 1664, in 10 tomes; which contains Euclid's Elements and Data, Apollonius, Theodosius, &c; with the modern Elements of arithmetic, algebra, trigonometry, architecture, geography, navigation, optics, spherics, astronomy, music, perspective, &c. The work is remarkable for this, that a kind of real and universal characters are used throughout; so that the demonstrations may be understood by such as only remember the characters, without any dependence on language or words at all.

Since Herigon, the Elements of the several parts of mathematics have been also delivered by others; particularly the Jesuit Schottus, in his *Cursus Mathematicus*, in 1674; De Chales, in his *Cursus*, 1674; Sir Jonas Moore, in his *New System of Mathematics*, in

1681; Ozanam, in his *Cours de Mathématique*, in 1699; Jones, in his *Synopsis Palmariorum Matheseos*, in 1706; and many others, but above all, Christ. Wolffius, or Wolf, in his *Elementa Matheseos Universæ*, in 2 vols 4to, the 1st published in 1713, and the 2d in 1715; a very excellent work of the kind. Another edition of the work was published at Geneva, in 5 vols 4to, of the several dates 1732, 1733, 1735, 1738, and 1741.

The ELEMENTS of Euclid, as they were the first, so they continue still the best system of geometry, are in 15 books. There have been numerous editions and commentaries of this work. Proclus wrote a commentary on it. Orontius Fineus first gave a printed edition of the first 6 books, in 1530, with notes, to explain Euclid's sense. Peletarius did the same in 1557. Nic. Tartaglia, about the same time, made a comment on all the 15 books, with the addition of many things of his own. And the same was also done by Billingsley in 1570; and by Fluffates Candalla, a noble Frenchman, in the year 1578, with considerable additions as to the comparison and inscriptions of solid bodies; which work was afterwards republished with a prolix commentary, by Clavius. Commandine gave also a good edition of it. In 1703, Dr. Gregory published an edition of the whole works of Euclid, in Greek and Latin, including his Elements. But it would be endless to relate all the other editions of these Elements, either the whole, or in part, that have been given; some of the best of which are those of De Chales, Tacquet, Ozanam, Whiston, Stone, and most especially that of Dr. Rob. Simson, of Glasgow.

Other writers on the Elements of Geometry are almost out of number, in all nations.

ELEMENTS, in the Higher or Sublime Geometry, are the infinitely small parts, or differentials, of a right line, curve, surface, or solid.

ELEMENTS, in Astronomy, are those principles deduced from astronomical observations and calculations, and those fundamental numbers, which are employed in the construction of tables of the planetary motions. Thus, the Elements of the theory of the sun, or rather of the earth, are his mean motion and eccentricity, with the motion of the aphelia. And the Elements of the theory of the moon, are her mean motion, that of the node and apogee, the eccentricity, the inclination of her orbit to the plane of the ecliptic; &c.

ELEVATION, the height or altitude of any thing.

ELEVATION, in Architecture, denotes a draught or description of the principal face or side of a building; called also its upright or orthography.

ELEVATION, in Astronomy and Geography, is various; as Elevation of the equator, of the pole, of a star, &c.

ELEVATION of the Equator, is the height of the equator above the horizon; or an arc of the meridian intercepted between the equator and the horizon of the place.—The Elevation of the equator and of the pole together always make up a quadrant; the one being the complement of the other. Therefore, the Elevation of the pole being found, and subtracted from 90°, leaves the elevation of the equator.

ELEVATION of the Pole, is its height above the horizon;



horizon; or an arc of the meridian comprehended between the equator and the horizon of the place.

The elevation of the pole is always equal to the latitude of the place; that is, the arc of the meridian intercepted between the pole and the horizon, is every where equal to the arc of the same meridian intercepted between the equator and the zenith. Thus the north pole is elevated  $51^{\circ} 31'$  above the horizon of London; and the distance, or number of degrees, is the same between London and the equator; so that London is also in  $51^{\circ} 31'$  of north latitude.

**ELEVATION of a Star**, or of any other point in the sphere, is the angular height above the horizon; or an arc of the vertical circle intercepted between the star and the horizon. The meridian altitude of any such point, or its altitude when in the meridian, is the greatest of all.

**ELEVATION of a Cannon, or Mortar**, is the angle which the bore or the axis of the piece makes with the horizontal plane.

**Angle of ELEVATION**, is the angle which any line of direction makes above a horizontal line.

**ELEVATION** is also used by some writers on Perspective, for the scenography, or perspective representation of the whole body or building.

**ELLIPSE**, or **ELLIPSIS**, is one of the conic sections, popularly called an oval; being called an Ellipse or Ellipsis by Apollonius, the first and principal author on the conic sections, because in this figure the squares of the ordinates are *less* than, or *defective* of, the rectangles under the parameters and abscisses.

This figure is differently defined by different authors; either from some of its properties, or from mechanical construction, or from the section of a cone, which is the best and most natural way. Thus;

Fig. 1.





Fig. 2.



1. An Ellipse is a plane figure made by cutting a cone by a plane obliquely through the opposite sides of it; or so as that the plane makes a less angle with the base than the side of the cone makes with it; as ABD, fig. 1.

The line AB connecting the uppermost and lowest points of the section, is the transverse axis; the middle of it C, is the centre; and the perpendicular to it DCE, through the centre, is the conjugate axis. The parameter, or latus rectum, is a 3d proportional to the transverse and conjugate axes; and the foci are two points in the transverse axis, at such equal distances from the centre, that the double ordinates passing through those points, and perpendicular to the transverse, are equal to the parameter.

2. The Ellipse is also variously described from some of its properties. As first, That it is a figure of such a

nature that if two lines be drawn from two certain points C and D in the axis, fig. 2, to any point E in the circumference, the sum of those two lines CE and DE will be every where equal to the same constant quantity, viz, the axis AB. Or, secondly, that it is

Fig. 3.





Fig. 4.



a figure of such a nature, that the rectangle  $AG \times GB$  (fig. 3) of the abscisses, tending contrary ways, is to  $GH^2$  the square of the ordinate, as  $AB^2$  to  $IK^2$ , the square of the transverse axis to the square of the conjugate, or, which is the same thing, as the transverse axis is to the parameter. And so of other properties.

3. Or the Ellipse is also variously described from its mechanical constructions, which also depend on some of its chief properties. Thus; 1st, If in the axis AB, there be taken any point I (fig. 4); and if with the radii AI, BI, and centres F and f, the two foci, arcs be described, these arcs will intersect in certain points E, E, e, e, which will be in the curve or circumference of the figure: and thus several points I being taken in the axis AB, as many more points E, e, &c, will be found; then the curve line drawn through all these points E, e, will be an Ellipse. Or, thus; if there be taken a thread of the exact length of the transverse axis AB, and the ends of the thread be fixed by pins in the two foci F and f; (fig. 5) then moving a pen or pencil within the thread, so as to keep it always stretched out, it will describe the curve called an Ellipse.

Fig. 5.





Fig. 6.



**To Construct an ELLIPSE.** There are many other ways of describing or constructing an Ellipse, besides those just now given: as

1st. If upon the given transverse axis there be described a circle AGB (fig. 6), to which draw any ordinate DG, and DE a 4th proportional to the transverse, the conjugate, and the ordinate DG; then E is a point in the curve. Or if the circle *agb* be described on the conjugate axis *ab*, to which any ordinate *dg* is drawn, in which taking *dE* in like manner a 4th proportional to the conjugate, the transverse, and ordinate *dg*, then shall



shall E be in the curve. Or, having described the two circles, and drawn the common radius CgG cutting them in G and g; then dgE drawn parallel to the transverse, and DGE parallel to the conjugate, the intersection E of these two lines will be in the curve of the ellipse. And thus several points E being found, the curve may be drawn through them all with a steady hand.

Fig. 7.





Fig. 8.



2. If there be provided three rulers, of which the two GH and FI (fig. 7) are of the length of the transverse axis LK, and the third FG equal to HI the distance between the foci; then connecting these rulers so as to be moveable about the foci H and I, and about the points F and G, their intersection E will always be in the curve of the ellipse; so that by moving the rulers about the joints, with a pencil passed through the slits made in them, it will trace out the Ellipse.

Fig. 9.





Fig. 10.



3. If one end A of any two equal rulers AB, DB, (fig. 9 and 10) which are moveable about the point B, like a carpenter's joint-rule, be fastened to the ruler LK, so as to be moveable about the point A; and if the end D of the ruler DB be drawn along the side of the ruler LK; then any point E, taken in the side of the ruler DB, will describe an ellipse, whose centre is A, conjugate axis = 2DE, and transverse = 2AB + 2BE.

Another method of description is by the Elliptical Compass. See that article, below.

*Some of the more Remarkable Properties of the Ellipse.*

—1. The rectangles under the abscissas are proportional to the squares of their ordinates; or as the square of any axis, or any diameter, is to the square of its conjugate, so is the rectangle under two abscissas of the former, to the square of their ordinate parallel to the latter; or again, as any diameter is to its parameter, so is the said rectangle under two abscissas of that diameter, to the square of their ordinate. So that if  $d$  be

any diameter,  $c$  its conjugate,  $p$  its parameter =  $\frac{c^2}{d}$ ,  $x$  the one abscissa,  $d-x$  the other, and  $y$  the ordinate; then,


as  $d^2 : c^2 :: x \cdot \overline{d-x} : y^2$ , or  $d^2 y^2 = c^2 x \cdot \overline{d-x}$ ;

or as  $d : p :: x \cdot \overline{d-x} : y^2$ , or  $dy^2 = px \cdot \overline{d-x}$ .

From either of which equations, called the equation of the curve, any one of the quantities may be found, when the other three are given.

2. The sum of two lines drawn from the foci to meet in any point of the curve, is always equal to the transverse axis; that is,  $CE + DE = AB$ , in the 2d fig. Consequently the line CG drawn from the focus to the end of the conjugate axis, is equal to AI the semi-transverse.

Fig. 11.



3. If from any point of the curve, there be an ordinate to either axis, and also a tangent meeting the axis produced; then half that axis will be a mean proportional between the distances from the centre to the two points of intersection; viz, CA a mean proportional between CD and CT. And consequently all the tangents TE, TE, meet in the same point of the axis produced, which are drawn from the extremities E, E, of the common ordinates DE, DE, of all Ellipses described on the same axis AB.

4. Two lines drawn from the foci to any point of the curve, make equal angles with the tangent at that point: that is, the  $\angle FET = \angle fEt$ .

5. All the parallelograms are equal to each other, that are circumscribed about an Ellipsis; and every such parallelogram is equal to the rectangle of the two axes.

6. The sum of the squares of every pair of conjugate diameters, is equal to the same constant quantity, viz, the sum of the squares of the two axes.

7. If a circle be described upon either axis, and from any point in that axis an ordinate be drawn both to the circle and ellipse; then shall the ordinate of the circle be to the ordinate of the Ellipse, as that axis is to the other axis: viz,

$$AB : ab :: DG : DE,$$

and  $ab : AB :: dg : dE$ . (in the 6th fig.)

And in the same proportion is the area of the circle to the area of the Ellipse, or any corresponding segments ADG, ADE. Also the area of the Ellipse is a mean proportional between the areas of the inscribed and circumscribed circles. Hence therefore,

8. *To find the Area of an Ellipse.* Multiply the two axes together, and that product by .7854, for the area. Or

9. *To find the Area of any Segment ADE.* Find the  
3 1 2 area



area of the corresponding segment ADG of a circle on the same diameter AB; then say, as the axis AB : its conj.  $ab ::$  circ. seg. ADG : elliptic seg. ADE.

10. To find the length of the whole circumference of the Ellipse. Multiply the circumference of the circumscribing circle by the sum of the series

$$1 - \frac{d}{2^2} - \frac{3d^2}{2^2 \cdot 4^2} - \frac{3^2 \cdot 5d^3}{2^2 \cdot 4^2 \cdot 6^2} - \frac{3^2 \cdot 5^2 \cdot 7d^4}{2^2 \cdot 4^2 \cdot 6^2 \cdot 8^2} \&c,$$

for the area: where  $d$  is the difference between an unit and the square of the less axis divided by the square of the greater.

Or, for a near approximation, take the circumference of the circle whose diameter is an arithmetical mean between the two axes, or half their sum: that is,

$\frac{t+c}{2} \times 3.1416 =$  the perimeter nearly; being about the 200th part too little; where  $t$  denotes the transverse, and  $c$  the conjugate axis.

Or, again, take the circumference of the circle, the square of whose diameter is half the sum of, or an arithmetical mean between the squares of the two axes: that is,  $\sqrt{\frac{t^2+c^2}{2}} \times 3.1416 =$  the perimeter nearly; being about the 200th part too great.

Hence combining these two approximate rules together, the periphery of the Ellipse will be very nearly equal to half their sum, or equal to

$\frac{3.1416}{2} \times \left( \frac{t+c}{2} + \sqrt{\frac{t^2+c^2}{2}} \right)$ , within about the 30000th part of the truth.

For the length of any particular arc, and many other parts about the Ellipse, see my *Mensuration*, pa. 283, &c, 2d edit. See also my *Conic Sections*, for many other properties of the Ellipse, especially such as are common to the hyperbola also, or to the conic sections in general.

*Infinite ELLIPSES.* See ELLIPTOIDE.

ELLIPSOID, is an elliptical spheroid, being the solid generated by the revolution of an ellipse about either axis. See SPHEROID.

ELLIPTIC or ELLIPTICAL, something relating to an ellipse.

ELLIPTIC *Compasses*, or ELLIPTICAL *Compass*, is an instrument for describing ellipses at one revolution of the index. It consists of a cross ABGH (fig. 8.) with grooves in it, and an index CE, sliding in dove-tail grooves; by which motion the end E describes the curve of an ellipse.

ELLIPTICAL *Conoid*, is sometimes used for the spheroid.

ELLIPTICAL *Dial*, an instrument usually made of brass, with a joint to fold together, and the gnomon to fall flat, for the convenience of the pocket. By this instrument are found the meridian, the hour of the day, the rising and setting of the sun, &c.

ELLIPTOIDE, an infinite or indefinite Ellipsis, defined by the indefinite equation  $ay^{m+n} = bx^m \cdot a - x^n$  when  $m$  or  $n$  are greater than 1: for when they are each = 1, it denotes the common ellipse.

There are several kinds or degrees of Elliptoides, denominated from the exponent  $m+n$  of the ordinate  $y$ .

As the cubical Elliptoide, expressed by  $ay^3 = bx^2 \cdot a - x$ ; the biquadratic, or surfolid  $ay^4 = bx^2 \cdot a - x^2$ ; &c.

ELONGATION, in Astronomy, the distance of a planet from the sun, with respect to the earth; or the angle formed by two lines drawn from the earth, the one to the sun, and the other to the planet; or the arc measuring that angle: Or it is the difference between the sun's place and the geocentric place of the planet.

The Greatest Elongation, is the greatest distance to which the planets recede from the sun, on either side. This is chiefly considered in the inferior planets, Venus and Mercury; the Greatest Elongation of Venus being about 48 degrees, and of Mercury only about 28 degrees; which is the reason that this planet is so rarely seen, being usually lost in the light of the sun.

EMBER-Days, are certain days observed by the church at four different seasons of the year; viz, the Wednesday, Friday, and Saturday next after Quadregesima Sunday, or the 1st Sunday in Lent; after Whitunday; after Holyrood, or Holycross, the 14th day of September; and after St. Lucy, the 13th day of December. The name, it seems, is derived from Embers, or ashes, which it is supposed were strewed on the head, on these solemn fasts.

EMBER-Weeks, are those weeks in which the Ember-days fall. These Ember-weeks are now chiefly noticed on account of the ordination of priests and deacons; because the canon appoints the Sundays next after the Ember-weeks for the solemn times of ordination; though the bishops, if they please, may ordain on any Sunday or holiday.

EMBOLIMÆAN, and EMBOLISMIC, *Intercalary*, is chiefly used in speaking of the additional months inserted by chronologists to form the lunar cycle of 19 years.

The 19 solar years consisting of 6939 days and 18 hours, and the 19 lunar years only making 6726 days, it was found necessary to intercalate or insert 7 lunar months, containing 209 days; which, with the 4 bissextile days happening in the lunar cycle, make 213 days, and the whole 6939 days, the same as the 19 solar years, which make the lunar cycle.

In the course of 19 years there are 228 common moons, and 7 Embolismic moons, which are distributed in this manner, viz, the 3d, 6th, 9th, 11th, 14th, 17th, and 19th years, are Embolismic, and so contain 384 days each. And this was the method of computing time among the Greeks: though they did not keep regularly to it, as it seems the Jews did. And the method of the Greeks was followed by the Romans till the time of Julius Cæsar.

The Embolismic months, like other lunar months, are sometimes of 30 days, and sometimes only 29 days.

The *Embolismic Epacts* are those between 19 and 29; which are so called, because, with the addition of the epact 11, they exceed the number 30: or rather, because the years which have these epacts, are Embolismic; having 13 moons each, the 13th being the Embolismic.

EMBOLISMUS, in Chronology, signifies intercalation. As the Greeks used the lunar year, which contains only 354 days, that they might bring it to the solar



solar year, of 365 days, they had an Embolism every two or three years, when they added a 13th lunar month.

EMBOLUS, the moveable part of a pump or syringe; called also the piston, and popularly the sucker. The pipe or barrel of a syringe, &c, being close shut, the embolus cannot be drawn up without a very considerable force; which force being withdrawn, the embolus returns again with violence; owing to the greater pressure of air above than below it.

EMBRASURE, in Architecture, an enlargement of the aperture or opening of a door, or window, within side the wall, sloping back inwards, to give the greater play for the opening of the door, casement, &c, or to take in the more light.

EMBRASURES, in Fortification, are the apertures or holes through which the cannon are pointed, whether in casemates, batteries, or in the parapets of walls. In the navy, these are called port-holes. The Embrasures are placed 12 or 15 feet apart from each other; being made sloping or opening outwards, from 6 to 9 feet wide on the outside of the wall, and from 2 to 3 within, to allow the gun to traverse from side to side. Their base is about  $2\frac{1}{2}$  or 3 feet above the platform on the inside of the wall, but sloping down outwards, so as to be only about  $1\frac{1}{2}$  above it on the outside; in order that the muzzle on occasion may be depressed, and so the gun shoot low, or downwards.

EMERGENT *Year*, in Chronology, is the epoch, or date, from whence any people begin to compute their time or dates. So, our Emergent year is sometimes the year of the creation, but more usually the year of the birth of Christ. The Jews used that of the Deluge, or the Exodus, &c. The Emergent year of the Greeks, was the beginning of the Olympic games; while that of the Romans was the date of the building of their city.

EMERSION, in Astronomy, is the re-appearance of the sun, moon, or other planet, after having been eclipsed, or hid by the interposition of the moon, earth, or other body.

The Emersions and immerisions of Jupiter's first satellite, are particularly useful for finding the longitudes of places; the immerisions being observed from Jupiter's conjunction with the sun, till his opposition; and the Emersions from the opposition till the conjunction. But within 15 days of the conjunction, both before and after it, they cannot be observed, because the planet and his satellites are then lost in the sun's light.

EMERSION is also used when a star, after being hid by the sun, begins to re-appear, and to get out of his rays.

*Minutes or Scruples of EMERSION*, an arc of the moon's orbit, which her centre passes over, from the time she begins to emerge out of the earth's shadow, to the end of the eclipse.

EMERSION, in Physics, the rising of any solid above the surface of a fluid that is specifically heavier than the solid, into which it had been violently immersed, or pushed.

It is one of the known laws of hydrostatics, that a lighter solid, being forced down into a heavier fluid, immediately endeavours to emerge; and that with a force equal to the excess of the weight of a quantity of

the fluid above that of an equal bulk of the solid. Thus, if the body be immersed in a fluid of double its specific gravity, it will emerge again till half its bulk be above the surface of the fluid.

EMERSON (WILLIAM), a late eminent mathematician, was born in June 1701, at Hurworth, a village about three miles south of Darlington, on the borders of the county of Durham; at least it is certain that he resided here from his childhood. His father Dudley Emerson taught a school, and was a tolerable proficient in the mathematics; and without his books and instructions, perhaps his son's genius, though eminently fitted for mathematical studies, might never have been unfolded. Beside his father's instructions, our author was assisted in the learned languages by a young clergyman, then curate of Hurworth, who was boarded at his father's house. In the early part of his life he attempted to teach a few scholars: but whether from his concise method (for he was not happy in explaining his ideas), or the warmth of his natural temper, he made no progress in his school; he therefore soon left it off; and satisfied with a moderate competence left him by his parents, he devoted himself to a studious retirement, which he thus closely pursued, in the same place, through the course of a long life, being mostly very healthy, till towards the latter part of his days, when he was much afflicted with the stone. Toward the close of the year 1781, being sensible of his approaching dissolution, he disposed of the whole of his mathematical library to a bookseller at York; and on May the 20th, 1782, his lingering and painful disorder put an end to his life at his native village, being near 81 years of age.

Mr. Emerson, in his person, was rather short, but strong and well made, with an open countenance and ruddy complexion, being of a healthy and hardy disposition. He was very singular in his behaviour, dress, and conversation. His manner and appearance were that of a rude and rather boorish country man; he was of very plain conversation, and indeed seemingly rude, commonly mixing oaths in his sentences, though without any ill intention. He had strong good natural mental parts, and could discourse sensibly on any subject, but was always positive and impatient of any contradiction. He spent his whole life in close study, and writing books, from the profits of which, he redeemed his little patrimony from some original incumbrance. In his dress he was as singular as in every thing else. He possessed commonly but one suit of clothes at a time, and those very old in their appearance. He seldom used a waistcoat; and his coat he wore open before, except the lower button; and his shirt quite the reverse of one in common use, the hind-side turned foremost, to cover his breast, and buttoned close at the collar behind. He wore a kind of rusty coloured wig, without a crooked hair in it, which probably had never been tortured with a comb from the time of its being made. A hat he would make to last him the best part of a life time; gradually lessening the flaps, bit by bit, as it lost its elasticity and hung down, till little or nothing but the crown remained.

He often walked up to London when he had any book to be published, revising sheet by sheet himself:—trusting no eye but his own was always a favourite

maxim



maxim with him. In mechanical subjects, he always tried the propositions practically, making all the different parts himself on a small scale; so that his house was filled with all kinds of mechanical instruments, together or disjointed. He would frequently stand up to his middle in water while fishing; a diversion he was remarkably fond of. He used to study incessantly for some time, and then for relaxation take a ramble to any pot-alehouse where he could get any body to drink with and talk to. The late Mr. Montagu was very kind to Mr. Emerson, and often visited him, being pleased with his conversation, and used often to come to him in the fields where he was working, and accompany him home, but could never persuade him to get into a carriage: on these occasions he would sometimes exclaim, "Damn your whim-wham! I had rather walk." He was a married man, and his wife used to spin on an old-fashioned wheel, of his own making, a drawing of which is given in his *Mechanics*.

Mr. Emerson, from his strong vigorous mind and close application, had acquired a deep knowledge of all the branches of mathematics and physics, upon all parts of which he wrote good treatises, though in a rough and unpolished style and manner. He was not remarkable however for genius or discoveries of his own, as his works shew hardly any traces of original invention. He was well skilled in the science of music, the theory of sounds, and the various scales both ancient and modern; but he was a very poor performer, though he could make and repair some instruments, and sometimes went about the country tuning harpsichords.

The following is a list of Mr. Emerson's works; all of them printed in 8vo, excepting his *Mechanics* and his *Increments* in 4to, and his *Navigation* in 12mo. 1. The Doctrine of Fluxions.—2. The Projection of the Sphere, orthographic, stereographic, and gnomonical.—3. The Elements of Trigonometry.—4. The Principles of Mechanics.—5. A Treatise of Navigation on the Sea.—6. A Treatise on Arithmetic.—7. A Treatise on Geometry.—8. A Treatise of Algebra, in 2 books.—9. The Method of Increments.—10. Arithmetic of Infinites, and the Conic Sections, with other Curve Lines.—11. Elements of Optics and Perspective.—12. Astronomy.—13. Mechanics, with Centripetal and Centrifugal Forces.—14. Mathematical Principles of Geography, Navigation, and Dialling.—15. Commentary on the Principia, with the Defence of Newton.—16. Tracts.—17. Miscellanies.

*EMINENTIAL Equation*, a term used by some algebraists, in the investigation of the areas of curvilinear figures, for a kind of assumed equation that contains another equation Eminently, the latter being a particular case of the former. *Hayes's Flux.* pa. 97.

*ENCEINTE*, a French term, in Fortification, signifying the whole inclosure, circumference, or compass of a fortified place, whether built with stone or brick, or only made of earth, and whether with or without bastions, &c.

*ENCYCLOPÆDIA*, the circle or chain of arts and sciences; sometimes also written *Cyclopædia*.

*ENDECAGON*, a plane geometrical figure of eleven sides and angles, otherwise called Undecagon. If each side of this figure be 1, its area will be 9.3656399

$= \frac{1}{4}$  of the tang. of  $73\frac{7}{11}$  degrees, to the radius 1. See my *Mensuration*, pa. 114, &c, 2d edit. See also *REGULAR Figure*.

*ENFILADE*, a French term, applied to those trenches, and other lines, that are ranged in a right line, and so may be scoured or swept by the cannon lengthways, or in the direction of the line.

*To ENFILADE*, is to sweep lengthways by the firing of cannon, &c.

*A Battery d'ENFILADE*, is that where the cannon sweep a right line.

*A Post, or Command d'ENFILADE*, is a height from whence a whole line may be swept at once.

*ENGINE*, in Mechanics; a compound machine, consisting of several simple ones, as wheels, screws, levers, or the like, combined together, in order to lift, cast, or sustain a weight, or produce some other considerable effect, so as to save either force or time.

There are numberless kinds of engines; of which some are for war, as the Balista, Catapulta, Scorpio, Aries or Ram, &c.; others for the arts of peace, as Mills, Cranes, Presses, Clocks, Watches, &c, &c.

*ENGINEER*, or *INGINEER*, is applied to a contriver or maker of any kind of useful engines or machines; or who is particularly skilled or employed in them. And he is denominated either a civil or military Engineer, according as the objects of his profession respect civil or military purposes.

A military Engineer should be an expert mathematician and draughtsman, and particularly versed in fortification and gunnery, being the person officially employed to direct the operations both for attacking and defending works. When at a siege the Engineers have narrowly surveyed the place, they are to make their report to the general, or commander, by acquainting him which part they judge the weakest, and where approaches may be made with most success. It is their business also to draw the lines of circumvallation and contravallation; also to mark out the trenches, places of arms, batteries, and lodgments, and in general to direct the workmen in all such operations.

*ENGONASIS*, in Astronomy, the same as *Hercules*, one of the northern constellations; which see.

*ENGYSCOPE*, the same as *MICROSCOPE*.

*ENHARMONIC*, the last of the three kinds of music. It abounds in dieses, or the least sensible divisions of a tone. See *Philos. Trans.* number 481; also *Wallis Appendix ad Ptolom.* pa. 165, 166.

*ENNEADECATERIS*, in Chronology, a cycle or period of 19 solar years, being the same as the Golden Number and Lunar Cycle, or Cycle of the Moon; which see; as also *EMBOLISMIC*.

*ENNEAGON*, a plane geometrical figure of 9 sides and angles; and is otherwise called a Nonagon. If each side of this figure be 1, its area will be 6.1818242  $= \frac{1}{4}$  of the tang. of 70 degrees, to the radius 1. See my *Mensuration*, pa. 114, 2d edit. See also the article *REGULAR Figure*.

*ENTABLATURE*, in Architecture, is that part of an order of column which is over the capital, comprehending the Architrave, Frize, and Corniche.

*ENTABLATURE*, or *ENTABLEMENT*, is sometimes also used for the last row of stones on the top of the wall of a building; on which the timber and covering



ing rest; sometimes also called the Drip, because it projects a little, to throw the water off.

**ENVELOPE**, in Fortification, is a mound of earth, sometimes raised in the ditch of a place, and sometimes beyond it, being either in form of a single parapet, or of a small parapet bordered with a parapet. These Envelopes are made only to cover weak parts with single lines, without advancing towards the field, which cannot be done without works that require a great deal of room, such as horn-works, half-moons, &c. Envelopes are sometimes called Sillons, Contregards, Conferves, Lunettes, &c.

**ENUMERATION**, a numbering or counting. Sir Isaac Newton wrote an ingenious treatise, being an Enumeration of the lines of the 3d order.

**EOLIPILE**. See **ÆOLIPILE**.

**EPACT**, in Chronology, the excess of the solar month above the lunar synodical month; or of the solar year above the lunar year of 12 synodical months; or of several solar months above as many synodical months; or of several solar years above as many dozen of synodical months.

The Epacts then are either Annual or Menstrual.

**Menstrual EPACTS**, are the excesses of the civil calendar month above the lunar month. Suppose, for example, it were new moon on the first day of January: then since the month of January contains 31 days,

and the lunar month - 29<sup>ds</sup> 12<sup>h</sup> 44<sup>m</sup> 3<sup>s</sup>;  
the menstrual Epact is - 1 11 15 57

**Annual EPACTS**, are the excesses of the solar year above the lunar. Hence,

as the Julian solar year is 365<sup>ds</sup> 6<sup>h</sup> 0<sup>m</sup> 0<sup>s</sup>,  
and the Julian lunar year 354 8 48 38,  
the annual Epact will be 10 21 11 22,  
that is, almost 11 days. Consequently the Epact of 2 years, is 22 days; of 3 years, 33 days; or rather 3, since 30 days make an embolismic, or intercalary month. Then, adding still 11, the Epact of 4 years is 14 days; and so of the rest as in the following table, where they do not become 30, or 0 again, till the 19th year; so that at the 20th year the Epact is 11 again; and hence the cycle of Epacts expires with the Golden Number, or Lunar Cycle of 19 years, and begins with the same again.

TABLE of Julian Epacts.

Golden Numb.	Epacts	Golden Numb.	Epacts	Golden Numb.	Epacts
I	11	VIII	28	XV	15
II	22	IX	9	XVI	26
III	3	X	20	XVII	8
IV	14	XI	1	XVIII	19
V	25	XII	12	XIX	30
VI	6	XIII	23		or 0
VII	17	XIV	4		

Again, as the new moons are the same, or fall on the same day, every 19 years, so the difference between the solar and lunar years is the same every 19 years. And because the said difference is always to be added to the lunar year, to adjust or make it equal to the solar

year; hence the said difference respectively belonging to each year of the moon's cycle, is called the *Epact of the said year*, that is, the number to be added to the said year, to make it equal to the solar year. Upon this mutual respect between the cycle of the moon and the cycle of the Epacts, is founded this

*Rule for finding the Julian Epact, belonging to any year of the Moon's Cycle.*

Multiply the Golden Number, or the given year of the Moon's Cycle, by 11, and the product will be the Epact if it be less than 30; but if it exceed 30, then throw out as many 30's as the product contains, and the remainder will be the Epact.

*Rule to find the Gregorian Epact.*

1st, The difference between the Julian and Gregorian years being equal to the difference between the solar and lunar year, or 11 days, therefore the Gregorian Epact for any year is the same with the Julian Epact for the preceding year; and hence the Gregorian Epact will be found, by subtracting 1 from the golden number, multiplying the remainder by 11, and rejecting the 30's. This rule will serve till the year 1900; but after that year, the Gregorian Epact will be found by this rule: Divide the centuries of the given year by 4; multiply the remainder by 17; then to this product add 43 times the quotient, and also the number 86, and divide the whole sum by 25, reserving the quotient: next multiply the golden number by 11, and from the product subtract the reserved quotient, so shall the remainder, after rejecting all the 30's contained in it, be the Epact sought.

The following table contains the Golden Numbers, with their corresponding Epacts, till the year 1900.

TABLE of Gregorian Epacts.

Golden Numb.	Epacts	Golden Numb.	Epacts	Golden Numb.	Epacts
I	0	VIII	17	XV	4
II	11	IX	28	XVI	15
III	22	X	9	XVII	26
IV	3	XI	20	XVIII	7
V	14	XII	1	XIX	18
VI	25	XIII	12	I	0
VII	6	XIV	23		

On the subject of Epacts, see Wollius's *Elementa Chronologiae*, apud Opera, tom. 4, pa. 133; also *Philos. Trans.* vol. 46, pa. 417, or numb. 495, art. 5.

**EPAULE**, or **ESPAULE**, in Fortification, the shoulder of the bastion, or the angle made by the face and flank, otherwise called the Angle of the Epaulé.

**EPAULEMENT**, in Fortification, a side-work hastily thrown up, to cover the cannon or the men; and is made either of earth thrown up, or bags filled with earth or sand, or of gabions, or fascines, &c, with earth: of which latter sort are commonly the Epaulements of the places of arms for the cavalry behind the trenches.

**EPAULEMENT**, is also used for a demi-bastion, consisting



sisting of a face and flank, placed at the point of a horn-work or crown-work. Also for a little flank added to the sides of horn-works, to defend them when they are too long. Also for the redoubts made on a right line, to fortify it. And lastly, for an orillon, or mass of earth almost square, faced and lined with a wall, and designed to cover the cannon of a casemate.

EPHEMERIS, EPHEMERIDES, tables calculated by astronomers, shewing the present state of the heavens for every day at noon; that is, the places in which all the planets are found at that time; differing but little from an Astronomical Almanac. It is from such tables as these that the eclipses, conjunctions, and aspects of the planets are made out; as also horoscopes, or celestial schemes, constructed, &c.

There have been Ephemerides of Origan, Kepler, Argoli, Heckerus, Mezzarachis, Wing, Gadbury, Parker, De la Hire, &c.

In France the Academy of Sciences have published annually, from the beginning of the present century, a kind of Ephemeris, under the title of *Connoissance des Temps*, which is still continued, and is in great esteem; as are also the Ephemerides, published there every ten years, by M. Desplaces, and De la Lande.

There are now published such Ephemerides by the Academies of several other nations; but that which is in most esteem for its accuracy and use in finding the longitude, is the Nautical Almanac, or Astronomical Ephemeris, published in England by the Board of Longitude, under the direction of the Rev. Dr. Maskelyne, Astronomer Royal, which commenced with the year 1767.

EPICARMUS, an ancient poet and philosopher, born in Sicily, was a scholar of Pythagoras, and flourished in the time of Hiero, in whose reign it is said he introduced comedy at Syracuse. He wrote also treatises concerning philosophy and medicine; but none of his works have been preserved. He died at 90 years of age, according to Laertius, who has preserved four verses inscribed on his statue.

EPICURUS, a celebrated ancient philosopher, was born at Gargettium in Attica, in the 109th Olympiad, or about 340 years before Christ. He settled at Athens in a fine garden he had bought; where he lived with his friends in much tranquillity, and educated a great number of disciples; who lived all in common with their master. His school was never divided, but his doctrine was followed as an oracle; and the respect which his followers paid to his memory is admirable; his birth-day being still kept in Pliny's time, and even the very month he was born in observed as a continued festival, and his picture placed every where. He wrote a great many books, and valued himself upon making no quotations. He raised the atomical system to great reputation, though he was not the inventor of it, but only made some change in that of Democritus. As to his doctrine concerning the supreme good or happiness, it was very liable to be misrepresented, and some ill effects proceeded from thence, which discredited his sect, though undeservedly. He was charged with perverting the worship of the gods, and inciting men to debauchery. But he did not forget himself on this occasion: he published his opinions to the whole world; wrote some books of devotion; recommended the ve-

neration of the gods, sobriety, and chastity, living in an exemplary manner, and conformably to the rules of philosophical wisdom and frugality. He died of a suppression of urine, at 72 years of age.—Gassendus has given us all he could collect from the ancients concerning the person and doctrine of this philosopher.

EPICUREAN *Philosophy*, the doctrine, or system of philosophy maintained by Epicurus and his followers. This consisted of three parts; canonical, physical, and ethical. The first respected the canons or rules of judging; in which soundness and simplicity of sense, assisted by some natural reflections, chiefly formed his art. His search after truth proceeded only by the senses; to the evidence of which he gave so great a certainty, that he considered them as an infallible rule of truth, and termed them the First natural light of mankind.

In the 2d part of his philosophy he laid down atoms, space, and gravity, as the first principles of all things. He asserted the existence of God, whom he accounted a blessed immortal being, but who did not concern himself with human affairs.

As to his ethics, he made the supreme good of man to consist in pleasure, and consequently supreme evil in pain. Nature itself, says he, teaches us this truth; and prompts us from our birth to procure whatever gives us pleasure, and avoid what gives us pain. To this end he proposes a remedy against the sharpness of pain, which was to divert the mind from it, by turning our whole attention upon the pleasures we have formerly enjoyed. He held that the wise man must be happy, as long as he is wise: the pain, not depriving him of his wisdom, cannot deprive him of his happiness: from which it would seem that his pleasure consisted rather in intellectual than in sensual enjoyments: though this is a point strongly contested.

EPICUREANS, the sect of philosophers holding or following the principles and doctrine of Epicurus. As the nature of the pleasure, in which the chief happiness is supposed to be seated, is a great problem in the morals of Epicurus, there hence arise two kinds of Epicureans, the rigid and the remiss: the first were those who understood Epicurus's notion of pleasure in the best sense, and placed all their happiness in the pure pleasures of the mind, arising from the practice of virtue: while the loose or remiss Epicureans, taking the words of that philosopher in a gross sense, placed all their happiness in bodily pleasures or debauchery.

EPICYCLE, in the ancient astronomy, a little circle having its centre in the circumference of a greater one: or a small orb or sphere, which being fixed in the deferent of a planet, is carried along with it, and yet, by its own peculiar motion, carries the planet fastened to it round its proper centre.

It was by means of Epicycles that Ptolomy and his followers solved the various phenomena of the planets, but more especially their stations and retrogradations.

EPICYCLOID, is a curve generated by the revolution of a point of the periphery of a circle, which rolls along or upon the circumference of another circle, either on the convex or concave side of it.


When a circle rolls along a straight line, a point in its circumference describes the curve called a cycloid.

But



But if, instead of the right line, the circle roll along the circumference of another circle, either equal to the former or not, then the curve described by any point in its circumference is what is called the Epicycloid.

If the generating circle roll along the convexity of the circumference, the curve is called an *Upper*, or *Exterior Epicycloid*; but if along the concavity, it is called a *Lower*, or *Interior Epicycloid*. Also the circle that revolves is called the *Generant*; and the arc of the other circle along which it revolves, is called the *Base* of the Epicycloid. Thus, ABC or BLV is the Generant; DPVE the Exterior Epicycloid, its axis BV; DPUE the Interior Epicycloid; and DBE their common Base.



*For the Length of the Curve.*

The length of any part of the curve of an Epicycloid, which any given point in the revolving circle has described, from the position where it touched the circle upon which it revolved, is to double the versed sine of half the arc which all the time of revolving touched the quiescent circle, as the sum of the diameters of the circles, is to the semidiameter of the quiescent circle in the Exterior Cycloid; or as the difference of the diameters is to that semidiameter, for the Interior one.

*For the Area of the Epicycloid.*

Dr. Halley has given a general proposition for the measuring of all cycloids and Epicycloids: thus, the area of a cycloid, or Epicycloid, either primary, or contracted, or prolate, is to the area of the generating circle; and also the areas of the parts generated in those curves, to the areas of analogous segments of the circle; as the sum of double the velocity of the centre and the velocity of the circular motion, is to this velocity of the circular motion. See the Demonstr. in the Philos. Transf. number 218.

*Spherical Epicycloids* are formed by a point of the revolving circle, when its plane makes a constant angle with the plane of the circle on which it revolves. Messrs. Bernoulli, Maupertuis, Nicole, and Clairaut, have demonstrated several properties of these Epicycloids, in Hist. Acad. Sci. for 1732.

*Parabolic, Elliptic, &c. Epicycloids.*

If a parabola roll upon another equal to it; its focus will describe a right line perpendicular to the axis of the quiescent parabola: also the vertex of the rolling parabola will describe the cissoid of Diocles; and any other point of it will describe some one of Newton's defective hyperbolas, having a double point in the like point of the quiescent parabola.

In like manner, if an ellipse revolve upon another ellipse, equal and similar to it, its focus will describe a circle, whose centre is in the other focus, and consequently the radius is equal to the axis of the ellipse; and any other point in the plane of the ellipse will describe a line of the 4th order.

The same may be said also of an hyperbola, revolving upon another, equal and similar to it; for one of the foci will describe a circle, having its centre in the other focus, and the radius will be the principal axis of the hyperbola; and any other point of the hyperbola will describe a line of the 4th order.

Concerning these lines, see Newton's Principia, lib. 1; also De la Hire's Memoires de Mathematique &c, where he shews the nature of this line, and its use in Mechanics; see also Maclaurin's Geometria Organica.

**EPIPHANY**, a christian festival, otherwise called the Manifestation of Christ to the Gentiles, observed on the 6th of January, in honour of the appearance of our Saviour to the three magi or wise men, who came to adore him and bring him presents.

**EPISTYLE**, in the ancient Architecture, a term used by the Greeks for what we call Architrave, viz a massive stone, or a piece of wood, laid immediately over the capital of a column.

**EPOCH**, or **EPOCH**, a term or fixed point of time, from whence the succeeding years are numbered or reckoned.

Different nations make use of different Epochs. The christians chiefly use the Epoch of the nativity or incarnation of Jesus Christ; the Mahometans, that of the Hegira; the Jews, that of the creation of the world, or that of the Deluge; the ancient Greeks, that of the Olympiads; the Romans, that of the building of their city; the ancient Persians and Assyrians, that of Nabonassar; &c.

The doctrine and use of Epochs is of very great extent in chronology. To reduce the years of one Epoch to those of another, i. e. to find what year of one corresponds to a given year of another; a period of years has been invented, which, commencing before all the known Epochs, is, as it were, a common receptacle of them all, called the Julian Period. To this period all the Epochs are reduced; i. e. the year of this period when each Epoch commences, is determined. So that, adding the given year of one Epoch to the year of the period corresponding with its rise, and from the sum subtracting the year of the same period corresponding to the other Epoch, the remainder is the year of that other Epoch.

*Epoch of Christ*, is the common Epoch throughout Europe, commencing at the supposed time of our Saviour's nativity, December 25; or rather, according to the usual account, from his circumcision, or the 1st of January. The author of this Epoch was an Abbot of Rome, one Dionysius Exiguus, a Scythian, about the year 507 or 527. Dionysius began his account from the conception or incarnation, usually called the Annunciation, or Lady Day; which method obtained in the dominions of Great Britain till the year 1752, before which time the Dionysian was the same as the English Epoch: but in that year the Gregorian calendar having been admitted by act of parliament, they now reckon from the first of January, as in the other parts of Europe, except in the court of Rome, where the Epoch of the Incarnation still obtains for the date of their bulls.



*A TABLE of the Years of the most remarkable Epochs or Eras and Events.*

N. B. The years before Christ, are those before the reputed year of his birth, and not reckoned back from the first year of his age, as is generally done in such tables.	Julian Period.	Year of the World.	Years before Christ.
The Creation of the World	706	0	4007
The Deluge, or Noah's flood	2362	1656	2351
Assyrian monarchy founded by Nimrod	2537	1831	2176
The birth of Abraham	2714	2008	1999
Kingdom of Athens founded by Cecrops	3157	2451	1556
Entrance of the Israelites into Canaan	3262	2556	1451
The destruction of Troy	3529	2823	1184
Solomon's temple founded	3701	2995	1012
The Argonautic expedition	3776	3070	937
Lycurgus formed his laws	3829	3103	884
Arbaces, 1st king of the Medes	3838	3132	875
Olympiads of the Greeks began	3938	3232	775
Rome built, or Roman Era	3961	3255	752
Era of Nabonassar	3967	3261	746
First Babylonish captivity, by Nebuchadnezzar	4107	3401	606
The 2d ditto, and birth of Cyrus	4114	3408	599
Solomon's temple destroyed	4125	3419	588
Cyrus began to reign in Babylon	4177	3471	536
Peloponnesian war began	4282	3576	431
Alexander the great died	4390	3684	323
Captivity of 100,000 Jews by Ptolomy	4393	3687	320
Archimedes killed at Syracuse	4506	3800	207
Julius Cæsar invaded Britain	4659	3953	54
He corrected the calendar	4667	3961	46
The true year of Christ's birth	4709	4003	4
<i>The Christian Era begins here.</i>			Years since Christ.
Dionysian, or vulgar era of Christ's birth	4717	4007	0
Christ crucified, Friday April 3d	4746	4040	33
Jerusalem destroyed	4783	4077	70
Adrian's wall built in Britain	4833	4127	120
Dioclesian Epoch, or that of Martyrs	4997	4291	284
The council of Nice	5038	4332	325
Constantine the great died	5050	4344	337
The Saxons invited into Britain	5158	4452	445
Hegira, or flight of Mohammed	5335	4629	622
Death of Mohammed	5343	4637	630
The Persian Yefdegird	5344	4638	631
Sun, Moon, and Planets $\frac{1}{2}$ , $\frac{1}{4}$ , $\frac{1}{8}$ , $\frac{1}{16}$ , $\frac{1}{32}$ in $\frac{1}{2}$ , seen from the earth	5899	5193	1186
Art of printing discovered	6153	5447	1440
The reformation begun by Martin Luther	6230	5524	1517
The Calendar corrected by pope Gregory	6295	5589	1582
Oliver Cromwell died	6371	5665	1658
Sir Isaac Newton born, Dec. 25	6355	5649	1642
Made President of the Royal Society	6416	5710	1703
Died, March 20th	6440	5734	1727
New Planet discovered by Herschel	6494	5788	1781



**EQUABLE Motion**, Celerity, Velocity, &c, is that which is uniform, or without alteration, or by which equal spaces are passed over in equal times. Hence, the spaces, passed over in Equable motions, are proportional to the times. So that if a body pass over 20 feet in 1 second of time, it will pass over 40 feet in 2 seconds, and so on.

**EQUABLY Accelerated or Retarded**, &c, is when the motion or change is increased or decreased by equal quantities or degrees in equal times.

**EQUAL**, a term of relation between different things, but of the same kind, magnitude, quantity, or quality.—Wolfius defines Equals to be those things that may be substituted for each other, without any alteration of their quantity.—It is an axiom in mathematics &c, that two things which are equal to the same third, are also equal to each other. And if Equals be equally altered, by Equal addition, subtraction, multiplication, division, &c, the results will be also Equal.

**EQUAL Circles**, are those whose diameters are equal.

**EQUAL Angles**, are those whose sides are equally inclined, or which are measured by similar arcs of circles.

**EQUAL Lines**, are lines of the same length.

**EQUAL Plane Figures**, are those whose areas are equal; whether the figures be of the same form or not.

**EQUAL Solids**, are such as are of the same space, capacity, or solid content; whether they be of the same kind or not.

**EQUAL Curvatures**, are such as have the same or equal radii of curvature.

**EQUAL Ratios**, are those whose terms are in the same proportion.

**EQUAL**, in Optics, is said of things that are seen under equal angles.

**EQUALITY**, the exact agreement of two things in respect of quantity. Those figures are equal which may occupy the same space, or may be conceived to possess the same space, by the flexion or transposition of their parts. See a learned discourse upon this, by Dr. Barrow, in the 11th and 12th of his Mathematical Lectures.

**EQUALITY**, in Algebra, the relation or comparison between two quantities that are really or effectually equal. See EQUATION.

Equality, in Algebra, is usually denoted by two equal parallel lines, as  $=$ : thus  $2 + 3 = 5$ , i. e. 2 plus 3, are equal to 5. This character  $=$ , was first introduced by Robert Recorde. Des Cartes, and some others after him, use the mark  $\propto$  instead of it: as  $2 + 3 \propto 5$ .

**EQUALITY**, in Astronomy. *Circle of EQUALITY*, or the **EQUANT**. See CIRCLE and EQUANT.

*Ratio or Proportion of EQUALITY*, is that between two equal numbers or quantities.

*Proportion of EQUALITY evenly ranged*, or *ex aequo ordinata*, is that in which two terms, in a rank or series, are proportional to as many terms in another series, compared to each other in the same order, i. e. the first of one rank to the first of another, the 2d to the 2d, &c.

*Proportion of EQUALITY evenly disturbed*, called also

*ex aequo perturbata*, is that in which more than two terms of one rank, are proportional to as many terms of another, compared to each other in a different and interrupted order; viz, the 1st of one rank to the 2d of another, the 2d to the 3d, &c.

**EQUANT**, or **ÆQUANT**, in Astronomy, a circle formerly conceived by astronomers, in the plane of the deferent, or eccentric; for regulating and adjusting certain motions of the planets, and reducing them more easily to a calculus.

**EQUATED Anomaly**. See ANOMALY.

**EQUATED Bodies**. On Gunter's Sector there are sometimes placed two lines, answering to one another, and called the Lines of Equated Bodies. They are situated between the lines of superficies and solids, and are marked with the letters D, I, C, S, O, T, to signify dodecahedron, icosaedron, cube, sphere, octahedron, and tetrahedron.

The uses of these lines are, 1st, When the diameter of the sphere is given, to find the sides of the five regular bodies, each equal to that sphere; 2d, From the side of any one of those bodies being given, to find the diameter of the sphere, and the sides of the other bodies, which shall be each equal to the first given body. So, when the sphere is given, take its diameter, and apply it over on the sector in the points S, S; but when one of the other five bodies is given, apply its side over in its proper points; then the parallels taken from between the points of the other bodies, or sphere, shall be the sides or diameter, equal severally to the sphere or body first given.

**EQUATION**, in Algebra, an expression of equality between two different quantities; or two quantities, whether simple or compound, with the mark of equality between them: as  $2 + 3 = 7 - 2$ ; or  $2 \times 3 = 6$ ; or  $5 \times 3 = \frac{30}{2}$ ; or  $a + b = c$ ; or  $x^2 + ax = b$ ;

&c. When the things, or two sides of the equation, are the same, the expression becomes an Identity, as  $5 = 5$ , or  $a = a$ , &c. Sometimes the quantities are placed all on one side, and made equal to 0 or nothing on the other side; as  $5 - 5 = 0$ , or  $a - b = 0$ : which is no more than setting down the difference of two equal quantities equal to nothing.

The character or sign usually employed to denote an Equation, is  $=$ , which is placed between the two equal quantities, called the two sides of the Equation.

The *Terms* of an EQUATION, are the several quantities or parts of which it is composed. Thus, of the Equation  $a + b = c$ , the terms are  $a$ ,  $b$ , and  $c$ : and the tenor or import of the expression is, that some quantity represented by  $c$ , is equal to two others represented by  $a$  and  $b$ .

Equations are either Simple or Affected.

A *Simple EQUATION* is that which has only one power of the unknown quantity: as  $a + x = 3b$ , or  $ax^2 = bc$ ,  $a^3 + 2x^3 = 5b$ , &c; where  $x$  denotes the unknown quantity, and the other letters known ones. But

An *Affected*, or *Adaffected EQUATION* contains two or more different powers: as  $x^2 + ax = b$ , or  $3x - 4x^2 + x^3 = 25$ , &c.



Again, Equations are denominated from the highest power contained in them; as quadratic, cubic, biquadratic, &c. Thus,

A *Quadratic EQUATION*, is that in which the unknown quantity rises to two dimensions, or to the square or 2d power: as  $x^2 + 20x = 200$ , or  $ax - x^2 = b$ .

A *Cubic EQUATION*, is that in which the unknown quantity is of three dimensions, or rises to the cube or 3d power: as  $x^3 = 25$ , or  $x^3 - 2x^2 = 27$ , or  $x^3 - ax^2 + bx = c$ .

A *Biquadratic EQUATION*, is that in which the unknown quantity is of 4 dimensions, or rises to the 4th or biquadratic power: as  $x^4 = 25$ , or  $x^4 - 20x = 10$ , or  $x^4 - ax^3 + bx^2 - cx = d$ .

And so for other higher orders of Equations.

The *Root* of an EQUATION, is the value of the unknown letter or quantity contained in it. And this value being substituted in the terms of the Equation instead of that letter or quantity, will cause both sides to vanish, or will make the one side exactly equal to

the other. So the root of the Equation  $\frac{3x}{5} + 18 = 24$ , is 10; because that using 10 for  $x$ , it becomes  $\frac{3 \times 10}{5} + 18 = 6 + 18 = 24$ .

Every Equation has as many roots as it has dimensions, or as it contains units in the index of the highest power, when the powers are all reduced to integral exponents. So the simple Equation of the 1st power, has only one root; but the quadratic has 2, the cubic 3, the biquadratic 4, &c. Thus the two roots of this equation  $x^2 - 4x = -3$  are 1 and 3; for either of these substituted for  $x$  makes  $x^2 - 4x$  come out equal to  $-3$ . Also the three roots of  $x^3 - 4x^2 - 11x = -30$ , or  $x^3 - 4x^2 - 11x + 30 = 0$ , are 2, 5, and  $-3$ ; as will appear by substituting each of these instead of  $x$  in the equation, which will make all the terms on one side equal to the other side. And so of others.

*The Relation between the Roots of Equations, and the Coefficients of their Terms.*—In every Equation, when the terms are ranged in order according to the order of the powers, the greater before the less; the first term or highest power freed from its coefficient, by dividing all the terms by it, and all brought to one side, and made equal to nothing on the other side, when it will appear in this form,

$$x^n + ax^{n-1} + bx^{n-2} + cx^{n-3} \dots = 0;$$

then the relations between the roots and coefficients, are as follow:

1st, The coefficient  $a$  of the 2d term, is equal to the sum of all the roots.

2d, The coefficient  $b$  of the 3d term, is equal to the sum of all the products of the roots that can be made by multiplying every two of them together. In like manner,

3d, The coefficients  $c, d, e$ , &c, of the following terms, are respectively equal to the sum of the products of the roots made by multiplying every three together, or every four together, or every five together, &c, the signs of all the roots being changed. All which will appear below, in the Generation of Equations.

The Roots of Equations are Positive or Negative,

and Real or Imaginary. Thus, the two roots of the Equation  $x^2 - 4x = -3$ , are 1 and 3, real and both positive; but the roots of the Equation  $x^3 - 4x^2 - 11x = -30$ , are 2, 5, &  $-3$ , are real, two positive and one negative; and the roots of the equation  $x^3 + 9x = 10$ , are 1 and  $-\frac{1}{2} \pm \frac{1}{2} \sqrt{-39}$ , one real and two imaginary.

The *Generation* of EQUATIONS, is the multiplying of certain assumed simple equations together, to produce compound ones, with intent to shew the nature of these; a method invented by Harriot, which is this: Suppose  $x$  to denote the unknown quantity of any equation, and let the roots of that equation, or the values of  $x$ , be,  $a, b, c, d$ , &c; that is  $x = a$ , and  $x = b$ , and  $x = c$ , &c; or  $x - a = 0$ , and  $x - b = 0$ , and  $x - c = 0$ , &c; then multiply these last equations together, thus,

$$x - a = 0$$

$$x - b = 0$$

$$\begin{array}{r} x^2 - ax \\ -bx + ab \end{array}$$

$$\left. \begin{array}{r} x^2 - a \\ -b \end{array} \right\} \cdot x + ab = 0, \text{ the quadratic Equation,}$$

$$x - c \dots \dots = 0$$

$$\begin{array}{r} x^3 - a + b \cdot x^2 + abx \\ -c \cdot x^2 + ac + bc \cdot x - abc \end{array}$$

$$\begin{array}{r} x^3 - a \cdot x^2 + ab \cdot x - abc = 0, \text{ the cubic Equ.} \\ -b \quad + ac \\ -c \quad + bc \\ x - d \quad \quad \quad = 0 \end{array}$$

$$\begin{array}{r} x^4 - a \cdot x^3 + ab \cdot x^2 - abc \cdot x + abcd = 0, \text{ the} \\ -b \quad + ac \quad -abd \quad \text{biquadratic Equ.} \\ -c \quad + bc \quad -acd \\ -d \quad + ad \quad -bcd \\ \quad + bd \\ \quad + cd \end{array}$$

Now the roots of these equations are  $a, b, c, d$ , &c; and it is obvious that the sum of all the roots is the coefficient of the 2d term, the sum of all the products of every two is the coefficient of the 3d term, the sum of all the products of every three that of the 4th term, and so on, to the last term, which is the continual product of all the roots.

*Reduction* of EQUATIONS, is the transforming or changing them to their simplest and most commodious form, to prepare them for finding or extracting their roots. The most convenient form is, that the terms be ranged according to the powers of the unknown letter, the highest power foremost next the left hand, and that term to have only  $+1$  for its coefficient; also all the terms containing the unknown letter to be on one side of the equation, and the absolute known term only on the other side.

Now this reduction chiefly respects the first term, or that which contains the highest power of the unknown quantity; and the general rule for reducing it is, to consider in what manner it is involved or connected with other quantities, and then perform the counter or opposite



posite relation or operation; for every operation is undone or counteracted by the reverse of it; as addition by subtraction, multiplication by division, involution by evolution, &c: then bring all the unknown terms to one side, and the known term to the other side, changing the signs, from + to -, or from - to +, of those terms which are changed from one side to the other; and lastly divide by the coefficient of the first term, with its sign.

Thus, for ex. if  $5x - 12 = 3x + 4$  :  
then is  $5x - 3x = 12 + 4$ , or  $2x = 16$ ;  
and so  $x = \frac{16}{2} = 8$ .

Again, if  $ax - b = cx + d$  :  
then  $ax - cx = b + d$ ,  
and  $x = \frac{b + d}{a - c}$

And, if  $6x^2 - 20 = 12x + 2x^2$  :  
then  $6x^2 - 2x^2 = 12x + 20$ ,  
or  $4x^2 - 12x = 20$ ,  
and  $x^2 - 3x = 5$ .

Also, if  $\frac{a}{a+x} + \frac{b}{x} = 3$  :  
then  $ax + ab + bx = 3ax + 3x^2$ ,  
and  $-3x^2 + ax + bx = 3ax - ab$ ,  
or  $-3x^2 + bx - 2ax = -ab$ ,  
and  $x^2 + \frac{2a-b}{3}x = \frac{ab}{3}$ .

Also, if  $\sqrt{a^2 + x^2} - 3b = \frac{2ab}{x}$  :

then  $\sqrt{a^2 + x^2} = 3b + \frac{2ab}{x} = \frac{3bx + 2ab}{x}$ ,  
and  $a^2 + x^2 = \frac{9b^2x^2 + 12ab^2x + 4a^2b^2}{x^2}$ ,  
and  $a^2x^2 + x^4 = 9b^2x^2 + 12ab^2x + 4a^2b^2$ ,  
and  $x^4 + a^2 - 9b^2x^2 - 12ab^2x = 4a^2b^2$ .

#### Extracting or finding the Roots of EQUATIONS.

This is finding the value or values of the unknown letter in an Equation, the rules for which are various, according to the degree of the Equation.

##### 1. For the Root of a Simple EQUATION.

Having reduced the equation as above, by bringing the unknown terms to one side, and the known ones to the other, freeing the former from radicals and fractions, by their counter operations, and lastly dividing by the coefficients of the unknown quantity, the value of it is then found: as in the first and 2d examples of reduction above given.

##### 2. For the Roots of Quadratic EQUATIONS.

These are usually found by what is called completing the square; which consists in squaring half the coefficient of the 2d term, and adding it to both sides of the equation; for then the unknown side is a complete square of a binomial, and the other side consists only of known quantities. Therefore, extract the root on both sides, so shall the root of the first side be a binomial, one part of which is the unknown letter, and the

other a known or given quantity, and the root of the other side is taken either + or -, since the square of either of these is the same given quantity: lastly, bringing over the known part of the binomial root to the other side, with a contrary sign, gives the two roots or values of the unknown letter sought.

Thus, if  $x^2 + 2ax = b^2$  be a general quadratic Equation,  $2a$  being the coefficient of the 2d term, and  $b^2$  the absolute known term, both with their signs. Then,  $a$  is half that coefficient, and  $a^2$  its square; which being added, gives  $x^2 + 2ax + a^2 = a^2 + b^2$ ; and the root extracted gives  $x + a = \pm \sqrt{a^2 + b^2}$ ; then, transposing  $a$ , it is  $x = -a \pm \sqrt{a^2 + b^2}$ , the two roots, or values of  $x$ .

And if  $x^2 - 4x = 5$  : then, adding  $2^2$  or 4,  
it is  $x^2 - 4x + 4 = 9$ ;  
the root is  $x - 2 = \pm 3$ ,  
then  $x = 2 \pm 3 = 5$  or  $-1$ ,  
the two roots, or values of  $x$ .

And if  $x^2 + 4x = 5$  :  
then  $x^2 + 4x + 4 = 9$ ;  
and  $x + 2 = \pm 3$ ;  
or  $x = -2 \pm 3 = 1$ , or  $-5$ , the two roots, which are the same numbers as before, but with the signs changed.

Also, if  $x^2 - 4x = -5$  :  
then  $x^2 - 4x + 4 = 4 - 5 = -1$ ;  
and  $x^2 - 2 = \pm \sqrt{-1}$ ;  
or  $x = 2 \pm \sqrt{-1}$ , two imaginary roots.

##### 3. For the Roots of Cubic EQUATIONS.

A Cubic EQUATION is that in which the unknown letter ascends to the 3d power; as  $x^3 + a^3 = b^3$ , or  $x^3 + ax^2 + bx = c$ .

The 2d term of every Cubic Equation being taken away, those equations may all be reduced to this form,  $x^3 + ax = b$ ; and the general value of one root is

$x = \sqrt[3]{\frac{1}{2}b + \sqrt{\frac{1}{4}b^2 + \frac{1}{27}a^3}} + \sqrt[3]{\frac{1}{2}b - \sqrt{\frac{1}{4}b^2 + \frac{1}{27}a^3}}$ . This rule is usually called Cardan's, because first published by him, but it was invented both by Scipio Ferreus, and Nich. Tartalea, by whom it was communicated to Cardan. See the article ALGEBRA.

When the 2d term is negative, or the equation of this form,  $x^3 - ax = \pm b$ , the radical  $\sqrt{\frac{1}{4}b^2 + \frac{1}{27}a^3}$  becomes  $\sqrt{\frac{1}{4}b^2 - \frac{1}{27}a^3}$ , which will be imaginary or impossible when  $\frac{1}{27}a^3$  is greater than  $\frac{1}{4}b^2$ , for  $\sqrt{\frac{1}{4}b^2 - \frac{1}{27}a^3}$  will then be the square root of a negative quantity, which is impossible: and yet, in this case, the root  $x$  is a real quantity; though algebraists have never been able to find a real finite general expression for it. And this is called the Irreducible or Impracticable Case.

This case may indeed be resolved by the trisection of an arc or angle; or by any of the usual methods of converging; or by general expressions in infinite series. See Saunderson's Algebra, pa. 713; Philos. Transf. vol. 18, pa. 136, or Abr. vol. 1, pa. 87; also vol. 70, pa. 415. See also the article Cubic Equations.

Mr. Cotes observes, in his Logometria, pa. 29, that the solution of all cubic Equations depends either upon the trisection of a ratio, or of an angle. See this method explained in Saunderson's Alg. p. 718.

*Biquadratic*



*Biquadratic EQUATIONS*, or those that are of 4 dimensions, are resolved after various methods. The first rule was given by Lewis Ferrari, the companion of Cardan, which is one of the best. A 2d method was given by Des Cartes, and another by Mr. Simpson and Dr. Waring. For the explanation of which, see *Biquadratic Equations*.

*EQUATIONS of the Higher Degrees or Orders.*

There is no general rule to express algebraically the roots of Equations above those of the 4th degree; and therefore methods of approximation are here made use of, which, though not accurately, are yet practically true. Some of these excel in ease and simplicity, and others in quickness of converging. Among these may be reckoned first, Double Position, or Trial-and-Error, both in respect of ease and universality, as it applies in the simplest manner to all sorts of Equations whatever, not excepting even exponential ones, radical expressions of ever so complex a form, expressions of logarithms, of arches by the sines or tangents, of arcs of curves by the abscissas, or any other fluents, or roots of fluxional Equations. For an explanation of this and other methods of converging to the roots of equations, by Halley, Newton, Raphson, &c, &c, see *APPROXIMATION*, and *CONVERGING*.

Besides the methods above adverted to, there have been some others, given in the Memoirs of several Academies, and elsewhere. As, by M. Daniel Bernoulli, in the *Acta Petropolitana*, tom. 3, p. 92; and by M. L. Euler, in the same, vol. 6, New Series, and tom. 5, p. 63 & 82; by Mr. Thos. Simpson, in his *Essays*, p. 82; in his *Dissertations*, p. 102; in his *Algebra*, p. 158; and in his *Select Exercises*, p. 215.

*Absolute EQUATION*. See *ABSOLUTE*.

*Affected, or Affected EQUATION*. See *AFFECTED*.

*Differential EQUATION*, is the Equation of Differences or Fluxions.

*Eminential EQUATION*. See *EMINENTIAL*.

*Exponential EQUATION*, one in which the exponents of the powers are variable or unknown quantities. See *EXPONENTIAL*.

*Fluential EQUATION*, is the Equation of the fluents.

*Fluxional EQUATION*, is the Equation of the fluxions.

*Literal EQUATION*, is a general Equation expressed in letters, as contradistinguished from a

*Numeral Equation*, one expressed in numbers.

*Transcendental EQUATION*. See *TRANSCENDENTAL*.

*EQUATION*, in Astronomy, as *Annual EQUATION*, is either of the mean motion of the sun and moon, or of the moon's apogee and nodes.

The Annual Equation of the sun's centre being given, the other three corresponding annual Equations, will be also given, and therefore a table of the first will serve for all of them. Thus, if the annual Equation of the sun's centre, taken from such a table for any time, be called  $s$ ; and if  $\frac{1}{10}s = A$ , and  $\frac{1}{6}s = B$ ; then shall the other annual Equations for that time be thus,

$A + \frac{1}{30}A = m$ , that of the moon's mean motion; and  $B + \frac{1}{30}B = a$ , that of the moon's apogee; and  $\frac{1}{2}B - \frac{1}{120}B = n$ , that of her nodes.

And here note, that when  $s$ , or the Equation of the sun's centre, is additive; then  $m$  is negative,  $a$  is positive, and  $n$  is negative. But on the contrary, when  $s$

is negative or subductive; then  $m$  is positive,  $a$  negative, and  $n$  positive.

There is also an *Equation of the moon's mean motion*, depending on the situation of her apogee in respect of the sun; which is greatest when the moon's apogee is in an octant with the sun, and is nothing at all when it is in the quadratures or syzygies. This Equation when greatest, and the sun in perigee, is  $3' 56''$ . But it is never above  $3' 34''$  when the sun is in apogee. At other distances of the sun from the earth, this Equation when greatest, is reciprocally as the cube of that distance. But when the moon's apogee is any where out of the octants, this Equation grows less, and is mostly, at the same distance between the earth and sun, as the sine of double the distance of the moon's apogee from the next quadrature or syzygy, is to radius. This is to be added to the moon's motion while her apogee passes from a quadrature with the sun to a syzygy; but is to be subtracted from it, while the apogee moves from the syzygy to the quadrature.

There is moreover another *Equation of the moon's motion*, which depends on the aspect of the nodes of the moon's orbit with respect to the sun: and this is greatest when her nodes are in octants to the sun, and quite vanishes when they come to their quadratures or syzygies. This Equation is proportional to the sine of double the distance of the node from the next syzygy or quadrature; and at the greatest is only  $47''$ . This must be added to the moon's mean motion while the nodes are passing from the syzygies with the sun to their quadratures; but subtracted while they pass from the quadratures to the syzygies.

From the sun's true place subtract the equated mean motion of the lunar apogee, as was shewn above, the remainder will be the annual argument of the said apogee; from whence the eccentricity of the moon and the 2d Equation of her apogee may be compared. See *Theory of the MOON's motions*, &c.

*EQUATION of the Centre*, called also *Prosthapheresis*, and *Total Prosthapheresis*, is the difference between the true and mean place of a planet, or the angle made by the lines of the true and mean place; or, which amounts to the same, between the mean and equated anomaly.

The greatest Equation of the Centre may be obtained by finding the sun's longitude at the times when he is near his mean distances, for then the difference will give the true motion for that interval of time: next find the sun's mean motion for the same interval of time; then half the difference between the true and mean motions will shew the greatest Equation of the Centre.

For Example, by observations made at the Royal Observatory at Greenwich, it appears that at the following mean times the sun's longitudes were as annexed; viz,

Mean times.	Sun's longitudes.
1769 Oct. 1 at $23^h 49^m 12^s$	$6^s 9^o 32' 0.6''$
1770 Mar. 29 at $0 4 50$	$0 8 50 27.5$

diff. of time  $178^d 0 15 38$ ; true dif. lon.  $5 29 18 27$

tropical year  $= 365^d 5^h 48^m 42^s = 365.2421527$ ;  
observed interval  $= 178 0 15 38 = 178.0108565$ ;

Then



Then

365.2421527 : 178.01085648 :: 360° : 175.455948

or 175° 27' 21" the mean motion.

Therefore - 175° 27' 21" of mean motion,

answers to - 179 18 27 of true motion;

their difference is 3 51 6

and its half - 1 55 33

is the greatest Equation of the Centre according to these observations.

To find the Equation of the Centre, or to resolve Kepler's problem, is a very troublesome operation, especially in the more eccentric orbits. How this is to be done, has been shewn by Newton, Gregory, Keil, Machin, La Caille, and others, by methods little differing from one another; which consist chiefly in finding a certain intermediate angle, called the eccentric anomaly; having known the mean anomaly, and the dimensions of the sun's orbit. The mean anomaly is easily found, by determining the exact time when the sun is in the aphelion, and using the following proportion, viz,

As the time of a tropical revolution, or solar year,  
Is to the interval between the aphelion and given time,  
So is 360 degrees, to the degrees of the mean anomaly.  
Or it may be found by taking the sun's mean motion, at the given time out of tables.

To find the Eccentric Anomaly, say,

As the aphelion distance;

Is to the perihelion distance;

So is the tangent of half the mean anomaly,

To the tangent of an arc.

Which arc added to half the mean anomaly, gives the eccentric anomaly. Then,

To find the True Anomaly, say,

As the square root of the aphelion distance,

Is to the square root of the perihelion distance;

So is the tangent of half the eccentric anomaly,

To the tangent of half the true anomaly.

Then, the difference between the true and mean anomaly, gives the Equation of the Centre, sought. Which is subtractive, from the aphelion to the perihelion, or in the first 6 signs of anomaly; and additive, from the perihelion to the aphelion, or in the last 6 signs of anomaly; and hence called *Prosthaphæresis*.

By this problem a table may easily be formed. When the Equations of the Centre for every degree of the first 6 signs of mean anomaly are found, they will serve also for the degrees of the last 6 signs, because equal anomalies are at equal distances on both sides of their apses. Then set these equations orderly to their signs and degrees of anomaly; the first 6 being reckoned from the top of the table downwards, and signed *subtract*; the last 6, for which the same Equations serve, in a contrary order, being reckoned from the bottom upwards, and marked *add*. Let also the difference between every adjacent two Equations, called Tabular Differences, be set in another column. Hence, from these Equations of the Centre, augmented or diminished by the proportional parts of their respective tabular differences, for any given minutes and seconds, may easily be deduced Equations of the Centre to any mean anomaly proposed. Robertson's Elem. of Navig. book 5, p. 286, 290, 295, and 308, where such a table of Equations is given.

The late excellent Mr. Euler has particularly considered this subject, in the Mem. de l'Acad. de Berlin, tom. 2, p. 225 & seq. where he resolves the following problems:

1. To find the true and mean anomaly corresponding to the planet's mean distance from the sun; that is, where the planet is in the extremity of the conjugate axis of its orbit.

2. The eccentricity of a planet being given, to find the eccentric anomaly corresponding to the greatest Equation.

3. The eccentricity being given, to find the mean anomaly corresponding to the greatest Equation.

4. From the same data, to find the true anomaly corresponding to this Equation.

5. From the same data, to find the greatest Equation.

6. The greatest Equation being given, to find the eccentricity.

Mr. Euler observes, that this problem is very difficult, and that it can only be resolved by approximation, and tentatively, in the manner he mentions: but if the eccentricity be not great, it may then be found directly from the greatest Equation. Thus, if the greatest Equation be  $= m$ , and the eccentricity  $= n$ ;

then is 
$$m = 2n + \frac{11}{2^2 \cdot 3} n^3 + \frac{599}{2^2 \cdot 3 \cdot 5 \cdot 7} n^5 + \&c.$$

Whence by reversion

$$n = \frac{1}{2}m - \frac{11}{2^2 \cdot 3} m^3 - \frac{587}{2^1 \cdot 3 \cdot 5} m^5 - \&c.$$

Where the greatest equation  $m$  must be expressed in parts of the radius, which may be done by reducing the angle  $m$  into seconds, and adding 4.6855749 to the log. of the resulting number, which will be the log. of the number  $m$ .

The mean anomaly to which this greatest equation corresponds, will be

$$x = 90^\circ + \frac{5}{8} m - \frac{5}{2^2 \cdot 3} m^3 - \frac{1}{2^2 \cdot 5} m^5 - \&c. \quad \text{Whence,}$$

if  $90^\circ$  be added to  $\frac{5}{8}$  of the greatest Equation, the sum will be the mean anomaly sufficiently exact.

Mr. Euler subjoins a table, by which may be found the greatest Equations, with the mean and eccentric anomalies corresponding to these greatest Equations for every 100th part of unity, which he supposes equal to the greatest eccentricity, or when the transverse and distance of the foci become infinite. The last column of the table gives also the logarithm of that distance of the planet from the sun where its Equation is greatest. By means of this table, any eccentricity being given, by interpolation will be found the corresponding greatest Equation. But the chief use of the table is to determine the eccentricity when the greatest Equation is known; and without this help Mr. Euler thinks the problem cannot be resolved.

EQUATION of Time, denotes the difference between mean and apparent time, or the reduction of the apparent unequal time, or motion of the sun or a planet, to equal and mean time, or motion; or the Equation of time is the difference between the sun's mean motion, and his right ascension. Apparent time is that which takes its beginning from the passage of the sun's centre over the meridian of any place; and had the sun no motion



motion in the ecliptic, or was his motion reduced to the equator or in right ascension uniform, he would always return to the meridian after equal intervals of time. But his apparent motion in the ecliptic being continually varying, and his motion in right ascension being rendered farther unequal on account of the obliquity of the ecliptic to the equator, from these causes it arises that the intervals of his return to the meridian become unequal, and the sun will gradually come too slow or too soon to the meridian for an equable motion, such as that of clocks and watches ought to be; and this retardation or acceleration of the sun's coming to the meridian, is called the equation of time.

Now, computing the celestial motions according to equal time, it is necessary to turn that time back again into apparent time, that they may correspond to observation: on the contrary, any phenomenon being observed, the apparent time of it must be converted into equal time, to have it correspond with the times marked in the astronomical tables.

The Equation of time is nothing at four different times in the year, when the whole mean and unequal motions exactly agree; viz, about the 15th of April,

the 15th of June, the 31st of August, and the 24th of December: at all other times the sun is either too fast or too slow for mean, equal, or clock time, by a certain number of minutes and seconds, which at the greatest is 16' 14'', and happens about the 1st of November; every other day throughout the year having a certain quantity of this difference belonging to it; which however is not exactly the same every year, but only every 4th year; for which reason it is necessary to have 4 tables of this Equation, viz, one for each of the four years in the period of leap years. Instead of these, may be here inserted, as follows, one general equation of time, according to the place of the sun, in every point of the ecliptic: where it is to be observed, that the sign of the ecliptic is placed at the tops of the columns, and the particular degree of the sun's place, in each sign, in the first and last columns; and in the angle of meeting in all the other columns, is the equation of time, in minutes and seconds, when the sun has any particular longitude: supposing the obliquity of the ecliptic 23° 28', and the sun's apogee in 9° of ♊.

A Table of the Equation of Time, for every Degree of the Sun's Longitude.

De- g.	♈ 0 m s	♈ 1 m s	♈ 2 m s	♈ 3 m s	♈ 4 m s	♈ 5 m s	♈ 6 m s	♈ 7 m s	♈ 8 m s	♈ 9 m s	♈ 10 m s	♈ 11 m s	De- g.
0	7+36	1-9	3-51	1+13	5+57	2+20	7-38	15-31	13-33	1-11	11+28	14+19	0
1	7 17	1 23	3 47	1 26	5 59	2 4	7 58	15 39	13 17	0 42	11 45	14 13	1
2	6 58	1 36	3 41	1 40	6 0	1 48	8 19	15 46	13 0	0 12	12 1	14 6	2
3	6 39	1 48	3 37	1 53	6 1	1 31	8 40	15 52	12 42	0+17	12 17	13 59	3
4	6 20	2 0	3 32	2 7	6 1	1 14	9 1	15 57	12 23	0 46	12 32	13 51	4
5	6 1	2 11	3 26	2 20	6 0	0 56	9 21	16 2	12 4	1 16	12 46	13 43	5
6	5 42	2 22	3 19	2 33	5 59	0 38	9 41	16 6	11 44	1 45	12 59	13 34	6
7	5 24	2 32	3 12	2 45	5 57	0 20	10 1	16 9	11 23	2 14	13 12	13 24	7
8	5 5	2 42	3 4	2 58	5 54	0 1	10 20	16 11	11 1	2 43	13 24	13 14	8
9	4 47	2 51	2 56	3 11	5 51	0-18	10 39	16 13	10 39	3 11	13 35	13 3	9
10	4 28	3 0	2 47	3 23	5 47	0 37	10 57	16 13	10 16	3 39	13 45	12 51	10
11	4 9	3 8	2 38	3 35	5 42	0 57	11 15	16 13	9 53	4 7	13 54	12 39	11
12	3 50	3 16	2 29	3 46	5 37	1 17	11 33	16 12	9 29	4 35	14 2	12 27	12
13	3 32	3 23	2 19	3 58	5 31	1 38	11 51	16 10	9 5	5 2	14 9	12 14	13
14	3 13	3 30	2 8	4 9	5 24	1 58	12 8	16 7	8 40	5 29	14 16	12 0	14
15	2 55	3 36	1 57	4 19	5 17	2 19	12 25	16 4	8 14	5 56	14 22	11 46	15
16	2 37	3 41	1 46	4 29	5 9	2 40	12 41	16 0	7 48	6 22	14 27	11 31	16
17	2 19	3 46	1 35	4 39	5 1	3 1	12 57	15 55	7 22	6 48	14 31	11 16	17
18	2 1	3 50	1 23	4 48	4 52	3 22	13 12	15 49	6 55	7 13	14 35	11 1	18
19	1 43	3 53	1 11	4 57	4 43	3 44	13 27	15 42	6 28	7 37	14 38	10 46	19
20	1 26	3 56	0 59	5 5	4 33	4 5	13 42	15 35	6 0	8 1	14 40	10 30	20
21	1 9	3 58	0 46	5 13	4 22	4 26	13 56	15 26	5 32	8 24	14 41	10 14	21
22	0 52	4 0	0 34	5 20	4 11	4 47	14 9	15 17	5 4	8 47	14 42	9 58	22
23	0 36	4 1	0 21	5 27	3 59	5 9	14 21	15 7	4 36	9 9	14 41	9 41	23
24	0 20	4 1	0 8	5 33	3 46	5 30	14 33	14 56	4 8	9 31	14 40	9 24	24
25	0 4	4 1	0+5	5 39	3 33	5 52	14 44	14 44	3 39	9 53	14 39	9 6	25
26	0-11	4 0	0 19	5 44	3 19	6 13	14 53	14 31	3 10	10 14	14 37	8 48	26
27	0 26	3 59	0 31	5 48	3 4	6 35	15 5	14 17	2 41	10 34	14 34	8 30	27
28	0 40	3 57	0 46	5 52	2 50	6 56	15 14	14 3	2 11	10 53	14 30	8 12	28
29	0 53	3 54	0 59	5 55	2 35	7 17	15 23	13 48	1 41	11 11	14 25	7 54	29
30	1 9	3 51	1 13	5 57	2 20	7 28	15 31	13 33	1 11	11 28	14 19	7 36	30



The Equations with +, are to be added to the apparent time, to have the mean time; those with -, are to be subtracted from apparent time, to give the mean time.

The preceding mark, whether + or -, at the top of any column, belongs to all the numbers or equations in that column till the sign changes; after which, the remainder of the column belongs to the contrary sign.

The Equation answering to any point of longitude between one degree and another, or any number of minutes or parts of a degree, is to be found by proportion in the usual way, viz, as 1° or 60', to that number of minutes, so is the whole difference in the Equation from the given whole degree of longitude to the next degree, to the proportional part of it answering to the given number of minutes.

See Tables of the Equation of Time computed for every year, in the Nautical Almanac, by a method proposed and illustrated by Dr. Maskelyne, the astronomer royal, viz, by taking the difference between the sun's true right ascension and his mean longitude, corrected by the Equation of the equinoxes in right ascension, and turning it into time at the rate of 1 minute of time to 15' of right ascension. Philos. Trans. vol. 54, p. 336.

EQUATION of a Curve, is an Equation shewing the nature of a curve by expressing the relation between any absciss and its corresponding ordinate, or else the relation of their fluxions, &c. Thus, the Equation to the circle, is  $ax - x^2 = y^2$ , where  $a$  is its diameter,  $x$  any absciss, or part of that diameter, and  $y$  the ordinate at that point of the diameter; the meaning being, that whatever absciss is denoted by  $x$ , then the square of its corresponding ordinate will be  $ax - x^2$ . In like manner the Equation

of the ellipse is  $\frac{p}{a} \cdot \overline{ax - x^2} = y^2$ ,

of the hyperbola is  $\frac{p}{a} \cdot \overline{ax + x^2} = y^2$ ,

of the parabola is  $- - - px = y^2$ .

Where  $a$  is an axis, and  $p$  the parameter.

And in like manner for any other curves.

This method of expressing the nature of curves by algebraical equations, was first introduced by Des Cartes, who, by thus connecting together the two sciences of algebra and geometry, made them mutually assisting to each other, and so laid the foundation of the greatest improvements that have been made in every branch of them since that time. See Des Cartes's Geometry; also Newton's Lines of the 3d Order, and many other similar works on curve lines, by several authors.

EQUATOR, in Geography, a great circle of the earth, equally distant from its two poles, and dividing it into two equal parts, or hemispheres, the northern and southern. The Equator is sometimes simply called the Line.

The circle in the heavens conceived directly over the Equator, is the Equinoctial. See EQUINOCTIAL.

The greatest height of the Equator above the horizon, is equal to the latitude of the place.

TABLE for turning Degrees and Minutes into Time, and the Contrary.

D	H	M	D	H	M	D	H	M
M	M	S	M	M	S	M	M	S
1	0	4	61	4	4	121	8	4
2	0	8	62	4	8	122	8	8
3	0	12	63	4	12	123	8	12
4	0	16	64	4	16	124	8	16
5	0	20	65	4	20	125	8	20
6	0	24	66	4	24	126	8	24
7	0	28	67	4	28	127	8	28
8	0	32	68	4	32	128	8	32
9	0	36	69	4	36	129	8	36
10	0	40	70	4	40	130	8	40
11	0	44	71	4	44	131	8	44
12	0	48	72	4	48	132	8	48
13	0	52	73	4	52	133	8	52
14	0	56	74	4	56	134	8	56
15	1	0	75	5	0	135	9	0
16	1	4	76	5	4	136	9	4
17	1	8	77	5	8	137	9	8
18	1	12	78	5	12	138	9	12
19	1	16	79	5	16	139	9	16
20	1	20	80	5	20	140	9	20
21	1	24	81	5	24	141	9	24
22	1	28	82	5	28	142	9	28
23	1	32	83	5	32	143	9	32
24	1	36	84	5	36	144	9	36
25	1	40	85	5	40	145	9	40
26	1	44	86	5	44	146	9	44
27	1	48	87	5	48	147	9	48
28	1	52	88	5	52	148	9	52
29	1	56	89	5	56	149	9	56
30	2	0	90	6	0	150	10	0
31	2	4	91	6	4	151	10	4
32	2	8	92	6	8	152	10	8
33	2	12	93	6	12	153	10	12
34	2	16	94	6	16	154	10	16
35	2	20	95	6	20	155	10	20
36	2	24	96	6	24	156	10	24
37	2	28	97	6	28	157	10	28
38	2	32	98	6	32	158	10	32
39	2	36	99	6	36	159	10	36
40	2	40	100	6	40	160	10	40
41	2	44	101	6	44	161	10	44
42	2	48	102	6	48	162	10	48
43	2	52	103	6	52	163	10	52
44	2	56	104	6	56	164	10	56
45	3	0	105	7	0	165	11	0
46	3	4	106	7	4	166	11	4
47	3	8	107	7	8	167	11	8
48	3	12	108	7	12	168	11	12
49	3	16	109	7	16	169	11	16
50	3	20	110	7	20	170	11	20
51	3	24	111	7	24	171	11	24
52	3	28	112	7	28	172	11	28
53	3	32	113	7	32	173	11	32
54	3	36	114	7	36	174	11	36
55	3	40	115	7	40	175	11	40
56	3	44	116	7	44	176	11	44
57	3	48	117	7	48	177	11	48
58	3	52	118	7	52	178	11	52
59	3	56	119	7	56	179	11	56
60	4	0	120	8	0	180	12	0



As one whole revolution of the earth, or of the  $360^\circ$  of the Equator, is performed in 24 hours, which is at the rate of  $15^\circ$  to the hour; hence the number of degrees of the Equator, answering to any other given time, or the time answering to any given degrees of the Equator, will be easily found by proportion, viz,

as 1hr. :  $15^\circ$  :: any time : its degrees,

or as  $15^\circ$  : 1hr. :: any degrees : their time.

And thus is computed the foregoing Table for turning time into degrees of the Equator, and the contrary.

**EQUATORIAL, Universal, or PORTABLE OBSERVATORY**, is an instrument intended to answer a number of useful purposes in practical astronomy, independent of any particular observatory. It may be employed in any steady room or place, and it performs most of the useful problems in the science of astronomy. The following is the description of one lately invented, and named the *Universal Equatorial*.

The principal parts of this instrument (fig. 2, plate viii.) are, 1st, The azimuth or horizontal circle A, which represents the horizon of the place, and moves on a long axis B, called the vertical axis. 2d, The Equatorial or hour-circle C, representing the Equator, placed at right angles to the polar axis D, or the axis of the earth, upon which it moves. 3d, The semicircle of declination E, on which the telescope is placed, and moving on the axis of declination, or the axis of motion of the line of collimation F. Which circles are measured and divided as in the following Table :

Measures of the several circles and divisions on them:	Radius Inches	Limb divided to	Non. of 30 gives seconds	Divid. on limb into pts. of Inc.	Divid. by Non. into pts. of Inc.
Azimuth or horizontal circle	5.1	15'	30"	45th	1350th
Equatorial or hour circle	5.1	15', or 1 m. in time	30" 2"	45th	1350th
Vertical semicircle for declination or latitude	5.	15'	30"	42d	1260th

4th, The telescope, which is an achromatic refractor with a triple object-glass, whose focal distance is 17 inches, and its aperture 2.45 inc., and it is furnished with 6 different eye-tubes; so that its magnifying powers extend from 44 to 168. The telescope in this Equatorial may be brought parallel to the polar axis, as in the figure, so as to point to the pole-star in any part of its diurnal revolution; and thus it has been observed near noon, when the sun has shone very bright. 5th, The apparatus for correcting the error in altitude occasioned by refraction, which is applied to the eye-end of the telescope, and consists of a slide G moving in a groove or dove-tail, and carrying the several eye-tubes of the telescope, on which slide there is an index corresponding

to five small divisions engraved on the dove-tail; a very small circle, called the refraction circle H, moveable by a finger-screw at the extremity of the eye-end of the telescope; which circle is divided into half minutes, one whole revolution of it being equal to  $3' 18''$ , and by its motion it raises the centre of the cross-hairs on a circle of altitude; and also a quadrant I of  $1\frac{1}{2}$  inc. radius, with divisions on each side, one expressing the degree of altitude of the object viewed, and the other expressing the minutes and seconds of error occasioned by refraction, corresponding to that degree of altitude. To this quadrant is joined a small round level K, which is adjusted partly by the pinion that turns the whole of this apparatus, and partly by the index of the quadrant; for which purpose the refraction circle is set to the same minute &c, which the index points to on the limb of the quadrant; and if the minute &c, given by the quadrant, exceed the  $3' 18''$  contained in one entire revolution of the refraction circle, this must be set to the excess above one or more of its entire revolutions; then the centre of the cross-hairs will appear to be raised on a circle of altitude to the additional height which the error of refraction will occasion at that altitude.

The principal adjustment in this instrument, is that of making the line of collimation to describe a portion of an hour circle in the heavens: in order to which, the azimuth circle must be truly level; the line of collimation, or some corresponding line represented by the small brass rod M parallel to it, must be perpendicular to the axis of its own proper motion; and this last axis must be perpendicular to the polar axis. On the brass rod M there is occasionally placed a hanging level N, the use of which will appear in the following adjustments:

The azimuth circle may be made level by turning the instrument till one of the levels be parallel to an imaginary line joining two of the feet screws; then adjust that level with these two feet screws; turn the circle  $180^\circ$ , or half round; and if the bubble be not then right, correct half the error by the screw belonging to the level, and the other half error by the two foot screws, repeating this operation till the bubble come right; then turn the circle  $90^\circ$  from the two former positions, and set the bubble right, if it be wrong, by the foot screw at the end of the level; when this is done, adjust the other level by its own screw, and the azimuth circle will be truly level. The hanging level must then be fixed to the brass rod by two hooks of equal length, and made truly parallel to it: for this purpose, make the polar axis perpendicular or nearly perpendicular to the horizon; then adjust the level by the pinion of the declination semicircle: reverse the level, and if it be wrong, correct half the error by a small steel screw that lies under one end of the level, and the other half error by the pinion of the declination semicircle, repeating the operation till the bubble be right in both positions. To make the brass rod, on which the level is suspended, at right angles to the axis of motion of the telescope, or line of collimation, make the polar axis horizontal, or nearly so; set the declination semicircle to  $0^\circ$ , and turn the hour-circle till the bubble come right; then turn the declination-circle



circle to  $90^\circ$ ; adjust the bubble by raising or depressing the polar axis (first by hand till it be nearly right, afterwards tighten with an ivory key the socket which runs on the arch with the polar axis, and then apply the same ivory key to the adjusting screw at the end of the said arch till the bubble come quite right); then turn the declination-circle to the opposite  $90^\circ$ ; if the level be not then right, correct half the error by the aforefaid adjusting screw at the end of the arch, and the other half error by the two screws that raise or depress the end of the brass rod. The polar axis remaining nearly horizontal as before, and the declination-femicircle at  $0^\circ$ , adjust the bubble by the hour-circle; then turn the declination-femicircle to  $90^\circ$ , and adjust the bubble by raising or depressing the polar axis; then turn the hour-circle 12 hours; and if the bubble be wrong, correct half the error by the polar axis, and the other half error by the two pair of capstan screws at the feet of the two supports on one side of the axis of motion of the telescope; and thus this axis will be at right angles to the polar axis. The next adjustment, is to make the centre of the cross hairs remain on the same object, while the eye-tube is turned quite round by the pinion of the refraction apparatus: for this adjustment, set the index on the slide to the first division on the dove-tail; and set the division marked 18" on the refraction-circle to its index; then look through the telescope, and with the pinion turn the eye-tube quite round; then if the centre of the hairs does not remain on the same spot during that revolution, it must be corrected by the four small screws, 2 and 2 at a time, which will be found upon unscrewing the nearest end of the eye-tube that contains the first eye-glass; repeating this correction till the centre of the hairs remain on the spot looked at during a whole revolution. To make the line of collimation parallel to the brass rod on which the level hangs, set the polar axis horizontal, and the declination-circle to  $90^\circ$ , adjust the level by the polar axis; look through the telescope on some distant horizontal object, covered by the centre of the cross hairs: then invert the telescope, which is done by turning the hour-circle half round; and if the centre of the cross hairs does not cover the same object as before, correct half the error by the uppermost and lowermost of the 4 small screws at the eye-end of the large tube of the telescope; this correction will give a second object now covered by the centre of the hairs, which must be adopted instead of the first object; then invert the telescope as before; and if the second object be not covered by the centre of the hairs, correct half the error by the same two screws as were used before: this correction will give a third object, now covered by the centre of the hairs, which must be adopted instead of the second object; repeat this operation till no error remain; then set the hour-circle exactly to 12 hours, the declination-circle remaining a  $90^\circ$  as before; and if the centre of the cross hairs do not cover the last object fixed on, set it to that object by the two remaining small screws at the eye-end of the large tube, and then the line of collimation will be parallel to the brass rod. For rectifying the nonius of the declination and Equatorial circles, lower the telescope as many degrees &c below  $0^\circ$  or  $\text{Æ}$  on the decli-

nation-femicircle as are equal to the complement of the latitude; then elevate the polar axis till the bubble be horizontal; and thus the Equatorial circle will be elevated to the co-latitude of the place: set this circle to 6 hours; adjust the level by the pinion of the declination-circle; then turn the Equatorial circle exactly 12 hours from the last position; and if the level be not right, correct one half of the error by the Equatorial circle, and the other half by the declination-circle: then turn the Equatorial circle back again exactly 12 hours from the last position; and if the level be still wrong, repeat the correction as before, till it be right, when turned to either position: that being done, set the nonius of the Equatorial circle exactly to 6 hours, and the nonius of the declination-circle exactly to  $0^\circ$ .

The chief uses of this Equatorial are,

1st, To find the meridian by one observation only: for this purpose, elevate the Equatorial circle to the co-latitude of the place, and set the declination-femicircle to the sun's declination for the day and hour of the day required; then move the azimuth and hour-circles both at the same time, either in the same or contrary direction, till you bring the centre of the cross hairs in the telescope exactly to cover the centre of the sun; when that is done, the index of the hour-circle will give the apparent or solar time at the instant of observation; and thus the time is gained, though the sun be at a distance from the meridian; then turn the hour-circle till the index points precisely at 12 o'clock, and lower the telescope to the horizon, in order to observe some point there in the centre of the glass; and that point is the meridian mark, found by one observation only. The best time for this operation is 3 hours before, or 3 hours after 12 at noon.

2d, To point the telescope on a star, though not on the meridian, in full day-light. Having elevated the equatorial circle to the co-latitude of the place, and set the declination-femicircle to the star's declination, move the index of the hour-circle till it shall point to the precise time at which the star is then distant from the meridian, found in the tables of the right ascension of the stars, and the star will then appear in the glass.

Besides these uses, peculiar to this instrument, it may also be applied to all the purposes to which the principal astronomical instruments are applied; such as a transit instrument, a quadrant, and an equal-altitude instrument.

See the description and drawing of an Equatorial telescope, or portable observatory, invented by Mr. Short, in the *Philos. Trans.* number 493, or vol. 46, p. 242; and another by Mr. Nairne, vol. 61, p. 107.

**EQUIANGULAR Figure**, is one that has all its angles equal among themselves; as the square, and all the regular figures.

An equilateral figure inscribed in a circle, is always Equiangular. But an Equiangular figure inscribed in a circle, is not always equilateral, except when it has an odd number of sides: If the sides be of an even number, then they may either be all equal, or else half of them will always be equal to each other, and the other half to each other, the equals being placed alternately. See the demonstration in my *Mathematical Miscellany*, pa. 272.



**EQUIANGULAR**, is also said of any two figures of the same kind, when each angle of the one is equal to a corresponding angle in the other, whether each figure, separately considered in itself, be an equiangular figure or not, that is, having all its angles equal to each other. Thus, two triangles are Equiangular to each other, if, ex. gr. one angle in each be of  $30^\circ$ , a second angle in each of  $50^\circ$ , and the third angle of each equal to  $100^\circ$  degrees.

Equiangular triangles have not their like sides necessarily equal, but only proportional to each other; and such triangles are always similar to each other.

**EQUICRURAL Triangle**, is one that has two of its sides equal to each other; and is more usually called an Isosceles triangle.

**EQUICULUS**, **EQUULEUS**, or **EQUUS Minor**, a constellation of the northern hemisphere. See **EQUULEUS**.

**EQUIDIFFERENT**, are such things as have equal differences, or arithmetically proportional. If the terms have all the same difference, viz, the 1st and 2d, the 2d and 3d, the 3d and 4th, &c, they are said to be Continually Equidifferent; as the numbers 3, 6, 9, 12, &c, where the common difference is 3. But if the several different couplets only have the same difference, as the 1st and 2d, the 3d and 4th, the 5th and 6th, &c, they are said to be Discretely Equidifferent; as the terms 3 and 6, 7 and 10, 9 and 12, &c. See **ARITHMETICAL Progression** and **Proportion**.

**EQUILATERAL Figure**, is one that has all its sides equal to each other. Such as the square, and all the regular figures or polygons, or a triangle that has all its sides equal. See **EQUIANGULAR**.

**EQUILATERAL Hyperbola**, is that which has the two axes equal to each other, and every pair of conjugate diameters also equal to each other. The asymptotes also are at right angles to each other, and make each half a right angle with either axis. Also, such an hyperbola is equal to its opposite hyperbola, and likewise to its conjugate hyperbola; so that all the four conjugate hyperbolas are mutually equal to each other.

Moreover, as the 3d proportional to the two axes is the parameter, therefore in such a figure, the parameter and two axes are all three equal to one another. Hence, as the general equation to hyperbolas is

$y^2 = \frac{p}{t} \cdot tx + x^2$  or  $= \frac{c^2}{t^2} \cdot tx + x^2$ , where  $t$  is the transverse axis,  $c$  the conjugate,  $p$  the parameter,  $x$  the absciss, and  $y$  the ordinate; then making  $t$ ,  $c$ , and  $p$  all equal, the equation, for the Equilateral Hyperbola, becomes  $y^2 = tx + x^2$ ; differing from the equation of the circle only in the sign of the term  $x^2$ , which in the circle is  $-$ .

**EQUILIBRIUM**, is an equality between two equal forces acting in opposite directions; so that they mutually balance each other; like the two equal arms, or scales, of a balance, &c.

**EQUILIBRIUM**, in solid bodies, forms a considerable part of the science of Statics. And Equilibrium of fluids, a considerable part of the doctrine of Hydrostatics.

**EQUIMULTIPLES**, the products of quantities equally multiplied.

Thus,  $3a$  and  $3b$  are Equimultiples of  $a$  and  $b$ ; and  $ma$  and  $mb$  are Equimultiples of  $a$  and  $b$ .

Equimultiples of any quantities, have the same ratio as the quantities themselves.

Thus  $a : b :: 3a : 3b :: ma : mb$ .

**EQUINOCTIAL**, a great circle in the heavens under which the equator moves in its diurnal motion. The poles of this circle are the poles of the world. It divides the sphere into two equal parts, the northern and southern. It cuts the horizon, of any place, in the east and west points; and at the meridian its elevation above the horizon is equal to the co-latitude of the place. The Equinoctial has also various other properties; as,

1. Whenever the sun comes to this circle, he makes equal days and nights all round the globe; because he then rises due east, and sets due west. Hence it has the name Equinoctial. All stars which are under this circle, or have no declination, do also rise due east, and set due west.

2. All people living under this circle, or upon the equator, or line, have their days and nights at all times equal to each other.

3. From this circle, on the globe, is counted, upon the meridian, the declination in the heavens, and the latitude on the earth.

4. Upon the Equinoctial, or equator, is counted the longitude, making in all  $360^\circ$  quite round, or else  $180^\circ$  east, and  $180^\circ$  west.

5. And as the time of one whole revolution is divided into 24 hours; therefore 1 hour answers to  $15^\circ$ , or the 24th part of  $360^\circ$ . Hence,

$1^\circ$  of longitude answers to 4 min. of time,  
 $15'$  - - - - - to 1 min. of time,  
 $1'$  - - - - - to 4 sec. of time, &c.

6. The shadows of those people who live under this circle are cast to the southward of them for one half of the year, and to the northward of them during the other half; and twice in a year, viz, at the time of the equinoxes, the sun at noon casts no shadow, being exactly in their zenith.

**EQUINOCTIAL Colure**, is the great circle passing through the poles of the world and the Equinoctial points, or first points of Aries and Libra.

**EQUINOCTIAL Dial**, is one whose plane is parallel to the Equinoctial. The properties or principles of this dial are,

1. The hour lines are all equally distant from one another, quite round the circumference of a circle; and the style is a straight pin, or wire, set up in the centre of the circle, perpendicular to the plane of the dial.

2. The sun shines upon the upper part of this dial-plane from the 21st of March to the 22d of September, and upon the under part of the plane the other half of the year.

Some of these dials are made of brass, &c; and set up in a frame, to be elevated to any given latitude.

**EQUINOCTIAL Points**, are the two opposite points where the Ecliptic and Equinoctial cross each other; the



the one point being in the beginning of Aries, and called the Vernal point, or Vernal Equinox; and the other in the beginning of Libra, and called the Autumnal point, or Autumnal Equinox.

It is found by observation, that the Equinoctial points, and all the other points of the ecliptic, are continually moving backwards, or in antecedentia, i. e. westwards. This retrograde motion of the Equinoctial points, is that phenomenon called the Precession of the Equinoxes, and is made at the rate of 50'' every year nearly.

**EQUINOXES**, the times when the sun enters the Equinoctial points, or about the 21st of March and 22d of September: the former being the Vernal or Spring Equinox, and the latter time the Autumnal Equinox.

As the sun's motion is unequal, being sometimes quicker and sometimes slower, it hence happens that there are about 8 days more from the vernal to the autumnal Equinox, or while the sun is on the northern side of the equator, than while he is in moving through the southern signs from the autumnal to the vernal Equinox, or on the southern side of the equator. According to the observations of M. Cassini, the sun is 186<sup>d</sup> 14<sup>h</sup> 53<sup>m</sup> in the northern signs, and only 178 14 56 in the southern signs, so that 7 23 57 is the difference of them, or nearly 8 days.

**EQUINUS Barbatus**, a kind of comet. See **HIPPEUS**.

**EQUULEUS**, **EQUICULUS**, and **EQUUS Minor**, *Equi sedio*, the Horse's Head, one of the 48 old constellations, in the northern hemisphere. Its stars, in Ptolemy's catalogue, are 4, in Tycho's 4, in Hevelius's 6, and in Flamsteed's 10.

**ERECT Vision**. See **VISION**.

**ERECT Dials**, such as stand perpendicular to the horizon, and are of various kinds; as Erect Direct, when they face exactly one of the four cardinal points, east, west, north, south; and Erect Declining, when declined from the cardinal points. See **DIAL**.

To **ERECT a Perpendicular**, is a popular problem in practical geometry, and denotes to raise a perpendicular from a given line &c, as distinguished from Demitting or letting a perpendicular fall upon a line &c, from some point out of it. See **PERPENDICULAR**.

**ERIDANUS**, the *River*, a constellation of the southern hemisphere, and one of the 48 old asterisms. The stars in this constellation, in Ptolemy's catalogue, are 34, in Tycho's 19, and in the British catalogue 84.

**ERRATIC**, an epithet applied to the planets, which are called Erratic or wandering stars, in contradistinction to the fixed stars.

**ESCALADE**, or **SCALADE**, a furious attack of a wall or a rampart; carried on with ladders, to pass the ditch, or mount the rampart; without proceeding in form, breaking ground, or carrying on regular works to secure the men.

**ESPAULE**, or **EPAULE**. See **EPAULE**.

**ESPLANADE**, in Fortification, called also *Glacis*, a part which serves as a parapet to the counterscarp, or covert way; being a declivity or slope of earth, commencing from the top of the counterscarp, and losing itself insensibly in the level of the champaign.

**Esplanade** also means the ground which has been levelled from the glacis of the counterscarp, to the first houses; or the vacant space between the works and the houses of the town.

The term is also applied, in the general, to any piece of ground that is made flat or level, and which before had some eminence that incommoded the place.

**ESTIVAL** *Occident, Orient, or Solstice*. See **OCCIDENT, ORIENT, SOLSTICE**.

**EVAPORATION**, the act of dissipating the humidity of a body in fumes or vapour; differing from exhalation, which is properly a dispersion of dry particles issuing from a body.

Evaporation is usually produced by heat, and by the change of air: thus, common salt is formed by evaporating all the humidity in the brine or salt water; which evaporation is either performed by the heat of the sun, as in the salt-works on the sea-coast, &c; or by means of fire, as at the salt-springs, &c: and it is well known how useful a brisk wind is in drying wet clothes, or the surface of the ground; while in a calm, still atmosphere, they dry extreme slowly.

But, though Evaporation be generally considered as an effect of the heat and motion of the air, yet M. Gauteron, in the *Memoires de l'Acad. des Scienc.* an. 1705, shews, that a quite opposite cause may have the same effect, and that fluids lose more of their parts in the severest frost than when the air is moderately warm: thus, in the great frost of the year 1708, he found that the greater the cold, the more considerable the evaporation; and that ice itself lost full as much as the warmer liquors that did not freeze.

There are indeed few subjects of philosophical investigation that have occasioned a greater variety of opinion than the theory of Evaporation, or of the ascent of water in such a fluid as air, between 8 and 9 hundred times lighter than itself, to different heights according to the different densities of the air; in which case it must be specifically lighter than the air through which it ascends. The Cartesians account for it by supposing, that by the action of the sun upon the water, small particles of the water are formed into hollow spheres and filled with the *materia subtilis*, which renders them specifically lighter than the ambient air, so that they are buoyed up by it.

Dr. Nieuwentyt, in his *Religious Philosopher*, cont. 19, and several others, have alleged, that the sun emits particles of fire which adhere to those of water, and form molecule, or small bodies, lighter than an equal bulk of air, which consequently ascend till they come to a height where the air is of the same specific gravity with themselves; and that these particles being separated from the fire with which they are incorporated, coalesce and descend in dew or rain.

Dr. Halley has advanced another hypothesis, which has been more generally received: he imagined, that by the action of the sun on the surface of the water, the aqueous particles are formed into hollow spherules, that are filled with a finer air highly rarefied, so as to become specifically lighter than the external air. *Philos. Trans.* number 192, or *Abr.* vol. 2, p. 126.

Dr. Defaguliers, dissatisfied with these two hypotheses, proposes another in the *Philos. Trans.* number 407, or *Abr.* vol. 7, pa. 61. See also his *Course of Experimental*



Experimental Philosophy, vol. 2, p. 336. He supposes that heat acts more powerfully on water than on common air; that the same degree of heat which rarefies air two-thirds, will rarefy water near 14,000 times; and that a very small degree of heat will raise a steam or vapour from water, even in winter, whilst it condenses the air; and thus the particles of water are converted into vapour by being made to repel each other strongly, and, deriving electricity from the particles of air to which they are contiguous, are repelled by them and by each other, so as to form a fluid which, being lighter than the air, rises in it, according to their relative gravities. The particles of this vapour retain their repellent force for a considerable time, till, by some diminution of the density of the air in which they float, they are precipitated downwards, and brought within the sphere of each other's attraction of cohesion, and so join again into drops of water.

Many objections have been urged against this opinion, by Mr. Clare in his *Treatise of the Motion of Fluids*, pa. 294, and by Mr. Rowning in his *System of Philosophy*, part 2, diss. 6; to which Dr. Hamilton has added the two following, viz, that if heat were the only cause of evaporation, water would evaporate faster in a warm close room, than when exposed in a colder place, where there is a constant current of air; which is contrary to experience; and that the evaporation of water is so far from depending on its being rarefied by heat, that it is carried on even whilst water is condensed by the coldness of the air, till it freezes; and since it evaporates even when frozen into hard ice, it must also evaporate in all the lesser degrees of cold. And therefore heat does not seem to be the principal, much less the only cause of Evaporation.

Others have more successfully accounted for the phenomena of Evaporation on another principle, viz that of solution; and shewn, from a variety of experiments, that what we call Evaporation, is nothing more than a gradual solution of water in air, produced and supported by the same means, viz, attraction, heat, and motion, by which other solutions are effected.

It seems the Abbé Nollet first started this opinion, though without much pursuing it, in his *Leçons de Physique Experimentale*, first published in 1743: he offers it as a conjecture, that the air of the atmosphere serves as a solvent or sponge, with regard to the bodies that encompass it, and receives into its pores the vapours and exhalations that are detached from the masses to which they belong in a fluid state; and he accounts for their ascent on the same principles with the ascent of liquors in capillary tubes. On his hypothesis, the condensation of the air contributes, like the squeezing of a sponge, to their descent.

Dr. Franklin, in a paper of *Philosophical and Meteorological Observations, Conjectures and Suppositions*, delivered to the Royal Society about the year 1747, and read in 1756, suggested a similar hypothesis: he observes, that air and water mutually attract each other; and hence he concludes, that water will dissolve in air, as salt in water; every particle of air assuming one or more particles of water; and when too much is added, it precipitates in rain. But as there is not the same contiguity between the particles of air as of water, the solution of water in air is not carried on without

a motion of the air, so as to cause a fresh accession of dry particles. A small degree of heat so weakens the cohesion of the particles of water, that those on the surface easily quit it, and adhere to the particles of air: a greater degree of heat is necessary to break the cohesion between water and air; for its particles being by heat repelled to a greater distance from each other, thereby more easily keep the particles of water that are annexed to them from running into cohesions that would obstruct, refract, or reflect the heat: and hence it happens that when we breathe in warm air, though the same quantity of moisture may be taken up from the lungs as when we breathe in cold air, yet that moisture is not so visible. On these principles he accounts for the production and different appearances of fogs, mists, and clouds. He adds, that if the particles of water bring electrical fire when they attach themselves to air, the repulsion between the particles of water electrified, joins with the natural repulsion of the air to force its particles to a greater distance, so that the air being more dilated, rises and carries up with it the water; which mutual repulsion of the particles of air is increased by a mixture of common fire in the particles of water. When air, loaded with surrounding particles of water, is compressed by adverse winds, or by being driven against mountains, &c, or condensed by taking away the fire that assisted it in expanding, the particles will approach one another, and the air with its water will descend as a dew; or if the water surrounding one particle of air come in contact with the water surrounding another, they coalesce and form a drop, producing rain; and since it is a well-known fact, that vapour is a good conductor of electricity, as well as of common fire, it is reasonable to conclude with Mr. Henley, that Evaporation is one great cause of the clouds becoming at times surcharged with this fluid. *Philos. Trans.* vol. 67, pa. 134. See also vol. 55, p. 182, or Franklin's *Letters and Papers on Philosophical Subjects*, p. 42 &c, and pa. 182, ed. 1769.

M. le Roi, of the Acad. of Sciences at Paris, has also advanced the same opinion, and supported it by a variety of facts and observations in the *Memoirs* for the year 1751. He shews, that water does undergo in the air a real dissolution, forming with it a transparent mixture, and possessing the same properties with the solutions of most salts in water; and that the two principal causes which promote the solution of water in the air, are heat and wind; that the hotter the air is, within a certain limit, the more water it will dissolve; and that at a certain degree of heat the air will be saturated with water; and by determining at different times the degree of the air's saturation, he estimates the influence of those causes on which the quantity depends that is suspended in the air in a state of solution. Accordingly, the air, heated by evaporating substances to which it is contiguous, becomes more rare and light, rises and gives way to a denser air; and, by being thus removed, contributes to accelerate the Evaporation. The fixed air contained in the internal parts of evaporating bodies, put into action by heat, seems also to increase their Evaporation. The wind is another cause of the increase of Evaporation, chiefly by changing and renewing the air which immediately encompasses the evaporating substances; and from the consideration of



of these two causes combined, it appears why the quantity of vapour raised in the night is less than that of the day, since the air is then both less heated and less agitated. To the objection urged against this hypothesis, on account of the Evaporation of water in a vacuum, this ingenious writer replies, that the water itself contains a great quantity of air, which gradually disengages itself, and causes the Evaporation; and that it is impossible that a space containing water which evaporates should remain perfectly free from air. To this objection a late writer, Dr. Dobson of Liverpool, replies, that though air appears, by unquestionable experiments, to be a chemical solvent of water, and as such, is to be considered as one cause of its Evaporation, heat is another cause, acting without the intervention of air, and producing a copious Evaporation in an exhausted receiver; agreeably to an experiment of Dr. Irving, who says, that in an exhausted receiver water rises in vapour more copiously at  $180^{\circ}$  of Fahrenheit's thermometer, than in the open air at  $212^{\circ}$ , its boiling point. Dr. Dobson farther adds, that water may exist in air in three different states; in a state of perfect solution, when the air will be clear, dry, and heavy, and its powers of solution still active; in a state of beginning precipitation, when it becomes moist and foggy, its powers of solution are diminished, and it becomes lighter in proportion as its water is deposited; and also, when it is completely precipitated, which may happen either by a slower process, when the dissolved water falls in a drizzling rain, or by a more sudden process, when it descends in brisk showers. *Philos. Transf.* vol. 67, p. 257, and Phipps's Voyage towards the North Pole, p. 211.

Dr. Hamilton, professor of philosophy in the university of Dublin, transmitted to the Royal Society in 1765, a long Dissertation on the nature of Evaporation, in which he proposes and establishes this theory of solution; and though other writers had been prior in their conjectures, and even in their reasoning on this subject, Dr. Hamilton assures us, that he has not represented any thing as new which he was conscious had ever been proposed by any one before him, even as a conjecture. Dr. Hamilton having evinced the agreement between Solution and Evaporation, concludes, that Evaporation is nothing more than a gradual solution of water in air, produced and promoted by attraction, heat, and motion, just as other solutions are effected.

To account for the ascent of aqueous vapours into the atmosphere, this ingenious writer observes, that the lowest part of the air being pressed by the weight of the upper against the surface of the water, and continually rubbing upon it by its motion, attracts and dissolves those particles with which it is in contact, and separates them from the rest of the water. And since the cause of solution in this case is the stronger attraction of the particles of water towards the air, than towards each other, those that are already dissolved and taken up, will be still farther raised by the attraction of the dry air that lies over them, and thus will diffuse themselves, rising gradually higher and higher, and so leave the lowest air not so much saturated but that it will still be able to dissolve and take up fresh particles of water; which process is greatly promoted by

the motion of the wind. When the vapours are thus raised and carried by the winds into the higher and colder parts of the atmosphere, some of them will coalesce into small particles, which slightly attracting each other, and being intermixed with air, will form clouds; and these clouds will float at different heights, according to the quantity of vapour borne up, and the degree of heat in the upper parts of the atmosphere: and thus clouds are generally higher in summer than in winter. When the clouds are much increased by a continual addition of vapours, and their particles are driven close together by the force of the winds, they will run into drops heavy enough to fall down in rain. If the clouds be frozen before their particles are gathered into drops, small pieces of them being condensed and made heavier by the cold, fall down in thin flakes of snow. When the particles are formed into drops before they are frozen, they become hailstones. When the air is replete with vapours, and a cold breeze springs up, which checks the solution of them, clouds are formed in the lower parts of the atmosphere, and compose a mist or fog, which usually happens in a cold morning, and is dispersed when the sun has warmed the air, and made it capable of dissolving these watry particles. Southerly winds commonly bring rain, because, being warm and replete with aqueous vapours, they are cooled by coming into a colder climate; and therefore they part with some of them, and suffer them to precipitate in rain: whereas northerly winds, being cold, and acquiring additional heat by coming into a warmer climate, are ready to dissolve and receive more vapour than they before contained; and therefore, by long continuance, they are dry and parching, and commonly attended with fair weather.

Changes of the air, with respect to its density and rarity, as well as its heat and cold, will produce contrary effects in the solution of water, and the consequent ascent or fall of vapours. Several experiments prove that air, when rarefied, cannot keep so much water dissolved as it does in a more condensed state; and therefore when the atmosphere is saturated with water, and changes from a denser to a rarer state, the high and colder parts of it will let go some of the water before dissolved, forming new clouds, and disposing them to fall down in rain: but a change from a rarer to a denser state will stop the precipitation of the water, and enable the air to dissolve, either in whole or in part, some of those clouds that were formed before, and render their particles less apt to run into drops and fall down in rain: on this account, we generally find that the rarefied and condensed states of the atmosphere are respectively attended with rain or fair weather. See more on this subject in the *Philos. Transf.* vol. 55, pa. 146, or Hamilton's Philosophical Essays, p. 33.

Dr. Halley, before mentioned, has furnished some experiments on the Evaporation of water; the result of which is contained in the following articles: 1. That water salted to about the same degree as sea-water, and exposed to a heat equal to that of a summer's day, did, from a circular surface of about 8 inches diameter, evaporate at the rate of 6 ounces in 24 hours: whence by a calculus he finds that, in such circumstances, the water evaporates 1-10th of an inch deep in 12 hours: which



which quantity, he observes, will be found abundantly sufficient to furnish all the rains, springs, dews, &c. By this experiment, every 10 square inches of surface of the water yield in vapour *per diem* a cubic inch of water: and each square foot half a wine pint; every space of 4 feet square, a gallon; a mile square, 6914 tuns; and a square degree, of 69 English miles, will evaporate 33 millions of tuns a day; and the whole Mediterranean, computed to contain 160 square degrees, at least 5280 millions of tuns each day. *Philos. Transf.* number 189, or *Abridg.* vol. 2, pa. 108.—2. A surface of 8 square inches, evaporated purely by the natural warmth of the weather, without either wind or sun, in the course of a whole year, 16292 grains of water, or 64 cubic inches; consequently, the depth of water thus evaporated in one year, amounts to 8 inches. But this being too little to answer the experiments of the French, who found that it rained 19 inches of water in one year at Paris; or those of Mr. Townley, who found the annual quantity of rain in Lancashire above 40 inches; he concludes, that the sun and wind contribute more to Evaporation than any internal heat or agitation of the water. In effect, Dr. Halley fixes the annual Evaporation of London at 48 inches; and Dr. Dobson states the same for Liverpool at  $36\frac{1}{4}$  inches. *Philos. Transf.* vol. 67, p. 252.

3. The effect of the wind is very considerable, on a double account; for the same observations shew a very odd quality in the vapours of water, viz, that of adhering and hanging to the surface that exhaled them, which they clothe as it were with a fleece of vaporous air; which once investing the vapour, it afterwards rises in much less quantity. Whence, the quantity of water lost in 24 hours, when the air is very still, was very small, in proportion to what went off when there was a strong gale of wind abroad to dissipate the fleece, and make room for the emission of vapour; and this, even though the experiment was made in a place as close from the wind as could be contrived. Add, that this fleece of water, hanging to the surface of waters in still weather, is the occasion of very strange appearances, by the refraction of the vapours differing from and exceeding that of common air: whence every thing appears raised, as houses like steeples, ships as on land above the water, the land raised, and as it were lifted from the sea, &c.

4. The same experiments shew that the Evaporation in May, June, July, and August, which are nearly equal, are about three times as great as those in the months of November, December, January, and February. *Philos. Transf.* numb. 212, or *Abr.* vol. 2, pa. 110.

Dr. Brownrigg, in his *Art of making common salt*, pa. 189, fixes the Evaporation of some parts of England at 73.8 inches during the months of May, June, July, and August; and the Evaporation of the whole year at more than 140 inches. But the Evaporation of the four summer months at Liverpool, on a medium of 4 years, was found to be only 18.88 inches. Also Dr. Hales calculates the greatest annual Evaporation from the surface of the earth in England at 6.66 inches; and therefore the annual Evaporation from a surface of water, is to the annual Evaporation from the surface of the earth at Liverpool, nearly as 6 to 1. *Philos. Transf.* vol. 67, ubi supra.

In the *Transactions of the American Philosophical Society*, vol. 3, pa. 125, there is an ingenious paper on Evaporation, by Dr. Wistar. It is there shewn, that evaporation arises when the moist body is warmer than the medium it is inclosed in. And, on the contrary, it acquires moisture from the air, when the body is the colder. This carrying off, and acquiring of moisture, it is shewn, is by the passage of heat out of the body, or into it.

EUCLID, of Megara, a celebrated philosopher and logician; he was a disciple of Socrates, and flourished about 400 years before Christ. The Athenians having prohibited the Megarians from entering their city on pain of death, this philosopher disguised himself in women's clothes to attend the lectures of Socrates. After the death of Socrates, Plato and other philosophers went to Euclid at Megara, to shelter themselves from the tyrants who governed Athens.—This philosopher admitted but one chief good; which he at different times called *God*, or the *Spirit*, or *Providence*.

EUCLID, the celebrated mathematician, according to the account of Pappus and Proclus, was born at Alexandria, in Egypt, where he flourished and taught mathematics, with great applause, under the reign of Ptolemy Lagos, about 280 years before Christ. And here, from his time, till the conquest of Alexandria by the Saracens, all the eminent mathematicians were either born, or studied; and it is to Euclid, and his scholars, we are beholden for Eratosthenes, Archimedes, Apollonius, Ptolemy, Theon, &c, &c. He reduced into regularity and order all the fundamental principles of pure mathematics, which had been delivered down by Thales, Pythagoras, Eudoxus, and other mathematicians before him, and added many others of his own discovering: on which account it is said he was the first who reduced arithmetic and geometry into the form of a science. He likewise applied himself to the study of mixed mathematics, particularly to astronomy and optics.

His works, as we learn from Pappus and Proclus, are the *Elements*, *Data*, *Introduction to Harmony*, *Phenomena*, *Optics*, *Catoptrics*, a *Treatise of the Division of Superficies*, *Porisms*, *Loci ad Superficiem*, *Fallacies*, and *Four books of Conics*. The most celebrated of these, is the first work, the *Elements of Geometry*; of which there have been numberless editions, in all languages; and a fine edition of all his works, now extant, was printed in 1703, by David Gregory, Savilian professor of astronomy at Oxford.

The *Elements*, as commonly published, consist of 15 books, of which the two last it is suspected are not Euclid's, but a comment of Hypsicles of Alexandria, who lived 200 years after Euclid. They are divided into three parts, viz, the Contemplation of Superficies, Numbers, and Solids: The first 4 books treat of planes only; the 5th of the proportions of magnitudes in general; the 6th of the proportion of plane figures; the 7th, 8th, and 9th give us the fundamental properties of numbers; the 10th contains the theory of commensurable and incommensurable lines and spaces; the 11th, 12th, 13th, 14th, and 15th, treat of the doctrine of solids.

There is no doubt but, before Euclid, *Elements of Geometry* were compiled by Hippocrates of Chius, Eudoxus,



**Eudoxus**, Leon, and many others, mentioned by Proclus, in the beginning of his second book ; for he affirms that Euclid new ordered many things in the Elements of Eudoxus, completed many things in those of Theætetus, and besides strengthened such propositions as before were too slightly, or but superficially established, with the most firm and convincing demonstrations.

History is silent as to the time of Euclid's death, or his age. He is represented as a person of a courteous and agreeable behaviour, and in great esteem and familiarity with king Ptolomy ; who once asking him, whether there was any shorter way of coming at geometry than by his Elements, Euclid, as Proclus testifies, made answer, that there was no royal way or path to geometry.

**EUUDIOMETER**, an instrument for determining the purity of the air, or the quantity of pure and dephlogisticated or vital air contained in it, chiefly by means of its diminution on a mixture with nitrous air.

Instruments of this kind have been but lately made, and that in consequence of the experiments and discoveries of Dr. Priestley, for determining the salubrity of different kinds of air. That writer having discovered, that when nitrous air is mixed with any other air, their original bulk is diminished ; and that the diminution is nearly, if not exactly, in proportion to its salubrity ; he was hence led to adopt nitrous air as a true test of the purity of respirable air ; and nothing more seemed to be necessary but an easy, expeditious, and accurate method of estimating the degree of diminution in different cases ; and for this purpose, the Eudiometer was contrived ; of which several kinds have been invented, the principal of which are the following.

I. The Eudiometer originally used by Dr. Priestley is a divided glass tube, into which, after having filled it with common water, and inverted it into the same, one measure or more of common air, and an equal quantity of nitrous air, are introduced by means of a small phial, which is called the measure ; and thus the diminution of the volume of the mixture, which is seen at once by means of the graduations of the tube, instantly discovers the purity of the air required.

II. The discovery of Dr. Priestley was announced to the public in the year 1772 ; and several persons, both at home and abroad, presently availed themselves of it, by framing other more accurate instruments. The first of these was contrived by M. Landriani ; an account of which is published in the 6th volume of *Rosier's Journal*, for the year 1775. It consists of a glass tube, fitted by grinding to a cylindrical vessel, to which are joined two glass cocks and a small basin ; the whole being fitted to a wooden frame. In this instrument quicksilver is used instead of water ; though that is attended with an inconvenience, because the nitrous air acts upon the metal, and renders the experiment ambiguous.

III. In 1777, Mr. Magellan published an account of three Eudiometers invented by himself, consisting of glass vessels of rather difficult construction, and trouble some use. Mr. Cavallo observes, that the construction of all the three is founded on a supposition, that the mixture of nitrous and atmospheric air, having continued for some time to diminish, afterwards increases

again ; which it seems is a mistake : neither do they give accurate or uniform results in any two experiments made with nitrous and common air of precisely the same quality.

IV. A preferable method of discovering the purity of the air by means of an Eudiometer, is recommended by M. Fontana, of very great accuracy. The instrument is originally nothing more than a divided glass tube, though the inventor afterwards added to it a complicated apparatus, perhaps of little or no use. The first simple Eudiometer consisted only of a glass tube, uniformly cylindrical, about 18 inches long, and  $\frac{3}{4}$ ths of an inch diameter within side, the outside being marked with a diamond at such distances as are exactly filled by equal measures of elastic fluids : and when any parts of these divisions are required, the edge of a ruler, divided into inches and smaller parts, is held against the tube, so as that the first division of the ruler may coincide with one of the marks on the tube. The nitrous and atmospheric air are introduced into this tube, in order to be diminished, and thence the purity of the atmospheric air ascertained.

V. M. Saussure of Geneva has also invented an Eudiometer, which he thinks is more exact than any of those before described ; the apparatus of which is as follows : 1. A cylindrical glass bottle, with a ground stopple, containing about  $5\frac{1}{2}$  ounces, which serves as a receiver for mixing the two airs.—2. A small glass phial, to serve as a measure, and is about one-third the size of the receiver.—3. A small pair of scales that may weigh very exactly.—4. Several glass bottles, for containing the nitrous or other air to be used, and which may supply the place of the recipient when broken. The method of using it is as follows : The receiver is to be filled with water, closed exactly with its glass stopper, wiped dry on the outside, and then weighed very nicely. Being then immersed in a vessel of water, and held with the mouth downwards, the stopple is removed, and, by means of a funnel, two measures of common and one of nitrous air are introduced into it, one after another : these diminish as soon as they come into contact ; in consequence of which the water enters the recipient in proportionable quantity. After being stopped and well shaken, to promote the diminution, the receiver is to be opened again under water ; then stopped and shaken again, and so on for three times successively, after which the bottle is stopped for the last time under water, then taken out, wiped very clean and dry, and exactly weighed as before. It is plain that now, the bottle being filled partly with elastic fluid and partly with water, it must be lighter than when quite full of water ; and the difference between those two weights, shows nearly what quantity of water would fill the space occupied by the diminished elastic fluid. Now, in making experiments with airs of different degrees of purity, the said difference will be greater when the diminution is less, or when the air is less pure, and vice versa ; by which means the comparative purity between two different kinds of air is determined.

VI. But as this method, notwithstanding the encomiums bestowed on it by the inventor, is subject to several errors and inconveniences ; to remedy all these, another instrument was invented by Mr. Cavallo ; the description



tion of which, being long, may be seen in his *Treatise on the Nature and Properties of Air*, pa. 344.

Other constructions of the Eudiometer have also been given by Mr. Cavendish and Mr. Scheele. For farther information, see Magellan's Letter to Dr. Priestley, containing the Description of a Glass Apparatus, &c, and of New Eudiometers &c, 1777, pa. 15 &c; Priestley's Exp. and Obs. on Air, vol. 3, preface and appendix; the methods of Dr. Ingenhoufz in *Philos. Transf.* vol. 66, art. 15; see also the *Philos. Transf.* vol. 73; and Cavallo's *Treatise on Air*, pa. 274, 315, 316, 317, 328, 340, 344, and 834.

EUDOXUS, of Cnidus, a city of Caria in Asia Minor, flourished about 370 years before Christ. He learned geometry from Archytas, and afterwards travelled into Egypt to learn astronomy and other sciences. There he and Plato studied together, as Laertius informs us, for the space of 13 years; and afterwards came to Athens, fraught with all sorts of knowledge, which they had imbibed from the mouths of the priests. Here Eudoxus opened a school; which he supported with so much glory and renown, that even Plato, though his friend, is said to have envied him. Eudoxus composed *Elements of Geometry*, from whence Euclid liberally borrowed, as mentioned by Proclus. Cicero calls Eudoxus the greatest astronomer that had ever lived: and Petronius says, he spent the latter part of his life upon the top of a very high mountain, that he might contemplate the stars and the heavens with more convenience and less interruption: and we learn from Strabo, that there were some remains of his observatory at Cnidus, to be seen even in his time. He died in the 53d year of his age.

EVECTION, is used by some astronomers for the Libration of the moon; being an inequality in her motion, by which, at or near the quadratures, she is not in a line drawn through the centre of the earth to the sun, as she is at the syzygies, or conjunction and opposition, but makes an angle with that line of about  $2^{\circ} 51'$ . The motion of the moon about her axis only is equable, which rotation is performed exactly in the same time as she revolves about the earth; for which reason it is that she turns always the same face towards the earth nearly, and would do so exactly were it not that her menstrual motion about the earth, in an elliptic orbit, is not equable; on which account the moon, seen from the earth, appears to librate a little upon her axis, sometimes from east to west, and sometimes from west to east; or some parts in the eastern limb of the moon go backwards and forwards a small space, and some that were conspicuous, are hid, and then appear again.

The term EVECTION is used by some astronomers to denote that equation of the moon's motion, which is proportional to the sine of double the distance of the moon from the sun, diminished by the moon's anomaly: this equation is not yet accurately determined; some state it at  $1^{\circ} 30'$ , others at  $1^{\circ} 16'$ , &c. It is the greatest of all the moon's equations, except the Equation of the Centre.

EVEN *Number*, is that which can be divided into two equal whole numbers; such as the series of alternate numbers 2, 4, 6, 8, 10, &c.

EVENLY *Even Number*, is that which an even num-

ber measures by an even number; as 16, which the even number 8 measures by the even number 2.

EVENLY *Odd Number*, is that which an even number measures by an odd one; as 30, which the even number 6 measures by the odd number 5.

EVERARD's *Sliding Rule*, a particular sort of one invented by Mr. Thomas Everard, for the purpose of gauging. See SLIDING RULE.

EULER (LEONARD), one of the most extraordinary, and even prodigious, mathematical geniuses, that the world ever produced. He was a native of Basil, and was born April 15, 1707. The years of his infancy were passed at Riehen, where his father was minister. He was afterwards sent to the university of Basil; and as his memory was prodigious, and his application regular, he performed his academical tasks with great rapidity; and all the time that he saved by this, was consecrated to the study of mathematics, which soon became his favourite science. The early progress he made in this study, added fresh ardour to his application; by which too he obtained a distinguished place in the attention and esteem of professor John Bernoulli, who was then one of the chief mathematicians in Europe.

In 1723, M. Euler took his degree as master of arts; and delivered on that occasion a Latin discourse, in which he drew a comparison between the philosophy of Newton and the Cartesian system, which was received with the greatest applause. At his father's desire, he next applied himself to the study of theology and the oriental languages: and though these studies were foreign to his predominant propensity, his success was considerable even in this line: however, with his father's consent, he afterward returned to mathematics as his principal object. In continuing to avail himself of the counsels and instructions of M. Bernoulli, he contracted an intimate friendship with his two sons Nicholas and Daniel; and it was chiefly in consequence of these connections that he afterwards became the principal ornament of the philosophical world.

The project of erecting an academy at Petersburg, which had been formed by Peter the Great, was executed by Catharine the 1st; and the two young Bernoullis being invited to Petersburg in 1725, promised Euler, who was desirous of following them, that they would use their endeavours to procure for him an advantageous settlement in that city. In the mean time, by their advice, he made close application to the study of philosophy, to which he made happy applications of his mathematical knowledge, in a dissertation on the nature and propagation of sound, and an answer to a prize question concerning the masting of ships; to which the Academy of Sciences adjudged the *accessit*, or second rank, in the year 1727. From this latter discourse, and other circumstances, it appears that Euler had very early embarked in the curious and useful study of naval architecture, which he afterward enriched with so many valuable discoveries. The study of mathematics and philosophy however did not solely engage his attention, as he in the mean time attended the medical and botanical lectures of the professors at Basil.

Euler's merit would have given him an easy admission to honourable preferment either in the magistracy or university of his native city, if both civil and academic



mical honours had not been there distributed by lot. The lot being against him in a certain promotion, he left his country, set out for Petersburg, and was made joint professor with his countrymen Hermann and Daniel Bernoulli in the university of that city.

At his first setting out in his new career, he enriched the academical collection with many memoirs, which excited a noble emulation between him and the Bernoullis; an emulation that always continued, without either degenerating into a selfish jealousy, or producing the least alteration in their friendship. It was at this time that he carried to new degrees of perfection the integral calculus, invented the calculation by fines, reduced analytical operations to a greater simplicity, and thus was enabled to throw new light on all the parts of mathematical science.

In 1730, M. Euler was promoted to the professorship of natural philosophy; and in 1733 he succeeded his friend D. Bernoulli in the mathematical chair. In 1735, a problem was proposed by the academy, which required expedition, and for the calculation of which some eminent mathematicians had demanded the space of some months. The problem was undertaken by Euler, who completed the calculation in three days, to the great astonishment of the academy: but the violent and laborious efforts it cost him threw him into a fever, which endangered his life, and deprived him of the use of his right eye, which afterward brought on a total blindness.

The Academy of Sciences at Paris, which in 1738 had adjudged the prize to his memoir Concerning the Nature and Properties of Fire, proposed for the year 1740 the important subject of the Tides of the Sea; a problem whose solution comprehended the theory of the solar system, and required the most arduous calculations. Euler's solution of this question was adjudged a master-piece of analysis and geometry; and it was more honourable for him to share the academical prize with such illustrious competitors as Colin Maclaurin and Daniel Bernoulli, than to have carried it away from rivals of less magnitude. Seldom, if ever, did such a brilliant competition adorn the annals of the academy; and perhaps no subject, proposed by that learned body, was ever treated with such force of genius and accuracy of investigation, as that which here displayed the philosophical powers of this extraordinary triumvirate.

In the year 1741, M. Euler was invited to Berlin to direct and assist the academy that was there rising into fame. On this occasion he enriched the last volume of the Miscellanies (*Melanges*) of Berlin with five memoirs, which form an eminent, perhaps the principal, figure in that collection. These were followed, with amazing rapidity, by a great number of important researches, which are dispersed through the memoirs of the Prussian academy; a volume of which has been regularly published every year since its establishment in 1744. The labours of Euler will appear more especially astonishing, when it is considered, that while he was enriching the academy of Berlin with a profusion of memoirs, on the deepest parts of mathematical science, containing always some new points of view, often sublime truths, and sometimes discoveries of great importance; he still continued his philosophical contributions to the Petersburg academy, whose memoirs display the

marvellous fecundity of his genius, and which granted him a pension in 1742.

It was with great difficulty that this extraordinary man, in 1766, obtained permission from the king of Prussia to return to Petersburg, where he wished to pass the remainder of his days. Soon after his return, which was graciously rewarded by the munificence of Catharine the 2d, he was seized with a violent disorder, which ended in the total loss of his sight. A cataract, formed in his left eye, which had been essentially damaged by the loss of the other eye, and a too close application to study, deprived him entirely of the use of that organ. It was in this distressing situation that he dictated to his servant, a tailor's apprentice, who was absolutely devoid of mathematical knowledge, his *Elements of Algebra*; which by their intrinsic merit in point of perspicuity and method, and the unhappy circumstances in which they were composed, have equally excited wonder and applause. This work, though purely elementary, plainly discovers the proofs of an inventive genius; and it is perhaps here alone that we meet with a complete theory of the analysis of Diophantus.

About this time M. Euler was honoured by the Academy of Sciences at Paris with the place of one of the foreign members of that learned body; after which, the academical prize was adjudged to three of his memoirs, *Concerning the Inequalities in the Motions of the Planets*. The two prize questions proposed by the same Academy for 1770 and 1772 were designed to obtain from the labours of astronomers *a more perfect Theory of the Moon*. M. Euler, assisted by his eldest son, was a competitor for these prizes, and obtained them both. In this last memoir, he reserved for farther consideration several inequalities of the moon's motion, which he could not determine in his first theory, on account of the complicated calculations in which the method he then employed had engaged him. He afterward revised his whole theory, with the assistance of his son and Messrs Krafft and Lexell, and pursued his researches till he had constructed the new tables, which appeared, together with the great work, in 1772. Instead of confining himself, as before, to the fruitless integration of three differential equations of the second degree, which are furnished by mathematical principles, he reduced them to the three ordinates, which determine the place of the moon: he divided into classes all the inequalities of that planet, as far as they depend either on the elongation of the sun and moon, or upon the eccentricity, or the parallax, or the inclination of the lunar orbit. All these means of investigation, employed with such art and dexterity as could only be expected from a genius of the first order, were attended with the greatest success; and it is impossible to observe without admiration, such immense calculations on the one hand, and on the other the ingenious methods employed by this great man to abridge them, and to facilitate their application to the real motion of the moon. But this admiration will become astonishing, when we consider at what period and in what circumstances all this was effectuated. It was when our author was totally blind, and consequently obliged to arrange all his computations by the sole powers of his memory and his genius: it was when he was embarrassed in his domestic affairs by a dreadful fire,



that had consumed great part of his substance, and forced him to quit a ruined house, every corner of which was known to him by habit, which in some measure supplied the want of sight. It was in these circumstances that Euler composed a work which alone was sufficient to render his name immortal.

Some time after this, the famous oculist Wentzell, by couching the cataract, restored sight to our author ; but the joy produced by this operation was of short duration. Some instances of negligence on the part of his surgeons, and his own impatience to use an organ, whose cure was not completely finished, deprived him a second time and for ever of his sight : a relapse which was also accompanied with tormenting pain. With the assistance of his sons, however, and of Messrs Krafst and Lexell, he continued his labours : neither the infirmities of old age, nor the loss of his sight, could quell the ardour of his genius. He had engaged to furnish the academy of Petersburg with as many memoirs as would be sufficient to complete its acts for 20 years after his death. In the space of 7 years he transmitted to the Academy above 70 memoirs, and above 200 more, left behind him, were revised and completed by a friend. Such of these memoirs as were of ancient date were separated from the rest, and form a collection that was published in the year 1783, under the title of *Analytical Works*.

The general knowledge of our author was more extensive than could well be expected in one who had pursued, with such unremitting ardour, mathematics and astronomy as his favourite studies. He had made a very considerable progress in medical, botanical, and chemical science. What was still more extraordinary, he was an excellent scholar, and possessed in a high degree what is generally called *erudition*. He had attentively read the most eminent writers of ancient Rome; the civil and literary history of all ages and all nations was familiar to him; and foreigners, who were only acquainted with his works, were astonished to find in the conversation of a man, whose long life seemed solely occupied in mathematical and physical researches and discoveries, such an extensive acquaintance with the most interesting branches of literature. In this respect, no doubt, he was much indebted to a very uncommon memory, which seemed to retain every idea that was conveyed to it, either from reading or from meditation. He could repeat the *Æneid* of Virgil, from the beginning to the end, without hesitation, and indicate the first and last line of every page of the edition he used.

Several attacks of a vertigo, in the beginning of September, 1783, which did not prevent his computing the motions of the aerostatic globes, were however the forerunners of his mild passage out of this life. While he was amusing himself at tea with one of his grandchildren, he was struck with an apoplexy, which terminated his illustrious career at 76 years of age.


M. Euler's constitution was uncommonly strong and vigorous. His health was good; and the evening of his long life was calm and serene, sweetened by the fame that follows genius, the public esteem and respect that are never withheld from exemplary virtue, and several domestic comforts which he was capable of feeling, and therefore deserved to enjoy.

The catalogue of his works has been printed in 50 pages, 14 of which contain the manuscript works.—The printed ones consist of works published separately, and works to be found in the memoirs of several Academies, viz, in 38 volumes of the Petersburg Acts, (from 6 to 10 papers in each volume);—in several volumes of the Paris Acts;—in 26 volumes of the Berlin Acts, (about 5 papers to each volume);—in the Acta Eruditorum, in 2 volumes;—in the Miscellanea Taurinensia;—in vol. 9 of the Society of Ulyssingue;—in the Ephemerides of Berlin;—and in the Memoires de la Société Oeconomique for 1766.

EVOLVENT, in the Higher Geometry, a term used by some writers for the Involute, or curve resulting from the evolution of a curve, in contradistinction to that evolute, or curve supposed to be opened or evolved. See EVOLUTE, and INVOLUTE.

EVOLUTE, in the Higher Geometry, a curve first proposed by M. Huygens, and since much studied by the later mathematicians. It is any curve supposed to be evolved or opened, by having a thread wrapped close upon it, fastened at one end, and beginning to evolve or unwind the thread from the other end, keeping the part evolved, or wound off, tight stretched; then this end of the thread will describe another curve called the Involute. Or the same involute is described the contrary way, by wrapping the thread upon the Evolute, keeping it always stretched.

Thus, if EFGH be any curve, and AE either a part of the curve, or a right line; then if a thread be wound close upon the curve from A to H, where it is fixed, and then be unwound from A; the curve AEFHG, from which it is evolved, is called the Evolute; and the other curve ABCD described by the end of the thread, as it evolves or unwinds, is the Involute. Or, if the thread HD, fixed at H, be wound or wrapped upon the Evolute HGFEA, keeping it always tight, as at the several positions of it HD, GC, FB, EA, the extremity will describe the Involute curve DCBA.



From this description it appears, 1. That the parts of the thread at any positions, as EA, FB, GC, HD, &c, are radii of curvature, or osculatory radii, of the involute curve, at the points A, B, C, D.

2. The same parts of the thread are also equal to the corresponding lengths  $AE$ ,  $AEF$ ,  $AEFG$ , &c, of the Evolute; that is,

$\text{AE} = \text{AF}$  is the rad. of curvature to the point A,  
 $\text{BF} = \text{AF}$  " " " " B,  
 $\text{CG} = \text{AG}$  " " " " C,  
 $\text{DH} = \text{AH}$  " " " " D.

3. Any



3. Any radius of curvature BF, is perpendicular to the involute at the point B, and is a tangent to the Evolute curve at the point F.

4. The Evolute is the locus of the centre of curvature of the involute curve.

The finding the radii of Evolutes, is a matter of great importance in the higher speculations of geometry; and is even sometimes useful in practice; as is shewn by Huygens, the inventor of this theory, in applying it to the pendulum. Horol. Oscil. part 3. The doctrine of the Oscula of Evolutes is owing to M. Leibnitz, who first shewed the use of Evolutes in the measuring of curvatures.

To find the EVOLUTE and Involute Curves, the one from the other.

For this purpose, put

$x = AD$  the absciss of the involute,

$y = DB$  its ordinate,

$z = AB$  the involute curve,

$r = BC$  its radius of curvature,

$v = EF$  the absciss of the Evolute,

$u = FC$  its ordinate, and

$a = AE$  a given line, (fig. 2 above).

Then, by the nature of the radius of curvature, it is

$$r = \frac{z^3}{y\dot{x} - x\dot{y}} = BC = AE + EC; \text{ also by sim. triangles,}$$

$$\dot{z} : \dot{x} :: r : GB = \frac{r\dot{x}}{z} = \frac{\dot{x}\dot{z}^2}{y\dot{x} - x\dot{y}},$$

$$\dot{z} : \dot{y} :: r : GC = \frac{r\dot{y}}{\dot{z}} = \frac{y\dot{z}^2}{y\dot{x} - x\dot{y}}.$$

$$\text{Hence } EF = GB - DB = \frac{\dot{x}\dot{z}^2}{y\dot{x} - x\dot{y}} - y = v; \text{ and}$$

$$FC = AD - AE + GC = x - a + \frac{y\dot{z}^2}{y\dot{x} - x\dot{y}} = u;$$

which are the values of the absciss and ordinate of the Evolute curve EC; and therefore these may be found when the involute is given.

On the other hand, if  $v$  and  $u$ , or the Evolute be given: then, putting the given curve  $EC = s$ ; since  $CB = AE + EC$ , or  $r = a + s$ , this gives  $r$  the radius of curvature. Also, by similar triangles, there result these proportions, viz,

$$\dot{s} : \dot{v} :: r : \frac{r\dot{v}}{s} = \frac{a + s}{s} \dot{v} = GB,$$

$$\dot{s} : \dot{u} :: r : \frac{r\dot{u}}{s} = \frac{a + s}{s} \dot{u} = GC; \text{ theref.}$$

$$AD = AE + FC - GC = a + u - \frac{a + s}{s} \dot{u} = x,$$

$$\text{and } DB = GB - EF = \frac{a + s}{s} \dot{v} - v = y;$$

which are the absciss and ordinate of the involute curve, and which may therefore be found when the Evolute is given. Where it may be noted that  $\dot{s}^2 = \dot{v}^2 + \dot{u}^2$ , and  $\dot{z}^2 = \dot{x}^2 + \dot{y}^2$ . Also either of the quantities  $x, y$ , may be supposed to flow equably, in which case the respective second fluxion  $\ddot{x}$  or  $\ddot{y}$  will be nothing, and the corresponding term in the denominator  $y\dot{x} - x\dot{y}$  will vanish, leaving only the other term in it; which

will have the effect of rendering the whole operation simpler.

For Ex. Suppose it were required to find the Evolution EC when the given involute AB is the common parabola, whose equation is  $px = y^2$ , the parameter being  $p$ .

Here  $y = \sqrt{px}$ ,  $\dot{y} = \frac{1}{2}p^{\frac{1}{2}}x^{-\frac{1}{2}}\dot{x}$ , and  $\ddot{y} = -\frac{1}{4}p^{\frac{1}{2}}x^{-\frac{3}{2}}\dot{x}$ , making  $\ddot{x} = 0$ . Then, to find first AE the radius of curvature of the parabola AB at the vertex, when  $\dot{x} = 0$ , the general value of the radius of curvature

$$\text{above given becomes } r = \frac{z^3}{-x\ddot{y}} = \frac{\overline{\dot{x}^2 + \dot{y}^2}^{\frac{3}{2}}}{-x\ddot{y}} = (\text{by sub-}$$

$$\text{stituting the value of } \dot{y} \text{ and } \ddot{y} \text{ \&c}) \frac{p + 4x}{2\sqrt{p}}$$

which is the general value of  $r$  or BC, the radius of curvature, for any value of  $x$  or AD; and when  $x$  or AD is  $= 0$  or nothing, the value of  $r$ , or AE, becomes then

$$a = \frac{p^{\frac{3}{2}}}{2\sqrt{p}} = \frac{1}{2}p \text{ only; that is half the parameter of the axis is the radius of curvature at the vertex of the parabola.}$$

Again, in the general values of  $v$  and  $u$  above given, by substituting the values of  $\dot{y}$ ,  $\ddot{y}$ , and  $\dot{z}$ , also 0 for  $\ddot{x}$ , and  $\frac{1}{2}p$  for  $a$ ; those quantities become

$$v = \frac{\dot{z}^2}{-\ddot{y}} - y = \frac{\dot{x}^2 + \dot{y}^2}{-\ddot{y}} - y = \frac{4x^{\frac{3}{2}}}{p^{\frac{1}{2}}} = 4x\sqrt{\frac{x}{p}}; \text{ and}$$

$$u = x - a + \frac{y\dot{z}^2}{-x\ddot{y}} = 3x + \frac{1}{2}p - a = 3x.$$

Hence then, comparing the values of  $v$  and  $u$ , there is found  $3p^{\frac{1}{2}}v = 4x^{\frac{1}{2}}u$ , and  $27pv^2 = 16u^3$ ; which is the equation between the absciss and ordinate of the Evolute curve EC, shewing it to be the semicubical parabola.

In like manner the Evolute to any other curve is found.—The Evolute to the common cycloid, is an equal cycloid; a property first demonstrated by Huygens, and which he used as a contrivance to make a pendulum vibrate in the curve of a cycloid. See his Horolog. Oscil. See also, on the subject of Evolute and Involute Curves, the Fluxions of Newton, Maclaurin, Simpson, De l'Hôpital, &c, Wolf. Elem. Math. tom. I, &c, &c.

M. Varignon has applied the doctrine of the radius of the Evolute, to that of central forces; so that having the radius of the Evolute of any curve, there may be found the value of the central force of a body; which, moving in that curve, is found in the same point where that radius terminates; or reciprocally, having the central force given, the radius of the Evolute may be determined. Hist. de l'Acad. an. 1706.

The variation of curvature of the line described by the Evolution of a curve, is measured by the ratio of the radius of curvature of the Evolute, to the radius of curvature of the line described by the Evolution. See Maclaurin's Flux. art. 402, prop. 36.

Imperfect EVOLUTE, a name given by M. Reaumur to a new kind of Evolute. The mathematicians had hitherto only considered the perpendiculars let fall from the Involute on the convex side of the Evolute: but if other



other lines not perpendicular be drawn upon the same points, provided they be all drawn under the same angle, the effect will still be the same; that is, the oblique lines will all intersect in the curve, and by their intersections form the infinitely small sides of a new curve, to which they would be so many tangents.—Such a curve is a kind of Evolute, and has its radii; but it is an Imperfect one, since the radii are not perpendicular to the first curve, or Involute. Hist. de l'Acad. &c, an. 1709.

EVOLUTION, in Arithmetic and Algebra, denotes the Extraction of the roots out of powers. In which sense it stands opposed to Involution, which is the raising of powers. The note or character that has been used by some Algebraists, to denote Evolution, is  $\omega$ ; as the sign of involution is  $\oslash$ : characters I think first used by Dr. Pell.

EVOLUTION, in Geometry, the opening, or unfolding of a curve, and making it describe an Evolvent.

The equable Evolution of the periphery of a circle, or other curve, is such a gradual approach of the circumference to rectitude, as that its parts do all concur, and equally evolve or unbend; so that the same line becomes successively a less arc of a reciprocally greater circle; till at last they change into a straight line.—In the Philos. Trans. N<sup>o</sup> 260, a new quadratrix to the circle is found by this means, being the curve described by the equable Evolution of its periphery.

EURYTHMY, in Architecture, Painting, and Sculpture, is a kind of majesty, elegance, and easiness appearing in the composition of certain members or parts of a body, building, or painting, and resulting from the fine and exact proportions of them.

EUSTYLE, is the best manner of placing columns, with regard to their distance; which, according to Vitruvius, should be four modules, or two diameters and a quarter.

EXAGON. See HEXAGON.

EXALTATION, in Astrology, is a dignity which a planet acquires in certain signs of the zodiac; which dignity, it is supposed, gives the planet an extraordinary virtue, efficacy, and influence. The opposite side of the zodiac is called the Dejection of the planet.—Thus, the 15th degree of Cancer is the Exaltation of Jupiter, according to Albumazar, because it was the ascendant of that planet at the time of the creation; that of the sun is in the 19th degree of Aries; and its dejection in Libra; that of the moon is in Taurus, &c. Ptolemy gives the reason of this in his first book De Quadripartita.


EXCENTRIC, is applied to such figures, circles, spheres, &c, as have not the same centre; as opposed to Concentric, which have the same centre.

EXCENTRIC, or *Excentric Circle*, in the ancient Ptolemaic astronomy, was the very orbit of the planet itself, which it was supposed to describe about the earth, and which was conceived Excentric with it; called also the Deferent.

Instead of these Excentric Circles round the earth, the moderns make the planets describe elliptic orbits about the sun; which accounts for all the irregularities of their motions, and their various distances from the earth, &c, more justly and naturally.

EXCENTRIC, or *Excentric Circle*, in the New Astronomy, is the circle described from the centre of the orbit of a planet, with half the greatest axis as a radius; or it is the circle that circumscribes the elliptic orbit of the planet; as the circle AQB.

EXCENTRIC Anomaly, or *Anomaly of the Centre*, is an arc AQ of the Excentric circle, intercepted between the aphelion A, and the right line QH, drawn through the centre P of the planet perpendicular to the line of the apses AB.



EXCENTRIC Equation, in the Old Astronomy, is an angle made by a line drawn from the centre of the earth, with another line drawn from the centre of the Excentric, to the body or place of any planet. This is the same with the prosthapheresis; and is equal to the difference, accounted in an arch of the ecliptic, between the real and apparent place of the sun or planet. See EQUATION of the Centre.

EXCENTRIC Place of a planet, in its orbit, is the Heliocentric place, or that in which it appears as seen from the sun.

EXCENTRIC Place in the ecliptic, is the point of the ecliptic to which the planet is referred as viewed from the sun; and which coincides with the heliocentric longitude.

EXCENTRICITY, is the distance between the centres of two circles, or spheres, which have not the same centre.

EXCENTRICITY, in the Old Astronomy, is the distance between the centre of a planet and the centre of the earth.—That the planets have such an Excentricity, is allowed on all sides, and may be evinced from various circumstances; and especially this, that the planets at some times appear larger, and at others less; which can only proceed from hence, that their orbits being Excentric to the earth, in some parts of those orbits the planets are nearer to us, and in others more remote. And as to the Excentricities of the sun and moon, it is thought they are sufficiently proved, both from eclipses, from the moon's greater and less parallax at the same distance from the zenith, and from the sun's continuing longer by 8 days in the northern hemisphere than in the southern one.

EXCENTRICITY, in the New Astronomy, is the distance CS between the sun S and the centre C of a planet's orbit; or the distance of the centre from the focus of the elliptic orbit; called also the *Simple* or *Single Excentricity*.

When the greatest Equation of the centre is given, the Excentricity of the earth's orbit may be found by the following proportion; viz,

As the diameter of a circle in degrees,

Is to the diameter in equal parts;

So the greatest equat. of the centre in degrees,

To the Excentricity in equal parts. Thus,

Greatest equat. of the cent.  $1^{\circ} 55' 33'' = 1^{\circ} 9258333$  &c.

The diam. of a circ. being 1, its circumf. is  $3.1415926$ .

Then  $3.1415926 : 1 :: 360^{\circ} : 114^{\circ} 59' 15609$  diam. in deg.

And  $114.5915609 : 1 :: 1.9258333 : 0.016806$ , the Ex

Hence,



Hence, by adding this to 1, and subtracting it from 1, gives  $1.016806 = AS$  the aphelion distance, and  $0.983194 = BS$  the perihelion distance.

See Robertfon's Elem. of Navig. book 5, pa. 286.

*Otherwise*, thus: Since it is found that the sun's greatest apparent semi-diameter is to his least, as  $32' 43''$  to  $31' 38''$ , or as  $1963''$  to  $1898''$ ; the sun's greatest distance from the earth will be to his least, or AS to SB, as  $1963$  to  $1898$ ; of which,

the half dif. is  $32\frac{1}{2} = CS$ ,

and half sum  $1930\frac{1}{2} = CB$ ; wherefore,

as  $1930\frac{1}{2} : 32\frac{1}{2} :: 1 : .016835 = CS$  the Excentricity, to the mean distance or semi-axis 1; which is nearly the same as before.

The Excentricities of the orbits of the several planets, in parts of their own mean distances 1000, and also in English miles, are as below, viz, the Excentricity of the orbit of

	Parts.	Miles.
Mercury - -	210 - -	7,730,000
Venus - -	7 - -	482,000
Earth - -	17 - -	1,618,000
Mars - -	93 - -	13,486,000
Jupiter - -	48 - -	23,760,000
Saturn - -	55 - -	49,940,000
Georgian - -	$47\frac{1}{2}$ - -	86,000,000

*Double EXCENTRICITY*, is the distance between the two foci of the elliptic orbit, and is equal to double the Single Excentricity above given.

*EXCHANGE*, in Arithmetic, is the bartering or exchanging the money of one place for that of another; or the finding what quantity of the money of one place is equal to a given sum of another, according to a given course of exchange.

The several operations in this case are only different applications of the Rule of Three. See most books of Arithmetic.

*Arbitration of EXCHANGE*, is the method of remitting to, and drawing upon, foreign places, in such a manner as shall turn out the most profitable.

*Arbitration* is either Simple or Compound.

*Simple Arbitration* respects three places only. Here, by comparing the par of arbitration between a first and second place, and between the 1st and a 3d, the rate between the 2d and 3d is discovered; from whence a person can judge how to remit or draw to the most advantage, and to determine what that advantage is.

*Compound Arbitration* respects the cases in which the exchanges among three, four, or more places are concerned. A person who knows at what rate he can draw or remit directly, and also has advice of the course of exchange in foreign parts, may trace out a path for circulating his money, through more or fewer of such places, and also in such order, as to make a benefit of his skill and credit: and in this lies the great art of such negotiations. See my Arithmetic, pa. 105, &c.

*EXCURSION*, in Astronomy. See *ELONGATION*.

*Circles of EXCURSION*. See *CIRCLES*.

*EXEGESIS*, or *EXEGETICA*, in Algebra, is the finding, either in numbers or lines, the roots of the equation of a problem, according as the problem is either numeral or geometrical.

*EXHALATION*, a fume or steam Exhaling, or issuing, from a body, and diffusing itself in the atmosphere.

The terms Exhalation and Vapour are often used indifferently; but the more accurate writers distinguish them, appropriating the term Vapour to the moist fumes raised from water and other liquid bodies; and the term Exhalation to the dry ones emitted from solid bodies; as earth, fire, minerals, &c. In this sense, Exhalations are dry and subtle corpuscles, or effluvia, loosened from hard terrestrial bodies, either by the heat of the sun, or the action of the air, or some other cause: being emitted upwards to a certain height in the atmosphere, where, mixing with the vapours, they help to constitute clouds, and return back in dews, mists, rains, &c.

Sir Isaac Newton thinks, that true and permanent air is formed from the Exhalations raised from the hardest and most compact bodies.

*EXHAUSTED Receiver*, is a glass, or other vessel, applied on the plate of an air-pump, to have the air extracted out of it by the working of the pump.— Things placed in such an Exhausted Receiver, are said to be *in vacuo*.

*EXHAUSTIONS*, or the *Method of EXHAUSTIONS*, a method of demonstration founded upon a kind of Exhausting a quantity by continually taking away certain parts of it.

The method of Exhaustions was of frequent use among the ancient mathematicians; as Euclid, Archimedes, &c. It is founded on what Euclid says in the 10th book of his Elements; viz, that those quantities are equal, whose difference is less than any assignable quantity. Or thus, two quantities A and B are equal, when, if to or from one of them as A, any other quantity as *d* be subtracted, however small it be, then the sum or difference is respectively greater or less than the other quantity B: viz, *d* being an indefinitely small quantity,

if  $A + d$  be greater than B,

and  $A - d$  less than B,

then is A equal to B.

This principle is used in the 1st prop. of the 10th book, which imports, that if from the greater of two quantities be taken more than its half, and from the remainder more than its half, and so on; there will at length remain a quantity less than either of those proposed. On this foundation it is demonstrated, that if a regular polygon of infinite sides be inscribed in a circle, or circumscribed about it; then the space, which is the difference between the circle and the polygon, will by degrees be quite exhausted, and the circle become ultimately equal to the polygon. And in this way it is that Archimedes demonstrates, that a circle is equal to a right-angled triangle, whose two sides about the right angle, are equal, the one to the semidiameter, and the other to the perimeter of the circle. Prop. 1 De Dimensione Circuli.

Upon the Method of Exhaustions depends the Method of Indivisibles introduced by Cavalierius, which is but a shorter way of expressing the method of Exhaustions; as also Wallis's Arithmetic of Infinites, which is a farther improvement of the Method of Indivisibles; and hence also the Methods of Increments, Differentials, Fluxions, and



and Infinite Series. See some account of the Method of Exhaustions in Wallis's Algebra, chap. 73, and in Ronayne's Algebra, part 3, pa. 395.

EXPANSION, is the dilating, stretching, or spreading out of a body; whether from any external cause, as the cause of rarefaction, or from an internal cause, as elasticity. Bodies naturally expand by heat beyond their dimensions when cold; and hence it happens that their dimensions and specific gravities are different in different temperatures and seasons of the year. Air compressed or condensed, as soon as the compressing or condensing force is removed, expands itself by its elastic power to its former dimensions.

In some few cases indeed bodies seem to expand as they grow cold, as water in the act of freezing: but it seems this is owing to the extrication of a number of air bubbles from the fluid at a certain time; and is not at all a regular and gradual expansion like that of metals, &c, by means of heat. Mr. Boyle, in his History of Cold, says that ice takes up one 12th part more space than water; but by Major Williams's experiments on the force of freezing water, I have found it occupies but about the 17th or 18th part more space. Transac. of the R. Soc. of Edinb. vol. 2, pa. 28. In certain metals also, an Expansion takes place when they pass from a fluid to a solid state: but this too is not to be accounted any proper effect of cold, but of the arrangement of the parts of the metal in a certain manner; and is therefore to be accounted a kind of crystallization, rather than any thing else.

The Expansion of different bodies by heat is very various; and many experiments upon it are to be met with in the volumes of the Philos. Transf. and elsewhere. In the 48th vol. in particular, Mr. Smeaton has given a table of the Expansion of many different substances, as determined by experiment, from which the following particulars are extracted. Where it is to be noted, that the quantities of Expansion which answer to 180 degrees of Fahrenheit's thermometer, are expressed in ten-thousandth parts of an English inch, each substance being 1 foot or 12 inches in length.

White glass barometer tube	-	-	-	100
Martial regulus of antimony	-	-	-	130
Blistered steel	-	-	-	138
Hard steel	-	-	-	147
Iron	-	-	-	151
Bismuth	-	-	-	167
Copper hammered	-	-	-	204
Copper 8 parts, mixed with 1 of tin	-	-	-	218
Cast brass	-	-	-	225
Brass 16 parts, with tin 1	-	-	-	229
Brass wire	-	-	-	232
Speculum metal	-	-	-	232
Spelter folder, viz. brass 2 parts, zink 1	-	-	-	247
Fine pewter	-	-	-	274
Grain tin	-	-	-	298
Soft folder, viz. lead 2, tin 1	-	-	-	301
Zink 8 parts, tin 1, a little hammered	-	-	-	323
Lead	-	-	-	344
Zink or Spelter	-	-	-	353
Zink hammer'd $\frac{1}{2}$ an inch per foot	-	-	-	373

By other experiments too it has been found that, for each degree of heat of the thermometer, mercury, wa-

ter, and air, expand by the following parts of their own bulk, viz,

Mercury	the 9600th	} part of its bulk.
Water	the 6666th	
Air	the 435th	

From the foregoing table it appears, that there is no general rule for the degree of Expansion to which bodies are subject by the same degree of heat, either from their specific gravity or otherwise. Zink, which is much lighter than lead, expands more with heat; while glass, which is lighter than either, expands much less; and copper, which is heavier than a mixture of brass and tin, expands less.

It seems too that metals observe a proportion of Expansion in a fluid state, quite different from what they do in a solid one: For regulus of antimony seemed to shrink in fixing, after being melted, considerably more than zink.

But of all known substances, those of the aerial kind expand most by an equal degree of heat; and in general the greater quantity of latent heat that any substance contains, the more easily is it expanded; though even here no general rule can be formed. It is indeed certain that the densest fluids, such as mercury, oil of vitriol, &c, are less expandible than water, spirit of wine, or ether. Which last is so easily expanded, that were it not for the pressure of the atmosphere it would be in a continual state of vapour. And indeed this is the case, in some measure, with perhaps all fluids; as it has been found, by experiments with the best air-pumps, that water, and other fluids, ascend in vapours the more as the exhaustion is the more perfect; from which it would seem that water would wholly rise in vapour, in any temperature, if the pressure of the atmosphere was entirely taken off.

After bodies are reduced to a vaporous state, their Expansion seems to go on without any limitation, in proportion to the degree of heat applied; though it may be impossible to say what would be the ultimate effects of that principle upon them in this way. The force with which these vapours expand on the application of high degrees, is very great; nor does it appear that any obstacle whatever is insuperable by them.

On this principle depend the steam engines, so much used in various mechanical operations; likewise some hydraulic machines; and the instruments called manometers, which shew the variation of gravity in the external atmosphere, by the expansion or condensation of a small quantity of air confined in a proper vessel. On this principle also, perpetual movements might be constructed similar to those invented by Mr. Cox, on the principle of the barometer. And a variety of other curious machines may be constructed on the principle of aerial expansion; an account of some of which is given under HYDROSTATICS and PNEUMATICS.

On the principle of the Expansion of fluids are constructed Thermometers. And for the effects of the different Expansions of metals in correcting the errors of machines for measuring time, see the article PENDULUM.

The Expansion of solid bodies is measured by an instrument called the Pyrometer; and the force with which they expand is still greater than that of aerial vapours; the flame of a farthing candle produces an Expansion



Expansion in a bar of iron capable of counteracting a weight of 500 pounds. The quantity of expansion however is so small, that it has never been applied to the movement of any mechanical engine.

EXPECTATION, in the Doctrine of Chances, is applied to any contingent event, upon the happening of which some benefit &c is expected. This is capable of being reduced to the rules of computation: for a sum of money in Expectation when a particular event happens, has a determinate value before that event happens. Thus, if a person is to receive any sum, as 10l, when an event takes place which has an equal chance or probability of happening and failing, the value of the Expectation is half that sum or 5l.: but if there are 3 chances for failing, and only 1 for its happening, or one chance only in its favour out of all the 4 chances; then the probability of its happening is only 1 out of 4, or  $\frac{1}{4}$ , and the value of the Expectation is but  $\frac{1}{4}$  of 10l. which is only 2l. 10s. or half the former sum. And in all cases, the value of the Expectation of any sum is found by multiplying that sum by the fraction expressing the probability of obtaining it. So the value of the Expectation on 100l. when there are 3 chances out of 5 for obtaining it, or when the probability of obtaining it is  $\frac{3}{5}$ , is  $\frac{3}{5}$  of 100l. which is 60l. And if  $s$  be any sum expected on the happening of an event,  $b$  the chances for that event happening, and  $f$  the chances for its failing; then, there being  $b$  chances out of  $f + b$  for its happening, the probability will be  $\frac{b}{f + b}$ , and the

value of the expectation is  $\frac{b}{f + b} \times s$ . See Simpson's or De Moivre's Doctrine of Chances.

EXPECTATION of Life, in the Doctrine of Life Annuities, is the share, or number of years of life, which a person of a given age may, upon an equality of chance, expect to enjoy.

By the Expectation or share of life, says Mr. Simpson (Select Exercices pa. 273), is not here to be understood that particular period which a person hath an equal chance of surviving; this last being a different, and more simple consideration. The Expectation of a life, to put it in the most familiar light, may be taken as the number of years at which the purchase of an annuity, granted upon it, without discount of money, ought to be valued. Which number of years will differ more or less from the period above-mentioned, according to the different degrees of mortality to which the several stages of life are incident. Thus it is much more than an equal chance, according to the table of the probability of the duration of life (p. 254 ut supra), that an infant, just come into the world, arrives not to the age of 10 years; yet the Expectation or share of life due to it, upon an average, is near 20 years. The reason of which wide difference, is the great excess of the probability of mortality in the first tender years of life, above that respecting the more mature and stronger ages. Indeed if the numbers that die at every age were to be the same, the two quantities above specified would also be equal; but when the said numbers become continually less and less, the Expectation must of consequence be the greater of the two.

VOL. I.

Mr. Simpson has given a table and rule for finding this Expectation, pa. 255 and 273 as above. Thus,

*A Table of the EXPECTATIONS of Life in London.*

Age	Expectation	Age	Expectation	Age	Expectation
1	27.0	28	24.6	55	14.2
2	32.0	29	24.1	56	13.8
3	34.0	30	23.6	57	13.4
4	35.6	31	23.1	58	13.1
5	36.0	32	22.7	59	12.7
6	36.0	33	22.3	60	12.4
7	35.8	34	21.9	61	12.0
8	35.6	35	21.5	62	11.6
9	35.2	36	21.1	63	11.2
10	34.8	37	20.7	64	10.8
11	34.3	38	20.3	65	10.5
12	33.7	39	19.9	66	10.1
13	33.1	40	19.6	67	9.8
14	32.5	41	19.2	68	9.4
15	31.9	42	18.8	69	9.1
16	31.3	43	18.5	70	8.8
17	30.7	44	18.1	71	8.4
18	30.1	45	17.8	72	8.1
19	29.5	46	17.4	73	7.8
20	28.9	47	17.0	74	7.5
21	28.3	48	16.7	75	7.2
22	27.7	49	16.3	76	6.8
23	27.2	50	16.0	77	6.4
24	26.6	51	15.6	78	6.0
25	26.1	52	15.2	79	5.5
26	25.6	53	14.9	80	5.0
27	25.1	54	14.5		

For Example, if it be required to find the Expectation or share of life, due to a person of 30 years old. Opposite the given age in the first column of the table, stands 23.6 in the second col. for the years in the Expectation sought.

See De Moivre's Doctrine of Chances applied to the Valuation of Annuities, p. 288; or Dr. Price's Observations on Reverfionary Payments, p. 168, 364, 374, &c; or Philos. Transf. vol. 59, p. 89.

EXPERIMENT, in Philosophy, a trial of the effect or result of certain applications and motions of natural bodies, in order to discover something of their laws and relations, &c.

The making of experiments is grown into a kind of art; and there are now many collections of them, mostly under the denomination of Courses of Experimental Philosophy. Sturmius made a curious collection of the principal Discoveries and Experiments of the last age, under the title of Collegium Experimentale. Other Courses of Experiments have been published by Gravefande, Defaguliers, Helsham, Cotes, and others.

EXPERIMENTAL Philosophy, is that which proceeds on Experiments, or which deduces the laws of nature and the properties and powers of bodies, and their actions upon each other, from sensible Experiments and observations.

Experiments are of the utmost importance in philosophy; and the great advantages the modern physics have



have over the ancient, is chiefly owing to this, viz, that we abound much more in Experiments, and that we make more use of the Experiments we have. The method of the ancients, was chiefly to begin with the causes of things, and thence argue to the phenomena and effects; on the contrary, that of the moderns proceeds from Experiments and Observations, from whence the properties and laws of natural things are deduced, and general theories are formed.

Several of the ancients indeed thought as highly of Experiments as the moderns, and practised them also. Plato omits no occasion of speaking of the advantages of them; and Aristotle's history of animals bears ample testimony for *him*. Democritus's great employment was to make experiments; and even Epicurus himself owes part of his glory to the same cause.

Among the moderns, the making of Experiments was chiefly begun by Friar Bacon, in the 13th century, who it seems spent a great deal of money and labour in this way. After him, the lord chancellor Bacon is looked upon as the founder of the present mode of philosophising by Experiments. And his method has been prosecuted with laudable emulation by the Academy del Cimento, the Royal Society, the Royal Academy at Paris; by Mr. Boyle, and, over all, by Sir Isaac Newton, with many other illustrious names.

Indeed, Experiments, within the last century, are come so much into vogue, that nothing will pass in philosophy, but what is either founded on Experiments, or confirmed by them; so that the new philosophy is almost wholly Experimental.

Yet there are some, even among the learned, who speak of Experiments in a different manner, or perhaps rather of the abuse of them, and in derision of the pretenders to this practice. Thus, though Dr. Keil allows that philosophy has received very considerable advantages from the makers of Experiments; yet he complains of their dissingenuity, in too often wresting and distorting their Experiments and Observations to favour some darling theories they had espoused. Nay more, M. Hartsoeker, in his *Recueil de plusieurs Pieces de Physique*, undertakes to shew, that such as employ themselves in the making of Experiments, are not properly philosophers, but as it were the labourers or operators of philosophers, that work under them, and for them, furnishing them with materials to build their systems and hypotheses upon. And the learned M. Dacier, in the beginning of his discourse on Plato, at the head of his translation of the works of that philosopher, deals still more severely with the makers of Experiments. He breaks out with a kind of indignation at a tribe of idly curious people, whose sole employment consists in making Experiments on the gravity of the air, the equilibrium of fluids, the loadstone, &c, and yet arrogate to themselves the noble title of philosophers. But his honest indignation would have exceeded all bounds, had he lived to see the contemptible fall of one of the principal societies above-mentioned; while its members first amuse themselves with magnetical conundrums, spinning electrical wheels, torturing the unseen and unknown phlogistic particles; and finally polluting the source of science, and the streams of wisdom, with the folly of hunting after cockle-shells, caterpillars, and butterflies!

EXPLOSION, a sudden and violent expansion of an elastic fluid, by which it instantly throws off any obstacle that happens to be in the way, sometimes with astonishing force and rapidity, as the Explosion of fired gun-powder, &c.

Explosion differs from expansion, in that the latter is a gradual and continued power, acting uniformly for some certain time; whereas the former is always sudden, and only of momentary or immensurably short duration. The expansions of solid substances do not terminate in violent explosions, on account of their slowness, and the small space through which the expanding substance moves; though their strength may be equally great with that of the most active aerial fluids. Thus we find that though wedges of wood, when wetted, will cleave solid blocks of stone, they never throw them to any distance, as is the case with gun-powder. On the other hand, it is seldom that the expansion of any elastic fluid bursts a solid substance without throwing the fragments of it to a considerable distance, with effects that are often very terrible.

The most part of explosive substances are either aerial, or convertible into such, and raised into an elastic fluid. Thus gun-powder, whose essence seems to consist in common air fixed in the nitre, or at least an air of similar elasticity, where it is condensed into a bulk many hundred times less than the natural state of the atmosphere; which air being suddenly disengaged by the firing of the gun-powder, and the decomposition of its parts, it rapidly expands itself again with a force proportioned to the degree of its condensation when fixed in the gun-powder, and so explodes, and produces all those terrible effects that attend the explosion. The elastic fluid generated by the fired gun-powder expands itself with a velocity of about 10,000 feet per second, and with a force more than 1000 times greater than the pressure of the atmosphere on the same base.

The Electric Explosions seem to be still much more strong and astonishing; as in the cases of lightning, earthquakes, and volcanoes; and even in the artificial electricity produced by the ordinary machines. The astonishing strength of electric explosions, which is beyond all possible means of measuring it, manifests itself by the many tremendous effects we hear of fire-balls and lightning.

In cases where the electric matter acts like common fire, the force of the explosions, though very great, is capable of measurement, by comparing the distances to which bodies are thrown, with their weight. This is most evident in volcanoes, where the projections of the burning rocks and lava manifest the greatness of the power, at the same time that they afford a method of measuring it: and these explosions are owing to the extrication of aerial vapours, and their rarefaction by intense heat.

Next in strength to the aerial vapours, are those of aqueous and other liquids. Very remarkable effects of these are observed in steam-engines; and there is one case from which it has been inferred that aqueous steam is even vastly stronger than fired gun-powder. This is when water is thrown upon melted copper: for here the explosion is so strong as almost to exceed imagination; and the most terrible accidents have happened, even from so slight a cause as one of the workmen spit-  
ting







present. Also Albert Girard, in 1629, used the Exponents to roots, thus;  $\sqrt{\quad}$ ,  $\sqrt[2]{\quad}$ ,  $\sqrt[3]{\quad}$ , &c.

The notation of powers and roots by the present way of Exponents, has introduced a new and general arithmetic of Exponents or powers; for hence powers are multiplied by only adding their Exponents, divided by subtracting the Exponents, raised to other powers, or roots of them extracted, by multiplying or dividing the Exponent by the index of the power or root.—

$$\text{So } a^2 \times a^3 = a^5, \text{ and } a^{\frac{1}{2}} \times a^{\frac{1}{4}} = a^{\frac{3}{4}};$$

$$a^5 \div a^3 = a^2, \text{ and } a^{\frac{3}{4}} \div a^{\frac{1}{4}} = a^{\frac{1}{2}};$$

the 2d power of  $a^3$  is  $a^6$ ,

and the 3d root of  $a^6$  is  $a^2$ .

This algorithm of powers led the way to the invention of logarithms, which are only the indices or Exponents of powers: and hence the addition and subtraction of logarithms, answer to the multiplication and division of numbers; while the raising of powers, and extracting of roots, is effected by multiplying the logarithm by the index of the power, or dividing the logarithm by the index of the root.

EXPONENT of a Ratio, is, by some, understood as the quotient arising from the division of the antecedent of the ratio by the consequent: in which sense, the Exponent of the ratio of 3 to 2 is  $\frac{3}{2}$ ; and that of the ratio of 2 to 3 is  $\frac{2}{3}$ .

But others, and those among the best mathematicians, understand logarithms as the Exponents of ratios; in which sense they coincide with the idea of measures of ratios, as delivered by Kepler, Mercator, Halley, Cotes, &c.

EXPONENTIAL Calculus, the method of differencing, or finding the fluxions of, Exponential quantities, and of summing up those differences, or finding their fluents. See CALCULUS, FLUXIONS, and FLUENTS.

EXPONENTIAL Curve, is that whose nature is defined or expressed by an Exponential equation; as the curve denoted by  $a^x = y$ , or by  $x^a = y$ .

EXPONENTIAL Equation, is one in which is contained an exponential quantity: as the equation  $a^x = b$ , or  $x^a = ab$ , &c.

Exponential Equations are commonly best resolved by means of logarithms, viz, first taking the log. of the given equation: thus, taking the log. of the equation  $a^x = b$ , it is  $x \times \log. \text{ of } a = \log. \text{ of } b$ ; and hence

$$x = \frac{\log. b}{\log. a}$$

Also, the log. of the equation  $x^a = ab$ , is  $x \times \log. x = \log. ab$ ; and then  $x$  is easily found by trial-and-error, or the double rule of position.

EXPONENTIAL Quantity, is that whose power is a variable quantity; as the expression  $a^x$ , or  $x^a$ .

Exponential quantities are of several degrees, and orders, according to the number of exponents or powers, one over another. Thus,

$a^x$  is an Exponential of the 1st order,

$a^{x^2}$ , is one of the 2d order,

$a^{x^3}$  is one of the 3d order, and so on.

See Bernoulli Oper. tom. 1, pa. 182, &c.

EXPRESSION, in Algebra, is any algebraical quantity, simple or compound: as the expression,  $3a$ , or  $2ab$ , or  $\sqrt{a^2 + c^2}$ .

EXTENSION, one of the common and essential properties of body; or that by which it possesses or takes up some part of universal space, called the place of that body.

The extension of a body, is properly in every direction whatever; but it is usual to consider it as extended only in length, breadth, and thickness.

EXTERIOR Polygon, or Talus, is the outer or circumscribing one. See POLYGON and TALUS.

EXTERMINATION, or EXTERMINATING, in Algebra, is the taking away, or expelling of something from an expression, or from an equation: as to Exterminate surds, fractions, or any particular letter or quantity out of equations.

Thus, to take away the fractional form from this equation  $\frac{a^2 + x^2}{2c} = \frac{ab}{d}$ ; multiply each numerator by

the other's denominator, and the equation becomes  $a^2d + dx^2 = 2abc$ , out of fractions.

Also, to take away the radicality from the equation  $3\sqrt{a^2 - x^2} = 2c$ , raise each to the 2d power, and it becomes  $9a^2 - 9x^2 = 4c^2$ .

For Exterminating any quantity out of equations, there are various rules and methods, according to the form of the equations; of which many excellent specimens may be seen in Newton's Algebra, pa. 60, ed. 1738; or in Maclaurin's Algebra, part 1, chap. 12. For example, to Exterminate  $y$  out of these two equations,  $a + x = b + y$ ,

$$\text{and } 3b = 2x + y;$$

subtract the upper equation from the under, so shall there arise  $3b - a - x = 2x - b$ ; then, by the known methods of transposition &c, there is obtained  $4b - a = 3x$ ,

$$\text{and hence } x = \frac{4b - a}{3}.$$

EXTERNAL Angles, are the angles formed with-outside of a figure, by producing its sides out.

In a triangle, any External angle is equal to the sum of both the two internal opposite angles taken together: and, in any right-lined figure, the sum of all the external angles, is always equal to 4 right angles.

EXTRA-Constellary Stars, such as are not properly included in any constellation.

EXTRA-Mundane Space, is the infinite, empty, void space, which is by some supposed to be extended beyond the bounds of the universe, and consequently in which there is really nothing at all.

EXTRACTION of Roots, is the finding the roots of given numbers, or quantities, or equations.

The roots of quantities are denominated from their powers; as the square or 2d root, the cubic or 3d root, the biquadratic or 4th root, the 5th root, &c; which are the roots of the 2d, 3d, 4th, 5th, &c powers. The Extraction of roots has always made a part of arithmetical calculation, at least as far back as the composition of powers has been known: for the composition of powers always led to their resolution, or Extraction of roots, which is performed by the rules exactly reverse of the former. Thus, if any root be considered



sidered as consisting of two parts  $a + x$ , of which the former  $a$  is known, and the latter  $x$  unknown, then the square of this root being  $a^2 + 2ax + x^2$ , which is its composition, this indicated the method of resolution, so as to find out the unknown part  $x$ ; for having subtracted the nearest square  $a^2$  from the given quantity, there remains  $2ax + x^2$  or  $2a + x \times x$ ; therefore divide this remainder by  $2a$ , the double of the first member of the root, the quotient will be nearly  $x$  the other member; then to  $2a$  add this quotient  $x$ , and multiply the sum  $2a + x$  by  $x$ , and the product will make up the remaining part  $2ax + x^2$  of the given power.

The composition of the cubic or 3d power next presented itself, which consists of these four terms  $a^3 + 3a^2x + 3ax^2 + x^3$ ; by means of which the cubic roots of numbers have been extracted: viz, by subtracting the nearest cube  $a^3$  from the given power, dividing the remainder by  $3a^2$ , which gives  $x$  nearly for the quotient; then completing the divisor up to  $3a^2 + 3ax + x^2$ , multiply it by  $x$  for the other part of the power to be subtracted. And this was the extent of the Extraction of roots in the time of Lucas de Burgo, who, from 1470 to 1500, wrote several pieces on arithmetic and algebra, which were the first works of this kind printed in Europe.

It was not long however before the nature and composition of all the higher powers became known, and general tables of coefficients formed for raising them; the first of which is contained in Stifelius's arithmetic, printed at Norimberg in 1543, where he fully explains their use in Extracting the roots of all powers whatever, by methods similar to those for the square and cubic roots, as above described; and thus completed the Extraction of all sorts of roots of numbers, at least so far as respects that method of resolution. Since that time, however, many new methods of Extraction have been devised, as well as improvements made in the old way.

The Extraction of roots of equations followed closely that of known numbers. In De Burgo's time they extracted the roots of quadratic equations, the same way as at present. Ferreus, Tartalea, and Cardan extracted the roots of cubic equations, by general rules. Soon afterwards the roots of higher equations were extracted, at least in numbers, by approximation. And the late improvements in analytics have furnished general rules for Extracting the roots, in infinite series, of all equations whatever. All which methods may be seen in most books of arithmetic and algebra. Of which it may suffice to give here a short specimen of some of the easiest rules for Extracting the roots of quantities and equations, as they here follow.

I. *To Extract the Square Root of any Number.*—Point off, or divide the number, from the place of units, into portions of two figures each, as here of the number 99856, setting a point or mark over the space between each portion of two figures. Then, beginning at the left hand, take the greatest root 3, of the first part 9, placing it on the right hand for the first figure of the root, and subtracting its square 9 from the said first part; to the re-

$$\begin{array}{r} \text{root} \\ 99856 \text{ ( } 316 \\ 9 \\ \hline 61 \overline{) 98} \\ 1 \overline{) 61} \\ \hline 626 \overline{) 3756} \\ 6 \overline{) 3756} \end{array}$$

mainder, which here is 0, bring down the 2d part 98, and on the left hand of it place 6 the double of the first figure 3, for a divisor; conceive a cipher added to this, making it 60, and then divide the 98 by the 60, the quotient is 1 for the second figure of the root, which is accordingly placed there, after the 3, also in the divisor after the 6 and below the same; then multiply these as they stand, the 61 by the 1, and the product 61 set below the 98, and subtract it from the same, which leaves 37 for the next remainder; to this bring down the 3d period 56, making 3756 for the next resolvend: then form its divisor as before, viz, doubling the root 31, or adding, as they stand in the divisor, the 1 to the 61, either way making 62, which with a cipher makes 620, by which divide the resolvend 3756; the quotient of this division is 6, to be placed, as before, both as the next figure of the root, and at the end of the divisor 62, and below itself there; then multiply as they stand the whole divisor 626 by the 6, the product 3756 is exactly the same as the resolvend, and therefore the number 316 is accurately the square root of the given number 99856, as required.

When the root is to be carried into decimals, couplets of ciphers are to be added, instead of figures, as far as may be wanted. In which case too, a good abbreviation is made, after the work has been carried on to half the number of figures, by continuing it to the other half only by the contracted way of division; as here in the annexed example for the square root of 2 to eight decimals, or nine places of figures in all.

$$\begin{array}{r} 2 \text{ ( } 1.41421356 \\ 1 \\ \hline 24 \overline{) 100} \\ 4 \overline{) 96} \\ \hline 281 \overline{) 400} \\ 1 \overline{) 281} \\ \hline 2824 \overline{) 11900} \\ 4 \overline{) 11296} \\ \hline 28282 \overline{) 60400} \\ 2 \overline{) 56564} \\ \hline 28284 \overline{) 38361356} \\ \dots \dots 2828 \end{array}$$

II. *To extract the cubic root, or any other root whatever.* This is easiest done by one general rule, which I have invented, and published in my Tracts, vol. 1, pa. 49, which is to this effect: Let  $N$  be any number or power, whose  $n$ th root is to be extracted; and let  $R$  be the nearest rational root of  $N$ , of the same kind, or  $R^n$  the nearest rational power to  $N$ , either greater or less than it; then shall the true root be very nearly equal to

$$\frac{n+1 \cdot N + n-1 \cdot R^n}{n-1 \cdot N + n+1 \cdot R^n} \times R; \text{ which rule is general for}$$

any root whose index is denoted by  $n$ . And by expounding  $n$  successively by all the numbers 2, 3, 4, 5, &c, this theorem will give the following particular rules for the several roots, viz, the

$$\text{2d or squ. root, } \frac{3N + R^2}{N + 3R^2} \times R;$$

$$\text{3d or cube root, } \frac{4N + 2R^3}{2N + 4R^3} \times R, \text{ or } \frac{2N + R^3}{N + 2R^3} \times R;$$

$$\text{4th root: } \frac{5N + 3R^4}{3N + 5R^4} \times R;$$



5th root  $\frac{6N+4R^5}{4N+6R^5} \times R$ , or  $\frac{3N+2R^5}{2N+3R^5} \times R$ ;  
 6th root  $\frac{7N+5R^6}{5N+7R^6} \times R$ ;  
 7th root  $\frac{8N+6R^7}{6N+8R^7} \times R$ , or  $\frac{4N+3R^7}{3N+4R^7} \times R$ ;  
 &c. &c.

Or the theorem may be stated in the form of a proportion, thus:

as  $n-1 . N + n+1 . R^n : n+1 . N + n-1 . R^n ::$   
 $R : \text{the root sought very nearly.}$

For ex. suppose the problem proposed, of doubling the cube, or to find the cube root of the number 2. Here  $N = 2$ ,  $n = 3$ , and the nearest power, and root too, is 1:

Hence  $2N + R^3 = 4 + 1 = 5$ ,

and  $N + 2R^3 = 2 + 2 = 4$ ;

then  $4 : 5 :: 1 : \frac{5}{4} = 1.25$  the first approximation.

Again, taking  $R = \frac{5}{4}$ , and conseq.  $R^3 = \frac{125}{64}$ :

Hence  $2N + R^3 = 4 + \frac{125}{64} = \frac{381}{64}$ ,

and  $N + 2R^3 = 2 + \frac{250}{64} = \frac{378}{32}$ ;

then  $378 : 381 :: \frac{5}{4} : \frac{6385}{4096} = 1.259921$ , for the cube root of 2, which is exact in the very last figure.

And again by taking  $\frac{6385}{4096}$  for the value of  $R$ , a great many more figures may be found.

III. To Extract the Roots of Algebraic Quantities.—This is done by the same rules, and in the same manner as for the roots of numbers in arithmetic, as above taught. Thus, to Extract the square root of  $4a^2 + 12ax + 9x^2$ .

$$\begin{array}{r} 4a^2 + 12ax + 9x^2 \text{ (} 2a + 3x \text{ the root)} \\ 4a^2 \\ \hline 12ax + 9x^2 \\ 12ax + 9x^2 \\ \hline \end{array}$$

So also the root is carried out in an infinite series, in imitation of the like Extraction of numbers in infinite decimals: thus, for the square root of  $a^2 + x^2$ .

$$\begin{array}{r} a^2 + x^2 \left( a + \frac{x^2}{2a} - \frac{x^4}{8a^3} + \frac{x^6}{16a^5} \&c. \right) \\ a^2 \\ \hline 2a + \frac{x^2}{2a} \quad x^2 \\ \frac{x^2}{2a} \quad x^2 + \frac{x^4}{4a^2} \\ \hline 2a + \frac{x^2}{a} - \frac{x^4}{8a^3} \quad -\frac{x^4}{4a^2} \\ \frac{x^4}{8a^3} \quad -\frac{x^4}{4a^2} - \frac{x^6}{8a^4} + \frac{x^8}{4a^6} \\ \hline 2a + \frac{x^2}{a} - \frac{x^4}{4a^3} + \frac{x^6}{16a^5} \quad \frac{x^6}{8a^4} - \frac{x^8}{4a^6} \\ \frac{x^6}{16a^5} \quad \frac{x^6}{8a^4} + \frac{x^8}{16a^5} \&c. \\ \hline \end{array}$$

To extract the cube root of  $a^3 - x^3$  by the general rule in the 2d article.—Here  $N = a^3 - x^3$ ,  $R = a$ , or  $R^3 = a^3$ ; hence  $N + 2R^3 = 3a^3 - x^3$ , and  $2N + R^3 = 3a^3 - 2x^3$ ; therefore, by the rule,

$$3a^3 - x^3 : 3a^3 - 2x^3 :: a : a - \frac{x^3}{3a^2} - \frac{x^6}{9a^5} - \frac{x^9}{27a^8} \&c,$$

which is the cube root of  $a^3 - x^3$  very nearly.

But these sorts of roots are best extracted by the Binomial Theorem; which see.

IV. To Extract the Roots of Equations.—This is the same thing as to find the value of the unknown quantity in an equation; which is effected by various means, depending on the form of the equation, and the height of the highest power of the unknown quantity in it: for which, see the respective terms, EQUATION, ROOT, QUADRATIC, CUBIC, &c.

The most general, as well as the most easy, method of Extracting the roots of all equations, is by Double Position, or Trial-and-Error; as it easily applies to all sorts of equations whatever, be they ever so complex, even logarithmic and exponential ones. There are also several other good methods of approximating to the roots of equations, given by Newton, Halley, Raphson, De Moivre, &c; of which the most general is a rule for Extracting the root of the following indefinite equation,



$$\text{viz, } ax + bx^2 + cx^3 + dx^4 + ex^5 \&c \\ = gy + hy^2 + iy^3 + ky^4 + ly^5 \&c,$$

given by M. De Moivre in the Philos. Transf. vol. 20. p. 190, or Abr. vol. 1, pa. 101.

EXTRADOS, the outside of an arch of a bridge, vault, &c.

EXTREME-and-Mean Proportion, is when a line, or any quantity is so divided, as that the whole line is to the greater part, as that greater part is to the less part. Hence, in any line so divided, the rectangle of the whole line and the less segment, is equal to the square of the greater segment.

Euclid shews how to divide a line in Extreme-and-mean ratio, in his Elements, book 2, prop. 11, to this effect: Let AB be the given line; to which draw AE perpendicular and equal to half AB; in EA produced take EF = EB, so shall AF be equal to the greater part; consequently if AG be taken equal to AF, the line AB will be divided in G as required.



The same may be done otherwise thus:

As before, make AE (fig. 2.) perpendicular and =  $\frac{1}{2}AB$ ; join EB, on which take EC = EA, and then take BD = BC, so shall the line be divided in D as required.

No number can be divided into extreme and mean proportion, so that its two parts shall be rational; as is well demonstrated by Clavius, in his Commentary upon the 9th book of Euclid's Elements; and the same thing will also appear from the following algebraical solution of the same problem: Let  $a$  denote the whole line, and  $x$  the greater part; then shall  $a - x$  be the less part, and the




the rectangle of the whole and less part being put equal to the square of the greater part, gives this equation,  $x^2 = a \times a - x = a^2 - ax$ ; hence  $x^2 + ax = a^2$  and by completing the square, and extracting the root, &c, there is at last  $x = \frac{\sqrt{5}-1}{2}a$  the greater part; conse-

quently  $a - x = \frac{3-\sqrt{5}}{2}a$  is the less part. And as the square root of 5, which cannot be exactly extracted, makes a portion of both these parts, it is manifest that neither of them can be obtained in rational numbers.

Euclid makes great use of this problem, viz, in several parts of the 13th book of the Elements; and by means of it he constructs that notable proposition, viz the 10th of the 4th book, which is to construct an isosceles triangle having each angle at the base double the angle at the vertex.

EXTREMES *Conjunct*, and EXTREMES *Disjunct*, in Spherical Trigonometry, are the former the two circular parts that lie next the assumed middle part, and the latter are the two that lie remote from the middle part. These were terms applied by lord Napier, in his universal theorem for resolving all right-angled and quadrantal spherical triangles, and published in his *Logarithmorum Canonis Descriptio*, an. 1614. In this theorem, Napier condenses into one rule, in two parts, the rules for all the cases of right-angled spherical triangles, which had been separately demonstrated by Pitiscus, Lansbergius, Copernicus, Regiomontanus, and others. In this theorem, neglecting the right angle, Napier calls the other five parts, circular parts, which are, the two legs about the right angle, and the complements of the other three, viz of the hypotenuse, and the two oblique angles. Then, taking any three of these five parts, one of them will be in the middle between the other two, and these two are the Extremes *Conjunct* when they are immediately adjacent to that middle part, or they are the Extremes *Disjunct* when they are each separated from the middle one by another part. Thus, the five parts being AB, AC, and the complements of BC and of the two angles B and C: then if the three parts be AB, and the complements of the angle B and hypotenuse BC be taken, these three are contiguous to each other, the angle B lying in the middle between the other two: therefore the comp. of B is middle part, and AB with the comp. of BC the Extremes *Conjunct*. But if the three sides be taken; BC is equally separated from the two legs AB and AC, by two angles B and C; and therefore these two legs AB and AC are Extremes *Disjunct*, and the comp. of BC the middle part.



Napier's rule for resolving each case is in two parts, as below:

The rectangle contained by radius and the sine of the middle part, is equal to the rectangle of the tangents of the Extremes *conjunct*, or equal to the rectangle of the sines of the Extremes *disjunct*. Which rule comprehends all the cases that can happen in right-angled spherical triangles; in the application of which rule, the equal rectangles are divided into a proportion or analogy, in such manner that the term sought may be the last of

the four terms that are concerned, and consequently its corresponding term in the same rectangle must be the first of those terms.

EYE, the organ of sight, consisting of several parts, and of such forms as best to answer the purpose for which it was formed.

As vision or sight is effected by a refraction of light through the humours of the eye to the bottom or farther internal part of it, where the images of external objects are formed on a fine expansion of the optic nerve, called the *Retina*, and therefore the fore-part of the eye must be of a convex figure, and of such a precise degree of convexity as the particular refractive power of the several humours require for forming the image of an object at a given focal distance, viz, the diameter of the eye. Hence we find,

1st; The external part of the eye-ball CD (Plate 2, fig. 8.) is a strong pellucid substance, properly convex, and which, when dried, has some resemblance to a piece of transparent horn, for which reason it is called the *Cornea*, or horny coat of the eye.

2dly; Immediately behind this coat there is a fine clear humour which, from its likeness to water, is called the *Aqueous* or watery humour, and is contained in the space between CD and GFE.

3dly; In this space there is a membrane or diaphragm, called the *Uvea*, with a hole in the middle as at F, called the *Pupil*, of a muscular contexture for altering the dimensions of that hole, for the adjusting or admitting a due quantity of light.

4thly; Just behind this diaphragm is placed a lenticular-formed substance GE, of a considerable consistence, called from its transparency the *Crystalline* humour. This is contained in a fine tunic called the *Choroides*, and is suspended in the middle of the eye by a ring of muscular fibres called the *Ligamentum Ciliare*, as at G and E; by which means it is moved a little nearer to, or farther from, the bottom of the eye, to alter the focal distance.

5thly; All the remaining interior part of the eye, constituting the great body of it, from GHE to IMK, is made up of a large quantity of a jelly-like substance, called the *Vitreous* or glassy humour; though it resembles glass in nothing except its transparency; it being most like the white of an egg of any thing.

6thly; On one side of the hinder part of the Eye, as at K, the optic nerve enters it from the brain, and is expanded over all the interior part of the eye to G and E quite around, the expansion being named the *Retina*. On this delicate membrane, the image IM of every external object OB, is formed according to the optic laws of nature, in the following manner.

Let OB be any very distant object. Then a pencil of rays proceeding from any point L, will fall on the cornea CD, and be refracted by the aqueous humour under it, to a point in the axis of that pencil continued out. Then the radius of convexity of the cornea being nearly  $\frac{1}{3}$  of an inch; and the sine of incidence in air to that of refraction in the aqueous humour, being nearly as 4 to 3; supposing the rays parallel, or the object very far distant, the focal distance after the first refraction, by the proper theorem  $\frac{mr}{m-n}$ , will be found  $1\frac{1}{3}$  inch from the cornea:  $r$  being the radius  $\frac{1}{3}$ , and  $m$  to  $n$  as 4 to 3.

The



The rays thus refracted by the cornea, fall converging on the crystalline humour, and tend to a point 1.228 inch behind it; also the radii of convexity in the said humour are  $\frac{1}{3}$  and  $\frac{1}{4}$  respectively; and the sine of incidence to that of refraction of the aqueous into the crystalline humour, being as 13 to 12; therefore, by this theorem  $\frac{m d r}{m d - n d + n r}$ , the focal distance after refraction in the crystalline, will be 1.02 inch from the fore part of it: where  $m = 13$ ,  $n = 12$ ,  $r = \frac{1}{3}$ , and  $d = 1.228$ .

The rays now pass from the crystalline to the vitreous humour still in a converging state, and the sines of incidence and refraction being here as 12 to 13, as found by experiment; and since the surface of the vitreous humour is concave which receives the rays, and is the same with the convexity of the hinder surface of the crystalline, the radius will be the same, viz  $\frac{1}{4}$  of an inch. Therefore the focal distance after this third

refraction will be found, by this theorem,  $\frac{m d r}{n d - m d + n r}$ , to be 6 tenths of an inch nearly from the hinder part of the cornea: where  $m = 12$ ,  $n = 13$ ,  $r = \frac{1}{4}$ , and  $d = .82$ ; the thickness of the lens of the cornea being nearly  $\frac{1}{2}$  of an inch.

Now experience shews that the distance of the retina in the back part of the eye, behind the cornea, is nearly equal to that focal distance; and therefore it follows that all objects at a great distance have their images formed on the retina in the bottom or hinder part of the eye, and thus distinct vision is produced by this wonderful organ of optic sensation.

When the distance of objects is not very great, the focal distance, after the last refraction in the vitreous humour, will be a little increased; and to do this we can move the crystalline a little nearer the cornea by means of the ligamentum ciliare, and thus on all occasions it may be adjusted for a due focal distance for every distance of objects, excepting that which is less than 6 or 7 inches, in good eyes. Many are of opinion, however, that this is effected by a power in the eye to alter the convexity of the crystalline humour as occasion requires; though this is rather doubtful.

By what has been said it appears, that rays of light flowing from every part of an object OB, placed at a proper distance from the eye, will have an image IM thereby formed on the retina in the bottom of the eye; and since the rays OM, BI, which come from the extreme parts of the object, cross each other in the middle of the pupil, the position of the image IM will be contrary to that of the object, or inverted, as in the case of a lens.

The apparent place of any part of an object is in the axis, and conjugate focus of that pencil of rays by which that part or point is formed in the image. Thus, OM is the axis, and O the focus proper to the rays by which the point M in the image is made; therefore the sensation of the place of that part will be conceived in the mind to be at O; in like manner the idea of place belonging to the point I, will be referred in the axis IB, to the proper focus B; therefore the apparent place of the whole image IM, will be conceived in the mind to occupy all the space between O and B, and at the distance AL from the eye.

Hence likewise appears the reason, why we see an object upright by means of an inverted image; for since the apparent place of every point M will be in the axis MO at O; and this axis crossing the axis of the eye HL in the pupil, it follows, that the sensible place O of that point will lie, without the eye, on the contrary side of the axis of the eye, to that of the point in the eye; and since this is true of all other parts or points in the image, it is evident that the position of every part of the object will be on the contrary side of the axis to every corresponding part in the image, and therefore the whole object OB will have a contrary position to that of the image IM, or will appear upright.

If the convexity of the cornea CD happens not exactly to correspond to the diameter of the eye, considered as the natural focal distance, then the image will not be formed on the retina, and consequently no distinct vision can be effected in such an eye.

If the cornea be too convex, the focal distance in the eye will be less than its diameter, and the image will be formed short of the retina. Hence the reason why people having such eyes are obliged to hold things very near them, to lengthen the focal distances; and also why they use concave glasses to counteract or remedy the excess of convexity, in order to view distant objects distinctly.

When the eye has less than a just degree of convexity, or is too flat, as is generally the case with old eyes, by a natural deficiency of the aqueous humour, then the rays tend to a point or focus beyond the retina or bottom of the eye; and to supply this want of convexity in the cornea, we use convex lenses in those frames called spectacles, or visual glasses.

Since the rays of light OA, BA, which constitute the visual angle OAB, will, when they are intercepted by a lens, be refracted sooner to the axis; the said angle will thereby be enlarged, and the object of course become magnified; which is the reason why those lenses are called magnifiers, or reading-glasses.

The dimensions, or magnitude, of an object OB, are judged of by the quantity of the angle OAB which it subtends at the eye. For if the same object be placed at two different distances L and N, the angles OAB, oAb, which in these two places it subtends at the eye, will be of different magnitude; and the lineal dimensions, viz length and breadth, will be at N and at L, as the angle oAb is to the angle OAB. But the surfaces of the objects will be as the squares of those angles, and the solidities as the cubes of them.

It is found by experience, that two points O, L, in any object, will not be distinctly seen by the Eye, till they are near enough to subtend an angle OAL of one minute. And hence when objects, however large they may be, are so remote as not to be seen under an angle of one minute, they cannot properly be said to have any apparent dimensions or magnitude at all; such as is the case of the large bodies of the planets, comets, and fixed stars. But the optic science has supplied means of enlarging this natural small angle under which most distant objects appear, and thereby increasing their apparent magnitudes to a very surprising degree, in the instance of that noble instrument the telescope.

On the other hand, there are in creation an infinity



of objects, of such small dimensions, that they will not subtend the requisite angle, if brought to the nearest limits of distinct vision, viz 6, 7, or 8 inches from the Eye, as found by experience; and therefore to render them visible at a very near distance, we have a variety of glasses, and instruments of different constructions, usually called microscopes, by which those minute objects appear many thousand times larger than to the naked Eye; and thereby enrich the mind with discoveries of the sublimest nature, in regard to creative power, wisdom, and œconomy.

EYE-glass, in Optical Instruments, is that which is next the Eye in using the machine. This is usually a lens convex on both sides; but Eustachia Divini long

since invented a microscope of this kind, the power of which he places very greatly above that of the common sort; and this chiefly depending on the Eye-glass, which was double, consisting of two plano-convex glasses, so placed as to touch one another in the middle of their convex surface. This instrument is well spoken of by Fabri in his Optics, and as possessing this peculiar excellence, that it shews all the objects flat, and not crooked, and takes in a large area, though it magnifies very much.

*Bull's EYE*, a star of the first magnitude, in the Eye of the constellation Taurus, the bull, and by the Arabs called *Aldebaran*.

## F.

## F A C

**F**ACE, or FAÇADE, in Architecture, is sometimes used for the front or outward part of a building, which immediately presents itself to the eye; or the side where the chief entrance is, or next the street, &c.

FACE, FACIA, or FASCIA, also denotes a flat member, having a considerable breadth, and but a small projecture. Such are the bands of an architrave, larmier, &c.

FACE, in Astrology, is used for the 3d part of a sign.—Each sign is supposed to be divided into three faces, of 10 degrees each: the first 10 degrees compose the first face; the next 10 degrees, the 2d face; and the last 10, the 3d face.—Venus is in the 3d face of Taurus; that is, in the last 10 degrees of it.

FACES of a Bastion, in Fortification, are the two foremost sides, reaching from the flanks to the outermost point of the bastion, where they meet, and form the salient angle of the bastion. These are usually the first parts that are undermined, or beaten down; because they reach the farthest out, are the least flanked, and are therefore the weakest.

FACE of a Place, is the extent between the outermost points of two adjacent bastions; containing the curtain, the two flanks, and the two faces of those bastions that look towards each other. This is otherwise called the Tenaille of the place.

FACE Prolonged, is that part of a line of defence rasant, which is between the angle of the epaule or shoulder of a bastion and the curtain; or the line of a defence rasant diminished by the face of the bastion.

VOL. I.

## F A C

FACIA, in Architecture. See FACE, and FASCIA.

FACTORS, in Multiplication, in Arithmetic, a name given to the two numbers that are multiplied together, viz, the multiplicand and multiplier; so called because they are to *facere productum*, make or constitute the factum or product.

FACTUM, the product of two quantities multiplied together. As, the factum of 3 and 4 is 12; and the factum of  $2a$  and  $5b$  is  $10ab$ .

FACULÆ, in Astronomy, a name given by Scheiner, and others after him, to certain bright spots on the sun's disc, that appear more bright and lucid than the rest of his body.

Hevelius assures us that, on July 20, 1634, he observed a facula whose breadth was equal to a 3d part of the sun's diameter. He says too that the maculæ often change into Faculæ; but these seldom or never into maculæ. And some authors even contend that all the maculæ degenerate into Faculæ before they quite disappear. Many authors, after Kircher and Scheiner, have represented the sun's body full of bright, fiery spots, which they conceive to be a sort of volcanos in the body of the sun: but Huygens, and others of the latest and best observers, finding that the best telescopes discover nothing of the matter, agree entirely to explode the phenomena of Faculæ. All the foundation he could see for the notion of Faculæ, he says, was, that in the darkish clouds which frequently surround the maculæ, there are sometimes seen little points or sparks.



brighter than the rest. Their cause is attributed by these authors to the tremulous agitation of the vapours near our earth; the same as sometimes shews a little unevenness in the circumference of the sun's disc when viewed through a telescope. Strictly then, the Faculæ are not eruptions of fire and flame, but refractions of the sun's rays in the rarer exhalations, which, being condensed near that shade, seem to exhibit a light greater than that of the sun.

FACULTY, denotes the several parts of an university, divided according to the arts or sciences taught or professed there. In most universities there are four Faculties; that of arts, which includes philosophy and the humanities or languages, and is the most ancient and extensive; the 2d is that of theology; the 3d, that of medicine; and the 4th, jurisprudence, or laws.

FALCATED, one of the phases of the planets, vulgarly called horned. The astronomers say, the moon, or any planet, is Falcated, when the enlightened part appears in form of a crescent, like a sickle, or reaping-hook, which by the Latins is called *falk*. The moon is Falcated while she moves from the 3d quarter to the conjunction, and so on from hence to the first quarter; the bright part appearing then like a crescent, viz during the first and last quarters. But during the 2d and 3d quarters, the light part appears gibbous, and the dark part Falcated.

FALCON or FAUCON, and FALCONET or FAUCONET, certain old species of cannon, now long disused.

FALL, the descent or natural motion of bodies towards the centre of the earth, &c. Galileo first discovered the ratio of the acceleration of falling bodies; viz, that the spaces descended from rest are as the squares of the times of descent; or, which comes to the same thing, that if the whole time of Falling be divided into any number of equal parts, whatever space it falls through in the first part of the time, it will Fall 3 times as far in the 2d part of time, and 5 times as far in the 3d portion of time, and so on, according to the uneven numbers 1, 3, 5, 7, &c. See ACCELERATION, DESCENT, GRAVITY, &c.

FALSE-BRAYE, in Fortification. See FAUSSE-BRAYE.

FALSE Position, in Arithmetic. See POSITION.

FALSE Root, a name given by Cardan, to the negative roots of equations, and numbers. So the root of 9 may be either 3 or - 3, the former he calls the true, and the latter the false or fictitious root; also of this equation  $x^2 - x = 6$ , the two roots are 3 and - 2, the former true, and the latter False.

FASCIA, in Architecture. See FACIA and FACE.

FASCIÆ, in Astronomy, are certain stripes or rows of bright parts, observed on the bodies of some of the planets, like swathes, bands, or belts; especially on the planet Jupiter.

The Fasciæ, or belts of Jupiter, are more lucid than the rest of the disc, and are terminated by parallel lines. They are sometimes broader and sometimes narrower; nor do they always possess the same part of the disc.

M. Huygens also observed a very large kind of Fascia in Mars, in the year 1656; but it was darker than the rest of the disc, and occupied the middle part of it.

FASCINES, in Fortification, are faggots made of

the twigs and small branches of trees and brush wood, bound up in bundles; these, being mixed with earth, serve to fill up ditches, to make the parapets of trenches, batteries, &c.

FATHOM, an English measure of the length of 6 feet or 2 yards; and is taken from the utmost extent of both arms when stretched into a right line.

FATUUS *Ignis*. See *IGNIS Fatuus*.

FAUCON, and FAUCONET, the same as Falcon and Falconet; the old names of certain species of ordnance; which, as well as many other names, are now no longer in use, as it has been for some time the practice to denominate the several sizes of cannon from the weight of their ball, instead of calling them by those fanciful and unmeaning names.

FAUSSE-BRAYE, in Fortification, an elevation of earth, about three feet above the level ground; round the foot of the rampart on the outside, defended by a parapet about four or five fathoms distant from the upper parapet, which parts it from the berme, and the edge of the ditch. The Fausse-braye is the same with what is otherwise called *Chemin des rondes*, and *Basse enceinte*; and its use is for the defence of the ditch.

FEATHER-EDGED, is a term used by workmen, for such boards as are thicker on one edge, or side, than on the other.

FEBRUARY, the 2d month of the year, containing 28 days for three years, and every fourth year 29 days.—In the first ages of Rome, February was the last month of the year, and preceded January, till the *Decemviri* made an order that February should be the 2d month of the year, and come after January.

FELLOWSHIP, COMPANY, or PARTNERSHIP, is a rule in arithmetic, of great use in balancing accounts among merchants, and partners in trade, teaching how to assign to every one of them his due share of the gain or loss, in proportion to the stock he has contributed, and the time it has been employed, or according to any other conditions. Or, more generally, it is a method of dividing a given number, or quantity, into any number of parts, that shall have any assigned ratios to one another. And hence comes this general rule: Having added into one sum the several numbers that express the proportions of the parts, it will be,

As that sum of the proportional numbers :

Is to the given quantity that is to be divided ::

So is each proportional number :

To the corresponding share of the given quantity.

*For Ex.* Suppose it be required to divide the number 120 into three parts that shall be in proportion to each other as the numbers 1, 2, 3.—Here 120 is the quantity to be divided, and 6 is the sum of the numbers 1, 2, and 3, which express the proportions of the parts; therefore as

$$6 : 120 :: \begin{cases} 1 : 20 \text{ the 1st part,} \\ 2 : 40 \text{ the 2d part,} \\ 3 : 60 \text{ the 3d part.} \end{cases}$$

This rule is usually distinguished into two cases, one in which time is concerned, or in which the stocks of partners are continued for different times; and the other in which time is not considered; this latter being called *Single Fellowship*, and the former *Double Fellowship*.



*Single FELLOWSHIP, or FELLOWSHIP without Time,* is the case in which the times of continuance of the shares of partners are not considered, because they are all the same; and in this case, the rule will be as above, viz,

As the whole stock of the partners :

Is to the whole gain or loss ::

So is each one's particular stock :

To his share of the gain or loss.

*Ex.* Two partners, A and B, form a joint stock, of which A contributed 75l, and B 45l; with which they gain 30l: how much of it must each one have?

As 120 : 30 ::  $\begin{cases} 75 : 18l. 15s. = A's \text{ share,} \\ 45 : 11l. 5s. = B's \text{ share.} \end{cases}$

*Double FELLOWSHIP, or FELLOWSHIP with Time,* is the case in which the times of the stocks continuing are considered, because they are not all the same.

In this case, the shares of the gain or loss must be proportional, both to the several shares of the stock, and to the times of their continuance, and therefore proportional to the products of the two. Hence this Rule: Multiply each particular share of the stock by the time of its continuance, and add all the products together into one sum; then say,

As that sum of the products :

Is to the whole gain or loss ::

So is each several product :

To the corresponding share of the gain or loss.

*For Ex.* A had in company 50l. for 4 months, and B 60l. for 5 months; and their gain was 24l: how must it be divided between them?

50	60
4	5
200	300
300	

As 500 : 24 ::  $\begin{cases} 200 : 9l. 12s. = A's \text{ share,} \\ 300 : 14l. 8s. = B's \text{ share.} \end{cases}$

FERGUSON (JAMES), an eminent experimental philosopher, mechanist, and astronomer, was born in Bamfshire, in Scotland, 1710, of very poor parents. At the very earliest age his extraordinary genius began to unfold itself. He first learned to read, by overhearing his father teach his elder brother: and he had made this acquisition before any one suspected it. He soon discovered a peculiar taste for mechanics, which first arose on seeing his father use a lever. He pursued this study a considerable length, while he was yet very young; and made a watch in wood-work, from having once seen one. As he had at first no instructor, nor any help from books, every thing he learned had all the merit of an original discovery; and such, with inexpressible joy, he believed it to be.

As soon as his age would permit, he went to service; in which he met with hardships, which rendered his constitution feeble through life. While he was servant to a farmer (whose goodness he acknowledges in the modest and humble account of himself which he prefixed to one of his publications), he contemplated and learned to know the stars, while he tended the sheep; and began the study of astronomy, by laying down, from his

own observations only, a celestial globe. His kind master, observing these marks of his ingenuity, procured him the countenance and assistance of some neighbouring gentlemen. By their help and instructions he went on gaining farther knowledge, having by their means been taught arithmetic, with some algebra, and practical geometry. He had got some notion of drawing, and being sent to Edinburgh, he there began to take portraits in miniature, at a small price; an employment by which he supported himself and family for several years, both in Scotland and England, while he was pursuing more serious studies. In London he first published some curious astronomical tables and calculations; and afterwards gave public lectures in experimental philosophy, both in London and most of the country towns in England, with the highest marks of general approbation. He was elected a fellow of the Royal Society, and was excused the payment of the admission fee and the usual annual contributions. He enjoyed from the king a pension of 50 pounds a year, besides other occasional presents, which he privately accepted and received from different quarters, till the time of his death; by which, and the fruits of his own labours, he left behind him a sum to the amount of about six thousand pounds, instead of which all his friends had always entertained an idea of his great poverty. He died in 1776, at 66 years of age, though he had the appearance of many more years.

Mr. Ferguson must be allowed to have been a very uncommon genius, especially in mechanical contrivances and executions, for he executed many machines himself in a very neat manner. He had also a good taste in astronomy, with natural and experimental philosophy, and was possessed of a happy manner of explaining himself in an easy, clear, and familiar way. His general mathematical knowledge, however, was little or nothing. Of algebra he understood but little more than the notation; and he has often told me he could never demonstrate one proposition in Euclid's Elements; his constant method being to satisfy himself, as to the truth of any problem, with a measurement by scale and compasses. He was a man of a very clear judgment in any thing that he professed, and of unwearied application to study: benevolent, meek, and innocent in his manners as a child: humble, courteous, and communicative: instead of pedantry, philosophy seemed to produce in him only diffidence and urbanity.

The list of Mr. Ferguson's public works, is as follows:

1. Astronomical Tables and Precepts, for calculating the true times of New and Full Moons, &c; 1763.
- 2. Tables and Tracts, relative to several arts and sciences; 1767.—3. An Easy Introduction to Astronomy, for Young Gentlemen and Ladies; 2d edit. 1769.—4. Astronomy explained upon Sir Isaac Newton's Principles; 5th edit. 1772.—5. Lectures on Select Subjects in Mechanics, Hydrostatics, Pneumatics, and Optics; 4th edit. 1772.—6. Select Mechanical Exercises; with a short Account of the life of the author, by himself; 1773.—7. The Art of Drawing in Perspective made easy; 1775.—8. An Introduction to Electricity; 1775.—9. Two Letters to the Rev. Mr. John Kennedy; 1775.—10. A Third Letter to the Rev. Mr. John Kennedy; 1775.



**FERMAT (PETER)**, counsellor of the parliament of Toulouse, in France, who flourished in the 17th century, and died in 1663. He was a general scholar, and a universal genius, cultivating jurisprudence, poetry, and mathematics, but especially the latter, for his amusement. He was contemporary and intimately connected with Merfenne, Torricelli, Des Cartes, Pascal, Roberval, Huygens, Frenicle, and Carcavi, and several others the most celebrated philosophers of their time. He was a first-rate mathematician, and possessed the finest taste for pure and genuine geometry, which he contributed greatly to improve, as well as algebra.

Fermat was author of 1. A Method for the Quadrature of all sorts of Parabolas.—2. Another on Maximums and Minimums: which serves not only for the determination of plane and solid problems; but also for drawing tangents to curve lines, finding the centres of gravity in solids, and the resolution of questions concerning numbers: in short a method very similar to the Fluxions of Newton.—3. An Introduction to Geometric Loci, plane and solid.—4. A Treatise on Spherical Tangencies: where he demonstrates in the solids, the same things as Vieta demonstrated in planes.—5. A Restoration of Apollonius's two books on Plane Loci.—6. A General Method for the dimension of Curve Lines. Besides a number of other smaller pieces, and many letters to learned men; several of which are to be found in his *Opera Varia Mathematica*, printed at Toulouse, in folio, 1679.

**FERMENTATION**, an intestine motion, arising spontaneously among the small and insensible particles of a mixed body, thereby producing a new disposition, and a different combination of those parts. Fermentation differs from dissolution, as the cause from its effect, the latter being only a result or effect of the former.

**FESTOON**, in Architecture &c, a decoration in form of a garland or cluster of flowers.

**FICHANT FLANK**. See **FLANK**.

**FICHANT Line of Defence**. See **FIXED Line of Defence**.

**FIELD-FORT**. See **FORTINE**.

**FIELD-Pieces**, are small cannon, usually carried along with an army in the field: such as, one pounders, one and a half, two, three, four, six, nine, and 12 pounders; which, being light and small, are easily carried.

**FIELD-Staff**, is a staff carried by the gunners, in which they screw lighted matches, when they are on service; which is called arming the Field-staffs. See **LINSTOCK**.

**FIELD of View, or of Vision**, is the whole space or extent within which objects can be seen through an optical machine, or at one view of the eye without turning it.

The precise limits of this space are not easily ascertained, for the natural view of the eye. In looking at a small distance, we have an imperfect glimpse of objects through almost the extent of a hemisphere, or at least for above 60 degrees each way from the optic axis; but towards the extremity of this space, objects are very imperfectly seen; and the diameter of the field of distinct vision does not subtend an angle of more than 5 degrees at most, so that the diameter of a distinct image on the retina is less than  $\frac{1}{100}$  of an inch; but it is probably much less.

**FIELD-Book**, in Surveying, a book used for setting down angles, distances, and other things, remarkable in taking surveys.

The pages of the Field-book may be conveniently divided into three columns. In the middle column are to be entered the angles taken at the several stations by the theodolite, with the distances measured from station to station. And the offsets, taken with the offset-staff, on either side of the station line, are to be entered in the columns on either side of the middle column, according to their position, on the right or left, with respect to that line: also on the right or left of these are to be set down the names and characters of the objects, with proper remarks, &c. See a specimen in my Treatise on Mensuration, pa. 517, ed. 2d.

**FIFTH**, in Music, one of the harmonical intervals or concords; called by the ancients Diapente.

The Fifth is the 3d in order of the concords, and the ratio of the chords that produce it, is that of 3 to 2. It is called Fifth, because it contains five terms, or sounds, between its extremes, and four degrees; so that in the natural scale of music it comes in the 5th place, or order, from the fundamental.—The imperfect, or defective Fifth, by the ancients called Semidiapente, is less than the Fifth by a mean semitone.

**FIGURAL**, the same as **FIGURATE** numbers; which see.

**FIGURATE Numbers**, such as do or may represent some geometrical figure, in relation to which they are always considered; as triangular, pentagonal, pyramidal, &c, numbers.

Figurate numbers are distinguished into orders, according to their place in the scale of their generation, being all produced one from another, viz, by adding continually the terms of any one, the successive sums are the terms of the next order, beginning from the first order which is that of equal units 1, 1, 1, 1, &c; then the 2d order consists of the successive sums of those of the 1st order, forming the arithmetical progression 1, 2, 3, 4, &c; those of the 3d order are the successive sums of those of the 2d, and are the triangular numbers 1, 3, 6, 10, 15, &c; those of the 4th order are the successive sums of those of the 3d, and are the pyramidal numbers 1, 4, 10, 20, 35, &c; and so on, as below:

Order.	Name.	Numbers.
1.	Equals.	1, 1, 1, 1, 1, &c.
2.	Arithmeticals,	1, 2, 3, 4, 5, &c.
3.	Triangulars,	1, 3, 6, 10, 15, &c.
4.	Pyramidal,	1, 4, 10, 20, 35, &c.
5.	2d Pyramidal,	1, 5, 15, 35, 70, &c.
6.	3d Pyramidal,	1, 6, 21, 56, 126, &c.
7.	4th Pyramidal,	1, 7, 28, 84, 210, &c.

The above are all considered as different sorts of triangular numbers, being formed from an arithmetical progression whose common difference is 1. But if that common difference be 2, the successive sums will be the series of square numbers: if it be 3, the series will be pentagonal numbers, or pentagons; if it be 4, the series will be hexagonal numbers, or hexagons; and so on. Thus:

*Arith-*




Arithme- ticals.	1st Sums, or Polygons.	2d Sums, or 2d Polygons.
1, 2, 3, 4,	Tri. 1, 3, 6, 10	1, 4, 10, 20
1, 3, 5, 7,	Sqrs. 1, 4, 9, 16	1, 5, 14, 30
1, 4, 7, 10,	Pent. 1, 5, 12, 22	1, 6, 18, 40
1, 5, 9, 13, &c.	Hex. 1, 6, 15, 28	1, 7, 22, 50

And the reason of the names triangles, squares, pentagons, hexagons, &c, is, that those numbers may be placed in the form of these regular figures or polygons, as here below :

#### Triangles.




#### Squares.



#### Pentagons.



#### Hexagons.



But the Figurative numbers of any order may also be found without computing those of the preceding orders ; which is done by taking the successive products of as many of the terms of the arithmetics 1, 2, 3, 4, 5, &c, in their natural order, as there are units in the number which denominates the order of Figurates required, and dividing those products always by the first product : thus, the triangular numbers are found by dividing the products  $1 \times 2$ ,  $2 \times 3$ ,  $3 \times 4$ ,  $4 \times 5$ , &c, each by the 1st pr.  $1 \times 2$  ; the first pyramids by dividing the products  $1 \times 2 \times 3$ ,  $2 \times 3 \times 4$ ,  $3 \times 4 \times 5$ , &c, by the first  $1 \times 2 \times 3$ . And, in general, the figurate numbers of any order  $n$ , are found by substituting successively 1, 2, 3, 4, 5, &c, instead of  $x$  in this general expression  $\frac{x \cdot x + 1 \cdot x + 2 \cdot x + 3 \cdot \&c}{1 \cdot 2 \cdot 3 \cdot 4 \cdot \&c}$  ; where the factors in the numerator and denominator are supposed to be multiplied together, and to be continued till the number in each be less by 1 than that which expresses the

order of the Figurates required. See Maclaurin's Fluxions, art. 351, in the notes ; also Simpson's Algebra, pa. 213 ; or Malcolm's Arithmetic, pa. 396, where the subject of Figurates is treated in a very extensive and perspicuous manner.

FIGURE, in general, denotes the surface or terminating extremes of a body.—All finite bodies have some figure, form, or shape ; whence, figurability is reckoned among the essential properties of body, or matter : a body without Figure, would be an infinite body.

FIGURES, in Architecture and Sculpture, denote representations of things made in solid matter ; such as statues, &c.

FIGURES, in Arithmetic, are the numeral characters, by which numbers are expressed or written, as the ten digits, 1, 2, 3, 4, 5, 6, 7, 8, 9, 0. These are usually called the Arabic, and Indian figures, from which people it is supposed they have been derived. They were brought into Europe by the Moors of Spain, and into England about 1130, as Dr. Wallis apprehends : see his Algebra, pa. 9. However, from some ancient dates, supposed to consist wholly or in part of Arabian figures, some have concluded that these Figures, originally Indian, were known and used in this country at least as early as the 10th century. The oldest date discovered by Dr. Wallis, was on a chimney piece, at Helmdon, in Northamptonshire, thus M133, that is 1133. Other dates discovered since, are 1090, at Colchester, in Essex ; M16 or 1016, at Widgel-hall, near Buntingford, in Hertfordshire ; 1011 on the north front of the parish church of Rumsey in Hampshire ; and 975 over a gate-way at Worcester.

Dr. Ward, however, has urged several objections against the antiquity of these dates. As no example occurs of the use of these figures in any ancient manuscript, earlier than some copies of Johannes de Sacro Bosco, who died in 1256, he thinks it strange that these Figures should have been used by artificers so long before they appear in the writings of the learned ; and he also disputes the fact. The Helmdon date, according to him, should be 1233 ; the Colchester date 1490 ; that at Widgel-hall has in it no Arabic Figures, the 1 and 6 being I and G, the initial letters of a name ; and the date at Worcester consists, he supposes, of Roman numerals, being really MXV. Martyn's Abridg. Philos. Transf. vol. 9, pa. 420.

Mr. Gibbon observes (in his History of the Decline and Fall of the Roman Empire, vol. v. pa. 321). that “ under the reign of the caliph Waled, the Greek language and characters were excluded from the accounts of the public revenue. If this change was productive of the invention or familiar use of our present numerals, the Arabic characters or cyphers, as they are commonly styled, a regulation of office has promoted the most important discoveries of arithmetic, algebra, and the mathematical sciences.”

On the other hand it may be observed that ; “ according to a new, though probable notion, maintained by M. de Villafion) Anecdota Græca, tom. ii. p. 152, 157), our cyphers are not of Indian or Arabic invention. They were used by the Greek and Latin arithmeticians long before the age of Boethius. After the extinction of science in the west, they were adopted in the Arabic versions from the original manuscripts, and



and restored to the Latins about the eleventh century.

FIGURE, in Astrology, a description, draught, or construction of the state and disposition of the heavens, at a certain point of time; containing the places of the planets and stars, marked down in a Figure of 12 triangles, called houses. This is also called a Horoscope, and Theme.

FIGURE, of an Eclipse, in Astronomy, denotes a representation upon paper &c, of the path or orbit of the sun or moon, during the time of the eclipse; with the different phases, the digits eclipsed, and the beginning, middle, and end of darkness, &c.

FIGURE, or Delineation, of the full moon, such as, viewed through a telescope with two convex glasses, is of considerable use in observations of eclipses, and conjunctions of the moon with other luminaries. In this Figure are usually represented the maculæ or spots of the moon, marked by numbers; beginning with the spots that usually enter first within the shade at the time of the eclipses, and also emerge the first.

FIGURE, in Conic Sections, according to Apollonius, is the rectangle contained under the latus-rectum and the transverse axis, in the ellipse and hyperbola.

FIGURE, in Fortification, is the plan of any fortified place; or the interior polygon, &c. When the sides, and the angles, are all equal, it is called a regular Figure; but when unequal, an irregular one.

FIGURE, in Geomancy, is applied to the extremes of points, lines, or numbers, thrown or cast at random: on the combinations or variations of which, the fates of this art found their divinations.

FIGURE, in Geometry, denotes a surface or space inclosed on all sides; and is either superficial or solid; superficial when it is inclosed by lines, and solid when it is inclosed or bounded by surfaces.

Figures are either straight, curved, or mixed, according as their bounds are straight, or curved, or both.—The exterior bounds of a Figure, are called its sides; the lowest side, its base; and the angular point opposite the base, the vertex of the Figure; also its height, is the distance of the vertex from the base, or the perpendicular let fall upon it from the vertex.

For Figures, equal, equiangular, equilateral, circumscribed, inscribed, plane, regular, irregular, similar, &c; see the respective adjectives.

Apparent FIGURE, in Optics, that Figure, or shape, under which an object appears, when viewed at a distance. This is often very different from the true figure; for a straight line viewed at a distance may appear but as a point; a surface as a line; a solid as a surface; and a crooked figure as a straight one. Also, each of these may appear of different magnitudes, and some of them of different shapes, according to their situation with regard to the eye. Thus an arch of a circle may appear a straight line; a square or parallelogram, a trapezium, or even a triangle; a circle, an ellipsis; angular magnitudes, round; a sphere, a circle; &c.

Also any small light, as a candle, seen at a distance in the dark, will appear magnified, and farther off than it really is. Add to this, that when several objects are seen at a distance, under angles that are so small as to be insensible, as well as each of the angles subtended by

any one of them, and that next to it; then all these objects appear not only as contiguous, but as constituting and seeming but one continued magnitude.

FIGURE of the Sines, Cosines, Versed-sines, Tangents, or Secants, &c, are Figures made by conceiving the circumference of a circle extended out in a right line, upon every point of which are erected perpendicular ordinates equal to the Sines, Cosines, &c, of the corresponding arcs; and then drawing the curve line through the extremity of all these ordinates; which is then the Figure of the Sines, Cosines, &c.

It would seem that these Figures took their rise from the circumstance of the extension of the meridian line by Edward Wright, who computed that line by collecting the successive sums of the secants, which is the same thing as the area of the Figure of the secants, this being made up of all the ordinates, or secants, by the construction of the Figure. And in imitation of this, the Figures of the other lines have been invented. By means of the Figure of the secants, James Gregory shewed how the logarithmic tangents may be constructed, in his Exercitationes Geometricæ, 4to, 1668.


#### Construction of the Figures of Sines, Cosines, &c.

Let ADB &c (fig. 1) be the circle, AD an arc, DE its sine, CE its cosine, AE the versed sine, AF the tangent, GH the cotangent, CF the secant, and CH the cosecant. Draw a right line *aa* equal to the whole circumference ADGBA of the circle, upon which lay off also the lengths of several arcs, as the arcs at every  $10^\circ$ , from 0 at *a*, to  $360^\circ$  at the other end at *a*; upon these points raise perpendicular ordinates, upwards or downwards, according as the sine, cosine, &c, is affirmative or negative in that part of the circle; lastly, upon these ordinates set off the length of the sines, cosines, &c, corresponding to the arcs at those points of the line or circumference *aa*, drawing a curve line through the extremities of all these ordinates; which will be the Figure of the sines, cosines, versed-sines, tangents, cotangents, secants, and cosecants, as in the annexed Figures. Where it may be observed, that the following curves are the same, viz, those of the sines and cosines, those of the tangents and cotangents, and those of the secants and cosecants; only some of their parts a little differently placed.

Fig. 1.




Sines.



Cosines.



Versed-sines.




Tangents.



Tangents.

Cotangents.



Secants.

Cofecants.



It may be known when any of these lines, viz, the sines, cosines, &c, are affirmative or negative, i. e. to be set upwards or downwards, by observing the following general rules for those lines in the 1st, 2d, 3d, and 4th quadrants of the circle.

The sines	in the 1st and 2d are affirmative,	in the 3d and 4th negative:
The cosines	in the 1st and 4th are affirmative,	in the 2d and 3d negative:
The tangents	in the 1st and 3d are affirmative,	in the 2d and 4th negative:
The cotangents	in the 1st and 3d are affirmative,	in the 2d and 4th negative:
The secants	in the 1st and 4th are affirmative,	in the 2d and 3d negative:
The cofecants	in the 1st and 2d are affirmative,	in the 3d and 4th negative:

And all the versed sines are affirmative.

To find the Equation and Area, &c, to each of these Curves.

Draw any ordinate  $de$ ; putting  $r$  = the radius  $AC$  of the given circle,  $x = ad$  or  $AD$  any absciss or arc, and  $y = de$  its ordinate, which will be either the sine  $DE = s$ , cosine  $CE = c$ , versed sine  $AE = v$ , tangent  $AF = t$ , cotangent  $GH = \tau$ , secant  $CF = f$ , or cofecant  $CH = \sigma$ , according to the nature of the particular construction. Now, from the nature of the circle, are obtained these following general equations, expressing the relations between the fluxions of a circular arc and its sine, or cosine, &c.

$$\dot{x} = \frac{rs}{\sqrt{r^2 - s^2}} = \frac{-rc}{\sqrt{r^2 - c^2}} = \frac{rv}{\sqrt{2rv - vv}} = \frac{r^2 \dot{t}}{r^2 + t^2}$$

$$= \frac{-r^2 \dot{\tau}}{r^2 + \tau^2} = \frac{r^2 \dot{f}}{f \sqrt{f^2 - r^2}} = \frac{-r^2 \dot{\sigma}}{\sigma \sqrt{\sigma^2 - r^2}}.$$

And these also express the relation between the absciss and ordinate of the curves in question, each in the order in which it stands; where  $x$  is the common absciss to all of them, and the respective ordinates are  $s$ ,  $c$ ,  $v$ ,  $t$ ,  $\tau$ ,  $f$ , and  $\sigma$ . And hence the area &c, of any of these curves may be found, as follows:

1. In the FIGURE of Sines.—Here  $x = ad$ , and  $s =$  the ordinate  $de$ ; and the equation of the curve, as

above, is  $\dot{x} = \frac{rs}{\sqrt{r^2 - s^2}}$ . Hence the fluxion of the

area, or  $s\dot{x}$  is  $\frac{rs\dot{s}}{\sqrt{r^2 - s^2}}$ ; the correct fluent of which is

$r^2 \mp r\sqrt{r^2 - s^2} = r^2 - rc = rv$  the rectangle of radius and vers. i. e.  $-$  or  $+$  as  $s$  is increasing or decreasing; which is a general expression for the area  $ade$  in the Figure of sines. When  $s = 0$ , as at  $a$  or  $b$ , this expression becomes 0 or  $2r^2$ ; that is 0 at  $a$ , and  $2r^2 =$  the area  $acb$ ; or  $r^2 =$  the area of  $afg$  when  $ad$  becomes a quadrant  $af$ .

2. In the FIGURE of Cosines.—Here  $x = ad$ , and  $c = de$ ; and the equation of the curve is

$\dot{x} = \frac{-rc}{\sqrt{r^2 - c^2}}$ . Hence the fluxion of the area is

$c\dot{x} = \frac{-rcc}{\sqrt{r^2 - c^2}}$ ; and the fluent of this is  $r\sqrt{r^2 - c^2}$

$= rs$ , the rectangle of radius and sine, for the general area  $adec$ . When  $s = r$ , or  $c = 0$ , this becomes  $r^2 =$  the area  $afc$ , whose absciss  $af$  is equal to a quadrant of the circumference; the same as in the Figure of the sines, upon an equal absciss.

3. In the FIGURE of Versed sines.—Here  $x = ad$ , and  $v = de$ ; and the equation of the curve is

$\dot{x} = \frac{rv}{\sqrt{2rv - vv}}$ . Hence the fluxion of the area is

$v\dot{x} = \frac{rv\dot{v}}{\sqrt{2rv - vv}} = \frac{r\dot{v}\sqrt{v}}{\sqrt{2r - v}}$ ; and the fluent of

this is  $rx - rs = r \times AD - DE$  for the area  $ade$  in the Figure of versed sines. When  $AD$  or  $ad$  is a quadrant  $AG$  or  $af$ , this becomes  $\frac{3 \cdot 1416}{2} r^2 - r^2 =$  the area  $afg$ . And when  $AD$  or  $ad$  is a semicircle  $ab$ , it becomes  $3 \cdot 1416 r^2 =$  the area  $abg$  in the Figure of versed sines.

4. In the FIGURE of Tangents.—Here  $x = ad$ , and

$t = de$ ; and the equation of the curve is  $\dot{x} = \frac{r^2 \dot{t}}{r^2 + t^2}$ .

Hence the fluxion of the area is  $t\dot{x} = \frac{r^2 t \dot{t}}{r^2 + t^2}$ ; and

the correct fluent of this is  $\frac{1}{2} r^2 \times \text{hyp. log. of } \frac{r^2 + t^2}{r^2} = r^2 \times \text{hyp. log. of } \frac{\sqrt{r^2 + t^2}}{r}$

of  $\frac{f}{r}$ . And hence the Figure of the tangents may be used for constructing the logarithmic secants; a property that was remarked by Gregory at the end of his Exercit. Geomet.

When  $ad$  becomes a quadrant  $af$ ,  $t$  being then infinite, this becomes infinite for the area  $afg$ . And the same for the Figure of cotangents, beginning at  $f$  instead of  $a$ .



5. For the *FIGURE of the Secants*.—Here  $x = ad$ , and  $f = de$ ; and the equation of the curve is  $\dot{x} = \frac{r^2 \dot{f}}{f \sqrt{f^2 - r^2}}$ .

Hence the fluxion of the area is  $f \dot{x} = \frac{r^2 \dot{f}}{\sqrt{f^2 - r^2}}$ ; the

fluent of which is  $r^2 \times \text{hyp. log. of } \frac{f + \sqrt{f^2 - r^2}}{r}$

for the general area  $ade$ . And when  $ad$  becomes the quadrant  $af$ , this expression becomes infinite for the area  $afg$ .

The same process will serve for the Figure of cosecants, beginning at  $f$  instead of  $a$ .

From hence the meridional parts in Mercator's chart may be calculated for any latitude  $AD$  or  $ad$ : For the merid. parts : are to the arc of latitude  $AD$  :: as the sum of the secants : to the sum of as many radii or :: as the area  $ade$  : to  $ad \times \text{radius } ac$  or  $AD \times AC$  in the first figure.

FILLET, in Architecture, any little square member or ornament used in crowning a larger moulding.

FINÆUS (ORONTIUS), in French *Finé*, professor of mathematics in the Royal-college of Paris, was the son of a physician, and was born at Briançon in Dauphiné in 1494. He went young to Paris, where his friends procured him a place in the college of Navarre. He applied himself there to philosophy and polite literature; but more especially to mathematics, in which, having a natural propensity, he made a considerable proficiency. Particularly he made a good progress in mechanics; in which, having both a genius to invent instruments, and a skilful hand to make them, he gained much reputation by the specimens he gave of his ingenuity.

Finæus first made himself publicly known by correcting and publishing Siliceus's *Arithmetic*, and the *Margarita Philosophica*. He afterwards read private lectures in mathematics, and then taught that science publicly in the college of Gervais: from the reputation of which, he was recommended to Francis the 1st, as the properest person to teach mathematics in the new college which that prince had founded at Paris. And here, though he spared no pains to improve his pupils, he yet found time to publish a great many books upon most parts of the mathematics. But neither his genius, his labours, his inventions, and the esteem which numberless persons shewed him, could secure him from that fate which so often befalls men of letters. He was obliged to struggle all his life-time with poverty; and when he died, left a numerous family deeply in debt. However, as merit must always be esteemed in secret, though it seldom has the luck to be rewarded openly; so Finæus's children found Mecænaſes, who for their father's sake assisted his family.—He died in 1555, at 61 years of age.

Like all other mathematicians and astronomers of those times, he was greatly addicted to astrology; and had the misfortune to be a long time imprisoned for having predicted some things, that were not acceptable to the court of France. He was also one of those, who vainly boasted of having found out the quadrature of the circle. An edition of his works, translated into the

Italian language, was published in 4to, at Venice, 1587; consisting of Arithmetic, Practical Geometry, Cosinography, Astronomy, and Dialling.

FINITE, the property of any thing that is bounded or limited, either in its power, or extent, or duration, &c; as distinguished from the property of infinite, or without bounds.

FINITOR, the horizon; being so called, because it finishes or bounds the sight or prospect.

FIRE, is that subtle invisible cause by which bodies are made hot to the touch, and expanded or enlarged in bulk; by which fluids are rarefied into vapour; or solid bodies become fluid, and at last either dissipated and carried off in vapour, or else melted into glass. It seems also to be the chief agent in nature on which animal and vegetable life have an immediate dependence.

The disputes concerning fire, which long existed among philosophers, have now in a great measure subsided. Those celebrated philosophers of the last century, Bacon, Boyle, and Newton, were of opinion, that Fire was not a substance distinct from other bodies, but that it entirely consisted in the violent motion of the parts of any body, which was produced by the mechanical force of impulsion, or of attrition. So Boyle says, when a piece of iron becomes hot by hammering, "there is nothing to make it so, except the forcible motion of the hammer impressing a vehement and variously determined agitation on the small parts of the iron." And Bacon defines heat, which he makes synonymous with Fire, an "expansive undulatory motion in the minute particles of a body, whereby they tend with some rapidity from a centre towards a circumference, and at the same time a little upwards." And according to Newton, Fire is a body heated so hot as to emit light copiously; for what else, says he, is red-hot iron, but Fire? and what else is a fiery coal than red-hot wood? by which he suggests, that bodies which are not Fire, may be changed and converted into it.

On the other hand, the chemists strenuously contended that Fire was a fluid of a certain kind, distinct from all others, and universally present throughout the whole globe. Boerhaave particularly maintained this doctrine; and in support of it brought this argument, that flint and steel would strike fire, and produce the same degree of heat in Nova Zembla as they would do under the equator. Other arguments were drawn from the increased weight of metallic calces, which they thought proceeded from the fixing of the element of Fire in the substance whose weight was thus increased. For a long time however, the matter was most violently disputed; but the mechanical philosophers at last prevailed through the deference paid to the principles of Newton, though he himself had scarcely taken any active part in the contest.

The experiments of Dr. Black however seemed to bring the dispute to a decision, and that in favour of the chemists, concerning what he called latent-heat. From these discoveries it appears, that Fire may exist in bodies in such a manner, as not to discover itself in any other way than by its action on the minute parts of the body; but that suddenly this action may be changed in such a manner, as no longer to be directed upon the particles of the body itself, but upon external objects; in which case we then perceive its action by our sense of feeling,



feeling, or discover it by the thermometer, and call it heat, or sensible Fire.

From this discovery, and others in electricity, it is now pretty generally allowed, that Fire is a distinct fluid, capable of being transferred from one body to another. But when this was discovered, another question no less perplexing arose, viz, what kind of a fluid it was; or whether it bears any analogy to those with which we are better acquainted. Now there are found two fluids, viz, the solar light, and the electric matter, both of which occasionally act as fire, and which therefore seem likely to be all the same at the bottom; and popularly the matter has been long since determined; the solar rays and the electric fluid having been indifferently accounted elementary Fire. Some indeed have imagined both these fluids to be mere phlogiston itself, or at least containing a large portion of it; and Mr. Scheele went so far in this way as to form an hypothesis, which he endeavoured to support by experiments, that Fire is composed of phlogiston and dephlogisticated air. But it is now ascertained beyond doubt, that the result of such a combination is not Fire, but fixed air.

It was long since observed by Newton, that heat was certainly conveyed by a medium more subtle than the common air; for two thermometers, one included in the vacuum of an air-pump, the other placed in the open air, at an equal distance from the fire, would grow equally hot in nearly the same time. This and other experiments shew, that Fire exists and acts where there is no other matter, and of consequence it is a fluid per se, independent of every terrestrial substance, without being generated or compounded of any thing we are yet acquainted with. To determine the nature of the fluid, we have only to consider whether any other can be discovered which will pass through the perfect vacuum just mentioned, and act there as Fire. Such a fluid is found in the solar light, which is well known to act even in vacuo as the most violent Fire. The solar light will likewise act in the very same manner in the most intense cold; for M. de Saussure has found, that on the cold mountain top the sun-beams are equally powerful as on the plain below, if not more so. It appears therefore, that the solar light will produce heat independent of any other substance whatever; that is, where no other body is present, at least as far as we can judge, except the light itself, and the body to be acted upon. We cannot therefore avoid concluding, that a certain modification of the solar light is the cause of heat, expansion, vapour, &c, and answers to the rest of the characters given in the foregoing definition of Fire, and that independent of any other substance whatever.

It is very probable too, that the electric matter is no other than the solar light absorbed by the earth, and thus becoming subject to new laws, and assuming many properties apparently different from what it has when it acts as light. Even in this case it manifests its identity with Fire or light, viz, by producing a most intense heat where a large quantity of it passes through a small space. So that at any rate, the experiments which have already been made, and the proofs drawn from the phenomena of nature, shew such a strong affinity between the elements of Fire, light and electricity, that we may not only assert their identity upon the

most probable grounds, but lay it down as a position against which at present no argument of any weight has an existence.

*FIRE-Arrow*, is a small iron dart, furnished with springs and bars, and also a match impregnated with sulphur and powder, which is wound about its shaft. It is chiefly used by privateers and pirates to fire the sails of the enemy's ship, and for this purpose it is discharged from a musketoon, or a swivel gun. The match being kindled by the explosion, it communicates the flame to the sail against which it is directed, where the arrow fastens itself by means of its bars and springs. This weapon is peculiar to hot climates, particularly the West Indies; the sails being very dry, are quickly set on Fire, and the Fire is soon conveyed to the masts, rigging, and finally to the vessel itself.

*FIRE-Balls*, in Artillery, are certain balls composed of combustible matters, such as fine or mealed powder, sulphur, saltpetre, rosin, pitch, &c. These are thrown into the enemy's works in the night time, to discover where they are; or to set Fire to houses, galleries, or blinds of the besiegers, &c. They are sometimes armed with spikes or hooks of iron, that they may not roll off, but stick or hang where they are to take effect.

*FIRE-Balls*, or *Fiery-Meteors*, in Meteorology, a kind of luminous bodies usually appearing at a great height above the earth, with a splendor surpassing that of the moon, and sometimes apparently as large. They have not been found to observe any regular course or motion, but, on the contrary, moving in all directions, and with very different degrees of celerity; frequently breaking into several smaller ones; sometimes making a strong hissing sound, sometimes bursting or vanishing with a loud report, and sometimes not.

These luminous appearances doubtless constitute one part of the ancient prodigies, blazing stars, or comets, which last they sometimes resemble in being attended with a train; but more often they appear round. The first of these of which we have any accurate account, was observed by Dr. Halley, and some other philosophers at different places, in the year 1719, the height of which above the surface of the earth was computed at more than 70 miles. Many others have been pretty accurately observed since that time, and described by philosophers, as in the French Memoirs, and in the Philos. Trans. vols. 30, 41, 42, 43, 46, 47, 48, 51, 53, 54, 63, 74, &c. The velocities, directions, appearances, and heights of all these were found to be very various; though the height of all of them was supposed above the limits assigned to our atmosphere, or where it loses its refractive power. The most remarkable of those on record, appeared on the 18th of August 1783, about 9 o'clock in the evening. It was seen to the northward of Shetland, and took a south-easterly direction for an immense space, being observed as far as the southern provinces of France, and by some it was said to have been seen at Rome, passing over a space of 1000 miles in about half a minute of time, and at a very great height. During its course it appeared several times to change its shape; sometimes appearing in the form of one ball, sometimes of two or more; sometimes with a train, and sometimes without one.

There are divers opinions concerning the nature and origin



origin of these Fire-balls. The first thing that occurred to philosophers on this subject was, that the meteors in question were burning bodies rising from the surface of the earth, and flying through the atmosphere with great rapidity. But this hypothesis was soon rejected, on considering that there was no power known by which such bodies could either be raised to a sufficient height, or projected with the velocity of the meteors. The next hypothesis was, that, instead of one single body, they consist of a train of sulphureous vapours, extending a vast way through the atmosphere, and being kindled at one end, display the luminous appearances in question by the fire running from one end of the train to the other. But it is not easy to conceive how such matters can exist and be disposed in such lines in so rare a part of the atmosphere, and even to burn there, in an almost perfect vacuum. For which reason this hypothesis was abandoned, for another, which was, that those meteors are permanent solid bodies, not rising from the earth, but revolving round it in very excentric orbits, and thus in their perigeon moving with vast rapidity. But as the various appearances of one and the same meteor, to observers at different places, are not compatible with the idea of a single body so revolving, this hypothesis has also been given up in its turn. Another hypothesis that has sometimes been advanced is this; viz, that these meteors are a kind of bodies that take fire as soon as they come within the atmosphere of the earth. But this cannot be supposed without implying a previous knowledge of these bodies, which it is impossible we can have. The only opportunity we can have of seeing them, is when they are on fire. Before that time they are in an invisible and unknown state; and it is surely improper to argue concerning them in this state, or pretend to determine any one of their properties, when it is not in our power at all to see or investigate them. As these meteors therefore never manifest themselves to our senses but when they are on fire, the only rational conclusion we can draw from thence is, that they have no existence in any other state; and consequently that their substance must be composed of that fluid which, when acting after a certain manner, becomes luminous and shews itself as fire; remaining invisible and eluding our researches in every other case. On this hypothesis, it is now pretty generally concluded, that the Fire-balls are great bodies of electric matter, moving from one part of the heavens where, to our conception, it is superabundant, to another where it is deficient: a conclusion attended with much probability, from the analogy observed between electricity and the phenomena of these meteors: and hence these Fire-balls appear to be of the same family with shooting-stars, lightning, the aurora-borealis, &c, being all referred to the same origin, viz, the electricity of the atmosphere.

*FIRE-Engine*, is a machine for extinguishing accidental Fires by means of a stream or jet of water. The common squirting Fire-engine consists of a lifting pump placed in a vessel of water, and wrought by two levers that act always together. During the stroke, the water raised by the piston of the pump spouts forcibly through a pipe joined to the pump-barrel, and made capable of any degree of elevation by means of a yield-

ing leather pipe, or by a ball and socket turning every way, screwed on the top of the pump. The vessel containing the water is covered with a strainer, to prevent the mud, &c, which is poured into it with the water, from choking the pump-work. Between the strokes of this engine the stream is discontinued, for want of an air vessel. However, in some cases, Engines of this construction have their use, because the stream, though interrupted, is much smarter than when the engine is made to throw water in a continued stream. See these Engines particularly described in Desaguliers's Exper. Philos. vol. 2, pa. 505; or Martin's Philos. Britan. vol. 2, pa. 69. See also the figure of them, plate viii. fig. 3.

*FIRE-Engine*, is also sometimes used for the machine employed in raising water by steam, and more properly called *STEAM-Engine*; which see.

*FIRE-Lock*, or *Fusil*, a small gun or musket, which fires with a flint and steel; as distinguished from the old musket, or match-lock, which was fired with a match. The Fire-lock is now in common use with the European armies, and carried by the foot-soldiers. It is usually about 3 feet 8 inches in the barrel; and, including the stock, 4 feet 8 inches, carrying a leaden bullet, of which 29 make 2lb. The diameter of the ball is .55, and that of the barrel .56 parts of an inch. The time of the invention of Fire-locks is uncertain; but they were used in 1690.

*FIRE-Places*, are contrivances for communicating heat to rooms, and also for answering various purposes of art and manufacture.

The general properties of air and fire, on which their construction chiefly depends, are the following, viz, that air is rarefied by heat, and condensed by cold; i. e. the same quantity of air takes up more space when warm than when cold. Air rarefied and expanded by heat, is specifically lighter than it was before, and will rise in other air of greater density: so that a fire being made in any chimney, the air about and over the fire is rarefied by the heat, thence becomes lighter, and so rises in the funnel, and goes out at the top of the chimney: the other air in the room, flowing towards the chimney, supplies its place, is then rarefied in its turn, and rises likewise; and the place of the air thus carried out of the room, is supplied by fresh air coming in through doors and windows, or, if they be shut, through every crevice with violence; or if the avenues to the room be so closed up, that little or no fresh supply of air can be obtained, the current up the funnel must flag; and the smoke, no longer driven up, float about in the room.

Upon these principles, various contrivances and kinds of Fire-grates and stoves have been devised, from the old very open and wide chimney places, down to the present modish ones, which are much narrowed in the front, opening, by side and back jambs, and a low breast or mantle, besides the convenience of a flap, called a register, that covers the top of the Fire-stove, but opening to any degree with a small winch, which lifts the back part sloping upwards, and so throws the smoke freely up the funnel, and yet admitting as little air to pass as you please; by which simple means the warm air is kept very much in the room, while the very narrow and sloping orifice promotes the brisk ascent of



the smoke, and yet prevents its return down again, for the same reason.

Another very ingenious, but more complex, apparatus, called the Pennsylvania Fire-place, was invented by Dr. Franklin, by which a room is kept very warm by a constant supply of fresh hot air, that passes into it through the stove itself. See the description in his Letters and Papers on Philosophical Subjects.

*FIRE-Pot*, in the Military Art, is a small earthen pot, into which is put a grenade, filled with fine powder till the grenade be covered; the pot is then covered with a piece of parchment, and two pieces of match laid across and lighted. This pot being thrown where it is designed to do execution, breaks and fires the powder, and this again fires the powder in the grenade, which ought to have no fuze, that its operation may be the quicker.

*Rasant*, or *Razant FIRE*, is a fire from the artillery and small arms, directed parallel to the horizon, or to those parts of the works of a place that are defended.

*Running FIRE*, is when ranks of men fire one after another; or when the lines of an army are drawn out to fire on account of a victory; in which case each squadron or battalion takes the fire from that on its right, from the right of the first line to the left, and from the left to the right of the second line, &c.

*FIRE-Ships*, in the Navy, are vessels charged with combustible materials or artificial Fire-works; which having the wind of an enemy's ship, grapple her, and set her on fire.

Anderson, in his History of Commerce, vol. 1, p. 432, ascribes the invention to the English, in this instance, viz, some vessels being filled with combustible matter, and sent among the Spanish ships composing the Invincible Armada in 1588; and hence arose it is said the terrible invention of Fire-ships.

But Livy informs us, that the Rhodians had invented a kind of Fire-ships, which were used in junction with the Roman fleet in their engagement with the Syrians, in the year 190 before Christ: cauldrons of combustible and burning materials were hung out at their prows, so that none of the enemies' ships durst approach them: for these fell on the enemies' galleys, struck their beaks into them, and at the same time set them on fire. Livy, lib. 37, cap. 30.

*Wild-FIRE*, is a kind of artificial or factitious fire, that burns even under water, and that with greater violence than out of it. It is composed of sulphur, naphtha, pitch, gum, and bitumen, and it is only extinguishable by vinegar, mixed with sand and urine, or by covering it with raw hides. It is said its motion is contrary to that of natural fire, always following the direction in which it is thrown, whether it be downwards, sideways, or otherwise.

The French call it Greek Fire, or Feu Gregeois, because first used by the Greeks about the year 660, as is observed by the Jesuit Petavius, on the authority of Nicetas, Theophanes, Cedrenus, &c. The inventor, according to the same author, was an engineer of Heliopolis, in Syria, named Callinicus, who first applied it in the sea-fight commanded by Constantine Pogonates, against the Saracens, near Cyzicus, in the Hellespont; and with such effect, that he burnt the whole fleet, which contained 30,000 men. But others refer it to

a much older date, and ascribe the invention to Marcus Gracchus; an opinion which is supported by several passages, both in the Greek and Roman writers; which shew that it was anciently used by both these nations in their wars. See Scaliger against Cardan.

The successors of Constantine used it on several occasions, with great advantage: and it is remarkable that they were able to keep the secret of the composition to themselves; so that no other nation knew it in the year 960.

It is recorded by Chorier, in his Hist. de Dauph. that Hugh, king of Burgundy, demanding ships of the emperor Leo for the siege of Fresne, desired also the Greek Fire.

And F. Daniel gives a good description of the Greek Fire, in his account of the siege of Damietta, under St. Louis. Every body, says he, was astonished with the Greek Fire, which the Turks then prepared; and the secret of which is now lost. They threw it out of a kind of mortar, and sometimes shot it with an odd sort of cross-bow, which was strongly bent by means of a handle, or winch, of much greater force than the bare arm. That which was thrown from the mortar sometimes appeared in the air of the size of a tun, with a long tail, and a noise like that of thunder. The French, by degrees, got the secret of extinguishing it; in which they succeeded several times.

After all, perhaps the invention of the Wild-Fire is to be ascribed to other nations, and to a still older date, and that it was the same as that used among the Indians in Alexander's invasion, when it was said they fought with thunder and lightning, or shot Fire with a terrible noise.

*FIRE-Works*, otherwise called Pyrotechnia, are artificial Fires, or preparations made of gunpowder, sulphur, and other inflammable and combustible ingredients, used on occasion of public rejoicings, and other solemnities. The principal of these are rockets, serpents, stars, hail, mines, bombs, garlands, letters, and other devices.

The invention of Fire-works is attributed, by M. Mahudel, to the Florentines and people of Sienna; who found out likewise the method of adding to them decorations of statues, with fire issuing from their eyes and mouths.

*FIRKIN*, an English measure of capacity; being the 4th part of a barrel; and containing 8 gallons of ale, soap, butter, or herrings; or 9 gallons of beer.

*FIRLOT*, a dry measure used in Scotland. The oat-firLOT contains 21½ pints of that country, or about 85 English pints; and the barley-firLOT, 31 standard pints. The wheat firLOT contains about 2211 cubic inches; and therefore exceeds the English bushel by 60 cubic inches, or almost an English quart.

*FIRMAMENT*, by some old astronomers, is the orb of the fixed stars, or the highest of all the heavens. But in scripture and common language it is used for the middle regions, or the space or expanse appearing like an arch quite around or above us in the heavens. Many ancients and moderns also accounted the Firmament a fluid matter; but those who gave it the name of Firmament must have taken it for a solid one.

*FIRMNESS*, is the consistence of a body; or that state when its sensible parts cohere, or are united together, so that the motion of one part induces a motion



tion of the rest. In which sense firmness stands opposed to fluidity.

The firmness of bodies then depends on the connexion or cohesion of their particles; and the cause of cohesion the Newtonians hold to be an attractive force, inherent in bodies, which binds their small particles together; exerting itself only at the points of contact, or extremely near them, and vanishing at greater distances.

FIRST *Mover*, in the old Astronomy, is the Primum Mobile, or that which gives motion to the other parts of the universe.

FISSURES, in the History of the Earth, are certain interruptions, mostly parallel to each other, that divide or separate the strata of it from one another, in nearly horizontal directions; and the parts of the same stratum in nearly vertical directions.

FIXED *Line of Defence*, a line drawn along the face of the bastion, and terminating in the curtain.

FIXED *Signs of the Zodiac*, according to some, are the four signs Taurus, Leo, Scorpio, Aquarius. They are so called because the sun passes them respectively in the middle of each quarter, when that season is more settled and Fixed than under the signs which begin and end it.

FIXED *Stars*, are such as constantly retain the same position and distance with respect to each other; by which they are contradistinguished from erratic or wandering stars, which are continually varying their situation and distance.—The Fixed Stars only are properly and absolutely called stars; the rest having their peculiar denomination of planet or comet.

FIXITY, or FIXEDNESS, the quality of a body which determines it fixed; or a property which enables it to endure the fire and other violent agents.

A body may be said to be fixed in two respects: 1st, When on being exposed to the fire, or a corrosive menstruum, its particles are indeed separated, and the body rendered fluid, but without being resolved into its first elements. The 2d, when the body sustains the active force of the fire or menstrooms whilst its integral parts are not carried off in fumes. Each kind of Fixity is the result of a strong or intimate cohesion between the particles.

FLAME, the subtlest and brightest part of the fuel, ascending above it in a pyramidal or conical figure, and heated red-hot. Sir Isaac Newton defines Flame as only red-hot smoke, or the vapour of any substance raised from it by fire, and heated to such a degree as to emit light copiously. Is not Flame, says he, a vapour, fume, or exhalation, heated red-hot; that is, so hot as to shine? For bodies do not Flame without emitting a copious fume; and this fume burns in the Flame. The ignis fatuus is a vapour shining without heat; and is there not the same difference between this vapour and Flame, as between rotten wood shining without heat, and burning coals of fire? In distilling hot spirits, if the head of the still be taken off, the vapour which ascends will take fire at the Flame of a candle, and turn into Flame. Some bodies, heated by motion or fermentation, if the heat grow intense, fume copiously; and if the heat be great enough, the fumes will shine, and become Flame. Metals in fusion do not Flame, for want of a copious fume. All flaming bodies, as oil, tallow, wax, wood, fossil coal, pitch, sul-

phur, &c, by burning, waste in smoke, which at first is lucid; but at a little distance from the body ceases to be so, and only continues hot. When the Flame is put out, the smoke is thick, and frequently smells strongly: but in the Flame it loses its smell; and, according to the nature of the fuel, the Flame is of divers colours. That of sulphur and spirit of wine is blue; that of copper opened with sublimite, green; that of tallow, yellow; of camphire, white; &c. Newton's Optics, p. 318.

FLAMSTEED (JOHN), an eminent English astronomer, being indeed the first astronomer royal, for whose use the royal observatory was built at Greenwich, thence called Flamsteed House. He was born at Denby in Derbyshire the 19th of August 1646. He was educated at the free school of Derby, where his father lived; and at 14 years of age was afflicted with a severe illness, which rendered his constitution tender ever after, and prevented him then from going to the university, for which he was intended. He nevertheless prosecuted his school education with the best effect; and then, in 1662, on quitting the grammar school, he pursued the natural bent of his genius, which led him to the study of astronomy, and closely perused Sacrobosco's book *De Sphæra*, which fell in his way, and which laid the groundwork of all that mathematical and astronomical knowledge, for which he became afterward so justly famous. He next procured other more modern books of the same kind, and among them Streete's *Astronomia Carolina*, then lately published, from which he learned to calculate eclipses and the planets' places. Some of these being shewn to a Mr. Halton, a considerable mathematician, he lent him Riccioli's *Almagestum Novum*, and Kepler's *Tabulæ Rudolphinæ*, which he profited much by. In 1669, having calculated some remarkable eclipses of the moon, he sent them to lord Brouncker, president of the Royal Society, which were greatly approved by that learned body, and procured him a letter of thanks from Mr. Oldenburg their secretary, and another from Mr. John Collins, with whom, and other learned men, Mr. Flamsteed for a long time afterwards kept up a correspondence by letters on literary subjects. In 1670, his father, observing he held correspondence with these ingenious gentlemen, advised him to take a journey to London, to make himself personally acquainted with them; an offer which he gladly embraced, and visited Mr. Oldenburg and Mr. Collins, who introduced him to Sir Jonas Moore, which proved the means of his greatest honour and preferment. He here got the knowledge and practice of astronomical instruments, as telescopes, micrometers, &c. On his return, he called at Cambridge, and visited Dr. Barrow, Mr. Isaac Newton, and other learned men there, and entered himself a student of Jesus College. In 1672 he extracted several observations from Mr. Gascoigne's and Mr. Crabtree's letters, which improved him greatly in dioptrics. In this year he made many celestial observations, which, with calculations of appulses of the moon and planets to fixed stars for the year following, he sent to Mr. Oldenburg, who published them in the *Philos. Transf.*

In 1673, Mr. Flamsteed wrote a small tract concerning the true diameters of all the planets, when at their greatest and least distances from the earth; which he lent to Mr. Newton in 1685, who made some use of it



it in the 4th book of his *Principia*.—In 1674 he wrote an ephemeris, to shew the falsity of Astrology, and the ignorance of those who pretended to it: with calculations of the moon's rising and setting; also occultations and appulses of the moon and planets to the fixed stars. To which, at Sir Jonas Moore's request, he added a table of the moon's southings for that year; from which, and from Philips's theory of the tides, the high-waters being computed, he found the times come very near. In 1674 too, he drew up an account of the tides, for the use of the king. Sir Jonas also shewed the king, and the duke of York, some barometers and thermometers that Mr. Flamsteed had given him, with the necessary rules for judging of the weather; and otherwise took every opportunity of speaking favourably of Flamsteed to them, till at length he brought him a warrant to be the king's astronomer, with a salary of 100l. per annum, to be paid out of the office of Ordnance, because Sir Jonas was then Surveyor General of the Ordnance. This however did not abate our author's propensity for holy orders, and he was accordingly ordained at Ely by bishop Gunning.

On the 10th of August, 1675, the foundation of the Royal Observatory at Greenwich was laid; and during the building of it, Mr. Flamsteed's temporary observatory was in the queen's house, where he made his observations of the appulses of the moon and planets to the fixed stars, and wrote his *Doctrine of the Sphere*, which was afterward published by Sir Jonas, in his *System of the Mathematics*.

About the year 1684 he was presented to the living of Burflow in Surry, which he held as long as he lived. Mr. Flamsteed was equally respected by the great men his contemporaries, and by those who have succeeded since his death. Dr. Wotton, in his *Reflections upon Ancient and Modern Learning*, styles our author one of the most accurate Observers of the Planets and Stars, and says he calculated tables of the eclipses of the several satellites, which proved very useful to the astronomers. And Mr. Molyneux, in his *Dioptrica Nova*, gives him a high character; and, in the admonition to the reader prefixed to the work, observes, that the geometrical method of calculating a ray's progress is quite new, and never before published; and for the first hint of it, says he, I must acknowledge myself obliged to my worthy friend Mr. Flamsteed. He wrote several small tracts, and had many papers inserted in the *Philosophical Transactions*, viz, several in almost every volume, from the 4th to the 29th, too numerous to be mentioned in this place particularly.

But his great work, and that which contained the main operations of his life, was the *Historia Cælestis Britannica*, published in 1725, in 3 large folio volumes. The first of which contains the observations of Mr. William Gascoigne, the first inventor of the method of measuring angles in a telescope by means of screws, and the first who applied telescopical sights to astronomical instruments, taken at Middleton, near Leeds in Yorkshire, between the years 1638 and 1643; extracted from his letters by Mr. Crabtree; with some of Mr. Crabtree's observations about the same time; and also those of Mr. Flamsteed himself, made at Derby between the years 1670 and 1675; besides a multitude of curious observations, and necessary tables to be used with

them, made at the Royal Observatory, between the years 1675 and 1689.—The 2d volume contains his observations, made with a mural arch of near 7 feet radius, and 140 degrees on the limb, of the meridional zenith distances of the fixed stars, sun, moon, and planets, with their transits over the meridian; also observations of the diameters of the sun and moon, with their eclipses, and those of Jupiter's satellites, and variations of the compass, from 1689 to 1719: with tables shewing how to render the calculation of the places of the stars and planets easy and expeditious. To which are added, the moon's place at her oppositions, quadratures, &c; also the planets' places, derived from the observations.—The 3d volume contains a catalogue of the right-ascensions, polar-distances, longitudes, and magnitudes of near 3000 fixed stars, with the corresponding variations of the same. To this volume is prefixed a large preface, containing an account of all the astronomical observations made before his time, with a description of the instruments employed; as also of his own observations and instruments; with a new Latin version of Ptolemy's catalogue of 1026 fixed stars; and Ulegh-beig's places annexed on the Latin page, with the corrections: a small catalogue of the Arabs: Tycho Brahe's of about 780 fixed stars: the Landgrave of Hesse's of 386: Hevelius's of 1534: and a catalogue of some of the southern fixed stars not visible in our hemisphere, calculated from the observations made by Dr. Halley at St. Helena, adapted to the year 1726.

This work he prepared in a great measure for the press, with much care and accuracy: but through a natural weakness of constitution, and the declines of age, he died of a strangury before he had finished it, December the 19th, 1719, at 73 years of age; leaving the care of finishing and publishing his work to his friend Mr. Hodgson.—A less perfect edition of the *Historia Cælestis* had before been published, without his consent, viz, in 1712, in one volume folio, containing his observations to the year 1705.

Thus then, as Dr. Keil observed, our author, with indefatigable pains, for more than 40 years watched the motions of the stars, and has given us innumerable observations of the sun, moon, and planets, which he made with very large instruments, accurately divided, and fitted with telescopic sights; whence we may rely much more on the observations he has made, than on former astronomers, who made their observations with the naked eye, and without the like assistance of telescopes.

**FLANK**, in Fortification, is that part of the bastion which reaches from the curtain to the face; and it defends the curtain, with the opposite face and flank.

*Oblique or Second FLANK*, or *FLANK of the Curtain*, is that part of the curtain from whence the face of the opposite bastion can be seen, being contained between the lines rasant and fichant, or the greater and less lines of defence; or the part of the curtain between the Flank and the point where the fichant line of defence terminates.

*Covered, Low, or Retired FLANK*, is the platform of the casemate, which lies hid in the bastion; and is otherwise called the Orillon.

*Fichant FLANK*, is that from whence a cannon playing,



playing, fires directly on the face of the opposite bastion.

*Rasant* or *Razant* FLANK, is the point from whence the line of defence begins, from the conjunction of which with the curtain, the shot only raseth the face of the next bastion, which happens when the face cannot be discovered but from the Flank alone.

*Simple* FLANKS, are lines going from the angle of the shoulder to the curtain; the chief office of which is for the defence of the moat and place.

FLANK, is also a term of war, used by way of analogy for the side of a battalion, army, &c, in contradistinction to the front and rear. So, to attack the enemy in Flank, is to discover and fire upon them on one side.

FLANKED, is used of something that has Flanks, or may be approached on the Flank. As

FLANKED *Angle*, which is that formed by the two faces of the bastion, and forming its point or angle. Also,

FLANKED *Line of Defence*, FLANKED *Tenaille*, &c.

FLANKING, in general, is the discovering and firing upon the side of a place, body, battalion, &c. To Flank a place, or other work, is to dispose it in such a manner as that every part of it may be played upon both in front and rear.

FLANKING means also defending. Any fortification that has no defence, but just right forwards, is faulty; and to render it complete, one part ought to be made to Flank the other. Hence the curtain is always the strongest part of any place, because it is flanked at each end.

Battalions also are said to be Flanked by the wings of the cavalry. And a house is sometimes said to be Flanked with two pavilions, or two galleries; meaning it has a gallery, &c, on each side.—There are also Flanking Angle, Flanking Line of Defence, &c.

FLAT *Bastion*, is that which is built on a right line, as on the middle of the curtain, &c.

FLEXIBLE, is the property or quality of a body that may be bent.

FLEXION, the same as Flexure.

FLEXURE, or FLEXION, is the bending or curving of a line or figure.

When a line first bends one way, and then gradually changes to a bend the contrary way, the point where the two parts join, or where the bending changes to the other side, is called the point of inflexion, or of contrary Flexure.

FLIE, or FLY, that part of the mariner's compass, on which the 32 points of the wind are drawn, and over which the needle is placed, and fastened underneath.

FLOAT-Boards, the boards fixed to the outer rim of undershot water wheels, serving to receive the impulse of the stream, by which the wheel is carried round.—There may be too many of these boards on a wheel. It is thought to be the best rule, to have their distance asunder such, that each of them may come out of the water as soon as possible, after it has received and acted with its full impulse; or, which comes to the same thing, when the succeeding one is in a direction perpendicular to the surface of the water.

FLOATING *Bridge*, is a bridge of boats, casks, &c, covered with planks, firmly bound together for the passage of men, horses, or carriages, &c.

FLOOD, a Deluge or inundation of water.

FLOOD, is also used in speaking of the tide, when it is rising or flowing up; in contradistinction to the Ebb, which is when it is decreasing or running out.

FLOORING, in Carpentry, is commonly understood of the boarding of the Floors. The measurement of Flooring is estimated in squares, of 100 square feet each, or of 10 feet on each side every way; for 10 times 10 are 100. Hence the length of the floor being multiplied by its breadth, in feet, and two figures cut off on the right-hand, gives the squares, and feet, or decimals cut off. Thus, a Floor being 22 feet long, and 16 wide;

then 22 length  
16 breadth

132  
22

3,52 the content is therefore 3 squares, and 52 feet, or decimals.

FLUENT, or *Flowing Quantity*, in the Doctrine of Fluxions, is the variable quantity which is considered as increasing or decreasing; or the Fluent of a given fluxion, is that quantity whose fluxion being taken, according to the rules of that doctrine, shall be the same with the given fluxion. See FLUXIONS.

*Contemporary* FLUENTS, are such as flow together or for the same time. And the same is to be understood of Contemporary Fluxions.—When Contemporary Fluents are always equal, or in any constant ratio; then also are their fluxions respectively either equal, or in that same constant ratio. That is, if  $x = y$ , then is  $\dot{x} = \dot{y}$ ; or if  $x : y :: n : 1$ , then is  $\dot{x} : \dot{y} :: n : 1$ ; or if  $x = ny$ , then is  $\dot{x} = n\dot{y}$ .

It is easy to find the fluxions to all the given forms of Fluents; but, on the contrary, it is difficult to find the Fluents of many given fluxions; and indeed there are numberless cases in which this cannot at all be done, excepting by the quadrature and rectification of curve lines, or by logarithms, or infinite series.

This doctrine, as it was first invented by Sir Isaac Newton, so it was carried by him to a considerable degree of perfection, at least as to the most frequent, and most useful forms of fluents; as may be seen in his Fluxions, and in his Quadrature of Curves. Maclaurin, in his Treatise of Fluxions, has made several inquiries into Fluents, reducible to the rectification of the ellipse and hyperbola: and D'Alembert has pursued the same subject, and carried it farther, in the Memoires de l'Acad. de Berlin, tom. 2, p. 200. To the celebrated Mr. Euler this doctrine is greatly indebted, in many parts of his various writings, as well as in the Institutio Calculi Integralis, in 3 vols 4to, Petr. 1768. The ingenious Mr. Cotes contributed very much to this doctrine, in his Harmonia Mensurarum, concerning the measures of ratios and angles, in a large collection of different forms of fluxions, with their corresponding Fluents. And this subject was farther prosecuted in the same way by Walmesley, in his Analyse des Mesures des Rapports et des Angles, a large vol. in 4to, 1749. Besides many other Authors who, by their ingenious labours, have greatly contributed to facilitate and extend the doctrine of Fluents; as Emerson, Simpson, Landen, Waring, &c, in this country; with l'Hôpital,



pital, and many other learned foreigners. Lastly, in 1785 was published at Vienna, by M. Paccassi, a German nobleman, *Udhandlung uber eine neue Methode zu Integriren*, being a method of integrating, or finding the Fluents of given fluxions, by the rules of the direct method, or by taking again the fluxion of the given fluxion, or the 2d fluxion of the fluent sought; and then making every flowing quantity its fluxion, and 2d fluxion, in geometrical progression; a method however, which, it seems, only holds true in the easiest cases or forms, whose fluents are easily had by the most common methods. See this method farther explained in the rules following.

As it is only in certain particular forms and cases that the Fluents of given fluxions can be found; there being no method of performing this universally a priori, by a direct investigation; like finding the fluxion of a given fluent quantity; we can do little more than lay down a few rules for such forms of fluxions as are known, from the direct method, to belong to such and such kinds of Fluents or flowing quantities: and these rules, it is evident, must chiefly consist in performing such operations as are the reverse of those by which the fluxions are found to given flowing quantities. The principal cases of which are as follow:

I. *To find the Fluent of a simple fluxion; or that in which there is no variable quantity, and only one fluxional quantity.* This is done by barely substituting the variable or flowing quantity instead of its fluxion, and is the result or reverse of the notation only. Thus,

The Fluent of  $ax$  is  $ax$ .

The Fluent of  $ay + 2y$  is  $ay + 2y$ .

The Fluent of  $\sqrt{a^2 + x^2}$  is  $\sqrt{a^2 + x^2}$ .

II. *When any power of a flowing quantity is multiplied by the fluxion of the root.* Then, having substituted, as before, the flowing quantity for its fluxion, divide the result by the new index of the power. Or, which is the same thing, take out, or divide by, the fluxion of the root; add 1 to the index of the power; and divide by the index so increased.

So if the fluxion proposed be  $3x^5\dot{x}$ ;

Strike out  $\dot{x}$ , then it is  $3x^5$ ;

add 1 to the index, and it is  $3x^6$ ;

divide by the index 6, and it is  $\frac{3}{6}x^6$  or  $\frac{1}{2}x^6$ ;

which is the Fluent of the proposed fluxion  $3x^5\dot{x}$ .

In like manner, the Fluent

of  $4ax\dot{x}$  is  $2ax^2$ ;

of  $3x^{\frac{1}{2}}\dot{x}$  is  $2x^{\frac{3}{2}}$ ;

of  $ax^n\dot{x}$  is  $\frac{a}{n+1}x^{n+1}$ ;

of  $\frac{\dot{z}}{z^2}$  or  $z^{-2}\dot{z}$  is  $-z^{-1}$  or  $-\frac{1}{z}$ .

of  $(a^3 + z^3)^4 z^2 \dot{z}$  is  $\frac{1}{15}(a^3 + z^3)^5$ .

III. *When the root under a vinculum is a compound quantity; and the index of the part or factor without the vinculum increased by 1, is some multiple of that under the vinculum:* Put a single variable letter for the compound root; and substitute its powers and fluxion instead of those, of the same value, in the given quantity; so will it be reduced to a simpler form, to which the preceding rule can then be applied.

3

So, if the given fluxion be  $\dot{F} = (a^2 + x^2)^{\frac{2}{3}} x^3 \dot{x}$ ; where 3, the index of the quantity without the vinculum, increased by 1, makes 4, which is double of 2, the exponent of  $x^2$  within the same; therefore putting  $z = a^2 + x^2$ , thence  $x^2 = z - a^2$ , and its fluxion is  $2x\dot{x} = \dot{z}$ ; hence then  $x^3\dot{x} = \frac{1}{2}x^2\dot{z} = \frac{1}{2}\dot{z}(z - a^2)$ , and

the given quantity  $\dot{F}$  or  $(a^2 + x^2)^{\frac{2}{3}} x^3 \dot{x}$  is  $= \frac{1}{2}z^{\frac{2}{3}}\dot{z}(z - a^2)$  or  $= \frac{1}{2}z^{\frac{5}{3}}\dot{z} - \frac{1}{2}a^2 z^{\frac{2}{3}}\dot{z}$ ; and the Fluent of each term gives

$F = \frac{3}{16}z^{\frac{5}{3}} - \frac{3}{16}a^2 z^{\frac{2}{3}} = 3z^{\frac{5}{3}}(\frac{1}{16}z - \frac{1}{16}a)$ ; or, by substituting the value of  $z$  instead of it, the same Fluent is  $3(a^2 + x^2)^{\frac{5}{3}} \times (\frac{1}{16}x^2 - \frac{3}{16}a^2)$ , or  $\frac{3}{16} \cdot \overline{a^2 + x^2}^{\frac{5}{3}} \cdot \overline{x^2 - 3a^2}$ .

IV. *When there are several terms involving two or more variable quantities, having the fluxion of each multiplied by the other quantity or quantities:* Take the Fluent of each term, as if there was only one variable quantity in it, namely that whose fluxion is contained in it, supposing all the others to be constant in that term; then if the Fluents of all the terms so found, be the very same quantity, that quantity will be the Fluent of the whole.

Thus, if the given fluxion be  $\dot{x}y + x\dot{y}$ . Then, the Fluent of  $\dot{x}y$  is  $xy$ , supposing  $y$  constant; and the Fluent of  $x\dot{y}$  is also  $xy$ , when  $x$  is constant; therefore the common resulting quantity  $xy$  is the required Fluent of the given fluxion  $\dot{x}y + x\dot{y}$ .

And, in like manner, the Fluent of  $\dot{x}yz + x\dot{y}z + xy\dot{z}$  is  $xyz$ .

V. *When the given fluxional expression is in this form  $\frac{\dot{x}y - x\dot{y}}{y^2}$ , viz, a fraction including two quantities, being the fluxion of the former drawn into the latter, minus the fluxion of the latter drawn into the former, and divided by the square of the latter:* then the Fluent is the fraction  $\frac{x}{y}$ , or of the former quantity divided by the latter. That is,

The Fluent of  $\frac{\dot{x}y - x\dot{y}}{y^2}$  is  $\frac{x}{y}$ ;

and the Fluent of  $\frac{2x\dot{x}y^2 - 2x^2y\dot{y}}{y^4}$  is  $\frac{x^2}{y^2}$ .

Though the examples of this case may be performed by the foregoing one. Thus the given fluxion

$\frac{\dot{x}y - x\dot{y}}{y^2}$  reduces to  $\frac{\dot{x}}{y} - \frac{x\dot{y}}{y^2}$  or  $\frac{\dot{x}}{y} - xy\dot{y}^{-2}$ ;

of which the Fluent of  $\frac{\dot{x}}{y}$  is  $\frac{x}{y}$  when  $y$  is constant;

and the Fluent of  $xy\dot{y}^{-2}$  is  $+\frac{x}{y}$  or  $\frac{x}{y}$  when  $x$  is con-

stant; and therefore, by that case,  $\frac{x}{y}$  is the Fluent of

the whole  $\frac{\dot{x}y - x\dot{y}}{y^2}$ .

VI. *When the fluxion of a quantity is divided by the quantity itself:* Then the Fluent is equal to the hyperbolic logarithm of that quantity; or, which is the same thing, the Fluent is equal to 2.30258509 &c, multiplied by the common log. of the same quantity.

So,



So, the Fluent  
of  $\frac{\dot{x}}{x}$  or  $x^{-1}\dot{x}$  is the hyp. log. of  $x$ ;  
of  $\frac{2\dot{x}}{x}$  is  $2 \times$  hyp. log. of  $x$ , or  $=$  h. l. of  $x^2$ ;  
of  $\frac{a\dot{x}}{x}$  is  $a \times$  h. l. of  $x$ , or h. l. of  $x^a$ ;  
of  $\frac{\dot{x}}{a+x}$  is the h. l. of  $a+x$ ;  
of  $\frac{3x^2\dot{x}}{a+x^3}$  is the h. l. of  $a+x^3$ .

VII. Many Fluents may be found by the direct method of fluxions, thus: Take the fluxion again of the given fluxional expression, or the 2d fluxion of the Fluent sought; into which substitute  $\frac{\dot{x}^2}{x}$  for  $\ddot{x}$ , and  $\frac{y^2}{y}$  for  $\ddot{y}$ , &c, that is, make  $x, \dot{x}, \ddot{x}$ , as also  $y, \dot{y}, \ddot{y}$ , &c, in continual proportion, or  $x:\dot{x}::\dot{x}:\ddot{x}$ , and  $y:\dot{y}::\dot{y}:\ddot{y}$ , &c; then divide the square of the given fluxional expression by the 2d fluxion, just found, and the quotient will be the Fluent sought in many cases.

Or the same rule may be otherwise delivered thus: In the given fluxion  $\dot{F}$ , write  $x$  for  $\dot{x}$ ,  $y$  for  $\dot{y}$ , &c, and call the result  $G$ , taking also the fluxion of this quantity,  $\dot{G}$ ; then make  $\dot{G}:\dot{F}::G:F$ , so shall the 4th proportional  $F$  be the Fluent as before. And this is the rule of M. Paccassi.

It may be proved if this be the true Fluent, by taking the fluxion of it again, which, if it agree with the proposed fluxion, will shew that the Fluent is right; otherwise, it is wrong.

Thus, if it be proposed to find the Fluent of  $nx^{n-1}\dot{x}$ . Here  $\dot{F} = nx^{n-1}\dot{x}$ ; write first  $x$  for  $\dot{x}$ , and it is  $nx^{n-1}x$  or  $nx^n = G$ ; the fluxion of this is  $\dot{G} = n^2x^{n-1}\dot{x}$ ; therefore  $\dot{G}:\dot{F}::G:F$  becomes  $n^2x^{n-1}\dot{x}:nx^{n-1}\dot{x}::nx^n:x^n = F$ , the Fluent sought.

For a 2d ex. suppose it be proposed to find the Fluent of  $\dot{x}y + x\dot{y}$ . Here  $\dot{F} = \dot{x}y + x\dot{y}$ ; then, writing  $x$  for  $\dot{x}$ , and  $y$  for  $\dot{y}$ , it is  $xy + xy$  or  $2xy = G$ ; the fluxion of which is  $2\dot{x}y + 2x\dot{y} = \dot{G}$ ; then  $\dot{G}:\dot{F}::G:F$  becomes  $2\dot{x}y + 2x\dot{y}:\dot{x}y + x\dot{y}::2xy:xy = F$ , the Fluent sought.

VIII. To find Fluents by means of a table of forms of Fluxions and Fluents.

In the following table are contained the most usual forms of fluxions that occur in the practical solution of problems, with their corresponding Fluents set opposite to them; by means of which, viz, comparing any proposed fluxion with the corresponding form here, the Fluent of it will be found.

Where it is to be noted, that the logarithms in the said forms, are the hyperbolic ones, which are found by multiplying the common logs. by 2.3025850929940 &c. Also the arcs whose sine, or tangent, &c, are mentioned, have the radius 1, and are those in the common tables of sines, tangents, &c.—And the numbers  $m, n$ , &c. are to be some quantities, as the forms fail when  $n = 0$ , or  $m = 0$ , &c.

Forms	Fluxions.	Fluents.
I	$x^{n-1}\dot{x}$	$\frac{1}{n}x^n$
II	$(a \pm x^n)^{m-1}x^{n-1}\dot{x}$	$\pm \frac{1}{mn}(a \pm x^n)^m$
III	$\frac{x^{mn-1}\dot{x}}{(a \pm x^n)^{m+1}}$	$\frac{1}{mna} \times \frac{x^{mn}}{(a \pm x^n)^m}$
IV	$\frac{(a \pm x^n)^{m-1}\dot{x}}{x^{mn+1}}$	$\frac{-1}{mna} \times \frac{(a \pm x^n)^m}{x^{mn+1}}$
V	$(m\dot{x}y + nxy)\dot{x}^{m-1}y^{n-1}$ , or $\left(\frac{m\dot{x}}{x} + \frac{n\dot{y}}{y}\right)x^m y^n$	$x^m y^n$
VI	$m\dot{x}x^{m-1}\dot{y}y^{n-1}\dot{z}z^{r-1} + n\dot{x}x^{m-1}y^{n-1}\dot{z}z^{r-1} + r\dot{x}x^{m-1}y^{n-1}\dot{z}z^{r-1}$ , or $(m\dot{x}y\dot{z} + n\dot{x}y\dot{z} + r\dot{x}y\dot{z})x^{m-1}y^{n-1}z^{r-1}$ , or $\left(\frac{m\dot{x}}{x} + \frac{n\dot{y}}{y} + \frac{r\dot{z}}{z}\right)x^m y^n z^r$	$x^m y^n z^r$
	$\frac{\dot{x}}{x}$ or $x^{-1}\dot{x}$	log. of $x$



Forms.	Fluxions.	Fluents.
VIII	$\frac{x^{n-1} \dot{x}}{a \pm x^n}$	$\pm \frac{1}{n} \log. \text{ of } a \pm x^n$
IX	$\frac{x^{-1} \dot{x}}{a \pm x^n}$	$\frac{1}{na} \log. \text{ of } \frac{x^n}{a \pm x^n}$
X	$\frac{\frac{1}{2}n-1}{x} \frac{\dot{x}}{a - x^n}$	$\frac{1}{n\sqrt{a}} \log. \text{ of } \frac{\sqrt{a} + \sqrt{x^n}}{\sqrt{a} - \sqrt{x^n}}$
XI	$\frac{\frac{1}{2}n-1}{x} \frac{\dot{x}}{a + x^n}$	$\frac{2}{n\sqrt{a}} \times \text{arc to tang. } \sqrt{\frac{x^n}{a}}, \text{ or } \frac{1}{n\sqrt{a}} \times \text{arc to cofine } \frac{a - x^n}{a + x^n}$
XII	$\frac{\frac{1}{2}n-1}{x} \frac{\dot{x}}{\sqrt{\pm a + x^n}}$	$\frac{2}{n} \log. \text{ of } \sqrt{x^n} + \sqrt{\pm a + x^n}$
XIII	$\frac{\frac{1}{2}n-1}{x} \frac{\dot{x}}{\sqrt{a - x^n}}$	$\frac{2}{n} \times \text{arc to fin. } \sqrt{\frac{x^n}{a}}, \text{ or } \frac{1}{n} \times \text{arc to verf. } \frac{2x^n}{a}$
XIV	$\frac{x^{-1} \dot{x}}{\sqrt{a \pm x^n}}$	$\frac{1}{n\sqrt{a}} \log. \text{ of } \frac{\pm \sqrt{a \mp x^n} \pm \sqrt{a}}{\sqrt{a \pm x^n} + \sqrt{a}}$
XV	$\frac{x^{-1} \dot{x}}{\sqrt{-a + x^n}}$	$\frac{2}{n\sqrt{a}} \times \text{arc to secant } \sqrt{\frac{x^n}{a}}, \text{ or } \frac{1}{n\sqrt{a}} \times \text{arc to cofin. } \frac{2a - x^n}{x^n}$
XVI	$\dot{x} \sqrt{dx - x^2}$	$\frac{1}{2} \text{ circ. seg. to diam. } d \text{ and verf. } x$
XVII	$c^{nx} \dot{x}$	$\frac{c^{nx}}{n \log. c}$
XVIII	$\dot{x} y^x \log. y + x y^{x-1} \dot{y}$	$y$
XIX	$(a + x^n)^m x^{rn-1} \dot{x}$ Putting $s = m + r$	$\frac{(a + x^n)^{m+1} x^{rn-n}}{ns} \times \left( \frac{1}{1} - \frac{r-1.a}{s-1.x^n} + \frac{r-1.r-2.a^2}{s-1.s-2.x^{2n}} - \frac{r-1.r-2.r-3.a^3}{s-1.s-2.s-3.x^{3n}} \&c \right)$ or $\frac{(a + x^n)^{m+1} x^{rn}}{rna} \times \left( \frac{1}{1} - \frac{s+1.x^n}{r+1.a} + \frac{s+1.s+2.x^{2n}}{r+1.r+2.a^2} - \frac{s+1.s+2.s+3.x^{3n}}{r+1.r+2.r+3.a^3} \&c \right)$

*The Use of the foregoing Table of Forms of Fluxions and Fluents.*—In the use of this table, it is to be observed, that the first column serves only to shew the number of the form, as a mark of reference; in the 2d column are the several forms of fluxions, which are of different kinds or classes; and in the 3d or last column are the corresponding Fluents.

The method of using the table is this. Having any fluxion given, whose Fluent it is proposed to find: First, compare the given fluxion with the several forms

of fluxions in the 2d column of the table, till one of the forms be found that agrees with it; which is done by comparing the terms of the given fluxion with the like parts of the tabular fluxion, viz, the radical quantity of the one, with that of the other; and the exponents of the variable quantities of each, both within and without the vinculum; all which, being found to agree or correspond, will give the particular values of the general quantities in the tabular form. Then substitute these particular values, for the same quantities



in the general or tabular form of the Fluent, and the result will be the particular Fluent sought; after it is multiplied by any coefficient the proposed fluxion may have.

*For Ex.* To find the Fluent of the given fluxional expression  $3x^{\frac{5}{3}}\dot{x}$ . This agrees with the first form; and by comparing the fluxions, it appears that  $x = x$ , and  $n - 1 = \frac{5}{3}$ , or  $n = \frac{8}{3}$ ; which being substituted in the tabular Fluent, or  $\frac{1}{n}x^n$ , gives, after multiplying by 3 the coefficient,  $3 \times \frac{3}{8}x^{\frac{8}{3}}$  or  $\frac{9}{8}x^{\frac{8}{3}}$  for the Fluent sought.

Again, To find the Fluent of  $5x^2\dot{x}\sqrt{c^3 - x^3}$ , or  $5x^2\dot{x} \cdot (c^3 - x^3)^{\frac{1}{2}}$ . This belongs to the 2d form; for  $a = c^3 - x^3 = -x^3$ ,  $n = 3$  under the vinculum,  $m - 1 = \frac{1}{2}$ , or  $m = \frac{3}{2}$ , and the exponent  $n - 1$  of  $x^{n-1}$  without the vinculum, by using 3 for  $n$ , is  $n - 1 = 2$ , which agrees with  $x^2$  in the fluxion given; and therefore all the parts of the form are found to answer. Then, substituting these values into the general Fluent,  $-\frac{1}{mn}(a - x^n)^m$ , it becomes  $-\frac{5 \times 2}{3 \times 3}(c^3 - x^3)^{\frac{3}{2}} = -\frac{10}{9}(c^3 - x^3)^{\frac{3}{2}}$ .

Thirdly, To find the Fluent of  $\frac{x^2\dot{x}}{1 + x^3}$ . This agrees with the 8th form; where  $\pm x^n = +x^3$  in the denominator, or  $n = 3$ ; and the numerator  $x^{n-1}$  then becomes  $x^2$ , which agrees with the numerator in the given fluxion; also  $a = 1$ . Hence then, by substituting in the general form of the Fluent  $\frac{1}{n}$  logarithm of  $a + x^n$ , it becomes  $\frac{1}{3}$  logarithm of  $1 + x^3$ .

**IX. To find Fluents by means of Infinite Series.**—When a finite form cannot be found to agree with a proposed fluxion, it is then usual to throw it into an infinite series, either by division, or extraction of roots, or by the binomial theorem, &c; after which, the Fluents of all the terms are taken separately.

*For Ex.* To find the Fluent of  $\frac{1-x}{1+x-x^2}\dot{x}$ . Here, by dividing the numerator by the denominator, this becomes  $\dot{x} - 2x\dot{x} + 3x^2\dot{x} - 5x^3\dot{x} + 8x^4\dot{x}$  &c; and, the Fluents of all the terms being taken, give  $x - x^2 + x^3 - \frac{5}{2}x^4 + \frac{8}{3}x^5$  &c, for the Fluent sought.

**To Correct a FLUENT.**—The Fluent of a given fluxion, found as above, sometimes wants a correction, to make it contemporary with that required by the problem under consideration, &c: for the Fluent of any given fluxion, as  $\dot{x}$ , may be either  $x$  (which is found by the rule) or it may be  $x \pm c$ , that is  $x$  plus or minus some constant quantity  $c$ ; because both  $x$  and  $x \pm c$  have the same fluxion  $\dot{x}$ : and the finding of this constant quantity, is called correcting the Fluent. Now this correction is to be determined from the nature of the problem in hand, by which we come to know the relation which the Fluent quantities have to each other at some certain point or time. Reduce therefore the general Fluential equation, found by the rules above, to that point or time; then if the equation be true at that point, it is correct; but if not, it wants a correction, and the quantity of that correction is the dif-

ference between the two general sides of the equation when reduced to that particular state. Hence the general rule for the correction is this:

Connect the constant, but indeterminate, quantity  $c$  with one side of the Fluential equation, as determined by the foregoing rules; then, in this equation, substitute for the variable quantities such values as they are known to have at any particular state, place, or time; and then from that particular state of the equation find the value of  $c$ , the constant quantity of the correction.

*Ex.* To find the Correct Fluent of  $\dot{z} = ax^3\dot{x}$ . First the general Fluent of this is  $z = ax^4$ , or  $z = ax^4 + c$ , taking in the correction  $c$ .

Now if it be known that  $z$  and  $x$  begin together, or that  $z = 0$ , when  $x = 0$ ; then writing 0 both for  $x$  and  $z$ , the general equation becomes  $0 = 0 + c$ , or  $c = 0$ ; so that, the value of  $c$  being 0, the Correct Fluents are  $z = ax^4$ .

But if  $z$  be  $= 0$ , when  $x$  is  $= b$ , any known quantity; then substituting 0 for  $z$ , and  $b$  for  $x$ , in the general equation, it becomes  $0 = ab^4 + c$ , from which is found  $c = -ab^4$ ; and this being written for it in the general equation, this becomes  $z = ax^4 - ab^4$ , for the correct, or contemporary Fluents.

Or lastly, if it be known that  $z$  is  $=$  some quantity  $d$ , when  $x$  is equal some other quantity, as  $b$ ; then substituting  $d$  for  $z$ , and  $b$  for  $x$ , in the general Fluential equation  $z = ax^4 + c$ , it becomes  $d = ab^4 + c$ ; and hence is deduced the value of the correction, viz,  $c = d - ab^4$ ; consequently, writing this value for  $c$  in the general equation, it becomes  $z = ax^4 - ab^4 + d$ , for the Correct equation of the Fluents in this case.

And hence arises another easy and general way of correcting the Fluents, which is this: In the general equation of the Fluents, write the particular values of the quantities which they are known to have at any certain time; then subtract the sides of the resulting particular equation, from the corresponding sides of the general one, and the remainders will give the Correct equation of the Fluents sought. So, as above,

the general equation being  $z = ax^4$ ;  
write  $d$  for  $z$ , and  $b$  for  $x$ , then  $d = ab^4$ ;  
hence by subtraction  $z - d = ax^4 - ab^4$ ,  
or  $z = ax^4 - ab^4 + d$ ,

the Correct Fluents as before.

**FLUID**, or **FLUID Body**, according to Newton, is that whose parts yield to the smallest force impressed, and by yielding are easily moved among each other; in which sense it stands opposed to a solid. This is the definition of a perfect fluid: if the Fluid require some sensible force to move its parts, it is imperfect in proportion to that force; such as perhaps all the fluids we know of in nature.

That Fluids have vacuities in their substance is evident, because certain bodies may be dissolved in them without increasing their bulk. Thus, water will dissolve a certain quantity of salt; after which it will receive a little sugar, and after that a little alum; and all this without increasing its first dimensions. Which shews that the particles of these solids are so far separated as to become smaller than those of the Fluid, and to be received and contained in the interstices between them.

Fluids are either elastic, such as air; or non-elastic,



as water, mercury, &c. These latter occupy the same space, or are of the same bulk, under all pressures or forces; but the former dilate and expand themselves continually by taking off the external pressure from them; for which reason it is that the density and elasticity of such fluids, are proportional to the force or weight that compresses them. The doctrine and laws of Fluids are of the greatest extent in philosophy: the properties of elastic Fluids constituting the doctrine of Pneumatics; those of the non-elastic ones, that of Hydrostatics; and their motions, Hydraulics. For which see these respective articles. Also,

For the laws of the pressure and gravitation in Fluids specifically heavier or lighter than the bodies immersed, see *SPECIFIC Gravity*.

For the laws of the resistance of Fluids, or the retardation of solid bodies moving in Fluids, see *RESISTANCE*. And

For the ascent of Fluids in capillary tubes, or between glass planes, &c, see *ASCENT*, and *CAPILLARY Tubes*.

*FLUTES*, or *FLUTINGS*, are certain channels or cavities cut along the shaft of a column, or pilaster.

*FLUIDITY*, that state or affection of bodies, which denominates or renders them Fluid; or that property by which they yield to the smallest force impressed: in contradistinction to Solidity or Firmness.

Fluidity is to be carefully distinguished from Liquidity or Humidity, which latter implies wetting or adhering. Thus, air, ether, mercury, and other melted metals, and even smoke and flame itself, are Fluid bodies, but not Liquid ones; whilst water, beer, milk, urine, &c, are both Fluids and Liquids at the same time.

The nature and causes of Fluidity have been variously assigned. The Gassendists, and ancient corpuscularians, require only three conditions as necessary to it; viz, a smallness and smoothness of the particles of the body, vacuities interspersed between them, and a spherical figure. The Cartesians, and after them Dr. Hook, Mr. Boyle, &c, beside these circumstances, require also a certain internal or intestine motion of the particles as chiefly contributing to Fluidity. Thus, Mr. Boyle, in his History of Fluidity, argues from various experiments: for example, a little dry powder of alabaster, or plaister of Paris, finely sifted, being put into a vessel over the fire, soon begins to boil like water; exhibiting all the motions and phenomena of a boiling liquor: it will tumble variously in great waves like that; will bear stirring with a stick or ladle like that, without resisting; and if strongly stirred near the side of the vessel, its waves will apparently dash against it: yet it is all the while a dry parched powder.

The like is observed in sand; a dish of which being set on a drum-head, briskly beaten by the sticks, or on the upper stone of a mill, it in all respects emulates the properties of a Fluid body. A heavy body, ex. gr. will immediately sink in it to the bottom, and a light one emerge to the top: each grain of sand has a constant vibratory and dancing motion; and if a hole be made in the side of the dish, the sand will spin out like water.

The Cartesians bring divers considerations to prove that the parts of Fluids are in continual motion: as 1st,

The change of solids into Fluids, ex. gr. ice into water, and vice versa; the chief difference between the body in those two states consisting in this, that the parts, being fixed and at rest in the one, resist the touch; whereas in the other, being already in motion, they give way to the slightest impulse. 2dly, The effects of Fluids, which commonly proceed from motion: such are the insinuation of Fluids among the pores of bodies; the softening and dissolving hard bodies; the actions of corrosive menstrua; &c: Add, that no solid can be brought to a state of Fluidity, without the intervention of some moving or moveable body, as fire, air, or water. Air, the same gentlemen hold to be the first spring of these causes of Fluidity, it being this that gives motion to fire and water, though itself receives its motion and action from the ether, or subtile medium.

But Boerhaave pleads strenuously that fire is the first mover, and the cause of all Fluidity in other bodies, as air, water, &c: without this, he shews that the atmosphere itself would fix into one solid mass. And in like manner, Dr. Black, of Edinburgh, mentions Fluidity as an effect of heat. The different degrees of heat which are required to bring different bodies into a state of Fluidity, he supposes may depend on some particulars in the mixture and composition of the bodies themselves: which is rendered farther probable from considering that the natural state of bodies in this respect is changed by certain mixtures; thus, when two metals are compounded, the mixture is commonly more fusible than either of them separately.

Newton's idea of the cause of Fluidity is different: he makes it to be the great principle of attraction. The various intestine motion and agitation among the particles of Fluid bodies, he thinks is naturally accounted for, by supposing it a primary law of nature, that as all the particles of matter attract each other when within a certain distance; so at all greater distances, they avoid and fly from one another. For then, though their common gravity, together with the pressure of other bodies upon them, may keep them together in a mass, yet their continual endeavour to avoid one another singly, and the adventitious impulses of heat and light, or other external causes, may make the particles of Fluids continually move round about one another, and so produce this quality.

As therefore the cause of cohesion of the parts of solid bodies appears to be their mutual attraction; so, on this principle, the chief cause of Fluidity seems to be a contrary motion, impressed on the particles of Fluids; by which they avoid and fly from one another, as soon as they come at, and as long as they keep at, such a distance from each other.

It is observed also in all Fluids, that the direction of their pressure against the vessels which contain them, is in lines perpendicular to the sides of such vessels; which property, being the necessary result of the spherical figure of the particles of any Fluid, shews that the parts of all Fluids are so, or of a figure very nearly approaching to it.

*FLUX*, in Hydrography, a regular and periodical motion of the sea, happening twice in 24 hours and 48 minutes, nearly; in which time the water is raised, and driven violently against the shores. The Flux, or Flow,



is one of the motions of the tide: the other, by which the water sinks and retires, being called the Reflux, or Ebb. See TIDE.

Between the Flux and Reflux there is a kind of rest or cessation, of about half an hour; during which time the water is at its greatest height, called High-water.

The Flux of the sea follows chiefly the course of the moon; and is always highest and greatest at new and full moons, particularly near the time of the equinoxes. In some parts, as at Mount St. Michael, it rises 80 or 90 feet, though in the open sea it never rises above a foot or two; and in some places, as about the Morea, there is no flux at all. It runs up some rivers above 120 miles: though up the river Thames it goes only about 80, viz, near to Kingston in Surry. Above London-bridge, the water flows 4 hours, and ebbs 8; and below the bridge, it flows 5 hours, and ebbs 7.

FLUXION, in the Newtonian analysis, denotes the rate or proportion at which a flowing or varying quantity increases its magnitude or quantity; and it is proportional to the magnitude by which the flowing quantity would be uniformly increased, in a given time, by the generating quantity continuing of the invariable magnitude it has at the moment of time for which the Fluxion is taken: by which it stands contradistinguished from fluent, or the flowing quantity, which is gradually, and indefinitely increasing, after the manner of a space which a body in motion describes.

Mr. Simpson observes, that there is an advantage in thus considering Fluxions, not as mere velocities of increase at a certain point, but as the magnitudes which would be uniformly generated in a given finite time: the imagination is not here confined to a single point, and the higher orders of Fluxions are rendered much more easy and intelligible. And though Sir Isaac Newton defines Fluxions to be the velocities of motions, yet he has recourse to the moments or increments, generated in equal particles of time, to determine those velocities, which he afterwards directs to expound by finite magnitudes of other kinds.

As to the illustration of this definition, and the rules for finding the Fluxions of all sorts of fluent quantities, see the following article, or the Method of Fluxions.

*Method of FLUXIONS*, is the algorithm and analysis of Fluxions, and fluents or flowing quantities.

Most foreigners define this as the method of differences or differentials, being the analysis of indefinitely small quantities. But Newton, and other English authors, call these infinitely small quantities, moments; considering them as the momentary increments of variable quantities; as of a line considered as generated by the flux or motion of a point, or of a surface generated by the flux of a line. Accordingly, the variable quantities are called Fluents, or flowing quantities; and the method of finding either the Fluxion, or the fluent, the method of Fluxions.

M. Leibnitz considers the same infinitely small quantities as the differences, or differentials, of quantities; and the method of finding those differences, he calls the Differential Calculus.

Besides this difference in the name, there is another in the notation. Newton expresses the Fluxion of a quantity, as of  $x$ , by a dot placed over it, thus  $\dot{x}$ ;

while Leibnitz expresses his differential of the same  $x$ , by prefixing the initial letter  $d$ , as  $dx$ . But, setting aside these circumstances, the two methods are just alike.

The Method of Fluxions is one of the greatest, most subtle, and sublime discoveries of perhaps any age: it opens a new world to our view, and extends our knowledge, as it were, to infinity; carrying us beyond the bounds that seemed to have been prescribed to the human mind, at least infinitely beyond those to which the ancient geometry was confined.

The history of this important discovery, recent as it is, is a little dark, and embroiled. Two of the greatest men of the last age have both of them claimed the invention, Sir I. Newton, and M. Leibnitz; and nothing can be more glorious for the method itself, than the zeal with which the partisans of either side have asserted their title.

To exhibit a just view of this dispute; and of the pretensions of each party, we may here advert to the origin of the discovery, and mark where each claim commenced, and how it was supported.

The principles upon which the Method of Fluxions is founded, or which conducted to it, had been laying, and gradually developing, from the beginning of the last century, by Fermat, Napier, Barrow, Wallis, Slusius, &c, who had methods of drawing tangents, of maxima and minima, of quadratures, &c, in certain particular cases, as of rational quantities, upon nearly the same principles. And it was not wonderful that such a genius as Newton should soon after raise those faint beginnings into a regular and general system of science, which he did about the year 1665, or sooner.

The first time however that the method appeared in print, was in 1684, when M. Leibnitz gave the rules of it in the *Leipfic Acts* of that year; but without the demonstrations. The two brothers however, John and James Bernoulli, being greatly struck with this new method, applied themselves diligently to it, found out the demonstrations, and applied the calculus with great success.

But before this, M. Leibnitz had proposed his Differential Method, viz, in a letter, dated Jan. 21, 1677, in which he exactly pursues Dr. Barrow's method of tangents, which had been published in 1670: and Newton communicated his method of drawing tangents to Mr. Collins, in a letter dated Dec. 10, 1672; which letter, together with another dated June 13, 1676, were sent to Mr. Leibnitz by Mr. Oldenburgh, in 1676. So that there is a strong presumption that he might avail himself of the information contained in these letters, and other papers transmitted with them, and also in 1675, before the publication of his own letter, containing the first hint of his differential method. Indeed it sufficiently appears that Newton had invented his method before the year 1669, and that he actually made use of it in his *Compendium of Analysis and Quadrature of Curves* before that time. His attention seems to have been directed this way, even before the time of the plague which happened in London in 1665 and 1666, when he was about 28 years of age.

This is all that is heard of the method, till the year 1687, when Newton's admirable *Principia* came out, which



which is almost wholly built on the same calculus. The common opinion then was, that Newton and Leibnitz had each invented it about the same time: and what seemed to confirm it was, that neither of them made any mention of the other; and that, though they agreed in the substance of the thing, yet they differed in their ways of conceiving it, calling it by different names, and using different characters. However, foreigners having first learned the method through the medium of Leibnitz's publication, which spread the method through Europe, those geometers were insensibly accustomed to look upon him as the sole, or principal inventor, and became ever after strongly prejudiced in favour of his notation and mode of conceiving it.

The two great authors themselves, without any seeming concern, or dispute, as to the property of the invention, enjoyed the glorious prospect of the progress continually making under their auspices, till the year 1699, when the peace began to be disturbed.

M. Facio, in a treatise on the Line of Swiftest Descent, declared, that he was obliged to own Newton as the first inventor of the differential calculus, and the first by many years; and that he left the world to judge, whether Leibnitz, the second inventor, had taken any thing from him. This precise distinction between first and 2d inventor, with the suspicion it insinuated, raised a controversy between M. Leibnitz, supported by the editors of the *Leipfic Acts*, and the English mathematicians, who declared for Newton. Sir Isaac himself never appeared on the scene; his glory was become that of the nation; and his adherents, warm in the cause of their country, needed not his assistance to animate them.

Writings succeeded each other but slowly, on either side; probably on account of the distance of places; but the controversy grew still hotter and hotter: till at length M. Leibnitz, in the year 1711, complained to the Royal Society, that Dr. Keil had accused him of publishing the Method of Fluxions invented by Sir I. Newton, under other names and characters. He insisted that nobody knew better than Sir Isaac himself, that he had stolen nothing from him; and required that Dr. Keil should disavow the ill construction which might be put upon his words.

The Society, thus appealed to as a judge, appointed a committee to examine all the old letters, papers, and documents, that had passed among the several mathematicians, relating to the point; who, after a strict examination of all the evidence that could be procured, gave in their report as follows: "That Mr. Leibnitz was in London in 1673, and kept a correspondence with Mr. Collins by means of Mr. Oldenburgh, till Sept. 1676, when he returned from Paris to Hanover, by way of London and Amsterdam: that it did not appear that Mr. Leibnitz knew any thing of the differential calculus before his letter of the 21st of June, 1677, which was a year after a copy of a letter, written by Newton in the year 1672, had been sent to Paris to be communicated to him, and above 4 years after Mr. Collins began to communicate that letter to his correspondents; in which the Method of Fluxions was sufficiently explained, to let a man of his sagacity into the whole matter: and that Sir I. New-

ton had even invented his method before the year 1669, and consequently 15 years before M. Leibnitz had given any thing on the subject in the *Leipfic Acts*." From which they concluded that Dr. Keil had not at all injured M. Leibnitz in what he had said.

The Society printed this their determination, together with all the pieces and materials relating to it, under the title of *Commercium Epistolicum de Analyfi Promota*, 8vo, Lond. 1712. This book was carefully distributed through Europe, to vindicate the title of the English nation to the discovery; for Newton himself, as already hinted, never appeared in the affair: whether it was that he trusted his honour with his compatriots, who were zealous enough in the cause; or whether he felt himself even superior to the glory of it.

M. Leibnitz and his friends however could not shew the same indifference: he was accused of a theft; and the whole *Commercium Epistolicum* either expresses it in terms, or insinuates it. Soon after the publication therefore, a loose sheet was printed at Paris, in behalf of M. Leibnitz, then at Vienna. It is written with great zeal and spirit; and it boldly maintains that the Method of Fluxions had not preceded the Method of Differences; and even insinuates that it might have arisen from it. The detail of the proofs however, on each side, would be too long, and could not be understood without a large comment, which must enter into the deepest geometry.

M. Leibnitz had begun to work upon a *Commercium Epistolicum*, in opposition to that of the Royal Society; but he died before it was completed.

A second edition of the *Commercium Epistolicum* was printed at London in 1722; when Newton, in the preface, account, and annotations, which were added to that edition, particularly answered all the objections which Leibnitz and Bernoulli were able to make since the *Commercium* first appeared in 1712; and from the last edition of the *Commercium*, with the various original papers contained in it, it evidently appears that Newton had discovered his Method of Fluxions many years before the pretensions of Leibnitz. See also Raphson's History of Fluxions.

There are however, according to the opinion of some, strong presumptions in favour of Leibnitz; i. e. that he was no plagiarist: for that Newton was at least the first inventor, is past all dispute; his glory is secure; the reasonable part, even among the foreigners, allow it: and the question is only, whether Leibnitz took it from him, or fell upon the same thing with him; for, in his theory of abstract notions, which he dedicated to the Royal Academy in 1671, before he had seen any thing of Newton's, he already supposed infinitely small quantities, some greater than others; which is one of the great principles of his system.

Before prosecuting farther the history and improvements of this science, it will be proper to premise somewhat of the principles and practice of it, according to the ideas of the inventor.

#### *Principles of the Method of Fluxions.*

1. In the doctrine of Fluxions, magnitudes or quantities, of all kinds, are considered, not as made up of a number



number of small parts, but as generated by continued motion, by means of which they increase or decrease: as a line by the motion of a point; a surface by the motion of a line; and a solid by the motion of a surface: which is no new principle in geometry; having been used by Euclid and Archimedes. So likewise, time may be considered as represented by a line, increasing uniformly by the motion of a point. And quantities of all kinds whatever, which are capable of increase and decrease, may in like manner be represented by lines, surfaces, or solids, considered as generated by motion.

2. Any quantity, thus generated, and variable, is called a *Fluent*, or a *flowing quantity*. And the rate or proportion according to which any flowing quantity increases, at any position or instant, is the *Fluxion* of the said quantity, at that position or instant: and it is proportional to the magnitude by which the flowing quantity would be uniformly increased, in a given time, with the generating celerity uniformly continued during that time.

3. The small quantities that are actually generated or described, in any small given time, and by any continued motion, either uniform or variable, are called *Increments*.


4. Hence, if the motion of increase be uniform, by which increments are generated, the increments will in that case be proportional, or equal, to the measures of the Fluxions: but if the motion of increase be accelerated, the increments so generated, in a given finite time, will exceed the Fluxion; and if it be a decreasing motion, the increment so generated, will be less than the Fluxion. But if the time be indefinitely small, so that the motion be considered as uniform for that instant; then these nascent increments will always be proportional or equal to the Fluxions, and may be substituted for them, in any calculation.

5. To illustrate these definitions: Suppose a point  $m$  be conceived to move from the position  $A$ , and to



generate a line  $AP$ , with a motion any-how regulated; and suppose the celerity of the point  $m$ , at any position  $P$ , to be such, as would, if from thence it should become, or continue, uniform, be sufficient to describe, or pass uniformly over, the distance  $Pp$ , in the given time allowed for the Fluxion: then will the said line  $Pp$  represent the Fluxion of the said fluent or flowing line  $AP$ , at that position.


6. Again, suppose the right line  $mn$  to move, from the position  $AB$ , continually parallel to itself,



with any continued motion, so as to generate the fluent, or flowing rectangle  $ABQP$ , whilst the point  $m$  describes the line  $AP$ ; also let the distance  $Pp$  be taken,


as above, to express the Fluxion of the line or base  $AP$ ; and complete the rectangle  $PQqp$ . Then, like as  $Pp$  is the Fluxion of the line  $AP$ , so is the small parallelogram  $Pq$  the Fluxion of the flowing parallelogram,  $AQ$ ; both these Fluxions or increments being uniformly described in the same time.

7. In like manner, if the solid  $AERP$  be conceived as generated by the plane  $PQR$  moving,



from the position  $ABE$ , always parallel to itself, along the line  $AD$ ; and if  $Pp$  denote the Fluxion of the line  $AP$ . Then, like as the parallelogram  $Pq$ , or  $Pp \times PQ$ , expresses the Fluxion of the flowing rectangle  $AQ$ , so likewise shall the Fluxion of the variable solid or prism  $AR$  be expressed by the prism  $Pr$ , or  $Pp \times$  the plane  $PR$ . And in both these last two cases, it appears that the Fluxion of the generated rectangle, or prism, is equal to the product of the generant, whether line or plane, drawn into the Fluxion of the line along which it moves.

8. Hitherto the generant, or generating line or plane, has been considered as of a constant or invariable magnitude; in which case the fluent, or quantity generated, is a parallelogram, or a prism, the former being described by the motion of a line, and the latter by the motion of a plane. In like manner are other figures, whether plane or solid, conceived to be described, by the motion of a variable magnitude, whether it be a line or a plane. Thus, let a variable line  $PQ$  be carried with a parallel motion along  $AP$ , or whilst a point  $P$  is carried along, and describes, the line  $AP$ ,



suppose another point  $Q$  to be carried by a motion perpendicular to the former, and to describe the line  $PQ$ : let  $pq$  be another position of  $PQ$ , indefinitely near to the former; and draw  $Qr$  parallel to  $AP$ . Now in this case there are several fluents or flowing quantities, with their respective Fluxions: viz, the line or fluent  $AP$ , the Fluxion of which is  $Pp$ , or  $Qr$ ; the line or fluent  $PQ$ , the Fluxion of which is  $qr$ ; the curve, or oblique line  $AQ$ , described by the oblique motion of the point, the Fluxion of which is  $Qq$ ; and lastly the surface  $APQ$ , described by the variable line  $PQ$ , and the Fluxion of which is the rectangle  $PQrp$ , or  $PQ \times Pp$ . And in the same manner may any solid be conceived to be described, by the motion of a variable plane parallel to itself, substituting the variable plane for the variable line; in which case, the Fluxion of the solid, at any position, is represented by the variable



riable plane, at that position, drawn into the Fluxion of the line along which it is carried.

9. Hence then it follows generally, that the Fluxion of any figure, whether plane or solid, at any position, is equal to the section of it, at that position, drawn into the Fluxion of the axis, or line along which the variable section is supposed to be perpendicularly carried; i. e. the Fluxion of the figure  $AQP$ , is equal the plane  $PQ \times Pp$  when that figure is a solid, or to the ordinate  $PQ \times Pp$  when the figure is a surface.

10. It also follows, from the same premises, that, in any curve, or oblique line,  $AQ$ , whose absciss is  $AP$ , and ordinate is  $PQ$ , the Fluxions of these three form a small right-angled plane triangle  $Qqr$ ; for  $Qr = Pp$  is the Fluxion of the absciss  $AP$ ,  $qr$  the Fluxion of the ordinate  $PQ$ , and  $Qq$  the Fluxion of the curve or right line  $AQ$ . And consequently that, in any curve, the square of the Fluxion of the curve, is equal to the sum of the squares of the Fluxions of the absciss and ordinate, when these two lines are at right angles to each other.

11. From the premises it also appears, that contemporaneous fluents, or quantities that flow or increase together, which are always in a constant ratio to each other, have their Fluxions also in the same constant ratio at every position. For, let  $AP$  and  $BQ$  be two

A ————— P p B ————— Q q

contemporaneous fluents, described in the same time by the motion of the points  $P$  and  $Q$ , the contemporaneous positions being  $P, Q$ , and  $p, q$ ; and let  $AP$  be to  $BQ$ , or  $Ap$  to  $Bq$ , in the constant ratio of  $n$  to 1.

Then is ————  $AP = n \times BQ$ ,  
and ————  $Ap = n \times Bq$ ;  
therefore by subtraction, ————  $Pp = n \times Qq$ ;  
that is, the Fluxion  $Pp$  : Fluxion  $Qq :: n : 1$ ,  
the same as Fluent  $AP$  : Fluent  $BQ :: n : 1$ ;  
or the Fluxions and Fluents are in the same constant ratio.

But if the ratio of the fluents be variable, so will that of the Fluxions be also, though not in the same variable ratio with the former, at every position.

#### *The Notation, &c, in FLUXIONS.*

12. To apply the foregoing principles to the determination of the Fluxions of algebraic quantities, by means of which those of all other kinds are determined, it will be necessary first to premise the notation used in this science, with some observations. As, first, that the final letters of the alphabet  $z, y, x, v$ , &c, are used to denote variable or flowing quantities; and the initial letters  $a, b, c, d$ , &c, constant or invariable ones: Thus, the variable base  $AP$  of the flowing rectangular figure  $ABQP$ , at art. 6, may be represented by  $x$ ; and the invariable altitude  $PQ$ , by  $a$ : also the variable base or absciss  $AP$ , of the figures in art. 8, may be represented by  $x$ ; the variable ordinate  $PQ$ , by  $y$ ; and the variable curve or line  $AQ$ , by  $z$ .

Secondly, that the Fluxion of a quantity denoted by a single letter, is represented by the same letter with a point over it: Thus the Fluxion of  $x$  is expressed

by  $\dot{x}$ , that of  $y$  by  $\dot{y}$ , and that of  $z$  by  $\dot{z}$ . As to the Fluxions of constant or invariable quantities, as of  $a, b, c$ , &c, they are equal to 0 or nothing, because they do not flow, or change their magnitude.

Thirdly, that the increments of variable or flowing quantities, are also denoted by the same letters with a small ( $'$ ) over them: So the increments of  $x, y, z$ , are  $x', y', z'$ .

13. From these notations, and the foregoing principles, the quantities and their Fluxions, there considered, will be denoted as below.

In all the foregoing figures, put

the variable or flowing line ————  $AP = x$ ,  
in art. 6, the constant line ————  $PQ = a$ ,  
in art. 8, the variable ordinate ————  $PQ = y$ ,  
the variable curve or right line ————  $AQ = z$ ;

Then shall the several Fluxions be thus represented, viz,

$\dot{x} = Pp$  the Fluxion of the line  $AP$ ,  
 $a\dot{x} = PQ \cdot qp$  the Fluxion of  $ABQP$  in art. 6,  
 $y\dot{x} = PQRp$  the Fluxion of  $APQ$  in art. 8,


$\dot{z} = Qq = \sqrt{\dot{x}^2 + \dot{y}^2}$  the Fluxion of  $AQ$ ,  
and  $a\dot{x} = Pr$  the Flux. of the solid in art. 7, if

$a$  denote the constant generating plane  $PQR$ .  
Also  $n\dot{x} = BQ$  in the figure to art. 11,  
and  $n\dot{v} = Qq$  the Fluxion of the same.

14. The principles and notation being now laid down, we may proceed to the practice and rules of this doctrine, which consists of two principal parts, called the direct and inverse method of Fluxions; viz, the direct method, which consists in finding the Fluxion of any proposed fluent or flowing quantity; and the inverse method, which consists in finding the fluent of any proposed Fluxion. As to the former of these two problems, it can always be determined, and that in finite algebraic terms; but the latter, or finding of fluents, only in some certain cases, except by means of infinite series.—First then, of

#### *The Direct Method of FLUXIONS.*

15. To find the Fluxion of the product or rectangle of two variable quantities; let  $ARQP = xy$  be the flowing or variable rectangle, generated by two lines  $RQ$  and  $PQ$  moving always perpendicular to each other, from the positions  $AP$  and  $AR$ ; denoting the one by  $x$ , and the other by  $y$ ; and suppose  $x$  and  $y$  to be so related, that the curve  $AQ$  always passes through their intersection  $Q$ , or the opposite angle of the rectangle.



Now this rectangle consists of the two trilineal spaces  $APQ$ ,  $ARQ$ , of which the Fluxion of the former is  $PQ \times Pp$  or  $\dot{x}y$ , and that of the latter is  $RQ \times Rr$  or  $x\dot{y}$ , by art. 8; therefore the sum of the two,  $\dot{x}y + x\dot{y}$ , is the Fluxion of the whole rectangle  $xy$  or  $ARQP$ .

*The same otherwise.*—Let the sides of the rectangle,  $x$  and  $y$ , by flowing, become  $x + x'$  and  $y + y'$ : then the product of the two, or  $xy + x'y' + x'y + y'y$  will be the new or contemporaneous value of the flowing rectangle  $PR$  or  $xy$ ; subtract the one value from the other, and



and the remainder  $\dot{x}y + \dot{a}y + \dot{a}'y$ , will be the increment generated in the same time as  $\dot{x}$  or  $\dot{y}$ ; of which the last term  $\dot{a}'y$  is nothing, or indefinitely small in respect of the other two terms, because  $\dot{x}$  and  $\dot{y}$  are indefinitely small in respect of  $x$  and  $y$ ; which term being therefore omitted, there remains  $\dot{x}y + \dot{a}y$  for the value of that increment: and hence, by substituting  $\dot{x}$  and  $\dot{y}$  for  $\dot{a}$  and  $\dot{a}'$ , to which they are proportional, there arises  $\dot{x}y + \dot{a}y$  for the value of the Fluxion of  $xy$ ; the same as before.

17. Hence may be derived the Fluxions of all powers and products, and of all other forms of algebraic quantities whatever. And first for the continual products of any number of quantities, as  $xyz$ , or  $wxyz$ , or  $vwxxyz$ , &c. For  $xyz$  put  $q$  or  $pz$ , so that  $p = xy$ , and  $xyz = pz = q$ . Now, taking the Fluxion of  $q = pz$ , by the last article, it is  $\dot{q} = \dot{p}z + p\dot{z}$ ; but  $p = xy$ , and so  $\dot{p} = \dot{x}y + x\dot{y}$  by the same article; substituting therefore these values of  $p$  and  $\dot{p}$  instead of them, in the value of  $q$ , this becomes  $\dot{q} = \dot{x}yz + x\dot{y}z + xy\dot{z}$ , the Fluxion of  $xyz$  required; which is therefore equal to the sum of the products arising from the Fluxion of each letter or quantity multiplied by the product of the other two.

Again, to determine the Fluxion of  $wxyz$ , the continual product of four variable quantities; put this product, viz,  $wxyz$  or  $qw = r$ , where  $q = xyz$  as above; then, taking the Fluxion by the last article,  $\dot{r} = \dot{q}w + q\dot{w}$ ; and this, by substituting for  $q$  and  $\dot{q}$  their values as above, becomes

$\dot{r} = \dot{w}xyz + w\dot{x}y + w\dot{y}z + wxy\dot{z}$ , the Fluxion of  $wxyz$  as required; consisting of the Fluxion of each quantity drawn into the products of the other three.

In the very same manner it is found that the Fluxion of  $vwxxyz$  is

$\dot{v}wxyz + v\dot{w}xyz + vw\dot{x}y + vw\dot{y}z + vwxy\dot{z}$ ; and so on, for any number of quantities whatever; in which it is always found that there as many terms as there are variable quantities in the proposed fluent, and that these terms consist of the Fluxion of each variable quantity multiplied by the product of all the rest of the quantities.

18. From hence is easily derived the Fluxion of any power of a variable quantity, as of  $x^2$ , or  $x^3$ , or  $x^4$ , &c. For, in the rectangle or product  $xy$ , if  $x = y$ , then is the product  $xy = xx$  or  $x^2$ , and also its Fluxion  $\dot{x}y + x\dot{y} = \dot{x}x + x\dot{x}$  or  $2x\dot{x}$ , the Fluxion of  $x^2$ .

Again, if all the three  $x, y, z$  be equal; then is the product of the three  $xyz = xxx$  or  $x^3$ ; and its Fluxion  $\dot{x}yz + x\dot{y}z + xy\dot{z} = \dot{x}xx + x\dot{x}x + xx\dot{x}$  or  $3x^2\dot{x}$ , the Fluxion of  $x^3$ .

And in the same manner it will appear that the Fluxion of  $x^4$  is  $= 4x^3\dot{x}$ ,

that of  $x^5$  is  $= 5x^4\dot{x}$ ,

that of  $x^n$  is  $= nx^{n-1}\dot{x}$ ;

where  $n$  is any positive whole number. That is, the Fluxion of any positive integral power, is equal to the exponent of the power ( $n$ ), multiplied by the next less power of the same quantity ( $x^{n-1}$ ), and by the Fluxion of the root ( $\dot{x}$ ).

19. Next, for the Fluxion of any fraction, as  $\frac{x}{y}$  of one variable quantity divided by another; put the proposed fraction  $\frac{x}{y} = q$ ; then multiplying by the denominator,  $x = qy$ ; and, taking the Fluxions,

$\dot{x} = \dot{q}y + q\dot{y}$ , or  $\dot{q}y = \dot{x} - q\dot{y}$ ; and, by division,

$\dot{q} = \frac{\dot{x}}{y} - \frac{q\dot{y}}{y}$  (by substituting the value of  $q$ , or  $\frac{x}{y}$

$\frac{\dot{x}}{y} - \frac{xy}{y^2} = \frac{\dot{x}y - xy\dot{y}}{y^2}$  the Fluxion of  $\frac{x}{y}$ , as required.

That is, the Fluxion of any fraction, is equal to the Fluxion of the numerator drawn into the denominator, minus the Fluxion of the denominator drawn into the numerator, and the remainder divided by the square of the denominator.

20. Hence too is easily derived the Fluxion of any negative integer power of a variable quantity, as of  $x^{-n}$  or  $\frac{1}{x^n}$ , which is the same thing. For here the

numerator of the fraction is 1, whose Fluxion is nothing; and therefore, by the last article, the Fluxion of such a fraction, or negative power, is barely equal to minus the Fluxion of the denominator, divided by the square of the said denominator. That is, the Fluxion of

$x^{-n}$ , or  $\frac{1}{x^n}$ , is  $-\frac{n x^{n-1} \dot{x}}{x^{2n}}$  or  $-\frac{n \dot{x}}{x^{n+1}}$

or  $-n x^{-n-1} \dot{x}$ ; which is the same rule as before for integral powers.

Or, the same thing is otherwise derived immediately from the Fluxion of a rectangle or product, thus: put

the proposed fraction, or quotient,  $\frac{1}{x^n} = q$ ; then is

$qx^n = 1$ ; and, taking the Fluxions,

$\dot{q}x^n + qn x^{n-1} \dot{x} = 0$ ; hence  $\dot{q}x^n = -qn x^{n-1} \dot{x}$ , and (dividing by  $x^n$ ),

$\dot{q} = -\frac{qn \dot{x}}{x} =$  (by substituting  $\frac{1}{x^n}$  for  $q$ ),  $-\frac{n \dot{x}}{x^{n+1}}$  or

$-n x^{-n-1} \dot{x}$ ; the same as before.

21. Much in the same manner is obtained the Fluxion of any surd, or fractional power of a fluent

quantity, as of  $x^{\frac{m}{n}}$  or  $\sqrt[n]{x^m}$ . For, putting the proposed quantity

$x^{\frac{m}{n}} = q$ , then,

raising each to the  $n$  power,  $x^m = q^n$ ;

take the Fluxions,  $mx^{m-1} \dot{x} = nq^{n-1} \dot{q}$ ;

divide by  $nq^{n-1}$ ,  $\dot{q} = \frac{mx^{m-1} \dot{x}}{nq^{n-1}} = \frac{m^{m-1} \dot{x}}{nx^{\frac{m}{n}} - \frac{m}{n}}$


$\frac{m}{n} x^{\frac{m}{n}-1} \dot{x}$ : which is still the same rule as before, for

finding the Fluxion of any power of a fluent quantity, and which is therefore general, whether the exponent be positive or negative, or integral or fractional.

22. For



22. For the Fluxions of Logarithms: Let A be the principal vertex of an hyperbola, having its asymptotes CD, CP, with the ordinates DA, BA, PQ, &c, parallel to them. Then, from the nature of the hyperbola, and of logarithms, it is known that any space ABPQ is the log. of the ratio of CB to CP, to the modulus ABCD. Now put  $1 = CB$  or BA the side of the square or rhombus DB;  $m =$  the modulus, or area of DB, or sine of the angle C to the radius 1; also the absciss  $CP = x$ , and the ordinate  $PQ = y$ . Then, by the nature of the hyperbola,  $CP \times PQ$  is always equal to DB, that is,  $xy = m$ ; hence  $y = \frac{m}{x}$ , and the fluxion of the space,



or  $\dot{xy}$  is  $\frac{m\dot{x}}{x} = PQqp$  the fluxion of the log. of  $x$ , to the modulus  $m$ . And in the ordinary hyp. logarithms the modulus  $m$  being 1, therefore  $\frac{\dot{x}}{x}$  is the fluxion of the hyp. log. of  $x$ ; which is therefore equal to the Fluxion of the quantity, divided by the quantity itself. And the same might be brought out in several other ways, independent of the figure of the hyperbola.

23. By means of the Fluxions of logarithms, are determined those of exponential quantities, i. e. quantities which have their exponent also a flowing or variable quantity. These exponentials are of two kinds, viz, when the root is a constant quantity, as  $e^x$ ; and when the root is variable, as  $y^x$ .

In the former case, put the proposed exponential  $e^x = z$ , a single variable quantity; then take the logarithm of each, so shall log. of  $z = x \times \log. \text{ of } e$ ; take the fluxions of these, so shall  $\frac{\dot{z}}{z} = \dot{x} \times \log. \text{ of } e$ ; hence  $\dot{z} = z\dot{x} \times \log. \text{ of } e = e^x \dot{x} \times \log. \text{ of } e$ , the fluxion of the proposed exponential  $e^x$ ; and which therefore is equal to the said proposed quantity, drawn into the fluxion of the exponent, and also into the log. of the root.

24. Also in the 2d case, put the exponential  $y^x = z$ ; then the logarithms give log.  $z = x \times \log. y$ , and the fluxions give  $\frac{\dot{z}}{z} = \dot{x} \times \log. y + x \times \frac{\dot{y}}{y}$ ; hence  $\dot{z} =$

$z\dot{x} \times \log. y + \frac{zxy\dot{y}}{y} =$  (by substituting  $y^x$  for  $z$ )

$y^x \dot{x} \times \log. y + xy^{x-1}\dot{y}$ , is the fluxion of the proposed exponential  $y^x$ ; which therefore consists of two terms, of which the one is the fluxion of the proposed quantity considering the exponent only as constant, and the other is the fluxion of the same quantity considering the root as constant.

Of Second, Third, &c. Fluxions.—Having explained the manner of considering and determining the first fluxions of flowing or variable quantities; it remains now to consider those of the higher orders, as 2d, 3d, 4th, &c. fluxions.

25. If the rate or celerity with which any flowing quantity changes its magnitude, be constant, or the same, at every position; then is the fluxion of it also

constantly the same. But if the variation of magnitude be continually changing, either increasing or decreasing; then will there be a certain degree of fluxion peculiar to every point or position; and the rate of variation or change in the fluxion, is called the *Fluxion of the Fluxion*, or the *second Fluxion* of the given fluent quantity. In like manner, the variation or fluxion of this 2d fluxion is called the *third Fluxion* of the first proposed fluent quantity; and so on.

And these orders of fluxions are denoted by the fluent letter or quantity, with the corresponding number of points over it; viz, 2 points for the 2d fluxion, 3 for the 3d fluxion, 4 for the 4th fluxion, and so on. So the different orders of the fluxions of  $x$ , are  $\dot{x}$ ,  $\ddot{x}$ ,  $\dddot{x}$ ,  $\ddot{\ddot{x}}$ , &c; where each is the fluxion of the one next before it.

26. This description of the higher orders of fluxions may be illustrated by the three figures at the 8th article; where, if  $x$  denote the absciss AP, and  $y$  the ordinate PQ; and if the ordinate PQ or  $y$  flow along the absciss AP or  $x$ , with an uniform motion; then the fluxion of  $x$ , viz  $\dot{x} = Pp$  or  $Qr$  is a constant quantity, or  $\dot{x} = 0$ , in all the figures. Also, in fig. 1, in which AQ is a right line,  $y$  is  $= r q$ , or the fluxion of PQ, is a constant quantity, or  $\dot{y} = 0$ ; for, the angle Q, = the angle A, being constant,  $Qr$  is to  $r q$ , or  $\dot{x}$  to  $\dot{y}$ , in a constant ratio. But in the 2d figure,  $r q$ , or the fluxion of PQ, continually increases more and more; and in fig. 3 it continually decreases more and more; and therefore in both these cases  $y$  has a 2d fluxion, being positive in fig. 2, but negative in fig. 3: and so on for the other orders of fluxions.

27. Thus, if for instance, the nature of the curve be such, that  $x^3$  is everywhere equal to  $a^2 y$ ; then, taking the fluxions, it is  $a^2 \dot{y} = 3x^2 \dot{x}$ ; and, considering  $\dot{x}$  always as a constant quantity, and taking always the fluxions, the equations of the several orders of fluxions will be as below; viz,

the 1st fluxions  $a^2 \dot{y} = 3x^2 \dot{x}$ ,

the 2d fluxions  $a^2 \ddot{y} = 6x\dot{x}\dot{x}$ ,

the 3d fluxions  $a^2 \dddot{y} = 6\dot{x}^2$ ,

the 4th fluxions  $a^2 \ddot{\ddot{y}} = 0$ ,

and all the higher fluxions  $= 0$  or nothing.

Also the higher orders of fluxions are found in the same manner as the lower ones. Thus,

The 1st flux. of  $y^3$  is  $- - - 3y^2 \dot{y}$ ;


its 2d flux. or flux. of  $3y^2 \dot{y}$ , considered as the rectangle of  $3y^2$  and  $\dot{y}$ , is  $- - - 3y^2 \ddot{y} + 6y\dot{y}\dot{y}$ ;

and the flux. of this again, or the 3d fluxion of  $y^3$ , is  $- - - 3y^2 \ddot{\dot{y}} + 18y\dot{y}\ddot{y} + 6\dot{y}^3$ .

28. In the foregoing articles, it has been supposed that the fluents increase; or that their fluxions are positive; but it often happens that some fluents decrease, and that therefore their fluxions are negative: and whenever this is the case, the sign of the fluxion must be changed, or made contrary to that of the fluent. So, of the rectangle  $xy$ , when both  $x$  and  $y$  increase together, the fluxion is  $\dot{xy} + x\dot{y}$ ; but if one of them, as  $y$ , decrease, while the other,  $x$ , increases; then the fluxion of  $y$  being  $-\dot{y}$ , the fluxion of  $xy$  will in that case be  $\dot{xy} - x\dot{y}$ . This may be illustrated by the annexed rectangle.



tangle  $APQR = xy$ , supposed to be generated by the motion of the line  $PQ$  from  $A$  towards  $C$ , and by the motion of the line  $RQ$  from  $B$  towards  $A$ : For, by the motion of  $PQ$ , from  $A$  towards  $C$ , the rectangle is increased, and its fluxion is  $+xy$ ; but by the motion of  $RQ$ , from  $B$  towards  $A$ , the rectangle is decreased, and the fluxion of the decrease is  $-xy$ ; therefore, taking the fluxion of the decrease from that of the increase, the fluxion of the rectangle  $xy$ , when  $x$  increases and  $y$  decreases, is  $xy - xy$ .



For the Inverse Method, or the finding of fluents, see *FLUENT*. And for the several applications of this science to *MAXIMA* and *MINIMA*, the drawing of *TANGENTS*, &c, see the respective articles.

An idea of the principles of Fluxions being now delivered, as above, we may next consider somewhat of the chief writings and improvements that have been made by divers authors, since the first discovery of them: indeed some of the chief improvements may be learned by consulting the preface to Dr. Waring's *Meditationes Analyticae*.

The inventor himself brought the doctrine of Fluxions to a considerable degree of perfection; as may be seen by many specimens of this science, given by him; particularly in his *Principia*, in his *Treatise on Quadratures*, and in his *Treatise on Fluxions*, published by Mr. Colson; from all which it will appear, that he not only laid down the whole theory of this method, both direct and inverse; but also applied it in practice, to the solution of many of the most useful and important problems in mathematics and philosophy.

Various improvements however have been made by many illustrious authors on this science; particularly by John Bernoulli, who treated of the fluents belonging to the fluxions of exponential expressions; James Bernoulli, Craig, Cheyne, Cotes, Manfredi, Riccati, Taylor, Fagnanus, Clairaut, D'Alembert, Euler, Condorcet, Walmesley, Le Grange, Emerson, Simpson, Landen, Waring, &c. There are several other treatises also on the principles of Fluxions, by Hayes, Newyentyt, L'Hôpital, Hodson, Rowe, &c, &c, delivering the elements of this science in an easy and familiar way.

*FLY*, in Mechanics, a heavy weight applied to certain machines, to regulate their motions, as in a jack, or to increase their effect, as in the coining engine, &c; by means of which the force of the power is not only preserved, but equally distributed in all the parts of the revolution.

The Fly is either like a cross, with heavy weights at the ends of its arms; or like a heavy wheel at right-angles to the axis of motion. It may be applied to various sorts of engines, whether moved by men, horses, wind, or water; and is of great use in those parts of an engine having a quick circular motion, and where the power or resistance acts unequally in the different parts of a revolution. In this case the Fly becomes a moderator, making the motion of revolution almost everywhere equal.

*FLYERS*, in Architecture, such stairs as go straight, and do not wind round, nor have the steps made taper-

ing, but equally broad at both ends. Hence, if one flight do not rise to the top of the story &c, there is a broad half pace, and then commonly another set of flyers.

*FLYING*, the progressive motion of a bird, or other winged animal, through the air.

The parts of birds chiefly concerned in Flying, are the wings and tail: by the former, the bird sustains and wafts himself along; and by the latter he is assisted in ascending and descending, to keep his body poised and upright, and steady. The wings are extended or stretched quite out, and then struck forcibly downwards against the air, which by its resistance raises the bird upwards; then to make another stroke, the wing, by means of its joints, readily closes in some degree, presenting the sharp edge of the pinion foremost to cut the air, and drawing the collapsed feathers after it like a flag, to diminish the resistance to the ascent as much as possible; the wing and feathers are then stretched out horizontally again, and another downward stroke made, which raises the bird still higher; and so on as far as he pleases, or as the density of the air will sustain him; performing those motions of the wings very rapidly, that the flight may be the quicker.

*Artificial FLYING*, is that attempted by men, &c, by the assistance of mechanics.

The art of flying has been attempted by several persons in all ages. Friar Bacon, about 500 years ago, not only asserts the possibility of flying, but affirms that he himself knew how to make a machine with which a man might be able to convey himself through the air like a bird; and further adds, that it had been tried with success. Though the fact is to be doubted, if, as it was said, it consisted in the following method; viz, in a couple of large thin hollow copper globes, exhausted of air; which being much lighter than air, would sustain a chair on which a person might sit. Father Francisco Lana, in his *Prodromo*, proposes the same thing, as his own thought. He computes, that a round vessel of plate-brass, 14 feet in diameter, weighing 3 ounces the square foot, will only weigh 1848 ounces; whereas a quantity of air of the same bulk will weigh near 2156 ounces; so that the globe will not only be sustained in the air, but will carry with it a weight of 304 ounces; and by increasing the size of the globe, the thickness of the metal remaining the same, he adds, a vessel might be made to carry a much greater weight. But the fallacy is obvious: a globe of the dimensions he describes, as Dr. Hook observes, would not sustain the pressure of the air, but be crushed inwards. Indeed it is not probable that such a globe can be made of a thinness sufficient to float in the atmosphere after it is exhausted of air, and yet be strong enough to sustain the compressing force of the atmosphere. But for this purpose it seems that the globe should be filled with an air as elastic or strong as the atmosphere, and yet be very much lighter; such as has lately been used in the *Mongolfiers* and *Balloons*; the former of which is filled with common air heated, so as to be more elastic, and less heavy; and the latter with inflammable air, which is as elastic as the common air, with only about one tenth of its weight. And thus the idea of flying, or rather floating, in the air, has been lately realized by the moderns, using however a different sort of air. See *AEROSTATION*.



The same author describes a machine for Flying, invented by the Sieur Besnier, a smith of Sable, in the county of Main. See Philos. Collec. numb. 1.

By the foregoing method however, at best, only a method of floating can be obtained, like a log floating in a current; but not of Flying, which consists in moving through the air, independent of any current; and which must be effected by something in the nature of wings. Attempts of this latter kind also have indeed been made by several persons of late years; but it does not appear that any of them have been attended with such success as to induce the authors of those attempts to make them public. The philosophers of king Charles the second's reign were much busied about this art; and the celebrated bishop Wilkins was so confident of success in it, that he says, he does not question but in future ages it will be as usual to hear a man call for his wings, when he is going a journey, as it is now to call for his boots.

The story of the flight of Dædalus is well known.

**FLYING Pinion**, is part of a clock having a fly or fan with which to gather the air, and so bridle the rapidity of the clock's motion, when the weight descends in the striking part.

**FOCAL Distance**, the Distance of the Focus, which is sometimes understood as its distance from the vertex, as in the parabola; and sometimes its distance from the centre, as in the ellipse or hyperbola.

**FOCUS**, in Geometry and the Conic Sections, is applied to certain points in the Ellipse, Hyperbola, and Parabola, where the rays reflected from all parts of these curves do concur or meet; that is, rays issuing from a luminous point in the one focus, and falling on all points of the curves, are reflected into the other Focus, or into the line directed to the other Focus, viz, into the other Focus in the ellipse and parabola, and directly from it in the hyperbola. Which is the reason of the name Focus, or Burning-point. Hence, as the one Focus of the parabola is at an infinite distance; and consequently all rays drawn from it, to any finite part of the curve about the vertex, are parallel to one another; therefore if rays from the sun, or any other object so distant as that those rays may be accounted parallel, fall upon the curve of a parabola or concave surface of a paraboloidal figure, those rays will all be reflected into its Focus.

Thus, the rays  $Pf$ , from the Focus  $f$ , are reflected in the direction  $PF$ , into the other Focus  $F$ , in the ellipse and parabola, and form the Focus  $F$ , into  $FQ$ , in the hyperbola.

In all the three curves, the double ordinate  $CD$  drawn through the Focus  $F$ , is the parameter of the axis, or a 3d proportional to  $AB$  and  $ab$ , the transverse and conjugate axes.

In the ellipse and parabola, the transverse axis is equal to the sum of the two lines  $PF + Pf$ , drawn from the Foci to any point  $P$  in the curve; but in the hyperbola, the transverse is equal to the difference of those two lines. That is,

$AB = PF + Pf$  in the ellipse and parabola,


$AB = PF - Pf$  in the hyperbola.

In the ellipse and parabola, the square of the distance between the Foci, is equal to the difference of the squares of the two axes; and in the hyperbola, it is equal to the sum of their squares: that is

$Ff^2 = AB^2 - ab^2$  in the ellipse and parabola.

$Ff^2 = AB^2 + ab^2$  in the hyperbola.

Therefore the two semi-axes, with the distance of the Focus from the centre, form always a right-angled triangle  $FaE$ , or  $AaE$ .



In all the curves, the conjugate semi-axis is a mean proportional between the distances of either Focus from either end of the transverse axis: that is,

$$AF : Ea :: Ea : FB,$$

$$\text{or } Ea^2 = AF \cdot FB.$$

If there be any tangent to these curves, and two lines drawn from the Foci to the point of contact; these two lines will make equal angles with that tangent.

So, if  $GPG$  touch the curve at  $P$ ;  
then is the angle  $FPG = \angle fPG$ .

If a line be drawn from either Focus, perpendicularly upon a tangent; the distance of their intersection from the centre will be equal to the semi-transverse axis. So, if  $FH$  or  $fH$  be perpendicular to the tangent  $PH$ ; then is  $EH = EA$  or  $EB$ . Consequently, the circle described on the diameter  $AB$ , will pass through all the points  $H$ .

The foregoing are the chief properties that are common to the Foci of all the three conic sections. To which may be added the following properties which are peculiar to the parabola: viz,

In the parabola, the distance from the Focus to the vertex, is equal to  $\frac{1}{4}$  of the parameter, or half the ordinate at the Focus: viz,  $AF = \frac{1}{2} FC$ .

Also, a line drawn from the Focus to any point in the curve, is equal to the sum of the Focal distance from the vertex and the absciss of the ordinate to that point: i. e.  $FP = AF + AL$ .

If from any point of a parabola there be drawn a tangent, and a perpendicular to it  $PK$ , both to meet the axis produced; then the focus will be equally distant from the two intersections and the point of contact: i. e.  $FG = FP = FK$ .




Hence also the subnormal  $IK$  is  $= 2AF$  or  $= FC$  the semi-parameter.

The line drawn from the Focus to any point of the curve, is equal to  $\frac{1}{4}$  the parameter of the diameter to that point: i. e.  $FP = \frac{1}{4}$  the parameter of the diameter  $Pf$ .

If an ordinate to any diameter pass through the Focus, it will be equal to half its parameter; and its absciss equal to  $\frac{1}{4}$  of the same parameter; or the absciss equal to half the ordinate: i. e.  $PL = \frac{1}{4}MN = \frac{1}{2}LM$  or  $\frac{1}{2}LN$ .

Focus, in Optics, is a point in which several rays meet, and are collected, after being either reflected or refracted. It is so called, because the rays being here brought together and united, their force and effect are increased, inasmuch as to be able to burn; and therefore it is that bodies are placed in this point to be burnt, or to shew the effect of burning glasses, or mirrors.—It is to be observed however, that in practice, the Focus is not an absolute point, but a space of some small breadth, over which the rays are scattered; owing to the different nature and refrangibility of the rays of light, and to the imperfections in the figure of the lens, &c. However, the smaller this space is, the better, or the nearer to perfection the machine approaches. Huygens shews that the Focus of a lens convex on both sides, has its breadth equal to  $\frac{5}{8}$  of the thickness of the lens.

*Virtual Focus, or Point of Divergence*, so called by Mr. Molyneux, is the point from whence rays tend, after refraction or reflection; being in this respect opposed to the ordinary Focus, or Point of Concurrence, where rays are made to meet after refraction or reflection. Thus, the Foci of an hyperbola are mutually Virtual Foci to each other; but, in an ellipse, they are common Foci to each other: for the rays are reflected from the other Focus in the hyperbola, but towards it in the ellipse; as appears by the figures at the beginning of this article.



And, in Dioptrics, let  $ABC$  be the concavity of a glass, whose centre is  $D$ , and axis  $DE$ : Let  $FG$  be a ray of light falling on the glass, parallel to the axis  $DE$ ; this ray  $FG$ , after it has passed through the glass, at its emergence at  $G$  will not proceed directly to  $H$ , but be refracted from the perpendicular  $DG$ , and will become the ray  $GK$ , which being produced to meet the axis in  $E$ , this point  $E$  is the Virtual Focus, as the ray is refracted directly from this point.

#### *Rules for the Foci of Lenses and Mirrors.*

##### *I. In Catoptrics, or Lenses.*

1. The Focus of a convex glass, i. e. the point where parallel rays transmitted through a convex glass,

whose surface is the segment of a sphere, do unite, is distant from the pole or vertex of the glass, almost a diameter and half of the convexity.—2. In a Plano-Convex glass, the Focus of parallel rays is distant from the pole of the glass a diameter of the convexity, if the segment do not exceed 30 degrees. Or the rule in Plano-Convex glasses is, As 107 : 193 :: so is the radius of convexity : to the refracted ray taken to its concurrence with the axis; which in glasses of larger spheres is almost equal to the distance of the Focus taken in the axis.—3. In Double-Convex glasses of the same sphere, the Focus is distant from the pole of the glass about the radius of the convexity, if the segment be but 30 degrees. But when the two convexities are unequal, or segments of different spheres, then the rule is, As the sum of the radii of both convexities : to the radius of either convexity alone :: so is double the radius of the other convexity : to the distance of the Focus.—Here observe, that the rays which fall nearer the axis of any glass, are not united with it so near the pole of the glass as those farther off: nor will the Focal distance be so great, in a plano-convex glass, when the convex side is towards the object, as when the plane side is towards it. And hence it is truly concluded, that, in viewing any object by a plano-convex glass, the convex side should always be turned outward; as also in burning by such a glass.

##### *II. For the Virtual Focus, observe*

1. That in Concave glasses, when a ray falls from air parallel to the axis, the Virtual Focus, by its first refraction, becomes at the distance of a diameter and a half of the concavity.—2. In Plano-Concave glasses, when the rays fall parallel to the axis, the Virtual Focus is distant from the glass, the diameter of the concavity.—3. In Plano-Concave glasses, as 107 : 193 :: so is the radius of the concavity : to the distance of the Virtual Focus.—4. In Double Concaves of the same sphere, the Virtual Focus of parallel rays is at the distance of the radius of the concavity. But, whether the concavities be equal or unequal, the Virtual Focus, or point of divergency of the parallel rays, is determined by this rule; As the sum of the radii of both concavities : is to the radius of either concavity :: so is double the radius of the other concavity : to the distance of the Virtual Focus.—5. In Concave glasses, exposed to converging rays, if the point to which the incident ray converges, be farther distant from the glass than the Virtual Focus of parallel rays, the rule for finding the Virtual Focus of this ray, is this; As the difference between the distance of this point from the glass, and the distance of the Virtual Focus from the glass : is to the distance of the Virtual Focus :: so is the distance of this point of convergence from the glass : to the distance of the Virtual Focus of this converging ray.—6. In Concave glasses, if the point to which the incident ray converges, be nearer to the glass than the Virtual Focus of parallel rays, the rule to find where it crosses the axis, is this; As the excess of the Virtual Focus, more than this point of convergency : is to the Virtual Focus :: so is the distance of this point of convergency from the glass : to the distance of the point where this ray crosses the axis.







are preserved at large in the *Anecdotes of Bowyer*, p. 562. There are many memoirs of Mr. Folkes's in the *Philos. Transf.* from vol. 30 to vol. 46, both inclusive; viz, 1. Account of an Aurora Borealis, vol. 30.—2. Of Lieuwenhoek's curious Microscope, vol. 32.—3. On the Standard Measures in the Capitol at Rome, vol. 39.—4. Observations of three Mock-suns, vol. 40.—5. On the fresh-water Polypus, vol. 42.—6. On human bones petrified, vol. 43.—7. On a passage in Pliny's Natural History, vol. 44.—8. On an Earthquake at London, vol. 46.—9. Ditto at Kensington, vol. 46.—10. Ditto at Newton, vol. 46.

FOMAHAUT, or FOMALHAUT, a star of the first magnitude in the water of the constellation Aquarius, or in the mouth of the southern fish. Its latitude is  $21^{\circ} 6' 28''$  south, and mean longitude to the beginning of 1760,  $11^{\circ} 0' 28' 55''$ .

FONTENELLE (BERNARD DE), a celebrated French author, was born at Rouen in 1657, and died in 1756, when he was near 100 years old. He was a universal genius: at a very early age he wrote several comedies and tragedies of considerable merit; and he did the same at a very advanced age. Voltaire declares him the most universal genius the age of Lewis the 14th produced; and compares him to lands situated in so happy a climate, as to produce all sorts of fruits. His last comedies, though they shewed the elegance of Fontenelle, were however, little fitted for the stage; he then also produced an *Apology for the Vortices of Des Cartes*; upon which Voltaire says, "We must excuse his comedies, on account of his great age; and his Cartesian opinions, as they were those of his youth, when they were universally received all over Europe."

In his poetical performances and Dialogues of the Dead, the spirit of Voiture was discerned, though more extended and more philosophical. His *Plurality of Worlds* is a work singular in its kind: his design in this was, to present that part of philosophy to view in a gay and pleasing dress.

Fontenelle applied himself also to mathematics and natural philosophy; in which he proved not less successful than he had been in polite literature. Having been appointed perpetual secretary to the Academy of Sciences, he discharged that trust above 40 years with universal applause: his History of the Academy often throws great light upon their memoirs, which are sometimes obscure; and it has been said, he was the first who introduced elegance into the sciences. The *Eloges*, which he pronounced on the deceased members of the Academy, have this peculiar merit, that they excite a respect for the sciences as well as for the authors.

Upon the whole, Fontenelle must be looked upon as the great master of the new art of treating abstract sciences, in a manner that make their study at once easy and agreeable: nor are any of his works of other kinds void of merit. All those talents which he possessed from nature, were assisted by a good knowledge of history and languages: and he perhaps surpasses all men of learning who have not had the gift of invention.

Beside his poetical and theatrical works, with those of *Belles lettres*, &c, he published *Elemens de Geometrie de l'Infini*, in 4to, 1727; also the *Theorie des Tourbillons Cartesiens*; and *Discours moraux & philosophiques*.

All his different works were collected in eleven volumes, 12mo, under the title of *Oeuvres Diverses*.

FOOT, a measure of length, divided into 12 inches, and each inch supposed to contain 3 barley-corns in length. Geometricians divide the Foot into 10 digits, and the digit into 10 lines, &c. The French divide their Foot, as we do, into 12 inches; but their inch they divide into 12 lines.

It seems this measure has been taken from the length of the human Foot; but it is of different lengths in different countries. The Paris royal Foot is to the English Foot, as 4263 to 4000, and exceeds the English by  $9\frac{1}{2}$  lines; the ancient Roman Foot of the Capitol consisted of 4 palms; equal to 11 inches and  $\frac{7}{8}$  English; the Rhinland, or Leyden Foot, used by the northern nations, is to the Roman Foot, as 19 to 20. For the proportions of the Foot of several nations, compared with the English, see the article MEASURE.

Square FOOT, is a square whose side is 1 foot, or 12 inches, and consequently its area is 144 square inches.

Cubic FOOT, is a cube whose side is one Foot, or 12 inches, and consequently it contains  $12^3$  or 1728 cubic inches.

Foot-bank, or Foot-step, in Fortification. See BANQUETTE.

FORCE, *Vis*, or *Power*, in Mechanics, Philosophy, &c, denotes the cause of the change in the state of a body, with respect to motion, rest, pressure, &c; as well as its endeavour to oppose or resist any change made in such its state. Thus, whenever a body, which was at rest, begins to move; or when its motion is either not uniform, or not direct; the cause of this change in the state of the body, is what is called *Force*, and is an external Force. Or, while a body remains in the same state, either of rest, or of uniform and rectilinear motion, the cause of its remaining in such state, is in the nature of the body, being an innate internal Force, and is called its *Inertia*.

Mechanical Forces may be reduced to two sorts; one of a body at rest, the other of a body in motion.

The Force of a body at rest, is that which we conceive to be in a body lying still on a table, or hanging by a rope, or supported by a spring, &c; and this is called by the names of Pressure, Tension, Force, or *Vis Mortua*, *Solicitatio*, *Conatus Movendi*, *Conamen*, &c; which kind of Force may be always measured by a weight, viz, the weight that sustains it. To this class of Forces may also be referred Centripetal and Centrifugal Forces, though they reside in a body in motion; because these Forces are homogeneous to weights, pressures, or tensions of any kind. The pressure, or Force of gravity in any body, is proportional to the quantity of matter in it.

The Force of a body in motion, is a power residing in that body, so long as it continues its motion; by means of which, it is able to remove obstacles lying in its way; to lessen, destroy, or overcome the Force of any other moving body, which meets it in an opposite direction; or to surmount any the largest dead pressure or resistance, as tension, gravity, friction, &c, for some time; but which will be lessened or destroyed by such resistance as lessens or destroys the motion of the body. This is called *Vis Motrix*, Moving Force, or Motive Force, and by some late writers *Vis Viva*, to distinguish it



it from the *Vis Mortua* spoken of before; and by these appellations, however different, the same thing is understood by all mathematicians; namely, that power of displacing, of withstanding opposite moving Forces, or of overcoming any dead resistance, which resides in a moving body, and which, in whole or in part, continues to accompany it, as long as the body moves; and may be otherwise called Percussive Force, or Momentum.

But concerning the measure of this kind of Force, mathematicians have been divided into two parties. It is allowed by both sides, that the measure of this Force depends partly upon the mass of matter in the body, or its weight, and partly upon the velocity of its motion; so that upon any increase of either weight or velocity, the moving Force will become greater. It is also agreed, that the velocity being given, or being the same in two moving bodies, their Forces will be in proportion to their masses or weights. But, when two bodies are equal, and the velocities with which they move are different, the two parties no longer agree about the measure of the moving Force.

The Cartesians and Newtonians maintain, that, in this case, the moving Force is in proportion simply as the velocity with which a body moves; so that with a double velocity it has a double Force, &c. But the Leibnitians assert, that the moving Force is proportional to the square of the velocity; so as, with a double velocity to have a quadruple Force, &c. Or, when the bodies are different, the former hold, that the momentum or moving Force of bodies, is in the compound ratio of their weights and velocities: But the Leibnitians hold, that it is in the compound ratio of the weights and squares of the velocities.

Though Leibnitz was the first who expressly asserted, that the Force of a body in motion is as the square of its velocity, which was in a paper inserted in the *Leipscic Acts* for the year 1686, yet it is thought that Huygens led him into that notion, by some demonstrations in the 4th part of his book *De Horologio Oscillatorio*, relating to the centre of oscillation, and by his dissertations, in answer to the objections of the abbot Catalan, one of which was published in 1684. This eminent mathematician had demonstrated, that in the collision of two bodies that are perfectly elastic, the sum of the products of each body multiplied by the square of its velocity, was the same after the shock as before; (though the same thing is true of the sums of the products of the bodies multiplied simply by their velocities). Now that proposition is so far general as to obtain in all collisions of bodies that are perfectly elastic: and it is also true, when bodies of a perfect elasticity strike any immoveable obstacle, as well as when they strike one another; or when they are constrained by any power or resistance to move in directions different from those in which they impel one another. These considerations might have induced Huygens to lay it down as a general rule, that bodies constantly preserve their Ascensional Force, i. e. the product of their mass by the height to which their centre of gravity can ascend, which is as the square of the velocity; and therefore, in a given system of bodies, the sum of the squares of their velocities will remain the same, and not be altered by the action of the bodies among

themselves, nor against immoveable obstacles. Leibnitz's metaphysical system led him to think that the same quantity of action or Force subsisted in the universe; and finding this impossible, if Force were estimated by the quantity of motion, he adopted Huygens's principle of the preservation of the Ascensional Force, and made it the measure of moving Forces. But it is to be observed that Huygens's principle, above-mentioned, is general only when bodies are perfectly elastic; and in some other cases which Maclaurin has endeavoured to distinguish: shewing at the same time that no useful conclusion in mechanics is affected by the disputes concerning the measure of the Force of bodies in motion, which have been objected to mathematicians. *Analyst*, Query 9. See Maclaurin's *Fluxions*, vol. 2, art. 533; Huygens *Oper.* tom. 1, pa. 248; &c.

Leibnitz's principle was adopted by several persons; as Wolfius, the Bernoullis, &c. Mr. Dan. Bernoulli, in his *Treatise*, has assumed the preservation of the *Vis Ascendens* of Huygens, or, as others express it, the *Conservatio Virium Vivarum*; and, in Bernoulli's own expression, *æqualitas inter descensum actualem ascensumque potentialem*, as an hypothesis of wonderful use in mechanics. But a late author contends, that the conclusions drawn from this principle are oftener false than true. See *De Conservat. Virium Vivarum Dissert.* Lond. 1744.

Catalan and Papin answered Leibnitz's paper published in 1686; and from that time the controversy became more general, and was carried on for several years by Leibnitz, John and Daniel Bernoulli, Poleni, Wolfius, s'Gravesande, Camus, Muschenbroek, &c, on one side; and Pemberton, Eames, Defaguliers, Dr. S. Clark, M. de Mairan, Jurin, Maclaurin, Robins, &c, on the other. See *Act. Erud.* 1686, 1690, 1691, 1695; *Nouv. de la Rep. des Let.* Sept. 1686, 1687, art. 2; *Comm. Epist. inter Leibn. et Bern.* Ep. 24, p. 143; *Discourses sur les Loix de la Comm. du Mouvement*, *Oper.* tom. 3, & *Diff. de vera Notione Virium Vivarum*, ib.; *Act. Petropol.* tom. 1, p. 131, &c; *Hydronamica*, sect. 1; Herman, in *Act. Petrop.* tom. 1, p. 2, &c; Polen. de Castellis; Wolf. in *Act. Petrop.* tom. 1, p. 217, &c, and in *Cosmol. Gener.*; Graves. in *Journ. Lit. and Phys. Elem. Math.* 1742, lib. 2, cap. 2 and 3; *Memoir. de l'Acad. des Sciences* 1728; Muschenbr. *Int. ad Phil. Nat.* 1762, vol. 1, p. 83 &c; Pemb. &c, in *Phil. Transf.* number 371, 375, 376, 396, 400, 401, or *Abridg.* vol. 6, p. 216 &c. Mairan in *Memo. de l'Acad. des Sc.* 1728, *Phil. Transf.* numb. 459, or *Abridg.* vol. 8, p. 236, *Philos. Transf.* vol. 43, p. 423 &c; Maclaurin's *Acc. of Newton's Discoveries*, p. 117 &c, *Flux.* ubi supr. & *Recueil des Pieces qui ont emporté le Prix* &c. tom. 1; Defagul. *Course Exp. Philos.* vol. 1, p. 393 &c, vol. 2, p. 49 &c; and Robins's *Traacts*, vol. 2, p. 135.

The nature and limits of this work will not admit of a full account of the arguments and experiments that have been urged on both sides of this question; but they may be found chiefly in the preceding references. A few of them however may be considered, as follows.

The defenders of Leibnitz's principle, beside the arguments above-mentioned, refer to the spaces that bodies ascend to, when thrown upwards, or the penetrations of bodies let fall into soft wax, tallow, clay, snow, and



and other soft substances, which spaces are always as the squares of the velocities of the bodies. On the other hand, their opponents retort, that such spaces are not the measures of the force in question, which is rather percussive and momentary, as those above are passed over in unequal times, and are indeed the joint effect of the Forces and times.

Desaguliers brings an argument from the familiar experiment of the balance, and the other simple mechanic powers, shewing that the effect is in proportion to the velocity multiplied by the weight; for example, 4 pounds being placed at the distance of 6 inches from the centre of motion of a balance, and 2 pounds at the distance of 12 inches; these will have a *Vis Viva* if the balance be put into a swinging motion. Now it appears that these Forces are equal, because, with contrary directions, they soon destroy each other; and they are to each other in the simple ratio of the velocity multiplied by the mass, viz  $4 \times 6 = 24$ , and  $2 \times 12 = 24$  also.

Mr. Robins, in his remarks on J. Bernoulli's treatise, entitled, *Discours sur les Loix de la Communication du Mouvement*, informs us, that Leibnitz adopted this opinion through mistake; for though he maintained that the quantity of Force is always the same in the universe, he endeavours to expose the error of Des Cartes, who also asserted, that the quantity of motion is always the same; and in his discourse on this subject in the *Acta Eruditorum* for 1686, he says that it is agreed on by the Cartesians, and all other philosophers and mathematicians, that there is the same Force requisite to raise a body of 1 pound to the height of 4 yards, as to raise a body of 4 pounds to the height of 1 yard; but being shewn how much he was mistaken in taking that for the common opinion, which would, if allowed, prove the force of the body to be as the square of the velocity it moved with, he afterwards, rather than own himself capable of such a mistake, endeavoured to defend it as true; since he found it was the necessary consequence of what he had once asserted; and maintained, that the force of a body in motion was proportional to the height from which it must fall, to acquire that velocity; and the heights being as the squares of the velocities, the Forces would be as the masses multiplied by them; whereas, when a body descends by its gravity, or is projected perpendicularly upwards, its motion may be considered as the sum of the uniform and continual impulses of the power of gravity, during its falling in the former case, and till they extinguish it in the latter. Thus when a body is projected upwards with a double velocity, these uniform impulses must be continued for a double time, in order to destroy the motion of the body; and hence it follows, that the body, by setting out with a double velocity, and ascending for a double time, must arise to a quadruple height, before its motion is exhausted. But this proves that a body with a double velocity moves with a double Force, since it is produced or destroyed by the same uniform power continued for a double time, and not with a quadruple force, though it rises to a quadruple height; so that the error of Leibnitz consisted in his not considering the time, since the velocities alone are not the causes of the spaces described, but the times and the velocities together; yet this is the fallacious argument on which he first built his new doctrine;

and those which have been since much insisted on, and derived from the indentings or hollows produced in soft bodies by others falling into them, are much of the same kind. Robins's *Tracts*, vol. 2, p. 178.

But many of the experiments and reasonings, that have been urged on both sides in this controversy, have been founded in the different senses applied to the term Force. The English and French philosophers, by the word Force, mean the same thing as they do by momentum, motion, quantity of motion, percussion, or instantaneous pressure, which is measured by the mass drawn into the velocity, and may be known by its effect; and when they consider bodies as moving through a certain space, they allow for the time in which that space is described: whereas the Dutch, Italian, and German philosophers, who have espoused the new opinion, mean by the word Force, or Force inherent in a body in motion, that which it is able to produce; or, in other words, the Force is always measured by the whole effect produced by the body in motion; until its whole Force be entirely communicated or destroyed, without any regard to the time employed in producing this total effect. Thus, say they, if a point runs through a determinate space, and presses with a certain given Force, or intensity of pressure, it will perform the same action whether it move fast or slow, and therefore the time of the action in this case ought not to be regarded. *s'Gravesande, Phys. El. Math. § 723—728.*

Mr. Euler observes, with regard to this dispute concerning the measure of vivid Force, or living Force, as it is sometimes called, that we cannot absolutely ascribe any Force to a body in motion, whether we suppose this Force proportional to the velocity, or to the square of the velocity: for the Force exerted by a body, striking another at rest, is different from that which it exerts in striking the same body in motion; so that this Force cannot be ascribed to any body considered in itself, but only relatively to the other bodies it meets with. There is no force in a body absolutely considered, but its inertia, which is always the same, whether the body be in motion or at rest. But if this body be forced by others to change its state, its inertia then exerts itself as a Force, properly so called, which is not absolutely determinable; because it depends on the changes that happen in the state of the body.

A second observation which has been made by some eminent writers, is, that the effect of a shock of two or more bodies, is not produced in an instant, but requires a certain interval of time. If this be so, the heterogeneity between the *Vis Viva* and *Mortua*, or Living and Dead Force, will vanish; since a pressure may always be assigned, which in the same time, however little, shall produce the same effect. If then the *Vis Viva* be homogeneous to the *Vis Mortua*, and having a perfect measure and knowledge of the latter, we need require no other measure of the former than that which is derived from the *Vis Mortua* equivalent to it.

Now that the change in the state of two bodies, by their shock, does not happen in an instant, appears evidently from the experiments made on soft bodies: in these, percussion forms a small cavity, visible after the shock, if the bodies have no elasticity. Such a cavity cannot certainly be made in an instant. And if the

shock



shock of soft bodies require a determinate time, we must certainly say as much of the hardest, though this time may be so small as to be beyond all our ideas. Neither can an instantaneous shock agree with that constant law of nature, by virtue of which nothing is performed *per saltum*. But it is needless to insist farther upon this, since the duration of any shock may be determined from the most certain principles.

There can be no shock or collision of bodies, without their making mutual impressions on each other: these impressions will be greater or less, according as the bodies are more or less soft, other circumstances being the same. In bodies called hard, the impressions are small; but a perfect hardness, which admits of no impression, seems inconsistent with the laws of nature; so that while the collision lasts, the action of bodies is the result of their mutually pressing each other. This pressure changes their state; and the Forces exerted in percussion are really pressures, and truly *Vires Mortuæ*, if we will use this expression, which is no longer proper, since the pretended infinite difference between the *Vires Vivæ* and *Mortuæ* ceases.

The Force of percussion, resulting from the pressures that bodies exert on each other, while the collision lasts, may be perfectly known, if these pressures be determined for every instant of the shock. The mutual action of the bodies begins the first moment of their contact; and is then least; after which this action increases, and becomes greatest when the reciprocal impressions are strongest. If the bodies have no elasticity, and the impressions they have received remain, the Forces will then cease. But if the bodies be elastic, and the parts compressed restore themselves to their former state, then will the bodies continue to press each other till they separate. To comprehend therefore perfectly the Force of percussion, it is requisite first to define the time the shock lasts, and then to assign the pressure corresponding to each instant of this time; and as the effect of pressures in changing the state of any body may be known, we may thence come at the true cause of the change of motion arising from collision. The Force of percussion therefore is no more than the operation of a variable pressure during a given time; and to measure this Force, we must have regard to the time, and to the variations according to which the pressure increases and decreases.

Mr. Euler has given some calculations relative to these particulars; and he illustrates their tendency by this instance: Suppose that the hardness of the two bodies, A and B, is equal; and such, that being pressed together with the force of 100lb, the impression made on each is of the depth of  $\frac{1}{800}$ th part of a foot. Suppose also that B is fixed, and that A strikes it with the velocity of 100 feet in a second; according to Mr. Euler, the greatest Force of compression will be equivalent to 400lb; and this Force will produce in each of these bodies an impression equal to  $\frac{1}{25}$  of a foot; and the duration of the collision, that is, till the bodies arrive at their greatest compression, will be about  $\frac{1}{800}$  of a second. Mr. Euler, in his calculations, supposes the hardness of a body to be proportional to the Force or pressure requisite to make a given impression on it; so that the Force by which a given impression is made on a body, is in a compound ratio of the hardness of

the body and of the quantity of the impression. But he observes, that regard must be had to the magnitude of the bodies, as the same impression cannot be made on the least bodies as on the greatest, from the defect of space through which their component particles must be driven: he considers therefore only the least impressions, and supposes the bodies of such magnitudes, that with respect to them the impressions may be looked upon as nothing. What he supposes concerning the hardness of bodies, neither implies elasticity nor the want of it, as elasticity only produces a restitution of figure and impression when the pressing Force ceases; but this restitution need not be here considered. It is also supposed, that the bodies which strike each other, have plane and equal bases, by which they touch each other in the collision; so that the impression hereby made diminishes the length of each body. It is farther to be observed, that in Mr. Euler's calculations, bodies are supposed so constituted, that they may not only receive impressions from the Forces pressing them, but that a greater Force is requisite to make a greater impression. This excludes all bodies, fluid or solid, in which the same Force may penetrate farther and farther, provided it have time, without ever being in equilibrio with the resistance: thus a body may continually penetrate farther into soft wax, although the force impelling it be not increased: in these, and the like cases, nothing is required but to surmount the first obstacles; which being once done, and the connection of parts broken, the penetrating body always advances, meeting with the same obstacles as before, and destroying them by an equal Force. But Mr. Euler only considers the first obstacles which exist before any separation of parts, and which are doubtless such, that a greater impression requires a greater Force. Indeed this chiefly takes place in elastic bodies; but it seems likewise to obtain in all bodies when the impressions made on them are small, and the contexture of their parts is not altered.

These things being premised, let the mass or weight of the body A be expressed in general by A, and let its velocity before the shock be that which it might acquire by falling from the height *a*. Farther, let the hardness of A be expressed by M, and that of B by N, and let the area of the base, on which the impression is made, be *cc*; then will the greatest compression be

made with the Force  $\sqrt{\frac{MNcc}{M+N}} \times Aa$ . Therefore

if the hardness of the two bodies, and the plane of their contact during the whole time of their collision be the same, this Force will be as  $\sqrt{Aa}$ , that is, as the square root of the *Vis Viva* of the striking body A. And as  $\sqrt{a}$  is proportional to the velocity of the body A, the Force of percussion will be in a compound ratio of the velocity and of the subduplicate ratio of the mass of the body striking; so that in this case neither the Leibnitian nor the Cartesian propositions take place. But this Force of percussion depends chiefly on the hardness of the bodies; the greater this is, the greater will the Force of percussion be. If  $M = N$ , this Force will be as  $\sqrt{Mcc} \times Aa$ , that is, in a compound subduplicate ratio of the *Vis Viva* of the striking body, of the hardness, and of the plane of contact. But if



$M$ , the hardness of one of the bodies, be infinite, the Force of percussion will be as  $\sqrt{Ncc \times Aa}$ ; at the same time, if  $M = N$ , this Force will be as  $\sqrt{\frac{1}{2}Ncc \times Aa}$ . Therefore, all other things being equal, the Force of percussion, if the striking body be infinitely hard, will be to the Force of percussion when both the bodies are equally hard, as  $\sqrt{2}$  to 1.

Mr. Euler farther deduces from his calculation, that the impression received by the bodies  $A$  and  $B$  will be as follows; viz, as

$$\sqrt{\frac{N \times Aa}{M + N \times Mcc}} \text{ and } \sqrt{\frac{M \times Aa}{M + N \times Ncc}} \text{ respectively.}$$

If therefore the hardness of  $A$ , that is  $M$ , be infinite, it will suffer no impression; whereas that on  $B$  will extend to the depth of  $\sqrt{\frac{Aa}{Ncc}}$ . But if the hardness of the two bodies be the same, or  $M = N$ , they will each receive equal impressions of the depth  $\sqrt{\frac{Aa}{2Ncc}}$ .

So that the impression received by the body  $B$  in this case, will be to the impression it receives in the former, as 1 to  $\sqrt{2}$ .

Mr. Euler has likewise considered and computed the case where the striking body has its anterior surface convex, with which it strikes an immoveable body whose surface is plane. He has also examined the case when both bodies are supposed immoveable; and from his formulæ he deduces the known laws of the collision of elastic and non-elastic bodies. He has also determined the greatest pressures the bodies receive in these cases; and likewise the impressions made on them. In particular he shews, that the impressions received by the body struck, or  $B$ , if moveable, is to the impression received by the same body when immoveable, as  $\sqrt{B}$  to  $\sqrt{A + B}$ .

There are several curious as well as useful observations in Defaguliers's Experimental Philosophy, concerning the comparative Forces of men and horses, and the best way of applying them. A horse draws with the greatest advantage when the line of direction is level with his breast; in such a situation, he is able to draw 200lb for 8 hours a day, walking about  $2\frac{1}{2}$  miles an hour. But if the same horse be made to draw 240lb, he can work only 6 hours a day, and cannot go quite so fast. On a carriage, indeed, where friction alone is to be overcome, a middling horse will draw 1000lb. But the best way to try the Force of a horse, is to make him draw up out of a well, over a single pulley or roller; and in that case, an ordinary horse will draw about 200lb, as before observed.

It is found that 5 men are of equal Force with one horse, and can with equal ease push round the horizontal beam of a mill, in a walk 40 feet wide; whereas 3 men will do it in a walk only 19 feet wide.

The worst way of applying the Force of a horse is to make him carry or draw up hill: for if the hill be steep, 3 men will do more than a horse, each man climbing up faster with a burden of 100lb weight, than a horse that is loaded with 300lb: a difference which is owing to the position of the parts of the human body being better adapted to climb, than those of a horse.

On the other hand, the best way of applying the Force of a horse, is the horizontal direction, in which a man can exert the least Force: thus, a man that weighs 140lb, when drawing a boat along by means of a rope coming over his shoulders, cannot draw above 27lb, or exert above  $\frac{1}{7}$ th part of the Force of a horse employed to the same purpose; so that in this way the Force of a horse is equal to that of 7 men.

The best and most effectual posture in a man, is that of rowing; when he not only acts with more muscles at once for overcoming the resistance, than in any other position; but also as he pulls backwards, the weight of his body assists by way of lever. See Defaguliers's Exp. Philos. vol. 1, p. 241, where several other observations are made relative to Force acquired by certain positions of the body; from which that author accounts for most feats of strength and activity. See also a Memoir on this subject by M. De la Hire, in the Mem. Roy. Acad. 1729; or in Defaguliers's Exp. &c. p. 267 &c., who has published a translation of part of it with remarks.

FORCE is distinguished into Motive and Accelerative or Retardive.

*Motive* FORCE, otherwise called Momentum, or Force of Percussion, is the absolute Force of a body in motion, &c; and is expressed by the product of the weight or mass of matter in the body multiplied by the velocity with which it moves. But

*Accelerative* FORCE, or *Retardive* FORCE, is that which respects the velocity of the motion only, accelerating or retarding it; and it is denoted by the quotient of the motive Force divided by the mass or weight of the body. So,

if  $m$  denote the motive Force,  
and  $b$  the body, or its weight,  
and  $f$  the accelerating or retarding Force,

then is  $f$  as  $\frac{m}{b}$ .

Again, Forces are either Constant or Variable.

*Constant* FORCES are such as remain and act continually the same for some determinate time. Such, for example, is the Force of gravity, which acts constantly the same upon a body while it continues at the same distance from the centre of the earth, or from the centre of Force, wherever that may be. In the case of a constant Force  $F$  acting upon a body  $b$ , for any time  $t$ , we have these following theorems; putting

$f$  = the constant accelerating Force =  $F \div b$ ,

$v$  = the velocity at the end of the time  $t$ ,

$s$  = the space passed over in that time, by the constant action of that Force on the body:

and  $g = 16\frac{1}{2}$  feet, the space generated by gravity in 1 second, and calling the accelerating Force of gravity 1; then is

$$s = \frac{1}{2}tv = gft^2 = \frac{v^2}{4gf};$$

$$v = 2gft = \frac{2s}{t} = \sqrt{4gfs};$$

$$t = \frac{v}{2gf} = \frac{2s}{v} = \sqrt{\frac{s}{gf}};$$

$$f = \frac{v}{2gt} = \frac{s}{gt^2} = \frac{v^2}{4gs}.$$

*Variable* FORCES are such as are continually changing



ing in their effect and intensity; such as the Force of gravity at different distances from the centre of the earth, which decreases in proportion as the square of the distance increases. In variable Forces, theorems similar to those above may be exhibited by using the fluxions of quantities, and afterwards taking the fluents of the given fluxional equations. And herein consists one of the great excellencies of the Newtonian or modern analysis, by which we are enabled to manage, and compute the effects of all kinds of variable Forces, whether accelerating or retarding. Thus, using the same notation as above for constant forces, viz,  $f$  the accelerating Force at any instant,  $t$  the time a body has been in motion by the action of the variable Force,  $v$  the velocity generated in that time,  $s$  the space run over in that time, and  $g = 16\frac{1}{2}$  feet; then is

$$\dot{s} = \frac{v\dot{v}}{2gf} = v\dot{t};$$

$$\dot{v} = \frac{2gfs}{v} = 2gft;$$

$$\dot{t} = \frac{\dot{s}}{v} = \frac{\dot{v}}{2gf};$$

$$f = \frac{v\dot{v}}{2gs} = \frac{\dot{v}}{2gt}.$$

In these four theorems, the Force  $f$ , though variable, is supposed to be constant for the indefinitely small time  $t$ ; and they are to be used in all cases of variable Forces, as the former ones in constant Forces; viz, from the circumstances of the problem under consideration, deduce a general expression for the value of the force  $f$ , at any indefinite time  $t$ ; then substitute it in one of these theorems, which shall be proper to the case in hand; and the equation thence resulting will determine the corresponding values of the other quantities in the problem.

It is also to be observed, that the foregoing theorems equally hold good for the destruction of motion and velocity, by means of retarding or resisting Forces, as for the generation of the same by means of accelerating Forces.

Many applications of these theorems may be seen in my *Select Exercises*, p. 172 &c.

There are many other denominations and kinds of Forces; such as attractive, central, centrifugal, &c, &c; for which see the respective words.

**FORCER**, in Mechanics, is properly a piston without a valve. For, by drawing up such a piston, the air is drawn up, and the water follows; then pushing the piston down again, the water, being prevented from descending by the lower valve, is forced up to any height above, by means of a side branch between the two.


See the ways of making these in *Desaguliers's Exper. Philos.* vol. 2, p. 161 &c. See also *Clare's Motion of Fluids*, p. 60.

**FORCING Pump**, one that acts, or raises water, by a Forcing piston. See above.

**FORELAND**, or **FORENESS**, in Navigation, a point of land jutting out into the sea.

**FORELAND**, in Fortification, is a small piece of ground between the wall of a place and the moat; called also *Berne* and *Liziere*.

**FORE-STAFF**, an instrument used at sea, for taking the altitudes of the heavenly bodies; being so called, because the observer, in using it, turns his face forward or towards the object, in contradistinction to the *Back-staff*, with which he turns his back to the object. It is also called the *Cross-staff*, because it consists of several pieces set across a staff.



The Fore-staff is formed of a straight square staff AB, of about 3 feet long, having each of its four sides graduated like a line of tangents, and four crosses, or vanes, FF, EE, DD, CC, sliding upon it, of unequal lengths, the halves of which represent the radii to the lines of tangents on the different sides of the staff. The first or shortest of these vanes, FF, is called the *Ten Cross*, or *Ten Vane*, and belongs to the 10 scale, or that side of the instrument on which divisions begin at 3 degrees, and end at 10. The next longer cross, EE, is called the *30 Cross*, belonging to that side of the staff where the divisions begin at 10 degrees, and end at 30, called the *30 scale*. The third vane DD, is called the *60 Cross*, and belongs to that side where the divisions begin at 20 degrees, and end at 60. The last or longest vane CC, called the *90 Cross*, belongs to the side where the divisions begin at 30 degrees, and end at 90.

The chief use of this instrument, is to take the height of the sun, and stars, or the distance between two stars: and the 10, 30, 60, or 90 Cross is to be used, according as the altitude is more or less; that is, if the altitude be less than 10 degrees, the 10 Cross is to be used; if above 10, but less than 30, the 30 Cross is to be used; and so on.

*To observe an Altitude with the Fore-staff.* Apply the flat end of the staff to the eye, and slide one of the crosses backwards and forwards upon it, till over the upper end of the Cross be just seen the centre of the sun or star, and over the under end the extreme horizon; then the degrees and minutes cut by the cross on the side of the staff proper to the vane in use, gives the altitude above the horizon.

*In like manner, for the Distance between two luminaries;* the Staff being set to the eye, bring the cross just to subtend or cover that distance, by having the one luminary just at the one end of it, and the other luminary at the other end of it; and the degrees and minutes, in the distance, will be cut on the proper side of the staff, as before.

**FORMULA**, a theorem or general rule, or expression, for resolving certain particular cases of some problem, &c. So  $\frac{1}{2}s + \frac{1}{2}d$  is a general Formula for the



the greater of two quantities whose sum is  $s$  and difference  $d$ ; and  $\frac{1}{2}s - \frac{1}{2}d$  is the Formula, or general value, for the less quantity. Also  $\sqrt{dx - x^2}$  is the Formula, or general value, of the ordinate to a circle, whose diameter is  $d$ , and absciss  $x$ .

**FORT**, a little castle or fortress; or a place of small extent, fortified either by nature or art.

The Fort is usually encompassed with a moat, rampart, and parapet, to secure some high ground, or passage of a river; to make good or strengthen an advantageous post; or to fortify the lines and quarters of a siege.

*Field Fort*, otherwise called *Fortin*, or *Fortlet*, and sometimes *Sconce*, is a small Fort, built in haste, for the defence of a pass or post; but particularly constructed for the defence of a camp in the time of a siege, where the principal quarters are usually joined, or made to communicate with each other, by lines defended by fortins and redoubts. Their figure and size are various, according to the nature of the situation, and the importance of the service for which they are intended; but they are most commonly made square, each side about 100 toises, the perpendicular 10, and the faces 25; the ditch about this Fort may be 10 or 12 toises wide; the parapet is made of turf, and fraised, and the ditch pallisadoed when dry. There may be made a covert-way about this Fort, or else a row of pallisades might be placed on the outside of the ditch. Some of these are fortified with bastions, and some with demi-bastions.—A Fort differs from a citadel, as this last is erected to command and guard some town; and from a redoubt, as it is closed on all sides, while the redoubt is open on one side.

*Royal Fort*, is one whose line of defence is at least 26 fathoms long.

*Star Fort*, is a sconce or redoubt, constituted by re-entering and salient angles, having commonly from five to eight points, and the sides flanking each other.

*Forts* are sometimes made triangular, only with half bastions; or of various other figures, regular or irregular; and sometimes in the form of a semicircle, especially when they are situated near a river, or the sea, as at the entrance of a harbour, for the convenience of firing at ships quite around them on that side. In the construction of all Forts, it should be remembered, that the figure of fewest sides and bastions, that can probably answer the proposed defence, is always to be preferred; as works on such a plan are sooner executed, and with less expence; besides, fewer troops will serve, and they are more readily brought together in case of necessity.

**FORTIFICATION**, called also *Military Architecture*, is the art of fortifying or strengthening a town or other place, by making certain works around it, to secure it and defend it from the attacks of enemies.

Fortification has been undoubtedly practised by all nations, and in all ages; being at first doubtless very rude and simple, and varying in its nature and manner, according to the mode of attack, and the weapons made use of. Thus, when villages and towns were first formed, it was found necessary, for the common safety, to encompass them with walls and ditches, to prevent all violence and sudden surprises from their

neighbours. When offensive and missive weapons came to be used, walls were made as a defence against the assailants, and look holes or loop holes made through the walls to annoy the enemy, by shooting arrows &c through them. But finding that as soon as the enemy got close to the walls, they could no longer be seen nor annoyed by the besieged, these added square towers along the wall, at proper distances from each other, so that all the intervening parts of the wall might be seen and defended from the adjacent sides of the towers. However, this manner of inclosing towns was found to be rather imperfect, because there remained still the outer face of the towers which fronted the field, that could not be seen and defended from any other part. To remedy this imperfection, they next made the towers round instead of square, as seeming better adapted both for strength to resist the battering engines, and for being defended from the other parts of the wall. Nevertheless, a small part of these towers still remained unseen, and incapable of being defended, for which reason they were again changed for square ones, as before, but with this difference, that now they presented an angle of the square outwards to the field, instead of a face or side; and thus such a disposition of the works was obtained, as that no part could be approached by the enemy without being seen and attacked.

Since the use of gun-powder, it has been found necessary to add thick ramparts of earth to the walls, and the towers have been enlarged into bastions, as well as many other things added, that have given a new appearance to the whole art of defence, and the name of Fortification, on account of the strength afforded by it, which was about the year 1500, when the round towers were changed into bastions.

But notwithstanding all the improvements made in this art since the invention of gun-powder, that of attacking is still superior to it: the superiority of the besieger's fire, together with the greater number of men, sooner or later compels the besieged to submit. A special advantage was added to the art of attacking by M. Vauban, at the siege of Ath, in the year 1697, viz, in the use of ricochet firing, or at a low elevation of the gun, by which the shot was made to run and roll a great way along the inside of the works, to the great annoyance of the besieged.

The chief authors who have treated of Fortification, since it has been considered as a particular art, are the following, and mostly in the order of time: viz, La Treille, Alghisi, Marchi, Pasino, Ramelli, Cataneo, and Speckle, who, as Mr. Robins says, was one of the greatest geniuses that has applied to this art: he was architect to the city of Stralsburgh, and died in the year 1589: he published a treatise on Fortification in the German language, which was reprinted at Leipzig in 1736. Afterwards, Errard, who was engineer to Henry the Great of France; Stevinus, engineer to the prince of Orange; with Marolois, the chevalier de Ville, Lorini, Coehorn, the count de Pagan, and the marshal de Vauban; which last two noble authors have contributed greatly to the perfection of the art; besides Scheiter, Mallet, Belidor, Blondel, Muller, Montalambert, &c. Also a list of several works on the art of Fortification may be added, as follows: viz,

Melder's



Melder's Praxis Fortificatoria: Les Fortifications du Comte de Pagan: L'Ingénieur Parfait du Sieur de Ville: Sturmy's Architectura Militaris Hypothetical.: Blondel's Nouvelle Maniere de Fortifier les Places: The Abbé de Fay's Veritable Maniere de Bien Fortifier: Vauban's Ingénieur François: Coehorn's Nouvelle Fortification tant pour un Terrain bas & humide, que sec & élevé: Alexander de Grotte's Fortification: Donatus Roselli's Fortification: Medrano's Ingénieur François: The chevalier de St. Julien's Architecture Militaire: Lansberg's Nouvelle Maniere de Fortifier les Places: An anonymous treatise in French, called Nouvelle Maniere de Fortifier les Places, tirée des Methodes du chevalier de Ville, &c: Ozanam's Traité de Fortification: Memoires de l'Artillerie de Surirey de St. Remy: Muller's treatises of Elementary and Practical Fortification: and Montalambert's Fortification Perpendiculaire.

*Maxims in FORTIFICATION.* From the nature and circumstances of this art, certain general rules, or maxims, have been drawn, and laid down. These may indeed be multiplied to any extent, but the principal of them, are the following: viz,

1. That the manner of fortifying should be accommodated to that of attacking. So that no one manner can be assured always to hold, unless it be assured that the manner of besieging is incapable of being altered. Also, to judge of the perfection of a Fortification, the method of besieging at the time when it was built must be considered.

2. All the parts of a Fortification should be equally strong on all sides, where there is equal danger; and they should be able to resist the most powerful machines used in besieging.

3. A Fortification should be so contrived, as to be defended with the fewest men possible: which consideration, when well attended to, saves a great deal of expence.

4. That the defendants may be in the better condition, they must not be exposed to the enemies' artillery; but the aggressors must be exposed to theirs. Hence,

5. All the parts of a Fortification should be so disposed, as that they may defend each other. In order to this, every part ought to be flanked, i. e. seen sideways, capable of being seen and defended from some other part; so that there be no place where an enemy can lodge himself, either unseen, or under shelter.

6. All the campaign around must lie open to the defendants; so that no hill or eminence must be allowed, behind which the enemy might shelter himself from the guns of the Fortification; or from which he might annoy them with his own. Hence, the fortress is to command all the place round about; and consequently the outworks must all be lower than the body of the place.

7. No line of defence must exceed the point-blank musket-shot, which is from 120 to 150 fathoms.

8. The more acute the angle at the centre, the stronger is the place; as consisting of the more sides, and consequently more defensible.

9. All the defences should be as nearly direct as possible.

10. The works that are most remote from the centre

of the place, ought always to be open to those that are more near.

These are the general laws and views of Fortification. As to the particular ones, or such as respect the several members or parts of the work, they are given under those articles respectively.

Fortification is either theoretical or practical.

*Theoretical FORTIFICATION*, consists in tracing the plans and profiles of a work on paper, with scales and compasses; and in examining the systems proposed by different authors, to discover their advantages and defects. And

*Practical FORTIFICATION*, consists in forming a project of a work according to the nature of the ground, and other necessary circumstances, tracing it on the ground, and executing the project, together with all the military buildings, such as magazines, storehouses, bridges, &c.

Again, Fortification is either Defensive or Offensive.

*Defensive FORTIFICATION* is the art of defending a town that is besieged, with all the advantages the Fortification of it will admit. And

*Offensive FORTIFICATION* is the same with the attack of a place, being the art of making and conducting all the different works in a siege, in order to gain possession of the place.

FORTIFICATION is also used for the place fortified; or the several works raised to defend and flank it, and keep off the enemy.

All Fortifications consist of lines and angles, which have names according to their various offices. The principal lines are those of circumvallation, of contravallation, of the capital, &c. The principal angles are those of the centre, the flanking angle, flanked angle, angle of the epaule, &c.

Fortifications are either durable or temporary.

*Durable FORTIFICATION*, is that which is built and intended to remain a long time. Such are the usual Fortifications of cities, frontier places, &c. And a

*Temporary FORTIFICATION*, is that which is erected on some emergent occasion, and only for a short time. Such are field-works, thrown up for the seizing and maintaining a post, or passage; those about camps, or in sieges; as circumvallations, contravallations, redoubts, trenches, batteries, &c.

Again, Fortifications are either regular or irregular. A

*Regular FORTIFICATION*, is that in which the bastions are all equal; or which is built in a regular polygon, the sides and angles of which are usually about a musket shot from each other. A Regular Fortification, having the parts all equal, has the advantage of being equally defensible; so that there are no weak places. And an

*Irregular FORTIFICATION*, is that in which the bastions are unequal, and unlike; or the sides and angles not all equal, and equidistant.

In an Irregular Fortification, the defence and strength being unequal, it is necessary to reduce the irregular shape of the ground, as near as may be, to a regular figure: i. e. by inscribing it in an oval, instead of a circle; so that one half may be similar and equal to the other half.

*Marine*



*Marine Fortification*, is sometimes used for the art of raising works on the sea coast &c, to defend harbours against the attacks of shipping.—See a neat treatise on Marine Fortification, at the end of Robertson's Elements of Navigation.

There are many modes of Fortification that have been much esteemed and used; a small specimen of a comparative view of the principal of these, is represented in plate x, viz, those of Count Pagan, and Mess. Vauban, Coehorn, Belidor, and Blondel; the explanation of which is as follows:

#### 1. Pagan's System.

- A Half Bastions.
- B Ravelin and Counterguard.
- C Counterguards before the bastions.
- D The Ditch.
- E The Glacis.
- G The place of Arms.
- H Retired Flanks.
- a Line of Defence.

#### 2. Vauban's System.

- b Angle of the Bastion, or Flanked angle.
- c Angle of the Shoulder.
- d Angle of the Flank.
- e Salient Angle.
- f Face of the bastion.
- g The Flank.
- h The Curtain.
- i Tenailles.
- k Traverses in the covert way.

#### 3. Coehorn's System.

- 1 Concave Flanks.
- 2 The Curtains.
- 3 Redoubts in the re-entering angles.
- 4 Traverses.
- 5 Stone lodgments.
- 6 Round Flanks.
- 7 Redoubt.
- 8 Coffers planked on the sides, and above covered overhead with a foot of earth.

#### 4. Belidor's System.

- I Cavaliers.
- K Rams-horns, or Tenailles.
- L Retrenchments within the detached bastions.
- M Circular Curtain.
- N The Ravelin.
- P Lunettes with retired batteries.
- Q Redoubt.
- R Detached Redoubt.
- S An Arrow.
- P Small Traverses.

#### 5. Blondel's System.

- I Retired Battery.
- m Lunettes.
- n Ravelin, with Retired Bastion.
- o Orillons.

Another, or new method of Fortification has lately been proposed by M. Montalambert, called *Fortification Perpendiculaire*, because the faces of the works are

made by a series of lines running zigzag perpendicular to one another.

*Profile of a Fortification*, is a representation of a vertical section of a work; serving to shew those dimensions which cannot be represented in plans, and are necessary in the building of a Fortification. The names and dimensions of the principal parts are as follow (see fig. 8, pl. vii), where the numbers or dimensions are all expressed in feet.

AB The level of the ground plane,

AC = 27,

CD = 18, and CW =  $16\frac{1}{2}$ ; also DN is parallel to AB.

DE = 30,

EF = 2, FG = 3, GH = 3, HI =  $4\frac{1}{2}$ , IL =  $1\frac{1}{2}$ ,

LK = 18, KM =  $2\frac{1}{2}$ , NP = 36, NO = 5, PR = 7,

RS = 1, ST = 12 or 18, OV = 9, Pn = 120,

mz = 3, nu = 3, mc = 30, cd = 2,

de = 3, ef = 3, fl =  $4\frac{1}{2}$ , rg = 120, lb = 1.

AW the interior talus or slope of the rampart,

WE, or DE the terre-plein of ditto,

FG the talus, and GH the upper part, of the banquette,

HL the interior side of the parapet,

LM the upper part of ditto,

N the cordon of 1 foot radius,

NP the depth, and Pn the breadth of the ditch,

OQ interior side of revetement,

NR the scarp or exterior side of ditto,

ST the depth of the foundation,

YZ revetement of the parapet,

nu the counterescarp,

mc the covert-way,

ce talus of the banquette,

ef the upper part of ditto,

fb parapet of the covert-way,

hg the glacis.

Other sections are at fig. 2, pl. 12.

*FORTIFIED Place*, a Fortrefs, or Fortification, i. e. a place well flanked, and sheltered with works.

*FORTIN*, or *Fortlet*, a diminutive of the word Fort, meaning a little fort, or sconce, called also Field Fort.

*Star FORTIN*, is that whose sides flank each other, &c.

*FORTRESS*, the same as Fort, or a Fortification.

*FOSTER (SAMUEL)*, an English mathematician, and astronomy professor of Gresham-college, was born in Northamptonshire, and admitted a Sizer at Emanuel-college Cambridge in 1616. He took the degree of bachelor of arts in 1619, and of master in 1623. He applied early to the mathematics, and attained a great proficiency in it, of which he gave the first specimen in 1624, in a treatise on *The Use of the Quadrant*.

On the death of Mr. Gellibrand, he was chosen to succeed him, in 1636, as astronomy professor in Gresham-college, London. He quitted it again however the same year, though for what reason does not appear, and was succeeded by Mr. Mungo Murray, professor of philosophy at St. Andrews in Scotland. But this gentleman marrying, the professorship again became vacant, and Mr. Foster was re-elected in 1641.

Mr. Foster was one of those gentlemen who held private meetings for cultivating philosophy and useful knowledge,



Knowledge, which afterwards gave rise to the Royal Society. In 1646, Dr. Wallis, who was one of those associating gentlemen, received from Mr. Foster a theorem *de triangulo sphaerico*, which he published in his *Mechanics*. Neither was it only in this branch of science that he excelled, but he was likewise well versed in the ancient languages; as appears from his revising and correcting the *Lemmata* of Archimedes, which had been translated into Latin from an Arabic manuscript by Mr. John Greaves. He made also several curious observations upon eclipses of the sun and moon, in various places; and was particularly noted for inventing, as well as improving, astronomical and other mathematical instruments. After a long declining state of health, he died at Gresham-college in 1652.

His printed works are as follow; of which the first two articles were published before his death, and the rest of them after it.

1. *The Description and Use of a small Portable Quadrant, for the easy finding the hour of Azimuth*; 4to, 1624. Originally published at the end of Gunter's Description of the Cross-Staffe, as an appendix to it.
2. *The Art of Dialling*; 4to, 1638. Reprinted, with additions, in 1675.
3. *Posthuma Fosteri*; by Wingate; 4to, 1652.
4. *Four Treatises of Dialling*; 4to, 1654.
5. *Miscellanies, or Mathematical Lucubrations*. Published by John Twysden, with additions of his own; and an appendix by Leybourne; folio, 1659.
6. *The Sector altered, and other Scales added, &c.* Published by Leybourne in 1661, in 4to.

There have been two other persons of the same name, who have published some mathematical pieces. The first was,

WILLIAM FOSTER, who was a disciple of Mr. Oughtred, and afterward a teacher of the Mathematics in London. He distinguished himself by a book, which he dedicated to Sir Kenelm Digby, entitled, *The Circles of Proportion, and the Horizontal Instrument, &c*; 4to, 1633.—The other was

MARK FOSTER, who lived later in point of time than either of the other two, and published a treatise entitled, *Arithmetical Trigonometry: being the solution of all the usual Cases in Plain Trigonometry by Common Arithmetic, without any tables whatsoever*. 12mo, 1690.

FOUNDATION, that part of a building which is underground: or the mass which supports a building, and upon which it stands: or it is the coffer or bed dug below the level of the ground, to raise a building upon.

FOUNTAIN, in Philosophy, a spring or source of water rising out of the Ground. See SPRING.

FOUNTAIN, or Artificial FOUNTAIN, in Hydraulics, a machine or contrivance by which water is violently spouted or darted up; called also a Jet d'eau.

There are various kinds of Artificial Fountains, but all formed by a pressure of one sort or another upon the water &c, viz, either the pressure or weight of a head of water, or the pressure arising from the spring and elasticity of the air, &c. When these are formed by the pressure of a head of water, or any other fluid of the same kind with the Fountain, or jet, then will this spout up nearly to the same height as that head, abating only a little for the resistance of the air, with that of

the adjutage &c, in the fluid's rushing through; but, when the Fountain is produced by any other force than the pressure of a column of the same fluid with itself, it will rise to such a height as may be nearly equal to the altitude of a column of the same fluid whose pressure is equal to the given force that produces the fountain.

*To Construct an Artificial Fountain, playing by the pressure of the water.* This is to be effected by making a close connection between a head or elevated piece of water, and the lower place, where the Fountain is to play; which may be done in this manner: Having a head of water, naturally, or, for want of such, make an artificial one, raising the water by pumps, or other machinery: from this head convey the water in close pipes, in any direction, down to the place where the Fountain is to play; and there let it issue through an adjutage, or small hole, turned upwards, by which means it will spout up nearly as high as the head of the water comes from, as above mentioned.

*To Construct an Artificial Fountain, playing by the spring or elasticity of the air.* A vessel proper for a reservoir, as AB, fig. 4, plate viii, is provided either of metal, or glass, or the like, ending in a small neck c, at the top: through this neck is put a tube cd, till the lower end come near the bottom of the vessel, this being about half full of water. The neck is so contrived, as that a syringe, or condensing pipe, may be screwed upon the tube; by means of which a large quantity of air may be intruded through the tube into the water, out of which it will disengage itself, and emerge into the vacant part of the vessel, and lie over the surface of the water CD.

Now, the water in the vessel being thus pressed by the air, which is, for ex. double the density of the external air; and the elastic force of air being proportional to its density, or to its gravitating force, the effect will be the same, as if the weight of the column of air, over the surface of the water, were double that of the column pressing in the tube; so that the water must be forced to spout up through, when the syringe is removed, with a force equal to the excess of pressure of the included air, above that of the external, that is, in this instance, with a force equal to the pressure of an entire column of the atmosphere; which being equal to the pressure of a column of 33 or 34 feet of water, it follows that the Fountain will play to nearly 33 or 34 feet high.

These aerial or aquatic Fountains may be applied in different ways, so as to exhibit various appearances; and from these alone arises the greatest part of our artificial water-works: even the engine for extinguishing fire, is a Fountain playing by the force of confined air.

*A Fountain spouting the water in various directions.* Suppose AB the vertical tube, or spout, in which the water rises, (fig. 5, pl. viii): into this let several other tubes be fitted; some horizontal, others oblique, or inclining, or reclining, &c, as at E, H, L, N, P, &c. Then, as all water retains the direction of the aperture through which it comes, that issuing through A will rise perpendicularly; and the rest will tend different ways, describing arches of different magnitudes.

Or thus: Suppose the vertical tube AB (fig. 6) through which the water rises, to be stopped at the top,



top, as in A; and, instead of pipes or cocks, let it be only perforated with little holes all around or only half round its surface: then will the water spin out, in all directions, through the little holes, to different distances.

And hence, if the tube AB be about the height of a man, and having a turn-cock at C; upon opening this cock, the spectators will be sprinkled unexpectedly with a shower.

*FOUNTAIN playing by drawing the breath.* Suppose AB (fig. 8, plate vi), a globe of glass, or metal, in which is fitted a tube CD, having a small orifice in C, and reaching almost to D, the bottom of the globe. Then if the air be sucked, or drawn with the mouth, out of the tube CD, and the orifice C be immediately immersed under cold water, the water will ascend through the tube into the sphere. Thus proceeding, by repeated exsuctions, till the vessel be above half full of water; then applying the mouth to C, and blowing air into the tube, upon removing the mouth, the water will spout forth.

Or, if the globe be put into hot water, the air being thus rarefied, will make the water spout as before.

And this kind of Fountain is called Pila Heronis, or Hero's Ball, from the name of its inventor.

*FOUNTAIN, whose stream raises and plays a brass ball.* Provide a hollow brass ball A (fig. 9, pl. vi), made very thin, that its weight may not be too great for the force of the water; and let the tube BC, through which the water rises, be exactly perpendicular to the horizon. Then the ball, being laid in the bottom of the cup or basin B, will be taken up in the stream, and sustained at a considerable height, playing a little up and down.

*FOUNTAIN which spouts water in form of a shower.* To the tube in which the water is to rise, fit a spherical, or lenticular head AB (fig. 1, pl. xi) made of a plate of metal, and perforated at the top with a great number of little holes. The water rising violently towards AB, will be there divided into innumerable little threads, and afterwards broken, and dispersed into the finest drops.

*FOUNTAIN which spreads the water in form of a tablecloth.* To the tube AB (fig. 2, pl. xi) solder two spherical segments, C and D, almost touching each other, with a screw E, to contract or amplify the interstice or chink, at pleasure. Some choose to make a smooth and even cleft, in a spherical or lenticular head, fitted upon the tube. The water spouting through this chink or cleft will expand itself like a cloth.—And thus, the fountain may be made to spout out in the figure of men, or other animals.

*FOUNTAIN, which, when it has done spouting, may be turned like an hour-glass.* Provide two glasses, A and B, (fig. 3, pl. xi) to be so much the larger as the fountain is to play the longer, and placed at so much the greater distance from each other as the water is desired to spout the higher. Let CDE be a crooked tube, furnished in E with a jet; and GHI another bent tube, furnished with a jet in I: GF and KL are to be other lesser tubes, open at both ends, and reaching near the

bottom of the vessels A and B, to which the tubes CD and GH are likewise to reach.

Now if the vessel A be filled with water, it will descend through the tube CD; and it will spout up through the jet E, by the pressure of the column of water CD. But unless the pipe GF were open at G, to let the air run up to F, and press at the top of the surface of the water in the cavity A, the water could not run down and spout at E. After its fall again, it will sink through the little tube KL, into the vessel B, and expel the air through the tube GI. At length, when all the water is emptied out of the vessel A, by turning the machine upside down, the vessel B will be the reservoir, and make the water spout up through the jet I, the pipe KL supplying B with air to let the water descend in the direction GHI.

Hence, if the vessels A and B contain just as much water as will be spouted up in an hour's time, we shall have a spouting clepsydra, or water-clock; which may be graduated or divided into quarters, minutes, &c.

*FOUNTAIN of Command.* This depends on the same principles with those of the former: CAE (fig. 4, pl. xi) is a vessel of water secured against the entrance of the air, except through the pipe GF, when the cock C, by which it is filled, is shut. There is another pipe EDHB, going from the bottom of the water to the jet B in the basin DB; but this is stopped by the cock H. At the lowest part of the basin DB, there is a small hole at I, to let the water of the basin DB run into the basin GH under it; there is also a small triangular hole or notch, in the bottom of the pipe FG, at G. Turn the cock H, and the fountain will play for some time, then stop, then play again alternately for several times together. When those times of playing and stopping are known before hand, you may command the Fountain to play or stop; whence its name. The cause of this phenomenon is as follows: the water coming down the pipe EDHB, would not come out at B, if the air Ss, above the water, were not supplied as it dilated: now it is supplied by the pipe GF, which takes it in at the notch G, and delivers it out at F; but after some time the water, which was spouted out at B, falling down into the basin DB, rises high enough to come above the notch G, which stops the passage of the air; so that the air Ss, above the water in the vessel CAE, wanting a supply, cannot sufficiently press, and the Fountain ceases playing. But when the water of DB has run down into the lower basin GH, through the hole I, till it falls below the top of the notch G, the air runs up into the upper receptacle, and supplies that at Ss, and the Fountain plays again. This is seen a little before hand, by a skin of water on the notch G, before the air finds a passage, and then you may command the Fountain to play. It is evident that the hole I must be less than the hole of the jet, or else all the water would run out into the lower basin, without rising high enough to stop the notch G.

*FOUNTAIN that begins to play upon the lighting of candles, and ceases as they go out.* Provide two cylindrical vessels, AB and CD, (fig. 7, pl. xi.) connect them by tubes, open at both ends, KL, BE, &c, so that the air may descend out of the higher into the lower:



to the tubes folder candlesticks, H, &c, and to the hollow cover of the lower vessel CE, fit a little tube or jet, FG, furnished with a cock G, and reaching almost to the bottom of the vessel. In G let there be an aperture, furnished with a screw, by which water may be poured into CD. Then, upon lighting the candles H, &c, the air in the contiguous tubes becoming rarefied by the heat, the water will begin to spout through GF.

By the same contrivance may a statue be made to shed tears upon the presence of the sun, or on the lighting of a candle, &c: all that is here required, being only to lay tubes from the cavity where the air is rarefied, to some other cavities placed near the eyes, full of water.

A FOUNTAIN by the Rarefaction of the air, may be made in the following manner: Let AB and CD, fig. 5, pl. xi, be two pipes fixed to a brass head C, made to screw into a glass vessel E, which having a little water in it, is inverted till the pipes are screwed on; then reverting it suddenly, so as to put A, the lower end of the spouting pipe AB, into a jar of water A, and the lower end of the descending pipe CD, into a receiving vessel D, the water will spout up from the jar A into the tall glass vessel E, from which it will go down at the mouth C, through CD, into the vessel D, till the water is wholly emptied out of A, making a Fountain in E, into D. The reason of the play of the Fountain is this: the pipe CD, being 2 feet 9 inches long, lets down a column of water, which rarefies the air 1-12th part in the vessel E, where it presses against the water spouting at B with 1-12th of the force by which the water is pushed up at the hole A, by the pressure of the common air on the water in the vessel A; so that the water spouts up into E, when the air is rarefied 1-12th, with the difference of the pressure of the atmosphere, and the forementioned rarefied air; i. e. of 33 to  $2\frac{1}{4}$ , or of 12 to 1. This would raise the water 2 feet 9 inches; but the length of the pipe A, of 9 inches, being deducted, the jet will only rise 2 feet. This, says Desaguliers, may be called a syphon Fountain, where AB is the driving leg, and CD the issuing leg.

FOUNTAIN of Hero of Alexandria, so called, because it was contrived by him. In the second Fountain above described, the air is compressed by a syringe; in this, (see fig. 6, pl. xi) the air, being only compressed by the concealed fall of water, makes a jet, which, after some continuance, is considered by the ignorant as a perpetual motion; because they imagine that the same water which fell from the jet rises again. The boxes CE and DYX, being close, we see only the basin ABW, with a hole at W, into which the water spouting at B falls; but that water does not come up again; for it runs down through the pipe WX into the box DYX, from whence it drives out the air, through the ascending pipe YZ, into the cavity of the box CE, where, pressing upon the water that is in it, it forces it out through the spouting pipe OB, as long as there is any water in CE; so that this whole play is only whilst the water contained in CE, having spouted out, falls down through the pipe WX, or of the boxes CE and DY above one another: the height of the water, measured from the basin ABW to the surface of the water in the lower box DYX, is always equal to the

VOL. I.

height measured from the top of the jet to the surface of the water in the middle cavity at CE. Now, since the surface CE is always falling, and the water in DY always rising, the height of the jet must continually decrease, till it is shorter by the depth of the cavity CE, which is emptying, added to the depth of the cavity DY, which is always filling; and when the jet is fallen so low, it immediately ceases. The air is represented by the points in this figure.

To prepare this Fountain for playing, which should be done unobserved, pour in water at W, till the cavity DXY is filled; then invert the Fountain, and the water will run from the cavity DXY into the cavity CE, which may be known to be full, when the water runs out at B held down. Set the Fountain up again, and, to make it play, pour in about a pint of water into the basin ABW; and as soon as it has filled the pipe WX it will begin to play, and continue as long as there is any water in CE. You may then pour back the water left in the basin ABW, into any vessel, and invert the Fountain, which, being set upright again, will be made to play, by putting back the water poured out into ABW; and so on as often as you please.

Spouting FOUNTAIN, or Jet d'Eau, is any Fountain whose water is darted forth impetuously through jets, or ajutages, and returns in form of rains, nets, folds, or the like.

FOUNTAIN-Pen, is a pen contrived to contain a quantity of ink, and let it flow very gently, so as to supply the writer a long time without the necessity of taking fresh ink.

The Fountain-pen, represented fig. 8, pl. xi, consists of divers pieces of metal, F, G, H, the middle piece F carrying the pen, which is screwed into the inside of a little pipe; and this again is soldered into another pipe of the same size as the lid G; in which lid is soldered a male screw, for screwing on the cover; as also for stopping a little hole at the place, and hindering the ink from passing through it: at the other end of the piece F is a little pipe, on the outside of which may be screwed the top cover H. A porte-craion goes in the cover, to be screwed into the last-mentioned pipe, to stop the end of the pipe into which the ink is to be poured by a funnel.

To use the Pen, the cover G must be taken off, and the pen a little shaken, to make the ink run more freely.

FOURTH, in Music, one of the harmonic intervals, or concords. It consists in the mixture of two sounds, which are in the ratio of 4 to 3; i. e. of two sounds produced by chords, whose lengths are to each other as 4 to 3.

FRACTION, or Broken Number, in Arithmetic and Algebra, is a part, or some parts, of another number or quantity considered as a whole, but divided into a certain number of parts; as  $\frac{3}{4}$ ths, which denotes 3 parts out of 4, of any quantity.

Fractions are usually divided into Vulgar, Decimal, Duodecimal, and Sexagesimal. For the last three sorts, see the respective words.

Vulgar FRACTIONS, called also simple Fractions, are usually denoted by two numbers, the one set under the other, with a small line between them: thus  $\frac{3}{4}$  denotes the Fraction three-fourths, or 3 parts out of 4, of some whole



whole quantity considered as divided into 4 equal parts.

The lower number 4, is called the Denominator of the Fraction, shewing into how many parts the whole or integer is divided; and the upper number 3, is called the Numerator, and shews how many of those equal parts are contained in the Fraction. Hence it follows, that as the numerator is to the denominator, so is the Fraction itself, to the whole of which it is a Fraction; or as the denominator is to the numerator, so is the whole or integer, to the Fraction: thus, the integer being denoted by 1, as  $4 : 3 :: 1 : \frac{3}{4}$  the Fraction.— And hence there may be innumerable Fractions all of the same value, as there may be innumerable quantities all in the same ratio, viz, of 4 to 3; such as 8 to 6, or 12 to 9, &c. So that if the two terms of any Fraction i. e. the numerator and denominator, be either both multiplied or both divided by any number, the resulting Fraction will still be of the same value: thus,  $\frac{3}{4}$  or  $\frac{6}{8}$  or  $\frac{9}{12}$  or  $\frac{12}{16}$  &c, are all of the same value with each other.

Fractional expressions are usually distinguished into Proper and Improper, Simple and Compound, and Mixt Numbers.

A *Proper* FRACTION, is that whose numerator is less than the denominator; and consequently the Fraction is less than the whole or integer; as  $\frac{3}{4}$ .

*Improper* FRACTION, is when the numerator is either equal to, or greater than, the denominator; and consequently the Fraction either equal to, or greater than, the whole integer, as  $\frac{4}{4}$ , which is equal to the whole; or  $\frac{5}{4}$ , which is greater than the whole.

*Simple* FRACTIONS, or *Single* FRACTIONS, are such as consist of only one numerator, and one denominator; as  $\frac{3}{4}$ , or  $\frac{5}{4}$ , or  $\frac{12}{5}$ .

*Compound* FRACTIONS are Fractions of Fractions, and consist of several Fractions, connected together by the word *of*: as  $\frac{2}{3}$  of  $\frac{3}{4}$ , or  $\frac{1}{2}$  of  $\frac{2}{3}$  of  $\frac{3}{4}$ .

A *Mixt Number* consists of an integer and a Fraction joined together: as  $1\frac{3}{4}$ , or  $12\frac{2}{3}$ .

The arithmetic of Fractions consists in the Reduction, Addition, Subtraction, Multiplication, and Division of them.

*Reduction of* FRACTIONS is of several sorts; as 1. *To reduce a given whole number into a Fraction of any given denominator.* Multiply the given integer by the proposed denominator, and the product will be the numerator. Thus, it is found that  $3 = \frac{6}{2}$ , and  $5 = \frac{20}{4}$ , or  $7 = \frac{35}{5}$ .

If no denominator be given, or it be only proposed to express the integer Fraction-wise, or like a Fraction; set 1 beneath it, for its denominator. So  $3 = \frac{3}{1}$ , and  $5 = \frac{5}{1}$ , and  $7 = \frac{7}{1}$ .

2. *To reduce a given Fraction to another Fraction equal to it, that shall have a given denominator.* Multiply the numerator by the proposed denominator, and divide the product by the former denominator, then the quotient set over the proposed denominator will form the Fraction required. Thus, if it be proposed to reduce  $\frac{3}{4}$  to an equal Fraction whose denominator shall be 8; then  $3 \times 8 = 24$ , and  $24 \div 4 = 6$  the numerator, so that  $\frac{6}{8}$  is the Fraction sought, being  $= \frac{3}{4}$ , and having 8 for its denominator.

3. *To Abbreviate, or reduce Fractions to lower terms.*

Divide their terms, i. e. numerator and denominator, by any number that will divide them both without a remainder, so shall the quotients be the corresponding terms of a new Fraction, equal to the former, but in smaller numbers. In like manner abbreviate these new terms again, and so on till there be no number greater than 1 that will divide them without a remainder, and then the Fraction is said to be in its least terms. Thus, to abbreviate  $\frac{15}{20}$ ; first divide both terms by 5, and the Fraction becomes  $\frac{3}{4}$ ; next divide these by 3, and it becomes  $\frac{1}{4}$ : so that  $\frac{15}{20} = \frac{3}{4} = \frac{1}{4}$ , which is in its least terms.

4. *To reduce Fractions to other equivalent ones of the same denominator.* Multiply each numerator, separately taken, by all the denominators except its own, and the products will be the new numerators; then multiply all the denominators continually together, for the common denominator, to these numerators. Thus,  $\frac{2}{3}$  and  $\frac{4}{5}$  reduce to  $\frac{10}{15}$  and  $\frac{12}{15}$ ; and  $\frac{2}{3}$ ,  $\frac{3}{4}$ , and  $\frac{4}{5}$  reduce to  $\frac{40}{60}$ ,  $\frac{45}{60}$ , and  $\frac{48}{60}$ .

5. *To find the value of a Fraction, in the known parts of its integer.* Multiply always the numerator by the number of parts of the next inferior denomination, and divide the products by the denominator. So, to find the value of  $\frac{9}{16}$  of a pound sterling; multiply 9 by 20 for shillings, and dividing by 16, gives 11 for the shillings; then multiply the remainder 4 by 12 pence, and dividing by 16 gives 3 for pence: so that 11s. 3d. is the value of  $\frac{9}{16}$ l. as required.

$$\begin{array}{r} 9 \\ 20 \\ 16 \overline{) 180} \quad (11s \\ 16 \\ \hline 20 \\ 16 \\ \hline 4 \\ 12 \\ 16 \overline{) 48} \quad (3d \\ 48 \\ \hline \end{array}$$

6. *To reduce a mixt number to an equivalent improper Fraction.* Multiply the integer by the denominator, and to the product add the numerator, for the new numerator, to be set over the same denominator as before. Thus  $3\frac{5}{8}$  becomes  $\frac{29}{8}$ .

7. *To reduce an improper Fraction to its equivalent whole or mixt number.* Divide the numerator by the denominator; so shall the quotient be the integral part, and the remainder set over the denominator will form the fractional part of the equivalent mixt number. Thus  $\frac{29}{8}$  reduces to  $3\frac{5}{8}$ , and  $\frac{32}{4} = 8$ .

8. *To reduce a compound Fraction to a simple one.* Multiply all the numerators together for the numerator, and all the denominators together for the denominator, of the simple Fraction sought. Thus,  $\frac{1}{2}$  of  $\frac{3}{4} = \frac{3}{8}$ , and  $\frac{2}{3}$  of  $\frac{4}{5}$  of  $\frac{7}{9} = \frac{56}{135}$ .

To reduce a Vulgar Fraction to a decimal. See DECIMALS. And for several other particulars concerning Reduction, as well as the other operations in Fractions; see my Arithmetic.

*Addition of* FRACTIONS. First reduce the Fractions to their simplest form, and reduce them also to a common denominator, if their denominators are different; then add all the numerators together, and set the sum over the common denominator, for the sum of all the Fractions as required.

$$\text{Thus, } \frac{2}{5} + \frac{3}{5} = \frac{5}{5} = 1;$$

$$\text{And } \frac{2}{7} + \frac{3}{4} = \frac{8}{28} + \frac{21}{28} = \frac{29}{28} = 1\frac{1}{28}.$$

*Subtraction of* FRACTIONS. Reduce the Fractions the same as for addition; then subtract the one numerator from



from the other, and set the difference over the common denominator.

$$\text{So } \frac{7}{5} - \frac{3}{5} = \frac{4}{5};$$

$$\text{And } \frac{7}{9} - \frac{4}{7} = \frac{49}{63} - \frac{36}{63} = \frac{13}{63}.$$

*To Multiply FRACTIONS together.* Reduce them all to the form of simple Fractions, if they are not so; then multiply all the numerators together for the numerator, and all the denominators together for the denominator of the product sought.

$$\text{Thus } \frac{2}{5} \times \frac{4}{5} = \frac{8}{25};$$

$$\text{And } \frac{3}{8} \times \frac{5}{8} \times \frac{7}{10} = \frac{105}{640} = \frac{21}{128} = \frac{7}{32}.$$

*To Divide FRACTIONS.* Divide the numerator by the numerator, and the denominator by the denominator, if they will exactly divide. Thus,  $\frac{8}{15} \div \frac{4}{5} = \frac{2}{3}$ .

But if they will not divide without a remainder, then multiply the dividend by the reciprocal of the divisor, that is, by the Fraction obtained by inverting or changing its terms. Thus,  $\frac{8}{15} \div \frac{3}{4} = \frac{8}{15} \times \frac{4}{3} = \frac{32}{45}$ .

*Algebraic FRACTIONS, or FRACTIONS in Species,* are exactly similar to vulgar Fractions, in numbers, and all the operations are performed exactly in the same way; therefore the rules need not be repeated, and it may be sufficient here to set down a few examples to the foregoing rules. Thus,

$$1. \text{ The Fraction } \frac{aab}{bc} \text{ abbreviates to } \frac{aa}{c}.$$

$$2. \frac{6a^3 - 9ax^2}{6a^2 + 3ax} = \frac{2a^2 - 3x^2}{2a + x}, \text{ by dividing by } 3a.$$

$$3. \frac{a^3 - a^2x + ax^2 - x^3}{a^2 - ax} = \frac{a^2 + x^2}{a}, \text{ by dividing by } a - x. \text{ See COMMON MEASURE.}$$

$$4. \frac{a}{b} \text{ and } \frac{c}{d} \text{ become } \frac{ad}{bd} \text{ and } \frac{bc}{bd}, \text{ when reduced to a common denominator.}$$

$$5. \frac{a}{b} + \frac{c}{d} = \frac{ad + bc}{bd}.$$

$$6. \frac{a}{x} - \frac{b}{z} = \frac{az - bx}{xz}.$$

$$7. \frac{a}{b} \times \frac{c}{d} = \frac{ac}{bd}.$$

$$8. \frac{a}{x} \div \frac{b}{z} = \frac{a}{x} \times \frac{z}{b} = \frac{az}{bx}.$$

*Continued FRACTION,* is used for a Fraction whose denominator is an integer with a Fraction, which latter Fraction has for its denominator an integer and a Fraction, and the same for this last Fraction again, and so on, to any extent, whether supposed to be infinitely continued, or broken off after any number of terms. Euler, *Analys. Inf.* vol. 1, p. 295.

$$\text{As } \frac{1}{2} + \frac{1}{3}, \text{ or } \frac{1}{2} + \frac{1}{3} + \frac{1}{4},$$

$$\text{or } \frac{1}{2} + \frac{2}{3} + \frac{3}{4} + \frac{4}{5} + \&c.$$

Or, using letters instead of numbers,

$$\frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d} + \frac{1}{e} + \&c.$$

$$\text{or } \frac{A}{a} + \frac{B}{b} + \frac{C}{c} + \frac{D}{d} + \frac{E}{e} + \&c.$$

When these series are not far extended, it is not difficult to collect them by common arithmetic.

Lord Brouncker, it seems, was the first who considered Continued Fractions, or at least, who applied them to the quadrature of curves, in Wallis's *Arith. Infin.* prop. 191, vol. 1, p. 469 &c, where this author explains the manner of forming them, giving several numeral examples, in approximating ratios, as well as the general series

$$\frac{a}{a} \frac{b}{\beta} \frac{c}{\gamma} \frac{d}{\delta} \&c, \text{ as he denotes it.}$$

Huygens also used it for the like purpose, viz, to approximate the ratios of large numbers, in his *Descrip. Autom. Planet. in Oper. Relig.* p. 173 &c, edit. Amst. 1728. And a special treatise on Continued Fractions was given by Euler, in his *Analys. Infin.* vol. 1, pa. 295 &c.

This subject is perhaps capable of much improvement, though it has been rather neglected, as very little use has been made of it, except, by those authors, in approximating to the value of Fractions, and ratios, that are expressed in large numbers; besides a method of Goniometry by De Lagny, explained in the Introduction to my *Logarithms*, pa. 78; as also some use I have made of it in summing very slowly converging series, in my *Tracts*, p. 38 & seq.

As to the reducing of common Fractions, and ratios, that are expressed in large numbers, to Continued Fractions, it is no more than the common method of finding the greatest common measure of those two numbers, by dividing the greater by the less, and the last divisor always by the last remainder; for then the several quotients are the denominators of the Fractions, the numerators being always 1 or unity. Thus, to find

$$\text{approximating values of the Fraction } \frac{31415926535}{10000000000},$$

or to the ratio of 31415926535 to 10000000000, being the ratio of the circumference of a circle to its diameter, by means of a Continued Fraction; or, to change the said Common Fraction to a Continued Fraction: Dividing the greater term always by the less, the same as to find the greatest common measure of the said numbers or terms, the several quotients will be 3, 7, 15, 1, 292, 1, 1, &c, which, after the first, will be the denominators, to the common numerator 1; and therefore the said Fraction will be changed into this Continued Fraction,

$$3 + \frac{1}{7 + \frac{1}{15 + \frac{1}{1 + \frac{1}{292 + \frac{1}{1 + \frac{1}{1 + \frac{1}{\&c}}}}}}}$$

Hence, stopping at any part of these single Fractions, one after another, will give several values of the proposed ratio, all successively nearer and nearer the truth, but alternately too great and too little. So, stopping



at  $\frac{1}{7}$ , it is  $3\frac{1}{7} = \frac{22}{7} = 3.142857$  too great, or 22 to 7, the ratio of the circumference to the diameter as given by Archimedes. Again, stopping at  $\frac{1}{15}$ , it is

$$3\frac{1}{7\frac{1}{15}} = 3\frac{15}{106} = \frac{333}{106} = 3.141509 \text{ \&c, too little.}$$

But stopping at  $\frac{1}{1}$ ,

$$\text{it is } 3\frac{1}{7\frac{1}{15} + \frac{1}{1}} = 3\frac{1}{7\frac{1}{15}} = 3\frac{16}{113} = \frac{355}{113} \text{ (the ratio of}$$

Metius) = 3.1415929 &c, which is rather too great. And so on, always nearer and nearer, but alternately too great and too little.

And, in like manner is any algebraic Fraction thrown into a Continued Fraction. As the Fraction

$$\frac{a\beta\gamma\delta + a\beta d + ac\delta}{a\beta\gamma\delta + a\beta d + ac\delta + b\gamma\delta + bd}, \text{ which being in like}$$

manner divided, the quotients are  $\frac{a}{\alpha}, \frac{\beta}{b}, \frac{\gamma}{c}, \frac{\delta}{d}$ ;

which single Fractions being considered as denominators to other Fractions whose common numerator is 1, these will be the reciprocals of the former, and so will

become  $\frac{a}{\alpha}, \frac{b}{\beta}, \frac{c}{\gamma}, \frac{d}{\delta}$ ; and hence the proposed common Fraction is equal to this terminate Continued Fraction,

$$\frac{a}{\alpha} + \frac{b}{\beta} + \frac{c}{\gamma} + \frac{d}{\delta}.$$

On the other hand, any Continued Fraction being given, its equivalent common Fraction will be found, by beginning at the last denominator, or lowest end of the given Continued Fraction, and gradually collecting the Fractions backwards, till we arrive at the first, when the whole will thus be collected together into one common Fraction; as was done above in collecting the Fractions

$$\frac{1}{7} + \frac{1}{15} + \frac{1}{1} + \frac{1}{292} + \&c.$$

And in like manner the Continued Fraction

$$\frac{a}{\alpha} + \frac{b}{\beta} + \frac{c}{\gamma} + \frac{d}{\delta} \text{ collects into the Fraction}$$

$$\frac{a\beta\gamma\delta + a\beta d + ac\delta}{a\beta\gamma\delta + a\beta d + ac\delta + b\gamma\delta + bd}.$$

When the given Continued Fraction is an infinite one, collect it successively, first one term, then two together, three together, &c, till the sum is sufficiently exact. Or, if these collected sums converge too slowly to the true value, having collected a few of the terms into successive sums, these being alternately too great and too little, the true value will be found as near as you please by the method of arithmetical means, explained in my Tracts, vol. 1, Tract 2, pa. 11.

*Vanishing Fractions.* Such Fractions as have both their numerator and denominator vanish, or equal to 0, at the same time, may be called *Vanishing Fractions*. We are not to conclude that such Fractions are equal to nothing, or have no value; for that they have a certain determinate value, has been shewn by the best ma-

thematicians. The idea of such Fractions as these, first originated in a very severe contest among some French mathematicians, in which Varignon and Rolle were the two chief opposite combatants, concerning the then new or differential calculus, of which the latter gentleman was a strenuous opponent. Among other arguments against it, he proposed an example of drawing a tangent to certain curves at the point where the two parts cross each other; and as the fractional expression for the subtangent, by that method, had both its numerator and denominator equal to 0 at the point proposed, Rolle looked upon it as an absurd expression, and as an argument against the method of solution itself. The seeming mystery however was soon explained, and first of all by John Bernoulli. See an account of this affair in Montucla, Hist. Math. vol. 2, pa. 366.

Since that time, such kind of fractions have often been contemplated by mathematicians. As, by MacLaurin, in his Fluxions, vol. 2, pa. 698: Saunderson, in his Algebra, vol. 2, art. 469: De Moivre, in Miscel. Anal. pa. 165: Emerson, in his Algebra, pa. 212: and by many others. The same fractions have also proved a stumbling-block to more mathematicians than one, and the cause of more violent controversies: witness that between Powell and Waring, when they were competitors for the professorship at Cambridge. In the specimen of a work published on occasion of that competition, by Waring, was the fraction  $\frac{p-p^5}{1-p}$ ,

which he said became 4 when  $p = 1$ . This was struck at by Powell, as absurd, because when  $p = 1$ , then the fraction  $\frac{p-p^5}{1-p} = \frac{1-1}{1-1} = \frac{0}{0}$ , which was one chief cause of his not succeeding to the professorship.

Waring replied that  $\frac{p-p^5}{1-p}$  is  $= p + p^2 + p^3 + p^4$  (by common division)  $= 1 + 1 + 1 + 1 = 4$ , when  $p = 1$ . See the controversial pamphlets that passed between those two gentlemen at that time.

There are two modes of finding the value of such fractions, that have been given by the gentlemen above quoted. The one is by considering the terms of the fraction as two variable quantities, continually decreasing, till they both vanish together; or finding the ultimate value of the ratio denoted by the fraction. In this way of considering the matter, it appears that, as the terms of the fraction are supposed to decrease till they vanish, or become only equal to their fluxions or their increments, the value of the fraction at that state, will be equal to the fluxion or increment of the numerator divided by that of the denominator. Hence then,

taking the example  $\frac{x-x^5}{1-x}$  when  $x = 1$ ; the fluxion of the numerator is  $x - 5x^4x$ , and of the denominator  $-x$ ; therefore

$$\frac{x-5x^4x}{-x} = \frac{1-5x^4}{-1} = \frac{5x^4-1}{1} = 5x^4-1 = 5-1 = 4,$$

the value of the fraction  $\frac{x-x^5}{1-x}$  when  $x = 1$ .—Or,

$$\text{thus, because } x = 1, \text{ therefore } \frac{x-x^5}{1-x} = \frac{1-x^4}{1-x};$$

then the fluxion of the numerator,  $-4x^3x$ , divided by the



the fluxion of the denominator, or  $-x$ , gives  $4x^3$  or 4, the same as before.

The other method is by reducing the given expression to another, or simple form, and then substituting the values of the letters. So in the above example

$\frac{x-x^5}{1-x}$ , or  $\frac{1-x^4}{1-x}$ , when  $x=1$ ; divide the numerator

by the denominator, and it becomes  $1+x+x^2+x^3$ , which when  $x=1$ , becomes 4, for the given fraction, the same as before.—Again, to find the value of

$\frac{a\sqrt{ax-x^2}}{a-\sqrt{ax}}$  when  $x=a$ , in which case both the numerator and denominator become  $=0$ . Divide the numerator by the denominator, and the quotient is

$\sqrt{ax}+x+x\sqrt{\frac{x}{a}}$ ; which when  $x=a$ , becomes

$a+a+a=3a$ , for the value of the fraction in that state of it.

**FRAISE**, in Fortification, a kind of defence, consisting of pointed stakes, driven almost parallel to the horizon into the retrenchments of a camp, &c, to ward off and prevent any approach or scalade.

**FRANKLIN (DR. BENJAMIN)**, one of the most celebrated philosophers and politicians of the 18th century, was born in Bolton in North America in the year 1706, being the youngest of 13 children. His father was a tallow-chandler in Boston, and young Franklin was taken from school at 10 years of age, to assist him in his business. In this situation he continued two years; but disliking that occupation, he was bound apprentice to an elder brother, who was then a printer in Boston, but had learned that business in London, and who in the year 1721 began to print a newspaper, being the second ever published in America; the copies of which our author was sent to distribute, after having assisted in composing and printing it. Upon this occasion, our young philosopher enjoyed the secret and singular pleasure of being the much admired author of many essays in this paper; a circumstance which he had the address to keep a secret even from his brother himself; and this when he was only 15 years of age. The frequent ill usage from his brother, induced young Franklin to quit his service, which he did, at the age of 17, and went to New York. But not meeting employment here, he went forward to Philadelphia, where he worked with a printer a short time; after which, at the instance of Sir William Keith, governor of the province, he returned to Boston to solicit pecuniary assistance from his father to set up a printing-house for himself at Philadelphia, upon the promise of great encouragement from Sir William, &c. His father however thought fit to refuse such aid, alleging that he was yet too young (18 years old) to be entrusted with such a concern; so our author again returned to Philadelphia without it. Upon this, Sir William said he would advance the sum that might be necessary, and our young philosopher should go to England, and purchase all the types and materials, for which purpose he would give him letters of credit. He could never however get these letters, yet by dint of fair promises of their being sent on board the ship after him, he sailed for England, expecting these letters of credit were in the governor's packet, which he was to receive upon its

being opened. In this however he was cruelly deceived, and thus he was sent to London without either money, friends, or credit, at 18 years of age.

He soon found employment, however, as a journeyman printer, first at a Mr. Palmer's, and afterward with Mr. Watts, with whom he worked a considerable time, and by whom he was greatly esteemed, being also treated with such kindness, that it was always most gratefully remembered by our philosopher.

After a stay of 18 months in London, he returned to Philadelphia, viz in 1726, along with a merchant of that town, as his clerk, on a salary of 50 pounds a year. But his master dying the year after, he again engaged to direct the printing business of the same person with whom he had worked before. After continuing with him the best part of a year, our philosopher, in partnership with another young man, at length set up a printing-house himself.

A little before this time, young Franklin had gradually associated a number of persons, like himself, of a rational and philosophical turn of mind, and formed them into a club or society, to hold meetings, to converse and communicate their sentiments together, for their mutual improvement in all kinds of useful knowledge, which was in high repute for many years after. Among many other useful regulations, they agreed to bring such books as they had into one place, to form a common library. This resource being found defective, at Franklin's persuasion they resolved to contribute a small sum monthly towards the purchase of books for their use from London. Thus their stock began to increase rapidly; and the inhabitants of Philadelphia, being desirous of having a share in their literary knowledge, proposed that the books should be lent out on paying a small sum for the indulgence. Thus in a few years the society became rich, and possessed more books than were perhaps to be found in all the other colonies: the collection was advanced into a public library; and the other colonies, sensible of its advantages, began to form similar plans; from whence originated the libraries at Boston, New York, Charlestown, &c; that of Philadelphia being now not inferior to any in Europe.

About 1728 or 1729, young Franklin set up a newspaper, the second in Philadelphia, which proved very profitable, and otherwise useful, as affording an opportunity of making himself known as a political writer, by inserting several of his writings of that kind into it. In addition to his printing-house, he set up a shop to sell books and stationery; and in 1730 he married his wife, who proved very useful in assisting to manage the shop, &c. He afterward began to have some leisure, both for reading books, and writing them, of which he gave many specimens from time to time. In 1732 he began to publish *Poor Richard's Almanac*, which was continued for many years. It was always remarkable for the numerous and valuable concise maxims which it contained, for the œconomy of human life, all tending to exhort to industry and frugality: and in the almanac for the last year, all the maxims were collected in an address to the reader, entitled, *The Way to Wealth*. This has been translated into various languages, and inserted in different publications. It has also been printed on a large sheet, proper to be framed, and hung up in conspicuous places in all houses, as it

very



very well deserves to be. Mr. Franklin became gradually more known for his political talents, and in the year 1736, he was appointed clerk to the General Assembly of Pennsylvania; and was re-elected by succeeding assemblies for several years, till he was chosen a representative for the city of Philadelphia; and in 1737 he was appointed post-master of that city. In 1738, he formed the first fire-company there, to extinguish and prevent fires and the burning of houses: an example which was soon followed by other persons, and other places. And soon after, he suggested the plan of an association for insuring houses and ships from losses by fire, which was adopted; and the association continues to this day. In the year 1744, during a war between France and Great Britain, some French and Indians made inroads upon the frontier inhabitants of the province, who were unprovided for such an attack: the situation of the province was at this time truly alarming, being destitute of every means of defence. At this crisis Franklin stepped forth, and proposed to a meeting of the citizens of Philadelphia, a plan of a voluntary association for the defence of the province. This was approved of, and signed by 1200 persons immediately. Copies of it were circulated through the province; and in a short time the number of signers amounted to 10,000. Franklin was chosen colonel of the Philadelphia regiment; but he did not think proper to accept of the honour.

Pursuits of a different nature now occupied the greatest part of his attention for some years. Being always much addicted to the study of natural philosophy; and the discovery of the Leyden experiment in electricity having rendered that science an object of general curiosity; Mr. Franklin applied himself to it, and soon began to distinguish himself eminently in that way. He engaged in a course of electrical experiments with all the ardour and thirst for discovery which characterized the philosophers of that day. By these, he was enabled to make a number of important discoveries, and to propose theories to account for various phenomena; which have been generally adopted, and which will probably endure for ages. His observations he communicated, in a series of letters, to his friend Mr. Collinson; the first of which is dated March 28, 1747. In these he makes known the power of points in drawing and throwing off the electric matter, which had hitherto escaped the notice of electricians. He also made the discovery of a *plus* and *minus*, or of a *positive* and *negative* state of electricity; from whence in a satisfactory manner he explained the phenomena of the Leyden phial, first observed by Cuneus or Muschenbroeck, which had much perplexed philosophers. He shewed that the bottle, when charged, contained no more electricity than before, but that as much was taken from one side as was thrown on the other; and that, to discharge it, it was only necessary to make a communication between the two sides, by which the equilibrium might be restored, and that then no signs of electricity would remain. He afterwards demonstrated by experiments, that the electricity did not reside in the coating, as had been supposed, but in the pores of the glass itself. After a phial was charged, he removed the coating, and found that upon applying a new coating the shock might still be received. In the year 1749, he first suggested his idea of explaining the phenomena

of thunder-gusts, and of the aurora borealis, upon electrical principles. He points out many particulars in which lightning and electricity agree; and he adduces many facts, and reasoning from facts, in support of his positions. In the same year he conceived the bold and grand idea of ascertaining the truth of his doctrine, by actually drawing down the forked lightning, by means of sharp-pointed iron rods raised into the region of the clouds; from whence he derived his method of securing buildings and ships from being damaged by lightning. It was not until the summer of 1752 that he was enabled to complete his grand discovery the experiment of the electrical kite, which being raised up into the clouds, brought thence the electricity or lightning down to the earth; and M. D'Alibard made the experiment about the same time in France, by following the track which Franklin had before pointed out. The letters which he sent to Mr. Collinson, it is said, were refused a place among the papers of the Royal Society of London; and Mr. Collinson published them in a separate volume, under the title of *New Experiments and Observations on Electricity, made at Philadelphia, in America*; which were read with avidity, and soon translated into different languages. His theories were at first opposed by several philosophers, and by the members of the Royal Society of London; but in 1755, when he returned to that city, they voted him the gold medal which is annually given to the person who presents the best paper on some interesting subject. He was also admitted a member of the Society, and had the degree of doctor of laws conferred upon him by different universities: but at this time, by reason of the war which broke out between Britain and France, he returned to America, and interested himself in the public affairs of that country. Indeed he had done this long before; for although philosophy was a principal object of Franklin's pursuit for several years, he did not confine himself to it alone. In the year 1747 he became a member of the General Assembly of Pennsylvania, as a Burgess for the city of Philadelphia. Being a friend to the rights of man from his infancy, he soon distinguished himself as a steady opponent of the unjust schemes of the proprietaries. He was soon looked up to as the head of the opposition; and to him have been attributed many of the spirited replies of the assembly, to the messages of the governors. His influence in the body was very great. This arose not from any superior powers of eloquence; he spoke but seldom; and he never was known to make any thing like an elaborate harangue. His speeches often consisted of a single sentence, or of a well-told story, the moral of which was always obviously to the point. He never attempted the flowery fields of oratory. His manner was plain and mild. His style in speaking was, like that of his writings, simple, unadorned, and remarkably concise. With this plain manner, and his penetrating and solid judgment, he was able to confound the most eloquent and subtle of his adversaries, to confirm the opinions of his friends, and to make converts of the unprejudiced who had opposed him. With a single observation, he has rendered of no avail a long and elegant discourse, and determined the fate of a question of importance.

In the year 1749, he proposed a plan of an academy,



to be erected in the city of Philadelphia, as a foundation for posterity to erect a seminary of learning, more extensive and suitable to future circumstances; and in the beginning of 1750, three of the schools were opened, namely, the Latin and Greek school, the Mathematical, and the English schools. This foundation soon after gave rise to another more extensive college, incorporated by charter May 27, 1755, which still subsists, and in a very flourishing condition. In 1752, he was instrumental in the establishment of the Pennsylvania Hospital, for the cure and relief of indigent invalids, which has proved of the greatest use to that class of persons. Having conducted himself so well as Postmaster of Philadelphia, he was, in 1753, appointed Deputy Postmaster-general for the whole British colonies.

The colonies being much exposed to depredations in their frontier by the Indians and the French, at a meeting of commissioners from several of the provinces, Mr. Franklin proposed a plan for the general defence, to establish in the colonies a general government, to be administered by a president-general, appointed by the crown, and by a grand council, consisting of members chosen by the representatives of the different colonies; a plan which was unanimously agreed to by the commissioners present. The plan however had a singular fate: It was disapproved of by the ministry of Great Britain, because it gave too much power to the representatives of the people; and it was rejected by every assembly, as giving to the president general, who was to be the representative of the crown, an influence greater than appeared to them proper, in a plan of government intended for freemen. Perhaps this rejection, on both sides, is the strongest proof that could be adduced of the excellence of it, as suited to the situation of Great Britain and America at that time. It appears to have steered exactly in the middle, between the opposite interests of both. Whether the adoption of this plan would have prevented the separation of America from Great Britain, is a question which might afford much room for speculation.

In the year 1755, General Braddock, with some regiments of regular troops, and provincial levies, was sent to dispossess the French of the posts upon which they had seized in the back settlements. After the men were all ready, a difficulty occurred, which had nearly prevented the expedition. This was the want of waggons. Franklin now stepped forward, and, with the assistance of his son, in a little time procured 150. After the defeat of Braddock, Franklin introduced into the assembly a bill for organizing a militia, and had the dexterity to get it passed. In consequence of this act a very respectable militia was formed; and Franklin was appointed colonel of a regiment in Philadelphia, which consisted of 1200 men; in which capacity he acquitted himself with much propriety, and was of singular service; though this militia was soon after disbanded by order of the English ministry.

In 1757, he was sent to England, with a petition to the king and council, against the proprietaries, who refused to bear any share in the public expences and assessments; which he got settled to the satisfaction of the state. After the completion of this business, Franklin remained at the court of Great Britain for some time, as agent for the province of Pennsylvania;

and also for those of Massachusetts, Maryland, and Georgia. Soon after this, he published his Canada pamphlet, in which he pointed out, in a very forcible manner, the advantages that would result from the conquest of this province from the French. An expedition was accordingly planned, and the command given to General Wolfe; the success of which is well known. He now divided his time indeed between philosophy and politics, rendering many services to both. Whilst here, he invented the elegant musical instrument called the *Armonica*, formed of glasses played on by the fingers. In the summer of 1762 he returned to America; on the passage to which he observed the singular effect produced by the agitation of a vessel, containing oil floating on water: the upper surface of the oil remained smooth and undisturbed, whilst the water was agitated with the utmost commotion. On his return he received the thanks of the Assembly of Pennsylvania, which having annually elected him a member in his absence, he again took his seat in this body, and continued a steady defender of the liberties of the people.

In 1764, by the intrigues of the proprietaries, Franklin lost his seat in the assembly, which he had possessed for 14 years; but was immediately appointed provincial agent to England, for which country he presently set out. In 1766 he was examined before the parliament relative to the stamp-act; which was soon after repealed. The same year he made a journey into Holland and Germany; and another into France; being everywhere received with the greatest respect by the literati of all nations. In 1773 he attracted the public attention by a letter on the duel between Mr. Whately and Mr. Temple, concerning the publication of Governor Hutchinson's letters, declaring that he was the person who had discovered those letters. On the 29th of January next year, he was examined before the privy-council on a petition he had presented long before as agent for Massachusetts Bay against Mr. Hutchinson: but this petition being disagreeable to ministry, it was precipitately rejected, and Dr. Franklin was soon after removed from his office of Postmaster-general for America. Finding now all efforts to restore harmony between Great Britain and her colonies useless, he returned to America in 1775; just after the commencement of hostilities. Being named one of the delegates to the Continental Congress, he had a principal share in bringing about the revolution and declaration of independency on the part of the colonies. In 1776 he was deputed by Congress to Canada, to negotiate with the people of that country, and to persuade them to throw off the British yoke; but the Canadians had been so much disgusted with the hot-headed zeal of the New Englanders, who had burnt some of their chapels, that they refused to listen to the proposals, though enforced by all the arguments Dr. Franklin could make use of. On his return to Philadelphia, Congress, sensible how much he was esteemed in France, sent him thither to put a finishing hand to the private negotiations of Mr. Silas Deane; and this important commission was readily accepted by the Doctor, though then in the 71st year of his age. The event is well known; a treaty of alliance and commerce was signed between France and America; and M. Le Roi asserts, that the Doctor had a great share in the transaction, by strongly advising M. Maurepas not



to lose a single moment, if he wished to secure the friendship of America, and to detach it from the mother-country. In 1777 he was regularly appointed plenipotentiary from Congress to the French court; but obtained leave of dismissal in 1780. Having at length seen the full accomplishment of his wishes by the conclusion of the peace in 1783, which gave independency to America, he became desirous of revisiting his native country. He therefore requested to be recalled; and, after repeated solicitations, Mr. Jefferson was appointed in his stead. On the arrival of his successor, he repaired to Havre de Grace, and crossing the channel, landed at Newport in the Isle of Wight; from whence, after a favourable passage, he arrived safe at Philadelphia in September 1785. He was received amidst the acclamations of a vast multitude who flocked from all parts to see him, and who conducted him in triumph to his own house; where in a few days he was visited by the members of the Congress and the principal inhabitants of Philadelphia. He was afterward twice chosen president of the Assembly of Philadelphia; but his increasing infirmities obliged him to ask permission to retire, and to spend the remainder of his life in tranquillity; which was granted, in 1788. After this, the infirmities of age increased fast upon him; he became more and more afflicted with the gout and the stone, till the time of his death, which happened the 17th of April 1790, about 11 at night, at 84 years of age; leaving one son, governor William Franklin, a zealous loyalist, who now resides in London; and a daughter, married to Mr. William Bache, merchant in Philadelphia.

Doctor Franklin was author of many tracts on electricity, and other branches of natural philosophy, as well as on politics, and miscellaneous subjects. He had also many papers inserted in the Philosophical Transactions, from the year 1757 to 1774.

**FREEZE, or FRIZE**, in Architecture, a large flat member, being that part of the entablature of columns that separates the architrave from the cornice.

**FREEZING**, or Congelation, the fixing of a fluid body into a firm or solid mass by the action of cold: in which sense the term is applied to water when it freezes into ice; to metals when they resume their solid form after being melted by heat; or to glass, wax, pitch, tallow, &c, when they harden again after having been rendered fluid by heat. But it differs from crystallization, which is rather a separation of the particles of a solid from a fluid in which it had been dissolved more by the moisture than the action of heat.

The process of congelation is always attended with the emission of heat, as is found by experiments on the freezing of water, wax, spermaceti, &c; for in such cases it is always found that a thermometer dipt into the fluid mass, keeps continually descending as this cools, till it arrive at a certain point, being the point of freezing, which is peculiar to each fluid, where it is rather stationary, and then rises for a little, while the congelation goes on. But by what means it is that fluid bodies should thus be rendered solid by cold, or fluid by heat, or what is introduced into the bodies by either of those principles, are matters the learned have never yet been able to discover, or to satisfy them-

selves upon. The following phenomena however are usually observed.

Water, and some other fluids, suddenly dilate and expand in the act of Freezing, so as to occupy a greater space in the form of ice than before, in consequence of which it is that ice is specifically lighter than the same fluid, and floats in it. And the degree of expansion of water, in the state of ice, is by some authors computed at about  $\frac{1}{8}$  of its volume. Oil however is an exception to this property, and quicksilver too, which shrinks and contracts still more after Freezing. Mr. Boyle relates several experiments of vessels made of metal, very thick and strong; in which, when filled with water, close stopped, and exposed to the cold, the water being expanded in Freezing, and not finding either room or vent, burst the vessels. A strong barrel of a gun, with water in it close stopped and frozen, was rent the whole length. Huygens, to try the force with which it expands, filled a cannon with it, whose sides were an inch thick, and then closed up the mouth and vent, so that none could escape; the whole being exposed to a strong Freezing air, the water froze in about 12 hours, and burst the piece in two places. Mathematicians have computed the force of the ice upon this occasion; and they say, that such a force would raise a weight of 27720 pounds. Lastly, Major Edward Williams, of the Royal Artillery, made many experiments on the force of it, at Quebec, in the years 1784 and 1785. He filled all sizes of iron bomb-shells with water, then plugged the fuze hole close up, and exposed them to the strong Freezing air of the winter in that climate; sometimes driving in the iron plugs as hard as possible with a sledge hammer; and yet they were always thrown out by the sudden expansion of the water in the act of Freezing, like a ball shot by gunpowder, sometimes to the distance of between 400 and 500 feet, though they weighed near 3 pounds; and when the plugs were screwed in, or furnished with hooks or barbs, to lay hold of the inside of the shell by, so that they could not possibly be forced out, in this case the shell was always split in two, though the thickness of the metal of the shell was about an inch and three quarters. It is farther remarkable, that through the circular crack, round about the shells, where they burst, there stood out a thin film or sheet of ice, like a fin; and in the cases when the plugs were projected by Freezing water, there suddenly issued out from the fuze-hole, a bolt of ice, of the same diameter, and stood over it to the height sometimes of 8 inches and a half. And hence we need not be surprised at the effects of ice in destroying the substance of vegetables and trees, and even splitting rocks, when the frost is carried to excess.

It is also observed that water loses of its weight by Freezing, being found lighter after thawing again, than before it was frozen. And indeed it evaporates almost as fast when frozen, as when it is fluid.

It is said too that water does not freeze in vacuo; requiring for that purpose the presence and contiguity of the air. But this circumstance is liable to some doubt, and it may be suspected that the degree of cold has not been carried far enough in these instances; as it is found that mercury in thermometers has even been frozen



frozen, though it requires a vastly greater degree of cold to freeze mercury, than water.

That water which has been boiled freezes more readily than that which has not been boiled; and that a slight disturbance of the fluid disposes it to freeze more speedily; having sometimes been cooled several degrees below the Freezing point, without congealing when kept quite still, but suddenly freezing into ice on the least motion or disturbance.

That the water, being covered over with a surface of oil of olives, does not freeze so readily as without it; and that nut oil absolutely preserves it under a strong frost, when olive oil would not.

That rectified spirit of wine, nut oil, and oil of turpentine, seldom freeze.

That the surface of the water, in Freezing, appears all wrinkled; the wrinkles being sometimes in parallel lines, and sometimes like rays, proceeding from a centre to the circumference.

*FREEZING Mixture*, a preparation for the artificial congelation of water, and other fluids.

According to Mr. Boyle, all kinds of salts, whether alkaline or acid; and even all spirits, as spirit of wine, &c; as also sugar, and saccharum saturni, mixed with snow, are capable of Freezing moist fluids; and the same effect is produced, in a very high degree, by a mixture of oil of vitriol, or spirit of nitre, with snow.

M. Homberg remarks the same of equal quantities of corrosive sublimate, and sal ammoniac, with four times the quantity of distilled vinegar.

Boerhaave gives a method of producing artificial frost without either snow or ice: we must have for this purpose, at any season of the year, the coldest water that can be procured; this is to be mixed with a proper quantity of any salt (sal ammoniac will answer the intention best), at the rate of about 3 ounces to a quart of water. Another quart of water must be prepared in the same manner with the first; the salt, by being dissolved in each, will make the water much colder than it was before. The two quarts are then to be mixed together, and this will make them colder still. Two quarts more of water prepared and mixed in the same manner are to be mixed with these, which will increase the cold to a much higher degree in all. The whole of this operation is to be carried on in a cold cellar; and a glass of common water is then to be placed in the vessel of the fluid thus artificially cooled, and it will be turned into ice in the space of 12 hours.

There is also a method of making artificial ice by means of snow, without any kind of salt. For this purpose fill a small pewter dish with water, and upon that set a common pewter plate filled, but not heaped, with snow. Bring this simple apparatus near the fire, and stir the snow in the plate: the snow will dissolve, and the ice will be formed on the back of the plate, which was set in the dish of water.

Mr. Reaumur tried the effect of several salts, and examined the various degrees of cold by an ice thermometer, which being placed in the fluid to be frozen, shewed very exactly the degree of cold by the descent of the spirit.

Nitre, or saltpetre, usually passes for a salt that may be very serviceable in these artificial congelations; but the experiments of this gentleman prove that this opi-

nion is erroneous. The most perfectly refined saltpetre employed in the operation sunk the spirit in the thermometer only 3 degrees and a half below the fixed point. Less refined nitre sunk the thermometer lower, and gave a greater degree of cold; owing to the common or sea-salt that it contains when less pure, which has a greater effect than the pure saltpetre itself.

Two parts of common salt being mixed with three parts of powdered ice in very hot weather, the spirit in the thermometer immediately descended 15 degrees, which is half a degree lower than it would have descended in the severest cold of our winters. Mr. Reaumur then tried the salts all round, determining with great regularity and exactness, what was the degree of cold occasioned by each in a given dose. Among the neutral salts, none produced a greater degree of cold than the common sea salt. Among the alkalies, sal ammoniac sunk the thermometer only to 13 degrees. Pot-ashes sunk it just as low as well refined saltpetre.

For the common uses of the table, the ice is not required to be very hard, or such as is produced by long continuance of violent cold: it is rather desired to be like snow. Saltpetre, which is no very powerful freezer, is therefore more fit for the purpose than a more potent salt. It is not necessary that the congelation should be very suddenly made; but that it may retain its form as long as may be, when made, is of great importance.

If it be desired to have ices very hard and firm, and very suddenly prepared, then sea salt is of all others most to be chosen for the operation. The ices thus made will be very hard, but they will soon run. Pot-ashes afford an ice of about the hardness that is usually required. This forms indeed very slowly, but then it will preserve a long time. And common wood ashes will perform the business very nearly in the same manner as the pot-ashes; but for this purpose, the wood which is burnt, ought to be fresh.

The strong acid spirits of the neutral salts act much more powerfully in these congelations than the salts themselves, or indeed than any simple salt can do. Thus, spirit of nitre mixed with twice its quantity of powdered ice, immediately sinks the spirit in the thermometer to 19 degrees, or 4 degrees more than that obtained by means of sea salt, the most powerful of all the salts in making artificial cold. A much greater degree of cold may be given to this mixture, by piling it round with more ice mixed with sea salt. This gives a redoubled cold, and sinks the thermometer to 24 degrees. If this whole matter be covered with a fresh mixture of spirit of nitre and ice, a still greater degree of cold is produced, and so on; the cold being by this method of fresh additions to be increased almost without bounds: but it is to be observed, that every addition gives a smaller increase than the former.

It is very remarkable in the acid spirits, that though sea salt is so much more powerful than nitre in substance in producing cold, yet the spirit of nitre is much stronger than that of sea salt; and another not less wonderful phenomenon is, that spirit of wine, which is little else than liquid fire, has as powerful an effect in congelations, or very nearly so, as the spirit of nitre itself.



The several liquid substances which produce cold, in the same manner as the dry salts on being mixed with ice, are much more speedy in their action than the salts: because they immediately and much more intimately come into contact with the particles of the ice, than the salts can. Of this nature are spirit of nitre, spirit of wine, &c. To produce the expected degree of cold, it is always necessary that the ice and the added matter, whatever it be, should both run together, and, intimately uniting, form one clear fluid. It is hence that no new cold is produced with oil, which, though it melts the ice, yet cannot mix itself into a homogeneous liquid with it, but must always remain floating on the surface of the water that is produced by the melting of the ice. *Mem. Acad. Scienc. Par. 1734.*

It has been discovered, that fluids standing in a current of air, grow by this means much colder than before. Fahrenheit had long since observed, that a pond, which stands quite calm, often acquires a degree of cold much beyond what is sufficient for Freezing, and yet no congelation ensued: but if a slight breath of air happens in such a case to brush over the surface of the water, it stiffens the whole in an instant. It has also been discovered, that all substances grow colder by the evaporation of the fluids which they contain, or with which they are mixed. If both these methods therefore be practised upon the same body at the same time, they will increase the cold to almost any degree of intenseness we please.

But the most extraordinary instances of artificial Freezing, have since been made in Russia, at Hudson's bay, and other parts, by which quicksilver was frozen into a solid mass of metal. And the same thing had before happened from the natural cold of the atmosphere alone, in Siberia. In the winter of 1733, Professor Gmelin, with two other gentlemen of the Russian Academy, were sent by Anne Ivanouna, the new empress, to explore and describe the different parts of her Asiatic dominions, with the communication of Asia and America. In the winter of 1734-5, Mr. Gmelin being at Yeneseisk in  $58^{\circ} 30'$  north lat. and  $92^{\circ}$  long. east from Greenwich, first observed such a descent of the mercury, as must have been attended with congelation, being far below its Freezing point, now fixed at  $-40$  of Fahrenheit's thermometer. "Here, says he, we first experienced the truth of what various travellers have related with respect to the extreme cold of Siberia; for, about the middle of December, such severe weather set in, as we were sure had never been known in our time at Petersburg. The air seemed as if it were frozen, with the appearance of a fog, which did not suffer the smoke to ascend as it issued from the chimneys. Birds fell down out of the air as dead, and froze immediately, unless they were brought into a warm room. Whenever the door was opened, a fog suddenly formed round it. During the day, short as it was, parhelia and haloes round the sun were frequently seen; and in the night mock moons, and haloes about the moon. Finally, our thermometer, not subject to the same deception as the senses, left us no doubt of the excessive cold; for the quicksilver in it was reduced, on the 5th of January, old style, to  $-120^{\circ}$  of Fahrenheit's scale, lower than it had ever been observed in nature."

The next instance of congelation happened at Yakutsk, in  $62^{\circ}$  north lat. and  $150^{\circ}$  east longitude. The weather here was unusually mild for the climate, yet the thermometer fell to  $-72^{\circ}$ ; and one person informed the professor by a note, that the mercury in his barometer was frozen. He hastened immediately to his house to behold such a surprising phenomenon; but though he was witness to the fact, observing that the mercury did not continue in one column, but was divided in different places as into little cylinders, which appeared frozen, yet the prejudice he had entertained against the possibility of the congelation, would not allow him to believe it.

Another set of observations, in the course of which the mercury must frequently have been congealed, were made by professor Gmelin at Kirenga fort, in  $57\frac{1}{2}$  north lat., and  $108$  east long.; his thermometer, at different times, standing at  $-108$ ,  $-86$ ,  $-100$ ,  $-113$ , and many other intermediate degrees; in the course of the winter of 1737-8. On the 27th of November, after the thermometer had been standing for two days at  $-46^{\circ}$ , he found it sunk at noon to  $-108$ . Suspecting some mistake, after he had noted down the observation, he instantly ran back, and found it at  $-102$ ; but ascending with such rapidity, that in the space of half an hour it had risen to  $-19^{\circ}$ . This phenomenon, which appeared so surprising, doubtless depended on the expansion of the mercury frozen in the bulb of the thermometer, and which now melting, forced upwards the small thread in the stem. And similar appearances were observed on other days afterwards, when the thread of quicksilver in the thermometer was separated about 6 degrees.

A second instance where a natural congelation of mercury has certainly been observed, is recorded in the transactions of the Royal Academy of Sciences at Stockholm, as made by Mr. Andrew Hellant. The weather, in January 1760, was remarkably cold in Lapland; so that on the 5th of that month, the thermometers fell to  $-76$ ,  $-128$ , or lower; on the 23d and following days they fell to  $-58$ ,  $-79$ ,  $-92$ , and below  $-238$  entirely into the ball. This was observed at four different places in Lapland, situated between the 65th and 78th degrees of north lat. and the 21st and 28th degree of east longitude.

But the congelation of quicksilver, by an artificial Freezing mixture, was first observed, and put beyond doubt, by Mr. Joseph Adam Braun, professor of philosophy at Petersburg. This gentleman wishing to try how many degrees of cold he could produce, availed himself of a good opportunity which offered for that purpose on the 14th of December 1759, when the mercury in the thermometer stood in the natural cold at  $-34$ , which it is now known is only 5 or 6 degrees above its point of congelation. Assisting this natural cold therefore with a mixture prepared of aquafortis and pounded ice, his thermometer was sunk to  $-69$ . Part of the quicksilver must now have been really congealed, but unexpected by him, and he only thought of pursuing his object of producing still greater degrees of cold; and having expended all his pounded ice, he was obliged to use snow instead of it. With this fresh mixture the mercury sunk to  $-100$ ,  $-240$ , and  $-350^{\circ}$ . Taking the thermometer out, he found it whole,



whole, but the quicksilver fixed, and it continued so for 12 minutes. On repeating the experiment, with another thermometer which had been graduated no lower than  $-220$ , all the mercury sunk into the ball, and became solid as before, and did not re-ascend till after a still longer interval of time. Mr. Braun now suspected that the quicksilver was really frozen, and prepared for making a decisive experiment. This was accomplished on the 25th of the same month, and the bulb of the thermometer broken as soon as the metal was congealed; when it appeared that the mercury was changed into a solid and shining metallic mass, which flattened and extended under the strokes of a pestle, being rather less hard than lead, and yielding a dull sound like that metal. Mr. *Æpinus* made similar experiments at the same time, employing as well thermometers as tubes of a larger bore; in which last he remarked, that the quicksilver fell sensibly on being frozen, assuming a concave surface, and likewise that the congealed pieces sunk in fluid mercury: also, in their farther experiments, they invariably found that the mercury sunk lower when the whole of it was congealed, than if any part of it remained fluid: all shewing that, contrary to water, mercury contracted in Freezing. It was farther observed, that the mercury when congealed looked like the most polished silver, and when beaten flat, it was easily cut with a penknife, like soft thin sheet lead.

The fact being thus established, and fluidity no longer to be considered as an essential property of quicksilver, Mr. Braun communicated an account of his experiments to the Petersburg academy, on the 6th of September 1760; of which a large extract was inserted in the *Philos. Trans.* vol. 52, pa. 156. He afterwards declared that he never suffered a winter to pass without repeating the experiment of Freezing quicksilver, and never failed of success when the natural cold was of a sufficient strength for the purpose; and this degree of natural cold he supposes at  $-10$  of Fahrenheit; though some commencement of the congelation might be perceived when the temperature of the air was as high as  $+2$ .

The results of all his experiments were, that with the abovementioned frigorific mixtures, and once with rectified spirits and snow, when the natural cold was at  $-28^{\circ}$ , he congealed the quicksilver, and discovered that it is a real metal that melts with a very small degree of heat. However, not perceiving the necessary consequence of its great contraction in Freezing, he always confounded its point of congelation with that of its greatest contraction in Freezing, and thus marked the former a great deal too low.

In the process of his observations, Mr. Braun found that double aquafortis was more effectual than spirit of nitre; but with this simple spirit, which seldom brings the mercury lower than  $-148$ , this metal may be frozen in the following manner: Six glasses being filled with snow as usual, and the thermometer put in one of them, the spirit of nitre was poured upon it; when the mercury would fall no lower in this, the thermometer was removed to the second, and so on to the third and fourth, in which fourth immersion the mercury was congealed.

Mr. *Æpinus* gives the following direction for using

the fuming spirit of nitre: Take some of this spirit, cooled as much as possible, and put it into a wine glass till it be about half full, filling it up with snow, and stirring them till the mixture become of the consistence of pap; by which means you obtain, almost in an instant, the necessary degree of cold for the Freezing of quicksilver.

It is remarked by Mr. Braun, that by the mixture of snow and spirit of nitre, which froze the mercury, he never was able to bring thermometers, filled with the most highly rectified spirit of wine, lower than  $-148$ : so that the cold which will freeze mercury, will not freeze spirit of wine; and therefore spirit thermometers are the most fit to determine the degree of coldness in frigorific mixtures, till we can construct solid metallic thermometers with sufficient accuracy. Mr. Braun tried the effects of different Fluids in his frigorific mixtures: he always found that Glauber's spirit of nitre and double aquafortis were the most powerful; and from a number of experiments made when the temperature of the air was between 21 and 28 of Fahrenheit, he concludes, that spirit of salt pounded upon snow increased the natural cold  $36^{\circ}$ ; spirit of sal ammoniac, 12; oil of vitriol, 42; Glauber's spirit of nitre, 70; aquafortis, 48; simple spirit of nitre, 36; dulcified spirit of vitriol, 24; Hoffman's anodyne liquor, 38; spirit of hartshorn, 12; spirit of sulphur, 12; spirit of wine rectified, 24; camphorated spirit, 18; French brandy, 14; and several kinds of wine increased the natural cold to 7, 8, or 9 degrees.

The most remarkable congelation of mercury, by natural cold, that has ever been observed, was that related by Dr. Peter Simon Pallas, who had been sent by the empress of Russia, with some other gentlemen, on an expedition similar to that of Mr. Gmelin. Being at Krasnoyarsk in the year 1772, in north lat.  $56^{\circ} 30'$ , and east long.  $93^{\circ}$ , he had an opportunity of observing the phenomenon we speak of. On the 6th and 7th of December that year, says he, there happened the greatest cold I have ever experienced in Siberia: the air was calm at the time, and seemingly thickened; so that, though the sky was in other respects clear, the sun appeared as through a fog. I had only one small thermometer left, in which the scale went no lower than  $-7^{\circ}$ ; and on the 6th in the morning, I remarked that the quicksilver in it sunk into the ball, except some short columns which stuck fast in the tube. When the ball of the thermometer, as it hung in the open air, was touched with the finger, the quicksilver rose; and it could plainly be seen that the solid columns stuck and resisted a good while, and were at length pushed upward with a sort of violence. He also placed upon the gallery, on the north side of his house, some quicksilver in an open bowl. Within an hour he found the edges and surface of it frozen solid; and some minutes afterward the whole was condensed by the natural cold into a soft mass very much like tin. While the inner part was still fluid, the frozen surface exhibited a great variety of branched wrinkles; but in general it remained pretty smooth in Freezing. The congealed mercury was more flexible than lead; but on being bent short, it was found more brittle than tin; and when hammered out thin, it seemed somewhat granulated. When the hammer was not perfectly cooled, the quicksilver



quickfilver melted away under it in drops; and the same thing happened when the metal was touched with the finger, by which also the finger was immediately benumbed. When the frozen mass was broken to pieces in the cold, the fragments adhered to each other and to the bowl in which they lay. In the warm room it thawed on its surface gradually, by drops, like wax on the fire, and did not melt all at once. Although the frost seemed to abate a little towards night; yet the congealed quickfilver remained unaltered, and the experiment with the thermometer could still be repeated. On the 7th of December he had an opportunity of making the same observations all day; but some hours after sunset, a northwest wind sprung up, which raised the thermometer to  $-46^{\circ}$ , when the mass of quickfilver began to melt.

The experiments of Mr. Braun were successfully repeated at Gottingen, in 1774, by Mr. John Frederick Blumenbach; being encouraged to this attempt by the excessive cold of the winter that year, especially the night of January the 11th, when he made the experiment, the thermometer standing at  $-10$  in the open air. Mr. Blumenbach at 5 in the evening, put 3 drachms of quickfilver into a small sugar glass, and covered it with a mixture of snow and Egyptian sal ammoniac, setting the glass out in the air upon a mixture also of sal ammoniac. At one the next morning, the mercury was found frozen quite solid, and hard to the glass; and did not melt again till 7 or 8 the next morning. The colour of the frozen mercury was a dull pale white with a blueish cast, like zinc, very different from the natural appearance of quickfilver.

In the year 1775, by similar means, quickfilver was twice frozen by Mr. Hutchins, governor of Albany fort, in Hudson's bay, viz, in the months of January and February of that year. And the same was done on the 28th of January 1776, by Dr. Lambert Bicker, secretary of Rotterdam. The temperature of the atmosphere was then at  $+2^{\circ}$ ; and the lowest it could reduce the thermometer by artificial cold, was  $-94$ ; when, on breaking the glass, the mercury was found frozen.

In the beginning of the year 1780 M. Von Elterlein of Vytegra, a town of Russia, in lat.  $61^{\circ}$  north, and long.  $36^{\circ}$  east, froze quickfilver by natural cold. On the 4th of January 1780, the cold being increased to  $-34$  that evening at Vytegra, he exposed to the open air 3 ounces of very pure quickfilver in a china teacup, covered with paper pierced full of holes. Next day, at 8 in the morning, he found it solid, and looking like a piece of cast lead, with a considerable depression in the middle. On attempting to loosen it in the cup, his knife raised shavings from it as if it had been lead, which remained sticking up; and at length the metal separated from the bottom of the cup in one mass. He then took it in his hand to try if it would bend: it was stiff like glue, and broke into two pieces; but his fingers immediately lost all feeling, and could scarcely be restored in an hour and a half by rubbing with snow. At 8 o'clock the thermometer stood at  $-57$ ; but half after 9 it was risen to  $-40$ ; and then the two pieces of mercury which lay in the cup had lost so much of their hardness, that they could no longer be broken, or cut into shavings, but resembled

a thick amalgam, which, though it became fluid when pressed by the fingers, immediately afterwards resumed the consistence of pap. With the thermometer at  $-39$ , the quickfilver became fluid. The cold was never less on the 5th than  $-28$ , and by 9 in the evening it had increased again to  $-33$ . This experiment seems to fix the Freezing point of mercury at  $-40$  of Fahrenheit's thermometer, or 40 below 0; which is  $72^{\circ}$  below the Freezing point of water.

In the winter of 1781 and 82, Mr. Hutchins resumed the subject of Freezing quickfilver by artificial cold, with such success, that from his experiments and those of M. Von Elterlein, last mentioned, the Freezing point of mercury is now almost as well settled, viz at  $-40$ , as that of water is at  $+32$ . Other philosophers indeed had not been altogether inattentive to this subject. Professor Braun himself had taken great pains to investigate it; but for want of a proper attention to the difference between the contraction of the fluid mercury by cold, and that of the congealing metal by Freezing, he could not determine any thing certain concerning it.

An instance of the natural congelation of quickfilver also occurred in Jemmland, one of the provinces of Sweden, on the 1st of January 1782; and lastly, on the 26th of the same month, Mr. Hutchins observed the same effect of the cold at Hudson's bay; when he found that at the point of its Freezing a mercurial thermometer stood at  $-40$ , and a spirit thermometer at  $-30$ .

On this subject, see the Philos. Transf. vol. 51, pa. 672; vol. 52, pa. 156; vol. 66, pa. 174; vol. 73, pa. 303 and 325; vol. 76, pa. 241; vol. 77, pa. 285; vol. 78, pa. 43; and several others, particularly vol. 79, pa. 199, &c, being experiments on the congelation of quickfilver in England, by Mr. Richard Walker, where he proves that mercury may be frozen not only in England in summer, but even in the hottest climate, at any season of the year, and without the use of ice or snow.

**FREEZING Point**, denotes the point or degree of cold, shewn by a mercurial thermometer, at which certain fluids begin to freeze, or, when frozen, at which they begin to thaw again. On Fahrenheit's thermometer, this point is at  $+32$  for water, and at  $-40$  for quickfilver, these fluids freezing at those two points respectively. It would also be well if the Freezing points for other fluids were ascertained, and the whole arranged in a table.

**FREEZING Rain**, or *Raining Ice*, a very uncommon kind of shower which sometimes falls, particularly one in December 1672, in the west of England: of which some accounts are given in the Philos. Transf. number 90.

This rain, as soon as it touched any thing above ground, as a bough, or the like, immediately settled into ice; and by enlarging and multiplying the icicles, it broke all down with its weight. The rain that fell on the snow, immediately froze into ice, without sinking in the snow at all.

It made an amazing destruction of trees, beyond any thing in all history. "Had it concluded with some gust of wind, says a gentleman on the spot, it might have been of terrible consequence. Having weighed the



the sprig of an ash tree, the wood of which was just three quarters of a pound, the ice upon it amounted to 16 pounds. Some were frightened with the noise in the air; till they discerned it was the clatter of icy boughs dashed against each other." Dr. Beale observes, that there was no considerable frost perceived on the ground during the whole; from which he concludes, that a frost may be very fierce and dangerous on the tops of some hills, while in other places it keeps at some feet above the ground; and may wander about very furious in some places, and be remiss in others not far off. This rain was followed by glowing heats, and a wonderful forwardness of vegetation.

**FRENICLE** (**BERNARD**), a celebrated French mathematician of the 17th century. He was the contemporary and companion of Des Cartes, Fermat, and the other learned mathematicians of their time. He was admitted Geometrician of the French Academy in 1666; and died in 1675.

He had many papers inserted in the *Ancient Memoirs* of the Academy, of 1666, particularly in vol. 5, of that collection, viz, 1. A method of resolving problems by Exclusions.—2. Treatise of right-angled Triangles in Numbers.—3. Short tract on Combinations.—4. Tables of Magic Squares.—5. General method of making Tables of Magic Squares.

**FRESCO**, is a sort of painting which is made upon the plastering of walls before it is dry.

**FRIABILITY**, the property of a body that is Friable.

**FRIABLE**, a quality of bodies by which they are rendered tender and brittle, easily crumbled or reduced to powder between the fingers; their force of cohesion being such as easily exposes them to such solution. Such are pumice, and all calcined stones, burnt allum, &c.

It is supposed that Friability arises from hence, that the body consists wholly of dry parts irregularly combined, and which are readily separated, as having nothing unctuous or glutinous to bind them together.

**FRICTION**, the act of rubbing or grating the surfaces of bodies against or over each other, called also Attrition.

The phenomena arising from the Friction of divers bodies, under different circumstances, are very numerous and considerable. Mr. Hawksbee gives a number of experiments of this kind; particularly of the attrition or Friction of glass, under various circumstances; the result of which was, that it yielded light, and became electrical. Indeed all bodies by Friction are brought to conceive heat; many of them to emit light; particularly a cat's back, sugar, beaten sulphur, mercury, sea water, gold, copper, &c, but above all diamonds, which when briskly rubbed against glass, gold, or the like, yield a light equal to that of a live coal when blown by the bellows.

**FRICTION**, in Mechanics, denotes the resistance a moving body meets with from the surface on which it moves.

Friction arises from the roughness or asperity of the surface of the body moved on, and that of the body moving: for such surfaces consisting alternately of eminences and cavities, either the eminences of the one must be raised over those of the other, or they must be

both broken and worn off: but neither can happen without motion, nor can motion be produced without a force impressed. Hence the force applied to move the body is either wholly or partly spent on this effect; and consequently there arises a resistance, or Friction, which will be greater as the eminences are greater, and the substance the harder; and as the body, by continual Friction, becomes more and more polished, the Friction diminishes.

As the Friction is less in a body that rolls, than when it slides, hence in machines, lest the Friction should employ a great part of the power, care is to be taken that no part of the machine slide along another, if it can be avoided; but rather that they roll, or turn upon each other. With this view it may be proper to lay the axes of cylinders, not in a groove or concave matrix, as usual, but between little wheels, called Friction wheels, moveable on their respective axes: for by this contrivance, the Friction is transferred from the circumference of those wheels to their pivots. And in like manner the Friction may be still farther diminished, by making the axis of those wheels rest upon other Friction wheels that turn round with them. This was long since recommended by P. Casabus; and experience confirms the truth of it. Hence also it is, that a pulley moveable on its axis resists less than if it were fixed, and the cord sliding over the circumference. And the same may be observed of the wheels of coaches, and other carriages. Indeed about 20 years ago Friction balls or rollers were placed within the naves of carriage wheels by some persons, particularly a Mr. Varlo; and lately Mr. Garnett had a patent for an improved manner of applying Friction wheels to any axis, as of carriages, blocks or pulleys, scale beams, &c, in which the inclosed wheels or rollers are kept always at the same distance by connecting rods or bars.

From these principles, with the assistance of the higher geometry, Claus Roemer determined the figure of the teeth of wheels, that should make the least resistance possible, which he found should be epicycloids. And the same was afterwards demonstrated by De la Hire, and Camus.

M. Amontons, by experiment, attempted to settle a foundation for the precise calculation of the quantity of Friction; which M. Parent endeavoured to confirm from reasoning and geometry. M. Amontons' principle is, that the Friction of two bodies depends only on the weight or force with which they press each other, being always more or less in proportion to that pressure; esteeming it a vulgar error, that the quantity of Friction has any dependence on the extent of the surface that is rubbed, or that the Friction increases with the surface: arguing that it will require the same weight to draw along a plane, a piece of wood on its narrow edge, as on its broad and flat side; because, though on the broad side there be 4 times the number of touching particles, yet each particle is pressed with but  $\frac{1}{4}$  of the weight bearing on those of the narrow side; and since 4 times the number multiplied by  $\frac{1}{4}$  of the weight is equal to  $\frac{1}{4}$  of the number multiplied by 4 times the weight, it is plain that the effect, that is, the resistance, is equal in both cases, and therefore requires the same force to overcome it.

On the first proposal of this paradox, M. De la Hire very



very properly had recourse to experiments, as the best test, had they been judiciously performed: such as they were however, they succeeded in favour of this system. He laid several pieces of rough wood on a rough table; their sizes were unequal; but he laid weights on them, so as to render them all equally heavy: and he found that the same precise force, or weight, applied to them by a little pulley, was required to put each in motion, notwithstanding all the inequality of the surfaces. The experiment succeeded in the same manner with pieces of marble, laid on a marble table. After this, by reasoning, M. de la Hire gave a physical solution of the effect. And M. Amontons settled a calculus of the value of Friction, with the loss sustained by it in machines, on the foundation of this new principle. In wood, iron, lead, and brass, which are the chief materials used in machines, he makes the resistance caused by Friction to be nearly the same in all, when those materials are anointed with oil or fat: and the quantity of this resistance, independent of the magnitude of the surface, he makes nearly equal to a third part of the weight of the body moved, or of the force with which the two bodies are pressed together. Others have observed, that if the surfaces be hard and well polished, the Friction will be less than a third part of the weight; but if the parts be soft or rugged, it will be much greater. It was farther observed, that in a cylinder moved on two small gudgeons, or on a small axis, the Friction would be diminished in the same proportion as the diameter of these gudgeons is less than the diameter of the cylinder; because in this case, the parts on which the cylinder moves and rubs, will have less velocity than the power which moves it in the same proportion, which is in effect making the Friction to be proportional to the velocity. So that, from the whole of their observations, this general proposition is deduced, viz, That the resistances arising from Friction, are to one another in a ratio compounded of the pressures of the rubbing parts, and the velocities of their motions. Principles which, it is now known from better experiments, are both erroneous; notwithstanding the hypothesis of M. Amontons has been adopted, and attempted to be confirmed by Camus, Desaguliers, and others.

M. Muschenbroek and the abbé Nollet, however, on the other hand, have concluded from experiments, that the Friction of bodies depends on the magnitude of their surface, as well as on their weight. Though the former says, that in small velocities the Friction varies very nearly as the velocity, but that in great velocities the proportion increases faster: he has also attempted to prove, that by increasing the weight of a body, the Friction does not always increase exactly in the same ratio. *Introd. ad Phil. Nat.* vol. 1, c. 9, and *Lect. Phys. Exp.* tom. 1, p. 241. Helsham and Ferguson, from the same kind of experiments, have endeavoured to prove, that the Friction does not vary by changing the quantity of surface on which the body moves; and the latter of these asserts, that the Friction increases very nearly as the velocity; and that by increasing the weight, the Friction is increased in the same ratio. Indeed there is scarce any subject of experiment, with regard to which, different persons have formed such various conclusions. Of those who have written on the theory, no one has established it altogether on true principles, till the experiments

lately made by Mr. Vince of Cambridge: Euler, whose theory is extremely elegant, and would have been quite satisfactory had his principles been founded on good experiments, supposes the Friction to vary in proportion to the velocity of the body, and its pressure upon the plane; neither of which is true: and others, though they have justly imagined that Friction is a uniformly retarding force, have yet retained the other supposition, and so rendered their solutions not at all applicable to the cases for which they were intended.

For these reasons a new and ingenious set of experiments was successfully instituted by the rev. Samuel Vince, A. M. of Cambridge, which are published in the 75th vol. of the *Philos. Trans.* p. 165. The object of these experiments was to determine,

1st, Whether Friction be a uniformly retarding force.

2d, The quantity of Friction.

3d, Whether Friction varies in proportion to the pressure or weight.

4th, Whether the Friction be the same on whichever of its surfaces a body moves.

Mr. Vince says, “the experiments were made with the utmost care and attention, and the several results agreed so very exactly with each other, that I do not scruple to pronounce them to be conclusive.”—“A plane was adjusted parallel to the horizon, at the extremity of which was placed a pulley, which could be elevated or depressed in order to render the string which connected the body and the moving force parallel to the plane or horizon. A scale accurately divided was placed by the side of the pulley perpendicular to the horizon, by the side of which the moving force descended; upon the scale was placed a moveable stage, which could be adjusted to the space through which the moving force descended in any given time, which time was measured by a well regulated pendulum clock vibrating seconds. Every thing being thus prepared, the following experiments were made to ascertain the law of Friction. But let me first observe, that if Friction be a uniform force, the difference between it and the given force of the moving power must be also uniform, and therefore the moving body must descend with a uniformly accelerated velocity, and consequently the spaces described from the beginning of the motion must be as the squares of the times, just as when there was no Friction, only they will be diminished on account of the Friction.” Accordingly the experiments are then related, which are performed agreeably to these ingenious and philosophical ideas, and from them are deduced these general conclusions, which may be considered as established and certain facts or maxims. viz,

1st, That Friction is a uniformly retarding force in hard bodies, not subject to alteration by the velocity; except when the body is covered with cloth, woollen, &c, and in this case the Friction increases a little with the velocity.

2dly, Friction increases in a less ratio than the quantity of matter, or weight of the body. This increase however is different for the different bodies, more or less; nor is it yet sufficiently known, for any one body,



body, what proportion the increase of Friction bears to the increase of weight.

3dly, The smallest surface has the least Friction; the weight being the same. But the ratio of the Friction to the surface is not yet accurately known.

Mr. Vince's experiments consisted in determining how far the sliding bodies would be drawn, in given times, by a weight hanging freely over a pulley. This method would both shew him if the Friction were a constant retarding force, and the other conclusions above stated. For as the spaces described by any constant force, in given times, are as the squares of the times; and as the weight drawing the body is a constant force, if the Friction, which acts in opposition to the weight, should also be a constant force, then their difference, or the force by which the body is urged, will also be constant, in which case the spaces described ought to be as the squares of the times; which happened accordingly in the experiments.

Mr. Vince adds some remarks on the nature of the experiments which have been made by others. These, he observes, the authors "have instituted, To find what moving force would *just* put a body at rest in motion: and they concluded from thence, that the accelerative force was then equal to the Friction; but it is manifest, that any force which will put a body in motion must be greater than the force which opposes its motion, otherwise it could not overcome it; and hence, if there were no other objection than this, it is evident, that the Friction could not be very accurately obtained; but there is another objection, which totally destroys the experiment, so far as it tends to shew the quantity of Friction, which is the strong cohesion of the body to the plane when it lies at rest." This he confirms by several experiments, and then adds, "From these experiments therefore it appears, how very considerable the cohesion was in proportion to the Friction when the body was in motion; it being, in one case almost  $\frac{2}{3}$ , and in another it was found to be very nearly equal to the whole Friction. All the conclusions therefore deduced from the experiments, which have been instituted to determine the Friction from the force necessary to *put* a body in motion (and I have never seen any described but upon such a principle) have manifestly been totally false; as such experiments only shew the resistance which arises from the cohesion and Friction conjointly." *Philos. Trans.* vol. 75, pa. 165.

Mr. Emerson, in his *Principles of Mechanics*, deduces from experiments the following remarks relating to the quantity of Friction: When a cubic piece of soft wood of 8 pounds weight, moves upon a smooth plane of soft wood, at the rate of 3 feet per second, its Friction is about  $\frac{1}{3}$  of the weight; but if it be rough, the Friction is little less than half the weight: on the same supposition, when both the pieces of wood are very smooth, the Friction is about  $\frac{1}{4}$  of the weight: the Friction of soft wood on hard, or of hard wood upon soft, is  $\frac{1}{5}$  or  $\frac{1}{6}$  of the weight; of hard wood upon hard wood,  $\frac{1}{7}$  or  $\frac{1}{8}$ ; of polished steel moving on steel or pewter,  $\frac{1}{4}$ ; moving on copper or lead,  $\frac{1}{7}$  of the weight. He observes in general, that metals of the same sort have more Friction than those of different sorts; that lead makes much resistance; that iron or steel running in brass makes the least Friction of any; and that metals oiled make the Friction less than when

polished, and twice as little as when unpolished. Desaguliers observes that, in M. Camus's experiments on small models of sledges in actual motion, there are more cases in which the Friction is less than where it is more than  $\frac{1}{3}$  of the weight. See a table, exhibiting the Friction between various substances, formed from his experiments in *Desag. Exp. Philos.* vol. 1, p. 193 &c. also p. 133 to 138, and p. 182 to 254, and p. 458 to 460. On the subject of Friction, see several vols. of the *Philos. Trans.* as vol. 1, p. 206; vol. 34, p. 77; vol. 37, p. 394; vol. 53, p. 139, &c.

FRIDAY, the 6th day of the week, so called from Friga, or Friya, a goddess worshipped by the Saxons on this day. It is a fast-day in the church of England, in memory of our Saviour's crucifixion, unless Christmas-day happen to fall on Friday, which is always a festival.

Good FRIDAY, the Friday next before Easter, representing the day of our Saviour's crucifixion.

FRIGID Zone, the space about either pole of the earth to which the sun never rises for one whole day at least in their winter. These two zones extend to about  $23\frac{1}{2}$  degrees every way from the pole, as their centre.

FRIGORIFIC, in Physics, something belonging to, or that occasions cold.—Some philosophers, as Gassendus, and other corpuscularians, denying cold to be a mere privation, or absence of heat, contend that there are actual Frigorific corpuscles or particles, as well as fiery ones: whence proceed cold and heat. But later philosophers allow of no other Frigorific particles beside those nitrous salts &c, which float in the air in cold weather, and occasion freezing.

FRIZE, FRIEZE, or FREEZE, in Architecture, a part of the entablature of columns, between the architrave and cornice.

FRONT, in Architecture, denotes the principal face or side of a building; or that presented to their chief aspect and view.

FRONT, in Perspective, a projection or representation of the face, or forepart of an object, or of that part directly opposite to the eye, called also more usually *orthography*.

FRONTISPIECE, in Architecture, the portale, or principal face of a fine building.

FRONTON, in Architecture, an ornament among us more usually called Pediment.

FROST, such a state of the atmosphere as causes the congelation or freezing of water or other fluids into ice.

The nature and effects of Frost in different countries, are mentioned under the articles Congelation, and Freezing. In the more northern parts of the world, even solid bodies are affected by Frost, though this is only or chiefly in consequence of the moisture they contain, which being frozen into ice, and so expanding as water is known to do when frozen, it bursts and rends any thing in which it is contained, as plants, trees, stones, and large rocks. Some fluids expand by Frost, as water, which expands about  $\frac{1}{12}$ th part, for which reason ice floats in water; but others again contract, as quicksilver, and hence frozen quicksilver sinks in the fluid metal.

Frost, being derived from the atmosphere, naturally proceeds from the upper parts of bodies downwards, as the water and the earth: so, the longer a Frost is continued,



continued, the thicker the ice becomes upon the water in ponds, and the deeper into the earth the ground is frozen. In about 16 or 17 days Frost, Mr. Boyle found it had penetrated 14 inches into the ground. At Moscow, in a hard season, the Frost will penetrate 2 feet deep in the ground; and Capt. James found it penetrated 10 feet deep in Charlton island, and the water in the same island was frozen to the depth of 6 feet. Scheffer assures us, that in Sweden the Frost pierces 2 cubits, or Swedish ells into the earth, and turns what moisture is found there into a whitish substance, like ice; and standing waters to 3 ells, or more. The same author also mentions sudden cracks or rifts in the ice of the lakes of Sweden, 9 or 10 feet deep, and many leagues long; the rupture being made with a noise not less loud than if many guns were discharged together. By such means however the fishes are furnished with air; so that they are rarely found dead.

The natural histories of Frosts furnish very extraordinary effects of them. The trees are often scorched, and burnt up, as with the most excessive heat; and split or shattered. In the great Frost in 1683, the trunks of oak, ash, walnut, &c, were miserably split and cleft, so that they might be seen through, and the cracks often attended with dreadful noises like the explosion of fire-arms. Philos. Transf. number 165.

The close of the year 1708, and the beginning of 1709, were remarkable, throughout the greatest part of Europe, for a severe Frost. Dr. Derham says, it was the greatest in degree, if not the most universal, in the memory of man; extending through most parts of Europe, though scarcely felt in Scotland or Ireland.

In very cold countries, meat may be preserved by the Frost 6 or 7 months, and prove tolerable good eating. See Capt. Middleton's observations made in Hudson's bay, in the Philos. Transf. no. 465, sect. 2. In that climate the Frost seems never out of the ground, it having been found hard frozen in the two summer months. Brandy and spirit of wine, set out in the open air, freeze to solid ice in 3 or 4 hours. Lakes and standing waters, not above 10 or 12 feet deep, are frozen to the ground in winter, and all their fish perish. But in rivers, where the current of the tide is strong, the ice does not reach so deep, and the fish are preserved. Id. ib.

Some remarkable instances of Frost in Europe, and chiefly in England, are recorded as below: In the year

- 220, Frost in Britain that lasted 5 months.
- 250, The Thames frozen 9 weeks.
- 291, Most rivers in Britain frozen 6 weeks.
- 359, Severe Frost in Scotland for 14 weeks.
- 508, The rivers in Britain frozen for 2 months.
- 558, The Danube quite frozen over.
- 695, Thames frozen 6 weeks; booths built on it.
- 759, Frost from Oct. 1 till Feb. 26, 760.
- 827, Frost in England for 9 weeks.
- 859, Carriages used on the Adriatic sea.
- 908, Most rivers in England frozen 2 months.
- 923, The Thames frozen 13 weeks.
- 987, Frost lasted 120 days: began Dec. 22.
- 998, The Thames frozen 5 weeks.

1035, Severe Frost on June 24: the corn and fruits destroyed.

1063, The Thames frozen 14 weeks.

1076, Frost in England from Nov. till April.

1114, Several wooden bridges carried away by ice.

1205, Frost from Jan. 14 till March 22.

1407, Frost that lasted 15 weeks.

1434, From Nov. 24 till Feb. 10: Thames frozen down to Gravesend.

1683, Frost for 13 weeks.

1708, Severe Frost for many weeks.

1715, The same for many weeks.

1739, One for 9 weeks. Began Dec. 24.

1742, Severe Frost for many weeks.

1747, Severe Frost in Russia.

1754, Severe one in England.

1760, The same in Germany.

1776, The same in England.

1788, Thames frozen below bridge; booths on it.

*Hoar Frost*, is the dew frozen or congealed, early in cold mornings; chiefly in autumn. Though many Cartesians will have it formed of a cloud; and either congealed in the cloud, and so let fall; or ready to be congealed as soon as it arrives at the earth.

*Hoar Frost*, M. Regis observes, consists of an assemblage of little parcels of ice crystals; which are of various figures, according to the different disposition of the vapours, when met and condensed by the cold.

**FRUSTUM**, in Geometry, is the part of a solid next the base, left by cutting off the top, or segment, by a plane parallel to the base: as the Frustum of a pyramid, of a cone, of a conoid, of a spheroid, or of a sphere, which is any part comprised between two parallel circular sections; and the Middle Frustum of a sphere, is that whose ends are equal circles, having the centre of the sphere in the middle of it, and equally distant from both ends.

*For the Solid Content of the Frustum of a cone, or of any pyramid, whatever figure the base may have.* Add into one sum, the areas of the two ends and the mean proportional between them; then  $\frac{1}{3}$  of that sum will be a mean area, or the area of an equal prism, of the same altitude with the Frustum; and consequently that mean area being multiplied by the height of the Frustum, the product will be the solid content of it.

That is, if  $A$  denote the area of the greater end,  $a$  that of the less, and  $h$  the height;

then  $A + a + \sqrt{Aa} \times \frac{1}{3}h$  is the solidity.

Other rules for pyramidal or conic Frustums may be seen in my Mensuration, p. 189, 2d edit. 1788.


*The curve Surface of the Zone or Frustum of a sphere,* is had by multiplying the circumference of the sphere by the height of the Frustum. Mensur. p. 197.

*And the Solidity of the same Frustum is found,* by adding together the squares of the radii of the two ends, and  $\frac{1}{3}$  of the square of the height of the Frustum, then multiplying the sum by the said height and by the number 1.5708. That is,  $R^2 + r^2 + \frac{1}{3}h^2 \times \frac{1}{2}ph$  is the solid content of the spheric Frustum, whose height is  $h$ , and the radii of its ends  $R$  and  $r$ ,  $p$  being = 3.1416. Mensur. p. 209.

For the Frustums of spheroids, and conoids, either parabolic or hyperbolic, see Mensur. p. 326, 328, 332, 382, 435. And in p. 486 &c, are general theorems concerning the Frustum of a sphere, cone, spheroid, or conoid,



conoid, terminated by parallel planes, when compared with a cylinder of the same altitude, on a base equal to the middle section of the Frustrum made by a parallel plane. The difference between the Frustrum and the cylinder is always the same quantity, in different parts of the same, or of similar solids, or whatever the magnitude of the two parallel ends may be; the inclination of those ends to the axis, and the altitude of the Frustrum being given; and the said constant difference is  $\frac{1}{4}$  part of a cone of the same altitude with the Frustrum, and the radius of its base is to that altitude, as the fixed axis is to the revolving axis of the Frustrum. Thus, if BEC be any conic section, or a right line, or a circle, whose axis, or a part of it, is AD; AB and CD the extreme ordinates, FE the middle ordinate, AF being = FD; then taking, as AD to DK, so is the whole fixed axis, of which AD is a



part, to its conjugate axis; and completing the parallelogram AGHD: then if the whole figure revolve about the axis AD, the line BEC will generate the Frustrum of the cone or conoid, according as it is a right line or a conic section, or it will generate the whole solid when AB vanishes, or A and B meet in the same point; likewise AGHD will generate a cylinder, and ADK a cone: then is the 4th part of this cone always equal to the difference between the said cylinder generated by AGHD and the solid or Frustrum generated by ABEC; having all the same altitude or axis AD.

In the parabolic conoid, this difference and the cone vanish, and the Frustrum, or whole conoid ABEC, is always equal to the cylinder AGHD, of the same altitude.

In the sphere, or spheroid, the Frustrum ABEC is less than the cylinder AGHD, by  $\frac{1}{4}$  of the cone AKD. And

In the cone or hyperboloid, that Frustrum is greater than the cylinder, by  $\frac{1}{4}$  of the said cone AKD, which is similar to the other cone IBCD.

It may be observed, that the same relations are true, whether the ends of the Frustrum are perpendicular or oblique to the axis. And the same will hold for the Frustrum of any pyramid, whether right or oblique; and such a Frustrum of a pyramid will exceed the prism, of the same altitude, and upon the middle section of the Frustrum, by  $\frac{1}{4}$  of the same cone.

It has been observed, that the difference, or  $\frac{1}{4}$  of the cone AKD, is the same, or constant, when the altitude and inclination of the ends of the Frustrum remain the same. But when the inclination of the ends varies, the altitude being constant; then the said difference varies so as to be always reciprocally as the cube of the conjugate to the diameter AD. And when both the altitude and inclination of the ends vary, the differential cone is as the cube of the altitude directly, and the cube of the said conjugate diameter reciprocally: but if they vary so, as that the altitude is always reciprocally as that diameter, then the difference is a constant quantity.

Another general theorem for Frustrums, is this. In the Frustrum of any solid, generated by the revolution of any conic section about its axis, if to the sum of the

two ends be added 4 times the middle section,  $\frac{1}{4}$  of the last sum will be a mean area, and being drawn into the altitude of the solid, will produce the content. That is,  $\overline{AE + DC + 4FE} \times \frac{1}{4} AD$  is the content of ABCD.

And this theorem is general for all Frustrums, as well as the complete solids, whether right or oblique to the axis, and not only of the solids generated from the circle or conic sections, but also of all pyramids, cones, and in short of any solid whose parallel sections are similar figures.

The same theorem also holds good for any parabolic area ABEC, and is very nearly true for the area of any other curve whatever, or for the content of any other solid than those above mentioned.

FUGUE, in Music, is when the different parts of a musical composition follow each other, each repeating in order what the first had performed.

FULCRUM, or *Prop*, in Mechanics, is the fixed point about which a lever &c turns and moves.

FULGURATING *Phosphorus*, a term used by some English writers, to express a substance of the phosphorus kind. It was prepared both in a dry and liquid state, but the preparation it seems was not well known to any but the inventor of it. This matter not only shone in the dark in both states, but communicated its light to any thing it was rubbed on. When inclosed in a glass vessel well stopped, it sometimes would Fulgurate, or throw out little flashes of light, and sometimes fill the whole phial with waves of flame. It does not need recruiting its light at the fire, or in the sunshine, like the phosphorus of the Bolognian stone, but of itself continues in a state of shining for several years together, and is seen as soon as exposed in the dark; the solid or dry matter always resembling a burning coal of fire, though not consuming itself. *Philos. Trans.* No 134.

FULIGINOUS, an epithet applied to thick smoke or vapour replete with soot or other crass matter.

In the first fusion of lead, there exhales a great deal of Fuliginous vapour, which being retained and collected, makes what is called Litharge. And Lamp-black is what is gathered from the Fuliginous vapours of pines, and other resinous wood, when burnt.

FULMINANT, FULMINANS, or FULMINATING, an epithet applied to something that thunders, or makes a noise like thunder.

*Aurum FULMINANS.* See *AURUM*.

*Pulvis FULMINANS*, is a composition of 3 parts of nitre, 2 parts of salt of tartar, and 1 of sulphur.—Both the *Aurum* and *Pulvis Fulminans* produce their effect chiefly downwards; in which they differ from gunpowder, which acts in orbem, or all around, but principally upwards. When the composition is laid in brass ladles, and so set on fire, after fulmination, the ladles are often found perforated. It differs also from gunpowder in this, that it does not require to be confined, in order to fulminate, and it must be slowly and gradually heated. Some instants before explosion, a light blue flame appears on its surface, proceeding from the vapours beginning to kindle. No more fire or flame is perceived during the fulmination, being suffocated and extinguished by the quickness and violence of the commotion. Nor does the Fulminating powder generally



generally kindle the combustible bodies in contact with it, because the time of its inflammation is too short.

**FULMINATING Damp.** See **DAMP**.

**FULMINATION**, or **FULGURATION**, a vehement noise or shock resembling thunder, caused by the sudden explosion and inflammation of divers preparations; as aurum fulminans, &c, when set on fire.

**FUNCTION**, a term used in analytics, for an algebraical expression any how compounded of a certain letter or quantity with other quantities or numbers; and the expression is said to be a Function of that letter or quantity. Thus  $a - 4x$ , or  $ax + 3x^2$ , or  $2x - a\sqrt{a^2 - x^2}$ , or  $x^c$ , or  $c^x$ , is each of them a Function of the quantity  $x$ .

On the subject of Functions, their divisions, transformations, explication by infinite series, &c, see Euler's *Analys. Infinitorum*, c. 1, where the subject is fully treated.

**FURLONG**, an English long measure, containing 660 feet, or 220 yards, or 40 poles or perches, or the 8th part of a mile.

**FURNITURE**, in Dialling, certain additional points and lines drawn on a dial, by way of ornament. Such as the signs of the zodiac, length of days, parallels of declination, azimuths, points of the compass, meridians of chief cities, Babylonian, Jewish, or Italian hours, &c.

**FUSAROLE**, in Architecture, a small round member cut in form of a collar, with oval beads, under the echinus, or quarter-round, in the Doric, Ionic, and Composite capitals.

**FUSEE**, or **FUSY**, in Watch-work, is that part resembling a low cone with its sides a little sunk or concave, which is drawn by the spring, and about which the chain or string is wound.


The spring of a watch is the first mover. It is rolled up in a cylindrical box, against which it acts, and which it turns round in unbending itself. The chain, which at one end is wound about the Fusee, and at the other fastened to the spring-box, disengages itself from the Fusee in proportion as the box is turned. And hence the motion of all the other parts of the spring-watch. Now the effort or action of the spring is continually diminishing from first to last; and unless that inequality was rectified, it would draw the chain with more force, and wind a greater quantity of it upon the box, at one time than another; so that the movement would never keep equal time.

To correct this irregularity of the spring, it was very happily contrived to have the spring applied to the arms of levers, which are continually longer as the force of the spring is weaker: this foreign assistance, always increasing as it is most needed, maintains the action and effect of the spring in an equality.

It is for this reason then that the Fusee is made tapering somewhat conical, its radius at every point of the axis answering to the corresponding strength of the spring.

Now if the action of the spring diminished equally, as the parallels to the base of a triangle do; the cone,

which is generated of a triangle, would be the precise figure required for the Fusee; but it is certain that the weakening of the spring is not in that proportion; and therefore the Fusee should not be exactly conical; and in fact experience shews that it should be a little hollowed about the middle, because the action of the spring is not there sufficiently diminished of itself.



Mr. Varignon has investigated the figure of the Fusee, or the nature of the curve by whose revolution about its axis, shall be produced the solid whose figure the Fusee is to have. This curve it may easily be shewn is an hyperbola whose asymptote is the axis of the Fusee. Thus, let DFE be the curve of the Fusee, its axis being ABC: let AD express the greatest strength of the spring when the watch is quite wound up, or when the spring acts at D, and BG the least strength when the watch is down, or when the spring acts at E; so as that  $BE : AD :: AD : BG$ , or  $BE \times BG = AD^2$ ; join DG, producing it to meet the axis produced in C; then shall HI denote the strength of the spring acting at the corresponding point F of the Fusee; and the nature of it must be such that the rectangle  $HI \times HF$  be equal to a constant quantity, or HF must be reciprocally as HI, or  $HI : BG :: BE : HF$ ;

but because AD, HI, BG, are directly proportional to CA, CH, CB, theref. these are reciprocally propor. to AD, HF, BE; and consequently the curve DFE is an hyperbola, whose centre is C, and asymptotes AC and KL: so that the figure of the Fusee is the solid generated by an equilateral hyperbola revolved about its asymptote. See also Martin's *Mathem. Instit.* vol. 2, p. 364.

**FUSEE**, **FUSE**, or **FUZE**, in Artillery, is a wooden tap or tube used to set fire to the powder in a bomb-shell. The bore of this tube is filled with a composition, of sulphur one part, saltpetre 3 parts, and mealed powder 3, 4, or 5 parts. The tube is driven hard into the hole in the shell, having first cut it to the exact length answering to the time of the intended flight of the shell, so that, the composition in the Fuse catching fire by the discharge of the shell from the mortar, it just burns down to its lower end, and so sets fire to the powder in the shell, and thereby bursts it, at the moment when it arrives at the end of its range or flight.

**FUST**, in Architecture, the shaft of a column, or the part comprehended between the base and the capital; called also the Naked.

**FUZE**, or **FUZEE**. See **FUSEE**.



## G.

## G A G

**GABIONS**, in Fortification, are large cylindrical baskets, open at both ends, made of osier twigs, of 3 or 4 feet in diameter, and from 3 to 6 feet high. These, being filled with earth, are sometimes used as merlons for the batteries, and sometimes as a parapet for the lines of approach, when the attacks are carried on through a stony or rocky ground, and to advance them with extraordinary vigour. They serve also to make lodgments in some posts, and to secure other places from the shot of the enemy; who, on their part, endeavour to burn and destroy the Gabions, by throwing pitched faggots among them.

**GABLE**, or **GABLE-end**, of a house, is the upright triangular end, from the cornice or eaves to the top of its roof.

**GAGE**, in Hydrostatics, Pneumatics, &c, is an instrument for ascertaining measures of various kinds. As

**GAGE of the Air-pump**, is adapted for shewing the degree to which the air is rarefied, or the receiver is exhausted, at any time by the air-pump. This is either the common barometer-gage, both long and short, or the pear gage, which at first was thought a great improvement, but afterwards it was discovered that its seeming accuracy was founded on a fallacy, which gave an erroneous indication of exhaustion. See *Air-pump*.

**GAGE of the Barometer**, is a contrivance for estimating the exact degree of the rise and fall of the mercury in the tube of that instrument. It is well known that whilst the mercury rises in the tube, it sinks in the cistern, and vice versa; and consequently the divisions on the scale fixed near the top of the tube had their distance from the surface of the mercury in the cistern always various; from which there must often happen errors in determining the height of the mercury in the tube. To remedy this inconvenience, a line is cut upon a round piece of ivory, which is fixed near the cistern: this line is accurately placed at a given distance from the scale; for example at 27 inches; and a small float of cork, with a cylindrical piece of ivory fixed to its upper surface, on which a line is cut at the exact distance of 2 inches from the under side of the cork, is left to play freely on the quicksilver, and the cylinder works in a groove made in the other piece. From this construction it appears, that if these marks are made to coincide, by raising or lowering the screw which acts on the quicksilver, then the divisions on the scale will express the true measure of the distance from the surface.

**GAGE of the Condenser**, is a glass tube of a particular construction, adapted to the condensing engine, and designed to shew the exact density and quantity of the

air contained at any time in the condenser. See Defaguliers's Exper. Philos. vol. 2, p. 394.

**Sea GAGE**, an instrument for finding the depth of the sea. Several sorts of these have been invented by Dr. Hales, Dr. Defaguliers, and others. Formerly, the machines for this purpose consisted of two bodies, the one specifically lighter, and the other specifically heavier than the water, so joined together, that as soon as the heavy one came to the bottom, the lighter should get loose from it, and emerge; and the depth was to be estimated by the time the compound was in falling from the top to the bottom of the water, together with the time the lighter body was in rising, reckoned from the disappearing of the machine, till the emergent body was seen again; but no certain conclusion could be drawn from so precarious and incomplete an experiment.

But that invented by Drs. Hales and Defaguliers was of a more exact nature, depending on the pressure of the fluid only. For as the pressure of fluids in all directions is the same at the same depth, a Gage which discovers what the pressure is at the bottom of the sea, will shew what the true depth of the sea is in that place, whether the time of the machine's descent be longer or shorter.

Dr. Hales, in his Vegetable Statics, describes his Gage for estimating the pressures made in opaque vessels; where honey being poured over the surface of mercury in an open vessel, rises upon the surface of the mercury as it is pressed up into a tube whose lower orifice is immersed into the honey and mercury, and whose top is hermetically sealed. Now as by the pressure, the air in the tube is condensed, and the mercury rises, so the mercury comes down again when the pressure is taken off, and would leave no mark of the height to which it had risen; but the honey (or treacle, which does better) which is upon the mercury, sticking to the inside of the tube, leaves a mark, which shews the height to which it had risen, and consequently gives the quantity of pressure, and the height of the surface of the fluid.

Defaguliers's addition to this machine, consisted in a contrivance to carry it down to the bottom of the sea by means of a heavy weight, which was immediately disengaged by striking the bottom, and the Gage, made very light for the purpose, re-ascended to the top.

Dr. Hales afterwards made more experiments of this sort, and proposed another Sea Gage for vast depths, which is described in the Philos. Transf. N<sup>o</sup> 405, and is to this effect. Suppose a pretty long tube of copper or iron, close at the upper end, to be let down into the sea,



sea, to any depth, the water will rise in the tube to a height bearing a certain proportion to the depth of the sea to which the machine is sunk. And this proportion is as follows: 33 feet of sea water being nearly equal to the mean pressure of the atmosphere, therefore at 33 feet deep, the air in the tube will be compressed into half the length of the tube, or the water will rise and fill half way up the tube; in like manner at 66 feet deep, the water will occupy  $\frac{2}{3}$  of the tube; at 99 feet deep it will fill  $\frac{3}{4}$  of the tube; at 132 feet deep it will fill  $\frac{4}{5}$  of the tube; and so on. Hence therefore, by knowing the height to which the water rises in the tube, there will be known the consequent depth of the sea.

But, in very great depths, the scale near the top of the tube would be so small, and the divisions so close, that there would be no accuracy in the experiment, unless the tube were of a very great length, and this again would render it both liable to be broken, and quite impracticable.

To remedy this inconvenience, he made the following contrivance: To the bottom of the tube he screwed a large hollow globe of copper, with a small orifice, or a short pipe at bottom of the globe, to let in the water; by which means he had a very great quantity of air, and the scale enlarged. See also Defagul. Exp. Phil. vol. 2, p. 224 and 241.

*Bucket Sea GAGE*, is an instrument contrived by Dr. Hales to find the different degrees of coolness and saltness of the sea at different depths. This Gage consists of a common pale or bucket, with two heads: these heads have each a round hole in the middle, about 4 inches in diameter, covered with square valves opening upward; and that they may both open and shut together, there is a small iron rod, having one end fixed to the upper side of the lower valve, and the other end to the lower side of the upper valve. So that, as the bucket descends with its sinking weight into the sea, both the valves may open by the force of the water, which by that means has a free passage through the bucket. But when the bucket is drawn up, then both the valves shut by the force of the water at the upper end of the bucket; so that the bucket is drawn up full of the lowest sea water to which it has descended, and immediately the mercurial thermometer, fixed within it, is examined, to see the degree of temperature; and the degree of saltness is afterwards examined at leisure. Philof. Transf. numb. 9, p. 149, and numb. 24, p. 447, or Abridg. vol. 2, p. 260.

Lord Charles Cavendish adapted a thermometer for the temperature of the sea water, at different depths. See Philof. Transf. vol. 50, p. 300, and Phipps's Voyage towards the North Pole, p. 142 &c.

*Aqueo-mercurial GAGE* is the name of an apparatus contrived by Dr. Hales, and applied, in various forms, to the branches of trees, to determine the force with which they imbibe moisture. Vegetable Statics, vol. 1, ch. 2, p. 84.

*Sliding GAGE*, a tool used by mathematical instrument makers, for measuring and setting off distances; consisting of a beam, tooth, sliding socket, and the shoulder of the socket.

*Tide GAGE*, an instrument used for determining the height of the tides by Mr. Bayly, in the course of a

voyage towards the south pole &c, in the Resolution and Adventure, in the years 1772, 1773, 1774, and 1775. This instrument consists of a glass tube, whose internal diameter was  $\frac{7}{10}$ ths of an inch, lashed fast to a 10 foot fir rod, divided into feet, inches, and parts; the rod being fastened to a strong post fixed firm and upright in the water. At the lower end of the tube was an exceeding small aperture, through which the water was admitted. In consequence of this construction, the surface of the water in the tube was so little affected by the agitation of the sea, that its height was not altered the 10th part of an inch when the swell of the sea was 2 feet; and Mr. Bayly was certain, that with this instrument he could discern a difference of the 10th of an inch in the height of the tide.

*Water GAGE*. See ALTITUDE, and HYDROMETER.

*Wind GAGE*, an instrument for measuring the force of the wind upon any given surface. Several have been invented formerly, and one was lately invented by Dr. Lind, which is described in the Philof. Transf. vol. 65. See several also under the article ANEMOMETER.

GAGER, see GAUGER.

GAGING, see GAUGING.

*GALAXY*, or *Milky-Way*, or *Via Lactea*, in Astronomy, that long, whitish, luminous track, which seems to encompass the heavens like a swath, scarf, or girdle; and which is easily seen in a clear night, especially when the moon is not up. It is of a considerable, though unequal breadth; being also in some parts double, but in others single.

The Galaxy passes through many of the constellations in its circuit round the heavens, and keeps its exact place or position with respect to them.

There have been various strange and fabulous stories and opinions concerning the Galaxy.

The ancient poets, and even some of the philosophers, speak of it as the road or way by which the heroes went to heaven. But the Egyptians called it the Way of Straw, from the story of its rising from burning straw, thrown behind the goddess Isis in her flight from the giant Typhon. While the Greeks, who affect to derive every thing in the heavens from some of their own fables, have two origins for it; the one, that Juno, without perceiving it, accidentally gave suck to Mercury when an infant, but that as soon as she turned her eyes upon him, she threw him from her, and as the nipple was drawn from his mouth, the milk ran about for a moment; and the other, that the infant Hercules, being laid by the side of Juno when asleep, on waking she gave him the breast; but soon perceiving who it was, she threw him from her, and the heavens were marked by the wasted milk.

Some other philosophers however gave it a different turn, and different origin: these esteemed it to be a tract of liquid fire, spread in this manner along the skies: and others again, supposing a celestial region beyond all that was visible, and imagining that fire, at some time let loose from thence, was to consume the world, made this a part of that celestial fire, and appealed to it as a presage of what would surely happen. This diffused brightness they considered as a crack in the vault or wall of heaven, and fancied this a glimmering of the celestial fire through it, and that there required nothing more than the undoing of this crack.



crack by some accident in nature, or by the will of the Gods, to make the whole frame start, and let out the fire of destruction.

Aristotle makes the Galaxy a kind of meteor, formed of a crowd of vapours, drawn into that part by certain large stars disposed in the region of the heavens answering to it. Others, finding that the Galaxy was seen all over the globe, that it always corresponded to the same fixed stars, and that it was far above the highest planets, set Aristotle's opinion aside, and placed the Galaxy in the firmament or region of the fixed stars; and concluded that it was nothing else but an assemblage of an infinite number of minute stars. And since the invention of telescopes, this opinion has been abundantly confirmed. For, by directing a good telescope to any part of the milky way, we perceive an innumerable multitude of very small stars, where before we only observed a confused whiteness, arising from the assemblage and union of their joint light; like as any thing powdered with fine white powder, at a distance we only observe the confused whiteness, but on examining it very near we perceive all the small particles of the powder separately; as Milton finely expresses it,

A broad and ample road, whose dust is gold,  
And pavement stars, as stars to thee appear,  
Seen in the Galaxy, that milky way,  
Which nightly, as a circling Zone thou see'st.  
Powder'd with stars.

There are other such marks in the heavens; as the nebulæ, or, nebulous stars, and certain whitish parts about the south pole, called Magellanic clouds, which are all of the same nature, appearing to be vast clusters of small stars when viewed through a telescope, which are too faint to affect the eye singly.

M. le Monnier however, not being able to discover more stars in this space than in other parts of the heavens, disputes the opinion above recited as to the reason of the whiteness, and supposes that this and the nebulous stars are occasioned by some other kind of matter. *Inst. Ast.* p. 60.

GALILEI (GALILEO,) a most excellent philosopher, mathematician and astronomer, was born at Pisa in Italy, in 1564. From his infancy he had a strong propensity to philosophy and mathematics, and soon made a great progress in these sciences. So that in 1592 he was chosen professor of mathematics at Padua. While he was professor there, visiting Venice, then famous for the art of glass-making, he heard that in Holland a glass had been invented, through which very distant objects were seen distinctly as if near at hand. This was sufficient for Galileo; his curiosity was raised, and put him upon considering what must be the form of such a glass, and the manner of making it. The result of his enquiry was the invention of the telescope, produced from this hint, without having seen the Dutch glass. All the discoveries he made in astronomy were easy and natural consequences of this invention, which opening a way, till then unknown, into the heavens, thence brought the finest discoveries. One of the first of these, was that of 4 of Jupiter's satellites, which he called the Medicean stars or planets, in honour of Cosmo the 2d, grand-duke of Tuscany, who was of

that family. Cosmo sent for our astronomer from Padua, and made him professor of mathematics at Pisa in 1611; and soon after inviting him to Florence, gave him the office and title of *principal philosopher and mathematician to his highness*.

He had been but a few years at Florence, before the Inquisition began to be very busy with him. Having observed some solar spots in 1612, he printed that discovery the following year at Rome; in which, and in some other pieces, he ventured to assert the truth of the Copernican system, and brought several new arguments to confirm it. For these he was cited before the Inquisition at Rome, in 1615: after some months imprisonment, he was released, and sentence pronounced against him, that he should renounce his heretical opinions, and not defend them by word or writing, or insinuate them into the minds of any persons. But having afterwards, in 1632, published at Florence his *Dialogues of the two Great Systems of the World, the Ptolomaic and Copernican*, he was again cited before the holy-office, and committed to the prison of that ecclesiastical court at Rome. The inquisitors convened in June that year; and in his presence pronounced sentence against him and his books, obliging him to abjure his errors in the most solemn manner; committed him to the prison of their office during pleasure; and enjoined him, as a saving penance, for three years to come, to repeat once a week the seven penitential psalms: reserving to themselves, however, the power of moderating, changing, or taking away altogether or in part, the said punishment and penance. On this sentence, he was detained in prison till 1634; and his *Dialogues of the System of the World* were burnt at Rome.

Galileo lived ten years after this; seven of which were employed in making still further discoveries with his telescope. But by the continual application to that instrument, added to the damage his sight received from the nocturnal air, his eyes grew gradually weaker, till he became totally blind in 1639. He bore this calamity with patience and resignation, worthy of a great philosopher. The loss neither broke his spirit, nor stopped the course of his studies. He supplied the defect by constant meditation; by which means he prepared a large quantity of materials, and began to arrange them by dictating his ideas; when, by a distemper of three months continuance, wasting away by degrees, he expired at Arcetri near Florence, in January 1642, being the 78th year of his age.

Galileo was in his person of small stature, though of a venerable aspect, and vigorous constitution. His conversation was affable and free, and full of pleasantry. He took great delight in architecture and painting, and designed extremely well. He played exquisitely on the lute; and whenever he spent any time in the country, he took great pleasure in husbandry. His learning was very extensive; and he possessed in a high degree a clearness and acuteness of wit. From the time of Archimedes, nothing had been done in mechanical geometry, till Galileo, who being possessed of an excellent judgment, and great skill in the most abstruse points of geometry, first extended the boundaries of that science, and began to reduce the resistance of solid bodies.



bodies to its laws. Besides applying geometry to the doctrine of motion, by which philosophy became established on a sure foundation, he made surprising discoveries in the heavens by means of his telescope. He made the evidence of the Copernican system more sensible, when he shewed from the phases of Venus, like to those of the moon, that Venus actually revolves about the sun. He proved the rotation of the sun on his axis, from his spots; and thence the diurnal rotation of the earth became more credible. The satellites that attend Jupiter in his revolution about the sun, represented, in Jupiter's smaller system, a just image of the great solar system; and rendered it more easy to conceive how the moon might attend the earth, as a satellite, in her annual revolution. By discovering hills and cavities in the moon, and spots in the sun constantly varying, he shewed that there was not so great a difference between the celestial bodies and the earth as had been vainly imagined.

He rendered no less service to science by treating, in a clear and geometrical manner, the doctrine of motion, which has justly been called the key of nature. The rational part of mechanics had been so much neglected, that hardly any improvement was made in it for almost 2000 years. But Galileo has given us fully the theory of equable motions, and of such as are uniformly accelerated or retarded, and of these two compounded together. He, first of any, demonstrated that the spaces described by heavy bodies, from the beginning of their descent, are as the squares of the times; and that a body, projected in any direction not perpendicular to the horizon, describes a parabola. These were the beginnings of the doctrine of the motion of heavy bodies, which has been since carried to so great a height by Newton. In geometry, he invented the cycloid, or trochoid; though the properties of it were afterwards chiefly demonstrated by his pupil Torricelli. He invented the simple pendulum, and made use of it in his astronomical experiments: he had also thoughts of applying it to clocks; but did not execute that design: the glory of that invention was reserved for his son Vicenzio, who made the experiment at Venice in 1649; and Huygens afterward carried this invention to perfection. Of Galileo's invention also, was the machine, with which the Venetians render their Laguna fluid and navigable. He also discovered the gravity of the air, and endeavoured to compare it with that of water; besides, opening up several other enquiries in natural philosophy. In short, he was not esteemed and followed by philosophers only, but was honoured by persons of the greatest distinction of all nations.

Galileo had scholars too that were worthy of so great a master, by whom the gravitation of the atmosphere was fully established, and its varying pressure accurately and conveniently measured, by the column of quicksilver of equal weight sustained by it in the barometrical tube. The elasticity of the air, by which it perpetually endeavours to expand itself, and, while it admits of condensation, resists in proportion to its density, was a phenomenon of a new kind (the common fluids having no such property), and was of the utmost importance to philosophy. These principles opened a vast field of new and useful knowledge, and explained a

great variety of phenomena, which had been accounted for before that time in a very absurd manner. It seemed as if the air, the fluid in which men lived from the beginning, had been then but first discovered. Philosophers were every where busy enquiring into its various properties and their effects: and valuable discoveries rewarded their industry. Of the great number who distinguished themselves on this occasion, may be mentioned Torricelli and Viviani in Italy, Pascal in France, Otto Guerick in Germany, and Boyle in England.

Galileo wrote a number of treatises, many of which were published in his life-time. Most of them were also collected after his death, and published by Mendessi in 2 vols 4to, under the title of *L'Opere di Galileo Galilei Lynceo*, in 1656. Some of these, with others of his pieces, were translated into English and published by Thomas Salisbury, in his Mathematical Collections, in 2 vols folio. A volume also of his letters to several learned men, and solutions of several problems, were printed at Bologna in 4to. His last disciple, Vincenzo Viviani, who proved a very eminent mathematician, methodized a piece of his master's, and published it under this title, *Quinto libro de gli Elementi d'Euclidi*, &c.; at Florence in 1674, 4to. Viviani published some more of Galileo's things, being Extracts from his letters to a learned Frenchman, where he gives an account of the works which he intended to have published, and a passage from a letter of Galileo dated at Arcetri, Oct. 30, 1635, to John Camillo, a mathematician of Naples, concerning the angle of contact. Besides all these, he wrote many other pieces, which were unfortunately lost through his wife's devotion; who, solicited by her confessor, gave him leave to peruse her husband's manuscripts; of which he tore and took away as many as he said were not fit to be published.

**GALLERY**, in Architecture, a covered place in a building, much longer than broad; which is usually placed in the wings of the building, and serving to walk in, and to place pictures in. It denotes a little aisle, or walk, serving as a common passage to several rooms placed in a line, or row.

**GALLERY**, in Fortification, a covered walk, or passage, made across the ditch of a besieged town, with timbers fastened in the ground and covered over.

**GALLON**, an English measure of capacity, for things both liquid and dry, containing 2 pottles, or 4 quarts, or 8 pints. But those pints and quarts, and consequently the Gallon itself, are different, according to the quality of the things measured: the wine Gallon, for instance, contains 231 cubic inches, and holds 8lb 5½ oz, avoirdupois, of pure water; the beer and ale Gallon contains 282 cubic inches, and holds 10lb 3¼ oz of water; and the Gallon dry measure, for grain, meal, &c, contains 268½ cubic inches, and holds 9lb 11½ oz of water.

**GALLOPER**, in Artillery, the name of a carriage serving for the very small guns, and having shafts so as to be drawn without a limber.

**GAMING**. See CHANCES, and LAWS of CHANCE.

**GARDECAUT**, or GUARD DU CORD, in a watch, is that which stops the fusee, when wound up, and for that end is driven up by the spring. Some call it Guard-cock; others Guard du Gut.

**GAR-**



**GARRISON GUNS**, such as are mounted and used in a Garrison, consisting of the following weights, viz. the 42, 32, 24, 18, 12, 9, and 6 pounders; being made either of brass or iron.

*Table of the Weight and Dimensions of Garrison Guns.*

Brass Garrison Guns.			Iron Garrison Guns.		
Shot	Length	Weight.	Shot	Length	Weight
lb	f. in.	Cw. qr. lb.	lb.	f. in.	Cw. qr. lb.
42	10 0	64 0 0	32	9 8	56 0 0
32	9 2	49 2 18	24	9 8	48 0 0
24	8 4	37 0 0	18	9 0	36 0 0
18	7 6	27 3 0	12	7 8	24 0 0
12	6 7	18 2 0	9	7 0	18 0 0
9	6 0	13 3 0	6	6 1	12 0 0
6	5 3	9 1 0	4	5 4	8 0 0

**GASSENDI (PETER)**, one of the most celebrated philosophers France has produced, was born at Chanterrier, about 3 miles from Digne in Provence, in the year 1592. When a child, he took great delight in gazing at the moon and stars whenever they appeared. This pleasure often drew him into bye-places, that he might feast his eyes freely and undisturbed; by which means his parents had him often to seek, not without many anxious fears and apprehensions. In consequence of this promising disposition, he was sent to the best schools, to cultivate it with the instructions of the first masters. He profited so well of these aids, that he was invited to be professor of rhetoric at Digne, before he was quite 16 years of age. After filling this office three years, upon the death of his master at Aix, he was appointed to succeed him as professor of philosophy. After a few years residence here, he composed his *Paradoxical Exercitations*; which coming to the hands of Nicholas Peiresc, that great patron of learning joined with Joseph Walter, prior of Valette, in promoting him; and, having entered into holy orders, he was first made canon of the church of Digne and doctor of divinity, and then warden or rector of the same church.

Gassendi's fondness for astronomy grew up with his years; and his reputation daily increasing, he was appointed the king's professor of mathematics at Paris in 1645. This institution being chiefly intended for astronomy, our author read lectures on that science to a crowded audience. However, he did not long enjoy this situation; for a dangerous cough and inflammation of the lungs obliged him, in 1647, to return to Digne for the benefit of his native air. Having thus, and by the intermission of his studies, recovered his health, he again returned to Paris in 1653; where, after first writing and publishing the lives of Tycho Brahe, Copernicus, Purbach, and Regiomontanus, in 1654 he again renewed his astronomical labours, with the design of completing the system of the heavens.

But while he was thus employed, too intensely for the feeble state of his health, he relapsed into his former disorder, under which, with the aid of too copious and numerous bleedings, by order of three physicians, he sunk in the year 1655, at 63 years of age.

Gassendi wrote against the metaphysical meditations of Des Cartes; and divided with that great man the philosophers of his time, almost all of whom were either Cartesians or Gassendists. To his knowledge in philosophy and mathematics, he joined profound erudition and deep skill in the languages. He wrote, 1. Three volumes on Epicurus's philosophy; and six others, which contain his own philosophy.—2. *Astronomical Works*.—3. The lives of Nicholas de Peiresc, Epicurus, Copernicus, Tycho Brahe, Purbach, and Regiomontanus.—4. *Epistles*, and other treatises. All his works were collected together, and printed at Lyons in 1658, in 6 volumes folio.

Gassendi was the first person that saw the transit of Mercury over the sun, viz. Nov. 7, 1631; as Horrox first predicted and shewed the transit of Venus.—His library was large and valuable: to which he added an astronomical and philosophical apparatus, which, for their accuracy and magnitude, were purchased by the emperor Ferdinand the 3d.—It appears by his letters, printed in the 6th volume of his works, that he was often consulted by the most celebrated astronomers of his time, as Kepler, Longomontanus, Snell, Hevelius, Galileo, Kircher, Bulliald, and others: and he has generally been esteemed one of the founders of the reformed philosophy, in opposition to the groundless hypotheses and empty subtleties of Aristotle and the schoolmen.

**GAUGE-Line**, a line on the common Gauging rod, used for the purpose of gauging liquids. See *GAUGING-Rod*.

**GAUGE-Point**, of a solid measure, is the diameter of a circle, whose area is expressed by the same number as the solid content of that measure. Or it is the diameter of a cylinder, whose altitude is 1, and its content the same as of that measure.

Thus, the solid content of a wine gallon being 231 cubic inches; if a circle be conceived to contain so many square inches, its diameter will be 17.15; which is therefore the Gauge-point for wine measure. And an ale gallon containing 282 cubic inches; by the same rule, the Gauge-point for ale measure will be found to be 18.95. And after the same manner may the Gauge-point for any other measure be determined.

Hence it follows, that when the diameter of a cylinder in inches is equal to the Gauge-point in any measure, given likewise in inches, every inch in its length will contain an integer of the same measure. So in a cylinder whose diameter is 17.15 inches, every inch in height contains one entire gallon in wine measure; and in another, whose diameter is 18.95, every inch in length contains one ale gallon.

**GAUGER**, an officer appointed by the commissioners of excise, to Gauge, measure, or examine, all casks, tuns, pipes, barrels, hogheads, of beer, wine, oil, &c.

**GAUGING**, the art or act of measuring the capacities or contents of all kinds of vessels, and determining the quantity of fluids, or other matters contained in them. These are principally pipes, tuns, barrels, rundlets, and other casks; also backs, coolers, vats, &c.



As to the solid contents of all prismatical vessels, as cubes, parallelopipedons, cylinders, &c, they are found by multiplying the area of the base by their altitude. And the contents of all pyramidal bodies, and cones, are equal to 1-3d of the same.

In Gauging, it has been usual to divide casks into four varieties or forms, denominated as follows, from the supposed resemblance they bear to the frustums of solids of the same names: viz,

1. The middle frustum of a spheroid,
2. The middle frustum of a parabolic spindle,
3. The two equal frustums of a paraboloid,
4. The two equal frustums of a cone.

And particular rules, adapted to each of these forms, may be found in most books of Gauging, and in my *Mensuration*, p. 575 &c. But as the form is imaginary, and only guessed at, it hardly ever happens that a true solution is brought out in this way; beside which, it is very troublesome and inconvenient to have so many rules to put in practice. I shall therefore give here one rule only, from p. 592 of that book, which is not only general for all casks that are commonly met with, but quite easy, and very accurate, as having been often verified and proved by filling the casks with a true gallon measure.

*General Rule.* Add into one sum, 39 times the square of the bung diameter, 25 times the square of the head diameter, and 26 times the product of those diameters; multiply the sum by the length of the cask, and the product by the number .00034; then this last product divided by 9 will give the wine gallons, and divided by 11 will give the ale gallons.

Or,  $39B^2 + 25H^2 + 26BH \times \frac{L}{114}$  is the content in inches; which being divided by 231 for wine gallons, or by 282 for ale gallons, will be the content.

*For Ex.* If the length of a cask be 40 inches, the bung diameter 32, and the head diameter 24.

Here	-	$32^2 \times 39$	=	39936
and	-	$24^2 \times 25$	=	14400
and	-	$32 \times 24 \times 26$	=	19968

the sum	-	74304
multiplied by	-	40

and divid. by 114) 2972160

gives - 26071 cubic inches;

this divided by 231 gives 112 wine gallons,

or divided by 282 gives 92 ale gallons.

But the common practice of Gauging is performed mechanically, by means of the Gauging or Diagonal Rod, or the Gauging Sliding Rule, the description and use of which here follow.

GAUGING, or *Diagonal Rod*, is a rod or rule adapted for determining the contents of casks, by measuring the diagonal only, viz the diagonal from the bung to the extremity of the opposite stave next the head. It is a square rule, having 4 sides or faces, being usually 4 feet long, and folding together by means of joints.

Upon one face of the rule is a scale of inches, for taking the measure of the diagonal; to these are adapted the areas, in ale gallons, of circles to the corresponding diameters, like the lines on the under sides

of the three slides in the sliding rule, described below. And upon the opposite face are two scales, of ale and wine gallons, expressing the contents of casks having the corresponding diagonals; and these are the lines which chiefly constitute the difference between this instrument and the sliding rule; for all the other lines upon it are the same with those in that instrument, and are to be used in the same manner.

*To use the Diagonal Rod.* Unfold the rod straight out, and put it in at the bung hole of the cask to be gauged, till its end arrive at the intersection of the head and opposite stave, or to the farthest possible distance from the bung-hole, and note the inches and parts cut by the middle of the bung; then draw out the rod, and look for the same inches and parts on the opposite face of it, and annexed to them are found the contents of the cask, both in ale and wine gallons.

*For Ex.* Let it be required to find, by this rod, the content of a cask whose diagonal measures 34.4 inches; which answers to the cask in the foregoing example, whose head and bung diameters are 32 and 24, and length 40 inches; for if to the square of 20, half the length, be added the square of 28, half the sum of the diameters, the square root of the sum will be 34.4 nearly.

Now, to this diagonal 34.4, corresponds, upon the rule, the content 91 ale gallons, or 111 wine gallons; which are but 1 less than the content brought out by the former general rule above given.

GAUGING Rule, or *Sliding Rule*, is a sliding rule particularly adapted to the purposes of Gauging. It is a square rule, of four faces or sides, three of which are furnished with sliding pieces running in grooves. The lines upon them are mostly logarithmic ones, or distances which are proportional to the logarithms of the numbers placed at the ends of them; which kind of lines was placed upon rulers, by Mr. Edmund Gunter, for expeditiously performing arithmetical operations, using a pair of compasses for taking off and applying the several logarithmic distances: but instead of the compasses, sliding pieces were added, by Mr. Thomas Everard, as more certain and convenient in practice, from whom this sliding rule is often called *Everard's Rule*. For the more particular description and uses of this rule, see my *Mensuration*, p. 564, 2d edition.

The writers on Gauging are, Beyer, Kepler, Dechaes, Hunt, Everard, Dougherty, Shettleworth, Shirtcliffe, Leadbetter, &c.

GAZONS, in Fortification, turfs, or pieces of fresh earth covered with grass, cut in form of a wedge, about a foot long, and half a foot thick, to line or face the outside of works made of earth, to keep them up, and prevent their mouldering.

GELLIBRAND (HENRY), professor of astronomy at Gresham-college, was born in London the 27th of Nov. 1597. He was sent to Trinity-college, Oxford, in 1615, and took his degree in arts 1619. He then entered into orders, and became curate of Chiddingstone in Kent. Afterwards, taking a great fancy to mathematics, by happening to hear one of Sir Henry Saville's lectures in that science, he immediately set himself to the close study of that noble science, and relinquished his fair prospects in the church. Contenting himself there-fore



fore with his private patrimony, which was now come into his hands by the death of his father, the same year he entered again a student at Oxford, making mathematics his sole employment. He made such proficiency in this science before he proceeded A. M., which was in 1623, that he drew the attention and intimate friendship of Mr. Henry Briggs, then lately removed from the geometry professorship in Gresham-college to that of Savilian professor of geometry at Oxford, by the founder Sir Henry Savile, and who, upon the death of Mr. Gunter, procured for our author the professorship of astronomy in Gresham-college, to which he was elected in the beginning of the year 1627. His friend Mr. Briggs dying in 1630, before he had finished the introduction to his *Trigonometria Britannica*, he recommended the completing and publishing of that work to our author. Gellibrand accordingly added a preface, and the application of the logarithms to plane and spherical trigonometry, &c, and the whole was printed at Gouda, under the care of Adrian Vlacq, in 1633.

While Mr. Gellibrand was preparing that work, he was brought into trouble in the high-commission court, by Dr. Laud, then bishop of London, on account of an almanac, published by William Beale, servant to Mr. Gellibrand, for the year 1631, with the approbation of his master. In this almanac, the popish saints, then usually put into calendars, were omitted, and the names of other saints and martyrs, mentioned in the Book of Martyrs, were placed in their stead, as they stand in Fox's calendars. This it seems gave offence to the bishop, and occasioned the prosecution. But when the cause came to be heard, it appeared that other almanacs of the same kind had formerly been printed; upon which, both master and man were acquitted by Abp. Abbot and the whole court, Laud only excepted; which was afterward made one of the articles against him on his own trial.

It seems Gellibrand was strongly attached to the old Ptolomaic system. For when he went over to Holland, about the printing of Briggs's book abovementioned, he had some discourse with Lansberg, an eminent brother astronomer in Zealand, who affirming that he was fully persuaded of the truth of the Copernican system; our author observes, "that this so styled a truth he should receive as an hypothesis; and so be easily led on to the consideration of the imbecility of man's apprehension, as not able rightly to conceive of this admirable opifice of God, or frame of the world, without falling foul of so great an absurdity:" so firmly was he fixed in his adherence to the Ptolomaic system. Gellibrand wrote several things after this, chiefly tending to the improvement of navigation, which would probably have been further advanced by him, had his life been continued longer; but he was untimely carried off by a fever, in 1636, at 39 years of age.

The character of Mr. Gellibrand is that of a plain, plodding, industrious, well-intentioned man, with little invention or genius. His writings are chiefly as below:

1. *Trigonometria Britannica*; or the Doctrine of Triangles, being the 2d part of Briggs's work abovementioned.

2. A small Treatise concerning the longitude.

VOL. I.

3. A Discourse on the *Variation of the Magnetic Needle*; annexed to Wright's Errors in Navigation detected.

4. *Institution Trigonometrical*, with its application to astronomy and navigation; 8vo, 1635.

5. *Epitome of Navigation*, with the necessary tables; 8vo.

6. Several manuscripts never published; as, The Doctrine of Eclipses.—A Treatise of Lunar Astronomy.—A Treatise of Ship-building, &c.

GEMINI, a constellation of the northern hemisphere, one of the 48 old constellations, and the 3d in order of the zodiacal signs, Aries, Taurus, Gemini, &c. This constellation consists of two children, twins, called Castor and Pollux, and denoted by the mark,  $\Pi$ , being a rude drawing of the same.

This constellation was, more anciently, depicted by a couple of young kids, by the Egyptians and eastern nations, as denoting that part of the spring when these animals appear; but the Greeks altered them to two children, which some of them make to be Castor and Pollux, some of them again Hercules and Apollo, and others Triptolemus and Jason; but the Arabians afterwards changed the figures into two peacocks, their religion not allowing them to paint or draw any human figure. Sir Isaac Newton thinks the figures had some reference to the Argonautic expedition.

The ancients attributed to every sign of the zodiac one of the principal deities for its tutelary power. Phœbus had the care of Gemini, and hence all the jargon of astrologers about the agreement of the sun and this constellation.

The stars in the sign Gemini are, in Ptolomy's catalogue 25, in Tycho's 25, in Hevelius's 38, and in the Britannic catalogue 85.

GENERATED, is used by some mathematical writers for whatever is produced by arithmetical operation, or in geometry by the motion of other magnitudes. Thus 20 is the product Generated of 4 and 5; *ab* that of *a* and *b*, 4, 8, 16, &c, the powers generated of or from the root 2, and  $a^2, a^3, a^4$ , &c, those from the root *a*. So also, a circle is Generated by the revolution of a line about one of its extremities, a cone by the rotation of a right-angled triangle about its perpendicular, a cylinder by the rotation of a rectangle about one of its sides, or, otherwise, by the motion of a circle in the direction of a right line, and keeping always parallel to itself.

GENERATING *Line* or *Figure*, in Geometry, is that which, by any kind of supposed motion, may generate, or produce, any other figure, plane, or solid.

Thus a line, according to Euclid, generates a circle; or a right-angled triangle, a cone; &c; and thus also Archimedes supposes his spirals to be generated by the motions of Generating points and lines; the figure thus generated, is called the *Generant*.

It is a general theorem in geometry, that the measure of any generant, or figure produced by any kind of motion of any other figure, or Generating quantity, is equal to the product of this Generating quantity drawn into the length of the path described by its centre of gravity, whatever the kind of motion may be, whether rotatory, or direct, &c.



In the modern analysis, or fluxions, all sorts of quantities are considered as Generated by some such motion, and the quantity hereby generated is called a *Fluent*.

**GENERATION**, in Mathematics, is used for the formation or production of any geometrical figure, or other quantities. Such as of the figures mentioned in the foregoing articles, or the Generation of equations, curves, solids, &c.

**GENESIS**, in Geometry, means much the same as Generation mentioned above, being the formation of a line, surface, or solid, by the motion or flux of a point, line, or surface; as of a globe by the rotation of a semi-circle about its diameter, &c.

In the Genesis of figures, the line or surface which moves, is called the *Describent*; and the line round which, or according to which, the revolution or other motion is made, the *Dirigent*.

**GENETHLIACI**, in Astrology, are persons who erect horoscopes, or pretend to foretell what shall befall a person, by means of the stars which presided at his nativity.

The ancients called them Chaldaei, and by the general name Mathematici: accordingly, the several civil and canon laws, which we find made against the mathematicians, only respect the Genethliaci, or astrologers.

They were expelled Rome, by a formal decree of the senate; and yet found so much protection from the credulity of the people, that they remained in the city unmolested.

Antipater and Archinapolus have shewn that Genethliology should rather be founded on the time of the conception than on that of the birth.

**GEOCENTRIC**, is said of a planet or its orbit, to denote its having the earth for its centre. The moon alone is properly geocentric. And yet the motions of all the planets may be considered in respect of the earth, or as they appear from the earth, and thence called their Geocentric motions.—Hence also the terms Geocentric place, or latitude, or longitude, &c, as explained below.

**GEOCENTRIC Place**, of a planet, is the place where it appears to us, from the earth; or it is a point in the ecliptic, to which a planet, seen from the earth, is referred.

**GEOCENTRIC Latitude**, of a planet, is its latitude as seen from the earth; or the inclination of a line, connecting the planet and the earth, to the plane of the earth's (or true) ecliptic. Or it is the angle which the said line (connecting the planet and the earth) makes with a line drawn to meet a perpendicular let fall from the planet to the plane of the ecliptic.

**GEOCENTRIC Longitude**, of a planet, is the distance measured on the ecliptic, in the order of the signs, between the Geocentric place and the first point of Aries.

**GEODESIA**, is properly that part of practical geometry that teaches how to divide or lay out lands and fields, among several owners.

**GEODESIA** is also applied, by some writers, to all measurements in the field, and as synonymous with surveying.

**GEODESIA** is defined by Vitalis, as the art of mea-

suring surfaces and solids, not by imaginary lines, as is done in geometry, but by sensible and visible things; or by the sun's rays, &c.

**GEOGRAPHER**, a person skilled in Geography.

**GEOGRAPHICAL**, something relating to Geography, as

**GEOGRAPHICAL Mile**, which is the sea-mile or minute, being the 60th part of a degree of a great circle.

**GEOGRAPHICAL Table**. See MAP.

**GEOGRAPHY**, the science that teaches and explains the nature and properties of the earth, as to its figure, place, magnitude, motions, celestial appearances, &c, with the various lines, real or imaginary, on its surface.

Geography is distinguished from Cosmography, as a part from the whole; this latter considering the whole visible world, both heaven and earth. And from Topography and Chorography, it is distinguished, as the whole from a part.

Golnitz considers Geography as either exterior or interior: but Varenus more justly divides it into General and Special; or Universal and Particular.

**General or Universal GEOGRAPHY**, is that which considers the earth in general, without any regard to particular countries, or the affections common to the whole globe: as its figure, magnitude, motion, land, sea, &c.

**Special or Particular GEOGRAPHY**, is that which contemplates the constitution of the several particular regions, or countries; their bounds, figure, climate, seasons, weather, inhabitants, arts, customs, language, &c.

**History of GEOGRAPHY**. The study and practice of Geography must have commenced at very early ages of the world. By the accounts we have remaining, it seems this science was in use among the Babylonians and Egyptians, from whom it passed to the Greeks first of any Europeans, and from these successively to the Romans, the Arabians, and the western nations of Europe. Herodotus says the Greeks first learned the pole, the gnomon, and the 12 divisions of the day, from the Babylonians. But Pliny and Diogenes Laertius assert, that Thales of Miletus, in the 6th century before Christ, first found out the passage of the sun from tropic to tropic, and it is said was the author of two books, the one on the tropic, and the other on the equinox; both probably determined by means of the gnomon; whence he was led to the discovery of the four seasons of the year, which are determined by the equinoxes and solstices; all which however it is likely he learned of the Egyptians, as well as his division of the year into 365 days. This it is said was invented by the second Mercury, surnamed Trismegistus, who, according to Eusebius, lived about 50 years after the Exodus. Pliny expressly says that this discovery was made by observing when the shadow returned to its marks; a clear proof that it was done by the gnomon. It is farther said that Thales constructed a globe, and represented the land and sea upon a table of brass. Farther that Anaximander, a disciple of Thales, first drew the figure of the earth upon a globe; and that Hecate, Democritus, Eudoxus, and others, formed Geographical



phical maps, and brought them into common use in Greece.

Meton and Euctemon observed the summer solstice at Athens, on the 27th of June 432 years before Christ, by watching narrowly the shadow of the gnomon, with the design of fixing the beginning of their cycle of 19 years.

Timocharis and Aristillus, who began their observations about 295 B. C., it seems first attempted to fix the latitudes and longitudes of the fixed stars, by considering their distances from the equator, &c. One of their observations gave rise to the discovery of the precession of the equinoxes, which was first remarked by Hipparchus about 150 years after: who also made use of their method, for delineating the parallels of latitude and the meridians, on the surface of the earth; thus laying the foundation of this science as it now appears.

The latitudes and longitudes, thus introduced by Hipparchus, were not however much attended to till Ptolemy's time. Strabo, Vitruvius, and Pliny, have all of them entered into a minute geographical description of the situation of places, according to the length of the shadows of the gnomon, without noticing the longitudes and latitudes.

Maps at first were little more than rude outlines, and topographical sketches of different countries. The earliest on record were those of Sesostris, mentioned by Eustathius; who says, that "this Egyptian king, having traversed great part of the earth, recorded his march in maps, and gave copies of them not only to the Egyptians, but to the Scythians, to their great astonishment." Some have imagined with much probability, that the Jews made a map of the Holy Land, when they gave the different portions to the nine tribes at Shiloh: for Joshua tells us that they were sent to walk through the land, and that they *described it in seven parts in a book*; and Josephus relates that when Joshua sent out people from the different tribes to measure the land, he gave them, as companions, persons well skilled in geometry, who could not be mistaken in the truth.

The first Grecian map on record, was that of Anaximander, mentioned by Strabo, lib. 1, p. 7, supposed to be the one referred to by Hipparchus under the designation of the *ancient map*. Herodotus minutely describes a map made by Aristagoras tyrant of Miletus, which will serve to give some idea of the maps of those times. He relates, that Aristagoras shewed it to Cleomenes king of Sparta, to induce him to attack the king of Persia at Susa, in order to restore the Ionians to their ancient liberty. It was traced upon brass or copper, and seems to have been a mere itinerary, containing the route through the intermediate countries which were to be traversed in that march, with the rivers Halys, the Euphrates, and Tigris, which Herodotus mentions as necessary to be crossed in that expedition. It contained one straight line called the *Royal Road* or *Highway*, which took in all the stations or places of encampment from Sardis to Susa; being 111 in the whole journey, and containing 13,500 stadia, or 1687½ Roman miles of 5000 feet each.

These itinerary maps of the places of encampment were indispensably necessary in all armies and marches;

and indeed war and navigation seem to be the two grand causes of the improvements both in Geography and astronomy. Athenæus quotes Bæton as author of a work intitled, *The encampments of Alexander's march*; and likewise Amyntas to the same purpose. Pliny observes that Diognetus and Bæton were the surveyors of Alexander's marches, and then quotes the exact number of miles according to their mensuration; which he afterwards confirms by the letters of Alexander himself. The same author also remarks that a copy of this great monarch's surveys was given by Xenocles his treasurer to Patrocles the geographer, who was admiral of the fleets of Seleucus and Antiochus. His book on geography is often quoted both by Strabo and Pliny; and it seems that this author furnished Eratosthenes with the principal materials for constructing his map of the oriental part of the world.

Eratosthenes first attempted to reduce Geography to a regular system, and introduced a regular parallel of latitude, which began at the straits of Gibraltar, passed eastwards through the isle of Rhodes, and so on to the mountains of India, noting all the intermediate places through which it passed. In drawing this line, he was not regulated by the same latitude, but by observing where the longest day was 14 hours and a half, which Hipparchus afterwards determined was the latitude of 36 degrees.

This first parallel through Rhodes was ever after considered with a degree of preference, in constructing all the ancient maps; and the longitude of the then known world was often attempted to be measured in stadia and miles, according to the extent of that line, by many succeeding geographers.

Eratosthenes soon after attempted not only to draw other parallels of latitude, but also to trace a meridian at right angles to these, passing through Rhodes and Alexandria, down to Syene and Meroë; and at length he undertook the arduous task of determining the circumference of the globe, by an actual measurement of a segment of one of its great circles. To find the magnitude of the earth, is indeed a problem which has engaged the attention of astronomers and geographers ever since the spherical figure of it was known. It seems Anaximander was the first among the Greeks who wrote upon this subject. Archytas of Tarentum, a Pythagorean, famous for his skill in mathematics and mechanics, also made some attempts in this way; and Dr. Long conjectures that these are the authors of the most ancient opinion that the circumference of the earth is 400,000 stadia: and Archimedes makes mention of the ancients who estimated the circumference of the earth at only 30,000 stadia.

As to the methods of measuring the circumference of the earth, it would seem, from what Aristotle says in his treatise *De Cælo*, that they were much the same as those used by the moderns, deficient only in the accuracy of the instruments. That philosopher there says, that different stars pass through our zenith, according as our situation is more or less northerly; and that in the southern parts of the earth stars come above our horizon, which are no longer visible if we go northward. Hence it appears that there are two ways of measuring the circumference of the earth; one by observing stars which pass through the zenith of one place, and do



not pass through that of another; the other, by observing some stars which come above the horizon of one place, and are observed at the same time to be in the horizon of another. The former of these methods, which is the best, was followed by Eratosthenes at Alexandria in Egypt, 250 years before Christ. He knew that at the summer solstice, the sun was vertical to the inhabitants of Syene, a town on the confines of Ethiopia, under the tropic of cancer, where they had a well made to observe it, at the bottom of which the rays of the sun fell perpendicularly the day of the summer solstice: he observed by the shadow of a wire set perpendicularly in an hemispherical basin, how far the sun was on that day at noon distant from the zenith of Alexandria; when he found that distance was equal to the 50th part of a great circle in the heavens. Then supposing Syene and Alexandria under the same meridian, he inferred that the distance between them was the 50th part of a great circle upon the earth; and this distance being by measure 5000 stadia, he concluded that the whole circumference of the earth was 250,000 stadia. But as this number divided by 360 would give  $694\frac{2}{3}$  stadia to a degree, either Eratosthenes himself or some of his followers assigned the round number 700 stadia to a degree; which multiplied by 360, makes the circumference of the earth 252,000 stadia; whence both these measures are given by different authors, as that of Eratosthenes.

In the time of Pompey the Great, Posidonius determined the measure of the circumference of the earth by the 2d method above hinted by Aristotle, viz, the horizontal observations. Knowing that the star called Canopus was but just visible in the horizon of Rhodes, and at Alexandria finding its meridian height was the 48th part of a great circle in the heavens, or  $7\frac{1}{2}$  deg., answering to the like quantity of a circle on the earth: Then, supposing these two places under the same meridian, and the distance between them 5000 stadia, the circumference of the earth will be 240,000 stadia; which is the first measure of Posidonius. But according to Strabo, Posidonius made the measure of the earth to be 180,000 stadia, at the rate of 500 stadia to a degree. The reason of this difference is thought to be, that Eratosthenes measured the distance between Rhodes and Alexandria, and found it only 3750 stadia: taking this for a 48th part of the earth's circumference, which is the measure of Posidonius, the whole circumference will be 180,000 stadia. This measure was received by Marinus of Tyre, and is usually ascribed to Ptolemy. But this measurement is subject to great uncertainty, both on account of the great refraction of the stars near the horizon, the difficulty of measuring the distance at sea between Rhodes and Alexandria, and by supposing those places under the same meridian, when they are really very different.

Several geographers afterwards made use of the different heights of the pole in distant places under the same meridian, to find the dimensions of the earth. About the year 800, the khalif Almamun had the distance measured between two places that were 2 degrees asunder, and under the same meridian in the plains of Sinjar near the Red Sea. And the result was, that the degree at one time was found equal to 56 miles, and at another  $56\frac{1}{2}$  or  $56\frac{2}{3}$  miles.

The next attempt to find out the circumference of the earth, was in 1525, by Fernelius, a learned philosopher of France. For this purpose, he took the height of the pole at Paris, going from thence directly northwards, till he came to the place where the height of the pole was one degree more than at that city. The length of the way was measured by the number of revolutions made by one of the wheels of his carriage; and after proper allowances for the declivities and turnings of the road, he concluded that 68 Italian miles were equal to a degree on the earth.

According to these methods many other measurements of the earth's circumference have since that time been made, with much greater accuracy: a particular account of which is given under the article DEGREE.

Though the maps of Eratosthenes were the best of his time, they were yet very imperfect and inaccurate. They contained little more than the states of Greece, and the dominions of the successors of Alexander, digested according to the surveys abovementioned. He had indeed seen, and has quoted, the voyages of Pythias into the great Atlantic ocean, which gave him some faint idea of the western parts of Europe; but so imperfect, that they could not be realized into the outlines of a chart. Strabo says he was very ignorant of Gaul, Spain, Germany, and Britain; and he was equally ignorant of Italy, the coasts of the Adriatic, Pontus, and all the countries towards the north.

Such was the state of Geography, and the nature of the maps, before the time of Hipparchus. He made a closer connection between Geography and astronomy, by determining the latitudes and longitudes from celestial observations.

War has usually been the occasion of making or improving the maps of countries; and accordingly Geography made great advances from the progress of the Roman arms. In all the provinces occupied by that people, camps were every where constructed at proper intervals, and good roads made for communication between them; and thus civilization and surveying were carried on according to system, through the whole extent of that large empire. Every new war produced a new survey and itinerary of the countries where the scenes of action passed; so that the materials of Geography were accumulated by every additional conquest. Polybius says, that at the beginning of the second Punic war, when Hannibal was preparing his expedition against Rome, the countries through which he was to pass were carefully measured by the Romans. And Julius Cæsar caused a general survey of the Roman Empire to be made, by a decree of the senate. Three surveyors had this task assigned them, which they completed in 25 years. The Roman itineraries that are still extant, also shew what care and pains they had been at in making surveys in all the different provinces of their empire; and Pliny has filled the 3d, 4th, and 5th books of his Natural History with the Geographical distances that were thus measured. Other maps are also still preserved, known by the name of the Pentingerian Tables, published by Welfer and Bertius, which give a good specimen of what Vegetius calls the *Itinera Picta*, for the better direction of their armies in their march.

The Roman empire had been enlarged to its greatest extent, and all its provinces well known and surveyed, when



when Ptolomy, about 150 years after Christ, composed his system of Geography. The chief materials he employed in composing this work, were the proportions of the gnomon to its shadow, taken by different astronomers at the times of the equinoxes and solstices; calculations founded on the length of the longest days; the measured or computed distances of the principal roads contained in their surveys and itineraries; and the various reports of travellers and navigators. All these were compared together, and digested into one uniform body or system; and afterwards were translated by him into a new mathematical language, expressing the different degrees of latitude and longitude, after the invention of Hipparchus, which had been neglected for 250 years.

Ptolomy's system of Geography, notwithstanding it was still very imperfect, continued in vogue till the last three or four centuries, within which time the great improvements in astronomy, the many discoveries of new countries by voyagers, and the progress of war and arms, have contributed to bring it to a very considerable degree of perfection; the particulars of which will be found treated under their respective articles in this work.

Among the moderns, the chief authors on the subject of Geography are Johannes de Sacrobosco, or John Hallifax, who wrote a treatise on the sphere; Sebastian Munster, in his *Cosmographia Universalis*, in 1559; Clavius, on the sphere of Sacrobosco; Piccioli's *Geographia et Hydrographia Reformata*; Weigelius's *Speculum Terræ*; De Chales's Geography, in his *Mundus Mathematicus*; Cellarius's Geography; Cluverius's *Introductio in Universam Geographiam*; Leibnecht's *Elementa Geographiæ generalis*; Stevenius's *Compendium Geographicum*; Wolfius's *Geographia*, in his *Elementa Matheseos*; Busching's *New System of Geography*; Gordon's, Salmon's, and Guthrie's Grammars; and above all, Varenius's *Geographia generalis*, with Jurin's additions, the most scientific and systematical of any.

GEOMETER, more properly GEOMETRICIAN; which see.

GEOMETRICAL, something that has a relation to Geometry, or done after the manner, or by the means of Geometry. As, a Geometrical construction, a Geometrical curve, a Geometrical demonstration, genius, line, method, Geometrical strictness, &c.

GEOMETRICAL *Construction*, of an equation, is the drawing of lines and figures, so as to express by them the same general property and relation, as are denoted by the algebraical equation. See CONSTRUCTION of Equations.

GEOMETRICAL *Curve* or *Line*, called also an *Algebraical one*, is that in which the relations between the abscissas and ordinates may be expressed by a finite algebraical equation. See ALGEBRAICAL Curves.

GEOMETRICAL *Lines*, as observed by Newton, are distinguished into classes, orders, or genera, according to the number of the dimensions of the equation that expresses the relation between the ordinates and abscissas; or, which comes to the same thing, according to the number of points in which they may be cut by a right line.

Thus, a line of the first order, is a right line, since it

can be only once cut by another right line, and is expressed by the simple equation  $y + ax + b = 0$ : those of the 2d, or quadratic order, will be the circle, and the conic sections, since all of these may be cut in two points by a right line, and expressed by the equation  $y^2 + ax + b \cdot y + cx^2 + dx + e = 0$ : those of the 3d or cubic order, will be such as may be cut in 3 points by a right line, whose most general equation is  $y^3 + ax + b \cdot y^2 + cx^2 + dx + e \cdot y + fx^3 + gx^2 + bx + i = 0$ ; as the cubical and Neilian parabola, the cissoid, &c. And a line of an infinite order, is that which a right line may cut in infinite points; as the spiral, the cycloid, the quadratrix, and every line that is generated by the infinite revolutions of a radius, or circle, or wheel, &c.

In each of those equations,  $x$  is the abscissa,  $y$  its corresponding ordinate, making any given angle with it; and  $a, b, c$ , &c, are given or constant quantities, affected with their signs  $+$  and  $-$ , of which one or more may vanish, be wanting or equal to nothing, provided that by such defect the line or equation does not become one of an inferior order.

It is to be noted that a curve of any kind is denominated by a number next less than the line of the same kind: thus, a curve of the 1st order, (because the right line cannot be reckoned among curves) is the same with a line of the 2d order; and a curve of the 2d kind, the same with a line of the 3d order; &c.

It is to be observed also, that it is not so much the equation, as the construction or description, that makes any curve, geometrical, or not. Thus, the circle is a geometrical line, not because it may be expressed by an equation, but because its description is a postulate: and it is not the simplicity of the equation, but the easiness of the description, that is to determine the choice of the lines for the construction of a problem. The equation that expresses a parabola, is more simple than that which expresses a circle; and yet the circle, by reason of its more simple construction, is admitted before it. Again, the circle and the conic sections, with respect to the dimensions of the equations, are of the same order; and yet the circle is not numbered with them in the construction of problems, but by reason of its simple description is depressed to a lower order, viz, that of a right line; so that it is not improper to express that by a circle, which may be expressed by a right line; but it is a fault to construct that by the conic sections, which may be constructed by a circle.

Either, therefore, the law must be taken from the dimensions of equations, as observed in a circle, and so the distinction be taken away between plane and solid problems: or the law must be allowed not to be strictly observed in lines of superior kinds, but that some, by reason of their more simple description, may be preferred to others of the same order, and be numbered with lines of inferior orders.

In constructions that are equally Geometrical, the most simple are always to be preferred: and this law is so universal as to be without exception. But algebraical expressions add nothing to the simplicity of the construction; the bare descriptions of the lines here are only to be considered; and these alone were considered by those geometers who joined a circle with a right line.



line. And as these are easy or hard, the construction becomes easy or hard: and therefore it is foreign to the nature of the thing, from any other circumstance to establish laws relating to constructions.

Either, therefore, with the ancients, we must exclude all lines beside the circle, and perhaps the conic sections, out of geometry; or admit all, according to the simplicity of the description. If the trochoid were admitted into geometry, by means of it we might divide an angle in any given ratio; would it be right therefore to blame those who would make use of this line to divide an angle in the ratio of one number to another; and contend, that you must make use only of such lines as are defined by equations, and therefore not of this line, which is not so defined? If, when an angle is proposed to be divided, for instance, into 10001 parts, we should be obliged to bring a curve defined by an equation of more than 100 dimensions to do the business; which nobody could describe, much less understand; and should prefer this to the trochoid, which is a line well known; and easily described by the motion of a wheel, or circle: who would not see the absurdity?

Either therefore the trochoid is not to be admitted at all in geometry; or else, in the construction of problems, it is to be preferred to all lines of a more difficult description; and the reason is the same for other curves. Hence the trisections of an angle by a conchoid, which Archimedes in his Lemmas, and Pappus in his Collections, have preferred to the inventions of all others in this case, must be allowed as good; because we must either exclude all lines, beside the circle and right line, out of geometry, or admit them according to the simplicity of their descriptions; in which case the conchoid yields to none, except the circle. Equations are expressions of arithmetical computation, and properly have no place in geometry, excepting so far as quantities truly Geometrical (that is, lines, surfaces, solids, and proportions) may be said to be some equal to others. Multiplications, divisions, and such like computations, are newly received into Geometry, and that unwarily, and contrary to the first design of this science. For whoever considers the construction of problems by a right line and a circle, found out by the first geometers, will easily perceive that geometry was introduced, that by drawing lines, we might easily avoid the tediousness of computation. For which reason the two sciences ought not to be confounded together: the ancients so carefully distinguished between them, that they never introduced arithmetical terms into geometry; and the moderns, by confounding them, have lost the simplicity in which all the elegance of geometry consists. In short, that is arithmetically more simple, which is determined by the more simple equations; but that is Geometrically more simple, which is determined by the more simple drawing of lines; and in geometry that ought to be reckoned best, which is geometrically most simple. Newton's Arith. Univerf. appendix. See CURVES.

**GEOMETRICAL Locus**, or *Place*, called also simply *Locus*, is the path or track of some certain Geometrical determination, in which it always falls. See LOCUS.

**GEOMETRICAL Medium**. See MEDIUM.

**GEOMETRICAL Method of the Ancients**. The ancients established the higher parts of their geometry on

the same principles as the elements of that science, by demonstrations of the same kind: and they were careful not to suppose any thing done, till by a previous problem they had shewn that it could be done by actually performing it. Much less did they suppose any thing to be done that cannot be conceived; such as a line or series to be actually continued to infinity, or a magnitude diminished till it become infinitely less than what it is. The elements into which they resolved magnitudes were finite, and such as might be conceived to be real. Unbounded liberties have of late been introduced; by which geometry, which ought to be perfectly clear, is filled with mysteries. Maclaurin's Fluxions, Introd. p. 39.

**GEOMETRICAL Pace**, is a measure of 5 feet long.

**GEOMETRICAL Plan**, in Architecture. See PLAN.

**GEOMETRICAL Plane**. See PLANE.

**GEOMETRICAL Progression**, a progression in which the terms have all successively the same ratio: as 1, 2, 4, 8, 16, &c, where the common ratio is 2.

The general and common property of a Geometrical progression is, that the product of any two terms, or the square of any one single term, is equal to the product of every other two terms that are taken at an equal distance on both sides from the former. So of these terms,

$$1, 2, 4, 8, 16, 32, 64, \&c,$$

$$1 \times 64 = 2 \times 32 = 4 \times 16 = 8 \times 8 = 64.$$

In any Geometrical Progression, if

$a$  denote the least term,

$z$  the greatest term,

$r$  the common ratio,

$n$  the number of the terms,

$s$  the sum of the series, or all the terms;

then any of these quantities may be found from the others, by means of these general values, or equations, viz,

$$r = \sqrt[n-1]{\frac{z}{a}},$$

$$z = a \times r^{n-1},$$

$$a = z \div r^{n-1},$$

$$n = \frac{\log. \frac{rz}{a}}{\log. r} = \frac{\log. r + \log. z - \log. a}{\log. r},$$

$$s = \frac{r^n - 1}{r - 1} \times a = \frac{r^n - 1}{r - 1} \times \frac{z}{r^{n-1}} = \frac{rz - a}{r - 1} =$$

$$\frac{\frac{n}{z^{\frac{n-1}{n-1}}} - a^{\frac{n-1}{n-1}}}{\frac{1}{z^{\frac{n-1}{n-1}}} - a^{\frac{n-1}{n-1}}}.$$

$$\frac{\frac{1}{z^{\frac{n-1}{n-1}}} - a^{\frac{n-1}{n-1}}}{\frac{1}{z^{\frac{n-1}{n-1}}} - a^{\frac{n-1}{n-1}}}.$$

When the series is infinite, then the least term  $a$  is nothing, and the sum  $s = \frac{rz}{r-1}$ . See also PROGRESSION.

**GEOMETRICAL Proportion**, called also simply *Proportion*, is the similitude or equality of ratios.

Thus, if  $a : b :: c : d$ , or  $a : b = c : d$ , the terms  $a, b, c, d$  are in Geometrical Proportion; also 6, 3, 14, 7, are in Geometrical Proportion, because  $6 : 3 :: 14 : 7$ , or  $6 : 3 = 14 : 7$ .



In a Geometrical Proportion, the product of the extremes, or 1st and 4th terms, is equal to the product of the means, or the 2d and 3d terms: so  $ad = bc$ , and  $6 \times 7 = 3 \times 14 = 42$ . See PROPORTION.

GEOMETRICAL *Solution*, of a problem, is when the problem is directly resolved according to the strict rules and principles of geometry, and by lines that are truly Geometrical. This expression is used in contradistinction to an arithmetical, or a mechanical, or instrumental solution; the problem being resolved only by a ruler and compasses.

The same term is likewise used in opposition to all indirect and inadequate kinds of solutions, as by approximation, infinite series, &c. So, we have no Geometrical way of finding the quadrature of the circle, the duplicature of the cube, or two mean proportionals; though there are mechanical ways, and others, by infinite series, &c.

Pappus informs us, that the ancients endeavoured in vain to trisect an angle, and to find out two mean proportionals, by means of the right line and circle. Afterwards they began to consider the properties of several other lines; as the conchoid, the cissoid, and the conic sections; and by some of these they endeavoured to resolve some of those problems. At length, having more thoroughly examined the matter, and the conic sections being received into geometry, they distinguished Geometrical problems and solutions into three kinds; viz,

1. *Plane* ones, which, deriving their origin from lines on a plane, may be properly resolved by a right line and a circle.—2. *Solid* ones, which are resolved by lines deriving their original from the consideration of a solid; that is, of a cone.—3. *Linear* ones, to the solution of which are required lines more compounded.

According to this distinction, we are not to resolve solid problems by other lines than the conic sections; especially if no other lines beside the right line, circle, and the conic sections, must be received into geometry.

But the moderns, advancing much farther, have received into geometry all lines that can be expressed by equations; and have distinguished, according to the dimensions of the equations, those lines into classes or orders; and have laid it down as a law, not to construct a problem by a line of a higher order, that may be constructed by one of a lower.

GEOMETRICIAN, a person skilled in Geometry.

GEOMETRY, the science or doctrine of local extension, as of lines, surfaces, and solids, with that of ratios, &c.

The name Geometry literally signifies measuring of the earth, as it was the necessity of measuring the land that first gave occasion to contemplate the principles and rules of this art; which has since been extended to numberless other speculations; inasmuch that together with arithmetic, Geometry forms now the chief foundation of all the mathematics.

Herodotus (lib. 2), Diodorus (lib. 1), Strabo (lib. 17), and Proclus, ascribe the invention of Geometry to the Egyptians, and assert that the annual inundations of the Nile gave occasion to it; for those waters bearing away the bounds and land-marks of estates and farms, covering the face of the ground uniformly with

mud, the people, say they, were obliged every year to distinguish and lay out their lands by the consideration of their figure and quantity; and thus by experience and habit they formed a method or art, which was the origin of Geometry. A farther contemplation of the draughts of figures of fields thus laid down, and plotted in proportion, might naturally lead them to the discovery of some of their excellent and wonderful properties; which speculation continually improving, the art continually gained ground, and made advances more and more towards perfection.

Josephus however seems to ascribe the invention to the Hebrews: and others of the ancients make Mercury the inventor. Polyd. Virgil, de Invent. Rer. lib. 1, cap. 18.

From Egypt, this science passed into Greece, being carried thither by Thales; where it was much cultivated and improved by himself, as also by Pythagoras, Anaxagoras of Clazomenae, Hippocrates of Chios, and Plato, who testified his conviction of the necessity and importance of Geometry to the successful study of philosophy, by this inscription over the door of his academy, *Let no one ignorant of Geometry enter here*. Plato thought the word Geometry too mean a name for this science, and substituted instead of it the more extensive name of *Mensuration*; and after him others gave it the title of *Pantometry*. But even these are now become too scanty in their import, fully to comprehend its extent; for it not only inquires into, and demonstrates the quantities of magnitudes, but also their qualities, as the species, figures, ratios, positions, transformations, descriptions, divisions, the finding of their centres, diameters, tangents, asymptotes, curvature, &c. Some again define it as the science of inquiring, inventing, and demonstrating all the affections of magnitude. And Proclus calls it the knowledge of magnitudes and figures, with their limitations; as also of their ratios, affections, positions, and motions of every kind.

About 50 years after Plato, lived Euclid, who collected together all those theorems which had been invented by his predecessors in Egypt and Greece, and digested them into 15 books, called the Elements of Geometry; demonstrating and arranging the whole in a very accurate and perfect manner. The next to Euclid, of those ancient writers whose works are extant, is Apollonius Pergæus, who flourished in the time of Ptolemy Evergetes, about 230 years before Christ, and about 100 years after Euclid. He was author of the first and principal work on Conic Sections; on account of which, and his other accurate and ingenious geometrical writings, he acquired from his patron the emphatical appellation of *The Great Geometrician*. Contemporary with Apollonius, or perhaps a few years before him, flourished Archimedes, celebrated for his mechanical inventions at the siege of Syracuse, and not less so for his very many ingenious Geometrical compositions.

We can only mention Eudoxus of Cnidus, Archytas of Tarentum, Philolaus, Eratosthenes, Aristarchus of Samos, Dinostratus, the inventor of the quadratrix, Menæchmus, his brother and the disciple of Plato, the two Aristotle's, Conon, Thracidius, Nicoteles, Leon, Theudius, Hermodotus, Hero, and Nicomedes the inventor



inventor of the conchoid : besides whom, there are many other ancient geometers, to whom this science has been indebted.

The Greeks continued their attention to it, even after they were subdued by the Romans. Whereas the Romans themselves were so little acquainted with it, even in the most flourishing time of their republic, that Tacitus informs us they gave the name of mathematicians to those who pursued the chimeras of divination and judicial astrology. Nor does it appear they were more disposed to cultivate Geometry during the decline, and after the fall of the Roman empire. But the case was different with the Greeks ; among whom are found many excellent Geometers since the commencement of the Christian era, and after the translation of the Roman empire. Ptolemy lived under Marcus Aurelius ; and we have still extant the works of Pappus of Alexandria, who lived in the time of Theodosius ; the commentary of Eutocius, the Ascalonite, who lived about the year of Christ 540, on Archimedes's mensuration of the circle ; and the commentary on Euclid, by Proclus, who lived under the empire of Anastasius.

The consequent inundation of ignorance and barbarism was unfavourable to Geometry, as well as to the other sciences ; and the few who applied themselves to this science, were calumniated as magicians. However, in those times of European darkness, the Arabians were distinguished as the guardians and promoters of science ; and from the 9th to the 14th century, they produced many astronomers, geometers, geographers, &c ; from whom the mathematical sciences were again received into Spain, Italy, and the rest of Europe, somewhat before the year 1400. Some of the earliest writers after this period, are Leonardus Pisanus, Lucas Pacioli or De Burgo, and others between 1400 and 1500. And after this appeared many editions of Euclid, or commentaries upon him : thus, Orontius Finæus, in 1530, published a commentary on the first 6 books ; as did James Peletarius, in 1557 ; and about the same time Nicholas Tartaglia published a commentary on the whole 15 books. There have been also the editions, or commentaries, of Commandine, Clavius, Billingsly, Scheubelius, Herlinus, Dasypodius, Ramus, Herigon, Stevinus, Saville, Barrow, Taquet, Dechaies, Furnier, Scarborough, Keill, Stone, and many others ; but the completest edition of all the works of Euclid, is that of Dr. Gregory, printed at Oxford 1703, in Greek and Latin : the edition of Euclid, by Dr. Robert Simson of Glasgow, containing the first 6 books, with the 11th and 12th, is much esteemed for its correctness. The principal other elementary writers, besides the editors of Euclid, are Borelli, Pardies, Marchetti, Wolfius, Simpson, &c. And among those who have gone beyond Euclid in the nature of the Elementary parts of Geometry, may be chiefly reckoned, Apollonius, in his Conics, his Loci Plani, De Sectione Determinata, his Tangencies, Inclinations, Section of a Ratio, Section of a Space, &c ; Archimedes, in his treatises of the Sphere and Cylinder, the Dimension of the Circle, of Conoids and Spheroids, of Spirals, and the Quadrature of the Parabola ; Theodosius, in his Spherics ; Serenius, in his Sections of the Cone and Cylinder ; Kepler's Nova Stereometria ; Cavalieri's

Geometria Indivisibilium ; Torricelli's Opera Geometrica ; Viviani, in his Divinationes Geometricæ, Exercitatio Mathematica, De Locis Solidis, De Maximis & Minimis, &c ; Vieta, in his Effectio Geometrica, Supplement. Geometriæ, Sectiones Angulares, Responsum ad Problema, Apollonius Gallus, &c ; Gregory St. Vincent's Quadratura Circuli ; Fermat's Varia Opera Mathematica ; Dr. Barrow's Lectiones Geometricæ ; Bulliald de Lineis Spiralibus ; Cavalieri ; Schooten and Gregory's Exercitationes Geometricæ, and Gregory's Pars Universalis, &c ; De Billy's treatise De Proportionibus Harmonicis ; La Lovera's Geometria veterum promota ; Slusius's Mesolabium, Problemata Solida, &c ; Wallis, in his treatises De Cycloide, Cissoide, &c ; De Proportionibus, De Sectionibus Conicis, Arithmetica Infinitorum, De Centro Gravitatis, De Sectionibus Angularibus, De Angulo Contactus, Cuno-Cuneus, &c, &c ; Hugo De Omerique, in his Analysis Geometrica ; Pascal on the Cycloid ; Step. Angeli's Problemata Geometrica ; Alex. Anderson's Suppl. Apollonii Redivivi, Variorum Problematum Practice, &c ; Baroni's Geomet. Prob. &c ; Guido Grandi Geometr. Demonstr. &c ; Ghetaldi Apollonius Redivivus, &c ; Ludolph van Colen or a Ceulen, de Circulo et Adscriptis, &c ; Snell's Apollonius Batavus, Cyclometricus, &c ; Herberstein's Diotome Circulorum ; Palma's Exercit. in Geometriam ; Guldini Centro-Baryca ; with several others equally eminent, of more modern date, as Dr. Rob. Simson, Dr. Mat. Stewart, Mr. Tho. Simpson, &c. Since the introduction of the new Geometry, or the Geometry of Curve Lines, as expressed by algebraical equations, in this part of Geometry, the following names, among many others, are more especially to be respected, viz, Des Cartes, Schooten, Newton, Maclaurin, Brackenridge, Cramer, Cotes, Waring, &c, &c.

As to the subject of Practical Geometry, the chief writers are Beyer, Kepler, Ramus, Clavius, Mallet, Tacquet, Ozanam, Wolfius, Gregory, with innumerable others.

Geometry is distinguished into Theoretical or Speculative, and Practical.

*Theoretical or Speculative* GEOMETRY, treats of the various properties and relations in magnitudes, demonstrating the theorems, &c. And

*Practical* GEOMETRY, is that which applies those speculations and theorems to particular uses in the solution of problems, and in the measurements in the ordinary concerns of life.

Speculative Geometry again may be divided into Elementary and Sublime.

*Elementary or Common* GEOMETRY, is that which is employed in the consideration of right lines and plane surfaces, with the solids generated from them. And the

*Higher or Sublime* GEOMETRY, is that which is employed in the consideration of curve lines, conic sections, and the bodies formed of them. This part has been chiefly cultivated by the moderns, by help of the improved state of Algebra, and the modern analysis or Fluxions.

GIBBOUS, is used for the shape of one state of the enlightened part of the moon, being that in which she appears



appears more than half full or enlightened, which is the time between the first quarter and the full moon, and from the full moon to the last quarter; appearing then Gibbous, that is, bunched out, or convex on both sides of the enlightened part; as contradistinguished from the state when she is less than half full, when she is said to be horned, or a crescent.

GIMBOLS, are the brass rings by which a sea-compass is suspended in its box that usually stands in the binnacle.

GIN, in Artillery and Mechanics, is a machine for raising great weights, usually composed of three long legs, &c.

GIRDERS, in Architecture, are the largest beams or pieces of timber supporting the floors. Their ends are usually fastened into the summers, or breast-summers; and the joists are framed in at one end to the Girders. By the statute for rebuilding London, no Girder is to lie less than 10 inches into the wall, and their ends to be always laid in loam, &c. The shorter bearings a Girder has, and the oftener it is supported by the internal or partition walls, so much the better. The established breadth and depth of a Girder, according to its length of bearing, are as in the following tablet:

Girders and Summers in length	From	to	must be in	
	10 feet	15 ft.	Breadth 11 inc.	Depth 8 inc.
	15	18	13	9
	18	21	14	10
	21	24	16	12
	24	26	17	14

GIRT, in Timber-measuring, is the circumference of a tree; though some use this word for the quarter or 4th part of the circumference only, on account of the great use that is made of it; for the square of this 4th part is esteemed and used as equal to the area of the section of the tree; which square therefore multiplied by the length of the tree, is accounted the solid content. This content however is always about one-fourth part less than the true quantity; being nearly equal to what this will be after the tree is hewed square in the usual way: so that it seems intended to make an allowance for the squaring of the tree.

GIRT-Line, is a line on the common or carpenter's sliding rule, employed in casting up the contents of trees by means of their Girt.

GIVEN, *Datum*, a term very often used in mathematics, and signifies something that is supposed to be known.

Thus, if a magnitude be known, or if we can find another equal to it, it is said to be Given in magnitude. Or when the position of any thing is known, it is said to be Given in position. And when the diameter of a circle is known, the circle is Given in magnitude. Or the circle is Given in position when its centre is Given in position. When the kind or species of a figure is known, or remains the same, it is Given in specie. And so on.

VOL. I.

Euclid wrote a book of *Data*, or concerning things Given, in 95 propositions, usually accompanying his *Elements*, in the best editions, and which Pappus reckons as one of the best specimens of the analytical works of the ancients.

GLACIS, in Fortification, a sloping bank reaching from the parapet of the counterscarp, or covered-way, to the level side of the field, commonly at the distance of about 40 yards.

GLOBE, a round or spherical body, more usually called a sphere, bounded by one uniform convex surface, every point of which is equally distant from a point within called its centre. Euclid defines the Globe, or sphere, to be a solid figure described by the revolution of a semi-circle about its diameter, which remains unmoved. Also, its axis is the fixed line or diameter about which the semi-circle revolves; and its centre is the same with that of the revolving semi-circle, a diameter of it being any right line that passes through the centre, and terminated both ways by the superficies of the sphere. *Elem.* 11. def. 14, 15, 16, 17.

Euclid, at the end of the 12th book, shews that spheres are to one another in the triplicate ratio of their diameters, that is, their solidities are to one another as the cubes of their diameters. And Archimedes determines the real magnitudes and measures of the surfaces and solidities of spheres and their segments, in his treatise de Sphæra et Cylindro: viz, 1, That the superficies of any Globe is equal to 4 times a great circle of it.—2, That any sphere is equal to  $\frac{2}{3}$  of its circumscribing cylinder, or of the cylinder of the same diameter and altitude.—3, That the curve surface of the segment of a globe, is equal to the circle whose radius is the line drawn from the vertex of the segment to the circumference of the base.—4, That the content of a solid sector of the Globe, is equal to a cone whose altitude is the radius of the Globe, and its base equal to the curve superficies or base of the sector. With many other properties. And from hence are easily deduced these practical rules for the surfaces and solidities of Globes and their segments; viz,

1. *For the Surface of a Globe*, multiply the square of the diameter by 3.1416; or multiply the diameter by the circumference.

2. *For the Solidity of a Globe*, multiply the cube of the diameter by .5236 (viz  $\frac{1}{6}$  of 3.1416); or multiply the surface by  $\frac{1}{6}$  of the diameter.

3. *For the Surface of a Segment*, multiply the diameter of the Globe by the altitude of the segment and the product again by 3.1416.

4. *For the Solidity of a Segment*, multiply the square of the diameter of the Globe by the difference between 3 times that diameter and 2 times the altitude of the segment, and the product again by .5236, or  $\frac{1}{6}$  of 3.1416.

Hence, if  $d$  denote the diameter of the Globe,  $c$  the circumference,  $a$  the altitude of any segment, and  $p = 3.1416$ ; then

$$\begin{array}{l} \text{In the Globe} \\ \text{In the Segt.} \end{array} \left| \begin{array}{l} \text{The surface.} \\ pd^2 = cd \\ pad \end{array} \right| \left| \begin{array}{l} \text{The solidity} \\ \frac{1}{6} pd^3 \\ \frac{1}{6} pa^2 \times 3d - 2a \end{array} \right|$$

See the art. SPHERE, and my *Mensuration*, p. 197 &c, 2d edit.



The **GLOBE**, or *Terraqueous GLOBE*, is the body or mass of the earth and water together, which is nearly globular.

**GLOBE**, or *Artificial GLOBE*, is more particularly used for a Globe of metal, plaister, paper, pasteboard, &c, on the surface of which is drawn a map, or representation of either the heavens or the earth, with the several circles conceived upon them. And hence

**GLOBES** are of two kinds, Terrestrial, and Celestial; which are of considerable use in geography and astronomy, by serving to give a lively representation of their principal objects, and for performing and illustrating many of their operations in a manner easy to be perceived by the senses, and so as to be conceived even without any knowledge of the mathematical grounds of those sciences.

*Description of the Globes.*

The fundamental parts that are common to both Globes, are an axis, representing the axis of the world, passing through the two poles of a spherical shell, representing those of the world, which shell makes the body of the Globe, upon the external surface of which is drawn the representation of the whole surface of the earth, sea, rivers, islands, &c, for the Terrestrial Globe, and the stars and constellations of the heavens, for the Celestial one; besides the equinoctial and ecliptic lines, the zodiac, the two tropics and polar circles, and a number of meridian lines. There is next a brazen meridian, being a strong circle of brass, circumscribing the Globe, at a small distance from it quite round, in which the globe is hung by its two poles, upon which it turns round within this circle, which is divided into 4 times 90 degrees, beginning at the equator on both sides, and ending with 90 at the two poles. There are also two small hour circles, of brass, divided into twice 12 hours, and fitted on the meridian round the poles, which carry an index pointing to the hour. The whole is set in a wooden ring, placed parallel to, and representing the horizon, in which the Globe slides by the brass meridian, elevating or depressing the pole according to any proposed latitude. There is also a thin slip of brass, called a Quadrant of Altitude, made to fit on occasionally upon the brass meridian, at the highest or vertical point, to measure the altitude of any thing above the horizon. A magnetic compass is sometimes set underneath. See the figure of the Globes so mounted, at fig. 1, plate xii.

Such is the plain and simple construction of the artificial Globe, whether celestial or terrestrial, as adapted to the time only for which it is made. But as the angle formed by the equator and ecliptic, as well as their point of intersection, is always changing; to remedy these inconveniences, several contrivances have been made, so as to adapt the same Globes to any other time, either past or to come; as well as other contrivances to answer particular purposes.

Thus, Mr. Senex, a celebrated maker of Globes, had a contrivance which, by means of a nut and screw, caused the pole of the equator to revolve about the pole of the ecliptic, by any quantity answering to the precession of the equinoxes, since the time for which the Globe was made. *Philos. Trans.* number 447, or *Abr.* vol. 8, p. 217, also *Philos. Trans.* vol. 46, p. 290.

Mr. Joseph Harris, late assay-master of the Mint,

made some contrivances to shew the effects of the earth's motions. He fixed two horary circles under the brass meridian, to the axis, one at each pole, so as to turn round with the Globe, and that meridian served as an index to cut the horary divisions. The Globe in this state serves equally for resolving problems in both north and south latitudes, as also in places near the equator; whereas, in the common construction, the axis and horary circle prevent the brass meridian from being moveable quite round in the horizon. This Globe is also adapted for shewing how the vicissitudes of day and night, and the alteration of their lengths, are really occasioned by the motion of the earth: for this purpose, he divides the brass meridian, at one of the poles, into months and days, according to the sun's declination, reckoning from the pole. Therefore, by bringing the day of the month to the horizon, and rectifying the Globe according to the time of the day, the horizon will represent the circle separating light and darkness, and the upper half of the Globe the illuminated hemisphere, the sun being in the zenith. Mr. Harris also gives an account of a cheap machine for shewing how the annual motion of the earth in its orbit causes the change of the sun's declination, without the great expence of an orrery. *Philos. Trans.* number 456, or *Abr.* vol. 8, p. 352.

The late Mr. George Adams made also some useful improvements in the construction of the Globes. Besides what is usual, his Globes have a thin brass semi-circle moveable about the poles, with a small thin sliding circle upon it. On the terrestrial Globe, the former of these is a moveable meridian, and the latter is the visible horizon of any particular place to which it is set. But on the celestial Globe, the semi-circle is a moveable circle of declination, and its small annexed circle an artificial sun or planet. Each Globe has a brass wire circle, placed at the limits of the twilight. The terrestrial Globe has many additional circles, as well as the rhumb-lines, for resolving all the necessary geographical and nautical problems: and on the celestial Globe are drawn, on each side of the ecliptic, 8 parallel circles, at the distance of one degree from each other, including the zodiac; which are crossed at right-angles by segments of great circles at every 5th degree of the ecliptic, for the more readily noting the place of the moon or of any planet upon the Globe. On the strong brass circle of the terrestrial Globe, and about  $23\frac{1}{2}$  degrees on each side of the north pole, the days of each month are laid down according to the sun's declination: and this brass circle is so contrived, that the Globe may be placed with the north and south poles in the plane of the horizon, and with the south pole elevated above it. The equator, on the surface of either Globe, serves the purpose of the horary circle, by means of a semi-circular wire placed in the plane of the equator, carrying two indices, one of which is occasionally to be used to point out the time. For a farther account of these Globes, with the method of using them, see Mr. Adams's treatise on their construction and use.

There are also what are called Patent Globes, made by Mr. Neale; by means of which he resolves several astronomical problems, which do not admit of solution by the common Globes.

Mr.



Mr. Fergufon likewise made feveral improvements of the Globes, particularly one for conſtructing dials, and another called a planetary Globe. See Philof. Tranf. vol. 44, p. 535, and Fergufon's Aſtron. p. 291, and 292.

Laſtly, in the Philof. Tranf. for 1789, vol. 79, p. 1, Mr. Smeaton has propoſed ſome improvements of the celeftial Globe, eſpecially with reſpect to the quadrant of altitude, for the reſolution of problems relating to the azimuth and altitude. The difficulty, he obſerves, that has occurred in fixing a ſemicircle, ſo as to have a centre in the zenith and nadir points of the Globe, at the ſame time that the meridian is left at liberty to raiſe the pole to its deſired elevation, I ſuppoſe, has induced the Globe-makers to be contented with the ſtrip of thin flexible braſs, called the quadrant of altitude; and it is well known how imperfectly it performs its office. The improvement I have attempted, is in the application of a quadrant of altitude of a more ſolid conſtruction; which being affixed to a braſs ſocket of ſome length, and this ground, and made to turn upon an upright ſteel ſpindle, fixed in the zenith, ſteadily directs the quadrant, or rather arc, of altitude to its true azimuth, without being at liberty to deviate from a vertical circle to the right hand or left: by which means the azimuth and altitude are given with the ſame exactneſs as the meaſure of any other of the great circles. For a more particular deſcription of this improvement, illuſtrated with figures, ſee the place above quoted.

#### *The uſe of the Terreſtrial GLOBE.*

PROB. I. *To find the latitude and longitude of any place.*—Bring the place to the graduated ſide of the firſt meridian: then the degree of the meridian it cuts is the latitude ſought; and the degree of the equator then under the meridian is the longitude.

II. *To find a place, having a given latitude and longitude.*—Find the degree of longitude on the equator, and bring it to the braſs meridian; then find the degree of latitude on the meridian, either north or ſouth of the equator, as the given latitude is north or ſouth; then the point of the Globe juſt under that degree of latitude is the place required.

III. *To find all the places on the Globe that have the ſame latitude, and the ſame longitude, or hour, with a given place, as ſuppoſe London.*—Bring the given place London to the meridian, and obſerve what places are juſt under the edge of it, from north to ſouth; and all thoſe places have the ſame longitude and hour with it. Then turn the Globe quite round; and all thoſe places which paſs juſt under the given degree of latitude on the meridian, have the ſame latitude with the given place.

IV. *To find the Antæci, Pericæci and Antipodes, of any given place, ſuppoſe London.*—Bring the given place London to the meridian, then count  $51\frac{1}{2}$  the ſame degree of latitude ſouthward, or towards the other pole, and the point thus arrived at will be the Antæci, or where the hour of the day or night is always the ſame at both places at the ſame time, and where the ſeaſons and lengths of days and nights are alſo equal, but at half a year diſtance from each other, becauſe their ſea-

ſons are oppoſite or contrary. London being ſtill under the meridian, ſet the hour index to 12 at noon, or pointing towards London; then turn the Globe juſt half round, or till the index point to the oppoſite hour, or 12 at night; and the place that comes under the ſame degree of the meridian where London was, ſhews where the Pericæci dwell, or thoſe people that have the ſame ſeaſons and at the ſame time as London, as alſo the ſame length of days and nights &c at that time, but only their time or hour is juſt oppoſite, or 12 hours diſtant, being day with one when night with the other, &c. Laſtly, as the Globe ſtands, count down by the meridian the ſame degree of latitude ſouth, and that will give the place of the Antipodes of London, being diametrically under or oppoſite to it; and ſo having all its times, both hours and ſeaſons oppoſite, being day with the one when night with the other, and ſummer with the one when winter with the other.

V. *To find the Diſtance of two places on the Globe.*—If the two places be either both on the equator, or both on the ſame meridian, the number of degrees in the diſtance between them, reduced into miles, at the rate of 70 Engliſh miles to the degree, (or more exact  $69\frac{1}{2}$ ), will give the diſtance nearly. But in any other ſituations of the two places, lay the quadrant of altitude over them, and the degrees counted upon it, from the one place to the other, and turned into miles as above, will give the diſtance in this caſe.

VI. *To find the Difference in the Time of the day at any two given places, and thence the Difference of Longitude.*—Bring one of the places to the meridian, and ſet the hour index to 12 at noon; then turn the Globe till the other place comes to the meridian, and the index will point out the difference of time; then by allowing  $15^{\circ}$  to every hour, or  $1^{\circ}$  to 4 minutes of time, the difference of longitude will be known.—Or the difference of longitude may be found without the time, thus:

Fiſt bring the one place to the meridian, and note the degree of longitude on the equator cut by it; then do the ſame by the other place; which gives the longitudes of the two places; then ſubtracting the one number of degrees from the other, gives the difference of longitude ſought.

VII. *The time being known at any given place, as ſuppoſe London, to find what hour it is in any other part of the world.*—Bring the given place, London, to the meridian, and ſet the index to the given hour; then turn the Globe till the other place come to the meridian, and look at what hour the index points, which will be the time ſought.

VIII. *To find the Sun's place in the ecliptic, and alſo on the Globe, at any given time.*—Look into the calendar on the wooden horizon for the month and day of the month propoſed, and immediately oppoſite ſtands the ſign and degree which the ſun is in on that day. Then in the ecliptic drawn upon the Globe, look for the ſame ſign and degree, and that will be the place of the ſun required.

IX. *To find at what place on the earth the ſun is vertical, at a given moment of time at another place, as ſuppoſe London.*—Find the ſun's place on the Globe by the laſt problem, and turn the Globe about till



that place come to the meridian, and note the degree of the meridian just over it. Then turn the Globe till the given place, London, come to the meridian, and set the index to the given moment of time. Lastly, turn the Globe till the index points to 12 at noon; then the place of the earth, or Globe, which stands under the before noted degree, has the sun at that moment in the zenith.

X. *To find how long the sun shines without setting, in any given place in the frigid zones.*—Subtract the degrees of latitude of the given place from 90, which gives the complement of the latitude, and count the number of this complement upon the meridian from the equator towards the pole, marking that point of the meridian; then turn the Globe round, and carefully observe what two degrees of the ecliptic pass exactly under the point marked on the meridian. Then look for the same degrees of the ecliptic on the wooden horizon, and just opposite to them stand the months and days of the months corresponding, and between which two days the sun never sets in that latitude.

If the beginning and end of the longest night be required, or the period of time in which the sun never rises at that place; count the same complement of latitude towards the south or farthest pole, and then the rest of the work will be the same in all respects as above.

Note, that this solution is independent of the horizontal refraction of the sun, which raises him rather more than half a degree higher, by that means making the day so much longer, and the night the shorter; therefore in this case, set the mark on the meridian half a degree higher up towards the north pole, than what the complement of latitude gives; then proceed with it as before, and the more exact time and length of the longest day and night will be found.

XI. *A place being given in the torrid zone, to find on what two days of the year the sun is vertical at that place.*—Turn the Globe about till the given place come to the meridian, and note the degree of the meridian it comes under. Next turn the Globe round again, and note the two points of the ecliptic passing under that degree of the meridian. Lastly, by the wooden horizon, find on what days the sun is in those two points of the ecliptic; and on these days he will be vertical to the given place.

XII. *To find those places in the torrid zone to which the sun is vertical on a given day.*—Having found the sun's place in the ecliptic, as in the 8th problem, turn the Globe to bring the same point of the ecliptic on the Globe to the meridian; then again turn the Globe round, and note all the places which pass under that point of the meridian; which will be the places sought.

After the same manner may be found what people are Ascii for any given day. And also to what place of the earth, the moon, or any other planet, is vertical on a given day; finding the place of the planet on the globe by means of its right ascension and declination, like finding a place from its longitude and latitude given.

XIII. *To rectify the Globe for the latitude of any place.*—By sliding the brass meridian in its groove, elevate

the pole as far above the horizon as is equal to the latitude of the place; so for London, raise the north pole  $51\frac{1}{2}$  degrees above the wooden horizon: then turn the Globe on its axis till the place, as London, come to the meridian, and there set the index to 12 at noon. Then is the place exactly on the vertex, or top point of the Globe, at  $90^\circ$  every way round from the wooden horizon, which represents the horizon of the place. And if the frame of the Globe be turned about till the compass needle point to  $22\frac{1}{2}$  degrees, or two points west of the north point (because the variation of the magnetic needle is nearly  $22\frac{1}{2}$  degrees west), so shall the Globe then stand in the exact position of the earth, with its axis pointing to the north pole.

XIV. *To find the length of the day or night, or the sun's rising or setting, in any latitude; having the day of the month given.*—Rectify the Globe for the latitude of the place; then bring the sun's place on the globe to the meridian, and set the index to 12 at noon, or the upper 12, and then the Globe is in the proper position for noon day. Next turn the Globe about towards the east till the sun's place come just to the wooden horizon, and the index will then point to the hour of sunrise; also turn the Globe as far to the west side, or till the sun's place come just to the horizon on the west side, and then the index will point to the hour of sunset. These being now known, double the hour of setting will be the length of the day, and double the rising will be the length of the night.—And thus also may the length of the longest day, or the shortest day, be found for any latitude.

XV. *To find the beginning and end of Twilight on any day of the year, for any latitude.*—It is twilight all the time from sunset till the sun is  $18^\circ$  below the horizon, and the same in the morning from the time the sun is  $18^\circ$  below the horizon till the moment of his rise. Therefore, rectify the Globe for the latitude of the place, and for noon by setting the index to 12, and screw on the quadrant of altitude. Then take the point of the ecliptic opposite the sun's place, and turn the Globe on its axis westward, as also the quadrant of altitude, till that point cut this quadrant in the  $18^{\text{th}}$  degree below the horizon, then the index will shew the time of dawning in the morning; next turn the Globe and quadrant of altitude towards the east, till the said point opposite the sun's place meet this quadrant in the same  $18^{\text{th}}$  degree, and then the index will shew the time when twilight ends in the evening.

XVI. *At any given day, and hour of the day, to find all those places on the Globe where the sun then rises, or sets, as also where it is noon day, where it is day light, and where it is in darkness.*—Find what place the sun is vertical to, at that time; and elevate the Globe according to the latitude of that place, and bring the place also to the meridian; in which state it will also be in the zenith of the Globe. Then is all the upper hemisphere, above the wooden horizon, enlightened, or in day light; while all the lower one, below the horizon, is in darkness, or night: those places by the edge of the meridian, in the upper hemisphere, have noon day, or 12 o'clock; and those by the meridian below, have it midnight: lastly, all those places by the eastern side of the horizon, have the sun just setting,



ting, and those by the western horizon have him just rising.

Hence, as in the middle of a lunar eclipse the moon is in that degree of the ecliptic opposite to the sun's place; by the present problem it may be shewn what places of the earth then see the middle of the eclipse, and what the beginning or ending; by using the moon's place instead of the sun's place in the problem.

XVII. *To find the bearing of one place from another, and their angle of position.*—Bring the one place to the zenith, by rectifying the Globe for its latitude, and turning the Globe till that place come to the meridian; then screw the quadrant of altitude upon the meridian at the zenith, and make it revolve till it come to the other place on the Globe; then look on the wooden horizon for the point of the compass, or number of degrees from the south, where the quadrant of altitude cuts it, and that will be the bearing of the latter place from the former, or the angle of position sought.

*The Use of the Celestial GLOBE.*

The Celestial Globe differs from the terrestrial only in this; instead of the several parts of the earth, the images of the stars and constellations are designed. The meridian circle drawn through the two poles and through the point Cancer, represents the solstitial colure; but that through the point Aries, represents the equinoctial colure.

PROB. XVIII. *To exhibit the true representation of the face of the heavens at any given time and place.*—Rectify for the lat. of the place, by prob. 13, setting the Globe with its pole pointing to the pole of the world, by means of a compass. Find the sun's place in the ecliptic, and turn the Globe to bring it to the meridian, and there set the index to 12 at noon. Again revolve the Globe on its axis, till the index point to the given hour of the day or night: so shall the Globe in this position exactly represent the face of the heavens as it appears at that time, every constellation and star, in the heavens, answering in position to those on the Globe; so that, by examining the Globe, it will immediately appear which stars are above or below the horizon, which on the east or western parts of the heavens, which lately risen, and which going to set, &c. And thus the positions of the several planets, or comets, may also be exhibited; having marked the places of the Globe where they are, by means of their declination and right ascension.

XIX. *To find the Declination and Right-ascension of any star upon the Globe.*—Turn the Globe till the star come to the meridian: then the number of degrees on the meridian, between the equator and the star, is its declination; and the degree of the equator cut by the meridian, is the right-ascension of the star.—In like manner are found the declination and right-ascension of the sun, or any other point.

XX. *To find the Latitude and Longitude of any star drawn upon the Globe.*—Bring the solstitial colure to the meridian, and there fix the quadrant of altitude over the pole of the ecliptic in the same hemisphere with the star, and bring its graduated edge to the star: then the degree on the quadrant cut by the star is its latitude, counted from the ecliptic; and the degree of the ecliptic cut by the quadrant its longitude.

XXI. *To find the place of a star, planet, comet, &c. on the Globe; its declination and right-ascension being known.*—Find the given point of right-ascension on the equinoctial, and bring it to the meridian; then count the degrees of declination upon the meridian from the equinoctial, and there make a mark on the Globe, which will be the place of the planet, &c, sought.

XXII. *To find the place of a star, planet, comet, or other object on the Globe; its latitude and longitude being given.*—Bring the pole of the ecliptic to the meridian, and there fix the quadrant of altitude, which turn round till its edge cut the given longitude on the ecliptic; then count the given latitude, from the ecliptic, upon the quadrant of altitude, and there make a mark on the Globe, which will be the place of the planet, &c, sought.—The place on the Globe, of any such planet, &c, being found by this or the foregoing problem, its rising, or setting, or any other circumstance concerning it, may then be found, the same as the sun, by the proper problems.

XXIII. *To find the rising, setting, and culminating of a star, planet, sun, &c; with its continuance above the horizon, for any place and day; as also its oblique ascension and descension, with its eastern and western amplitude and azimuth.*—Adjust the Globe to the state of the heavens at 12 o'clock that day. Bring the star, &c, to the eastern side of the horizon: which will give its eastern amplitude and azimuth, and the time of rising, as for the sun. Again, turn the Globe to bring the same star to the western side of the horizon: so will the western amplitude and azimuth, with the time of setting, be found. Then, the time of rising, subtracted from that of setting, leaves the continuance of the star above the horizon: this continuance above the horizon taken from 24 hours, leaves the time it is below the horizon. Lastly, bring the star to the meridian, and the hour to which the index then points is the time of its culmination, or southing.

XXIV. *To find the altitude of the sun, or star, &c, for any given hour of the day or night.*—Adjust the Globe to the position of the heavens, and turn it till the index point at the given hour. Then fix on the quadrant of altitude, at 90 degrees from the horizon, and turn it to the place of the sun or star: so shall the degrees of the quadrant, intercepted between the horizon and the sun or star, be the altitude sought.

XXV. *Given the altitude of the sun by day, or of a star by night, to find the hour of the day or night.*—Rectify the Globe as in the foregoing problem; and turn the Globe and quadrant, till such time as the star or degree of the ecliptic the sun is in, cut the quadrant in the given degree of altitude; then will the index point at the hour required.

XXVI. *Given the azimuth of the sun or a star, to find the time of the day or night.*—Rectify the Globe, and bring the quadrant to the given azimuth in the horizon; then turn the Globe till the sun or star come to the quadrant, and the index will then shew the time of the day or night.

GLOBULAR, relating to, or partaking the property or shape of, the Globe. As Globular chart, Globular projection, or Globular sailing, &c.

GLOBULAR Chart, is a representation of the surface,



or part of the surface, of the terraqueous Globe upon a plane ; in which the parallels of latitude are circles nearly concentric ; and the meridians are curves bending towards the poles ; the rhumb-lines being curves also.

The merits of this chart consist in these particulars, viz, that the distances between places on the same rhumb are all measured by the same scale of equal parts ; and the distance of any two places in the arch of a great circle, is nearly represented in this chart by a straight line.

Land maps also made according to this projection would have great advantages over those made in any other way. But for sea charts for the use of navigation, Mercator's are preferable, as both the meridians and parallels, as also the rhumbs, are all straight lines.

This projection is not new, though not much noticed till of late. It is mentioned by Ptolomy, in his Geography ; and also by Blundeville, in his Exercises.

For Globular projection of maps or charts, see MAP.

GLOBULAR Sailing, is the method of resolving the cases of sailing upon principles deduced from the spherical figure of the earth. Such as Mercator's sailing, or Great-circle sailing ; which see.

GLOSSOCOMON, in Mechanics, is a name given by Heron to a machine composed of divers dented wheels with pinions, serving to raise huge weights.

GNOMON, in Astronomy, is an instrument or apparatus for measuring the altitudes, declinations, &c, of the sun and stars. The Gnomon is usually a pillar, or column, or pyramid, erected upon level ground, or a pavement. For making the more considerable observations, both the ancients and moderns have made great use of it, especially the former ; and many have preferred it to the smaller quadrants, both as more accurate, easier made, and more easily applied.


The most ancient observation of this kind extant, is that made by Pytheas, in the time of Alexander the Great, at Marfeilles, where he found the height of the Gnomon was in proportion to the meridian shadow at the summer solstice, as  $213\frac{1}{8}$  to 600 ; just the same as Gassendi found it to be, by an observation made at the same place, almost 2000 years after, viz, in the year 1636. Ricciol. Almag. vol. 1, lib. 3, cap. 14.

Ulugh Beigh, king of Parthia, &c, used a Gnomon in the year 1437, which was 180 Roman feet high. That erected by Ignatius Dante, in the church of St. Petronius, at Bologna, in the year 1576, was 67 feet high. M. Cassini erected another of 20 feet high, in the same church, in the year 1655.

The Egyptian obelisks were also used as Gnomons ; and it is thought by some modern travellers that this was the very use they were designed and built for ; it has also been found that their four sides stand exactly facing the four cardinal points of the compass. It may be added, that the Spaniards in their conquest of Peru, found pillars of curious and costly workmanship, set up in several places, by the meridian shadows of which their amatas or philosophers had, by long experience and repeated observations, learned to determine the times of the equinoxes ; which seasons of the year were celebrated with great festivity and rich offerings, in honour of the sun. Garcillasso de la Vega, Hist. Peru. lib. 2, cap. 22.

USE of the GNOMON, in taking the meridian altitude of the Sun, and thence finding the Latitude of the place.—

A meridian line being drawn through the centre of the Gnomon, note the point where the shadow of the Gnomon terminates when projected along the meridian line, and measure the distance of that extreme point from the centre of the Gnomon, which will be the length of its shadow. Then having the height of the Gnomon, and the length of the shadow, the sun's altitude is thence easily found.



Suppose, ex. gr. AB the Gnomon, and AC the length of the shadow. Here in the right-angled triangle ABC, are given the base AC, and the perpendicular AB, to find the angle C, or the sun's altitude, which will be found by this analogy, as  $CA : AB :: \text{radius} : \text{the tang. of } \angle C$ , that is, as the length of the shadow is to the height of the Gnomon, so is radius to the tangent of the sun's altitude above the horizon.


The following example will serve to illustrate this proposition : Pliny says, Nat. Hist. lib. 2, cap. 72, that at Rome, at the time of the equinoxes, the shadow is to the Gnomon as 8 to 9 ; therefore as

$8 : 9 :: 1 \text{ or radius} : \frac{9}{8} = 1.125$  a tangent, to which answers the angle  $48^\circ 22'$ , which is the height of the equator at Rome, and its complement  $41^\circ 38'$  is therefore the height of the pole, or the latitude of the place.

Riccioli remarks the following defects in the observations of the sun's height, made with the Gnomon by the ancients, and some of the moderns : viz, that they neglected the sun's parallax, which makes his apparent altitude less, by the quantity of the parallax, than it would be, if the Gnomon were placed at the centre of the earth : 2d, they neglected also the refraction, by which the apparent height of the sun is a little increased : and 3dly, they made the calculations from the length of the shadow, as if it were terminated by a ray coming from the centre of the sun's disc, whereas the shadow is really terminated by a ray coming from the upper edge of the sun's disc ; so that, instead of the height of the sun's centre, their calculations gave the height of the upper edge of his disc. And therefore, to the altitude of the sun found by the Gnomon, the sun's parallax must be added, and from the sum must be subtracted the sun's semidiameter, and refraction, which is different at different altitudes ; which being done, the correct height of the equator at Rome will be  $48^\circ 4' 13''$ , the complement of which, or  $41^\circ 55' 46''$ , is the latitude. Ricciol. Geogr. Refor. lib. 7, cap. 4.




The preceding problem may be resolved more accurately by means of a ray of light let in through a small hole, than by a shadow, thus : Make a circular perforation in a brass plate, to transmit enough of the sun's rays to exhibit his image on the floor, or a stage ; fix the plate parallel to the horizon in a high place, proper for observation, the height of which above the floor let be accurately measured with a plummet. Let the floor, or stage, be perfectly plane and horizontal, and coloured over with some white substance, to shew the sun more distinctly. Upon this horizontal plane draw a meridian line passing through the foot or centre of the Gnomon, i. e. the point upon which the plummet falls from the centre of the hole ; and upon this line note the extreme points I and K of the sun's image or diameter, and from each end subtract the image of half the diameter of the aperture, viz KH and LI : then will HL be the image of the sun's diameter, which, when bisected in B, gives the point on which the rays fall from the centre of the sun.



Now having given the line AB, and the altitude of the Gnomon AG, beside the right angle A, the angle B, or the apparent altitude of the sun's centre, is easily found, thus : as AB : AG :: radius : tang. angle B.

GNOMON, in Dialling, is the style, pin, or cock of a dial, the shadow of which points out the hours. This is always supposed to represent the axis of the world, to which it is therefore parallel, or coincident, the two ends of it pointing straight to the north and south poles of the world.

GNOMON, in Geometry, is a figure formed of the two complements, in a parallelogram, together with either of the parallelograms about the diameter. Thus the parallelogram AC being divided into four parallelograms by the two lines DG, EF parallel to the sides, forming the two complements AB and BC, with the two DE, FG about the diameter HI : then the two Gnomons are AB + BC + DE, and AB + BC + FG.



GNOMONIC Projection of the Sphere, is the representation of the circles of an hemisphere upon a plane touching it in the vertex, by the eye in the centre, or by lines or rays issuing from the centre of the hemisphere, to all the points in the surface.

In this projection of the sphere, all the great circles are projected into right lines, on the plane, of an indefinite length ; and all lesser circles that are parallel to

the plane, into circles ; but if oblique to the plane, then are they projected either into ellipses or hyperbolas, according to their different obliquity. It has its name from Gnomonics, or Dialling, because the lines on the face of every dial are from a projection of this sort : for if the sphere be projected on any plane, and upon that side of it on which the sun is to shine ; also the projected pole be made the centre of the dial, and the axis of the globe the style or Gnomon, and the radius of projection its height ; you will have a dial drawn with all its furniture. See Emerson's Projection of the Sphere.

GNOMONICS, the same as DIALLING ; or the art of drawing sun and moon dials, on any given plane ; being so called, because it shews how to find the hour of the day or night by the shadow of a Gnomon or style.

GOLDEN Number, is the particular year of the Metonic or Lunar Cycle. See LUNAR CYCLE.

To find the Golden Number : Add 1 to the given year, and divide the sum by 19, and what remains will be the Golden Number ; unless 0 remain, for then 19 is the Golden Number.

Thus, the Golden Number for the year 1791, is 6 ; as by the operation in the margin.

1791
1
19)1792(94
171
82
76
Golden No. 6

GOLDEN Rule, a rule so called on account of its excellent use, in arithmetic, and especially in ordinary calculations, by which numbers are found in certain proportions, viz, having three numbers given, to find a 4th number, that shall have the same proportion to the 3d as the 2d hath to the 1st. On this account, it is otherwise called *The Rule of Three*, and *The Rule of Proportion*. See RULE of Three.

Having stated, or set down in a line, the three terms, in the order in which they are proportional, multiply the 2d and 3d together, and divide the product by the 1st, so shall the quotient be the answer, or the 4th term sought.

Thus, if 3 yards of cloth cost a guinea or 21 shillings, what will 20 yards cost. Here the two prices or values must bear the same proportion to each other as the two quantities, or number of yards of cloth, i. e. 3 must bear the same proportion to 20, as 21s, the value of the former, must bear to the value of the latter : and therefore the stating and operation of the numbers will be thus,

Then multiplying the 2d and 3d together, and dividing the product by the 1st, it gives 140s. or 7l. for the answer, being the cost of 20 yards.

3 : 20 :: 21 : 140s.
20
3)420
Quot. 140s.
or 7l.

GONIOMETRICAL Lines, are lines used for measuring or determining the quantity of angles : such as sines, tangents, secants, versed sines, &c.


Mr. Jones, in the Philos. Transf. number 483, sect. 26, gave a paper, containing a commodious disposition of equations for exhibiting the relations of Goniometrical Lines ; from which a multitude of curious theo-

rems



rems may be derived. See also Robertson's Elem. of Navigation, vol. 1, p 181, Edit. 4.

**GONIOMETRY**, a method of measuring angles, so called by M. de Lagny, who gave several papers, on this method, in the Memoirs of the Royal Acad. an. 1724, 1725, 1729. M. de Lagny's method of Goniometry consists in measuring the angles with a pair of compasses, and that without any scale whatever, except an undivided semicircle. Thus, having any angle drawn upon paper, to be measured; produce one of the sides of the angle backwards behind the angular point; then with a pair of fine compasses describe a pretty large semicircle from the angular point as a centre, cutting the sides of the proposed angle, which will intercept a part of the semicircle. Take then this intercepted part very exactly between the points of the compasses, and turn them successively over upon the arc of the semicircle, to find how often it is contained in it, after which there is commonly some remainder: then take this remainder in the compasses, and in like manner find how often it is contained in the last of the integral parts of the 1st arc, with again some remainder: find in like manner how often this last remainder is contained in the former; and so on continually, till the remainder become too small to be taken and applied as a measure. By this means he obtains a series of quotients, or fractional parts, one of another, which being properly reduced into one fraction, give the ratio of the first arc to the semicircle, or of the proposed angle to two right angles, or 180 degrees, and consequently that angle itself in degrees and minutes.



Thus, suppose the angle BAC be proposed to be measured. Produce BA out towards *f*; and from the centre A describe the semicircle *abcf*, in which *ab* is the measure of the proposed angle. Take *ab* in the compasses, and apply it 4 times on the semicircle, as at *b*, *c*, *d*, and *e*; then take the remainder *fe*, and apply it back upon *ed*, which is but once, viz at *g*; again take the remainder *gd*, and apply it 5 times on *ge*, as at *h*, *i*, *k*, *l*, and *m*; lastly, take the remainder *me*, and it is contained just 2 times in *ml*. Hence the series of quotients is 4, 1, 5, 2; consequently the 4th or last arc *em* is  $\frac{1}{2}$  the third *ml* or *gd*, and therefore the 3d arc *gd* is  $\frac{1}{5\frac{1}{2}}$  or  $\frac{2}{11}$  of the 2d arc *ef*; and therefore again this 2d arc *ef* is  $\frac{1}{1\frac{2}{11}}$  or  $\frac{11}{13}$  of the 1st arc *ab*; and consequently this 1st arc *ab* is  $\frac{1}{4\frac{11}{13}}$  or  $\frac{13}{63}$  of the whole semicircle *af*. But  $\frac{13}{63}$  of 180° are  $37\frac{1}{2}$  degrees, or  $37^{\circ} 8' 34''\frac{1}{2}$ , which therefore is the measure of the angle sought. When the operation is nicely performed, this angle may be within 2 or 3 minutes of the truth; though M. de Lagny pretends to measure much nearer than that.

It may be added, that the series of fractions forms what is called a continued fraction. Thus, in the example above, the continued fraction, and its reduction, will be as follow:

$$\frac{1}{4} + \frac{1}{1 + \frac{1}{5\frac{1}{2}}} = \frac{1}{4} + \frac{1}{1\frac{2}{11}} = \frac{1}{4\frac{11}{13}} = \frac{13}{63};$$

the quotients being the successive denominators, and 1 always for each numerator.

**GORGE**, or *Neck*, in Architecture, is the narrowest part of the Tuscan or Doric capitals, lying above the shaft of the pillar, between the astragal and the annulets.

It is also a kind of concave moulding, serving for compartments &c, larger than a scotia, but not so deep.

**GORGE**, in Fortification, is the entrance into a bastion, or a ravelin, or other out-work.

The **GORGE** of a Bastion, is what remains of the sides of the polygon of a place, after cutting off the curtains; in which case it makes an angle in the centre of the bastion, viz, the angle made by two adjacent curtains produced to meet within the bastion.

In flat bastions, the Gorge is a right line on the curtain, reaching between the two flanks.

**GORGE** of a Half-moon, or of a Ravelin, is the space between the two ends of their faces next the place.

**GORGE** of the other out-works, is the interval between their sides next the great ditch.

All the Gorges are to be made without parapets: otherwise the besiegers, having taken possession of a work, might make use of them to defend themselves from the shot of the place. So that they are only fortified with pallisadoes, to prevent a surprize.

The **Demi-GORGE**, or *Half* the **GORGE**, is that part of the polygon between the flank and the centre of the bastion.

**GOTHIC Architecture**, is that which deviates from the manner, character, proportions, &c, of the antique; having its ornaments wild and chimerical, and its profiles incorrect. This manner of building came originally from the North, whence it was brought, in the 5th century, by the Goths into Germany, and has since been introduced into other countries. The first or most ancient style of Gothic building was very solid, heavy, massive and simple, with semicircular arches, &c: but the more modern style of the Gothic is exceedingly rich, light, and delicate; having an abundance of little whimsical ornaments, with sharp-pointed arches formed by the intersections of different circular segments; also lofty and light spires and steeples, large ramified windows, clustered pillars, &c. Of this kind are our English cathedrals, and many other old buildings.

**GRADUATION**, is used for the act of Graduating, or dividing any thing into degrees.

For an account of the various methods of Graduating mathematical and astronomical instruments, by straight and circular diagonals, and by concentric arcs, &c; see *Plain Scale*, *Nonius*, and *Vernier*. And for an account of Mr. Bird's improved method of dividing astronomical instruments, see *MURAL Arch*.

Mr. Ramsden, an ingenious mathematical instrument-maker of London, has lately published, by encouragement of the commissioners of longitude, an explanation and



and description of an engine contrived by him for dividing mathematical instruments, accompanied with proper drawings; in consideration of which, the said commissioners have granted to him the sum of 615*l*. See his book, 4to, 1777.

On the subject of dividing a foot into many thousand parts, for mathematical purposes, see *Philos. Transf.* vol. 2, p. 457, 459, 541, or *Abr.* vol. 1, p. 218, 220, &c. And for an account of various other methods and Graduations, see a paper of Mr. Smeaton's in the *Philos. Transf.* vol. 76, for the year 1786, p. 1; being "Observations on the Graduation of astronomical instruments; with an explanation of the method invented by the late Mr. Henry Hindley, of York, clock-maker, to divide circles into any given number of parts."

GRAHAM (GEORGE), clock and watch maker, the most ingenious and accurate artist in his time, was born at Gratwick, a village in the north of Cumberland, in 1675. In 1688 he came up to London, and was put apprentice to a person in that profession; but after being some time with his master, he was received, purely on account of his merit, into the family of the celebrated Mr. Tompion, who treated him with a kind of parental affection as long as he lived. That Mr. Graham was, without competition, the most eminent of his profession, is but a small part of his character: he was the best general mechanic of his time, and had a complete knowledge of practical astronomy; so that he not only gave to various movements for measuring time a degree of perfection which had never before been attained, but invented several astronomical instruments, by which considerable advances have been made in that science: he made great improvements in those which had before been in use; and, by a wonderful manual dexterity, constructed them with greater precision and accuracy than any other person in the world.

A great mural arch in the observatory at Greenwich was made for Dr. Halley, under Mr. Graham's immediate inspection, and divided by his own hand: and from this incomparable original, the best foreign instruments of the kind are copies made by English artists. The sector by which Dr. Bradley first discovered two new motions in the fixed stars, was of his invention and fabric. He comprised the whole planetary system within the compass of a small cabinet; from which, as a model, all the modern orreries have been constructed. And when the French Academicians were sent to the north, to make observations for ascertaining the figure of the earth, Mr. Graham was thought the fittest person in Europe to supply them with instruments; by which means they finished their operations in one year; while those who went to the south, not being so well furnished, were very much embarrassed and retarded in their operations.

Mr. Graham was many years a member of the Royal Society, to which he communicated several ingenious and important discoveries, viz, from the 31st to the 42d volume of the *Philos. Transactions*, chiefly on astronomical and philosophical subjects; particularly a kind of horary alteration of the magnetic needle; a quicksilver pendulum, and many curious particulars relating to the true length of the simple pendulum, upon

VOL. I.

which he continued to make experiments till almost the year of his death, which happened in 1751, at 76 years of age.

His temper was not less communicative than his genius was penetrating; and his principal view was the advancement of science, and the benefit of mankind. As he was perfectly sincere, he was above suspicion; and as he was above envy, he was candid.

GRANADO, in Artillery, is a little shell or hollow globe of iron, or other matter, which, being filled with powder, is fired by means of a small fusee, and thrown either by the hand, or a piece of ordnance. As soon as it is kindled, the case flies in pieces, to the great danger of all that stand near it. Granadoes serve to set fire to close and narrow passages, and are often thrown with the hand among the soldiers, to disorder their ranks; more especially in those posts where they stand thickest, as in trenches, redoubts, lodgments, &c.

GRAVESANDE (WILLIAM JAMES), a very celebrated Dutch mathematician and philosopher, was born at Bois-le-duc, Sept. 27, 1688. He studied the civil law at Leyden, but mathematical learning was his favourite amusement. When he had taken his doctor's degree, in 1707, he went and settled at the Hague, where he practised at the bar, and cultivated an acquaintance with learned men; with a Society of whom he published a periodical Review, entitled *Le Journal Littéraire*, which was continued without interruption from the year 1713 to the year 1722. The parts of it written or extracted by Gravefande were chiefly those relating to geometry and physics. But he enriched it also with several original pieces entirely of his own composition; viz, Remarks on the Construction of Pneumatical Engines: A Moral Essay on Lying: And a celebrated Essay on the Collision of Bodies; which, as it opposed the Newtonian philosophy, was attacked by Dr. Clarke, and many other learned men.

In 1715, when the States sent to congratulate George the 1st, on his accession to the throne, Dr. Gravefande was appointed secretary to the embassy. During his stay in England he was admitted a member of the Royal Society, and became intimately acquainted with Sir Isaac Newton. On his return to Holland, he was chosen professor of mathematics and astronomy at Leyden; where he had the honour of first teaching the Newtonian philosophy, which was then in its infancy. He died in 1742, at 54 years of age.

Gravefande was a man amiable in his private character, and respectable in his public one; for few men of letters have rendered more eminent services to their country. The ministers of the republic consulted him on all occasions when his talents were requisite to assist them, which his skill in calculation often enabled him to do in money matters. He was of great service as a decypherer, in detecting the secret correspondence of their enemies. And in his own profession none ever applied the powers of nature with more success, or to more useful purposes.

Of his publications, the principal are,

1. An Introduction to the Newtonian Philosophy; or, a Treatise on the Elements of Physics, confirmed by Experiments. This performance, being only a more perfect



perfect copy of his public lectures, was first printed in 1720; and hath since gone through many editions, with considerable improvements: the 6th edit. is in English, in 2 large vols. 4to, by Dr. Desaguliers, in 1747, under the title of Mathematical Elements of Natural Philosophy, confirmed by Experiments.

2. A treatise on the Elements of Algebra, for the use of Young Students; to which is added a Specimen of a Commentary on Newton's Universal Arithmetic; as also, A New Rule for determining the Form of an Assumed Infinite Series.

3. An Essay on Perspective. This was written at 19 years of age.

4. A New Theory of the Collision of Bodies.

5. A Course of Logic and Metaphysics.

With several smaller pieces.

His whole mathematical and philosophical works, except the first article above, were collected and published at Amsterdam, in 2 vols. 4to, to which is prefixed a critical account of his life and writings, by Professor Allamand.

GRAVITATION, the exercise of gravity, or the pressure a body exerts on another body beneath it by the power of gravity.

This is sometimes distinguished from gravity. Thus M. Maupertuis, in his *Figure de la Terre*, takes gravity for that force by which a body would fall to the earth supposed at rest; and Gravitation for the same, but diminished by the centrifugal force. It is only Gravitation, or gravity thus blended with the centrifugal force, that we can usually measure by our experiments. Methods however have been found to distinguish what remains of the primitive gravity, and what has been destroyed by the centrifugal force.

It is one of the laws of nature, discovered by Newton, and now received by all philosophers, that every particle of matter in nature gravitates towards every other particle; which law is the main principle in the Newtonian philosophy. But what is called Gravitation with respect to the gravitating body, is usually called attraction with respect to the body gravitated to. The planets, both primary and secondary, as also the comets, do all gravitate towards the sun, and towards each other; as well as the sun towards them; and that in proportion to the quantity of matter in each of them.

The Peripatetics &c hold, that bodies only gravitate or weigh when out of their natural places, and that Gravitation ceases when they are restored to the same, the purpose of nature being then fulfilled: and they maintain that the final cause of this faculty is only to bring elementary bodies to their proper place, where they may rest. But the moderns shew that bodies exercise gravity even when at rest, and in their proper places. This is particularly shewn of fluids; and it is one of the laws of hydrostatics, demonstrated by Boyle and others, that fluids gravitate in proprio loco, the upper parts pressing on the lower, &c.

For the laws of Gravitation of bodies in fluids specifically lighter or heavier than themselves, see SPECIFIC GRAVITY. Also for the centre or line or plane of Gravitation, see CENTRE, LINE, or PLANE.

GRAVITY, in Physics, the natural tendency or inclination of bodies towards the centre. And in this sense Gravity agrees with Centripetal force.

Gravity however is, by some, defined more generally as the natural tendency of one body towards another; and again by others still more generally as the mutual tendency of each body, and each particle of a body, towards all others: in which sense the word answers to what is more usually called Attraction. Indeed the terms Gravity, weight, centripetal force, and attraction, denote in effect all the same thing, only in different views and relations; all which however it is very common to confound, and use promiscuously. But, in propriety, when a body is considered as tending towards the earth, the force with which it so tends is called Gravity, Force of Gravity, or Gravitating Force; when the body is considered as immediately tending to the centre of the earth, it is called Centripetal Force; but when we consider the earth, or mass to which the body tends, it is called Attraction, or Attractive Force; and when it is considered in respect of an obstacle or another body in the way of its tendency, upon which it acts, it is called Weight.

Philosophers think differently on the subject of Gravity. Some consider it as an inactive property or innate power in bodies, by which they endeavour to join their centre. Others hold Gravity in this sense to be an occult quality, and to be exploded as such out of all sound philosophy. Newton, though he often calls it a vis, power, or property in bodies, yet explains himself, that he means nothing more by the word but the effect or phenomenon: he does not consider the principle, the cause by which bodies tend downwards, but the tendency itself, which is no occult quality, but a sensible phenomenon, be its causes what they may; whether a property essential to body, as some make it, or superadded to it, as others; or even an impulse of some body from without, as others.

It is a law of nature long observed, that all bodies near the earth have a Gravity or weight, or a tendency towards its centre, or at least perpendicular to its surface; which law the moderns, and especially Sir I. Newton, from certain observations have found to be much more extensive, and holding universally with respect to all known bodies and matter in nature. It is therefore at present acknowledged as a principle or law of nature, that all bodies, and all the particles of all bodies, mutually gravitate towards each other: from which single principle it is that Newton has happily deduced all the great phenomena of nature.

Hence Gravity may be distinguished into Particular and General.

Particular GRAVITY, is that which respects the earth, or by which bodies descend, or tend towards the centre of the earth; the phenomena or properties of which are as follow:

1. All circumterrestrial bodies do hereby tend towards a point, which is either accurately or very nearly the centre of magnitude of the terraqueous globe. Not that it is meant that there is really any virtue or charm in the point called the centre, by which it attracts bodies; but because this is the result of the gravitation of bodies towards all the parts of which the earth consists.

2. This point or centre is fixed within the earth, or at least has been so far as any authentic history reaches. For a consequence of its shifting, though ever so little, would



would be the overflowing of the low lands on that side of the globe towards which it should approach. Dr. Halley suggests, that it would well account for the universal deluge, to have the centre of gravitation removed for a time towards the middle of the then inhabited world; for the change of its place but the 2000th part of the radius of the earth, or about 2 miles, would be sufficient to lay the tops of the highest hills under water.

3. In all places equidistant from the centre of the earth, the force of Gravity is nearly equal. Indeed all parts of the earth's surface are not at equal distances from the centre, because the equatorial parts are higher than the polar parts by about 17 miles; as has been proved by the necessity of making the pendulum shorter in those places, before it will swing seconds. In the new Petersburg Transactions, vol. 6 and 7, M. Krafft gives a formula for the proportion of Gravity in different latitudes on the earth's surface, which is this:

$$y = (1 + 0.0052848 \sin^2 \lambda) g;$$

where  $g$  denotes the Gravity at the equator, and  $y$  the Gravity under any other latitude  $\lambda$ . On this subject, see also the articles DEGREE, and EARTH.

4. Gravity equally affects all bodies, without regard either to their bulk, figure, or matter: so that, abstracting from the resistance of the medium, the most compact and loose, the greatest and smallest bodies would all descend through an equal space in the same time; as appears from the quick descent of very light bodies in an exhausted receiver. The space which bodies do actually fall, in vacuo, is  $16\frac{1}{2}$  feet in the first second of time, in the latitude of London; and for other times, either greater or less than that, the spaces descended from rest are directly proportional to the squares of the times, while the falling body is not far from the earth's surface.

5. This power is the greatest at the earth's surface, from whence it decreases both upwards and downwards, but not both ways in the same proportion; for upwards the force of Gravity is less, or decreases, as the square of the distance from the centre increases, so that at a double distance from the centre, above the surface, the force would be only 1-4th of what it is at the surface; but below the surface, the power decreases in such sort that its intensity is in the direct ratio of the distance from the centre; so that at the distance of half a semidiameter from the centre, the force would be but half what it is at the surface; at  $\frac{1}{3}$  of a semidiameter the force would be  $\frac{1}{3}$ , and so on.

6. As all bodies gravitate towards the earth, so does the earth equally gravitate towards all bodies; as well as all bodies towards particular parts of the earth, as hills, &c, which has been proved by the attraction a hill has upon a plumb line, insensibly drawing it aside.—Hence the gravitating force of entire bodies consists of those of all their parts: for by adding or taking away any part of the matter of a body, its Gravity is increased or decreased in the proportion of the quantity of such portion to the whole mass. Hence also the gravitating powers of bodies, at the same distance from the centre, are proportional to the quantities of matter in the bodies.

*General or Universal GRAVITY*, is that by which all the planets tend to one another, and indeed by which

all the bodies and particles of matter in the universe tend towards one another.

The existence of the same principle of Gravitation in the superior regions of the heavens, as on the earth, is one of the great discoveries of Newton, who made the proof of it as easy as that on the earth. . . At first it would seem this was only conjecture with him: he observed that all bodies near the earth, and in its atmosphere, had the property of tending directly towards it; he soon conjectured that it probably extended much higher than any distance to which we could reach, or make experiments; and so on, from one distance to another, till he at length saw no reason why it might not extend as far as to the moon, by means of which she might be retained in her orbit as a stone in a sling is retained by the hand; and if so, he next inferred why might not a similar principle exist in the other great bodies in the universe, the sun and all the other planets, both primary and secondary, which might all be retained in their orbits, and perform their revolutions, by means of the same universal principle of gravitation.

These conjectures he soon realized and verified by mathematical proofs. Kepler had found out, by contemplating the motions of the planets about the sun, that the area described by a line connecting the sun and planet, as this revolved in its orbit, was always proportional to the time of its description, or that it described equal areas in equal times, in whatever part of its orbit the planet might be, moving always so much the quicker as its distance from the sun was less. And it is also found that the satellites, or secondary planets, respect the same law in revolving about their primaries. But it was soon proved by Newton, that all bodies moving in any curve line described on a plane, and which, by radii drawn to any certain point, describe areas about the point proportional to the times, are impelled or acted on by some power tending towards that point. Consequently the power by which all these planets revolve, and are retained in their orbits, is directed to the centre about which they move, viz, the primary planets to the sun, and the satellites to their several primaries.

Again, Newton demonstrated, that if several bodies revolve with an equable motion in several circles about the same centre, and that if the squares of their periodical times be in the same proportion as the cubes of their distances from the common centre, then the centripetal forces of the revolving bodies, by which they tend to their central body, will be in the reciprocal or inverse ratio of the squares of the distances. Or if bodies revolve in orbits approaching to circles, and the apses of those orbits be at rest, then also the centripetal forces of the revolving bodies will be reciprocally proportional to the squares of the distances. But it had been agreed on by the astronomers, and particularly Kepler, that both these cases obtain in all the planets. And therefore he inferred that the centripetal forces of all the planets are reciprocally proportional to the squares of the distances from the centres of their orbits.

Upon the whole it appears, that the planets are retained in their orbits by some power which is continually acting upon them: that this power is directed towards the centre of their orbits: that the intensity or efficacy of this power increases upon an approach



towards the centre, and diminishes on receding from the same, and that in the reciprocal duplicate ratio of the distances: and that, by comparing this centripetal force of the planets with the force of gravity on the earth, they are found to be perfectly alike, as may easily be shewn in various instances. For example, in the case of the moon, the nearest of all the planets. The rectilinear spaces described in any given time by a falling body, urged by any powers, reckoning from the beginning of its descent, are proportional to those powers. Consequently the centripetal force of the moon revolving in her orbit, will be to the force of Gravity on the surface of the earth, as the space which the moon would describe in falling during any small time, by her centripetal force towards the earth, if she had no circular motion at all, to the space a body near the earth would describe in falling by its Gravity towards the same.

Now by an easy calculation of those two spaces, it appears that the former force is to the latter, as the square of the semi-diameter of the earth is to the square of that of the moon's orbit. The moon's centripetal force therefore is equal to the force of Gravity; and consequently these forces are not different, but they are one and the same: for if they were different, bodies acted on by the two powers conjointly would fall towards the earth with a velocity double to that arising from the sole power of Gravity.

It is evident therefore that the moon's centripetal force, by which she is retained in her orbit, and prevented from running off in tangents, is the very power of Gravity of the earth extended thither. See Newton's Princip. lib. 1, prop. 45, cor. 2, and lib. 3, prop. 3; where the numeral calculation may be seen at full length.

The moon therefore gravitates towards the earth, and reciprocally the earth towards the moon. And this is also farther confirmed by the phenomena of the tides.

The like reasoning may also be applied to the other planets. For as the revolutions of the primary planets round the sun, and those of the satellites of Jupiter and Saturn round their primaries, are phenomena of the same kind with the revolution of the moon about the earth; and as the centripetal powers of the primary are directed towards the centre of the sun, and those of the satellites towards the centres of their primaries; and lastly as all these powers are reciprocally as the squares of the distances from the centres, it may safely be concluded that the power and cause are the same in all.

Therefore, as the moon gravitates towards the earth, and the earth towards the moon; so do all the secondaries to their primaries, and these to their secondaries; and so also do the primaries to the sun, and the sun to the primaries. Newton's Princip. lib. 3, prop. 4, 5, 6; Greg. Astron. lib. 1, sect. 7, prop. 46 and 47.

The laws of Universal Gravity are the same as those of bodies gravitating towards the earth, before laid down.

*Cause of GRAVITY.* Various theories have been advanced by the philosophers of different ages to account for this grand principle of Gravitation. The ancients, who were only acquainted with particular Gravity, or the tendency of sublunar bodies towards the earth, aimed no farther than a system that might answer the

more obvious phenomena of it. However, some hints are found concerning the Gravitation of celestial bodies in the account given of the doctrine of Thales and his successors; and it would seem that Pythagoras was still better acquainted with it, to which it is supposed he had a view in what he taught concerning the Harmony of the Spheres.

Aristotle and the Peripatetics content themselves with referring Gravity or weight to a native inclination in heavy bodies to be in their proper place or sphere, the centre of the earth. And Copernicus ascribes it to an innate principle in all parts of matter, by which, when separated from their wholes, they endeavour to return to them again the nearest way. In answer to Aristotle and his followers, who considered the centre of the earth as the centre of the universe, he observed that it was reasonable to think there was nothing peculiar to the earth in this principle of Gravity; that the parts of the sun, moon, and stars tended likewise to each other, and that their spherical figure was preserved in their various motions by this power. Copern. Revol. lib. 1, cap. 9. But neither of these systems assigns any physical cause of this great effect: they only amount to this, that bodies descend because they are inclined to descend.

Kepler, in his preface to the commentaries concerning the planet Mars, speaks of Gravity as of a power that was mutual between bodies, and says that the earth and moon tend towards each other, and would meet in a point so many times nearer to the earth than to the moon, as the earth is greater than the moon, if their motions did not hinder it. He adds, that the tides arise from the Gravity of the waters towards the moon. To him we also owe the important discovery of the analogy between the distances of the several planets from the sun, and the periods in which they complete their revolutions, viz, that the squares of their periodic times are always in the same proportion as the cubes of their mean distances from the sun. However, Kepler, Gassendi, Gilbert, and others, ascribe Gravity to a certain magnetic attraction of the earth; conceiving the earth to be one great magnet continually emitting effluvia, which take hold of all bodies, and draw them towards the earth. But this is inconsistent with the several phenomena.

Des Cartes and his followers, Rohault &c, attribute Gravity to an external impulse or trusion of some subtle matter. By the rotation of the earth, say they, all the parts and appendages of it necessarily endeavour to recede from the centre of rotation; but whence they cannot all actually recede, as there is no vacuum or space to receive them. But this hypothesis, founded on the supposition of a plenum, is overthrown by what has been since proved of the existence of a vacuum.

Dr. Hook inclines to an opinion much like that of Des Cartes. Gravity he thinks deducible from the action of a most subtle medium, which easily pervades and penetrates the most solid bodies; and which, by some motion it has, detruses all earthly bodies from it, towards the centre of the earth. Vossius too, and many others, give partly into the Cartesian notion, and suppose Gravity to arise from the diurnal rotation of the earth round its axis.

Dr. Halley, despairing of any satisfactory theory, chooses



chooses to have immediate recourse to the agency of the Deity. So Dr. Clarke, from a view of several properties of Gravity, concludes that it is no adventitious effect of any motion, or subtle matter, but an original and general law impressed by God on all matter, and preserved in it by some efficient power penetrating the very solid and intimate substance of it; being found always proportional, not to the surfaces of bodies or corpuscles, but to their solid quantity and contents. It should therefore be no more inquired why bodies gravitate, than how they came to be first put in motion. Aunot. in Rohault. Phys. part 1, cap. 11.

Gravesande, in his *Introduct. ad Philos. Newton.* contends that the cause of Gravity is utterly unknown; and that we are to consider it no otherwise than as a law of nature originally and immediately impressed by the Creator, without any dependence on any second law or cause at all. Of this he thinks the three following considerations sufficient proof. 1. That Gravity requires the presence of the gravitating or attracting body: so the satellites of Jupiter, for ex. gravitate towards Jupiter, wherever he may be. 2. That the distance being supposed the same, the velocity with which bodies are moved by the force of Gravity, depends on the quantity of matter in the attracting body: and the velocity is not changed, whatever the mass of the gravitating body may be. 3. That if Gravity do depend on any known law of motion, it must be some impulse from an extraneous body; so that as Gravity is continual, a continual stroke must also be required. Now if there be any such matter continually striking on bodies, it must be fluid, and subtle enough to penetrate the substance of all bodies: but how shall a body subtle enough to penetrate the substance of the hardest bodies, and so rare as not sensibly to hinder the motion of bodies, be able to impel vast masses towards each other with such force? how does this force increase the ratio of the mass of the body, towards which the other body is moved? whence is it that all bodies move with the same velocity, the distance and body gravitated to being the same? can a fluid which only acts on the surface either of the bodies themselves, or their internal particles, communicate such a quantity of motion to bodies, which in all bodies shall exactly follow the proportion of the quantity of matter in them?

Mr. Cotes goes yet farther. Giving a view of Newton's philosophy, he asserts that Gravity is to be ranked among the primary qualities of all bodies; and deemed equally essential to matter as extension, mobility, or impenetrability. *Præfat. ad Newt. Princip.* But Newton himself disclaims this notion; and to shew that he does not take Gravity to be essential to bodies, he declares his opinion of the cause; choosing to propose it by way of query, not being yet sufficiently satisfied about its experiments. Thus, after having shewn that there is a medium in nature vastly more subtle than air, by whose vibrations sound is propagated, by which light communicates heat to bodies, and by the different densities of which the refraction and reflection of light are performed; he proceeds to inquire: "Is not this medium much rarer within the dense bodies of the sun, stars, planets, and comets, than

in the empty celestial spaces between them? and in passing from them to greater distances, doth it not grow denser and denser perpetually, and thereby cause the Gravity of those great bodies towards one another, and of their parts towards the bodies; every body endeavouring to recede from the denser parts of the medium towards the rarer?"

For if this medium be supposed rarer within the sun's body than at its surface, and rarer there than at the hundredth part of an inch from his body, and rarer there than at the fiftieth part of an inch from his body, and rarer there than at the orb of Saturn; I see no reason why the increase of density should stop any where, and not rather be continued through all distances from the Sun to Saturn, and beyond.

And though this increase of density may at great distances be exceeding slow; yet if the elastic force of this medium be exceeding great, it may suffice to impel bodies from the denser parts of the medium towards the rarer with all that power which we call Gravity.

And that the elastic force of this medium is exceeding great, may be gathered from the swiftness of its vibrations. Sounds move about 1140 English feet in a second of time, and in seven or eight minutes of time, they move about one hundred English miles: light moves from the Sun to us in about seven or eight minutes of time, which distance is about 70000000 English miles, supposing the horizontal parallax of the Sun to be about twelve seconds; and the vibrations, or pulses of this medium, that they may cause the alternate fits of easy transmission, and easy reflection, must be swifter than light, and by consequence above 700000 times swifter than sounds; and therefore the elastic force of this medium, in proportion to its density, must be above  $700000 \times 700000$  (that is, above 490000000000) times greater than the elastic force of the air is in proportion to its density: for the velocities of the pulses of elastic mediums are in a subduplicate ratio of the elasticities and the rarities of the mediums taken together.

As Magnetism is stronger in small loadstones than in great ones, in proportion to their bulk; and Gravity is stronger on the surface of small planets, than those of great ones, in proportion to their bulk; and small bodies are agitated much more by electric attraction than great ones: so the smallness of the rays of light may contribute very much to the power of the agent by which they are refracted; and if any one should suppose, that æther (like our air) may contain particles which endeavour to recede from one another (for I do not know what this æther is), and that its particles are exceedingly smaller than those of air, or even than those of light; the exceeding smallness of such particles may contribute to the greatness of the force, by which they recede from one another, and thereby make that medium exceedingly more rare and elastic than air, and of consequence, exceedingly less able to resist the motions of projectiles, and exceedingly more able to press upon gross bodies by endeavouring to expand itself." *Optics*, p. 325 &c.

GRAVITY, in Mechanics, denotes the conatus or tendency of bodies towards the centre of the earth.—That part of mechanics which considers the equilibrium  
or



or motion of bodies arising from Gravity or weight, is particularly called Statics.

Gravity in this view is distinguished into Absolute and Relative.

*Absolute GRAVITY* is that with which a body descends freely and perpendicularly through an unresisting medium. The laws of which see under DESCENT OF BODIES, ACCELERATION, MOTION, &c.

*Relative GRAVITY* is that with which a body descends on an inclined plane, or through a resisting medium, or as opposed by some other resistance. The laws of which see under the articles INCLINED PLANE, DESCENT, FLUID, RESISTANCE, &c.

*GRAVITY*, in Hydrostatics. The laws of bodies gravitating in Fluids make the business of Hydrostatics.

Gravity is here divided into Absolute and Specific.

*Absolute or True GRAVITY*, is the whole force with which the body tends downwards.

*Specific GRAVITY*, is the relative, comparative, or apparent Gravity in any body, in respect of that of an equal bulk or magnitude of another body; denoting that Gravity or weight which is peculiar to each species or kind of body, and by which it is distinguished from all other kinds.

In this sense a body is said to be Specifically Heavier than another, when under the same bulk it contains a greater weight than that other; and reciprocally the latter is said to be Specifically Lighter than the former. Thus, if there be two equal spheres, each one foot in diameter; the one of lead, and the other of wood: since the leaden one is found heavier than the wooden one, it is said to be Specifically, or in Specie, Heavier; and the wooden one Specifically Lighter.

This kind of Gravity is by some called Relative; in opposition to Absolute Gravity, which increases in proportion to the quantity or mass of the body.

#### *Laws of the SPECIFIC GRAVITY of bodies.*

I. If two bodies be equal in bulk, their specific gravities are to each other as their weights, or as their densities.

II. If two bodies be of the same specific gravity or density, their absolute weights will be as their magnitudes or bulks.

III. In bodies of the same weight, the specific gravities are reciprocally as their bulks.

IV. The specific gravities of all bodies are in a ratio compounded of the direct ratio of their weights, and the reciprocal ratio of their magnitudes. And hence again the specific gravities are as the densities.

V. The absolute gravities or weights of bodies are in the compound ratio of their specific gravities and magnitudes or bulks.

VI. The magnitudes of bodies are directly as their weights, and reciprocally as their specific gravities.

VII. A body specifically heavier than a fluid, loses as much of its weight when immersed in it, as is equal to the weight of a quantity of the fluid of the same bulk or magnitude.

Hence, since the Specific Gravities are as the abso-

lute gravities under the same bulk; the Specific Gravity of the fluid, will be to that of the body immersed, as the part of the weight lost by the solid, is to the whole weight.

And hence the Specific Gravities of fluids are as the weights lost by the same solid immersed in them.

VIII. *To find the Specific Gravity of a Fluid, or of a Solid.*—On one arm of a balance suspend a globe of lead by a fine thread, and to the other fasten an equal weight, which may just balance it in the open air. Immerse the globe into the fluid, and observe what weight balances it then, and consequently what weight is lost, which is proportional to the Specific Gravity as above. And thus the proportion of the Specific Gravity of one fluid to another is determined by immersing the globe successively in all the fluids, and observing the weights lost in each, which will be the proportions of the Specific Gravities of the fluids sought.

This same operation determines also the Specific Gravity of the solid immersed, whether it be a globe or of any other shape or bulk, supposing that of the fluid known. For the Specific Gravity of the fluid is to that of the solid, as the weight lost is to the whole weight.

Hence also may be found the Specific Gravity of a body that is lighter than the fluid, as follows:

IX. *To find the Specific Gravity of a Solid that is lighter than the fluid, as water, in which it is put.*—Annex to the lighter body another that is much heavier than the fluid, so as the compound mass may sink in the fluid. Weigh the heavier body and the compound mass separately, both in water and out of it; then find how much each loses in water, by subtracting its weight in water from its weight in air; and subtract the less of these remainders from the greater.

Then, As this last remainder,

Is to the weight of the light body in air,  
So is the Specific Gravity of the fluid,  
To the Specific Gravity of that body.

X. The Specific Gravities of bodies of equal weight, are reciprocally proportional to the quantities of weight lost in the same fluid. And hence is found the ratio of the Specific Gravities of solids, by weighing in the same fluids, masses of them that weigh equally in air, and noting the weights lost by each.

The Specific Gravities of many kinds of bodies, both solid and fluid, have been determined by various authors. Marinus Ghetaldus particularly tried the Specific Gravities of various bodies, especially metals; which were taken from thence by Oughtred. In the Philos. Transf. are several ample tables of them, by various authors, particularly those of Mr. Davis, vol. 45, p. 416, or Abr. vol. 10, p. 206. Some tables of them were also published by P. Merfenne, Muschenbroeck, Ward, Cotes, Emerson, Martin, &c.

It will be sufficient here to give those of some of the most usual bodies, that have been determined with the greater certainty. The numbers in this table express the number of Avoirdupois ounces in a cubic foot of each body, that of common water being just 1000 ounces, or  $62\frac{1}{2}$  lb.



## TABLE of SPECIFIC GRAVITIES.

## I. Solids.

Platina, pure	23000
Fine gold	19640
Standard gold	18888
Lead	11325
Fine Silver	11091
Standard Silver	10535
Copper	9000
Copper halfpence	8915
Gun metal	8784
Fine brass	8350
Cast brass	8000
Steel	7850
Iron	7645
Pewter	7471
Cast Iron	7425
Tin	7320
Lapis calaminaris	5000
Loadstone	4930
Mean of the whole Earth	4500
Crude Antimony	4000
Diamond	3517
Granite	3500
White lead	3160
Island crystal	2720
Marble	2705
Pebble stone	2700
Jasper	2666
Rock crystal	2650
Pearl	2630
Green glass	2600
Flint	2570
Onyx stone	2510
Common stone	2500
Crystal	2210
Clay	2160
Oyster shells.	2092
Brick	2000
Common earth	1984
Nitre	1900
Vitriol	1880
Alabaster	1874
Horn	1840
Ivory	1825
Sulphur	1810
Chalk	1793
Solid gunpowder	1745
Alum	1714
Dry bone	1660
Human calculus	1542
Sand	1520
Lignum vitæ	1327
Coal	1250
Jet	1238
Ebony	1177
Pitch	1150
Rosin	1100
Mahogany	1063
Amber	1040
Brazil wood	1031
Boxwood	1030
Common water	1000

Bees wax	955
Butter	940
Oak	925
Gunpowder, shaken	922
Logwood	913
Ice	908
Ash	800
Maple	755
Beech	700
Elm	600
Fir	550
Sassafras wood	482
Charcoal	
Cork	240
New fallen snow	86

## II. Fluids.

Quicksilver	13600
Oil of Vitriol	1700
Oil of Tartar	1550
Honey	1450
Spirit of Nitre	1315
Aqua Fortis	1300
Treacle	1290
Aqua Regia	1234
Human blood	1054
Urine	1032
Cow's milk	1031
Sea Water	1030
Ale	1028
Vinegar	1026
Tar	1015
Water	1000
Distilled Water	993
Red Wine	990
Proof Spirits	931
Olive Oil	913
Pure Spirits of Wine	866
Oil of Turpentine	800
Æther	726
Common Air	1.232
or very nearly	1.28

These numbers being the weight of a cubic foot, or 1728 cubic inches, of each of the bodies, in Avoirdupois ounces, by proportion the quantity in any other weight, or the weight of any other quantity, may be readily known.

For ex. Required the content of an irregular block of common stone which weighs 1 cwt, or 112lb, or 1792 ounces. Here, as  $2500 : 1792 :: 1728 : 1228\frac{4}{5}$  cubic inches the content.

Ex. 2. To find the weight of a block of granite, whose length is 63 feet, and breadth and thickness each 12 feet; being the dimensions of one of the stones, of granite, in the walls of Balbeck. Here,  $63 \times 12 \times 12 = 9072$  feet is the content of the stone; therefore as  $1 : 9072 :: 3500 \text{ oz} : 31752000 \text{ oz}$  or 885 tons 18 cwt. 3 qrs. the weight of the stone.

XI. A body descends in a fluid specifically lighter, or ascends in a fluid specifically heavier, with a force equal to the difference between its weight and that of an equal bulk of the fluid.

XII. A body sinks in a fluid specifically heavier, so far.



far as that the weight of the body is equal to the weight of a quantity of the fluid of the same bulk as the part immersed. Hence, as the Specific Gravity of the fluid is to that of the body, so is the whole magnitude of the body, to the magnitude of the part immersed.

XIII. The Specific Gravities of equal solids are as their parts immersed in the same fluid.

The several theorems here delivered, are both demonstrable from the principles of mechanics, and are also equally conformable to experiment, which answers exactly to the calculation; as is abundantly evident from the courses of philosophical experiments, so frequently exhibited; where the laws of specific gravitation are well illustrated.

GREAT BEAR, one of the constellations in the northern hemisphere. See URSA MAJOR.

GREAT CIRCLES, of the Globe or Sphere, are those whose planes pass through the centre, dividing it into two equal parts or hemispheres, and therefore having the same centre and diameter with the sphere itself. The principal of these are, the equator, the ecliptic, the horizon, the meridians, and the two colures.

GREAT-CIRCLE SAILING, is the art of conducting a ship along the arc of a great circle. And it is also that part of the theory of navigation which treats of sailing in the arc of a great circle. See NAVIGATION.

GREAVES (JOHN), an eminent astronomer, antiquary and linguist, was born in 1602, being the eldest son of John Greaves rector of Colemore, near Alresford in Hampshire, and master of a grammar school, where his son of course was well grounded in the primary rules of literature. He then went to Baliol college in Oxford, in 1617; but afterward, on account of his skill in philosophy and polite literature, he was the first of five that were elected into Merton college. Having read over all the ancient Greek and Latin writers, he applied to the study of natural philosophy and mathematics; and having contracted an intimacy with Mr. Briggs, Savilian professor of geometry at Oxford, and Dr. Bainbridge, Savilian professor of astronomy there, he was animated by their examples to prosecute that study with the greatest industry; and not content with reading the writings of Purbach, Regiomontanus, Copernicus, Tycho Brahe, Kepler, and other celebrated astronomers of that and the preceding age, he made the ancient Greek, Arabian, and Persian authors familiar to him, having before gained an accurate skill in the oriental languages. These accomplishments procured him the professorship of geometry in Gresham-college London, in 1630; and at the same time he held his fellowship of Merton-college.

In a journey to the Continent, in 1635, he visited the celebrated Golius, professor of Arabic at Leyden, and Claud Hardy at Paris, to converse about the Persian language. Hence he passed through Italy, and accurately surveyed the venerable remains of antiquity at Rome, visiting and corresponding everywhere with the most learned men of every nation. After visiting Padua, Florence, and Leghorn, he hence embarked for Constantinople, where he arrived in 1638. From thence he passed over to Rhodes, and Alexandria in Egypt, where he staid four or five months, and made a great number of curious observations. He next went to

Grand Cairo, measured the pyramids; and while there he adjusted the measure of the foot, observed by all nations. From hence he returned again through Italy, and arrived in England in the year 1640, after storing his mind with a variety of curious knowledge, and collecting many valuable oriental manuscripts and ancient curiosities; and while at Rome he made a particular inquiry into the true state of the ancient weights and measures.

On the death of Dr. John Bainbridge, in 1643, he was chosen Savilian professor of astronomy at Oxford, and principal reader of Linacre's lecture in Merton college; an appointment for which he was eminently qualified, from his critical acquaintance with the works of the ancient and modern astronomers. In 1645 he proposed a method of reforming the calendar, by omitting the intercalary day for 40 years to come: the paper relating to which, was published by Dr. Thomas Smith, in the Philos. Trans. for 1699. In 1646, he published his *Pyramidographia*, or a Description of the Pyramids of Egypt; and, in 1647, his *Discourse on the Roman Foot and Denarius*; from which, as from two principles, the measures and weights used by the Ancients may be deduced. He also published several other curious works concerning antiquities, &c.

Soon after publishing the last mentioned book, he was ejected, by the parliament visitors, from the professorship of astronomy and fellowship of Merton-college; and the soldiers committed many outrages, breaking open his chests, and destroying many of his manuscripts; which greatly affected him. On this occasion he retired to London, where he afterwards married, and prosecuted his studies with great vigour, as appears from several of his philosophical and theological writings. This however proved but a transient happiness to him; for he died at London, the 8th of October 1652, before he was quite 50 years of age; and left his astronomical instruments to the Savilian library in Oxford, where they are deposited.

GREEK ORDERS, in Architecture, are the Doric, Ionic and Corinthian; in contradistinction to the two Latin orders, viz the Tuscan and Composite.

GREEN, One of the original colours of the rays of light, or of the prismatic colours exhibited by the refraction of the rays of light.

*Green* is the pleasantest of all the colours to the sight. And hence it has been inferred as a proof of the wisdom and goodness of the Deity, that almost all vegetables, cloathing the surface of the earth, are green; which they are when growing in the open air; though those in subterraneous places, or places inaccessible to fresh air, are white or yellow. See CHROMATICS, and COLOURS.

GREGORIAN CALENDAR, so called from Pope Gregory the 13th, is the new or reformed Calendar, shewing the new and full moons, with the time of Easter, and the other moveable feasts depending upon it, by means of epacts disposed through the several months of the Gregorian year.

GREGORIAN *Epoch*, is the epoch or time, from which the Gregorian calendar, or computation, took place. This began in the year 1582; so that the year 1800 is the 218th of this epoch.



**GREGORIAN Telescope**, a particular sort of telescope, invented by Mr. James Gregory. See **TELESCOPE**.

**GREGORIAN Year**, the new account, or new style, introduced upon the reformation of the calendar, by Pope Gregory the 13th, in the year 1582, and from whom it took its name. This was introduced to reform the old, or Julian year, established by Julius Cæsar, which consisted of 365 days 6 hours, or 365 days and a quarter, that is three years of 365 days each, and the fourth year of 366 days. But as the mean tropical year consists only of 365ds 5hrs 48m 57sec. the former lost 11min. 3sec. every year, which in the time of Pope Gregory had amounted to 10 days, and who, by adding these 10 days, brought the account of time to its proper day again, and at the same time appointed that every century after, a day more should be added, thereby making the years of the complete centuries, viz 1600, 1700, 1800, &c, to be common years of 365 days each, instead of leap years of 366 days, which makes the mean Gregorian year equal to 365ds 5hrs 45m. 36sec.

This computation was not introduced into the account of time in England, till the year 1752, when the Julian account had lost 11 days, and therefore the 3d of September was in that year, by act of parliament, accounted the 14th, thereby restoring the 11 days which had thus been omitted. See **YEAR**.

**GREGORY (JAMES)**, professor of mathematics, first in the university of St. Andrews, and afterwards in that of Edinburgh, was one of the most eminent mathematicians of the 17th century. He was a son of the Rev. Mr. John Gregory minister of Drumoak in the county of Aberdeen, and was born at Aberdeen in November 1638. His mother was a daughter of Mr. David Anderson of Finzaugh, or Finshaugh, a gentleman who possessed a singular turn for mathematical and mechanical knowledge. This mathematical genius was hereditary in the family of the Andersons, and from them it seems to have been transmitted to their descendants of the names of Gregory, Reid, &c. Alexander Anderson, cousin-german of the said David, was professor of mathematics at Paris in the beginning of the 17th century, and published there several valuable and ingenious works; as may be seen in the memoirs of his life and writings, under the article **ANDERSON**. The mother of James Gregory inherited the genius of her family; and observing in her son, while yet a child, a strong propensity to mathematics, she instructed him herself in the elements of that science. His education in the languages he received at the grammar school of Aberdeen, and went through the usual course of academical studies in the Marischal college; but he was chiefly delighted with philosophical researches, into which a new door had lately been opened by the key of the mathematics. Galileo, Kepler, Des Cartes, &c, were the great masters of this new method: their works therefore became the principal study of young Gregory, who soon began to make improvements upon their discoveries in Optics. The first of these improvements was the invention of the reflecting telescope; the construction of which instrument he published in his *Optica Promota*, in 1663, at 24 years of age. This discovery soon attracted the attention of the mathematicians, both of our own and of foreign countries, who

immediately perceived its great importance to the sciences of optics and astronomy. But the manner of placing the two specula upon the same axis appearing to Newton to be attended with the disadvantage of losing the central rays of the larger speculum, he proposed an improvement on the instrument, by giving an oblique position to the smaller speculum, and placing the eye-glass in the side of the tube. It is observable however, that the Newtonian construction of that instrument was long abandoned for the original or Gregorian, which is now always used when the instrument is of a moderate size; though Herschel has preferred the Newtonian form for the construction of those immense telescopes, which he has of late so successfully employed in observing the heavens.

About the year 1664 or 1665, coming to London, he became acquainted with Mr. John Collins, who recommended him to the best optic glass-grinders there, to have his telescope executed. But as this could not be done for want of skill in the artists to grind a plate of metal for the object speculum into a true parabolic concave, which the design required, he was much discouraged with the disappointment; and after a few imperfect trials made with an ill-polished spherical one, which did not succeed to his wish, he dropped the pursuit, and resolved to make the tour of Italy, then the mart of mathematical learning, that he might prosecute his favourite study with greater advantage. And the university of Padua being at that time in high reputation for mathematical studies, Mr. Gregory fixed his residence there for some years. Here it was that he published, in 1667, *Vera Circuli et Hyperbolæ Quadratura*; in which he propounded another discovery of his own, the invention of an infinitely converging series for the areas of the circle and hyperbola. He sent home a copy of this work to his friend Mr. Collins, who communicated it to the Royal Society, where it met with the commendations of lord Brouncker and Dr. Wallis. He reprinted it at Venice the year following, to which he added a new work, entitled *Geometria Pars Universalis, inserviens Quantitatum Curvarum Transmutationi et Mensuræ*; in which he is allowed to have shewn, for the first time, a method for the transmutation of curves. These works engaged the notice, and procured the author the correspondence of the greatest mathematicians of the age, Newton, Huygens, Wallis, and others. An account of this piece was also read by Mr. Collins before the Royal Society, of which Mr. Gregory, being returned from his travels, was chosen a member the same year, and communicated to them an account of a controversy in Italy about the motion of the earth, which was denied by Riccioli and his followers.—Through this channel, in particular, he carried on a dispute with Mr. Huygens on the occasion of his treatise on the quadrature of the circle and hyperbola, to which that great man had started some objections; in the course of which our author produced some improvements of his series. But in this dispute it happened, as it generally does on such occasions, that the antagonists, though setting out with temper enough, yet grew too warm in the combat. This was the case here, especially on the side of Gregory, whose defence was, at his own request, inserted in the Philosophical Transactions. It is unnecessary to enter into particulars: suffice



suffice it therefore to say that, in the opinion of Leibnitz, who allows Mr. Gregory the highest merit for his genius and discoveries, M. Huygens has pointed out, though not errors, some considerable deficiencies in the treatise above mentioned, and shewn a much simpler method of attaining the same end.

In 1668, our author published at London another work, entitled, *Exercitationes Geometricæ*, which contributed still much farther to extend his reputation. About this time he was elected professor of mathematics in the university of St. Andrew's, an office which he held for six years. During his residence there, he married, in 1669, Mary, the daughter of George Jameson, the celebrated painter, whom Mr. Walpole has termed the Vandyke of Scotland, and who was fellow disciple with that great artist in the school of Rubens at Antwerp.

In 1672, he published "The Great and New Art of Weighing Vanity: or a Discovery of the Ignorance and Arrogance of the Great and New Artist, in his Pseudo-philosophical Writings. By M. Patrick Mathers, Arch-bedal to the University of St. Andrews. To which are annexed some Tentamina de Motu Penduli & Projectorum." Under this assumed name, our author wrote this little piece to expose the ignorance of Mr. Sinclair, professor at Glasgow, in his hydrostatical writings, and in return for some ill usage of that author to a colleague of Mr. Gregory's. The same year, Newton, on his wonderful discoveries in the nature of light, having contrived a new reflecting telescope, and made several objections to Mr. Gregory's, this gave birth to a dispute between those two philosophers, which was carried on during this and the following year, in the most amicable manner on both sides; Mr. Gregory defending his own construction, so far, as to give his antagonist the whole honour of having made the catoptric telescopes preferable to the dioptric; and shewing, that the imperfections in these instruments were not so much owing to a defect in the object speculum, as to the different refrangibility of the rays of light. In the course of this dispute, our author described a burning concave mirror, which was approved by Newton, and is still in good esteem. Several letters that passed in this dispute, are printed by Dr. Desaguliers, in an Appendix to the English edition of Dr. David Gregory's *Elements of Catoptrics and Dioptrics*.

In 1674, Mr. Gregory was called to Edinburgh, to fill the chair of mathematics in that university. This place he had held but little more than a year, when, in October 1675, being employed in shewing the satellites of Jupiter through a telescope to some of his pupils, he was suddenly struck with total blindness, and died a few days after, to the great loss of the mathematical world, at only 37 years of age.

As to his character, Mr. James Gregory was a man of a very acute and penetrating genius. His temper seems to have been warm, as appears from his conduct in the dispute with Huygens; and, conscious perhaps of his own merits as a discoverer, he seems to have been jealous of losing any portion of his reputation by the improvements of others upon his inventions. He possessed one of the most amiable characters of a true philosopher, that of being content with his fortune in his situation. But the most brilliant part of his character

is that of his mathematical genius as an inventor, which was of the first order; as will appear by the following list of his inventions and discoveries. Among many others may be reckoned, his Reflecting Telescope;—Burning Concave Mirror;—Quadrature of the Circle and Hyperbola, by an infinite converging series;—his method for the Transformation of Curves;—a Geometrical Demonstration of lord Brounker's series for Squaring the Hyperbola—his Demonstration that the Meridian Line is analogous to a scale of Logarithmic Tangents of the Half Complements of the Latitude;—he also invented and demonstrated geometrically, by help of the hyperbola, a very simple converging series for making the logarithms;—he sent to Mr. Collins the solution of the famous Keplerian problem by an infinite series;—he discovered a method of drawing Tangents to Curves geometrically, without any previous calculations;—a rule for the Direct and Inverse method of Tangents, which stands upon the same principle (of exhaustions) with that of fluxions, and differs not much from it in the manner of application; a Series for the length of the Arc of a Circle from the Tangent, and vice versa; as also for the Secant and Logarithmic Tangent and Secant, and vice versa:—These, with others, for measuring the length of the elliptic and hyperbolic curves, were sent to Mr. Collins, in return for some received from him of Newton's, in which he followed the elegant example of this author, in delivering his series in simple terms, independent of each other. These and other writings of our author are mostly contained in the following works, viz.,

1. *Optica Promota*; 4to, London 1663.
2. *Vera Circuli et Hyperbolæ Quadratura*; 4to, Padua 1667 and 1668.
3. *Geometriæ Pars Universalis*; 4to, Padua 1668.
4. *Exercitationes Geometricæ*; 4to, London 1668.
5. *The Great and New Art of Weighing Vanity, &c.* 8vo, Glasgow 1672.

The rest of his inventions make the subject of several letters and papers, printed either in the *Philos. Transf.* vol. 3; the *Commerc. Epistol. Joh. Collins et Aliorum*, 8vo, 1715; in the Appendix to the English edition of Dr. David Gregory's *Elements of Optics*, 8vo, 1735, by Dr. Desaguliers; and some series in the *Exercitatio Geometrica* of the same author, 4to, 1684, Edinburgh; as well as in his little piece on Practical Geometry.

GREGORY (*Dr. David*), Savilian professor of astronomy at Oxford, was nephew of the above-mentioned Mr. James Gregory, being the eldest son of his brother Mr. David Gregory of Kinardie, a gentleman who had the singular fortune to see three of his sons all professors of mathematics, at the same time, in three of the British Universities, viz., our author David at Oxford, the second son James at Edinburgh, and the third son Charles at St. Andrews. Our author David, the eldest son, was born at Aberdeen in 1661, where he received the early parts of his education, but completed his studies at Edinburgh; and, being possessed of the mathematical papers of his uncle, soon distinguished himself likewise as the heir of his genius. In the 23d year of his age, he was elected professor of mathematics in the university of Edinburgh; and, in the same year, he published *Exercitatio Geometrica de Dimensione Figurarum*,



*Figurarum, sive Specimen Methodi generalis Dimetiendi quasvis Figuras*, Edinb. 1684, 4to. He very soon perceived the excellence of the Newtonian philosophy; and had the merit of being the first that introduced it into the schools, by his public lectures at Edinburgh. "He had (says Mr. Wiston, in the Memoirs of his own Life, i. 32) already caused several of his scholars to keep acts, as we call them, upon several branches of the Newtonian philosophy; while we at Cambridge, poor wretches, were ignominiously studying the fictitious hypothesis of the Cartesian."

In 1691, on the report of Dr. Bernard's intention of resigning the Savilian professorship of astronomy at Oxford, our author went to London; and being patronised by Newton, and warmly befriended by Mr. Flamsteed the astronomer royal, he obtained the vacant professorship, though Dr. Halley was a competitor. This rivalry, however, instead of animosity, laid the foundation of friendship between these eminent men; and Halley soon after became the colleague of Gregory, by obtaining the professorship of geometry in the same university. Soon after his arrival in London, Mr. Gregory had been elected a fellow of the Royal Society; and, previously to his election into the Savilian professorship, had the degree of doctor of physic conferred on him by the university of Oxford.

In 1693, he published in the *Philos. Transf.* a resolution of the Florentine problem *de Testudine veliformi quadrabili*; and he continued to communicate to the public, from time to time, many ingenious mathematical papers by the same channel.

In 1695, he printed at Oxford, *Catoptrica et Dioptrica Spherica Elementa*; a work which, we are informed in the preface, contains the substance of some of his public lectures read at Edinburgh, eleven years before. This valuable treatise was republished in English, first with additions by Dr. William Brown, with the recommendation of Mr. Jones and Dr. Desaguliers; and afterwards by the latter of these gentlemen, with an appendix containing an account of the Gregorian and Newtonian telescopes, together with Mr. Hadley's tables for the construction of both those instruments. It is not unworthy of remark, that, in the conclusion of this treatise, there is an observation which shews, that the construction of achromatic telescopes, which Mr. Dollond has carried to such great perfection, had occurred to the mind of David Gregory, from reflecting on the admirable contrivance of nature in combining the different humours of the eye. The passage is as follows: "Perhaps it would be of service to make the object lens of a different medium, as we see done in the fabric of the eye; where the crystalline humour (whose power of refracting the rays of light differs very little from that of glass) is by nature, who never does any thing in vain, joined with the aqueous and vitreous humours (not differing from water as to their power of refraction) in order that the image may be painted as distinct as possible upon the bottom of the eye."

In 1702 our author published at Oxford, in folio, *Astronomiæ Physicæ et Geometricæ Elementa*; a work which is accounted his master-piece. It is founded on the Newtonian doctrines, and was esteemed by New-

ton himself as a most excellent explanation and defence of his philosophy. In the following year he gave to the world an edition, in folio, of the works of Euclid, in Greek and Latin; being done in prosecution of a design of his predecessor Dr. Bernard, of printing the works of all the ancient mathematicians. In this work, which contains all the treatises that have been attributed to Euclid, Dr. Gregory has been careful to point out such as he found reason, from internal evidence, to believe to be the productions of some inferior geometrician. In prosecution of the same plan, Dr. Gregory engaged soon after, with his colleague Dr. Halley, in the publication of the Conics of Apollonius; but he had proceeded only a little way in this undertaking, when he died at Maidenhead in Berkshire, in 1710, being the 49th year of his age only.

Besides those works published in our author's life time, as mentioned above, he had several papers inserted in the *Philos. Transf.* vol. 18, 19, 21, 24, and 25, particularly a paper on the catenarian curve, first considered by our author. He left also in manuscript, *A Short Treatise of the Nature and Arithmetic of Logarithms*, which is printed at the end of Keill's translation of Commandine's Euclid; and a *Treatise of Practical Geometry*, which was afterwards translated, and published in 1745, by Mr. Maclaurin.

Dr. David Gregory married, in 1695, Elizabeth, the daughter of Mr. Oliphant of Langtown in Scotland. By this lady he had four sons, of whom, the eldest, David, was appointed regius professor of modern history at Oxford by king George the 1st, and died at an advanced age in 1767, after enjoying for many years the dignity of dean of Christchurch in that university.

When David Gregory quitted Edinburgh, he was succeeded in the professorship at that university by his brother James, likewise an eminent mathematician; who held that office for 33 years, and, retiring in 1725, was succeeded by the celebrated Maclaurin. A daughter of this professor James Gregory, a young lady of great beauty and accomplishments, was the victim of an unfortunate attachment, that furnished the subject of Mallet's well known ballad of *William and Margaret*.

Another brother, Charles, was created professor of mathematics at St. Andrews by Queen Anne, in 1707. This office he held with reputation and ability for 32 years; and, resigning in 1739, was succeeded by his son, who eminently inherited the talents of his family, and died in 1763.

*Some farther Particulars of the Family of the Gregorys and Andersons, communicated by Dr. Thomas Reid, Professor of Moral Philosophy in the University of Glasgow, a Nephew of the late Dr. David Gregory Savilian Professor at Oxford.*

Some account of the family of the Gregorys at Aberdeen, is given in the Life of the late Dr. John Gregory prefixed to his works, printed at Edinburgh for A. Strahan and T. Cadell, London, and W. Creech, Edinburgh, 1788, in four small 8vo volumes.

Who was the author of that Life, or whence he had his information, I do not know. I have heard it ascribed to Mr. Tytler the younger, whose father was appointed



one of the guardians of Dr. John Gregory's children. Some additions to what is contained in it, and remarks upon it, is all I can furnish upon this subject.

Page 3. I know nothing of the education of David Anderson of Finzaugh. He seems to have been a self-taught Engineer. Every public work which surpassed the skill of common artists, was committed to the management of David. Such a reputation he acquired by his success in works of this kind, that with the vulgar he got the by-name of *Davie do a' thing*, that is in the Scottish dialect, *David who could do every thing*. By this appellation he is better known than by his proper name. He raised the great bells into the steeple of the principal church: he cut a passage for ships of burden through a ridge of rock under water, which crossed the entrance into the harbour of Aberdeen. In a long picture gallery at Cullen House, the seat of the earl of Findlater, the wooden ceiling is painted with several of the fables of Ovid's *Metamorphosis*. The colours are still bright, and the representation lively. The present earl's grandfather told me that this painting was the work of David Anderson my ancestor, whom he acknowledged as a friend and relation of his family.

Such works, while they gave reputation to David, suited ill with his proper business, which was that of a merchant in Aberdeen. In that he succeeded ill; and having given up mercantile business, from a small remainder of his fortune began a trade of making malt; and having instructed his wife in the management of it, left it to her care, and went into England to try his fortune as an engineer; an employment which in his own country he had practised gratuitously. Having in that way made a fortune which satisfied him, he returned to Aberdeen, where his wife had also made money by her malting business.

After making such provision for their children as they thought reasonable, they agreed that the longest liver of the two should enjoy the remainder, and at death should bequeath it to certain purposes in the management of the magistrates of Aberdeen.

The wife happened to live longest, and fulfilled what had been concerted with her husband. Her legacies, well known in Aberdeen, are called after her name *Jane Guild's Mortifications*, a mortification in Scots law signifying a bequeathment for some charitable purpose. They consist of sums for different purposes. For orphans, for the education of boys and girls, for unmarried gentlewomen, and for widows; and they still continue to be useful to many in indigent circumstances. She was the daughter of Dr. Guild a minister of Aberdeen. Besides her money, she bequeathed a piece of tapestry, wrought by her own hand, and representing the history of queen Esther, from a drawing made by her husband. The tapestry continues to ornament the wall of the principal church.

In the same page it is said that Alexander Anderson, professor of mathematics at Paris, was the cousin-german of David above-mentioned. I know not the writer's authority for this: I have always heard that they were brothers; but for this I have only family tradition.

P. 4. It is here said that James Gregory was in-

structed in the *Elements of Euclid* by his mother, the daughter of David Anderson.

The account I have heard differs from this. It is, that his brother David, being ten or eleven years older, had the direction of his education after their father's death, and, when James had finished his course of philosophy, was at a loss to what literary profession he should direct him. After some unsuccessful trials, he put Euclid's *Elements* into his hand, and finding that he applied to Euclid with great avidity and success, he encouraged and assisted him in his mathematical studies.

This tradition agrees with what James Gregory says in the preface to his *Optica Promota*; where after mentioning his advance to the 26th proposition, he adds, *Ubi diu hæsi omne spe progrediendi orbatus, sed continuis hortatibus et auxiliis fratris mei Davidis Gregorii, in Mathematicis non parum versati (cui si quid in hisce Scientiis præstitero, me illud debere non inficias ibo) animatus, tandem incidi &c.* Whether David had been instructed in mathematics by his mother, or had any living instructor, I know not.

P. 5, 6. In these two pages I think the merit of Gregory compared with that of Newton in the invention of the catoptric telescope, is put in a light more unfavourable to Newton than is just. Gregory believing that the imperfection of the dioptric telescope arose solely from the spherical figure of the glasses, invented his telescope to remedy that imperfection. Being less conversant in the practice of mechanics, he did not attempt to make any model. The specula of his telescope required a degree of polish and a figure which the best opticians of that age were unable to execute. Newton demonstrated that the imperfection of the dioptric telescope arose chiefly from the different refrangibility of the rays of light; he demonstrated also that the catoptric telescope required a degree of polish far beyond what was necessary for the dioptric. He made a model of his telescope; and finding that the best polish which the opticians could give, was insufficient, he improved the polish with his own hand, so as to make it answer the purpose, and has described most accurately the manner in which he did this. And, had he not given this example of the practicability of making a reflecting telescope, it is probable that it would have passed as an impracticable idea to this day.

P. 11. To what is said of this James Gregory might have been added, that he was led by analogy to the true law of Refraction, not knowing that it was discovered by Des Cartes before (see Preface to *Optica Promota*); and that in 1670 having received in a letter from Collins, a Series for the Area of the Zone of a Circle, and as Newton had invented an universal method by which he could square all Curves Geometrical and Mechanical by Infinite Series of that kind; Gregory after much thought discovered this universal method, or an equivalent one. Of this he perfectly satisfied Newton and the other mathematicians of that time, by a letter to Collins in Feb. 1671. He was strongly solicited by his brother David to publish his Universal Method of Series without delay, but excused himself upon a point of honour; that as Newton was the first inventor, and as he had

been.



been led to it by an account of Newton's having such a method, he thought himself bound to wait till Newton should publish his method. I have seen the letters that passed between the brothers on this subject.

With regard to the controversy between James Gregory and Huygens, I take the subject of that controversy to have been, not whether J. Gregory's Quadrature of the Circle by a converging series was just, but whether he had demonstrated, as in one of his propositions he pretended to do, That it is impossible to express perfectly the Area of a Circle in any known Algebraical form, besides that of an infinite converging series. Huygens excepted to the demonstration of this proposition, and Gregory defended it; neither of them convinced his antagonist, nor do I know that Leibnitz improved upon what Gregory had done.

P. 12. David Gregory of Kinardie deserved a more particular account than is here given.

It is true that he served an apprenticeship to a mercantile house in Holland, but he followed that profession no longer than he was under authority, having a stronger passion for knowledge than for money. He returned to his own country in 1655, being about 28 years of age, and from that time led the life of a philosopher. Having succeeded to the estate of Kinardie by the death of an elder brother, he lived there to the end of that century. There all his children were born, of whom he had thirty-two by two wives.

Kinardie is above 40 English miles north from Aberdeen, and a few miles from Bamf, upon the river Diveron. He was a jest among the neighbouring gentlemen for his ignorance of what was doing about his own farm, but an oracle in matters of learning and philosophy, and particularly in medicine, which he had studied for his amusement, and begun to practise among his poor neighbours. He acquired such a reputation in that science, that he was employed by the nobility and gentlemen of that county, but took no fees. His hours of study were singular. Being much occupied through the day with those who applied to him as a physician, he went early to bed, rose about two or three in the morning, and, after applying to his studies for some hours, went to bed again and slept an hour or two before breakfast.

He was the first man in that country who had a barometer; and by some old letters which I have seen, it appeared, that he had corresponded with some philosophers on the continent about the changes in the barometer and in the weather, particularly with Mariotte the French philosopher. He was once in danger of being prosecuted as a conjurer by the Presbytery on account of his barometer. A deputation of that body having waited upon him to enquire into the ground of certain reports that had come to their ears, he satisfied them so far as to prevent the prosecution of a man known to be so extensively useful by his knowledge of medicine.—About the beginning of this century he removed with his family to Aberdeen, and in the time of queen Anne's war employed his thoughts upon an improvement in artillery, in order to make the shot of great guns more destructive to the enemy, and executed a model of the engine he had conceived. I have conversed with a clock-maker in Aberdeen who was em-

ployed in making this model; but having made many different pieces by direction without knowing their intention, or how they were to be put together, he could give no account of the whole. After making some experiments with this model, which satisfied him, the old gentleman was so sanguine in the hope of being useful to the allies in the war against France, that he set about preparing a field equipage with a view to make a campaign in Flanders, and in the mean time sent his model to his son the Savilian professor, that he might have his and Sir Isaac Newton's opinion of it. His son shewed it to Newton, without letting him know that his own father was the inventor. Sir Isaac was much displeased with it, saying, that if it tended as much to the preservation of mankind as to their destruction, the inventor would have deserved a great reward; but as it was contrived solely for destruction, and would soon be known by the enemy, he rather deserved to be punished, and urged the professor very strongly to destroy it, and if possible to suppress the invention. It is probable the professor followed this advice. He died soon after, and the model was never found.

When the rebellion broke out in 1715, the old gentleman went a second time to Holland, and returned when it was over to Aberdeen, where he died about 1720, aged 93.

He left an historical manuscript of the Transactions of his own Time and Country, which my father told me he had read.

I was well acquainted with two of this gentleman's sons, and with several of his daughters, besides my own mother. The facts abovementioned are taken from what I have occasionally heard from them, and from other persons of his acquaintance.

P. 14. In confirmation of what is said in this page, that the two brothers David and James were the first who taught the Newtonian philosophy in the Scotch Universities; I have by me a *Thesis*, printed at Edinburgh in 1690, by James Gregory, who was at that time a professor of philosophy at St. Andrews, and succeeded his brother David in the profession of mathematics at Edinburgh. In this *Thesis*, after a dedication to Viscount Tarbet, follow the names of twenty-one of his scholars who were candidates for the degree of A. M. then twenty-five positions or *Theses*. The first three relate to logic, and the abuse of it in the Aristotelian and Cartesian philosophy. He defines logic to be the art of making a proper use of things granted, in order to find what is sought, and therefore admits only two *Categories* in logic, viz, *Data* and *Quæsitæ*. The remaining twenty-two positions are a compend of Newton's Principia. This *Thesis*, as was the custom at that time in the Scotch universities, was to be defended in a public disputation, by the candidates, previous to their taking their degree.

The famous Dr. Pitcairn was a fellow student and intimate companion of these two Gregories, and during the vacation of the college was wont to go north with them to Kinardie, their father's house.

David Gregory was appointed a preceptor to the duke of Gloucester, queen Anne's son; but his entering upon that office was prevented by the death of that prince in the eleventh year of his age.

P. 19. D. Gregory's Euclid is said to have been wrote in



in prosecution of a design of his predecessor Dr. Bernard, of printing the works of all the antient mathematicians. This design ought to have been ascribed to Savile, who left in charge to the two professors of his foundation, to print the mathematical works of the antients, and I think left a fund for defraying the expence. Wallis did something in consequence of this charge; Gregory and Halley did a great deal; but I think nothing has been done in this design by the Savilian professors since their time.

P. 20. Besides what is mentioned, Dr. Gregory left in manuscript a Commentary on Newton's Principia, which Newton valued, and kept by him for many years after the author's death. It is probable that in what relates to astronomy, this commentary may coincide in a great measure with the author's astronomy, which indeed is an excellent Commentary upon that part of the Principia.

P. 24. This David Gregory published in Latin, a very good compend of arithmetic and algebra, with the title *Arithmetica et Algebrae Compendium, in Usum Juventutis Academicae*. Edinb. 1736. He had a design of publishing his uncle's Commentary on the Principia, with extracts from the papers left by James Gregory his grand uncle; but the expence being too great for his fortune, and he too gentle a solicitor of the assistance of others, the design was dropped. His son David, yet alive, was master of an East India ship.

P. 40. To the projectors of the society at Aberdeen, ought to have been added John Stewart professor of mathematics in the Marischal college at Aberdeen. He published an explanation of two treatises of Sir Isaac Newton, viz, his Quadrature of Curves, and his Analysis by Equations of an infinite number of terms. He was an intimate friend of Dr. Reid's.

Another of the first members of that society was Dr. David Skene, who, besides his eminence in the practice of medicine, had applied much to all parts of natural history, particularly to botany, and was a correspondent of the celebrated Linnæus.

Dr. John Gregory and Dr. David Skene were the first who attempted a college of medicine at Aberdeen. The first gave lectures to his pupils in the theory and practice of medicine, and in chemistry; the last, in anatomy, materia medica, and midwifery, in order to prepare them for attending the medical college at Edinburgh.

T. R.

*The following additional lines by Mr. James Millar, Professor of Mathematics, Glasgow.*

Another instance of the prevalence of mathematical genius in the family of Gregory or Anderson, whether produced by an original and inexplicable determination of the mind, or communicated by the force of example, and the consciousness of an intimate connection with a reputation already acquired in a particular line, is the celebrated Dr. Reid, professor of moral philosophy in the university of Glasgow; a nephew, by his mother, of the late Dr. David Gregory, Savilian professor at Oxford.

This gentleman, well known to the public by his moral and metaphysical writings, and remarkable for that liberality, and that ardent spirit of enquiry, which neither overlooks nor undervalues any branch of science, is

peculiarly distinguished by his abilities and proficiency in mathematical learning. The objects of literary pursuit are often directed by accidental occurrences. And apprehension of the bad consequences which might result from the philosophy of the late Mr. Hume, induced Dr. Reid to combat the doctrines of that eminent author; and produced a work, which has excited universal attention, and seems to have given a new turn to speculations upon that subject. But it is well known to Dr. Reid's literary acquaintance, that these exertions have not diminished the original bent of his genius, nor blunted the edge of his inclination for mathematical researches; which, at a very advanced age, he still continues to prosecute with a youthful attachment, and with unremitting assiduity.

It may farther be observed, of the extraordinary family above mentioned, that Dr. James Gregory, the present learned professor of physic and medicine in the university of Edinburgh, is the son of the late Dr. John Gregory, upon the memoirs of whose life the above remarks have been written by Dr. Reid; the said James has lately published a most ingenious work, intitled, *Philosophical and Literary Essays*, in 2 volumes 8vo, Edinb. 1792; and he seems to be another worthy inheritor of the singular genius of his family.

GREGORY (St. Vincent), a very respectable Flemish geometrician, was born at Bruges in 1584, and became a Jesuit at Rome at 20 years of age. He studied mathematics under the learned Jesuit Clavius. He afterward became a reputable professor of those sciences himself, and his instructions were solicited by several princes: he was called to Prague by the emperor Ferdinand the 2d; and Philip the 4th, king of Spain, was desirous of having him to teach mathematics to his son the young prince John of Austria. He was not less estimable for his virtues than his skill in the sciences. His well-meant endeavours were very commendable, when his holy zeal, though for a false religion, led him to follow the army in Flanders one campaign, to confess the wounded and dying soldiers, in which he received several wounds himself. He died of an apoplexy at Ghent, in 1667, at 83 years of age.

As a writer, Gregory St. Vincent was very diffuse and voluminous, but he was an excellent geometrician. He published, in Latin, three mathematical works, the principal of which was his *Opus Geometricum Quadraturæ Circuli, et Sectionum Coni*, Antwerp, 1647, 2 vol. folio. Although he has not demonstrated, in this work, the Quadrature of the circle, as he pretends to have done, the book nevertheless contains a great number of truths and important discoveries; one of which is this, viz, that if one asymptote of an hyperbola be divided into parts in geometrical progression, and from the points of division ordinates be drawn parallel to the other asymptote, they will divide the space between the asymptote and curve into equal portions; from whence it was shewn by Merenne, that, by taking the continual sums of those parts, there would be obtained areas in arithmetical progression, adapted to abscisses in geometrical progression, and which therefore were analogous to a system of logarithms.

GRENADÉ, or GRENADO. See GRANADÉ.

GRUS, the Crane, one of the new constellations, in the



the southern hemisphere; containing, according to Mr. Sharp's catalogue, 13 stars.

GRUS is also one of the Arabian constellations, and answers to our Ophiucus, to which they changed this constellation, their religion prohibiting them from drawing any human figures.

GRY, a measure containing one-tenth of a line. A line is one-tenth of a digit, and a digit is one-tenth of a foot, and a philosophical foot, one-third of a pendulum, whose diadromes, or vibrations, in the latitude of 45 degrees, are each equal to one second of time, or one-sixtieth of a minute.

GUARDS, a name that has been sometimes applied to the two stars nearest the north pole; being in the hind part of the chariot, at the tail of Urfa Minor or little bear; one of them being also called the pole star.

GUERICKE (OTTO or OTHO), counsellor to the elector of Brandenburg and burgomaster of Magdebourg, was born in 1602, and died in 1686 at Ham-bourg. He was one of the greatest philosophers of his time. It was Guericke that invented the air-pump; the two brass hemispheres, which being applied to each other, and the air exhausted, 16 horses were not able to draw them asunder; the marmouset of glass which descended in a tube in rainy weather, and rose again on the return of serene weather. This last machine fell into disuse on the invention of the barometer, especially after Huygens and Amontons gave theirs to the world. Guericke made use of his marmouset to foretell storms; from whence he was looked upon as a forcerer by the people; so that the thunder having one day fallen upon his house, and shivered to pieces several machines which he had employed in his experiments, they failed not to say it was a punishment from heaven that was angry with him.—Guericke was author of several works in natural philosophy, the principal of which was his *Experimenta Magdeburgica*, in folio, which contains his experiments on a vacuum.

GUERITE, in Fortification, a centry-box; being a small tower of wood, or stone, usually placed on the point of a bastion, or on the angles of the shoulder, to hold a centinel, who is to take care of the ditch, and watch against a surprise.

GUEULE, in Architecture. See GULA.

GUINEA, a gold coin struck in England.—The value or rate of the guinea has varied. It was at first equal to 20 shillings; but by the scarcity of gold it was afterwards advanced to 21*s.* 6*d.*; though it is now sunk to 21*s.*

The pound weight troy of gold is cut into 44 parts and a half, and each part makes a guinea, which is therefore equal to  $\frac{2}{3}$  lb, or  $\frac{2}{3}$  oz, or 5 dwts  $9\frac{2}{3}$  gr.

This coin took its name, Guinea, from the circumstance of the gold of which it was first struck being brought from that part of Africa called Guinea, for which reason also it bore the impression of an elephant.

GULA, GUEULE, or GOLA, in Architecture, a wavy member whose contour resembles the letter S, commonly called an Ogee.

GULBE, in Architecture, the same as Gorge.

GULF, or GULPH, in Geography, a part of the

ocean running up into the land through a narrow passage, or strait, and forming a bay within. As, the Gulf of Venice, or Adriatic sea; the Gulf of Arabia, or of Persia, which is the Red Sea; the Gulf of Constantinople, or the Black Sea; the Gulf of Mexico; &c.

GUN, a fire-arm, or weapon of offence, which forcibly discharges a ball or other matter through a cylindrical tube, by means of inflamed gun-powder.

The word Gun now includes most of the species of fire-arms; mortars and pistols being almost the only ones excepted from this denomination. They are divided into great and small guns: the former including all that are usually called cannon, ordnance, or artillery; and the latter includes musquets, firelocks, carabines, musketoons, blunderbusses, fowling-pieces, &c.

It is not certainly known at what time these weapons were first invented. And though the introduction of guns into the western part of the world is but of modern date, comparatively speaking; yet it is certain that in some parts of Asia they have been used for many ages, though in a very rude and imperfect manner. Philostratus speaks of a city near the river Hyphasis in the Indies, which was said to be impregnable, and that its inhabitants were relations of the gods, because they threw thunder and lightning upon their enemies; and other Greek authors, as also Quintus Curtius, speak of the same thing having happened to Alexander the Great. Hence some have imagined that guns were used by the eastern nations in his time, while others suppose the thunder and lightning alluded to by those authors, were only certain artificial fire-works, or rockets, such as we know are used in the wars by the Indians even in the present day against the Europeans. Be this however as it may, it is asserted by many modern travellers, that Guns were used in China as far back as the year of Christ 85, and have continued in use ever since.

The first hint of the invention of Guns in Europe, is in the works of Roger Bacon, who flourished in the 13th century. In a treatise written by him about the year 1280, he proposes to apply the violent explosive force of gun-powder for the destruction of armies. And though it is certainly known that the composition of gun-powder is described by Bacon in the said work, yet the invention has usually, though improperly, been ascribed to Bartholdus Schwartz, a German monk, who it is said discovered it only in the year 1320; and the invention is related in the following manner. Schwartz having, for some purpose, pounded nitre, sulphur, and charcoal together, in a mortar, which he afterwards covered imperfectly with a stone, a spark of fire accidentally fell into the mortar, which setting the mixture on fire, the explosion blew the stone to a considerable distance. Hence it is probable that Schwartz might be taught the simplest method of applying it in war; for it rather seems that Bacon conceived the manner of using it to be by the violent effort of the flame unconfined, and which is indeed capable of producing astonishing effects. (See GUNPOWDER.) And the figure and name of mortars given to a species of old artillery, and their employment, in throwing large stone bullets at an elevation, very much favour this conjecture.

Soon



Soon after the time of Schwartz, we find Guns commonly used as instruments of war. Great guns were first used. These were originally made of iron-bars soldered together, and fortified with strong iron hoops or rings; several of which are still to be seen in the Tower of London, and in the Warren at Woolwich. Others were made of thin sheets of iron rolled up together and hooped: and on particular emergencies some have been made of leather, and of lead, with plates of iron or copper. These first pieces were executed in a rude and imperfect manner, like the first essays of most new inventions. Stone balls were thrown out of them, and a small quantity of powder used on account of their weakness. They were of a cylindrical form, without ornaments, and were placed on their carriages by rings.

When, or by whom they were first made, is uncertain. It is known however that the Venetians used cannon at the siege of Claudia Jessa, now called Chioggia, in 1366, which were brought thither by two Germans, with some powder and leaden balls; as likewise in their wars with the Genoese in 1379. But before that, king Edward the 3d made use of cannon at the battle of Cressy in 1346, and at the siege of Calais in 1347. Cannon were employed by the Turks at the siege of Constantinople, then in possession of the Christians, in 1394, and in that of 1452, which threw a weight of 100lb; but they commonly burst at the 1st, 2d, or 3d firing. Louis the 12th had one cast at Tours, of the same size, which threw a ball from the Bastile to Charenton: one of these extraordinary cannon was taken at the siege of Dieu in 1546, by Don John de Castro, and is now in the castle of St. Julian da Barra, 10 miles from Lisbon: the length of is 20 feet 7 inches, its diameter at the middle 6 feet 3 inches, and it threw a ball of 100lb weight. It has neither dolphins, rings, nor button; is of an unusual kind of metal; and it has a large Indoſtan inscription upon it, which says it was cast in 1400.

Formerly, cannon were dignified with uncommon names. Thus, Lewis the 12th, in 1503, had 12 brass cannon cast, of an extraordinary size, called after the names of the 12 peers of France. The Spanish and Portuguese called them after their saints. The emperor Charles the 5th, when he marched against Tunis, founded the 12 apostles. At Milan there is a 70 pounder, called the Pimontelle; and one at Bois-le-duc, called the Devil. A 60 pounder at Dover-castle, called Queen Elizabeth's pocket-pistol. An 80 pounder in the Tower of London, brought there from Edinburgh-castle, called Mounts-meg. An 80 pounder in the royal arsenal at Berlin, called the Thunderer. An 80 pounder at Malaga, called the Terrible. Two curious 60 pounders in the arsenal at Bremen, called the Messenger of bad news. And lastly an uncommon 70 pounder in the castle of St. Angelo at Rome, made of the nails that fastened the copper-plates which covered the ancient Pantheon, with this inscription upon it, *Ex clavis trabalibus porticus Agrippæ*.

In the beginning of the 15th century these uncommon names were generally abolished, and the following more universal ones took place, viz,

Names.	Wt. of ball, Pounders.	Wt. of piece in cwt, about	
		Cwt.	
Cannon royal, or carthoun,	48	-	90
Bastard cannon, or $\frac{3}{4}$ carthoun,	36	-	79
Demi-carthoun,	24	-	60
Whole culverins,	18	-	50
Demi-culverins,	9	-	30
Falcon,	6	-	25
Sacker	largest size	-	18
	ordinary	-	15
	lowest fort	-	13
Basilisk	48	-	85
Serpentine	4	-	8
Aspic	2	-	7
Dragon	6	-	12
Syren	60	-	81
Falconet	3, 2, and 1	-	15, 10, 5
Rabinet	1	-	
Moyens	10 or 12 oz.	-	

These curious names of beasts and birds of prey were adopted on account of their swiftness in motion, or of their cruelty; as the falconet, falcon, sacker, and culverin, &c, for their swiftness in flying; the basilisk, serpentine, aspic, dragon, syren, &c, for their cruelty.

But, at present, cannon take their names from the weight of their proper ball. Thus, a piece that discharges a cast-iron ball of 24 pounds, is called a 24 pounder; one that carries a ball of 12 pounds, is called a 12 pounder; and so of the rest, divided into the following forts; viz,

Ship-guns, consisting in 42, 36, 32, 24, 18, 12, 9, 6, and 3 pounders.

Garrison-guns, in 42, 32, 24, 18, 12, 9, and 6 pounders.

Battering-guns, in 24, 18, and 12 pounders.

Field-pieces, in 12, 9, 6, 3, 2,  $1\frac{1}{2}$ , 1, and  $\frac{1}{2}$  pounders.

*Mortars*, it is thought, have been at least as ancient as cannon. They were employed in the wars of Italy, to throw balls of red-hot iron, stones, &c, long before the invention of shells. These last, it is supposed, were of German invention, and the use of them in war shewn by the following accident; viz, a citizen of Venlo, at a festival celebrated in honour of the duke of Cleves, throwing a number of shells, one of them fell on a house and set it on fire, by which misfortune the greatest part of the town was reduced to ashes. The first account of shells used for military purposes, is in 1435, when Naples was besieged by Charles the 8th. History informs us, with more certainty, that shells were thrown out of mortars at the siege of Wachtendonk, in Guelderland, in 1588, by the earl of Mansfield; and Cyprian Lucar wrote upon the method of filling and throwing such shells in his Appendix to the Colloquies of Tartaglia, printed at London in 1588; where also the compounding and throwing of carcases and various sorts of fire-works are shewn.

Mr. Malter, an English engineer, first taught the French the art of throwing shells, which they practised at the siege of Motte in 1634. The method of throwing red-hot balls out of mortars was first certainly put in



in practice at the siege of Stralsund in 1675 by the elector of Brandenburg: though some say in 1653 at the siege of Bremen.

Another species of ordnance has been long in use, by the name of Howitzer, which is a kind of medium as to its length, between the cannon and the mortar, and is a very useful piece, for discharging either shells or large balls, which is done either at point-blanc, or at a small elevation.

A new species of ordnance has lately been introduced by the Carron company, and thence called a Carronade, which is only a very short howitzer, and which possesses the advantage of being very light and easy to work.

The species of Guns before mentioned, are now made chiefly of cast iron; except the howitzer, which is of brass, as well as some cannon and mortars.

Muskets were first used at the siege of Rhege in the year 1521. The Spaniards were the first who armed part of their foot with these weapons. At first they were very heavy, and could not be used without a rest. They had match-locks, and did execution at a great distance. On their march the soldiers carried only the rests and ammunition, having boys to bear their muskets after them. They were very slow in loading, not only by reason of the unwieldiness of their pieces, and because they carried the powder and ball separate, but from the time it took to prepare and adjust the match; so that their fire was not near so brisk as ours is now. Afterwards a lighter match-lock musket came in use: and they carried their ammunition in bandeliers, to which were hung several little cases of wood covered with leather, each containing a charge of powder. The muskets with rests were used as late as the beginning of the civil wars in the time of Charles the 1st. The lighter kind succeeded them, and continued till the beginning of the present century, when they also were disused, and the troops throughout Europe armed with firelocks. These are usually made of hammered iron. For the dimensions, construction, and practice of every species of Gun, &c, see the several articles CANNON, MORTAR, &c. See also GUNNERY.

GUNNERY, the art of charging, directing, and exploding fire-arms, as cannon, mortars, muskets, &c, to the best advantage.

Gunnery is sometimes considered as a part of the military art, and sometimes as a part of pyrotechny. To the art of Gunnery too belongs the knowledge of the force and effect of gunpowder, the dimensions of the pieces, and the proportions of the powder and ball they carry, with the methods of managing, charging, pointing, spunging, &c. Also some parts of Gunnery are brought under mathematical consideration, which among mathematicians are called absolutely by the name Gunnery, viz, the rules and method of computing the range, elevation, quantity of powder, &c, so as to hit a mark or object proposed, and is more particularly called PROJECTILES; which see.

#### HISTORY of GUNNERY.

Long before the invention of gunpowder, and of Gunnery, properly so called, the art of artillery, or projectiles, was actually in practice. For, not to mention the use of spears, javelins, or stones thrown with the hand, or of bows and arrows, all which are found among the most barbarous and ignorant people, ac-

counts of the larger machines for throwing stones, darts, &c, are recorded by the most ancient writers. Thus, one of the kings of Judah, 800 years before the christian æra, erected engines of war on the towers and bulwarks of Jerusalem, for shooting arrows and great stones for the defence of that city. 2 Chron. xxvi. 15. Such machines were afterwards known among the Greeks and Romans by the names of Ballista, Catapulta, &c, which produced effects by the action of a spring of a strongly twisted cordage, formed of tough and elastic animal substances, no less terrible than the artillery of the moderns. Such warlike instruments continued in use down to the 12th and 13th centuries, and the use of bows still longer; nor is it probable that they were totally laid aside till they were superseded by gunpowder and the modern ordnance.

The first application of gunpowder to military affairs, it seems, was made soon after the year 1300, for which the proposal of friar Bacon, about the year 1280, for applying its enormous explosion to the destruction of armies, might give the first hint; and Schwartz, to whom the invention of gunpowder has been erroneously ascribed, on account of the accident abovementioned under the article Gun, might have been the first who actually applied it in this way, that is in Europe; for as to Asia, it is probable that the Chinese and Indians had something of the kind many ages before. Thus, only to mention the prohibition of fire-arms in the code of Gentoo laws, printed by the East India Company in 1776, which seems to confirm the suspicion suggested by a passage in Quintus Curtius, that Alexander the Great found some weapons of that kind in India: Cannon in the Shanferit idiom is called shet-aghnee, or the weapon that kills a hundred men at once.

However, the first pieces of artillery, which were charged with gunpowder and stone bullets of a prodigious size, were of very clumsy and inconvenient structure and weight. Thus, when Mahomet the 2d besieged Constantinople in 1453, he battered the walls with stones of this kind, and with pieces of the calibre of 1200 pounds; which could not be fired more than four times a day. It was however soon discovered that iron bullets, of much less weight than stone ones, would be more efficacious if impelled by greater quantities of stronger powder. This occasioned an alteration in the matter and form of the cannon, which were now cast of brass. These were lighter and more manageable than the former, at the same time that they were stronger in proportion to their bore. This change took place about the close of the 15th century.

By this means came first into use such powder as is now employed over all Europe, by varying the proportion of the materials. But this change of the proportion was not the only improvement it received. The practice of graining it is doubtless of considerable advantage. At first the powder had been always used in the form of fine meal, such as it was reduced to by grinding the materials together. And it is doubtful whether the first graining of powder was intended to increase its strength, or only to render it more convenient for filling into small charges and the charging of small arms, to which alone it was applied for many years, whilst meal-powder was still used for cannon.



But at last the additional strength which the grained powder was found to possess, doubtless from the free passage of the air between the grains, occasioned the meal-powder to be entirely laid aside.

For the last 200 years, the formation of cannon has been very little improved; the best pieces of modern artillery differing little in their proportions from those used in the time of Charles the 5th. Indeed lighter and shorter pieces have been often proposed and tried; but though they have their advantages in particular cases, it is agreed they are not sufficient for general service. Yet the size of the pieces has been much decreased; the same purposes being now accomplished, by smaller pieces than what were formerly thought necessary. Thus the battering cannon now approved, are those that formerly were called demi cannon, carrying a ball of 24 pounds weight; this weight having been found fully sufficient. The method also of making a breach, by first cutting off the whole wall as low as possible before its upper part is attempted to be beaten down, seems to be a considerable modern improvement in the practical part of gunnery. But the most considerable improvement in the practice, is the method of firing with small quantities of powder, and elevating the piece but a little, so that the bullet may just go clear of the parapet of the enemy, and drop into their works, called ricochet firing: for by this means the ball, coming to the ground at a small angle, and with a small velocity, does not bury itself, but bounds or rolls along a great way, destroying all before it. This method was first practised by M. Vauban at the siege of Aeth, in the year 1692. A practice of this kind was successfully used by the king of Prussia at the battle of Rosbach in 1757. He had several six-inch mortars, made with trunnions, and mounted on travelling carriages, which were fired obliquely on the enemy's lines, and among their horse. These being charged with only 8 ounces of powder, and elevated at one degree and a quarter, did great execution: for the shells rolling along the lines with burning fuses made the stoutest of the enemy not wait for their bursting.

The use of fire-arms was however long known before any theory of projectiles was formed. The Italians were the first people that made any attempts at the theory, which they did about the beginning of the 16th century, and amongst them it seems the first who wrote professedly on the flight of cannon shot, was Nicholas Tartalia, of Brescia, the same author who had so great a share in the invention of the rules for cubic equations. In 1537 he published, at Venice, his *Nova Scientia*, and in 1546 his *Questi & Inventioni diversi*, in both which he treats professedly on these motions, as well as in another work, translated into English with additions by Cyprian Lucar, under the title of *Colloquies concerning the Art of Shooting in great and small Pieces of Artillery*, and published at London in 1588. He determined, that the greatest range of a shot was when discharged at an elevation of  $45^{\circ}$ : and he asserted, contrary to the opinion of his contemporaries, that no part of the path described by a ball is a right line; although the curvature in the first part of it is so small, that it need not be attended to. He compared it to the surface of the sea; which, though it appears to be a plane, is yet doubtless incurvated round the centre of

the earth. He says he invented the gunner's quadrant, for laying a piece of ordnance at any point or degree of elevation; and though he had but little opportunity of acquiring any practical knowledge by experiments, he yet gave shrewd guesses at the event of some untried methods.

The philosophers of those times also took part in the questions arising upon this subject; and many disputes on motion were held, especially in Italy, which continued till the time of Galileo, and probably gave rise to his celebrated Dialogues on Motion. These were not published till the year 1638; and in the interval there were published many theories of the motion of military projectiles, as well as many tables of their comparative ranges; though for the most part very fallacious, and inconsistent with the motions of these bodies.

It is remarkable however that, during these contests, so few of those who were intrusted with the care of artillery, thought it worth while to bring their theories to the test of experiment. Mr. Robins informs us, in the preface to his *New Principles of Gunnery*, that he had met with no more than four authors who had treated experimentally on this subject. The first of these is Collado, in 1642, who has given the ranges of a falconet, carrying a three-pound shot, to every point of the gunner's quadrant, each point being the 12th part, or  $7^{\circ}$  and a half. But from his numbers it is manifest that the piece was not charged with its usual allotment of powder. The result of his trials shews the ranges at the point-blanc, and the several points of elevation, as below.

*Collado's Experiments.*

Elevation at		Range in	
Points.	Deg.	paces.	
0	or 0	-	268
1	- $7\frac{1}{2}$	-	594
2	- 15	-	794
3	- $22\frac{1}{2}$	-	954
4	- 30	-	1010
5	- $37\frac{1}{2}$	-	1040
6	- 45	-	1053
7	- $52\frac{1}{2}$	between the 3d and 4th	
8	- 60	between the 2d and 3d	
9	- $67\frac{1}{2}$	between the 1st and 2d	
10	- 75	between the 0 and 1st	
11	- $82\frac{1}{2}$	fell very near the piece.	

The next was by Wm. Bourne, in 1643, in his *Art of Shooting in Great Ordnance*. His elevations were not regulated by the points of the Gunner's quadrant, but by degrees; and he gives the proportions between the ranges at different elevations and the extent of the point-blanc shot, thus: if the extent of the point-blanc shot be represented by 1, then the proportions of the ranges at several elevations will be as below, viz.

*Bourne's Proportion of Ranges.*

Elevation.	Range.
$0^{\circ}$	1
5	$2\frac{2}{3}$
10	$3\frac{1}{3}$
15	$4\frac{1}{3}$
20	$4\frac{2}{3}$
and the greatest random $5\frac{1}{2}$ ;	

which



which greatest random, he says, in a calm day is at  $42^\circ$  elevation; but according to the strength of the wind, and as it favours or opposes the flight of the shot, the elevation may be from  $45^\circ$  to  $36^\circ$ .—He does not say with what piece he made his trials; though from his proportions it seems to have been a small one. This however ought to have been mentioned, as the relation between the extent of different ranges varies extremely according to the velocity and density of the bullet.

After him, Eldred and Anderson, both Englishmen, also published treatises on this subject. The former of these was many years gunner of Dover Castle, where most of his experiments were made, the earliest of which are dated in 1611, though his book was not published till 1646, and was intitled *The Gunner's Glasse*. His principles were sufficiently simple, and within certain limits very near the truth, though they were not rigorously so. He has given the actual ranges of different pieces of artillery at small elevations, all under 10 degrees. His experiments are numerous, and appear to be made with great care and caution; and he has honestly set down some, which were not reconcilable to his method; upon the whole he seems to have taken more pains, and to have had a juster knowledge of his business, than is to be found in most of his practical brethren.

Galileo printed his Dialogues on Motion in the year 1646. In these he pointed out the general laws observed by nature in the production and composition of motion, and was the first who described the action and effects of gravity on falling bodies: on these principles he determined, that the flight of a cannon-shot, or of any other projectile, would be in the curve of a parabola, unless so far as it should be diverted from that track by the resistance of the air. He also proposed the means of examining the inequalities which arise from thence, and of discovering what sensible effects that resistance would produce in the motion of a bullet at some given distance from the piece.

Notwithstanding these determinations and hints of Galileo, it seems that those who came after him never imagined that it was necessary to consider how far the operations of Gunnery were affected by this resistance. Instead of this, they boldly asserted, without making the experiment, that no great variation could arise from the resistance of the air in the flight of shells or cannon shot. In this persuasion they supported themselves chiefly by considering the extreme rarity of the air, compared with those dense and ponderous bodies; and at last it became an almost generally established maxim, that the flight of these bodies was nearly in the curve of a parabola.

Thus, Robert Anderson, in his *Genuine Use and Effects of the Gunne*, published in 1674, and again in his book, *To hit a Mark*, in 1690, relates a great many experiments; but proceeding on the principles of Galileo, he strenuously asserts that the flight of all bullets is in the curve of a parabola; undertaking to answer all objections that could be brought to the contrary. The same thing was also undertaken by Blondel, in his *Art de jeter les Bombes*, published in 1683; where, after long discussion, he concludes, that the variations from the air's resistance are so slight as not to deserve any notice. The same subject is treated of in the Philos.

Transf. N<sup>o</sup> 216, p. 68, by Dr. Halley; who also, swayed by the very great disproportion between the density of the air and that of iron or lead, thought it reasonable to believe that the opposition of the air to large metal-shot is scarcely discernible; although in small and light shot he owns that it must be accounted for.

But though this hypothesis went on smoothly in speculation; yet Anderson, who made a great number of trials, found it impossible to support it without some new modification. For though it does not appear that he ever examined the comparative ranges of either cannon or musket shot when fired with their usual velocities, yet his experiments on the ranges of shells thrown with velocities that were but small, in comparison of those above mentioned, convinced him that their whole track was not parabolical. But instead of making the proper inferences from hence, and concluding that the resistance of the air was of considerable efficacy, he framed a new hypothesis; which was, that the shell or bullet at its first discharge flew to a certain distance in a right line, from the end of which line only it began to describe a parabola: and this right line, which he calls the line of the impulse of the fire, he supposes is the same for all elevations. So that, by assigning a proper length to this line of impulse, it was always in his power to reconcile any two shots made at any two different angles; though the same method could not succeed with three shots; nor indeed does he ever inform us of the event of his experiments when three ranges were tried at one time.

But after the publication of Newton's Principia, it might have been expected, that the defects of the theory would be ascribed to their true cause, which is the great resistance of the air to such swift motions; as in that work he particularly considered the subject of such motions, and related the result of experiments, made on slow motions at least; by which it appeared, that in such motions the resistance increases as the square of the velocities, and he even hints a suspicion that it will increase above that law in swifter motions, as is now known to be the case. So far however were those who treated this subject scientifically, from making a proper allowance for the resistance of the atmosphere, that they still neglected it, or rather opposed it, and their theories still differed most egregiously from the truth. Huygens alone seems to have attended to this principle: for in the year 1690 he published a treatise on gravity, in which he gave an account of some experiments tending to prove that the track of all projectiles, moving with very swift motions, was widely different from that of a parabola. The rest of the learned generally acquiesced in the justness and sufficiency of Galileo's doctrine, and accordingly very erroneous calculations concerning the ranges of cannon were given. Nor was any farther notice taken of these errors till the year 1716, at which time Mr. Reffons, a French officer of artillery, of great merit and experience, gave in a memoir to the Royal Academy, importing that, "although it was agreed that theory joined with practice did constitute the perfection of every art; yet experience had taught him that theory was of very little service in the use of mortars: That the works of M. Blondel had justly enough described the several parabolic lines, according



to the different degrees of the elevation of the piece ; but that practice had convinced him there was no theory in the effect of gunpowder ; for having endeavoured, with the greatest precision, to point a mortar according to these calculations, he had never been able to establish any solid foundation upon them.”——

One instance only occurs in which D. Bernoulli applies the doctrine of Newton to the motions of projectiles, in the Com. Acad. Petrop. tom. 2, pa. 338 &c. Besides which nothing farther was done in this business till the time of Mr. Benjamin Robins, who published a treatise in 1742, intitled *New Principles of Gunnery*, in which he treated particularly, not only of the resistance of the atmosphere, but also of the force of gunpowder, the nature and effects of different guns, and almost every thing else relating to the flight of military projectiles ; and indeed he carried the theory of gunnery nearly to its utmost perfection.

The first thing considered by Mr. Robins, and which is indeed the foundation of all other particulars relating to Gunnery, is the explosive force of gunpowder. M. De la Hire, in the Hist. of the Acad. of Sciences for the year 1702, supposed that this force may be owing to the increased elasticity of the air contained in, and between the grains, in consequence of the heat and fire produced at the time of the explosion : a cause not adequate to the 200th part of the effect. On the other hand, Mr. Robins determined, by irrefragable experiments, that this force was owing to an elastic fluid, similar to our atmosphere, existing in the powder in an extremely condensed state, which being suddenly freed from the powder by the combustion, expanded with an amazing force, and violently impelled the bullet, or whatever may oppose its expansion.

The intensity of this force of exploded gunpowder Mr. Robins ascertained in different ways, after the example of Mr. Hawksbee, related in the Philos. Trans. N° 295, and his Physico-Mechan. Exper. pa. 81. One of these is by firing the powder in the air thus : A small quantity of the powder is placed in the upper part of a glass tube, and the lower part of the tube is immersed in water, the water being made to rise so near the top, that only a small portion of air is left in that part where the powder is placed : then in this situation the communication between the upper part of the tube and the external air being closed, the powder is fired by means of a burning glass, or otherwise ; the water descends upon the explosion, and stands lower in the tube than before, by a space proportioned to the quantity of powder fired.

Another way was by firing the powder in vacuo, viz, in an exhausted receiver, by dropping the grains of powder upon a hot iron included in the receiver. By this means a permanent elastic fluid was generated from the fired gunpowder, and the quantity of it was always in proportion to the quantity of powder that was used, as was found by the proportional sinking of the mercurial gage annexed to the air pump. The result of these experiments was, that the weight of the elastic air thus generated, was equal to  $\frac{3}{5}$  of the compound mass of the gunpowder which yielded it ; and that its bulk, when cold and expanded to the rarity of common atmospheric air, was about 240 times the bulk of the powder ; and consequently in the same proportion

would such fluid at first, if it were cold, exceed the force or elasticity of the atmosphere. But as Mr. Robins found, by another ingenious experiment, that air heated to the extreme degree of the white heat of iron, has its elasticity quadrupled, or is 4 times as strong ; he thence inferred that the force of the elastic air generated as above, at the moment of the explosion, is at least 4 times 240, or 960, or in round numbers about 1000 times as strong as the elasticity or pressure of the atmosphere, on the same space.

Having thus determined the force of the gunpowder, or intensity of the agent by which the projectile is to be urged, Mr. Robins next proceeds to determine the effects it will produce, or the velocity with which it will impel a shot of a given weight from a piece of ordnance of given dimensions ; which is a problem strictly limited, and perfectly soluble by mathematical rules, and is in general this : Given the first force, and the law of its variation, to determine the velocity with which it will impel a given body in passing through a given space, which is the length of the bore of the gun.

In the solution of this problem, Mr. Robins assumes these two postulates, viz, 1, That the action of the powder on the bullet ceases as soon as the bullet is out of the piece ; and 2d, That all the powder of the charge is fired and converted into elastic fluid before the bullet is sensibly moved from its place : assumptions which for good reasons are found to be in many cases very near the truth. It is to be noted also, that the law by which the force of the elastic fluid varies, is this, viz, that its intensity is directly as its density, or reciprocally proportional to the space it occupies, being so much the stronger as the space is less : a principle well known, and common to all elastic fluids. Upon these principles then Mr. Robins resolves this problem, by means of the 39th prop. of Newton's Principia in a direct way, and the result is equivalent to this theorem, when the quantities are expressed by algebraic symbols ; viz, the velocity of the ball

$$v = 27130 \sqrt{\frac{10a}{cd} \times \log. \frac{b}{a}}$$

$$\text{or} = 100 \sqrt{\frac{223ad^2}{w} \times \log. \frac{b}{a}};$$

where  $v$  is the velocity of the ball,  
 $a$  the length of the charge of powder,  
 $b$  the whole length of the bore,  
 $c$  the spec. grav. of the ball, or wt. of a cubic foot  
of the same matter in ounces,  
 $d$  the diam. of the bore,  
 $w$  the wt. of the ball in ounces.

For example, Suppose  $a = 2\frac{5}{8}$  inc.,  $b = 45$  inches,  
 $c = 11345$  oz, for a ball of lead, and  $d = \frac{3}{4}$  inches ;

then  $v = 27130 \sqrt{\frac{7}{2269} \times \log. \frac{120}{7}} = 1674$  feet  
per second, the velocity of the ball.

Or, if the wt. of the bullet be  $w = 1\frac{9}{16}$  oz  $= \frac{25}{8}$  oz.

Then  $v = 100 \sqrt{\frac{1115 \times 189}{29 \times 32} \times \log. \frac{120}{7}} = 1674$ .  
feet, as before.

“ Having in this proposition, says Mr. Robins, shewn how the velocity, which any bullet acquires from the



the force of powder, may be computed upon the principles of the theory laid down in the preceding propositions; we shall next shew, that the actual velocities, with which bullets of different magnitudes are impelled from different pieces, with different quantities of powder, are really the same with the velocities assigned by these computations; and consequently that this theory of the force of powder, here delivered, does unquestionably ascertain the true action and modification of this enormous power.

“ But in order to compare the velocities communicated to bullets by the explosion with the velocities resulting from the theory by computation; it is necessary that the actual velocities with which bullets move, should be capable of being discovered, which yet is impossible to be done by any methods hitherto made public. The only means hitherto practised by others for that purpose, have been either by observing the time of the flight of the shot through a given space, or by measuring the range of the shot at a given elevation; and thence computing, on the parabolic hypothesis, what velocity would produce this range. The first method labours under this insurmountable difficulty, that the velocities of these bodies are often so swift, and consequently the time observed is so short, that an imperceptible error in that time may occasion an error in the velocity thus found, of 2, 3, 4, 5, or 600 feet in a second. The other method is so fallacious, by reason of the resistance of the air (to which inequality the first is also liable), that the velocities thus assigned may not be perhaps the 10th part of the actual velocities sought.

“ To remedy then these inconveniences, I have invented a new method of finding the real velocities of bullets of all kinds; and this to such a degree of exactness (which may be augmented too at pleasure), that in a bullet moving with a velocity of 1700 feet in 1'', the error in the estimation of it need never amount to its 500th part; and this without any extraordinary nicety in the construction of the machine.”

Mr. Robins then gives an account of the machine by which he measures the velocities of the balls, which machine is simply this, viz, a pendulous block of wood suspended freely by a horizontal axis, against which block are to be fired the balls whose velocities are to be determined.

“ This instrument thus fitted, if the weight of the pendulum be known, and likewise the respective distances of its centre of gravity, and of its centre of oscillation, from its axis of suspension, it will thence be known what motion will be communicated to this pendulum by the percussion of a body of a known weight moving with a known degree of celerity, and striking it in a given point; that is, if the pendulum be supposed at rest before the percussion, it will be known what vibration it ought to make in consequence of such a determined blow; and, on the contrary, if the pendulum, being at rest, is struck by a body of a known weight, and the vibration, which the pendulum makes after the blow, is known, the velocity of the striking body may from thence be determined.

“ Hence then, if a bullet of a known weight strikes the pendulum, and the vibration, which the pendulum makes in consequence of the stroke, be ascertained;

the velocity with which the ball moved, is thence to be known.”

Mr. Robins then explains his method of computing velocities from experiments with this machine; which method is rather troublesome and perplexed, as well as the rules of Euler and Antoni, who followed him in this business, but a much simpler rule is given in my Tracts, vol. 1, p. 119, where such experiments are explained at full length, and this rule is expressed by either of the two following formulas,

$$v = 5.6727cg \times \frac{p+b}{bir} \sqrt{o} = 614.58cg \times \frac{p+b}{birn}, \text{ the}$$

velocity; where  $v$  denotes the velocity of the ball when it strikes the pendulum,  $p$  the weight of the pendulum,  $b$  the weight of the ball,  $c$  the chord of the arc described by the vibration to the radius  $r$ ,  $g$  the distance below the axis of motion to the centre of gravity,  $o$  the distance to the centre of oscillation,  $i$  the distance to the point of impact, and  $n$  the number of oscillations the pendulum will perform in one minute, when made to oscillate in small arcs. The latter of these two theorems is much the easiest, both because it is free of radicals, and because the value of the radical  $\sqrt{o}$ , in the former, is to be first computed from the number  $n$ , or number of oscillations the pendulum is observed to make.

With such machines Mr. Robins made a great number of experiments, with musket barrels of different lengths, with balls of various weights, and with different charges or quantities of powder. He has set down the results of 61 of these experiments, which nearly agree with the corresponding velocities as computed by his theory of the force of powder, and which therefore establish that theory on a sure foundation.

From these experiments, as well as from the preceding theory, many important conclusions were deduced by Mr. Robins; and indeed by means of these it is obvious that every thing may be determined relative both to the true theory of projectiles, and to the practice of artillery. For, by firing a piece of ordnance, charged in a similar manner, against such a ballistic pendulum from different distances, the velocity lost by passing through such spaces of air will be found, and consequently the resistance of the air, the only circumstance that was wanting to complete the theory of Gunnery, or military projectiles; and of this kind I have since made a great number of experiments with cannon balls, and have thereby obtained the whole series of resistances to such a ball when moving with every degree of velocity, from 0 up to 2000 feet per second of time. In the structure of artillery, they may likewise be of the greatest use: For hence may be determined the best lengths of guns; the proportions of the shot and powder to the several lengths; the thickness of a piece, so as it may be able to confine, without bursting, any given charge of powder; as also the effect of wads, chambers, placing of the vent, ramming the powder, &c. For the many other curious circumstances relating to this subject, and the various other improvements in the theory and practice of Gunnery, made by Mr. Robins, consult the first vol. of his Tracts, collected and published by Dr. Wilson, in the year 1761, where ample information may be found.

Soon after the first publication of Robins's New Principles of Gunnery, in 1742, the learned in several other



other nations, treading in his steps, repeated and farther extended the same subject, sometimes varying and enlarging the machinery; particularly Euler in Germany, D'Antoni in Italy, and Messrs. D'Arcy and Le Roy in France. But most of these, like Mr. Robins, with small fire-arms, such as muskets, and fusils.

But in the year 1775, in conjunction with several able officers of the Royal Artillery, and other ingenious gentlemen, I undertook a course of experiments with the ballistic pendulum, in which we ventured to extend the machinery to cannon shot of 1, 2, and 3 pounds weight. An account of these experiments was published in the *Philos. Trans.* for 1778, and for which the Royal Society honoured me with the prize of the gold medal. "These were the only experiments that I know of which had been made with cannon balls for this purpose, although the conclusions to be deduced from such, are of the greatest importance to those parts of natural philosophy which are dependent on the effects of fired gunpowder; nor do I know of any other practical method of ascertaining the initial velocities within any tolerable degree of the truth. The knowledge of this velocity is of the utmost consequence in Gunnery: by means of it, together with the law of the resistance of the medium, every thing is determinable relative to that business; for, besides its being an excellent method of trying the strength of different sorts of powder, it gives us the law relative to the different quantities of powder, to the different weights of shot, and to the different lengths and sizes of guns. Besides these, there does not seem to be any thing wanting to answer any inquiry that can be made concerning the flight and ranges of shot, except the effects arising from the resistance of the medium. In these experiments the weights of the pendulums employed were from 300 to near 600 pounds. In that paper is described the method of constructing the machinery, of finding the centres of gravity and oscillation of the pendulum, and of making the experiments, which are all set down in the form of a journal, with all the minute and concomitant circumstances; as also the investigation of the new and easy rule, set down just above, for computing the velocity of the ball from the experiments. The charges of powder were varied from 2 to 8 ounces, and the shot from 1 to near 3 pounds. And from the whole were clearly deduced these principal inferences, viz,

"1. First, That gunpowder fires almost instantaneously.—2. That the velocities communicated to balls or shot, of the same weight, by different quantities of powder, are nearly in the subduplicate ratio of those quantities: a small variation, in defect, taking place when the quantities of powder became great.—3. And when shot of different weights are employed, with the same quantity of powder, the velocities communicated to them, are nearly in the reciprocal subduplicate ratio of their weights.—4. So that, universally, shot which are of different weights, and impelled by the firing of different quantities of powder, acquire velocities which are directly as the square roots of the quantities of powder, and inversely as the square roots of the weights of the shot, nearly.—5. It would therefore be a great improvement in artillery, to make use of shot of a long form, or of heavier matter; for thus the mo-

mentum of a shot, when fired with the same weight of powder, would be increased in the ratio of the square root of the weight of the shot.—6. It would also be an improvement to diminish the windage; for by so doing, one-third or more of the quantity of powder might be saved.—7. When the improvements mentioned in the last two articles are considered as both taking place, it is evident that about half the quantity of powder might be saved, which is a very considerable object. But important as this saving may be, it seems to be still exceeded by that of the article of the guns; for thus a small gun may be made to have the effect and execution of another of two or three times its size in the present mode, by discharging a shot of two or three times the weight of its natural ball or round shot. And thus a small ship might discharge shot as heavy as those of the greatest now made use of.

"Finally, as the above experiments exhibit the regulations with regard to the weights of powder and balls, when fired from the same piece of ordnance, &c; so by making similar experiments with a gun, varied in its length, by cutting off from it a certain part before each course of experiments, the effects and general rules for the different lengths of guns may be certainly determined by them. In short, the principles on which these experiments were made, are so fruitful in consequences, that, in conjunction with the effects resulting from the resistance of the medium, they seem to be sufficient for answering all the enquiries of the speculative philosopher, as well as those of the practical artillery.

In the year 1786 was published the first volume of my *Traacts*, in which is detailed, at great length, another very extensive course of experiments which were carried on at Woolwich in the years 1783, 1784, and 1785, by order of the Duke of Richmond, Master General of the Ordnance. The objects of this course were very numerous, but the principal of them were the following:

"1. The velocities with which balls are projected by equal charges of powder, from pieces of the same weight and calibre, but of different lengths.

"2. The velocities with different charges of powder, the weight and length of the gun being the same.

"3. The greatest velocity due to the different lengths of guns, to be obtained by increasing the charge as far as the resistance of the piece is capable of sustaining.

"4. The effect of varying the weight of the piece; every thing else being the same.

"5. The penetration of balls into blocks of wood.

"6. The ranges and times of flight of balls; to compare them with their initial velocities for determining the resistance of the medium.

"7. The effect of wads;

of different degrees of ramming;

of different degrees of windage;

of different positions of the vent;

of chambers, and trunnions, and every other circumstance necessary to be known for the improvement of artillery."

All these objects were obtained in a very perfect and accurate manner; excepting only the article of ranges, which were not quite so regular and uniform as might



be wished. The balls too were most of them of one pound weight; but the powder was increased from 1 ounce, up till the bore was quite full; and the pendulum was from 600 to 800lb. weight. The conclusions from the whole were as follow:

“ 1. That the former law, between the charge and velocity of ball, is again confirmed, viz, that the velocity is directly as the square root of the weight of powder, as far as to about the charge of 8 ounces: and so it would continue for all charges, were the guns of an indefinite length. But as the length of the charge is increased, and bears a more considerable proportion to the length of the bore, the velocity falls the more short of that proportion.

“ 2. That the velocity of the ball increases with the charge to a certain point, which is peculiar to each gun, where it is greatest; and that by farther increasing the charge, the velocity gradually diminishes, till the bore is quite full of powder. That this charge for the greatest velocity is greater as the gun is longer, but not greater however in so high a proportion as the length of the gun is; so that the part of the bore filled with powder bears a less proportion to the whole in the long guns, than it does in the short ones; the part of the whole which is filled being indeed nearly in the reciprocal subduplicate ratio of the length of the empty part. And the other circumstances are as in this table.

TABLE of Charges producing the Greatest Velocity.				
Gun Num.	Length of the bore.	Length filled.	Part of the whole.	Wt. of the powder.
	inches.	inches.		oz.
1	28.2	8.2	$\frac{3}{10}$	12
2	38.1	9.5	$\frac{3}{10}$	14
3	57.4	10.7	$\frac{3}{10}$	16
4	79.9	12.1	$\frac{3}{10}$	18

“ 3. It appears that the velocity continually increases as the gun is longer, though the increase in velocity is but very small in respect of the increase in length, the velocities being in a ratio somewhat less than that of the square roots of the length of the bore, but somewhat greater than that of the cube roots of the length, and is indeed nearly in the middle ratio between the two.

“ 4. The range increases in a much less ratio than the velocity, and indeed is nearly as the square root of the velocity, the gun and elevation being the same. And when this is compared with the property of the velocity and length of gun in the foregoing paragraph, we perceive that very little is gained in the range by a great increase in the length of the gun, the charge being the same. And indeed the range is nearly as the 5th root of the length of the bore; which is so small an increase, as to amount only to about  $\frac{1}{5}$ th part more range for a double length of gun.

“ 5. It also appears that the time of the ball's flight

is nearly as the range; the gun and elevation being the same.

“ 6. It appears that there is no sensible difference caused in the velocity or range, by varying the weight of the gun, nor by the use of wads, nor by different degrees of ramming, nor by firing the charge of powder in different parts of it.

“ 7. But a great difference in the velocity arises from a small degree of windage. Indeed with the usual established windage only, namely, about  $\frac{1}{20}$ th of the caliber, no less than between  $\frac{1}{3}$  and  $\frac{1}{4}$  of the powder escapes and is lost. And as the balls are often smaller than that size, it frequently happens that half the powder is lost by unnecessary windage.

“ 8. It appears that the resisting force of wood, to balls fired into it, is not constant. And that the depths penetrated by different velocities or charges, are nearly as the logarithms of the charges, instead of being as the charges themselves, or, which is the same thing, as the square of the velocity.

“ 9. These, and most other experiments, shew that balls are greatly deflected from the direction they are projected in; and that so much as 300 or 400 yards in a range of a mile, or almost  $\frac{1}{4}$ th of the range, which is nearly a deflection of an angle of 15 degrees.

“ 10. Finally, these experiments furnish us with the following concomitant data, to a tolerable degree of accuracy, namely, the dimensions and elevation of the gun, the weight and dimensions of the powder and shot, with the range and time of flight, and the first velocity of the ball. From which it is to be hoped that the measure of the resistance of the air to projectiles, may be determined, and thereby lay the foundation for a true and practical system of Gunnery, which may be as well useful in service as in theory.”

Since the publication of those Tracts, we have prosecuted the experiments still farther, from year to year, gradually extending our aim to more objects, and enlarging the guns and machinery, till we have arrived at experiments with the 6 pounder guns, and pendulums of 1800 pounds weight. One of the new objects of enquiry, was the resistance the atmosphere makes to military projectiles; to obtain which, the guns have been placed at many different distances from the pendulum, against which they are fired, to get the velocity lost in passing through those spaces of air; by which, and the use of the whirling machine, described near the end of the 1st vol. of Robins's Tracts, for the slower motions, I have investigated the resistance of the air to given balls moving with all degrees of velocity, from 0 up to 2000 feet per second; as well as the resistance for many degrees of velocity, to planes and figures of other shapes, and inclined to their path in all varieties of angles; from which I have deduced general laws and formulas for all such motions.

Mr. Robins made also similar experiments on the resistance of the air; but being only with musket bullets, on account of their smallness, and of their change of figure by the explosion of the powder, I find they are very inaccurate, and considerably different from those above mentioned, which were accurately made with pretty considerable cannon balls, of iron. For which reason we may omit here the rules and theory deduced from them by Mr. Robins, till others more correct shall have.



have been established. All these experiments indeed agree in evincing the very enormous resistance the air makes to the swift motions of military projectiles, amounting in some cases to 20 or 30 times the weight of the ball itself; on which account the common rules for projectiles, deduced from the parabolic theory, are of little or no use in real practice; for, from these experiments it is clearly proved, that the track described by the flight even of the heaviest shot, is neither a parabola, nor yet approaching any thing near it, except when they are projected with very small velocities; in so much that some balls, which in the air range only to the distance of one mile, would in vacuo, when projected with the same velocity, range above 10 or 20 times as far. For the common rules of the parabolic theory, see PROJECTILES. And for a small specimen of my experiments on resistances, see the 2d vol. of the Edinburgh Philos. Transf.; as also my Conic Sections and Select Exercises, at the end, also the articles FORCE, and RESISTANCE, in this Dictionary.

Mr. Benjamin Thompson instituted a very considerable course of experiments of the same kind as those of Mr. Robins, with musket barrels, which was published in the Philos. Transf. vol. 71, for the year 1781. In these experiments, the conclusions of Mr. Robins are generally confirmed, and several other curious circumstances in this business are remarked by Mr. Thompson. This gentleman also pursues a hint thrown out by Mr. Robins relative to the determining the velocity of a ball from the recoil of the pendulous gun itself. Mr. Robins, in prop. 11, remarks that the effect of the exploded powder upon the recoil of the gun, is the same, whether the gun is charged with a ball, or without one; and that the chord, or velocity, of recoil with the powder alone, being subtracted from that of the recoil when charged with both powder and ball, leaves the velocity which is due to the ball alone. From whence Mr. Thompson observes, that the inference is obvious, viz, that the momentum thus communicated to the gun by the ball alone, being equal to the momentum of the ball, this becomes known; and therefore being divided by the known weight of the ball, the quotient will be its velocity. Mr. Thompson sets a great value on this new rule, the velocities by means of which, he found to agree nearly with several of those deduced from the motion of the pendulum; and in the other cases in which they differed greatly from these, he very inconsistently supposes that these latter ones are erroneous. In the experiments however contained in my Tracts, a great multitude of those cases are compared together, and the inaccuracy of that new rule is fully proved.

Having in the 9th prop. compared together a number of computed and experimented velocities of balls, to verify his theory: in the 10th prop. Mr. Robins assigns the changes in the force of powder, which arise from the different state of the atmosphere, as to heat and moisture, both which he finds have some effect on it, but especially the latter. In prop. 11 he investigates the velocity which the flame of gunpowder acquires by expanding itself, supposing it fired in a given piece of artillery, without either a bullet or any other body before it. This velocity he finds is upwards of 7000 feet per second. But the celebrated Euler, in his commentary on this part of Mr. Robins's book, thinks it may be

still much greater. And in this prop. too it is that Mr. Robins declares his opinion, above alluded to, viz, that the effect of the powder upon the recoil of the gun is the same, in all cases, whether fired with a ball, or without one.—In prop. 12 he ascertains the manner in which the flame of powder impels a ball which is laid at a considerable distance from the charge; shewing here that the sudden accumulation and density of the fluid against the ball, is the reason that the barrel is so often burst in those cases.—In prop. 13 he enumerates the various kinds of powder, and describes the properest methods of examining its goodness. He here shews that the best proportion of the ingredients, is when the saltpetre is  $\frac{3}{4}$  of the whole compound mass of the powder, and the sulphur and charcoal the other  $\frac{1}{4}$  between them, in equal quantities. In this prop. Mr. Robins takes occasion to remark upon the use of eprouvettes, or methods of trying powder; condemning the practice of the English in using what is called the vertical eprouvette; as well as that of the French, in using a small mortar, with a very large ball, and a small charge of powder: and instead of these, he strongly recommends the use of his ballistic pendulum, for its great accuracy: But for still more dispatch, he says he should use another method, which however he reserves to himself, without giving any particular description of it. From what has been done by Mr. Robins upon this head, several persons have introduced his method of suspending the gun as a pendulum, and noting the quantity of its oscillating recoil when fired with a certain quantity of powder; and of this kind I have contrived a machine, which possesses several advantages over all others; being extremely simple, accurate, and expeditious; so much so indeed, that the weighing out of the powder is the chief part of the trouble. See GUNPOWDER, and EPROUVETTE.

The other or 2d chapter of Mr. Robins's work, in 8 propositions, treats "of the resistance of the air, and of the track described by the flight of shot and shells." And of these, prop. 1 describes the general principles of the resistance of fluids to solid bodies moving in them. Here Mr. Robins discriminates between continued and compressed fluids, which immediately rush into the space quitted by a body moving in them, and whose parts yield to the impulse of the body without condensing and accumulating before it; and such fluids as are imperfectly compressed, rushing into a void space with a limited velocity, as in the case of our atmosphere, which condenses more and more before the ball as this moves quicker, and also presses the less behind it, by following it always with only a given velocity: hence it happens that the former fluid will resist moving bodies in proportion to the square of the velocity, while the latter resists in a higher proportion.—Proposition 2 is "to determine the resistance of the air to projectiles by experiments." One of the methods for this purpose, is by the ballistic pendulum, placing the gun at different distances from it, by which he finds the velocity lost in passing through certain spaces of air, and consequently the force of resistance to such velocities as the body moves with in the several parts of its path. And another way was by firing balls, with a known given velocity, over a large piece of water, in which the fall and plunge of the ball could



could be seen, and consequently the space it passed over in a given time. By these means Mr. Robins determined the resistances of the air to several different velocities, all which shewed that there was a gradual increase of the resistance, over the law of the square of the velocity, as the body moved quicker.—In the remaining propositions of this chapter, he proceeds a little farther in this subject of the resistance of the air; in which he lays down a rule for the proportion of the resistance between two assigned velocities; and he shews that when a 24 pound ball, fired with its full charge of powder, first issues from the piece, the resistance it meets with from the air is more than 20 times its weight. He farther shews that “the track described by the flight of shot or shells is neither a parabola, nor nearly a parabola, unless they are projected with small velocities;” and that “bullets in their flight are not only depressed beneath their original direction by the action of gravity, but are also frequently driven to the right or left of that direction by the action of some other force: and in the 8th or last proposition, he pretends to shew that the depths of penetration of balls into firm substances, are as the squares of the velocities. But this is a mistake; for neither does it appear that his trials were sufficiently numerous or various, nor were his small leaden balls fit for this purpose; and I have found, from a number of trials with iron cannon balls, that the penetrations are in a much lower proportion, and that the resisting force of wood is not uniform. See my TRACTS.

In the following small tracts, added to the principles, in this volume, Mr. Robins prosecutes the subject of the resistance of the air much farther, and lays down rules for computing ranges made in the air. But these must be far from accurate, as they are founded on the two following principles, which I know, from numerous experiments, are erroneous: viz, 1st, “That till the velocity of the projectile surpasses that of 1100 feet in a second, the resistance may be esteemed to be in the duplicate proportion of the velocity. 2d, That if the velocity be greater than that of 11 or 1200 feet in a second, then the absolute quantity of that resistance in these greater velocities will be near 3 times as great, as it should be by a comparison with the smaller velocities.” For, instead of leaping at once from the law of the square of the velocities, and ever after being about 3 times as much, my experiments prove that the increase of the resistance above the law of the square of the velocity, takes place at first in the smallest motions, and increases gradually more and more, to a certain point, but never rises so high as to be 3 times that quantity, after which it decreases again. To render this evident, I have inserted the following table of the actual quantities of resistances, which are deduced from accurate experiments, and which shew also the nature of the law of the variations, by means of the columns of differences annexed; reserving the detail of the experiments themselves to another occasion. These resistances are, upon a ball of 1.965 inc. diameter, in avoirdupois ounces, and are for all velocities, from 0, up to that of 2000 feet per second of time.

VOL. I.

*The quantity of the resistance of the air to a ball of, 1.965 inc. diameter.*

Veloc. in feet	Resist. in ounces	1st Differences	2d Differences
0	0.000		
5	0.006		
10	0.025		
15	0.054		
20	0.100		
25	0.155		
30	0.23		
40	0.42		
50	0.67		
100	2 $\frac{3}{4}$	8 $\frac{1}{4}$	
200	11	14	5 $\frac{3}{4}$
300	25	20	6
400	45	27	7
500	72	35	8
600	107	44	9
700	151	54	10
800	205	66	12
900	271	79	13
1000	350	92	13
1100	442	104	12
1200	546	115	11
1300	661	124	9
1400	785	131	7
1500	916	135	4
1600	1051	135	0
1700	1186	133	2
1800	1319	128	5
1900	1447	122	6
2000	1569		

The additional tracts of Mr. Robins, in the latter part of this volume, which contain many useful and important matters, are numbered and titled as follows, viz, Number 1, “Of the resistance of the air. Number 2, Of the resistance of the air; together with the method of computing the motions of bodies projected in that medium. Number 3, An account of the experiments, relating to the resistance of the air; exhibited at different times before the Royal Society, in the year 1746. Number 4, Of the force of fired gunpowder, together with the computation of the velocities thereby communicated to military projectiles. Number 5, A comparison of the experimental ranges of cannon and mortars with the theory contained in the preceding papers.—Practical Maxims relating to the effects and management of artillery, and the flight of shells and shot.—A proposal for increasing the strength of the British navy, by changing all the guns, from the 18 pounders downwards, into others of equal weight, but of a greater bore.” With several letters, and other papers, “On pointing, or the directing of cannon to strike distant objects; Of the nature and advantage of rifled barrel pieces,” &c.

4 D

I have



I have dwelt thus long on Mr. Robins's New Principles of Gunnery, because it is the first work that can be considered as attempting to establish a practical system of gunnery, and projectiles, on good experiments, on the force of gunpowder, on the resistance of the air, and on the effects of different pieces of artillery. Those experiments are however not sufficiently perfect, both on account of the smallness of the bullets, and for want of good ranges, to form a proper theory upon. I have supplied some of the necessary desiderata for this purpose, viz, the resistance of the air to cannon balls moving with all degrees of velocity, and the velocities communicated by given charges of powder to different balls, and from different pieces of artillery. But there are still wanting good experiments with different pieces of ordnance, giving the ranges and times of flight, with all varieties of charges, and at all different angles of elevation. A few however of those I have obtained, as in the following small table, which are derived from experiments made with a medium one-pounder gun, the iron ball being nearly 2 inches in diameter.

Powder	Elev. of gun	Veloc. of ball	Range	Time of flight
oz	o	feet	feet	//
2	15	860	4100	9
4	15	1230	5100	12
8	15	1640	6000	14½
12	15	1680	6700	15½
2	45	860	5100	21

The celebrated Mr. Euler added many excellent dissertations on the subject of Gunnery, in his translation of Robins's Gunnery into the German language; which were again farther improved in Brown's translation of the same into English in the year 1777. See also Antoni's *Examen de la Poudre*; the experiments of MM. D'Arcy and Le Roy, in the *Memoirs of the Academy* in 1751; and D'Arcy's *Essai d'une theorie d'artillerie* in 1760: my *Traacts*; and paper on the force of fired gunpowder in the *Philos. Transf.* for 1778: and Thompson's paper on the same subject in 1781. Of the common or parabolic theory of Gunnery, Mr. Simpson gave a very neat and concise treatise in his *Select Exercises*. And other authors on this part, are Starrat, Gray, Williams, Glenie, &c.

**GUNPOWDER**, a composition of nitre, sulphur, and charcoal, mixed together, and usually granulated. This easily takes fire; and when fired, it rarefies or expands with great vehemence, by means of its elastic force.—It is to this powder that we owe all the effect and action of guns, and ordnance of all sorts. So that fortification, with the modern military art, &c, in a great measure depends upon it. The above definition however is not general, for instead of the nitre, it has lately been discovered that the marine acid answers much better.

The invention of Gunpowder is ascribed, by Polydore Virgil, to a chemist; who having accidentally put

some of this composition in a mortar, and covered it with a stone; it happened to take fire, and blew up the stone. Thevet says that the person here spoken of was a monk of Fribourg, named Constantine Anelzen; but Belleforet and other authors, with more probability, hold it to be Bartholdus Schwartz, or the black, who discovered it, as some say, about the year 1320; and the first use of it is ascribed to the Venetians, in the year 1380, during the war with the Genoese. But there are earlier accounts of its use, after the accident of Schwartz, as well as before it. For Peter Mexia, in his *Various Readings*, mentions that the Moors being besieged in 1343, by Alphonfus the 11th, king of Castile, discharged a kind of iron mortars upon them, which made a noise like thunder; and this is seconded by what is related by Don Pedro, bishop of Leon, in his chronicle of king Alphonfus, who reduced Toledo, viz, that in a sea-combat between the king of Tunis and the Moorish king of Seville, about that time, those of Tunis had certain iron tubs or barrels, with which they threw thunderbolts of fire.

Du-Cange adds, that there is mention made of gunpowder in the registers of the chambers of accounts in France as early as the year 1338.

But it appears that Roger Bacon knew of Gunpowder near 100 years before Schwartz was born. He tells us, in his *Treatise De Secretis Operibus Artis & Naturæ, & de Nullitate Magiæ*, cap. 6, (which is supposed by some to have been published at Oxford in 1216, and which was undoubtedly written before his *Opus Majus*, in 1267), "that from saltpetre, and other ingredients, we are able to make a fire that shall burn at what distance we please." And Dr. Plott, in his *History of Oxfordshire*, pa. 236, assures us that these "other ingredients were explained in a MS. copy of the same treatise, in the hands of Dr. G. Langbain, and seen by Dr. Wallis, to be sulphur and wood coal." Farther, in the life of Friar Bacon in *Biographia Britannica*, vol. 1, we are told that Bacon himself has divulged the secret of this composition in a cypher, by transposing the letters of the two words in chap. xi. of the said treatise; where it is thus expressed: *sed tamen salis petra LURA MOPE CAN UBRE* (i. e. *carbonum pulvere*) *et sulphuris; et sic facies tonitrum & corruscationem, si scias artificium*: and from hence the biographer apprehends the words *carbonum pulvere* were transferred to the 6th chapter of Langbain's MS. In this same chapter Bacon expressly says that sounds like thunder, and corruscations, may be formed in the air, much more horrible than those that happen naturally. And farther adds, that there are many ways of doing this, by which a city or an army might be destroyed: and he supposes that by an artifice of this kind Gideon defeated the Midianites with only 300 men: Judges, chap. 7. There is also another passage to the same purpose, in the treatise *De Scientia Experimentalis*. See Dr. Jebb's edition of the *Opus Majus*, p. 474.

Mr. Robins, in the preface to his *Gunnery*, apprehends that Bacon describes Gunpowder not as a new composition first proposed by himself, but as the application of an old one to military purposes, and that it was known long before his time.

But Mr. Dutens carries the antiquity of Gunpowder



der still much higher, and refers to the writings of the ancients themselves for the proof of it. "Virgil, says he, and his Commentator Servius (*Æneid*, lib. 6, v. 585), Hyginus (*Fabul.* 61 and 650), Eustathius (*ad Odyss.* λ 234, pa. 1682, lib. 1), La Cerda (in *Virgil.* loc. cit.), Valerius Flaccus (lib. i. 662), and many other authors (as Raphael Volatarran. in *Comentar.* Cornelius Agrippa poster. *Oper. de Verbo Dei*, c. 100, p. 237.—Gruteri *Fax Artium Liberal* tom. 2, p. 1236), speak in such a manner of Salmoneus's attempts to imitate thunder, as suggest to us that this prince used for that purpose a composition of the nature of gunpowder. Eustathius in particular speaks of him on this occasion, as being so very expert in mechanics, that he formed machines, which imitated the noise of thunder; and the writers of fable, whose surprise in this respect may be compared to that of the Mexicans when they first beheld the fire-arms of the Spaniards, give out that Jupiter, incensed at the audacity of this prince, slew him with lightning, as he was employing himself in launching his thunder. But it is much more natural to suppose that this unfortunate prince, the inventor of Gunpowder, gave rise to these fables, by having accidentally fallen a victim to his own experiments. Dion (*Hist. Rom. in Caligula*, p. 662) and Joannes Antiochenus (in *Chronico*, &c. a Valesio edita, Paris 1634, p. 804), report the very same thing of Caligula, assuring us that this emperor imitated thunder and lightning by means of certain machines, which at the same time emitted stones. Themistius informs us that the Brachmans encountered one another with thunder and lightning, which they had the art of launching from on high at a considerable distance; (*Themist. Oratio* 27, p. 337). And in another place he relates, that Hercules and Bacchus, attempting to assail them in a fort where they were entrenched, were so roughly received by reiterated strokes of thunder and lightning, launched upon them from on high by the besieged, that they were obliged to retire, leaving behind them an everlasting monument of the rashness of their enterprise. Agathias the historian reports of Anthemius Tralensis, that having fallen out with his neighbour Zeno the rhetorician, he set fire to his house with thunder and lightning. It appears from all these passages, that the effects ascribed to these engines of war, especially those of Caligula, Anthemius, and the Indians, could be only brought about by Gunpowder. And what is still more, we find in Julius Africanus a receipt for an ingenious composition to be thrown upon an enemy, which very nearly resembles that powder. But what places this beyond all doubt, is a clear and positive passage of an author called Marcus Græcus, whose work in manuscript is in the royal library at Paris, intitled *Liber Ignium*. Dr. Mead had the same also in manuscript, and a copy of that is now in my hands. (See above). The author describes several ways of encountering an enemy, by launching fire upon him; and among others gives the following. Mix together one pound of live sulphur, two of charcoal of willow, and 6 of saltpetre; reducing them to a very fine powder in a marble mortar. He adds, that a certain quantity of this is to be put into a long, narrow, and well compacted cover, and so discharged into the air. Here we have the description

of a rocket. The cover with which thunder is imitated, he represents as short, thick, but half-filled, and strongly bound with packthread; which is exactly the form of a cracker. He then treats of different methods of preparing the match, and how one squib may set fire to another in the air, by having it inclosed within it. In short, he speaks as clearly of the composition and effects of Gunpowder, as any person in our times could do. I own I have not yet been able precisely to determine when this author lived, but probably it was before the time of the Arabian physician Mesue, who speaks of him, and who flourished in the beginning of the 9th century. Nay, there is reason to believe that he is the same of whom Galen speaks; in which case he will be of antiquity sufficient to support what I advance." It appears too, from many authors, and many circumstances, that this composition has been known to the Chinese and Indians for thousands of years. See what is said on this head under the article *GUN*.

To this history of Gunpowder it may be added, that it has lately been discovered that saltpetre or nitre is not essential to this composition, but that its place may be supplied by other substances; for new Gunpowder, of double the strength of the old, has lately been made in France, by the chemists in that country, without any nitre at all; and in the year 1790 I tried some of this new powder, that was made at Woolwich, with my eprouvette, when I found it about double the strength of the ordinary sort. This is effected by substituting, instead of the nitre, the like quantity of the marine acid.

But perhaps this new composition may not come into common and general use; both because of the great expence in procuring or making the acid, and of the trouble and danger of preventing it from taking fire by the heat of making it; for it is found to catch fire and explode from a very small degree of heat, and without the aid of a spark.

*As to the Preparation of GUNPOWDER*; there are divers compositions of it, with respect to the proportions of the three ingredients, to be met with in pyrotechnical writings; but the process of making it up is much the same in all.

For some time after the invention of artillery, Gunpowder was of a much weaker composition than that now in use, or that described by Marcus Græcus; which was chiefly owing to the weakness of their first pieces. See *GUN* and *CANNON*. Of 23 different compositions, used at different times, and mentioned by Tartaglia in his *Quest. and Inv.* lib. 3, quest. 5, the first, which was the oldest, contained equal parts of the three ingredients. But when guns of modern structure were introduced, Gunpowder of the same composition as the present came also into use. In the time of Tartaglia the cannon powder was made of 4 parts of nitre, one of sulphur, and one of charcoal; and the musket powder of 48 parts of nitre, 7 parts of sulphur, and 8 parts of charcoal; or of 18 parts of nitre, 2 parts of sulphur, and 3 parts of charcoal. But the modern composition is 6 parts of nitre, to one of each of the other two ingredients. Though Mr. Napier says, he finds the strength commonly to be greatest when the proportions are, nitre 3lb, charcoal about 9oz, and sulphur about 3oz. See his paper on Gunpowder in the *Transactions of the Royal Irish Academy*, vol. 2. The can-



non powder was in meal, and the musket powder grained. And it is certain that the graining of powder, which is a very considerable advantage, is a modern improvement. See the preface to Robins's Math. Tracts, pa. 32.

To make Gunpowder duly, regard is to be had to the purity or goodness of the ingredients, as well as the proportions of them; for the strength of the powder depends much on that circumstance, and also on the due working or mixing of them together.

To purify the nitre, by taking away the fixt or common salt, and earthy part. Dissolve it in a quantity of hot water over the fire; then filtrate it through a flannel bag, into an open vessel, and set it aside to cool, and to crystallize. These crystals may in like manner be dissolved and crystallized again; and so on, till they become quite pure and white. Then put the crystals into a dry kettle over a moderate fire, which gradually increase till it begins to smoke, evaporate, lose its humidity, and grow very white: it must be kept continually stirring with a ladle, lest it should return to its former figure, by which its greasiness would be taken away: after that, so much water is to be poured into the kettle as will cover the nitre; and when it is dissolved, and reduced to the consistency of a thick liquor, it must be continually stirred with a ladle till all the moisture is again evaporated, and it be reduced to a dry and white meal.

The like regard is to be had to the sulphur; choosing that which is in large lumps, clear and perfectly yellow; not very hard, nor compact, but porous; nor yet too much shining; and if, when set on fire, it freely burns all away, it is a sign of its goodness: so likewise, if it be pressed between two iron plates that are hot enough to make it run, and in the running appear yellow, and that which remains of a reddish colour, it is then fit for the purpose. But in case it be foul, it may be purified in this manner: melt the sulphur in a large iron ladle, or pot, over a very gentle coal fire, well kindled, but not flaming; then scum off all that rises on the top, and swims upon the sulphur; take it presently after from the fire, and strain it through a double linen cloth, letting it pass leisurely; so will it be pure, the gross matter remaining behind in the cloth.

For the charcoal, the third ingredient, such should be chosen as is large, clear, and free from knots, well burnt, and cleaving. The charcoal of light woods is mostly preferred, as of willow, and that of the branches or twigs of a moderate thickness, as of an inch or two in diameter. Dogwood is now much esteemed for this purpose. And a method of charring the wood in a large iron cylinder has lately been recommended, and indeed proved, as yielding better charcoal than formerly.

The charcoal not only concurs with the sulphur in supplying the inflammable matter, which causes the detonation of the nitre, but also greatly adds to the explosive power of it by the quantity of elastic vapour expelled during its combustion.

These three ingredients, in their purest state, being procured, long experience has shewn that they are then to be mixed together in the proportion before mentioned, to have the best effect, viz, three-quarters of the composition to be nitre, and the other quarter made up

of equal parts of the other two ingredients; or, which is the same thing, 6 parts nitre, 1 part sulphur, and 1 part charcoal.

But it is not the due proportion of the materials only, which is necessary to the making of good powder; another circumstance, not less essential, is the mixing them well together: if this be not effectually done, some parts of the composition will have too much nitre in them, and others too little; and in either case there will be a defect of strength in the powder. Robins, pa. 119.

After the materials have been reduced to fine dust, they are mixed together, and moistened with water, or vinegar, or urine, or spirit of wine, &c, and then beaten together with wooden pestles for 24 hours, either by hand, or by mills, and afterwards pressed into a hard, firm, and solid cake. When dry, it is grained or corned; which is done by breaking the cake of powder into small pieces, and so running it through a sieve; by which means the grains may have any size given them, according to the nature of the sieve employed, either finer or coarser; and thus also the dust is separated from the grains, and again mixed with other manufacturing powder, or worked up into cakes again.

Powder is smoothed, or glazed, as it is called, for small arms, by the following operation: a hollow cylinder or cask is mounted on an axis, turned by a wheel; this cask is half filled with powder, and turned for 6 hours; and thus by the mutual friction of the grains of powder it is smoothed, or glazed. The fine mealy part, thus separated or worn off from the rest, is again granulated.

*The Nature, Effects, &c, of Powder.*—When the powder is prepared as above, if the least spark be struck upon it from a steel and flint, the whole will immediately inflame, and burst out with extreme violence.—The effect is not hard to account for: the charcoal part of the grain upon which the spark falls, catching fire like tinder, the sulphur and nitre are readily melted, and the former also breaks into flame; the contiguous grains at the same time undergoing the same fate.

Sir Isaac Newton reasons thus upon the point: The charcoal and sulphur in Gunpowder easily take fire, and kindle the nitre; and the spirit of the nitre, being thereby rarefied into vapour, rushes out with an explosion much after the manner that the vapour of water rushes out of an eolipile; the sulphur also, being volatile, is converted into vapour, and augments the explosion: add, that the acid vapour of the sulphur, namely that which distils under a bell into oil of sulphur, entering violently into the fixt body of the nitre, lets loose the spirit of the nitre, and excites a greater fermentation, by which the heat is farther augmented, and the fixt body of the nitre is also rarefied into fume; and the explosion is thereby made more vehement and quick.

For if salt of tartar be mixed with Gunpowder, and that mixture be warmed till it takes fire, the explosion will be far more violent and quick than that of Gunpowder alone; which cannot proceed from any other cause, than the action of the vapour of the Gunpowder upon the salt of tartar, by which that salt is rarefied.

The explosion of Gunpowder therefore arises from the violent action, by which all the mixture being quickly and



and vehemently heated, is rarefied and converted into fume and vapour; which vapour, by the violence of that action becoming so hot as to shine, appears in the form of a flame.

M. De la Hire, in the History of the French Academy for 1702, ascribes all the force and effect of Gunpowder to the spring or elasticity of the air inclosed in the several grains of it, and in the intervals or spaces between the grains: the powder being kindled, sets the springs of so many little parcels of air a-playing, and dilates them all at once, whence the effect; the powder itself only serving to light a fire which may put the air in action; after which the whole is done by the air alone.

But it appears from the experiments and observations of Mr. Robins, that if this air be in its natural state at the time when the powder is fired, the greatest addition its elasticity could acquire from the flame of the explosion, would not amount to five times its usual quantity, and therefore could not suffice for the 200th part of the effort which is exerted by fired powder.

To understand the force of Gunpowder, it must be considered that, whether it be fired in a vacuum or in air, it produces by its explosion a permanent elastic fluid. See *Philos. Trans.* number 295; also Hauksbee's *Phys. Mechan. Exp.* p. 31. It also appears from experiment, that the elasticity or pressure of the fluid produced by the firing of Gunpowder, is, *cæteris paribus*, directly as its density.

To determine the elasticity and quantity of this elastic fluid, produced from the explosion of a given quantity of Gunpowder, Mr. Robins premises, that the elasticity of this fluid increases by heat, and diminishes by cold, in the same manner as that of the air; and that the density of this fluid, and consequently its weight, is the same with the weight of an equal bulk of air, having the same elasticity and the same temperature. From these principles, and from the experiments by which they are established (for a detail of which we must refer to the book itself, so often cited in these articles), he concludes that the fluid produced by the firing of Gunpowder is nearly  $\frac{3}{10}$  of the weight of the generating powder itself; and that the volume or bulk of this air or fluid, when expanded to the rarity of common atmospheric air, is about 244 times the bulk of the said generating powder.—Count Saluce, in his *Miscel. Phil. Mathem. Soc. Priv. Taurin.* p. 125, makes the proportion as 222 to 1; which he says agrees with the computation of Mess. Hauksbee, Amon-ton, and Belidor.

Hence it appears, that any quantity of powder fired in any confined space, which it adequately fills, exerts at the instant of its explosion against the sides of the vessel containing it, and the bodies it impels before it, a force at least 244 times greater than the elasticity of common air, or, which is the same thing, than the pressure of the atmosphere; and this without considering the great addition arising from the violent degree of heat with which it is endued at that time; the quantity of which augmentation is the next head of Mr. Robins's enquiry. He determines that the elasticity of the air is augmented in a proportion somewhat greater than that of 4 to 1, when heated to the extremest heat of red hot iron; and supposing that the

flame of fired Gunpowder is not of a less degree of heat, increasing the former number a little more than 4 times, makes nearly 1000; which shews that the elasticity of the flame, at the moment of explosion, is about 1000 times stronger than the elasticity of common air, or than the pressure of the atmosphere. But, from the height of the barometer, it is known that the pressure of the atmosphere upon every square inch, is on a medium  $14\frac{3}{4}$  lb; and therefore 1000 times this, or 14750 lb, is the force or pressure of the flame of Gunpowder, at the moment of explosion, upon a square inch, which is very nearly equivalent to 6 tons and a half.

This great force however diminishes as the fluid dilates itself, and in that proportion, viz, in proportion to the space it occupies, it being only half the strength when it occupies a double space, one third the strength when triple the space, and so on.

Mr. Robins farther supposes the degree of heat above mentioned to be a kind of medium heat; but that in the case of large quantities of powder the heat will be higher, and in very small quantities lower; and that therefore in the former case the force will be somewhat more, and in the latter somewhat less, than 1000 times the force of the atmosphere.

He farther found that the strength of powder is the same in all variations in the density of the atmosphere. But that the moisture of the air has a great effect upon it; for the same quantity which in a dry season would discharge a bullet with a velocity of 1700 feet in one second, will not in damp weather give it a velocity of more than 12 or 1300 feet in a second, or even less, if the powder be bad, and negligently kept. Robins's *Tracts*, vol. 1, p. 101, &c. Farther, as there is a certain quantity of water which, when mixed with powder, will prevent its firing at all, it cannot be doubted but every degree of moisture must abate the violence of the explosion; and hence the effects of damp powder are not difficult to account for.

It is to be observed, that the moisture imbibed by powder does not render it less active when dried again. Indeed, if powder be exposed to very great damps without any caution, or when common salt abounds in it, as often happens through negligence in refining the nitre, in such cases the moisture it imbibes may perhaps be sufficient to dissolve some part of the nitre: which is a permanent damage that no drying can retrieve. But when tolerable care is taken in preserving powder, and the nitre it is composed of has been well purged from common salt, it will retain its force for a long time; and it is said that powder has been known to have been preserved for 50 years without any apparent damage from its age.

The velocity of expansion of the flame of Gunpowder, when fired in a piece of artillery, without either bullet or other body before it, is prodigiously great, viz, 7000 feet per second, or upwards, as appears from the experiments of Mr. Robins. But Mr. Bernoulli and Mr. Euler suspect it is still much greater. And I suspect it may not be less, at the moment of explosion, than 4 times as much.

It is this prodigious celerity of expansion of the flame of fired Gunpowder, which is its peculiar excellence, and the circumstance in which it so eminently surpasses



surpasses all other inventions, either ancient or modern : for as to the momentum of these projectiles only, many of the warlike machines of the ancients produced this in a degree far surpassing that of our heaviest cannon shot or shells ; but the great celerity given to these bodies cannot be in the least approached by any other means but the flame of powder.

*To prove Gunpowder.* There are several ways of doing this. 1, By sight : thus, if it be too black, it is a sign that it is moist, or else that it has too much charcoal in it ; so also, if rubbed upon white paper, it blackens it more than good powder does : but if it be of a kind of azure colour, somewhat inclining to red, it is a sign of good powder. 2, By touching : for if in crushing it with the fingers ends, the grains break easily, and turn into dust, without feeling hard, it has too much coal in it ; or if, in pressing it under the fingers upon a smooth hard board, some grains feel harder than the rest, it is a sign the sulphur is not well mixed with the nitre. Also by thrusting the hand into the parcel of powder, and grasping it, as if to take out a handful, you will feel if it is dry and equal grained, by its evading the grasp, and running mostly out of the hand. 3, By burning ; and here the method most commonly followed for this purpose with us, says Mr. Robins, is to fire a small heap of it on a clean board, and to attend nicely to the flame and smoke it produces, and to the marks it leaves behind on the board : but besides this uncertain method, there are other contrivances made use of, such as powder-tryers acting by a spring, commonly sold at the shops, and others again that move a great weight, throwing it upwards, which is a very bad sort of epreuve. But these machines, says Mr. Robins, though more perfect than the common powder-tryers, are yet liable to great irregularities ; for as they are all moved by the instantaneous stroke of the flame, and not by its continued pressure, they do not determine the force of the fired powder with sufficient certainty and uniformity. Another method is to judge from the range given to a large solid ball, thrown from a very short mortar, charged with a small quantity of powder ; which is also an uncertain way, both on account of the great disproportion between the weight of the ball and powder, and the unequal resistance of the air ; not to mention that it is too tedious to prove large quantities of powder in this way ; for, “ if each barrel of powder was to be proved in this manner, the trouble of charging the mortar, and bringing back the ball each time, would be intolerable, and the delay so great, that no business of this kind could ever be finished ; and if a number of barrels are received on the merit of a few, it is great odds, but some bad ones would be amongst them, which may prove a great disappointment in time of service.” These exceptions do “ noways hold, continues Mr. Robins, against the method by which I have tried the comparative strength of different kinds of powder, which has been by the actual velocity given to a bullet, by such a quantity of powder as is usually esteemed a proper charge for the piece : and as this velocity, however great, is easily discovered by the motion which the pendulum acquires from the stroke of the bullet, it might seem a good amendment to the method used by the French (viz, that of the small mortar above

mentioned) to introduce this trial by the pendulum instead of it. But though I am satisfied, that this would be much more accurate, less laborious, and readier than the other, yet, as there is some little attention and caution required in this practice, which might render it of less dispatch than might be convenient, when a great number of barrels were to be separately tried, I should myself choose to practise another method not less certain, but prodigiously more expeditious ; so that I could engage, that the weighing out of a small parcel of powder from each barrel should be the greatest part of the labour ; and, doubtless, three or four hands could, by this means, examine 500 barrels in a morning : besides, the machines for this purpose, as they might be made of cast iron, would be so very cheap, that they might be multiplied at pleasure.” Robins, page 123. It is not certainly known what might be the particular construction of the epreuve here hinted at, but it was probably a piece of ordnance suspended like a pendulum, as he had made several experiments with a barrel in that manner. Be this however as it may, several persons, from those ideas and experiments of Mr. Robins, have made epreuves on this principle, which seems to be the best of any ; and on this idea also I have lately made a machine for this purpose, which has several peculiar contrivances, and advantages over all others, both in the nature of its motion, and the divisions on its arc, &c. It is a small cannon, the bore of which is about one inch in diameter, and is usually charged with 2 ounces of powder, and with powder only, as a ball is not necessary, and the strength of the powder is accurately shewn by the arc of the gun's recoil. The whole machine is so simple, easy, and expeditious, that, as Mr. Robins observed above, the weighing of the powder is the chief part of the trouble ; and so accurate and uniform, that the successive repetitions or firings with the same quantity of the same sort of powder, hardly ever yield a difference in the recoil of the 100th part of itself.

*To recover damaged Powder.* The method of the powder merchants is this ; they put part of the powder on a sail-cloth, to which they add an equal weight of what is really good ; then with a shovel they mingle it well together, dry it in the sun, and barrel it up, keeping it in a dry and proper place.

Others again, if it be very bad, restore it by moistening it with vinegar, water, urine, or brandy ; then they beat it fine, sift it, and to every pound of powder add an ounce, or an ounce and a half, or two ounces (according as it is decayed), of melted nitre ; and afterwards these ingredients are to be moistened and well mixed, so that nothing may be discerned in the composition ; which may be known by cutting the mass, and then they granulate it as useful.

In case the powder be quite spoiled, the only way is to extract the saltpetre with water, in the usual way, by boiling, filtrating, evaporating, and crystallizing ; and then, with fresh sulphur and charcoal, to make it up afresh.

On the subject of Gunpowder, see also Euler-on Robins's Gunnery, Antoni Examen de la Poudre, Baumé's Chemistry, and Thompson's Experiments in the Philos. Trans. for 1781.

GUNTER (EDMUND), an excellent English mathematician,



thematically, was born in Hertfordshire in 1581. He was educated at Westminster school under Dr. Busby, and from thence was elected to Christ-church College, Oxford, in 1599, where he took the degree of master of arts in 1606, and afterwards entered into holy orders; and in 1615 he took the degree of bachelor of divinity. But being particularly distinguished for his mathematical talents, when Mr. Williams resigned the professorship of astronomy in Gresham College, London, Mr. Gunter was chosen to succeed him, the 6th of March, 1619; where he greatly distinguished himself by his lectures and writings, and where he died in 1626, at only 45 years of age, to the great loss of the mathematical world.

Mr. Gunter was the author of many useful inventions and works. About the year 1606, he merited the title of an inventor, by the new projection of his Sector, which he then described in a Latin treatise, not printed however till some time afterwards.—In 1618 he had invented a small portable quadrant, for the more easy finding the hour and azimuth, and other useful purposes in astronomy.—And in 1620 or 1623, he published his *Canon Triangulorum*, or Table of Artificial Sines and Tangents, to the radius, 10,000,000 parts, to every minute of the quadrant, being the first tables of this kind published; together with the first 1000 of Briggs's logarithms of common numbers, which were in later editions extended to 10,000 numbers.—In 1622, he discovered, by experiment made at Deptford, the variation or changeable declination of the magnetic needle; his experiment shewing that the declination had changed by 5 degrees in the space of 42 years; and the same was confirmed and established by his successor Mr. Gellibrand.—He applied the logarithms of numbers, and of sines and tangents, to straight lines drawn on a scale or rule; with which proportions in common numbers and trigonometry were resolved by the mere application of a pair of compasses; a method founded on this property, that the logarithms of the terms of equal ratios are equidifferent. This was called Gunter's Proportion, and Gunter's Line; and the instrument, in the form of a two-foot scale, is now in common use for navigation and other purposes, and is commonly called the Gunter. He also greatly improved the Sector and other instruments for the same uses; the description of all which he published in 1624.—He introduced the common measuring chain, now constantly used in land-surveying, which is thence called Gunter's Chain.—Mr. Gunter drew the lines on the dials in Whitehall-garden, and wrote the description and use of them, by the direction of prince Charles, in a small tract; which he afterwards printed at the desire of king James, in 1624.—He was the first who used the word *co-sine*, for the sine of the complement of an arc. He also introduced the use of Arithmetical Complements into the logarithmical arithmetic, as is witnessed by Briggs, cap. 15, Arith. Log. And it has been said that he first started the idea of the Logarithmic Curve, which was so called, because the segments of its axis are the logarithms of the corresponding ordinates.

His works have been collected, and various editions of them have been published; the 5th is by Mr. William Leybourn, in 1673, containing the Description and Use of the Sector, Cross-staff, Bow, Quadrant, and

other instruments; with several pieces added by Samuel Foster, Henry Bond, and William Leybourn.

GUNTER'S CHAIN, the chain in common use for measuring land, according to true or statute measure; so called from Mr. Gunter its reputed inventor.

The length of the chain is 66 feet, or 22 yards, or 4 poles of  $5\frac{1}{2}$  yards each; and it is divided into 100 links, of 7.92 inches each.

This Chain is the most convenient of any thing for measuring land, because the contents thence computed are so easily turned into acres. The reason of which is, that an acre of land is just equal to 10 square chains, or 10 chains in length and 1 in breadth, or equal to 100000 square links. Hence, the dimensions being taken in chains, and multiplied together, it gives the content in square chains; which therefore being divided by 10, or a figure cut off for decimals, brings the content to acres; after which the decimals are reduced to roods and perches, by multiplying by 4 and 40. But the better way is to set the dimensions down in links as integers, considering each chain as 100 links; then, having multiplied the dimensions together, producing square links, divide these by 100000, that is, cut off five places for decimals, the rest are acres, and the decimals are reduced to roods and perches, as before.

*Ex.* Suppose in measuring a rectangular piece of ground, its length be 795 links,  
and its breadth 480 links.

$$\begin{array}{r} 63600 \\ 3180 \\ \hline \text{Ac. } 3.81600 \\ \hline 4 \\ \hline \text{Ro. } 3.264. \\ \hline 40 \\ \hline \text{Per. } 10.560 \end{array}$$

So the content is 3 ac. 3 roods 10 perches.

GUNTER'S LINE, a Logarithmic line, usually graduated upon scales, sectors, &c; and so called from its inventor Mr. Gunter.

This is otherwise called the *line of lines*, or *line of numbers*, and consists of the logarithms transferred upon a ruler, &c, from the tables, by means of a scale of equal parts, which therefore serves to resolve problems instrumentally, in the same manner as logarithms do arithmetically. For, whereas logarithms resolve proportions, or perform multiplication and division, by only addition and subtraction, the same are performed on this line by turning a pair of compasses over this way or that, or by sliding one slip of wood by the side of another, &c.

This line has been contrived various ways, for the advantage of having it as long as possible. As, first, on the two-feet ruler or scale, by Gunter. Then, in 1627 the logarithms were drawn by Wingate, on two separate rulers, sliding against each other, to save the use of compasses in resolving proportions. They were also in 1627 applied to concentric circles by Oughtred. Then in a spiral form by Mr. Milburne of Yorkhire, about the year 1650. Also, in 1657, on the present common sliding rule, by Seth Partridge.

Lastly, Mr. William Nicholson has proposed another disposition of them, on concentric circles, in the *Philos. Trans.* an. 1787, pa. 251. His instrument is equivalent



to a straight rule of  $28\frac{1}{2}$  inches long. It consists of three concentric circles, engraved and graduated on a plate of about  $1\frac{1}{2}$  inch in diameter. From the centre proceed two legs, having right-lined edges in the direction of radii; which are moveable either singly, or together. To use this instrument; place the edge of one leg at the antecedent of any proportion, and the other at the consequent, and fix them to that angle: the two legs being then moved together, and the antecedent leg placed at any other number, the other leg gives its consequent in the like position or situation on the lines.

The whole length of the line is divided into two equal intervals, or radii, of 9 larger divisions in each radius, which are numbered from 1 to 10, the 1 standing at the beginning of the line, because the logarithm of 1 is 0, and the 10 at the end of each radius; also each of these 9 spaces is subdivided into 10 other parts, unequal according to the logarithms of numbers; the smaller divisions being always 10ths of the larger; thus, if the large divisions be units or ones, the smaller are tenth-parts; if the larger be tens, the smaller are ones; and if the larger be 100's, the smaller are 10's; &c.

*Use of Gunter's Line.* 1. *To find the product of two numbers.* Extend the compasses from 1 to either of the numbers, and that extent will reach the same way from the other number to the product. Thus, to multiply 7 and 5 together; extend the compasses from 1 to 5, and that extent will reach from 7 to 35, which is the product.

2. *To divide one number by another.* Extend the compasses from the divisor to 1, and that extent will reach the same way from the dividend to the quotient. Thus, to divide 35 by 5; extend the compasses from 5 to 1, and that extent will reach from 35 to 7, which is the quotient.

3. *To find a 4th Proportional to three given Numbers;* as suppose to 6, 9, and 10. Extend from 6 to 9, and that extent will reach from 10 to 15, which is the 4th proportional sought. And the same way a 3d proportional is found to two given terms, extending from the 1st to the 2d, and then from the 2d to the 3d.

4. *To find a Mean Proportional between two given numbers,* as suppose between 7 and 28. Extend from 7 to 28, and bisect that extent; then its half will reach from 7 forward, or from 28 backward, to 14, the mean proportional between them.—Also, to extract the square root, as of 25, which is only finding a mean proportional between 1 and the given square 25, bisect the distance between 1 and 25, and the half will reach from 1 to 5, the root sought.—In like manner the cubic or 3d root, or the 4th, 5th, or any higher root, is found, by taking the extent between 1 and the given power; then take such part of it as is denoted by the index of the root, viz, the 3d part for the cube root, the 4th part for the 4th root, and so on, and that part will reach from 1 to the root sought.

If the Line on the Scale or Ruler have a slider, this is to be used instead of the compasses.

GUNTER'S QUADRANT, is a quadrant made of wood, brass, or some other substance; being a kind of stereographic projection on the plane of the equinoctial, the eye being supposed in one of the poles: so that the tropic, ecliptic, and horizon, form the arches of circles,

but the hour circles other curves, drawn by means of several altitudes of the sun, for some particular latitude every day in the year.

The use of this instrument, is to find the hour of the day, the sun's azimuth, &c, and other common problems of the sphere or globe; as also to take the altitude of an object in degrees. See QUADRANT.

GUNTER'S SCALE, usually called by seamen *the Gunter*, is a large plain scale, having various lines upon it, of great use in working the cases or questions in Navigation.

This Scale is usually 2 feet long, and about an inch and a half broad, with various lines upon it, both natural and logarithmic, relating to trigonometry, navigation, &c.

On the one side are the natural lines, and on the other the artificial or logarithmic ones. The former side is first divided into inches and tenths, and numbered from 1 to 24 inches, running the whole length near one edge. One half the length of this side consists of two plane diagonal scales, for taking off dimensions to three places of figures. On the other half or foot of this side, are contained various lines relating to trigonometry, in the natural numbers, and marked thus, viz,

*Rumb*, the rumbs or points of the compass,

*Chord*, the line of chords,

*Sine*, the line of sines,

*Tang.* the tangents,

*S. T.* the semitangents,

and at the other end of this half are

*Leag.* leagues, or equal parts,

*Rumb.* another line of rumbs,

*M. L.* miles of longitude,

*Chor.* another line of chords.

Also in the middle of this foot are *L.* and *P.* two other lines of equal parts. And all these lines on this side of the scale serve for drawing or laying down the figures to the cases in trigonometry and navigation.

On the other side of the scale are the following artificial or logarithmic lines, which serve for working or resolving those cases; viz,

*S. R.* the fine rumbs,

*T. R.* the tangent rumbs,

*Numb.* line of numbers,

*Sine*, Sines,

*V. S.* the versed sines,

*Tang.* the tangents,

*Meri.* Meridional parts.

*E. P.* Equal parts.

The late Mr. John Robertson, librarian to the Royal Society, greatly improved this scale, both as to size and accuracy, for the use of mariners. He extended it to 30 inches long, 2 inches broad, and half an inch thick; upon which the several lines are very accurately laid down by Messrs. Nairne and Blunt, ingenious instrument makers. Mr. Robertson died before his improved scales were published; but the account and description of them were supplied and drawn up by his friend Mr. William Mountaine, and published in 1778.

GUTTÆ, or *Drops*, in Architecture, are ornaments in form of little bells or cones, used in the Doric order, on the architrave, below the tryglyphs. There are usually six of them.

HADLEY'S



## H.

## H A L

**H**ADLEY's *Quadrant, Sextant, &c*, an excellent instrument so called from its inventor John Hadley, Esq. See its description and use under the article **QUADRANT**.

**HAIL**, or **HAILSTONES**, an aqueous concretion, usually in form of white or pellucid spherules, descending out of the atmosphere.

Hailstones assume various shapes, being sometimes round, at other times pyramidal, crenated, angular, thin, and flat, and sometimes stellated, with six radii like the small crystals of snow.

It is very difficult to account for the phenomena of hail in a satisfactory manner; and there are various opinions upon this head. It is usually conceived that hail is formed of drops of rain, frozen in their passage through the middle region. Others, as the Cartesians, take it for the fragments of a frozen cloud, half melted, and thus precipitated and congealed again. Signior Beccaria supposes, that it is formed in the higher regions of the air, where the cold is intense, and where the electric matter is very copious. In these circumstances, a great number of particles of water are brought near together, where they are frozen, and in their descent they collect other particles; so that the density of the substance of the Hailstone grows less and less from the centre; this being formed first in the higher regions, and the surface being collected in the lower. Accordingly, in mountains, Hailstones as well as drops of rain, are very small; and both agree in this circumstance, that the more intense is the electricity that forms them, the larger they are.

It is frequently observed that Hail attends thunder and lightning; and hence Beccaria observes, that as motion promotes freezing, so the rapid motion of the electrified clouds may promote that effect in the air.

Natural histories furnish us with a great variety of curious instances of extraordinary showers of Hail. See the *Philos. Trans.* number 203, 229; and *Hist. de France*, tom. 2, pa. 339.

**HALF-MOON**, in Fortification, is an outwork having only two faces, forming together a salient angle, which is flanked by some part of the place, and of the other bastions. See **DEMILUNE** and **RAVELIN**.

**HALF-TANGENTS**, are the tangents of the half arcs. See **SCALE** and **SEMITANGENTS**.

**HALLEY** (Dr. EDMUND), a most eminent English mathematician, philosopher and astronomer, was born in the parish of St. Leonard, Shoreditch, near

VOL. I.

## H A L

London, Oct. 29, 1656. His father, a wealthy citizen and soap-boiler, resolving to improve the promising disposition observed in his son, put him first to St. Paul's school, where he soon excelled in all parts of classical learning, and made besides a considerable advance in the mathematics; so that, as Wood observes, he had perfectly learnt the use of the celestial globe, and could make a complete dial; and we are informed by Halley himself, that he observed the change of the variation of the magnetic needle at London in 1672, one year before he left school. In 1673 he was sent to Oxford, where he chiefly applied himself to mathematics and astronomy, in which he was greatly assisted by a curious apparatus of instruments, which his father, willing to encourage his son's genius, had purchased for him. At 19 years of age he began to oblige the world with new observations and discoveries (which he continued to do to the end of a very long life), by publishing "A Direct and Geometrical method of finding the Aphelia and Excentricity of the planets." Besides various particular observations, made from time to time upon the celestial phenomena; he had, from his first admission into college, pursued a general scheme for ascertaining the true places of the fixed stars, and so to correct the errors of Tycho Brahe. His original view in this was, to carry on the design of that first restorer of astronomy, by completing the catalogue of those stars from his own observations; but upon farther enquiry, finding this province taken up by Hevelius and Flamsteed, he dropped that pursuit, and formed another; which was, to perfect the whole scheme of the heavens, by the addition of the stars which lie so near the south pole, that they could not be observed by those astronomers, as never rising above the horizon either at Dantzick or at Greenwich. With this view he left the University, before he had taken any degree, and embarked for the island of St. Helena in Nov. 1676, when he was only 20 years of age, and arrived there after a voyage of three months. He immediately set about his task with such diligence, that he completed his catalogue, and, returning home, landed in England in Nov. 1678, after an absence of two years only. The university of Oxford immediately conferred upon him the degree of A. M. and the Royal Society of London elected him one of their members.

In 1679 he was pitched upon by the Royal Society to go to Dantzick, to endeavour to adjust a dispute between Hevelius and Mr. Hooke, concerning the preference



ference as to plain and glass sights in astroscopical instruments. He arrived at Dantzick the 26th of May, when he immediately, in conjunction with Hevelius, set about their astronomical observations, which they closely continued till the 18th of July, when Halley left Dantzick, and returned to England.

In the year 1680 he undertook what is called the grand tour, accompanied by his friend the celebrated Mr. Nelson. In the way from Calais to Paris, Mr. Halley had a sight of a remarkable comet, as it then appeared a second time that year, in its return from the sun. He had the November before seen it in its descent; and he now hastened to complete his observations upon it, by viewing it from the royal observatory of France. His design in this part of his tour was, to settle a friendly correspondence between the two royal astronomers of Greenwich and Paris; and in the mean time to improve himself under so great a master as Cassini. From thence he went to Italy, where he spent great part of the year 1681; but his affairs calling him home, he then returned to England.

Soon after his return, he married the daughter of Mr. Tooke, auditor of the exchequer, and took up his residence at Islington, where he set up his tube and sextant, and eagerly pursued his favourite study: in the society of this amiable lady he lived happily for five-and-fifty years. In 1683 he published his *Theory of the Variation of the Magnetical Compass*; in which he supposes the whole globe of the earth to be one great magnet, having four magnetical poles or points of attraction, &c. The same year also he entered upon a new method of finding out the longitude, by an accurate observation of the moon's motion. His pursuits it seems were now a little interrupted by the death of his father, who having suffered greatly by the fire of London, as well as by a second marriage, into which he had imprudently entered, was found to have wasted his fortunes. Our author soon resumed his pursuits however; for in the beginning of 1684 he turned his thoughts to the subject of Kepler's sesqui-alterate proportion; when, after some meditation, he concluded from it, that the centripetal force must decrease in proportion to the square of the distance reciprocally. He found himself unable to make it out in any geometrical way; and therefore, after applying in vain for assistance to Mr. Hooke and Sir Christopher Wren, he went to Cambridge to Mr. Newton, who fully supplied him with what he so ardently sought. But Halley having now found an immense treasure in Newton, could not rest, till he had prevailed with the owner to enrich the public with it; and to this interview the world is in some measure indebted for the *Principia Mathematica Philosophiæ Naturalis*. That great work was published in 1686; and Halley, who had the whole care of the impression, prefixed to it a discourse of his own, giving a general account of the astronomical part of the book; and also a very elegant copy of verses in Latin.

In 1687 he undertook to explain the cause of a natural phenomenon, which had till then baffled the researches of the ablest geographers. It is observed that the Mediterranean sea never swells in the least, although there is no visible discharge of the prodigious quantity of water that runs into it from nine large rivers, besides several small ones, and the constant setting

in of the current at the mouth of the Straights. His solution of this difficulty gave so much satisfaction to the Society, that he was requested to prosecute these enquiries. He did so; and having shewn, by accurate experiments, how that vast accession of water was actually carried off in vapours raised by the action of the sun and wind upon its surface, he proceeded with the like success to point out the method used by nature to return the said vapours into the sea. This circulation he supposes to be carried on by the winds driving these vapours to the mountains; where being collected, they form springs, which uniting become rivulets or brooks, and many of these again meeting in the valleys, grow into large rivers, emptying themselves at last into the sea: thus demonstrating, in the most beautiful manner, the way in which the equilibrium of receipt and expence is continually preserved in the universal ocean.

He next ranged in the field of speculative geometry, where, observing some imperfections in the methods before laid down for constructing solid problems, or equations of the 3d and 4th powers, he furnished new rules, which were both more easy and more elegant than any of the former; together with a new method of finding the number of roots of such equations, and the limits of the same.

Mr. Halley next undertook to publish a more correct Ephemeris for the year 1688, there being then great want of proper ephemerides of any tolerable exactness, the common ones being justly complained of by Mr. Flamsteed.—In 1691 he published exact tables of the conjunctions of Venus and Mercury; and he afterwards shewed one extraordinary use to be made of those tables, viz, for discovering the sun's parallax, and thence the true distance of the earth from the sun.—In 1692, our author produced his tables for shewing the value of annuities on lives, calculated from bills of mortality; and his universal theorem for finding the foci of optic glasses.

But it would be endless to enumerate all his valuable discoveries now communicated to the Royal Society, and published in the *Philos. Transf.* of which, for many years, his pieces were the chief ornament and support. Their various merit is thrown into one view by the writer of his eloge in the *Paris Memoirs*; who, having mentioned his *History of the Trade-winds and Monsoons*, proceeds in these terms: "This was immediately followed by his estimation of the quantity of vapours which the sun raises from the sea; the circulation of vapours; the origin of fountains; questions on the nature of light and transparent bodies; a determination of the degrees of mortality, in order to adjust the valuation of annuities on lives; and many other works, all the sciences relating to astronomy, geometry, and algebra, optics and dioptrics, ballistics and artillery, speculative and experimental philosophy, natural history, antiquities, philology, and criticism; being about 25 or 30 dissertations, which he produced during the 9 or 10 years of his residence at London; and all abounding with ideas new, singular, and useful."

In 1691, the Savilian professorship of astronomy at Oxford being vacant, he applied for that office, but without success. Whiston, in the *Memoirs of his own Life*, tells us from Dr. Bentley, that Halley "being thought of for successor to the mathematical chair at Oxford,



Oxford, bishop Stillingfleet was desired to recommend him at court; but hearing that he was a sceptic and a banterer of religion, the bishop scrupled to be concerned, till his chaplain Bentley should talk with him about it, which he did. But Halley was so sincere in his infidelity, that he would not so much as pretend to believe the christian religion, though he thereby was likely to lose a professorship; which he did accordingly, and it was then given to Dr. Gregory."

Halley had published his Theory of the Variation of the Magnetical Compass, as has been already observed, in 1683; which, though it was well received both at home and abroad, he found, upon a review, liable to great and insuperable objections. Yet the phenomena of the variation of the needle, upon which it is raised, being so many certain and indisputed facts, he spared no pains to possess himself of all the observations relating to it he could possibly come at. To this end he procured an application to be made to king William, who appointed him commander of the Paramour pink, with orders to search out by observations the discovery of the rule of variations, and to lay down the longitudes and latitudes of the English settlements in America.—He set out on this attempt on the 24th of November, 1698: but having crossed the line, his men grew sickly; and his first lieutenant mutinying, he returned home in June 1699. Having got the lieutenant tried and cashiered, he set sail a second time in September following, with the same ship, and another of less bulk, of which he had also the command. He now traversed the vast Atlantic ocean from one hemisphere to the other, as far as the ice would permit him to go; and having made his observations at St. Helena, Brazil, Cape Verde, Barbadoes, the Madeiras, the Canaries, the coast of Barbary, and many other latitudes, he arrived in England in September 1700; and the next year published a general chart, shewing at one view the variation of the compass in all those places.

Captain Halley, as he was now called, had been at home little more than half a year, when he was sent by the king, to observe the course of the tides, with the longitude and latitude of the principal head-lands in the British channel; which having executed with his usual expedition and accuracy, he published a large map of the channel.

Soon after, the emperor of Germany resolving to make a convenient harbour for shipping in the Adriatic, captain Halley was sent by queen Anne to view the two ports on the coast of Dalmatia. He embarked on the 22d of November 1702; passed over to Holland; and going through Germany to Vienna, he proceeded to Istria: but the Dutch opposing the design, it was laid aside; yet the emperor made him a present of a rich diamond ring from his finger, and honoured him with a letter of recommendation, written with his own hand, to queen Anne. Presently after his return, he was sent again on the same business; when passing through Hanover, he supped with the electoral prince, who was afterward king George the 1st, and his sister the queen of Prussia. On his arrival at Vienna, he was the same evening presented to the emperor, who sent his chief engineer to attend him to Istria, where they repaired the fortifications of Trieste, and added new

ones.

Mr. Halley returned to England in Nov. 1703; and the same year he was made professor of geometry in the university of Oxford, instead of Dr. Wallis then just deceased, and he was at the same time honoured by the university with the degree of doctor of laws. He was scarcely settled in Oxford, when he began to translate into Latin, from the Arabic, *Apollonius de Sectione Rationis*; and to restore the two books *De Sectione Spatii* of the same author, which are lost, from the account given of them by Pappus; and he published the whole work in 1706. He afterwards had a share in preparing for the press Apollonius's Conics; and ventured to supply the whole 8th book, the original of which is also lost. To this work he added Serenus on the Section of the Cylinder and Cone, printed from the original Greek, with a Latin translation, and published the whole in folio 1710. Beside these, the *Miscellanea Curiosa*, in 3 volumes 8vo, had come out under his direction in 1708.

In 1713, he succeeded Doctor, afterwards Sir, Hans Sloane, in the office of Secretary to the Royal Society. And, upon the death of Mr. Flamsteed in 1719, he was appointed to succeed him at Greenwich as Astronomer Royal; upon which occasion, that he might be more at leisure to attend the duties of this office, he resigned that of secretary to the Royal Society in 1721. Although he was 63 or 64 years of age when he entered upon his office at Greenwich, for the space of 18 years he watched the heavens with the closest attention, hardly ever missing an observation during all that time, and, without any assistant, performed the whole business of the observatory himself.

Upon the accession of the late king, his consort queen Caroline made a visit at the Royal Observatory; and being pleased with every thing she saw, took notice that Dr. Halley had formerly served the crown as a captain in the navy: and she soon after obtained a grant of his half-pay for that commission, which he accordingly enjoyed from that time during his life. An offer was also made him of being appointed mathematical preceptor to the duke of Cumberland; but he declined that honour, on account of his advanced age, and the duties of his office. In 1729 he was chosen a foreign member of the Academy of Sciences at Paris.

About 1737 he was seized with a paralytic disorder in his right hand, which, it is said, was the first attack he ever felt upon his constitution: however, he came as usual once a week, till within a very short time of his death, to meet his friends in town on Thursdays, before the meeting of the Royal Society, at what is yet called Dr. Halley's club. His paralytic disorder increasing, his strength gradually wore away, till he expired Jan. 14, 1742, in the 86th year of his age; and his corps was interred in the church-yard of Lee near Blackheath.—Beside the works before mentioned, his principal publications are, 1. *Catalogus Stellarum Australium*. 2. *Tabula Astronomica*. 3. The Astronomy of Comets. With a great multitude of Papers in the Philos. Transf. from vol. 11 to vol. 60.

HALIFAX (JOHN). See SACROBOSCO.

HALO, or CORONA, a coloured circle appearing round the body of the sun, moon, or any of the larger stars.

Naturalists conceive the Halo to arise from a refraction



tion of the rays of light in passing through the fine rare vesiculæ of a thin vapour towards the top of the atmosphere.

Des Cartes observes, that the Halo never appears when it rains; whence he concludes that this phenomenon is occasioned by the refraction of light in the round particles of ice, which are then floating in the atmosphere; and to the different protuberance of these particles he ascribes the variation in the diameter of the Halo. Gassendi supposes, that a Halo is occasioned in the same manner as the rainbow; the rays of light being, in both cases, twice refracted and once reflected within each drop of rain or vapour, and that the difference between them is wholly owing to their different situation with respect to the observer. Dechales also endeavours to shew that the generation of the Halo is similar to that of the rainbow; and that the reason why the colours of the Halo are more dilute than those of the rainbow, is owing chiefly to their being formed, not in large drops of rain, but in very small vapour. But the most considerable and generally received theory, relating to the generation of Halos, is that of Mr. Huygens. This celebrated author supposes Halos, or circles round the sun, to be formed by small round grains of hail, composed of two different parts, the one of which is transparent, inclosing the other, which is opaque; which is the general structure actually observed in hail. He farther supposes that the grains or globules, that form these Halos, consisted at first of soft snow, and that they have been rounded by a continual agitation in the air, and thawed on their outside by the heat of the sun, &c. And he illustrates his ideas of their formation by geometrical figures.

Mr. Weidler endeavours to refute Huygens's manner of accounting for Halos, by a vast number of small vapours, each with a snowy nucleus, coated round with a transparent covering. He says, that when the sun paints its image in the atmosphere, and by the force of its rays puts the vapours in motion, and drives them toward the surface, till they are collected in such a quantity, and at such a distance from the sun on each side, that its rays are twice refracted, and twice reflected, when they reach the eye they exhibit the appearance of a Halo, adorned with the colours of the rainbow; which may happen in globular pellucid vapours without snowy nuclei, as appears by the experiment of hollow glass spheres filled with water: therefore, whenever those spherical vapours are situated as before mentioned, the refractions and reflections will happen every where alike, and the figure of a circular crown, with the usual order of colours, will be the consequence. *Philos. Trans.* number 458.

Newton's theory of Halos may be seen in his *Optics*, p. 155. And this curious theory was confirmed by actual observation in June 1692, when the author saw by reflection, in a vessel of stagnated water, three Halos, crowns, or rings of colours, about the sun, like three little rainbows concentric to his body. These crowns inclosed one another immediately, so that their colours proceeded in this continual order from the sun outward: blue, white, red; purple, blue, green, pale yellow, and red; pale blue, pale red. The like crowns sometimes appear about the moon. The more equal the globules of water or ice are to one another, the

more crowns of colours will appear, and the colours will be the more lively. *Optics*, p. 288.

There are several ways of exhibiting phenomena similar to these. The flame of a candle, placed in the midst of a steam in cold weather, or placed at the distance of some feet from a glass window that has been breathed upon, while the spectator is also at the distance of some feet from another part of the window, or placed behind a glass receiver, when air is admitted into the vacuum within it to a certain density, in each of these circumstances will appear to be encompassed by a coloured Halo. Also, a quantity of water being thrown up against the sun, as it breaks and disperses into drops, forms a kind of Halo or iris, exhibiting the colours of the natural rainbow. Musschenbroek observed, that when the glass windows of his room were covered with a thin plate of ice on the inside, the moon seen through it was surrounded with a large and variously coloured Halo; which, upon opening the window, he found arose entirely from that thin plate of ice, because none was seen except through this plate. Musschenbroek concludes his account of coronas with observing, that some density of vapour, or some thickness of the plates of ice, divides the light in its transmission either through the small globules or their interstices, into its separate colours; but what that density is, or what the size of the particles which compose the vapour, he does not pretend to determine. *Introd. ad Phil. Nat.* p. 1037.

It has often been observed that a Halo about the sun or moon, does not appear circular and concentric to the luminary, but oval and excentric, with its longest diameter perpendicular to the horizon, and extended from the moon farther downward than upward. Dr. Smith ascribes this phenomenon to the apparent concave of the sky being less than a hemisphere. When the angle which the diameter of a Halo subtends at the eye is  $45^\circ$  or  $46^\circ$ , and the bottom of the Halo is near the horizon, and consequently its apparent figure is most oval, the apparent vertical diameter is divided by the moon in the proportion of about 2 to 3 or 4, and is to the horizontal diameter drawn through the moon, as 4 to 3, nearly.—See farther on the subject of this article, Priestley's *Hist. of Discoveries relating to Vision*, p. 596—613; and Smith's *Optics*, art. 167, 513, 526, 527, &c.

HAMEL (JOHN BAPTISTE DU), a very learned French philosopher and writer, in the 17th century, was born in lower Normandy in 1614. At 18 years of age he published a treatise, in which he explained, in a very simple manner, and by one or two figures, Theodorus's 3 books upon Spherics; to which he added a tract upon trigonometry extremely perspicuous, and designed as an introduction to astronomy. He published afterwards various other works on astronomy and philosophy. Natural philosophy, as it was then taught, was only a collection of vague, knotty, and barren questions; when our author undertook to establish it upon right principles, and published his *Astronomia Physica*.

In 1666 M. Colbert proposed to Louis the 14th a scheme, which was approved of, for establishing a royal academy of sciences; and appointed our author secretary of it. In 1678, his *Philosophia Vetus & Nova*

was



was printed at Paris in 4 vols, 12mo; and in 1681 it was enlarged and printed there in 6 vols. He wrote several other pieces; and his works in this way were collected and published at Nuremberg 1681, in 4 volumes 4to, under the title of *Opera Philosophica & Astronomica*. These were highly valued then, though the improvements in philosophy since that time have rendered them of little or no use now.

In 1697 he resigned his place of secretary of the Royal Academy of Sciences; in which he was succeeded by M. Fontenelle. However, he published, in 1698, *Regiæ Scientiarum Academiæ Historia*, 4to, in four books; which being much liked, he afterwards augmented with two books more. This work contains an account of the foundation of the Royal Academy of Sciences, and its transactions, from 1666 to 1700, and is now the most useful of all his works. He was Regius Professor of Philosophy, in which office he was succeeded by M. Varignon, at his death, which happened Aug. 6, 1706, in the 93d year of his age.

**HANCES, HANCHES, or HANSES**, in architecture, are certain small intermediate parts of arches between the key or crown and the spring at the bottom, being perhaps about one-third of the arch, and situated nearer the bottom than the top or crown; and are otherwise called the *spandrels*.

**HANDSPIKE, or HANDSPEC**, a lever or piece of ash, elm, or other strong wood, for raising by the hand great weights, &c. It is 5 or 6 feet long, cut thin and crooked at the lower end, that it may get the easier between things that are to be separated, or under any thing that is to be raised. It is better than a crow of iron, because its length allows a better poise.

**HARD Bodies**, are such as are absolutely inflexible to any pressure or percussion whatever: differing from soft bodies, whose parts yield and are easily moved amongst one another, without restoring themselves again; and from elastic bodies, the parts of which also yield and give way, but presently restore themselves again to their former state and situation. Hence, hard bodies do not bend, or indent, but break. It is probable however there are no bodies in nature that are absolutely or perfectly either hard, soft, or elastic; but all possessing these qualities, more or less, in some degree. M. Bernoulli goes so far as to say that Hardness, in the common sense, is absolutely impossible, being contrary to the law of continuity.

The laws of motion for hard bodies are the same as for soft ones, both being supposed to adhere together on their impact. And these two sorts of bodies might be comprized under the common name of Unelastic.

**HARDENING**, the giving a greater degree of hardness to bodies than they had before.

There are several ways of Hardening iron and steel; as by hammering them, quenching them in cold water, &c.

**Cold-HARDENING**, is a superficial conversion of iron into steel, as if it were casing it, or covering it with a thin coat of harder matter. It is thus performed: Take cow horn or hoof, dry it well in an oven, and beat it to powder; put equal quantities of this powder and of bay salt into stale urine, or white wine vinegar, and mix them well together; cover the iron or steel all

over with this mixture, and wrap it up in loam, or plate iron, so as the mixture touch every part of the work; then put it in the fire, and blow the coals to it, till the whole lump have a blood red heat, but no higher; lastly, take it out, and quench it.—See **STEEL**, under which article are described other processes for this purpose.

**HARDNESS**, or Rigidity, that quality in bodies by which their parts so cohere as not to yield inward, or give way to an external impulse, without instantly going beyond the distance of their mutual attraction; and therefore are not subject to any motion in respect of each other, without breaking the body.

There were many fanciful opinions among the ancients concerning the cause of hardness; such as, heat, cold, dryness, the hooked figure of the particles of matter. The Cartesians make the Hardness of bodies to consist in rest, as that of soft and fluid ones in the motion of their particles.

Newton shews that the primary particles of all bodies, whether solid or fluid, are perfectly hard; and are not capable of being broken or divided by any power in nature. These particles, he maintains, are connected together by an attractive power; and according to the circumstances of this attraction, the body is either hard, or soft, or even fluid. If the particles be so disposed or fitted for each other, as to touch in large surfaces, the body will be hard; and the more so as those surfaces are the larger. If, on the contrary, they only touch in small surfaces, the body, by the weakness of the attraction, will remain soft.

At present, many philosophers think that Hardness consists in the absence or want of the action of the universal fluid, or elementary fire, among the particles of the body, or a deficiency of what is called latent heat; while on the contrary, fluidity, according to them, consists in the motion of the particles, in consequence of the action of that elementary fire.

Hardness appears to diminish the cohesion of bodies, in some degree, though their frangibility or brittleness does not by any means keep pace with their hardness. Thus, though glass be very hard and very brittle; yet flint is still harder, though less brittle. Among the metals, these two properties seem to be more connected, though even here the connection is by no means complete: for though steel be both the hardest and most brittle of all the metals; yet lead, which is the softest, is not the most ductile. Neither is Hardness connected with the specific gravity of bodies; for a diamond, the hardest substance in nature, is little more than half the weight of the lightest metal. And as little is it connected with the coldness, or electrical properties, or any other quality with which we are acquainted. Some bodies are rendered hard by cold, and others by different degrees of heat.

Mr. Quist and others have constructed tables of the Hardness of different substances. And the manner of constructing these tables, was by observing the order in which they were able to cut or make any impression upon one another. The following table, extracted from Magellan's edition of Cronstedt's Mineralogy, was taken from Quist, Bergman, and Kirwan. The first column shews the Hardness, and the second the specific gravity.

Diamond



	<i>Hard- ness.</i>	<i>Spec. Grav.</i>
Diamond from Ormus	20	3.7
Pink diamond	19	3.4
Blueish diamond	19	3.3
Yellowish diamond	19	3.3
Cubic diamond	18	3.2
Ruby	17	4.2
Pale blue sapphire	17	3.8
Pale ruby from Brazil	16	3.5
Deep blue sapphire	16	3.8
Topaz	15	4.2
Whitish ditto	14	3.5
Ruby spinell	13	3.4
Emerald	12	2.8
Garnet	12	4.4
Agate	12	2.6
Onyx	12	2.6
Sardonyx	12	2.6
Bohemian topaz	11	2.8
Occid. amethyst	11	2.7
Crystal	11	2.6
Carnelian	11	2.7
Green jasper	11	2.7
Schoerl	10	3.6
Tourmaline	10	3.0
Quartz	10	2.7
Opal	10	2.6
Chrysolite	10	3.7
Reddish yellow jasper	9	2.6
Zeolyte	8	2.1
Fluor	7	3.5
Calcareous spar	6	2.7
Gypsum	5	2.3
Chalk	3	2.7

**HARMONICA, HARMONICS**, a branch or division of the ancient music; being that part which considers the differences and proportions of sounds, with respect to acute and grave; as distinguished from Rhythmica, and Metrica.

Mr. Malcolm has made a very industrious and learned enquiry into the Harmonica, or harmonic principles, of the ancients.

**HARMONICA**, the name of a musical instrument invented by Dr. Franklin, consisting of the glasses, called musical glasses.

It is said that the first hint of musical glasses is to be found in an old English book, in which a number of various amusements were described. That author directs his pupil to choose half a dozen drinking-glasses; to fill each of them with water in proportion to the gravity or acuteness of the sound which he intended it to give; and having thus adjusted them one to another, he might entertain the company with a church tune. These were perhaps the rude hints which Mr. Puckeridge, an Irish gentleman, afterwards improved, and after him, Mr. E. Delaval, an ingenious member of the Royal Society; and finally brought to perfection by the celebrated Franklin. See the history and description in his Letters, particularly in that to Beccaria.

**HARMONICAL, or HARMONIC**, something relating to Harmony. Thus,

**HARMONICAL Arithmetic**, is so much of the theory

and doctrine of numbers, as relates to making the comparisons and reductions of musical intervals, which are expressed by numbers, for finding their mutual relations, compositions, and resolutions.

**HARMONICAL Composition**, in its general sense, includes the composition both of harmony and melody; i. e. of music, or song, both in a single part, and in several parts.

**HARMONICAL Interval**, the difference between two sounds, in respect of acute and grave: or that imaginary space terminated with two sounds differing in acuteness or gravity.

**HARMONICAL Proportion, or Musical Proportion**, is that in which the first term is to the third, as the difference of the first and second is to the difference of the 2d and 3d; or when the first, the third, and the said two differences, are in geometrical proportion. Or, four terms are in Harmonical proportion, when the 1st is to the 4th, as the difference of the 1st and 2d is to the difference of the 3d and 4th. Thus, 2, 3, 6, are in harmonical proportion, because  $2 : 6 :: 1 : 3$ . And the four terms 9, 12, 16, 24 are in harmonical proportion, because  $9 : 24 :: 3 : 8$ .—If the proportional terms be continued in the former case, they will form an harmonical progression, or series.

1. The reciprocals of an arithmetical progression are in Harmonical progression; and, conversely, the reciprocals of Harmonicals are arithmeticals. Thus, the reciprocals of the Harmonicals 2, 3, 6, are  $\frac{1}{2}, \frac{1}{3}, \frac{1}{6}$ ,

which are arithmeticals; for  $\frac{1}{2} - \frac{1}{3} = \frac{1}{6}$ , and  $\frac{1}{3} - \frac{1}{6} = \frac{1}{6}$  also: and the reciprocals of the arith-

meticals 1, 2, 3, 4, &c, are  $\frac{1}{1}, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \&c$ , which

are Harmonicals; for  $\frac{1}{1} : \frac{1}{3} :: \frac{1}{2} - \frac{1}{2} : \frac{1}{2} - \frac{1}{3}$ ; and so on. And, in general, the reciprocals of the arithmeticals  $a, a + d, a + 2d, a + 3d, \&c$ , viz,  $\frac{1}{a}, \frac{1}{a + d}, \frac{1}{a + 2d}, \frac{1}{a + 3d}, \&c$ , are Harmonicals; et e contra.

2. If three or four numbers in Harmonical proportion be either multiplied or divided by some number, the products, or the quotients, will still be in Harmonical proportion. Thus,

the Harmonicals	-	6, 8, 12,
multiplied by 2 give		12, 16, 24,
or divided by 2 give		3, 4, 6,

which are also Harmonicals.

3. To find a Harmonical mean proportional between two terms: Divide double their product by their sum.

4. To find a 3d term in Harmonical proportion to two given terms: Divide their product by the difference between double the 1st term and the 2d term.

5. To find a 4th term in Harmonical proportion to three terms given: Divide the product of the 1st and 3d by the difference between double the 1st and the 2d term.

Hence,



Hence, of the two terms  $a$  and  $b$ ;

the Harmonical mean is  $\frac{2ab}{a+b}$ ,

the 3d Harmonical propor. is  $\frac{ab}{2a-b}$ ,

also to  $a, b, c$ , the 4th Harm. is  $\frac{ac}{2a-b}$ .

6. If there be taken an arithmetical mean, and a Harmonical mean, between any two terms, the four terms will be in geometrical proportion. Thus, between 2 and 6,

the arithmetical mean is 4, and

the Harmonical mean is 3;

and hence  $2 : 3 :: 4 : 6$ .

Also, between  $a$  and  $b$ ,

the arithmetical mean is  $\frac{a+b}{2}$ , and

the Harmonical mean is  $\frac{2ab}{a+b}$ ;

but  $a : \frac{2ab}{a+b} :: \frac{a+b}{2} : b$ .

HARMONY, in Music, the agreeable result of an union of several musical sounds, heard at one and the same time; or the mixture of divers sounds, which together have an agreeable effect on the ear.

As a continued succession of musical sounds produces melody, so a continued combination of them produces Harmony.

Among the ancients however, as sometimes also among the moderns, Harmony is used in the strict sense of consonance; and so is equivalent to the symphony.

The words *concord* and Harmony do really signify the same thing; though custom has made a little difference between them. Concord is the agreeable effect of two sounds in consonance; and Harmony the effect of any greater number of agreeable sounds in consonance.

Again, Harmony always implies consonance; but concord is also applied to sounds in succession; though never but where the terms can stand agreeably in consonance. The effect of an agreeable succession of several sounds, is called *melody*; as that of an agreeable consonance is called Harmony.

Harmony is well defined, the sum or result of the combination of two or more concords; that is, of three or more simple sounds striking the ear all together; and different compositions of concords make different Harmony.

The ancients seem to have been entirely unacquainted with Harmony, the soul of the modern music. In all their explications of the *melopœia*, they say not one word of the concert or Harmony of parts. We have instances, indeed, of their joining several voices, or instruments, in consonance; but then these were not so joined, as that each had a distinct and proper melody, so making a succession of various concords; but they were either unisons, or octaves, in every note; and so all performed the same individual melody, and constituted one song.

When the parts differ, not in the tension of the whole, but in the different relations of the successive notes, it is this that constitutes the modern art of Harmony.

HARMONY of the Spheres, or *Celestial Harmony*, a kind of music much spoken of by many of the ancient philosophers and fathers, supposed to be produced by the sweetly tuned motions of the stars and planets. This Harmony they attributed to the various proportionate impressions of the heavenly bodies upon one another, acting at proper intervals. They think it impossible that such prodigious bodies, moving with such rapidity, should be silent: on the contrary, the atmosphere, continually impelled by them, must yield a set of sounds proportionate to the impression it receives; and that consequently, as they run all in different circuits, and with various degrees of velocity, the different tones arising from the diversity of motions, directed by the hand of the Almighty, must form an agreeable symphony or concert.

They therefore supposed, that the moon, as being the lowest of the planets, corresponded to *mi*; Mercury, to *fa*; Venus, to *sol*; the Sun, to *la*; Mars, to *fi*; Jupiter, to *ut*; Saturn, to *re*; and the orb of the fixed stars, as being the highest of all, to *mi*, or the octave.

It is thought that Pythagoras had a view to the gravitation of celestial bodies, in what he taught concerning the Harmony of the spheres.

A musical chord gives the same note as one double in length, when the tension or force with which the latter is stretched is quadruple; and the gravity of a planet is quadruple of the gravity of a planet at a double distance. In general, that any musical chord may become unison to a lesser chord of the same kind, its tension must be increased in the same proportion as the square of its length is greater; and that the gravity of a planet may become equal to the gravity of another planet nearer the sun, it must be increased in proportion as the square of its distance from the sun is greater. If therefore we should suppose musical chords extended from the sun to each planet, that all these chords might become unison, it would be requisite to increase or diminish their tensions in the same proportions as would be sufficient to render the gravities of the planets equal; and from the similitude of those proportions, the celebrated doctrine of the Harmony of the spheres is supposed to have been derived.

Kepler wrote a large work, in folio, on the Harmonies of the world, and particularly of that of the celestial bodies. He first endeavoured to find out some relation between the dimensions of the five regular solids and the intervals of the planetary spheres; and imagining that a cube, inscribed in the sphere of Saturn, would touch by its six planes the sphere of Jupiter, and that the other four regular solids in like manner fitted the intervals that are between the spheres of the other planets, he became persuaded that this was the true reason why the primary planets were precisely six in number, and that the author of the world had determined their distances from the sun, the centre of the system, from a regard to this analogy. But afterwards finding that the disposition of the five regular solids amongst the planetary spheres, was not agreeable to the intervals between their orbits, he endeavoured to discover other schemes of Harmony. For this purpose he compared the motions of the same planet at its greatest and least distances, and of the different planets in their several orbits, as they would appear viewed from



from the sun; and here he fancied that he found a similitude to the divisions of the octave in music. Lastly, he imagined that if lines were drawn from the earth, to each of the planets, and the planets appended to them, or stretched by weights proportional to the planets, these lines would then sound all the notes in the octave of a musical chord.

See his *Harmonics*; also *Plin. lib. 2, cap. 22*; *Macrob. in Somn. Scip. lib. 2, cap. 1*; *Plutarch de Animal. Procreatione, à Timæo*; and *Maclaurin's View of Newton's Discov. book 1, chap. 2.*

**HARQUEBUSS**, a hand-gun, or a fire-arm of a proper length and weight to be borne in the arm. Hanzelet prescribes its proper length to be 40 calibres, or diameters of its bore; and the weight of its ball 1 oz. and  $\frac{7}{8}$ ; its charge of powder as much.

**HARRIOT (THOMAS)**, a very eminent English mathematician and astronomer, was born at Oxford in 1560, and died at London July 2, 1621, in the 61st year of his age. Harriot has hitherto been known to the world only as an algebraist, though a very eminent one; but from his manuscript papers, that have been but lately discovered by Dr. Zach, astronomer to the duke of Saxe-Gotha, it appears that he was not less eminent as an astronomer and geometrician. Dr. Zach has printed an account of those papers, in the *Astronomical Ephemeris of the Royal Academy of Sciences at Berlin*, for the year 1788; of which, as it is very curious, and contains a great deal of information, I shall here give a translation, to serve as memoirs concerning the life and writings of this eminent man; afterwards adding only some necessary remarks of my own.

"I here present to the world (says Dr. Zach), a short account of some valuable and curious manuscripts, which I found in the year 1784, at the seat of the earl of Egremont, at Petworth in Sussex, in hopes that this learned and inquisitive age will either think my endeavours about them worthy of its assistance, or else will be thereby induced to attempt some other means of publishing them. The only undeniable proof I can now produce of the usefulness of such an undertaking, is by giving a succinct report of the contents of these materials, and briefly shewing what may be effected by them. And although I come to the performance of such an enterprize with much less abilities than the different parts of it require, yet I trust that my love for truth, my design and zeal to vindicate the honour due to an Englishman, the author of these manuscripts, which are the chief reasons that have influenced me in this undertaking, will serve as my excuse.

"A predecessor of the family of lord Egremont, viz, that noble and generous earl of Northumberland, named Henry Percy, was not only a generous favourer of all good learning, but also a patron and Mæcenas of the learned men of his age. Thomas Harriot, the author of the said manuscripts, Robert Hues (well known by his *Treatise upon the Globes*), and Walter Warner, all three eminent mathematicians, who were known to the earl, received from him yearly pensions; so that when the earl was committed prisoner to the Tower of London in the year 1606, our author, with Hues and Warner, were his constant companions; and were usually called the earl of Northumberland's three Magi.

"Thomas Harriot is a known and celebrated mathematician among the learned of all nations, by his excellent work, *Artis Analytica Praxis, ad æquationes algebraicas nova expedita & generali methodo, resolvendas, Tractatus posthumus*; Lond. 1631: dedicated to Henry earl of Northumberland; published after his death by Walter Warner. It is remarkable, that the fame and the honour of this truly great man were constantly attacked by the French mathematicians; who could not endure that Harriot should in any way diminish the fame of their Vieta and Des Cartes, especially the latter, who was openly accused of plagiarism from our author. [See *Montucla's Histoire des Mathematiques, part 3, p. 485 & seq.*—*Lettres de M. Des Cartes, tom. 3, pa. 457, edit. Paris 1667, in 4to.*—*Dictionnaire de Moreri, word Harriot.*—*Encyclopedie, word Algebra.*—*Lettres de M. de Voltaire, sur la nation Angloise, lettre 14.*—*Memoire de l'Abbé de Gua dans les Mem. de l'Acad. des Sciences de Paris pour 1741.*—*Jer. Collier's great Historical Dictionary, word Harriot.*—*Dr. Wallis's preface to his Algebra.*—*To which may be added the article Algebra, in this dictionary.*]

"Des Cartes published his *Geometry* 6 years after Harriot's work appeared, viz, in the year 1637. Sir Charles Cavendish, then ambassador at the French court at Paris, when Des Cartes's *Geometry* made its first appearance in public, observed to the famous geometrician Roberval, that these improvements in Analysis had been already made these 6 years in England, and shewed him afterwards Harriot's *Artis Analytica Praxis*, which as Roberval was looking over, at every page he cried out, *Oui! oui! il l'a vu! Yes! yes! he has seen it!* Des Cartes had also been in England before Harriot's death, and had heard of his new improvements and inventions in Analysis. A critical life of this man, which his papers would enable me to publish, will shew more clearly what to think upon this matter, which I hope may be discussed to the due honour of our author.

"Now all this relates to Harriot the celebrated analyst; but it has not hitherto been known that Harriot was an eminent astronomer, both theoretical and practical, which first appears by these manuscripts; among which, the most remarkable are 199 observations of the Sun's Spots, with their drawings, calculations and determinations of the sun's rotation about his axis. There is the greatest probability that Harriot was the first discoverer of these spots, even before either Galileo or Scheiner. The earliest intelligence we have of the first discovered solar spots, is of one Joh. Fabricius Phrysius, who in the year 1611 published at Wittenberg a small treatise, intitled, *De Maculis in Sole observatis & apparente eorum cum Sole conversione narratio*. Galileo, who is commonly accounted the first discoverer of the Solar Spots, published his book, *Istoria e Dimostrazioni intorne alle Macchie Solare e loro accidenti*, at Rome, in the year 1613. His first observation in this work, is dated June 2, 1612. Angelo de Filiis, the editor of Galileo's work, who wrote the dedication and preface to it, mentions, pa. 3, that Galileo had not only discovered these spots in the month of April in the year 1611, at Rome, in the Quirinal Garden, but had shewn them several months before (*molti mesi innanzi*) to his friends in Florence. And that the observations of the disguised Apelles (the Jesuit Scheiner, a pretender



tender to this first discovery) were not later than the month of October in the same year; by which the epoch of this discovery was fixed to the beginning of the year 1611. But a passage in the first letter of Galileo's works, pa. 11, gives a more precise term to this discovery. Galileo there says in plain terms, that he had observed the Spots in the Sun 18 months before. The date of this letter is May 24, 1612; which brings the true epoch of this discovery to the month of November, 1610. However, Galileo's first produced observations are only from June 2, 1612, and those of father Scheiner of the month of October, in the same year. But now it appears from Harriot's manuscripts, that his first observations of these Spots are of Dec. 8, 1610. It is not likely that Harriot could have this notice from Galileo, for I do not find this mathematician's name ever quoted in Harriot's papers. But I find him mentioning Josephus a Costa's book 1, chap. 2, of his *Natural and Moral History of the West Indies*, in which he relates that in Peru there are Spots to be seen in the Sun which are not to be seen in Europe. It rather seems that Harriot had taken the hint from thence. Besides, it is very likely that Harriot, who lived with so generous a patron to all good learning and improvements, had got the new invention of telescopes in Holland much sooner in England, than they could reach Galileo, who at that time lived at Venice. Harriot's very careful and exact observations of these Spots, shew also that he was in possession of the best and most improved telescopes of that time; for it appears he had some with magnifying powers of 10, 20, and 30 times. At least there are no earlier observations of the Solar Spots extant than his: they run from December 8, 1610, till January 18, 1613. I compared the corresponding ones with these observed by Galileo, between which I found an exact agreement. Had Harriot had any notion about Galileo's discoveries, he certainly would have also known something about the phases of Venus and Mercury, and especially about the singular shape of Saturn, first discovered by Galileo; but I find not a word in all his papers concerning the particular figure of that planet.

“ *Of Jupiter's Satellites.* I found among his papers a large set of observations, with their drawing, position, and calculations of their revolutions and periods. His first observation of those discovered Satellites, I find to be of January 16, 1610; and they go till February 26, 1612. Galileo pretends to have discovered them January 7, 1610; so that it is not improbable that Harriot was likewise the first discoverer of these attendants of Jupiter.

“ Among his other observations of the Moon, of some Eclipses, of the planet Mars, of Solstices, of Refraction, of the Declination of the Needle; there are most remarkable ones of the noted Comets of 1607, and of 1618, the latter, for there were two this year (*see Kepler de Cometis, pa. 49*). They were all observed with a cross-staff, by measuring their distances from fixed stars; whence these observations are the more valuable, as comets had before been but grossly observed: Kepler himself observed the comet of 1607 only with the naked eye, pointing out its place by a coarse estimation, without the aid of an instrument; and the elements of their orbits could, in defect of better observa-

tions, be only calculated by them. The observations of the comet of the year 1607, are of the more importance, even now for modern astronomy, as this is the same comet that fulfilled Dr. Halley's prediction of its return in the year 1759. That prediction was only grounded upon the elements afforded him by these coarse observations; for which reason he only assigned the term of its return to the space of a year. The very intricate calculations of the perturbations of this comet, afterwards made by M. Clairaut, reduced the limits to a month's space. But a greater light may now be thrown upon this matter by the more accurate observations on this comet by Mr. Harriot. In the month of October 1785, when I conversed upon the subject of Harriot's papers, and especially on this comet, with the very celebrated mathematician M. de la Grange, director of the Royal Academy of Sciences at Berlin; he then suggested to me an idea, which, if brought into execution, will clear up an important point in astronomy. It is well known to astronomers how difficult a matter it is, to determine the mass, or quantity of matter, in the planet Saturn; and how little satisfactory the notions of it are, that have hitherto been formed. The whole theory of the perturbations of comets depending upon this uncertain datum, several attempts and trials have been made towards a more exact determination of it by the most eminent geometricians of this age, and particularly by la Grange himself; but never having been satisfied with the few and uncertain data heretofore obtained for the resolution of this problem, he thought that Harriot's observations on the comet of 1607, and the modern ones of the same comet in 1759, would suggest a way of resolving the problem *a posteriori*; that of determining by them the elements of its ellipsis. The retardation of the comet compared to its period, may clearly be laid to the account of the attraction and perturbation it has suffered in the region of Jupiter and Saturn; and as the part of it belonging to Jupiter is very well known, the remainder must be the share which is due to Saturn; from whence the mass of the latter may be inferred. In consequence of this consideration I have already begun to reduce most of Harriot's observations of this comet, in order to calculate by them the true elements of its orbit on an elliptical hypothesis, to complete M. la Grange's idea upon this matter.

“ I forbear to mention here any more of Harriot's analytical papers, which I found in a very great number. They contain several elegant solutions of quadratic, cubic, and biquadratic equations; with some other solutions and *locæ geometricæ*, that shew his eminent qualifications, and will serve to vindicate them against the attacks of several French writers, who refuse him the justice due to his skill and accomplishments, merely to save Des Cartes's honour, who yet, by some impartial men of his own nation, was accused of public plagiarism.

“ Thomas Harriot was born at Oxford, in the year 1560. After being instructed in the rudiments of languages, he became a commoner of St. Mary's-Hall, where he took the degree of bachelor of arts in 1579. He had then so distinguished himself by his uncommon skill in mathematics, as to be recommended soon after to Sir Walter Raleigh, as a proper preceptor to him in that science. Accordingly that noble knight became his first patron, took him into his family, and allowed



him a handsome pension. In 1584 he went over with Sir Walter's first colony to Virginia; where he was employed in discovering and surveying the country, &c; maps of which I have found (says Dr. Zach) very neatly done among his papers. After his return he published *A Brief and True Report of the Newfoundland of Virginia, of the Commodities there found to be raised, &c*; Lond. 1588. This was reprinted in the 3d volume of Hakluyt's Voyages: it was also translated into Latin, and printed at Frankfort in the year 1590. Sir Walter introduced him to the acquaintance of the earl of Northumberland, who allowed him a yearly pension; Wood says, of 120l. only; but by some of his receipts, which I have found among his papers, it appears that he had 300l, which indeed was a very large sum at that time. Wood, in his *Athen. Oxon.* mentions nothing of Harriot's papers, except a manuscript in the library at Sion College, London, entitled *Ephemeris Chyrometrica*. I got access to this library and manuscripts, and was indeed in hopes of finding something more of Harriot's; for most of his observations are dated from Sion College; but I could not find any thing from Harriot himself. I found indeed some other papers of his friends: he mentions, in his observations, one Mr. Standish, at Oxford, and Nicol. Torperly, who also was of the acquaintance of the earl of Northumberland, and had a yearly pension: from the former I found two observations of the same comet of 1618, made in Oxford, which he communicated to Mr. Harriot.

" Thomas Harriot died July 2, 1621. His disease was a cancerous ulcer in the lip, which some pretend he got by a custom he had of holding the mathematical brass instruments, when working, in his mouth. I found several of his letters, and answers to them, from his physician Dr. Alexander Rhead, who in his treatise mentions Harriot's disease. His body was conveyed to St. Christopher's church, in London. Over his grave was soon after erected a monument, with a large inscription, which was destroyed with the church itself by the dreadful fire of September 1666. He was but 60 years of age."

The peculiar nature and merits of Harriot's Algebra, we have spoken largely and particularly of, under the art. *Algebra*, page 89. As to his manuscripts lately discovered by Dr. Zach, as above mentioned, it is with pleasure I can announce, that they are in a fair train to be published: they have been presented to the university of Oxford, on condition of their printing them; with a view to which, they have been lately put into the hands of an ingenious member of that learned body, to arrange and prepare them for the press.

HARRISON (JOHN), a most accurate mechanic, the celebrated inventor of the famous *time-keeper* for ascertaining the longitude at sea, and also of the compound or *gridiron-pendulum*; was born at Foulby, near Pontefract in Yorkshire, in 1693. His father was a carpenter, in which profession the son assisted; occasionally also, according to the miscellaneous practice of country artists, surveying land, and repairing clocks and watches; and young Harrison always was, from his early childhood, greatly attached to any machinery moving by wheels. In 1700 he removed with his father to Barrow in Lincolnshire; where, though his opportunities of acquiring knowledge were very few, he eagerly improved every incident from which he might

collect information; frequently employing all or great part of his nights in writing or drawing: and he always acknowledged his obligations to a clergyman who came every Sunday to officiate in the neighbourhood, who lent him a MS. copy of professor Sanderson's lectures; which he carefully and neatly transcribed, with all the diagrams. His native genius exerted itself superior to these solitary disadvantages; for in the year 1726, he had constructed two clocks, mostly of wood, in which he applied the escapement and compound pendulum of his own invention: these surpassed every thing then made, scarcely erring a second in a month. In 1728, he came up to London with the drawings of a machine for determining the longitude at sea, in expectation of being enabled to execute one by the Board of Longitude. Upon application to Dr. Halley, the astronomer royal, he referred him to Mr. George Graham; who advised him to make his machine, before applying to that Board. He accordingly returned home to perform his task; and in 1735 came to London again with his first machine; with which he was sent to Lisbon the next year, to make trial of it. In this short voyage, he corrected the dead reckoning about a degree and a half; a success which procured him both public and private encouragement. About the year 1739 he completed his second machine, of a construction much more simple than the former, and which answered much better: this, though not sent to sea, recommended Mr. Harrison yet stronger to the patronage of his friends and the public. His third machine, which he produced in 1749, was still less complicated than the second, and more accurate, as erring only 3 or 4 seconds in a week. This he conceived to be the *ne plus ultra* of his attempts; but by endeavouring to improve pocket-watches, he found the principles he applied to surpass his expectations, so much, as to encourage him to make his fourth time-keeper, which is in the form of a pocket-watch, about 6 inches diameter. With this time-keeper his son made two voyages, the one to Jamaica, and the other to Barbadoes; in which experiments it corrected the longitude within the nearest limits required by the act of the 12th of queen Anne; and the inventor had therefore, at different times, more than the proposed reward, receiving from the Board of Longitude at different times almost 24,000l. besides a few hundreds from the East India Company, &c. These four machines were given up to the Board of Longitude. The three former were not of any use, as all the advantages gained by making them, were comprehended in the last: being worthy however of preservation, as mechanical curiosities, they are deposited in the Royal Observatory at Greenwich. The fourth machine, emphatically distinguished by the name of *The time-keeper*, was copied by the ingenious Mr. Kendal; and that duplicate, during a three years circumnavigation of the globe in the southern hemisphere by captain Cook, answered as well as the original.

The latter part of Mr. Harrison's life was employed in making a fifth improved time-keeper, on the same principles with the preceding one; which, after a ten-weeks trial, in 1772, at the king's private observatory at Richmond, erred only  $4\frac{1}{2}$  seconds. Within a few years of his death, his constitution visibly declined; and he had frequent fits of the gout, a disorder that



never attacked him before his 77th year. His constitution at last yielding to the infirmities of old age, he died at his house in Red-Lion Square, in 1776, at 83 years of age.

Like many other mere mechanics, Mr. Harrison found a difficulty in delivering his sentiments in writing (at least in the latter periods of his life, when his faculties were much impaired) in which he adhered to a peculiar and uncouth phraseology. This was but too evident in his *Description concerning such Mechanism as will afford a nice or true Mensuration of Time*, &c. 8vo, 1775. This small work includes also an account of his new musical scale; being a mechanical division of the octave, according to the proportion which the radius and diameter of the circle have respectively to the circumference. He had in his youth been the leader of a band of church-fingers; had a very delicate ear for music; and his experiments on sound, with a curious monochord of his own improvement, it has been said were not less accurate than those he was engaged in for the mensuration of time.

HAUTEFEUILLE (JOHN), an ingenious mechanic, born at Orleans in 1674. He made a great progress in mechanics in general, but had a particular taste for clock-work, and made several discoveries in it that were of singular use. It was he it seems who found out the secret of moderating the vibration of the balance by means of a small steel-spring, which has since been made use of. This discovery he laid before the members of the Academy of Sciences in 1694; and these watches are, by way of eminence, called *pendulum-watches*; not that they have real pendulums, but because they nearly approach to the justness of pendulums. M. Huygens perfected this happy invention; but having declared himself the inventor and obtained a patent for making watches with spiral springs, the abbé Feuille opposed the registering of it, and published a piece on the subject against Huygens. He died in 1724, at 50 years of age. Besides the above,

He wrote a great many other pieces, most of which are small pamphlets, but very curious: as, 1. His Perpetual Pendulum. 2. New Inventions. 3. The Art of Breathing under Water, and the means of preserving a Flame shut up in a Small Place. 4. Reflections on machines for Raising Water. 5. His Opinion on the different Sentiments of Mallebranche and Regis, relating to the Appearance of the Moon when seen in the Horizon. 6. The Magnetic Balance. 7. A Placet to the king on the Longitude. 8. Letter on the Secret of the Longitude. 9. A New System on the Flux and Reflux of the Sea. 10. The means of making Sensible Experiments that prove the Motion of the Earth: and many other pieces.

HAYES (CHARLES, Esq.), a very singular person, whose great erudition was so concealed by his modesty, that his name is known to very few, though his publications are many. He was born in 1678, and died in 1760, at 82 years of age. He became distinguished in 1704 by a Treatise of Fluxions, in folio, being we believe the first treatise on that science ever published in the English language; and the only work to which he ever set his name. In 1710 came out a small 4to pamphlet, in 19 pages, intitled, A New and easy Method to find out the Longitude from observing the Altitudes of the Celestial

bodies. Also, in 1723, he published, *The Moon, a Philosophical Dialogue*; tending to shew, that the moon is not an opaque body, but has native light of her own.

To a skill in the Greek and Latin, as well as the modern languages, he added the knowledge of the Hebrew: and he published several pieces relating to the translation and chronology of the scriptures. During a long course of years he had the chief management of the late African company, being annually elected sub-governor. But on the dissolution of that company in 1752, he retired to Down in Kent, where he gave himself up to study; from whence however he returned in 1758, to chambers in Gray's inn, London, where he died in 1760, as mentioned above.

He left a posthumous work, that was published in 8vo, under the title of *Chronographia Asiatica et Aegyptiaca &c.*

HEAT, the opposite to cold, being a relative term denoting the property of fire, or of those bodies we denominate hot; being in us a sensation excited by the action of fire.

Heat, as it exists in the hot body, or that which constitutes and denominates a body hot, and enables it to produce such effects on our organs, is variously considered by the philosophers: some making it a quality, others a substance, and others only a mechanical affection. The former principle is laid down by Aristotle and the Peripatetics. While the Epicureans, and other corpuscularians, define Heat not as an accident of fire, but as an essential power or property of it, the same in reality with it, and only distinguished from it in the manner of our conception. So that Heat, on their principles, is no other than the volatile substance of fire itself, reduced into atoms, and emitted in a continual stream from ignited bodies; so as not only to warm the objects within its reach, but also, if they be inflammable, to kindle them, turn them into fire, and conspire with them to make flame. In effect, these corpuscles, say they, flying off from the ignited body, constitute fire while yet contained within the sphere of its flame; but when fled, or got beyond the same, and dispersed every way, so as to escape the apprehension of the eye, and only to be perceived by the feeling, they take the denomination of Heat, inasmuch as they excite in us that sensation.

The Cartesians, improving on this doctrine, assert that Heat consists in a certain motion of the insensible particles of a body, resembling the motion by which the several parts of our body are agitated by the motion of the heart and blood.

Our latest and best writers of mechanical, experimental, and chemical philosophy, differ very considerably about Heat. The chief difference is, whether it be a peculiar property of one certain immutable body, called fire, or phlogiston, or electricity; or whether it may be produced mechanically in other bodies, by inducing an alteration in their particles. The former tenet, which is as ancient as Democritus, and the system of atoms, had given way to that of the Cartesians, and other mechanists; but is now with great address retrieved, and improved on, by some of the latest writers, particularly Homburg, the younger Lemery, Gravesande, Boerhaave in his lectures on fire, Black, Crawford, and other chemical philosophers.



The thing called fire, according to Boerhaave, is a body sui generis, created such ab origine, unalterable in its nature and properties, and not either producible de novo from any other body, nor capable of being reduced into any other body, or of ceasing to be fire. This fire, he contends, is diffused equably every where, and exists alike, or in equal quantity, in all the parts of space, whether void, or possessed by bodies; but that naturally, and in itself, it is perfectly latent and imperceptible; being only discovered by certain effects which it produces, and which are cognizable by our senses. These effects are Heat, light, colour, rarefaction, and burning, which are all indications of fire, as being none of them producible by any other cause: so that wherever we observe any of these, we may safely infer the action and presence of fire. But though the effect cannot be without the cause, yet the fire may remain without any of these effects; any, we mean, gross enough to affect our senses, or become objects of them: and this, he adds, is the ordinary case; there being a concurrence of other circumstances, which are often wanting, necessary to the production of such sensible effects.

The mechanical philosophers, particularly Bacon, Boyle, and Newton, conceive otherwise of Heat; considering it not as an original inherent property of any particular sort of body; but as mechanically producible in any body. The former, in an express treatise *De Forma Calidi*, from a particular enumeration of the several phenomena and effects of Heat, deduces several general properties of it; and hence he defines Heat, an expansive undulatory motion in the minute particles of the body; by which they tend, with some rapidity, towards the circumference, and at the same time incline a little upwards.

Mr. Boyle, in a Treatise on the Mechanical Origin of Heat and Cold, strongly supports the doctrine of the producibility of Heat, with new observations and experiments; as in the instance of a smith briskly hammering a small piece of iron, which, though cold before, soon becomes exceedingly hot.

This system is also farther supported by Newton, who does not conceive fire as any particular species of body, originally endued with such and such properties. Fire, according to him, is only a body much ignited, that is heated hot, so as to emit light copiously: what else, says he, is red-hot iron but fire? and what else is a burning charcoal but red-hot wood? or flame itself, but red-hot smoke? It is certain that flame is only the volatile part of the fuel heated red-hot, i. e. so hot as to shine; and hence only such bodies as are volatile, that is, such as emit a copious fume, will flame; nor will they flame longer than they have fume to burn. In distilling hot spirits, if the head of the still be taken off, the ascending vapours will catch fire from a candle, and turn into a flame. And in the same manner several bodies, much heated by motion, attrition, fermentation, or the like, will emit lucid fumes, which, if they be copious enough, and the heat sufficiently great, will be flame; and the reason why fused metals do not flame, is the smallness of their fume; this is evident, because spelter, which fumes most copiously, does likewise flame. Add, that all flaming bodies, as oil, tallow, wax, wood, pitch, sulphur, &c, by flaming, waste and vanish into burning smoke. And do not all fixed bodies, when heated beyond a certain degree, emit light,

and shine? and is not this emission performed by the vibrating motion of their parts? and do not all bodies, which abound with terrestrial and sulphureous parts, emit light as often as those parts are sufficiently agitated, whether that agitation be made by external fire, or by friction, or percussion, or putrefaction, or by any other cause? Thus, sea water, in a storm; quicksilver agitated in vacuo; the back of a cat, or the neck of a horse, obliquely rubbed in a dark place; wood, flesh, and fish, while they putrefy; vapours from putrefying waters, usually called *ignes fatui*; stacks of moist hay or corn; glow-worms; amber and diamonds by rubbing; fragments of steel struck off with a flint, &c, all emit light. Are not gross bodies and light convertible into one another? and may not bodies receive much of their activity from the particles of light which enter their composition? I know no body less apt to shine than water; and yet water, by frequent distillations, changes into fixed earth, which, by a sufficient Heat, may be brought to shine like other bodies.

Add, that the sun and stars, according to Newton's conjecture, are no other than great earths vehemently heated: for large bodies, he observes, preserve their Heat the longest, their parts heating one another; and why may not great, dense, and fixed bodies, when heated beyond a certain degree, emit light so copiously, as by the emission and reaction of it, and the reflections and refractions of the rays within the pores, to grow still hotter, till they arrive at such a period of Heat as is that of the sun? Their parts also may be farther preserved from fuming away, not only by their fixity, but by the vast weight and density of their atmospheres incumbent on them, thus strongly compressing them, and condensing the vapours and exhalations arising from them. Hence we see warm water, in an exhausted receiver, shall boil as vehemently as the hottest water open to the air; the weight of the incumbent atmosphere, in this latter case, keeping down the vapours, and hindering the ebullition, till it has conceived its utmost degree of Heat. So also a mixture of tin and lead, put on a red-hot iron in vacuo, emits a fume and flame; but the same mixture in the open air, by reason of the incumbent atmosphere, does not emit the least sensible flame.

Thus much for the system of the producibility of Heat.

On the other hand, M. Homberg, in his *Essai du Soufre Principe*, holds, that the chemical principle or element, sulphur, which is supposed one of the simple, primary, pre-existent ingredients of all natural bodies, is real fire; and consequently that fire is co-eval with body. *Mem. de l'Acad. an. 1705.*

Dr. Gravesande goes upon much the same principle. According to him, fire enters the composition of all bodies, is contained in all bodies, and may be separated or procured from all bodies, by rubbing them against each other, and thus putting their fire in motion. But fire, he adds, is by no means generated by such motion. *Elem. Phys. tom. 2, cap. 1.* Heat, in the hot body, he says, is an agitation of the parts of the body, made by means of the fire contained in it; by such agitation a motion is produced in our bodies, which excites the idea of Heat in our minds: so that Heat, in respect of us, is nothing but that idea, and in the hot body nothing but motion. If such motion expel

the



the fire in right lines, it may give us the idea of light; if in a various and irregular motion, only of Heat.

Lemery, the younger, agrees with these two authors, in asserting this absolute and ingenerable nature of fire; but he extends it farther. Not contented with confining it as an element to bodies, he endeavours to shew, that it is equably diffused through all space; that it is present in all places, even in the void spaces between the bodies, as well as in the insensible interstices between their parts. And this last sentiment falls in with that of Boerhaave above delivered. *Mem. de l'Acad. an. 1713.*

Philosophers have lately distinguished Heat into Absolute, and Sensible. By Absolute Heat, or fire, they mean that power or element which, when it is in a certain degree, excites in animals the sensation of Heat; and by Sensible Heat, the same power considered in its relation to the effects which it produces: thus, two bodies are said to have equal quantities of sensible Heat, when they produce equal effects upon the mercury in the thermometer; but as bodies of different kinds have different capacities for containing Heat, the absolute Heat in such bodies will be different, though the sensible Heat be the same. Thus, a pound of water and a pound of antimony, of the same temperature, have equal sensible Heat; but the former contains a much greater quantity of absolute Heat than the latter.

M. De Luc has evinced, by a variety of experiments, that the expansions of mercury between the freezing and boiling points of water, correspond precisely to the quantities of absolute Heat applied, and that its contractions are proportionable to the diminution of this element within these limits. And from hence it may be inferred, that if the mercury were to retain its fluid form, its contractions would be proportionable to the decrements of the absolute Heat, though the diminution were continued to the point of total privation. But the comparative quantities of absolute Heat, which are communicated to different bodies, or separated from them, cannot be determined in a direct manner by the thermometer.

Some philosophers have apprehended that the quantities of absolute Heat in bodies, are in proportion to their densities. While others, as Boerhaave, imagined that Heat is equally diffused through all bodies, the densest as well as the rarest, and therefore that the quantities of Heat in bodies are in proportion to their bulk or magnitude: and, at his desire, Fahrenheit attempted to determine the fact by experiment. For this purpose, he took equal quantities of the same fluid, and gave them different degrees of Heat, then upon mixing them intimately together, he found that the temperature of the mixture was a just medium, or arithmetical mean, between the two. But if this experiment be made with water and mercury, in the same circumstances, viz in equal bulks, the result will be different, as the temperature of the mixture will not be a mean between the two, but always nearer to that of the water than to the quicksilver; so that, when the water is the hotter, the temperature of the mixture is above the mean, and below it when the water is the colder. And from experiments of this kind it has been inferred, that the comparative quantities of the absolute Heats of these fluids, are reciprocally proportional to the changes which

are produced in their sensible Heats, when they are mixed together at different temperatures: and this fact has been publicly taught, for several years, by Dr. Black, and Dr. Irvine, in the universities of Edinburgh and Glasgow. This rule however does not apply to those substances which, in mixture, excite sensible Heat by chemical action.

From the experiments and reasoning employed by Dr. Crawford, it more fully appears, that the quantities of absolute Heat in different bodies, are not as their densities; or that equal weights of heterogeneous substances, as air and water, having the same temperature, may contain unequal quantities of absolute Heat: he also shews, that if phlogiston be added to a body, a quantity of the absolute Heat of that body will be extricated; and if the phlogiston be separated again, an equal quantity of Heat will be absorbed. So that Heat and phlogiston appear to be two opposite principles in nature. But this ingenious writer has not presumed absolutely to decide the question that has been long agitated, whether Heat be a substance or a quality.— He inclines to the former opinion however, and observes, that if we adopt the opinion, that Heat is a distinct substance, or an element sui generis, the phenomena will be found to admit of a simple and obvious interpretation, and to be perfectly agreeable to the analogy of nature. See Crawford's Experiments and Observations on Animal Heat and the Inflammation of Combustible Bodies.

*Animal HEAT.* The Heat of animals is very various, both according to the variety of their kinds, and the difference of the seasons: accordingly, zoologists have divided them into hot and cold blooded, reckoning those to be hot that are near or above our own temperature, and all others cold whose Heat is below ours, and consequently affect us with the sense of cold; thus making the human species a medium between the hot and cold blooded animals, or at least the lowest order of the hot blooded.

The Heat of the human body, in its natural state, according to Dr. Boerhaave, is such as to raise the mercury in the thermometer to  $92^{\circ}$  or at most to  $94^{\circ}$ ; and Dr. Pitcairn makes the heat of the human skin the same. Indeed it is evident that different parts of the human body, and its different states, as well as the different seasons, will make it shew of different temperatures. Thus, by various experiments at different times, the Heat of the human body is made various by the following authors:

Boerhaave and Pitcairn	-	-	$92^{\circ}$
Amontons	-	-	$91, 92$ ; or $93$
Sir Isaac Newton	-	-	$95\frac{1}{2}$
Fahrenheit and Musschenbroek, the blood,			$96$
Dr. Martine, the skin	-	-	$97$ or $98$
————, the urine	-	-	$99$
Dr. Hales, the skin	-	-	$97$
————, the urine	-	-	$103$
Mr. John Hunter, under his tongue,			$97$
————, in his rectum	-	-	$98\frac{1}{2}$
————, his urethra at 1 inch,			$92$
at 2 inches,			$93$
at 4 inches,			$94$
the ball of the thermom. at the bulb of the urethra			$97$



For the powers of animals to bear various degrees of Heat, see the *Philos. Trans.* vol. 65, 68, &c.

There is hardly any subject of philosophical investigation that has afforded a greater variety of hypotheses, conjectures, and experiments, than the cause of animal Heat. The first opinion which has very generally obtained, is, that the Heat of animal bodies is owing to the attrition between the arteries and the blood. All the observations and reasoning brought in favour of this opinion however, only shew that the Heat and the motion of the arteries are generally proportional to each other; without shewing which is the cause, and which the effect; or indeed that either is the cause or effect of the other, since both may be the effects of some other cause.

Dr. Douglas, in his *Essay on the Generation of Heat* in animals, ascribes it solely to the friction of the globules of blood in their circulation through the capillary vessels.

Another opinion is, that the lungs are the fountain of Heat in the human body: and this opinion is supported by much the same sort of arguments as the former, and seemingly to little better purpose.

A third opinion is, that the cause of animal Heat is owing to the action of the solid parts upon one another. And as the heart and arteries move most, it has been thought natural to expect that the Heat should be owing to this motion. But even this does not seem very plausible, from the following considerations: 1st, The moving parts, however we term them solid, are neither hard nor dry; which two conditions are absolutely requisite to make them fit to generate Heat by attrition. 2d, None of their motions are swift enough to promise Heat in this way. 3d, They have but little change of surface in their attritions. And 4thly, The moveable fibres have fat, mucilage, or liquors everyway surrounding them, to prevent their being destroyed, or heated by attrition.

A fourth cause assigned for the Heat of our bodies, is that process by which our aliment and fluids are perpetually undergoing some alteration. And this opinion is chiefly supported by Dr. Stevenson, in the *Edinburgh Medical Essays*, vol. 5, art. 77.

The late ingenious Dr. Franklin inclines to this opinion, when he says, that the fluid fire, as well as the fluid air, is attracted by plants in their growth, and becomes consolidated with the other materials of which they are formed, and makes a great part of their substance; that when they come to be digested, and to undergo a kind of fermentation in the vessels, part of the fire, as well as part of the air, recovers its fluid active state again, and diffuses itself on the body digesting and separating it; &c. *Exper. and Obs. on Electricity*, p. 346.

Dr. Mortimer thinks the Heat of animals explicable from the phosphorus and air they contain. Phosphorus exists, at least in a dormant state, in animal fluids; and it is also known that they all contain air; it is therefore only necessary to bring the phosphoreal and aerial particles into contact, and Heat must of consequence be generated; and were it not for the quantity of aqueous humours in animals, fatal accensions would frequently happen. *Philos. Trans.* number 476.

Dr. Black supposes, that animal Heat is generated

altogether in the lungs, by the action of the air on the principle of inflammability, and is thence diffused over the rest of the body by means of the circulation. But Dr. Leslie urges several arguments against this hypothesis, tending to shew that it is repugnant to the known laws of the animal machine; and he advances another hypothesis instead of it, viz, that the subtle principle by chemists termed phlogiston, which enters into the composition of natural bodies, is in consequence of the action of the vascular system gradually evolved through every part of the animal machine, and that during this evolution Heat is generated. This opinion, he candidly acknowledges, was first delivered by Dr. Duncan of Edinburgh; and that something similar to it is to be found in Dr. Franklin's works, and in a paper of Dr. Mortimer's in the *Philos. Trans.*

The last hypothesis we shall mention, is the very plausible one of Dr. Crawford, lately published in his *Experiments and Observations on Animal Heat*. This ingenious gentleman has inferred, from a variety of experiments, that Heat and phlogiston, so far from being connected, as most philosophers have imagined, act in some measure in opposition to each other. By the action of Heat on bodies, the force of their attraction of phlogiston is diminished, and by the action of phlogiston, a part of their absolute Heat is expelled. He has also demonstrated, that atmospherical air contains a greater quantity of absolute Heat than the air which is expired from the lungs of animals: he makes the proportion of the absolute Heat of atmospherical air to that of fixed air, as 67 to 1; and the Heat of dephlogisticated air to that of atmospherical air as 4.6 to 1; and observing that Dr. Priestley has proved, that the power of this dephlogisticated air in supporting animal life, is 5 times as great as that of atmospherical air, he concludes that the quantity of absolute Heat contained in any kind of air fit for respiration, is very nearly in proportion to its purity, or to its power of supporting animal life; and since the air exhaled by respiration, is found to contain only the 67th part of the Heat which was contained in the atmospherical air, previous to inspiration, it is very reasonably inferred, that the latter must necessarily deposit a very great proportion of its absolute Heat in the lungs. Dr. Crawford has also shewn, that the blood which passes from the lungs to the heart, by the pulmonary vein, contains more absolute Heat than that which passes from the heart to the lungs, by the pulmonary artery; the absolute Heat of florid arterial blood being to that of venous blood, as  $11\frac{1}{2}$  to 10: therefore, since the blood which is returned by the pulmonary vein to the heart has the quantity of its absolute Heat increased, it must have acquired this Heat in its passage through the lungs; so that in the process of respiration a quantity of absolute Heat is separated from the air, and absorbed by the blood. Dr. Priestley has also proved, that in respiration, phlogiston is separated from the blood, and combined with air.

This theory however has been contested and disputed, and, it has been said, Dr. Crawford's experiments repeated, with contrary results. Though no regular and systematical theory has yet been formed in its stead.

*HEAT of Combustible and Inflammable Bodies.* Dr. Crawford's theory with respect to the inflammation of com-



combustible bodies, is founded on the same principles as his doctrine concerning the Heat of animals. According to him, the Heat which is produced by combustion, is derived from the air, and not from the inflammable body. Inflammable bodies, he says, abound with phlogiston, and contain little absolute Heat: the atmosphere, on the contrary, abounds with absolute Heat, and contains little phlogiston. In the process of inflammation, the phlogiston is separated from the inflammable body, and combined with the air; the air is phlogistified, and gives off a great proportion of its absolute Heat, which, when extricated suddenly, bursts forth into flame, and produces an intense degree of sensible Heat. And since it appears by calculation, that the Heat produced by converting atmospherical into fixed air, is such, if it were not dissipated, as would be sufficient to raise the air so changed, to more than 12 times the Heat of red-hot iron, it follows, that in the process of inflammation a very great quantity of Heat is derived from the air. But, on the contrary, no part of the Heat can be derived from the combustible body; because this body, during the inflammation, being deprived of its phlogiston, undergoes a change similar to that of the blood by the process of respiration, in consequence of which its capacity of containing Heat is increased; and therefore it will not give off any part of its absolute Heat, but, like the blood in its passage through the lungs, it will absorb Heat.

A similar theory of Heat has lately been published by Mr. Elliot. See his Philosophical Observations on the senses of Vision and Hearing; to which is added an Essay on Combustion and Animal Heat. 8vo, 1780.

HEAT, in *Geography*, is that which relates to the earth. There is a great variety in the Heat of different places and seasons. Naturalists have commonly laid it down, that the nearer any place is to the centre of the earth, the hotter it is found; but this does not hold strictly true. And if it were, the effect might be otherwise accounted for, and more satisfactorily, than from their imagined central fire.

Mr. Boyle, who had been at the bottom of some mines himself, with more probability suspects that this degree of Heat, at least in some of them, may arise from the peculiar nature of the minerals there produced. And he instances a mineral of the vitriolic kind, dug up in large quantities, in several parts of England, which, by the bare effusion of common water, will grow so hot as almost to take fire. To which may be added, that such places, in the bowels of the earth, usually feel hot, from the confined and stagnant state of the air in them, in which the heat is retained, through the want of a current or change of air to carry the Heat off.

On the other hand, on ascending high mountains, the air grows more and more cold and piercing. Thus, the tops of the Pike of Teneriffe, the Alps, and several other mountains, even in the most sultry countries, are found always invested with snow and ice, which the Heat is never sufficient to thaw. In some of the mountains of Peru there is no such thing as running water, but all ice: plants vegetate a little about the bottom of the mountains, but near the top no vegetable can live, for the intenseness of the cold. This effect is attributed to

the thinness of the air, and the little surface of the earth there is to reflect the rays, as well as the great distance of the general surface of the earth which reflects the rays back into the atmosphere.

As to the diversity in the Heat of different climes and seasons, it arises from the different angles under which the sun's rays strike upon the surface of the earth. In the *Philos. Trans. Abr.* vol. 2, p. 165, Dr. Halley has given a computation of this Heat, on the principle, that the simple action of the sun's rays, like other impulses or strokes, is more or less forcible, according to the sines of the angles of incidence, or to the sines of the sun's altitudes, at different times or places.

Hence it follows, that, the time of continuance, or the sun's shining on any place, being taken for a basis, and the sines of the sun's altitudes perpendicularly erected upon it, and a curve line drawn through the extremities of those perpendiculars, the area thus comprehended will be proportional to the collection of all the Heat of the sun's beams in that space of time.

Hence it will likewise follow, that at the pole, the collection of all the Heat of a tropical day, is proportional to the rectangle or product of the sine of  $23\frac{1}{2}$  degrees into 24 hours, or the circumference of a circle, or as  $\frac{8}{15}$  into 12 hours, the sine of  $23\frac{1}{2}$  degrees being nearly  $\frac{4}{15}$  of radius. Or the polar Heat will be equal to that of the sun continuing 12 hours above the horizon at 53 degrees height; and the sun is not 5 hours more elevated than this under the equinoctial.

But as it is the nature of Heat to remain in the subject, after the luminary is removed, and particularly in the air, under the equinoctial the 12 hours absence of the sun abates but little from the effect of his Heat in the day; but under the pole, the long absence of the sun for 6 months has so chilled the air, that it is in a manner frozen, and after the sun has risen upon the pole again, it is long before his beams can make any impression, being obstructed by thick clouds and fogs.

From the foregoing principle Dr. Halley computes the following table, exhibiting the Heat to every 10th degree of latitude, for the equinoctial and tropical sun, and from which an estimate may easily be made for the intermediate degrees.

Lat.	Sign that the Sun is in.		
	♈ or ♎	♊	♋
0	20000	18341	18341
10	19696	20290	15834
20	18794	21737	13166
30	17321	22651	10124
40	15321	23048	6944
50	12855	22991	3798
60	10000	22773	1075
70	6840	23543	0
80	3473	24673	0
90	0	25055	0

From the same principles, and table, also are deduced the following corollaries, viz,

1, That



1, That the equatorial Heat, when the sun becomes vertical, is as twice the square of the radius.—2, That at the equator the Heat is as the sine of the sun's declination.—3, That in the frigid zones, when the sun sets not, the Heat is as the circumference of a circle into the sine of the altitude at 6: And consequently that in the same latitude, these aggregates of Heat are as the sines of the sun's declination; and at the same declination of the sun, they are as the sines of the latitudes; and generally they are as the sines of the latitudes into the sines of declination.—4, That the equatorial day's Heat is everywhere as the cosine of the latitude.—5, In all places where the sun sets, the difference between the summer and winter Heats, when the declinations are contrary, is equal to a circle into the sine of the altitude at 6, in the summer parallel; and consequently those differences are as the rectangles of the sines of the latitude and declination.—6, The tropical sun has the least force of any at the equator; and at the pole it is greatest of all.

Many objections have been urged against this theory of Dr. Halley. Some have objected, that the effect of the sun's Heat is not in the simple, but in the duplicate ratio of the sines of the angles of incidence; like the law of the impulse of fluids. And indeed, the quantity of the sun's direct rays received at any place, being evidently as the sine of the angle of incidence, or of the sun's altitude, *if* the Heat be also proportional to the force with which a ray strikes, like the mechanical action or impulse of any body, then it will follow that the Heat must be in the compound ratio of both, that is, as the square of the sine of the sun's altitude. But this last principle is here only assumed gratis, as we do not know a priori that the Heat is proportional to the force of a striking body; and it is only experiment that can determine this point.

It is certain that Heat communicated by the sun to bodies on the earth, depends also much upon other circumstances beside the direct force of his rays. These must be modified by our atmosphere, and variously reflected and combined by the action of the earth's surface itself, to produce any remarkable effects of Heat. So that if it were not for these additional circumstances, it is probable the naked Heat of the sun would not be very sensible.

Dr. Halley himself was well apprised, that many other circumstances, besides the direct force of the sun's rays, contributed to augment or diminish the effect of this, and the Heat resulting from it, in different climates; and therefore no calculation, formed on the preceding theory, can be supposed to correspond exactly with observation and experiment. It has also been objected that, according to the foregoing theory, the greatest Heat in the same place should be at the summer solstice, and the most extreme cold at the winter solstice; which is contrary to experience. To this objection it may be replied, that Heat is not produced in bodies by the sun instantaneously, nor do the effects of his Heat cease immediately when his rays are withdrawn; and therefore those parts which are once heated, retain the Heat for some time; which, with the additional Heat daily imparted, makes it continue to increase, though the sun declines from us: and this is the reason why July is hotter than June, although the sun has

withdrawn from the summer tropic: as we also find it is generally hotter at one, two, or three in the afternoon, when the sun has declined towards the west, than at noon, when he is on the meridian. As long as the heating particles, which are constantly received, are more numerous than those which fly away or lose their force, the Heat of bodies must continually increase. So, after the sun has left the tropic, the number of particles, which Heat our atmosphere and earth, constantly increases, because we receive more in the day than we lose at night, and therefore our Heat must also increase. But as the days decrease again, and the action of the sun becomes weaker, more particles will fly off in the night time than are received in the day, by which means the earth and air will gradually cool. Farther, those places which are well cooled, require time to be heated again; and therefore January is mostly colder than December, although the sun has withdrawn from the winter tropic, and begun to emit his rays more perpendicularly upon us.

But the chief cause of the difference between the Heat of summer and winter is, that in summer the rays fall more perpendicularly, and pass through a less dense part of the atmosphere; and therefore with greater force, or at least in greater number in the same place: and besides, by their long continuance, a much greater degree of Heat is imparted by day than can fly off by night.

For the calculations and opinions of several other philosophers on this head, see Keill's Astron. Lect. 8; Ferguson's Astron. chap. 10; Long's Astron. § 777; Memo. Acad. Scienc. 1719.

As to the temperature or Heat of our atmosphere, it may be observed that the mercury seldom falls under  $16^{\circ}$  in Fahrenheit's thermometer; but we are apt to reckon it very cold at  $24^{\circ}$ , and it continues coldish to  $40^{\circ}$  and a little above. However, such colds have been often known as bring it down to  $0^{\circ}$ , the beginning of the scale, or nearly the cold produced by a mixture of snow and salt, often near it, and in some places below it. Thus, the degree of the thermometer has been observed at various times and places as follows:

Places	Latit.	Year	Thermom.
Pennsylvania	$40^{\circ} 0'$	1732	$5^{\circ}$
Paris	$48 50$	1709 & 1710	8
Leyden	$52 10$	1729	5
Utrecht	$52 8$	—	4
London	$51 31$	1709 & 1710	0
Copenhagen	$55 43$	1709	0
Upsal	$59 56$	1732	-1
Petersburg	$59 56$	—	-28
Torneo	$65 51$	1736-7	-33
Hudson's Bay	$52 24$	1775	-37

The middle temperature of our atmosphere is about  $48^{\circ}$ , being nearly a medium of all the seasons. The French make it somewhat higher, reckoning it equal to the cave of their royal observatory, or  $53^{\circ}$ . In cold countries, the air is found agreeable enough to the inhabitants while it is between  $40$  and  $50^{\circ}$ . In our climate we are best pleased with the heat of the air from  $50$  to  $60^{\circ}$ ; while in the hot countries the air is generally at a medium about  $70^{\circ}$ . With us, the air is not reckoned



reckoned warm till it arrives at about  $64^{\circ}$ , and it is very warm and sultry at  $80^{\circ}$ . It is to be noted that the foregoing observations are to be understood of the state of the air in the shade; for as to the Heat of bodies acted upon by the direct rays of the sun, it is much greater: thus, Dr. Martine found dry earth heated to above  $120^{\circ}$ ; but Dr. Hales found a very hot sun-shine Heat in 1727 to be about  $140^{\circ}$ ; and Muffchenbroek once observed it so high as  $150^{\circ}$ ; but at Montpelier the sun was so very hot, on one day in the year 1705, as to raise M. Amontons's thermometer to the mark of boiling water itself, which is our  $212^{\circ}$ .

It appears from the register of the thermometer kept at London by Dr. Heberden for 9 years, viz, from the end of 1763 to the end of 1772, that the mean Heat at 8 in the morning was  $47^{\circ}4$ ; and by another register kept at Hawkhill, near Edinburgh, that the mean Heat in that place, during the same period of time, was  $46^{\circ}$ . Also by registers kept in London and at Hawkhill, for the three years 1772, 1773, 1774, it appears, that the mean Heat of these three years in London, at 8 in the morning, was  $48^{\circ}5$ , and at 2 in the afternoon  $56^{\circ}$ , but the mean of both morning and afternoon  $52^{\circ}2$ ; while the mean Heat at Hawkhill for the same time,

at 8 in the morning was	$45^{\circ}4$
and at 2 in the afternoon	$50^{\circ}1$
and the mean of both	$47^{\circ}7$

The mean Heat of springs near Edinburgh seems to be  $47^{\circ}$ , and at London  $51^{\circ}$ . *Philos. Transf.* vol. 65, art. 44.

Lastly, from the meteorological journals of the Royal Society, published in the *Philos. Transf.* it appears that the mean heights of the thermometer, for the whole years, kept without and within the house, are as below:

	Therm. Without	Therm. Within
For 1775	$51^{\circ}5$	$52^{\circ}7$
1776	$51^{\circ}1$	$52^{\circ}9$
1777	$51^{\circ}0$	$53^{\circ}0$
1778	$52^{\circ}0$	$53^{\circ}1$
mean of all	$51^{\circ}4$	$52^{\circ}9$

HEAVEN, an azure transparent orb investing our earth, where the celestial bodies perform their motions. It is of divers denominations, as the highest or empyrean Heaven, the ethereal or starry Heaven, the planetary Heaven, &c.

Formerly the Heavens were considered as solid substances, or else as spaces full of solid matter; but Newton has abundantly shewn that the Heavens are void of almost all resistance, and consequently of almost all matter: this he proves from the phenomena of the celestial bodies; from the planets persisting in their motions, without any sensible diminution of their velocity; and the comets freely passing in all directions towards all parts of the Heavens.

Heaven, taken in this general sense, or the whole expanse between our earth and the remotest regions of the fixed stars, may be divided into two very unequal parts, according to the matter occupying them; viz, the atmosphere or aerial Heaven, possessed by air; and the ethereal Heaven, possessed by a thin and unresisting medium, called ether.

VOL. I.

HEAVEN is more particularly used, in Astronomy, for an orb, or circular region, of the ethereal Heaven.

The ancient astronomers assumed as many different Heavens as they observed different celestial motions. All these they made solid, thinking they could not otherwise sustain the bodies fixed in them; and of a spherical form, as being the most proper for motion. Thus they had seven Heavens for the seven planets; viz, the Heavens of the Moon, Mercury, Venus, the Sun, Mars, Jupiter, and Saturn. The 8th was for the fixed stars, which they particularly called the firmament. Ptolemy added a 9th Heaven, which he called the primum mobile. After him two crystalline Heavens were added by king Alphonsus, &c, to account for some irregularities in the motions of the other Heavens. And lastly an empyrean Heaven was drawn over the whole for the residence of the Deity; which made the number 12.

But others admitted many other Heavens, according as their different views and hypotheses required. Eudoxus supposed 23, Calippus 30, Regiomontanus 33, Aristotle 47, and Fracastor no less than 70.

The astronomers however did not much concern themselves whether the Heavens they thus allowed, were real or not; provided they served a purpose in accounting for any of the celestial motions, and agreed with the phenomena.

HEAVINESS, the same as Gravity, which see.

Heavy bodies do not tend precisely to the very centre of the earth, except at the poles, and the equator, on account of the spheroidal figure of the earth; their direction being every where perpendicular to the surface of the spheroid.

HEGIRA, a term in Chronology, signifying the epoch, or account of time, used by the Mahomedans, who begin their computation from the day that Mahomet was forced to make his escape from the city of Mecca, which happened on Friday the 16th of July 622.

The years of the Hegira are lunar ones, consisting only of 354 days. Hence, to reduce these years to the Julian calendar, that is, to find what Julian year a given year of the Hegira answers to: reduce the year of the Hegira into days, by multiplying by 354, divide the product by 365 $\frac{1}{4}$ , and to the quotient add 622, the year the Hegira commenced.

HEIGHT, the third dimension of a body, considered with regard to its elevation above the ground.

HEIGHT, of a figure, the same as its altitude, being the perpendicular from its vertex to the base.

HEIGHT of the Pole, &c. See Altitude of the Pole, &c.

HELIACAL, something relating to the sun. Thus,

HELIACAL *Rising*, of a star or planet, is when it rises with, or at the same time, as the sun. And Heliacal setting, the same as the setting with the sun.

Or, a star rises Heliacally, when, after it has been in conjunction with the sun, and so invisible, it gets at such a distance from him as to be seen in the morning before the sun's rising. And it is said to set Heliacally, when it approaches so near the sun as to be hid by his beams. So that, in strictness, the Heliacal rising and setting are only an apparition and occultation.


HELICE *Major* and *Minor*; the same as Urfa Major and Minor.

4 G

HELICOID



**HELICOID Parabola**, or the *Parabolic Spiral*, is a curve arising from the supposition that the common or Apollonian parabola is bent or twisted, till the axis come into the periphery of a circle, the ordinates still retaining their places and perpendicular positions with respect to the circle, all these lines still remaining in the same plane. Thus, the axis of a parabola being bent into the circumference BCDM, and the ordinates CF, DG, &c, still perpendicular to it, then the parabola itself, passing through the extremities of the ordinates, is twisted into the curve BFG, &c, called the Helicoid, or Parabolic Spiral.



Hence all the ordinates CF, DG, &c, tend to the centre of the circle, being perpendicular to the circumference.

Also, the equation of the curve remains the same as when it was a parabola; viz, putting  $x$  = any circular absciss BC, and  $y$  = CF the corresponding ordinate, then is  $px = y^2$ , where  $p$  is the parameter of the parabola.

**HELIOCENTRIC Place of a Planet**, is the place in which a planet would appear to be when viewed from the sun; or the point of the ecliptic, in which a planet viewed from the sun would appear to be. And therefore the Heliocentric place coincides with the longitude of a planet viewed from the sun.

**HELIOCENTRIC Latitude of a Planet**, is the inclination of the line drawn between the centre of the sun and the centre of a planet, to the plane of the ecliptic. The greatest Heliocentric Latitude is equal to the inclination of the planet's orbit to the plane of the ecliptic.

**HELIOCOMETES**, Comet of the Sun, a phenomenon sometimes observed at the setting of the sun; thus denominated by Sturmius and Pylen, who had seen it, because it seems to make a comet of the sun, being a large tail, or column of light, fixed or hung to that luminary, and dragging after it, at its setting, like the tail of a comet.

**HELIOMETER**, or **ASTROMETER**; an instrument for measuring, with particular exactness, the diameters of the sun, moon, and stars.

This instrument was invented by M. Bouguer in 1747, and is a kind of telescope, consisting of two object glasses of equal focal distance, placed by the side of each other, so that the same eye-glass serves for both. The tube of this instrument is of a conical form, larger at the upper end, which receives the two object-glasses, than at the lower, which is furnished with an eye-glass and micrometer. By the construction of this instrument,

two distinct images of an object are formed in the focus of the eye-glass, whose distance, depending on that of the two object-glasses from one another, may be measured with great accuracy. Mem. Acad. Sci. 1748.

Mr. Servington Savery discovered a similar method of improving the micrometer, which was communicated to the Royal Society in 1743.

**HELIOSCOPE**, a kind of telescope peculiarly adapted for viewing and observing the sun without hurting the eye.

There are various kinds of this instrument, usually made by employing coloured glass for the object or eye-glass, or both; and sometimes only using an eye-glass blacked by holding it over the smoke or flame of a lamp or candle, which is Huygens's way.—See Dr. Hooke's treatise on Helioscopes.

**HELIOSTATA**, an instrument invented by Dr. Gravesande, and so called from its property of fixing the sun-beam in one position, viz, in a horizontal direction across the dark chamber while it is used. See Gravesande's *Physices Element. Mathematica*, tom. 2, p. 715 ed. 3tia 1742, for an account of the principles, construction and use of this instrument.

**HELISPHERICAL Line**, is the Rhumb-line in Navigation; being so called, because on the globe it winds round the pole helically or spirally, coming still nearer and nearer to it.

**HELIX**, a Spiral line. See **SPIRAL**.

**HEMISPHERE**, the half of a sphere or globe, when divided in two by a plane passing through its centre.

Hemisphere is also used for a map or projection of half the terrestrial globe, or of half the celestial sphere, on a plane; being more frequently called a planisphere.

The centre of gravity of a Hemisphere, is five-eighths of the radius distant from the vertex.

A glass Hemisphere unites the parallel rays at the distance of four-thirds of a diameter from the pole of the glass.

**HEMITONE**, in Music, a half note.


**HENDECAGON**, a figure of eleven sides, or the Endecagon; which see.

**HENIOCHAS**, or **HENIOCHUS**, a northern constellation, the same as Auriga, which see.

**HEPTAGON**, in Geometry, a figure of seven sides and seven angles.—When those sides and angles are all equal, the Heptagon is said to be regular, otherwise it is irregular.

In a regular Heptagon, the angle C at the centre is  $= 51^{\circ}\frac{2}{7}$ , the angle DAB of the polygon is  $= 128^{\circ}\frac{4}{7}$ , and its half CAB  $= 64^{\circ}\frac{2}{7}$ . Also the area is = the square of the side  $AB^2 \times 3.6339124$  or  $= AB^2 \times \frac{7}{4} t$ , where  $t$  is the tangent of the angle CAB of  $64^{\circ}\frac{2}{7}$  to the radius 1; or  $t$  is the root of the equation;

$$t^{12} - 26t^{10} + 143t^8 - 245t^6 + 143t^4 - 26t^2 + 1 = 0; \text{ or } t = \frac{\sqrt{1+x}}{1-x} = \frac{1 + \sqrt{1-y^2}}{y} = \frac{y}{1 - \sqrt{1-y^2}}, \text{ where}$$





where the value of  $x$  and  $y$  are the roots of the equations

$$x^6 - \frac{5}{4}x^4 + \frac{3}{8}x^2 - \frac{1}{48} = 0,$$

$$y^6 = \frac{7}{4}y^4 + \frac{7}{8}y^2 - \frac{7}{64} = 0.$$

See my *Mensuration*, p. 21, 114, and 116, 2d edition.

**HEPTAGON**, in Fortification, a place fortified or strengthened with seven bastions for its defence.

**HEPTAGONAL Numbers**, are a kind of polygonal numbers in which the difference of the terms of the corresponding arithmetical progression is 5. Thus,

Arithmetics, 1, 6, 11, 16, 21, 26, &c.

Heptagonals, 1, 7, 18, 34, 55, 81, &c.

where the Heptagonals are formed by adding continually the terms of the arithmetics, above them, whose common difference is 5.

One property, among many others, of these Heptagonal numbers is, that if any one of them be multiplied by 40, and to the product add 9, the sum will be a square number.

Thus  $1 \times 40 + 9 = 49 = 7^2$ ;

and  $7 \times 40 + 9 = 289 = 17^2$ ;

and  $18 \times 40 + 9 = 729 = 27^2$ ;

and  $34 \times 40 + 9 = 1369 = 37^2$ ; &c.

Where it is remarkable that the series of squares so formed is  $7^2, 17^2, 27^2, 37^2$ , &c, the common difference of whose roots is 10, the double of the common difference of the arithmetical series from which the Heptagonals are formed.—See **POLYGONALS**.

**HEPTANGULAR Figure**, in Geometry, is one that has seven angles; and therefore also seven sides.

**HERCULES**, in Astronomy, a constellation of the northern hemisphere, and one of the 48 old constellations mentioned by ancient writers.

It is not known by what name it was distinguished by the Egyptians and others before the Greeks. These latter, perhaps not knowing its real name, first called it simply the kneeling man, because he is drawn in that posture; but they afterwards successively ascribed it to, and called it by the names of, Cetheus, Theseus, and lastly Hercules, which it still retains.

The stars in this constellation, in Ptolemy's catalogue, are 29; in Tycho's 28; and in the Britannic catalogue, 113.

**HERISSON**, in Fortification, a beam armed with iron spikes, having their points turned outward. It is supported in the middle by a stake, having a pivot on which it turns; and serves as a barrier to block up a passage.—Herissons are frequently placed before gates, especially the posterns of a town or fortrefs, to secure those passages which must of necessity be often opened.

**HERMANN (JAMES)**, a learned mathematician of the Academy of Berlin, and member of the Academy of Sciences at Paris, was born at Basil in 1678. He was a great traveller; and for 6 years was professor of mathematics at Padua. He afterwards went to Russia, being invited thither by the Czar in 1724, as well as his compatriot Daniel Bernoulli. On his return to his native country, he was appointed professor of morality and natural law at Basil; where he died in 1733, at 55 years of age.

He wrote several mathematical and philosophical pieces, in the Memoirs of different Academies, and elsewhere; but his principal work, is the *Phoronomia*, or two Books on the Forces and Motions of both Solid and Fluid bodies; 4to, 1716: a very learned work on the new mathematical physics.

**HERMETIC**, or **HERMETICAL Art**, a name given to chemistry, on a supposition that Hermes Trismegistus was its inventor.

**HERMETICAL Philosophy**, is that which undertakes to solve and explain all the phenomena of nature, from the three chemical principles, salt, sulphur, and mercury.—A considerable addition was made to the ancient Hermetical Philosophy, by the modern doctrine of alkali and acid.

**HERMETICAL Seal**, or *Hermetical Sealing*, a manner of stopping or closing glass vessels, for chemical and other operations, so very closely, that no substance can possibly exhale or escape. This is usually done by heating the neck of the vessel in the flame of a lamp, with a blow-pipe, till it be ready to melt, and then with a pair of hot pincers twisting it close together.

**HERSCHEL**, the name by which the French, and most other European nations, call the new planet, discovered by Dr. Herschel in the year 1781. Its mark or character is  $\text{H}$ . The Italians call it Ouranos, or Urania, but the English, the *GEORGIAN Planet*, which see.

**HERSE**, in Fortification, a lattice or portcullice, in the form of a harrow, beset with iron spikes, to block up a gate way, &c.

**HERSILLON**, or little Herse, in Fortification, is a plank armed with iron spikes, for the same use as the Herse, and also to impede the march of the infantry or cavalry.

**HESSE (WILLIAM Prince of)**, rendered his name immortal by his encouragement of learning, by his studies, and by his observations, for many years, of the celestial bodies. For this purpose, he erected an observatory at Cassel, and furnished it with good instruments, well adapted to that design; calling also to his assistance two eminent artists, Christopher Rothmann and Juste Byrge. His observations, which are of a very curious nature, were published at Leyden, in the year 1618, by Willebrord Snell; and are in part mentioned by Tycho Brahe, as well in his epistles as in the 2d volume of his *Progymnasmata*; a signal example to all princely and heroic minds, to undertake the promoting the arts of peace, and advancing this truly noble and celestial science. This prince died in the year 1597.

**HETERODROMUS Vetus**, or *Lever*, in Mechanics, a lever in which the fulcrum, or point of suspension, is between the weight and the power; being the same as what is otherwise called a lever of the first kind.

**HETEROGENEAL**, the same as **HETEROGENEOUS**; which see.

**HETEROGENEOUS**, literally imports things of different natures, or something that consists of parts of different or dissimilar kinds; in opposition to Homogeneous. Thus,

**HETEROGENEOUS Bodies**, are such as have their parts of unequal density.

**HETEROGENEOUS Line**, is that which consists of parts or rays of different refrangibility, reflexivity, and colour.

**HETEROGENEOUS Numbers**, are mixed numbers, consisting of integers and fractions.

**HETEROGENEOUS Particles**, are such as are of different kinds, natures, and qualities; of which generally all bodies consist.

**HETEROGENEOUS Quantities**, in Mathematics, are those



those which cannot have proportion, or be compared together as to greater and less; being of such different kind and consideration, as that one of them taken any number of times, never equals or exceeds the other. As lines, surfaces, and solids in geometry.

**HETEROGENEOUS Surds**, are such as have different radical signs; as  $\sqrt{a}$  and  $\sqrt[3]{b^2}$ ; or  $\sqrt[5]{10}$  and  $\sqrt[7]{20}$ .

**HETEROSCI**, in Geography, are such inhabitants of the earth as have their shadows at noon projected always the same way with regard to themselves, or always contrary ways with respect to each other. Thus, all the inhabitants without the torrid zone are Heteroscii, with regard to themselves, since any one such inhabitant has his shadow at noon always the same way, viz, always north of him in north latitude, and always south of him in south latitude; or these two situations are Heteroscii to each other, having such shadows projected contrary ways at all times of the year.

**HEVELIUS (JOHN)**, a very celebrated astronomer, and a burgomaster of Dantzick, was born in that city in 1611. He studied mathematics under Peter Cruger, in which he made a wonderful progress. He afterwards spent several years on his travels through Holland, England, Germany, and France, for his improvement in the sciences. On his return, he constructed excellent telescopes himself, and began diligently to observe the heavens, an employment he closely followed during the course of a long life, which was terminated only in 1687, at 76 years of age. Hevelius was author of several notable discoveries in the heavens. He was the first that observed the phenomenon called the libration of the moon, and made several other important observations on the other planets. He also discovered several fixed stars, which he named the *firmament of Sobieski*, in honour of John the 3d, king of Poland. He framed a large catalogue of the stars, and collected multitudes of the unformed ones into new constellations of his own framing. His wife was also well skilled in astronomy, and made a part of the observations that were published by her husband. His principal publications are, his *Selenographia*, or an exact description of the moon; in which he has engraved all her phases, and remarkable parts, distinguished by names, and ascertained their respective bounds by the help of telescopes; containing also a delineation of the several visible spots, with the various motions, changes, and appearances, discovered by the telescope, as also in the sun and other planets, 1647.—In 1654, two epistles; one to the celebrated astronomer Riccioli, concerning the Libration of the Moon; and the other to Bulliald, on the Eclipses of both luminaries.—In 1656, a Dissertation *De Natura Saturni faciei*, &c.—In 1668, his *Cometographia*, representing the whole nature of comets, their situation, parallaxes, distances, diverse appearances, and surprising motions, with a history of all the comets from the beginning of the world down to the present time; being enriched with curious sculpture of his own execution: to which he added a treatise on the planet Mercury, seen in the sun at Dantzick, May 3, 1661; with the history of a new star appearing in the neck of Cetus, and another in the beak of Cygnus; besides an illustration of some astronomical discoveries of the late Mr. Horrox, in his treatise on Venus seen in the sun, Nov. 24, 1639; with a discourse of some curious Paraselena and Parhelia observed at Dantzick. He sent copies of this work to

several members of the Royal Society at London, and among them to Mr. Hooke, in return for which, this gentleman sent to Hevelius a description of the Dioptric Telescope, with an account of the manner of using it; and recommending it to him, as much preferable to telescopes with plain sights. This gave rise to a dispute between them, viz, “whether distances and altitudes could be taken with plain sights any nearer than to a minute.” Hooke asserted that they could not; but that, with an instrument of a span radius, by the help of a telescope, they might be determined to the exactness of a second. Hevelius, on the other hand, insisted that, by the advantage of a good eye and long practice, he was able with his instruments to come up even to that exactness; and, appealing to experience and facts, sent by way of challenge 8 distances, each between two different stars, to be examined by Hooke. Thus the affair rested for some time with outward decency, but not without some inward grudge between the parties.

In 1673, Hevelius published the first part of his *Machina Cœlestis*, as a specimen of the exactness both of his instruments and observations; and sent several copies as presents to his friends in England, but omitting Mr. Hooke. This, it is supposed, occasioned that gentleman to print, in 1674, *Animadversions on the First Part of the Machina Cœlestis*; in which he treated Hevelius with a very magisterial air, and threw out several unhand-some reflections, which were greatly resented; and the dispute grew afterwards to such a height, and became so notorious, that in 1679 Dr. Halley went, at the request of the Royal Society, to examine both the instruments and the observations made with them. Of both these, Halley gave a favourable account, in a letter to Hevelius; and Hooke managed the controversy so ill, that he was universally condemned, though the preference has since been given to telescopic sights. However, Hevelius could not be prevailed on to make use of them: whether he thought himself too experienced to be informed by a young astronomer, as he considered Hooke; or whether, having made so many observations with plain sights, he was unwilling to alter his method, lest he might bring their exactness into question; or whether, being by long practice accustomed to the use of them, and not thoroughly apprehending the use of the other, nor well understanding the difference, is uncertain. Besides Halley's letter, Hevelius received many others in his favour, which he took the opportunity of inserting among the astronomical observations in his *Annus Climactericus*, printed in 1685. In a long preface to this work, he speaks with more confidence and greater indignation than he had done before, and particularly exclaimed against Hooke's dogmatical and magisterial manner of assuming a kind of dictatorship over him. This revived the contest, and occasioned several learned men to engage in it. The book itself being sent to the Royal Society, at their request an account of it was given by Dr. Wallis; who, among other things, took notice, that “Hevelius's observations had been misrepresented, since it appeared from this book, that he could distinguish by plain sights to a small part of a minute.” About the same time Mr. Molyneux also wrote a letter to the society, in vindication of Hevelius, against Hooke's animadversions. To all which, Hooke drew up a letter in answer, which was read before the society, containing many qualifying and



and accommodating expressions, but still at least expressing the superiority of telescopic sights over plain ones, excellent as the observations were that had been made with these.

In 1679, Hevelius had published the second part of his *Machina Caelestis*; but the same year, while he was at a seat in the country, he had the misfortune to have his house at Dantzic burnt down. By this calamity it is said he sustained several thousand pounds damage; having not only his observatory and all his valuable instruments and astronomical apparatus destroyed, but also a great many copies of his *Machina Caelestis*, an accident which has made this second part very scarce, and consequently very dear.

In 1690, were published a description of the heavens, called, *Firmamentum Sobiescianum*, in honour of John the 3d, king of Poland, as above mentioned; and also *Prodromus Astronomiae, & Novae Tabulae Solares, una cum Catalogo Fixarum*; in which he lays down the necessary preliminaries for taking an exact catalogue of the stars.

But both these works were posthumous; for Hevelius died the 28th of January 1687, exactly 76 years of age, as above said, and universally admired and respected; abundant evidence of which appears in a collection of letters between him and many other persons, that was printed at Dantzic in 1683.

HEXACHORD, a certain interval or musical concord, usually called a sixth.

HEXAEDRON, or HEXAHEDRON, one of the five regular or Platonic bodies; being indeed the same as the cube; and is so called from its having 6 faces.—The square of the side or edge of a Hexahedron, is one-third of the square of the diameter of the circumscribing sphere: and hence the diameter of a sphere is to the side of its inscribed Hexahedron, as  $\sqrt{3}$  to 1.

In general, if A, B, and C be put to denote respectively the linear side, the surface, and the solidity of a Hexahedron or cube, also  $r$  the radius of the inscribed sphere, and R the radius of the circumscribed one; then we have these general equations or relations:

1.  $A = 2r = \frac{2}{3}R\sqrt{3} = \sqrt{\frac{1}{3}}B = \sqrt[3]{C}$ .
2.  $B = 24r^2 = 8R^2 = 6A^2 = 6\sqrt[3]{C^2}$ .
3.  $C = 8r^3 = \frac{8}{9}R^3\sqrt{3} = A^3 = \frac{1}{8}B\sqrt{\frac{1}{3}}B$ .
4.  $R = r\sqrt{3} = \frac{1}{2}A\sqrt{3} = \frac{1}{2}\sqrt{\frac{1}{3}}B = \frac{1}{2}\sqrt{3} \times \sqrt[3]{C}$ .
5.  $r = \frac{1}{3}R\sqrt{3} = \frac{1}{2}A = \frac{1}{2}\sqrt{\frac{1}{3}}B = \frac{1}{2}\sqrt[3]{C}$ .

HEXAGON, in Geometry, a figure of six sides, and consequently of as many angles. When these are equal, it is a regular Hexagon.—The angles of a Hexagon are each equal to  $120^\circ$ , and its sides are each equal to the radius of its circumscribing circle. Hence a regular Hexagon is inscribed in a circle, by setting the radius off 6 times upon the periphery. And hence also, to describe a regular Hexagon upon a given line, describe an equilateral triangle upon it, the vertex of which will be the centre of the circumscribing circle.

The side of a Hexagon being  $s$ , its area will be  $= 2.5980762s^2 = \frac{3}{2}s^2 \times \text{tang. } 60^\circ = \frac{3}{2}s^2\sqrt{3}$ .

HEXASTYLE, in the Ancient Architecture, a building with 6 columns in front.

HIERO's Crown, in Hydrostatics. The history of this crown, and of the important hydrostatical proposition which it gave occasion to, is as follows: Hiero,

king of Syracuse, having furnished a workman with a quantity of gold for making a crown, suspected that he had been cheated, by the workman using a greater alloy of silver than was necessary in making it; and he applied to Archimedes to discover the fraud, without defacing the crown.

This celebrated mathematician was led by chance to a method of detecting the imposture, and of determining precisely the quantities of gold and silver composing the crown: for he observed, when bathing in a tub of water, that the water ran over as his body entered it, and he presently concluded that the quantity so running over was equal to the bulk of his body that was immersed. He was so pleased with the discovery, that it is said he ran about naked crying out, *εὕρηκα, εὕρηκα, I have found it*; and some affirm that he offered a hetacomb to Jupiter for having inspired him with the thought.

On this principle he procured a ball or mass of gold, and another of silver, exactly of the same weight with the crown; considering, that, if the crown were of pure gold, it would be of equal bulk and expel an equal quantity of water as the golden ball; and if it were of silver, then it would be of equal bulk and expel an equal quantity of water with the ball of silver; but of intermediate quantity, if it consisted of a mixture of the two, gold and silver; which, upon trial, he found to be the case; and hence, by a comparison of the quantities of water displaced by the three masses, he discovered the exact portions of gold and silver in the crown.

Now, suppose, for example, that each of the three masses weighed 100 ounces; and that on immersing them severally in water, there were displaced 5 ounces of water by the golden ball, 9 ounces by the silver, and 6 ounces by the compound, or crown; that is, their respective or comparative bulks are as 5, 9, and 6, the sum of which is 20.

Then the method of operation is this:

From	9	6
Take	6	5
	—	—

rem. 3                      1, whose sum is 4.

Therefore  $4 : 100 :: 3 : 75$  oz. of gold,

and  $4 : 100 :: 1 : 25$  oz. of silver.

That is, the crown consisted of 75 ounces of gold, and 25 ounces of silver.

See Cotes Hydrof. Lect. p. 81; or Martin's Philos. Britan, vol. 1, p. 305, &c. See also SPECIFIC Gravity.

HIGH-Water, that state of the tides when they have flowed to their greatest height, or have ceased to flow or rise. At High-water the motion commonly ceases for a quarter or half an hour, before it begin to ebb again. The times of High-water of every day of the moon's age, is usually computed from that which is observed on the day of the full or change; viz, by taking  $\frac{4}{5}$ ths of the moon's age on any day of the month, and adding it to the time of High-water on the day of the full or change; then is the sum nearly equal to the time of High-water on the day of the month proposed. And as to the times of High-water, on the day of the full and change of the moon, at many different places; they have been observed as they are set down in the following table.

*Table*



TABLE of the Times of High-water on the Days of the New and Full Moons, at many different Places.

Names of Places.	Countries.	High-w.	Names of Places.	Countries.	High-w.
Aberdeen.	Scotland	0h 45 <sup>m</sup>	Dort	Holland	3h 0 <sup>m</sup>
Aldborough	England	9 45	Dover	England	11 30
Alderney I.	England	12 0	Downs	England	1 15
Amazons River	South America	6 0	Dublin	Ireland	9 15
Amsterdam	Holland	3 0	Dunbar	Scotland	2 30
Amsterdam I. of	South Seas	8 30	Dundee	Scotland	2 15
Andrew's St.	Scotland	2 15	Dungarvan	Ireland	4 30
Anholt I.	Denmark	0 0	Dungeness	England	9 45
Antwerp	Flanders	6 0	Dunkirk	France	0 0
Archangel	Russia	6 0	Dunnofe	I. of Wight	9 45
Arran I.	Ireland	11 0	Dusky Bay	N. Zealand	10 57
Ashley Riv.	Carolina	0 45	Easter Isle	Chili	2 0
Augustine St.	Florida	7 30	Edystone	English Channel	5 30
Bajador Ca.	Negroland	0 0	Elbe R.	Germany	0 0
Baltimore	Ireland	4 30	Embsen	Germany	0 0
Barfleur Ca.	France	7 30	Eftaples	France	11 0
Bayonne	France	3 30	Falmouth	England	5 30
Beachy-head	England	0 0	Flamborough H.	England	4 0
N. and S. Bear	Labradore	12 0	C. Florida	Florida	7 30
Belfast	Ireland	10 0	Flushing	Holland	0 45
Bellisle	France	3 30	N. Foreland	England	9 45
Bermudas I.	Bahama I.	7 0	Foulness	England	6 45
Berwick	England	2 30	Fowey	England	5 15
Blackney	England	6 0	Fayal Isl.	Azores	2 20
Blanco Cap.	Negroland	9 45	Garonne R.	France	3 0
Blavet	France	3 0	Gibraltar	Spain	0 0
Bordeaux	France	3 0	C. Good Hope	Caffers	3 0
Boulogne	France	10 30	Goree (Isle)	Atlantic Ocean	1 30
Bremen	Germany	6 0	Granville	France	7 0
Brest	France	3 45	Gravelines	Flanders	0 0
Bridlington B.	England	3 45	Gravesend	England	1 30
Brill	Holland	1 30	Groin	Spain	3 3
Bristol	England	6 45	Guernsey I.	English Channel	1 30
Buchaness	Scotland	3 0	Hague	Holland	8 15
Button's Isles	New Brit.	6 50	Halifax	Nova Scotia	7 30
Cadiz	Spain	4 30	Hamburgh	Germany	6 0
Caen	France	9 0	Hare Isle	Canada	3 30
Calais	France	11 30	Harlem	Holland	9 0
Canaria I.	Canaries	3 0	Hartlepool	England	3 0
C. Cantin	Barbary	0 0	Harwich	England	11 15
Cape Town	Caffers	2 30	Havre de Grace	France	9 0
Caskets	Guernsey	8 15	Holy Head	Wales	1 30
Cathness Po.	Scotland	9 0	Honfleur	France	9 0
Charles Town	Carolina	3 0	Hull	England	6 0
Q. Charlotte's S.	New Zealand	9 0	Humber R.	England	5 13
Cherbourg	France	7 30	St. John's	Newfoundland	6 0
Churchill R.	Hudson's Bay	7 20	St. Julian (Port)	Patagonia	4 45
Ca. Cleare	Ireland	4 30	Kentishnock	English coast	0 0
Concarneau	France	3 0	Kinfale	Ireland	5 15
Conquet	France	2 15	Land's End	England	7 30
Coquet Isle	England	3 0	Leith	Scotland	4 30
Corke	Ireland	6 30	Leostoff	England	9 45
C. Corfe	Guinea	3 30	Lisbon	Portugal	2 15
Cromer	England	7 0	Liverpool	England	11 15
Dartmouth	England	6 30	Lizard	England	7 30
St. David's H.	Wales	6 0	Loire (Riv.)	France	3 0
Dieppe	France	10 30	London	England	3 0



Names of Places.	Countries.	High-w.	Names of Places.	Countries.	High-w.
Lundy (Isle)	England	5 <sup>h</sup> 15 <sup>m</sup>	Sandwich	England	11 <sup>h</sup> 30 <sup>m</sup>
Madeira	Atl. Ocean	12 4	Scarborough H.	England	3 45
St. Maloes	France	6 0	Scilly Isles	England	3 45
Isle of Man	England	9 0	Senegal	Negroland	10 30
Margate	England	11 15	Severn, (Mouth.)	England	6 0
St. Mary's (Isle)	Scilly Isles	3 45	Sheerneys	England	0 0
Milford	Wales	5 15	Sierra Leona	Guinea	8 15
Mount's Bay	England	4 30	Shetland I.	Scotland	3 0
Nantes	France	3 0	Isle of Sky	Scotland	5 30
Naze	Norway	11 15	Spurn	England	5 15
Needles	England	10 15	Start Point	England	6 45
Newcastle	England	3 15	Stockton	England	5 15
Nieuport	Flanders	12 0	Sunderland	England	3 20
Nore	England	0 0	Tanna	Pacific Ocean	3 0
North Cape	Lapland	3 0	Teneriff	Canaries	3 0
Orfordness	England	9 45	Texel (Isle)	Holland	7 30
Orkneys	Scotland	3 0	Thames Mouth	England	1 30
Ostend	Flanders	12 0	Tinmouth	England	3 0
Placentia	Newfoundland	9 0	Torbay	England	5 15
Plymouth	England	6 0	St. Valery	France	10 30
Portland	England	8 15	Vannes	France	3 45
Porto Praya	Cape Verdes	11 0	Ushant	France	4 30
Portsmouth	England	11 15	Waterford	Ireland	6 30
Quebec	Canada	7 30	Wells	England	6 0
Rh�e (Isle)	France	3 0	Weymouth	England	7 20
Resolution (Bay)	Ohitahoo	2 30	Whitby	England	3 0
Robin Hood's B.	England	3 0	Isle of Wight	England	0 0
Rochefort	France	4 15	Winchelsea	England	0 45
Rochelle	France	3 45	Wintertones	England	9 0
Rochester	England	0 45	Yarmouth	England	9 45
Rotterdam	Holland	3 0	New York	America	3 0
Rouen	France	1 15	Youghall	Ireland	4 30
Rye	England	11 15	Zuric Sea	Holland	3 0

HIPS, in Architecture, are those pieces of timber placed at the corners of a roof. These are much longer than the rafters, because of their oblique position.

HIP means also the angle formed by two parts of the roof, when it rises outwards.

HIP-Roof, called also Italian Roof, is one in which two parts of the roof meet in an angle, rising outwards: the same angle being called a valley when it sinks inwards.

HIPPARCHUS, a celebrated astronomer among the ancients, was born at Nice in Bithynia, and flourished between the 154th and the 163d olympiads; that is, between 60 and 135 years before Christ; for in this space of time it is that his observations are dated. He is accounted the first, who from vague and scattered observations, reduced astronomy into a science, and prosecuted the study of it systematically. Pliny often mentions him, and always with great commendation. He was the first, he tells us, who attempted to count the number of the fixed stars; and his catalogue is preserved in Ptolemy's Almagest, where they are all noted according to their longitudes and apparent magnitudes. Pliny places him among those men of a sublime genius, who, by foretelling the eclipses, taught mankind, that they ought not to be frightened at these phenomena. Thales was the first among the Greeks, who could discover when there was to be an eclipse. Sulpitius Gallus among the Romans

began to succeed in this kind of prediction; and he gave an essay of his skill very seasonably, the day before a battle was fought. After these two, Hipparchus improved that science very much; making ephemerides, or catalogues of eclipses, for 600 years. He admires him for making a review of all the stars, acquainting us with their situations and magnitudes; for by these means, says he, posterity will be able to discover, not only whether they are born and die, but also whether they change their places, and whether they increase or decrease. He mentioned a new star which was produced in his days; and by its motion, at its first appearance, he began to doubt whether this did not frequently happen, and whether those stars, which we call fixed, do not likewise move. Hipparchus is also memorable for being the first who discovered the precession of the equinoxes, or a very slow apparent motion of the fixed stars from east to west, by which in a great number of years they will seem to have performed a complete revolution. He endeavoured also to reduce to rule the many discoveries he made, and invented new instruments, by which he marked their magnitudes and places in the heavens; so that by means of them it might be easily observed, not only whether they appear and disappear, but likewise whether they pass by one another, or move, and whether they increase or decrease.

The first observations he made, were in the isle of Rhodes;



Rhodes; whence he got the name Rhodius; but afterwards he cultivated this science in Bithynia and Alexandria only. One of his works is still extant, viz, his Commentary upon Aratus's Phenomena. He composed several other works; and upon the whole it is agreed, that astronomy is greatly indebted to him, for laying that rational and solid foundation, upon which all succeeding astronomers have since built their superstructure.

HIRCUS, in Astronomy, a fixed star of the first magnitude, the same with Capella.

HIRCUS is also used by some writers for a comet, encompassed as it were with a mane, seemingly rough and hairy.

HIRE (PHILIP DE LA), an eminent French mathematician and astronomer, was born at Paris in 1640. His father, who was painter to the king, intending him for the same occupation, taught him drawing and such branches of mathematics as relate to it: but died when the son was only 17 years of age. Three years after this, he travelled into Italy for improvement in that art, where he spent 4 years. He applied himself also to mathematics, which gradually engrossed all his attention. On his return to Paris, he continued his mathematical studies with great eagerness, and he afterwards published some works, which gained him so much reputation, that he was named a member of the Academy of Sciences in 1678.

The minister Colbert having formed a design for a better chart or map of France than any former ones, De la Hire was appointed, with Picard, to make the necessary observations for that purpose. This occupied him some years in several of the provinces; and, beside the main object of his peregrinations, he was not unmindful of other branches of knowledge, but philosophized upon every thing that occurred, and particularly upon the variations of the magnetic needle, upon refractions, and upon the height of mountains, as determined by the barometer.

In 1683, de la Hire was employed in continuing the meridian line, which Picard had begun in 1669. He continued it from Paris northward, while Cassini carried it on to the south: but Colbert dying the same year, the work was dropped before it was finished. De la Hire was next employed, with other members of the academy, in taking the necessary levels for the grand aqueducts, which Louis the 14th was about to make.

The great number of works published by our author, together with his continual employments, as professor of the Royal College and of the Academy of Architecture, give us some idea of the great labours he underwent. His days were always spent in study; his nights very often in astronomical observations; seldom seeking any other relief from his labours, than a change of one for another. In his manner, he had the exterior politeness, circumspection, and prudence of Italy; on which account he appeared too reserved in the eyes of his countrymen; though he was always esteemed as a very honest disinterested man. He died in 1718, at 78 years of age.

Of the numerous works which he published, the principal are, 1. *Traité de Mécanique*; 1665.—2. *Nouvelle Méthode en Géométrie pour les Sections des Surfaces Coniques & Cylindriques*; 1673, 4to.

—3. *De Cycloide*; 1677, 12mo.—4. *Nouveaux Elémens des Sections Coniques: les Lieux Géométriques: la Construction, ou Efection des Equations*; 1678, 12mo.—5. *La Gnomonique, &c*; 1682, 12mo.—6. *Traité du Nivellement de M. Picard, avec des additions*; 1684.—7. *Sections Conicæ in novem libros distributæ*; 1685, folio. This was considered as an original work, and gained the author great reputation all over Europe.—8. *Traité du Mouvement des Eaux, &c*; 1686.—9. *Tabulæ Astronomicæ*; 1687 and 1702, 4to.—10. *Ecole des Arpenteurs*; 1689.—11. *Veterum Mathematicorum Opera, Græcè & Latine, pleraque nunc primum edita*; 1693, folio. This edition had been begun by Thevenot; who dying, the care of finishing it was committed to de la Hire. It shews that our author's strong application to mathematical and astronomical studies had not hindered him from acquiring a very competent knowledge of the Greek tongue. Beside these, and other smaller works, there are a vast number of his pieces scattered up and down in Journals, and particularly in the *Memoirs of the Academy of Sciences*, viz, from 1666 till the year 1718.

HOBBS (THOMAS), a famous writer and philosopher, was born at Malmesbury in Wiltshire, in 1588, being the son of a clergyman of that place. He completed his studies at Oxford, and was afterwards governor to the eldest son of William Cavendish earl of Devonshire, with whom he travelled through France and Italy, applying himself closely to the study of polite literature. In 1626 his patron the earl of Devonshire died; and 1628 his son also; the same year Mr. Hobbes published his translation of Thucydides in English. He soon after went abroad a second time as governor to the son of Sir Gervase Clifton; but shortly after returned, to resume his concern for the hopes of the Devonshire family, to whom he had so early attached himself; the countess dowager having desired to put the young earl under his care, then about 13 years of age. This charge was very agreeable to Mr. Hobbes's inclinations, and he discharged the trust with great diligence and fidelity. In 1634 he accompanied his young pupil to Paris, where he employed his own vacant hours in the study of natural philosophy, frequently conversing with Father Merfenne, Gassendi, and other eminent philosophers there. From Paris he attended his pupil into Italy, where he became acquainted with the celebrated Galileo, who freely communicated his notions to him; and from hence he returned with his ward into England. But afterwards, foreseeing the civil wars, he went to seek a retreat at Paris; where he was soon made acquainted with Des Cartes and the other learned philosophers there, with whom he afterwards held a correspondence upon several mathematical subjects, as appears from the letters of Mr. Hobbes published in the works of Des Cartes.

In 1642, Mr. Hobbs printed his famous book *De Cive*, which raised him many adversaries, who charged him with instilling principles of a dangerous tendency. Among many illustrious persons who, from the troubles in England, retired to France for safety, was Sir Charles Cavendish, brother to the Duke of Newcastle: and this gentleman, being well skilled in the mathematics, proved a constant friend and patron to Mr. Hobbes; who, by embarking in 1645 in a controversy



about squaring the circle, was grown so famous by it, that in 1647 he was recommended to instruct Charles prince of Wales, afterwards king Charles the 2d, in mathematical learning. During this he employed his vacant time in composing his *Leviathan*, which was published in England in 1651. After the publication of this work, he returned to England, and passed the remainder of his long life in a very retired and studious manner, in the house of the Earl of Devonshire, mostly at his seat in Derbyshire, but accompanying the earl always to London, fearing to be left out of his immediate protection, lest he should be seized by officers from the parliament or government, on account of the freedom of his opinions in politics and religion. He received great marks of respect from king Charles the 2d at the restoration in 1660, with a pension of 100l. a year. From that time, till his death, he applied himself to his studies, and in opposing the attacks of his adversaries, who were very numerous: in mathematical subjects disputes rose to a great height between him and Dr. Wallis, on account of his pretended Quadrature of the Circle, Cubature of the Sphere, and Duplication of the Cube, which he obstinately defended without ever acknowledging his error.

His long life was that of a perfectly honest man; a lover of his country, a good friend, charitable and obliging. He accustomed himself much more to thinking, than reading; and was fond of a well-selected, rather than a large library. He had a hatred to the clergy, having been persecuted by them, on account of the freedom of his doctrine, and having a very indifferent opinion of their knowledge and their principles. In his last sickness he was very anxious to know whether his disease was curable; and when intimations were given, that he might have ease, but no remedy, he said, 'I shall be glad to find a hole to creep out of the world at.' He died the 4th of Dec. 1679, at 91 years of age.

His chief publications were,

1. An English translation of Thucydides's History of the Grecian war.
2. De Mirabilibus Pecci, and Memoirs of his own Life, both in Latin verse.
3. Elements of Philosophy.
4. Answer to Sir William Davenant's Epistle, or Preface to Gondibert.
5. Human Nature, or the Fundamental Elements of Policy.
6. Elements of Law.
7. *Leviathan*; or the Matter, Form, and Power of a Commonwealth.
8. A Compendium of Aristotle's Rhetoric.
9. A Letter on Liberty and Necessity.
10. The Questions, concerning Necessity and Chance, stated.
11. Six Lessons to the Professors of Mathematics, of the Institution of Sir Henry Saville.
12. The marks of Absurd Geometry, &c.
13. Dialogues of Natural Philosophy.

Besides many other pieces on Polity, Theology, Mathematics, and other miscellaneous subjects, to the number of 41.

HOBITS, in Gunnery. See HOWITZ.

HOGSHEAD, a measure, or vessel, of wine or  
VOL. I.

oil; containing the 4th part of a tun, the half of a pipe, or 63 gallons.

HOLDER (WILLIAM), a learned and philosophical Englishman, was born in Nottinghamshire, educated at Cambridge, and in 1642 became rector of Blechingdon in Oxfordshire. In 1660 he proceeded D. D. he became afterwards canon of Ely, Fellow of the Royal Society, canon of St. Paul's, sub-dean of the royal chapel, and sub-almoner to the king. He was a general scholar, a very accomplished person, and a great virtuoso.

Dr. Holder greatly distinguished himself, by giving speech to a young gentleman of the name of Popham, who was born deaf. This was effected at his own house at Blechingdon in 1659; but the young man losing what he had been taught by Holder after he was called home to his friends, he was sent to Dr. Wallis, who brought him to his speech again. Holder published a book, intitled "the Elements of Speech; an essay or inquiry into the natural Production of Letters: with an appendix, concerning persons that are deaf and dumb, 1669," 8vo. In the appendix he relates how soon, and by what methods, he brought young Popham to speak. In the *Philos. Transf.* for July 1670, was inserted a letter from Dr. Wallis, in which he claims to himself the honour of bringing that gentleman to speak. By way of answer to which, in 1678, Dr. Holder published in 4to, "A Supplement to the *Philos. Transf.* of July 1670, with some reflections on Dr. Wallis's letter there inserted." Upon which the latter soon after published "A Defence of the Royal Society, and the Philosophical Transactions, particularly those of July 1670, in answer to the cavils of Dr. William Holder, 1678," 4to.

Dr. Holder's accomplishments were very general. He was skilled in the theory and practice of music, and wrote "A Treatise of the Natural Grounds and Principles of Harmony, 1694," 8vo. He wrote also "A Treatise concerning Time, with applications of the Natural Day, Lunar Month, and Solar Year, &c, 1694," 8vo. He died at Amen Corner in London, Jan. 24, 1697, and was buried in St. Paul's.

HOLLOW, in Architecture, a concave moulding, about a quarter of a circle, by some called a Casement, by others an Abacus.

HOLLOW-Tower, in Fortification, is a rounding made of the remainder of two brisures, to join the curtain to the crillon, where the small shot are played, that they may not be so much exposed to the view of the enemy.

HOLY *Thursday*, otherwise called Ascension day, being the 39th day after Easter Sunday, and kept in commemoration of Christ's ascension up into heaven.

HOLY *Rood*, or *Holy Cross*, a festival kept on the 14th of September, in memory of the exaltation of our Saviour's cross.

HOLY *Week*, is the last week of Lent, called also Passion Week,

HOLYWOOD (JOHN), or HALIFAX, or *Sacro-bosco*, was, according to Leland, Bale, and Pitts, born at Halifax in Yorkshire: according to Stainhurst, at Holywood near Dublin; and according to Dempster and Mackenzie, in Nithsdale in Scotland. Though  
there



there may perhaps have been more than one of the name. Mackenzie informs us, that having finished his studies, he entered into orders, and became a canon regular of the order of St. Augustin in the famous monastery of Holywood in Nithsdale. The English biographers, on the contrary, tell us that he was educated at Oxford. They all agree however in asserting, that he spent most of his life at Paris; where, says Mackenzie, he was admitted a member of the university, June 5, 1221, under the syndics of the Scotch nation; and soon after was elected professor of mathematics, which he taught with applause for many years. According to the same author, he died in 1256, as appears from the inscription on his monument in the cloisters of the convent of St. Maturine at Paris.

Holywood was contemporary with Roger Bacon, but probably older by about 20 years. He was certainly the first mathematician of his time; and he wrote, 1. *De Sphæra Mundi*; a work often reprinted, and illustrated by various commentators.—2. *De Anni Ratione, seu de Computo Ecclesiastico*.—3. *De Algorismo*, printed with *Comm. Petri Cirvilli Hisp*: Paris, 1498.

HOMOCENTRIC, the same as Concentric.

HOMODROMUS *Vectis*, or *Lever*, in Mechanics, is a lever in which the weight and power are both on the same side of the fulcrum, as in the lever of the 2d and 3d kind; being so called because here the weight and power move both in the same direction, whereas in the Heterodromus they move in opposite directions.

HOMOGENEAL, or HOMOGENEOUS, consisting of similar parts, or of the same kind and nature, in contradistinction from heterogeneous, where the parts are of different kinds.—Natural bodies are usually composed of Homogeneous parts, as a diamond, a metal, &c. But artificial bodies, on the contrary, are assemblages of heterogeneous parts, or parts of different kinds; as a building, of stone, wood, &c.

HOMOGENEAL *Light*, is that whose rays are all of one and the same colour, refrangibility, &c.

HOMOGENEAL *Numbers*, are those of the same kind and nature.

HOMOGENEAL *Surds*, are such as have one common radical sign; as  $\sqrt{27}$  and  $\sqrt{30}$ , or  $\sqrt{a}$  and  $\sqrt{b}$ , or  $\sqrt[3]{c}$  and  $\sqrt[3]{d}$ .

HOMOGENEUM *Adfectionis*, a name given by Vieta to the second term of a compound or affected equation, being that which makes it affected.

HOMOGENEUM *Comparisonis*, in Algebra, a name given by Vieta to the absolute known number or term in a compound or affected equation. This he places on the right-hand side of the equation, and all the other terms on the left.

HOMOLOGOUS, in Geometry, is applied to the corresponding sides of similar figures, or those that are opposite to equal or corresponding angles, and are so called because they are proportional to each other. For all similar figures have their like sides Homologous, or proportional to one another, also their areas or surfaces are Homologous or proportional to the squares of the like sides, and their solid contents Homologous or proportional to the cubes of the same.

HOOKE (ROBERT), a very eminent mathematician

and philosopher, was born, 1635, at Freshwater in the Isle of Wight, where his father was minister. He was intended for the church; but being of a weakly constitution, and very subject to the head-ach, all thoughts of that nature were laid aside. Thus left to himself, the boy followed the bent of his genius, which was turned to mechanics; and employed his time in making little toys, which he did with wonderful art and dexterity. He had also a good turn for drawing; for which reason, after his father's death, which happened in 1648, he was placed with Sir Peter Lely; but the smell of the oil-colours increasing his head-ach, he quitted painting in a very short time. He was afterwards kindly taken by Dr. Busby into his house, and supported there, while he attended Westminster-school; where he not only acquired a competent share of Greek and Latin, together with an insight into Hebrew and some other Oriental languages, but also made himself master of a good part of Euclid's Elements; and, as Wood asserts, invented 30 different ways of flying.

About the year 1653 he went to Christ-church in Oxford; and in 1655 was introduced to the Philosophical Society there; where, discovering his mechanic genius, he was first employed to assist Dr. Willis in his chemical operations, and was afterwards recommended to Mr. Robert Boyle, whom he served several years in the same capacity. He was also instructed in astronomy about this time by Dr. Seth Ward, Savilian Professor of that science; and from henceforward distinguished himself by many noble inventions and improvements of the mechanic kind. He also invented several astronomical instruments, for making observations both at sea and land, and was particularly serviceable to Mr. Boyle in completing the invention of the air-pump. In 1662 he was appointed Curator of Experiments to the Royal Society; and when that body was established by royal charter, he was in the list of those, who were first named by the council in May 20, 1663; and he was admitted accordingly June 3, with a peculiar exemption from all payments. Sept. 28 of the same year, he was named by lord Clarendon, chancellor of Oxford, for the degree of M. A.; and Oct. 19 it was ordered that the repository of the Royal Society should be committed to his care: the white gallery in Gresham-college being appropriated to that use. In May 1664, he began to read the astronomy lecture at Gresham college for the professor Dr. Pope, then in Italy; and the same year he was made Professor of Mechanics to the Royal Society by Sir John Cutler, with a salary of 50l. per annum, which that gentleman, the founder, settled upon him for life. Jan. 11, 1665, that society granted a salary also of 30l. a year, for his office of Curator of Experiments for life; and the month of March the same year he was elected professor of geometry in Gresham-college.

In 1665 too, he published in folio, his "Micrographia, or some Philosophical Descriptions of Minute Bodies, made by Magnifying Glasses, with Observations and Enquiries thereupon." And the same year, during the recess of the Royal Society on account of the plague, he attended Dr. Wilkins and other ingenious gentlemen into Surry, where they made several experiments. In 1666 he produced to the Royal Society



ciety a model for rebuilding the city of London, then destroyed by the great fire, with which the Society was well pleased; and the Lord Mayor and Aldermen preferred it to that of the city surveyor, though it happened not to be carried into execution. The rebuilding of the city according to the act of parliament requiring able persons to set out the ground for the proprietors, Mr. Hooke was appointed one of the surveyors; an employment in which he got most part of his estate, as appeared from a large iron chest of money found after his death, locked down with a key in it, and a date of the time, which shewed it to have been so shut up above 30 years. From 1668 he was engaged for many years in a warm contest with Hevelius, concerning the difference in accuracy between observing with astronomical instruments with plain and telescopic sights; in which dispute many learned men afterwards engaged, and in which Hooke managed so ill, as to be universally condemned, though it has since been agreed that he had the better side of the question.—In 1771 he attacked Newton's "New Theory of Light and Colours;" where, though he was obliged to submit in respect to the argument, it is said he came off with more credit. The Royal Society having commenced their meetings at Gresham-college, November 1674, the Committee in December allowed him 40l. to erect a turret over part of his lodgings, for trying his instruments, and making astronomical observations: and the year following he published "A Description of Telescopes, and some other instruments made by R. H. with a Postscript," complaining of some injustice done him by their secretary Mr. Oldenburg, who published the Philosophical Transactions, in regard to his invention of pendulum watches. This charge drew him into a dispute with that gentleman, which ended in a declaration of the Royal Society in their secretary's favour.—Mr. Oldenburg dying in 1677, Mr. Hooke was appointed to supply his place, and began to take minutes at the meeting in October, but did not publish the Transactions.—Soon after this, he grew more reserved than formerly; and though he read his Cutlerian Lectures, often made experiments, and shewed new inventions before the Royal Society, yet he seldom left any account of them to be entered in their registers; designing, as he said, to publish them himself, which however he never performed.—In 1686, when Newton's work the Principia was published, Hooke laid claim to his discovery concerning the force and action of gravity, which was warmly refuted by that great philosopher. Hooke, though a great inventor and discoverer himself, was yet so envious and ambitious, that he would fain have been thought the only man who could invent and discover. This made him often lay claim to the inventions and discoveries of other persons; on which occasions however, as well as in the present case, the thing was generally carried against him.

In the beginning of the year 1687, his brother's daughter, Mrs. Grace Hooke, who had lived with him several years, died: and he was so affected with grief at her death, that he hardly ever recovered it, but was observed from that time to become less active, more melancholy, and more cynical than ever. At the same time, a chancery suit in which he was concerned with Sir John Cutler, on account of his salary for reading

the Cutlerian Lectures, made him uneasy, and increased his disorder.—In 1691, he was employed in forming the plan of the hospital near Hoxton, founded by Robert Ask, alderman of London, who appointed archbishop Tillotson one of his executors; and in December the same year, Hooke was created M. D. by a warrant from that prelate. In July 1696, the chancery suit with Sir John Cutler was determined in his favour, to his inexpressible satisfaction. His joy on that occasion was found in his diary thus expressed; DOMSHLGISSA; that is, *Deo, Optimo, Maximo, sit honor, laus, gloria, in secula seculorum, Amen.* "I was born on this day of July 1635, and God hath given me a new birth: may I never forget his mercies to me! while he gives me breath may I praise him!"—In the same year 1696, an order was granted to him for repeating most of his experiments at the expence of the Royal Society, upon a promise of his finishing the accounts, observations, and deductions from them, and of perfecting the description of all the instruments contrived by him: but his increasing illness and general decay rendered him unable to perform it. He continued some years in this wasting condition; and thus languishing till he was quite emaciated, he died March 3, 1702, in his 67th year, at his lodgings in Gresham college, and was buried in St. Helen's church, Bishopsgate street; his corps being attended by all the members of the Royal Society then in London.

As to Mr. Hooke's character, it is not in all respects one of the most amiable. In his person he made rather a despicable figure, being but of a short stature, very crooked, pale, lean, and of a meagre aspect, with dark-brown hair, very long, and hanging over his face lank and uncut. Suitable to his person, his temper was penurious, melancholy, and mistrustful: and, though possessed of great philosophical knowledge, he had so much ambition, that he would be thought the only man who could invent or discover; and hence he often laid claim to the inventions and discoveries of others, while he boasted of many of his own which he never communicated. In the religious part of his character, he was so exemplary, that he always expressed a great veneration for the Deity; and seldom received any remarkable benefit in life, or made any considerable discovery in nature, or invented any useful contrivance, or found out any difficult problem, without setting down his acknowledgment to God, as many places in his diary plainly shew.—His chief publications are,

1. *Lectiões Cutlerianæ*, or the Cutlerian Lectures.
2. *Micrographia*, or Descriptions of Minute Bodies made by Magnifying Glasses.
3. A Description of Helioscopes.
4. A Description of some Mechanical Improvements of Lamps and Water-pipes.
5. Philosophical Collections.
6. Posthumous Works, collected from his papers by Richard Waller secretary to the Royal Society. Besides a number of papers in the Philos. Transf. volumes 1, 2, 3, 5, 6, 9, 16, 17, 22.

HORARY, something relating to Hours. As, HORARY Circles, hour lines or circles, marking the hours, or drawn at the distance of hours from one another.

HORARY Motion, is the motion or space moved in an hour.



hour. Thus, the Horary motion of the earth on her axis, is  $15^\circ$ ; for, completing her revolution of  $360^\circ$ , in 24 hours, therefore the motion in one hour will be the 24th part of  $360^\circ$ , which is 15 degrees.

HORIZON, in Astronomy, a great circle of the sphere, dividing the world into two parts, or hemispheres; the one upper, and visible; the other lower, and hid.

The Horizon is either Rational or Sensible.


HORIZON, *Rational, True, or Astronomical*, called also simply and absolutely the Horizon, is a great circle having its plane passing through the centre of the earth, and its poles are the zenith and nadir. Hence all the points of the Horizon, quite around, are at a quadrant distance from the zenith and nadir. Also the meridian and vertical circles cut the Horizon at right angles, and into two equal parts.

HORIZON, *Apparent, Sensible, or Visible*, is a lesser circle of the sphere, parallel to the rational Horizon, dividing the visible part of the sphere from the invisible, and whose plane touches the spherical surface of the earth.

The sensible Horizon is divided into Eastern and Western; the Eastern or Orive being that in which the heavenly bodies rise; and the Western, or Occidial, being that in which they set.

HORIZON, in Geography, is a circle dividing the visible part of the earth and heavens from that which is invisible. This is peculiarly called Sensible or Apparent Horizon, to distinguish it from the Rational or True, which passes through the centre of the earth; as already observed. These two Horizons, though distant from one another by the semidiameter of the earth, will appear to coincide when continued to the sphere of the fixed stars; because the earth compared with this sphere is but a point.

By Sensible Horizon is also often meant a circle which determines the segment of the surface of the earth, over which the eye can reach; called also the Physical Horizon. And in this sense we say, a spacious Horizon, a narrow or scanty Horizon, &c; depending chiefly on the height the eye is elevated above the earth.



For, it is evident that the higher the eye is placed, the farther is the visible Horizon extended. Thus, if the eye be at A, at the height AD above the earth; draw the two tangents Ab, Ar; and let one of these lines Ab, be moved round the point A, and in its revolution always touch the surface of the earth; then

the other point b will describe the visible Horizon hor, &c. But if the eye be placed higher as at B, the tangents BH and BR will reach farther, and the visible Horizon HOR will be larger.

The visible Horizon is most accurately observed at sea, and is therefore sometimes called the Horizon of the sea. In observing this Horizon, the visual rays Ab and Ar will, on account of the curve surface of the sea, always point a little below the true sensible Horizon SS or EF, and consequently below the rational Horizon TT, which is parallel to it.

To find the Depression of the Horizon of the sea below the true Horizon, which varies with the height of the eye, and in a small degree with the variation of the refractive power of the atmosphere, see DEPRESSION.

As to the right-lined distance, or tangent Eb, it may be found thus; as radius : sin.  $\angle C :: CA : Ab$ , or thus; as radius : tan.  $\angle C :: Ch : Ab$ , either of which will be nearly the same as the arc or curved distance Db. Or, without finding the angle C, thus; the square of Ab is equal to the difference of the squares of CA and Ch, i. e.  $Ab^2 = CA^2 - Ch^2 = CA + Ch \times CA - Ch = CA + Ch \times AD$ , and hence  $Ab = \sqrt{CA + Ch \times AD}$ , which is also equal to Ab nearly.

The distance on a perfect globe, if the visual rays came to the eye in a straight line, would be as above stated; but by means of the refraction of the atmosphere, distant objects on the Horizon appear higher than they really are, or appear less depressed below the true Horizon SS, and may be seen at a greater distance, especially on the sea. M. Legendre, in his Memoir on Measurements of the Earth, in the Mem. Acad. Sci. for the year 1787, says that, from several experiments, he is induced to allow for refraction a 14th part of the distance of the place observed, expressed in degrees and minutes of a great circle. Thus, if the distance be 14000 toises, the refraction will be 1000 toises, equal to the 57th part of a degree, or  $1' 3''$ .

HORIZON of the Globe, a broad wooden circle. See GLOBE.

HORIZONTAL, something that relates to the Horizon, or that is taken in the Horizon, or on a level with or parallel to it. Thus, we say, a Horizontal plane, Horizontal line, Horizontal distance, &c.

HORIZONTAL Dial, is one drawn on a plane parallel to the horizon; having its gnomon or style elevated according to the altitude of the pole of the place it is designed for.

HORIZONTAL Distance, is that estimated in the direction of the horizon.

HORIZONTAL Line, in Perspective, is a right line drawn through the principal point, parallel to the horizon; or it is the intersection of the Horizontal and perspective planes.

HORIZONTAL Line, or base of a hill, in Surveying, a line drawn on the Horizontal plane of the hill, or that on which it stands.

HORIZONTAL Moon. See APPARENT MAGNITUDE.

HORIZONTAL Parallax. See PARALLAX.

HORIZONTAL Plane, is that which is parallel to the horizon of the place, or not inclined to it.

HORIZONTAL



**HORIZONTAL Plane**, in Perspective. See PLANE.

**HORIZONTAL Projection**. See PROJECTION, and MAP.

**HORIZONTAL Range**, of a piece of ordnance, is the distance at which it falls on, or strikes the horizon, or on a Horizontal plane, whatever be the angle of elevation or direction of the piece. When the piece is pointed parallel to the horizon, the range is then called the point-blank or point-blanc range.

The greatest Horizontal range, in the parabolic theory, or in a vacuum, is that made with the piece elevated to 45 degrees, and is equal to double the height from which a heavy body must freely fall to acquire the velocity with which the shot is discharged. Thus, a shot being discharged with the velocity of  $v$  feet per second; because gravity generates the velocity  $2g$  or  $32\frac{1}{2}$  feet in the first second of time, by falling  $16\frac{1}{2}$  or  $g$  feet, and because the spaces descended are as the

squares of the velocities, therefore as  $4g^2 : v^2 :: g : \frac{v^2}{4g}$  the space a body must descend to acquire the velocity  $v$  of the shot or the space due to the velocity  $v$ ; consequently the double of this, or  $\frac{v^2}{2g} = \frac{v^2}{32\frac{1}{2}}$  is the greatest

Horizontal range with the velocity  $v$ , or at an elevation of 45 degrees; which is nearly half the square of a quarter of the velocity.

In other elevations, the Horizontal range is as the sine of double the angle of elevation; so that, any other elevation being  $e$ , it will be,

as radius 1 : sin.  $2e :: \frac{v^2}{32\frac{1}{2}} : \frac{v^2}{32\frac{1}{2}} \times \text{sin. } 2e$ , the range at the elevation  $e$ , with the velocity  $v$ .

But in a resisting medium, like the atmosphere, the actual ranges fall far short of the above theorems, in so much that with the great velocities, the actual or real ranges may be less than the 10th part of the potential ranges; so that some balls, which actually range but a mile or two, would in vacuo range 20 or 30 miles. And hence also it happens that the elevation of the piece, to shoot farthest in the resisting medium, is always below 45°, and gradually the more below it as the velocity is greater, so that the greater velocities with which balls are discharged from cannon with gunpowder, require an elevation of the gun equal to but about 30°, or even less. And the less the size of the balls is too, the less must this angle of elevation be, to shoot the farthest with a given velocity. See PROJECTILE, and GUNNERY.

**HORIZONTAL Refraction**. See REFRACTION.

**HORIZONTAL Speculum**, one to find a horizon at Sea, &c, when the atmosphere is hazy near the horizon, by which the sight of it is prevented.

A speculum of this kind was invented by a Mr. Serfon, on the principle of a top spinning, which always keeps its upright position, notwithstanding the motion of the substance it spins upon. This curious instrument, as it has since been improved by Mr. Smeaton, consists of a well polished metal speculum, of about 3 inches and a half in diameter, inclosed within a circular rim of brass; so fitted that the centre of gravity of the whole shall fall near the point on which it spins. This is the end of a steel axis running through the

centre of the speculum, above which it finishes in a square, for the conveniency of fitting a roller on it, which sets it in motion by means of a piece of tape wound about the roller.


Various other contrivances to form artificial horizons have been invented by different persons, as glass planes floating on mercury, &c. See HADLEY'S QUADRANT, and several inventions of this sort in the Philos. Trans. by Elton, Halley, Leigh, &c. vol. xxxvii, p. 273, vol. xxxviii, p. 167, vol. xl, p. 413, 417, &c.

**HORN-Work**, in Fortification, a sort of out-work, advancing toward the field, to cover and defend a curtain, bastion, or other place, suspected to be weaker than the rest; as also to possess a height; carrying in the fore-part, or head, two demi-bastions, resembling horns: these horns, epaulments, or shoulderings, being joined by a curtain, shut up on the side by two wings, parallel to one another, are terminated at the gorge of the work, and so present themselves to the enemy.

**HOROGRAPHY**, the art of making or constructing dials; called also Dialling, Horologiography, Gnomonica, Sciatherica, Photosciatherica, &c.

**HOROLOGIUM**, a common name, among ancient writers, for any instrument or machine for measuring the hours. See CLOCK, WATCH, SUN-DIAL, CHRONOMETER, CLEPSYDRA, &c.

**HOROMETRY**, the art of measuring or dividing time by hours, and keeping the account of time.



**HOROPTER**, in Optics, is a right line drawn through the point where the two optic axes meet, parallel to that which joins the centres of the two eyes, or the two pupils. As the line AB drawn through C the point of concurrence of the optic axes of the eyes, and parallel to HI joining the centres of the eyes.— This line is called the Horopter, because it is found to be the limit of distinct vision. It has several properties in Optics, which are described at large in Aguillonius, Opt. lib. 2, diss. 10.

**HOROSCOPE**, in Astrology, is the ascendant or first house, being that part of the zodiac which is just rising in the eastern side of the horizon at any proposed time, when a scheme is to be set or calculated, or a prediction made of any event. See ASCENDANT.

**HOROSCOPE** is also used for a scheme or figure of the 12 celestial houses; i. e. the 12 signs of the zodiac, in which is marked the disposition of the heavens for any given time. Thus it is said, To draw or construct a Horoscope or scheme, &c. And it is more peculiarly called,



called, Calculating a nativity, when the life and fortune of a person are the subject of prediction.

*Lunar* HOROSCOPE, is the point the moon issues out of, when the sun is in the ascending point of the east; and is also called the Part of Fortune.

HOROSCOPE was also a mathematical instrument, in manner of a planisphere; but now disused. It was invented by J. Paduanus, who wrote a special treatise upon it.

HORROR of a *Vacuum*, an imaginary principle among the more ancient philosophers, to which they ascribed the ascent of water in pumps, and other similar phenomena, which are now known to be occasioned by the weight of the air.

HORROX (JEREMIAH), an eminent English astronomer, was born at Toxteth in Lancashire, about the year 1619. From a grammar school in the country, he was sent to Cambridge, where he spent some time in academical studies. About 1633 he began to apply himself to the study of astronomy: but living at that time with his father at Toxteth, in very moderate circumstances, and being destitute of books and other assistances for such studies, he could not make any considerable progress in it. About the year 1636, he formed an acquaintance with Mr. William Crabtree, of Broughton near Manchester, who was engaged in the same studies, with whom a mutual correspondence was carried on till his death; sometimes communicating their improvements to Mr. Samuel Foster, professor of geometry at Gresham College in London. Having now obtained a companion in his studies, Mr. Horrox assumed new vigour, procured other instruments and books, and was pursuing his studies and observations with great assiduity, when he was suddenly cut off by death, the 3d of January 1640, in the 22d year of his age.

What we have of his writings is sufficient to shew how great a loss the world had by his death. He had just finished his *Venus in Sole visa*, 1639, a little before, as appears by some of the letters to his friend Mr. Crabtree, by which also it appears that he made his observations on that phenomenon at Hool near Liverpool. This tract, of Venus seen in the Sun, was published at Dantzick in 1668, by Hevelius, together with his own *Mercurius in Sole visus* May 3, 1661. His other posthumous works, or rather his imperfect papers, were published by Dr. Wallis, in 1673, 4to, with some account of his life; in which we find he first asserts and promotes the Keplerian astronomy against the hypothesis of Lansberg; which he proves to be inconsistent with itself, and neither agreeing with observations nor theory. He likewise reasons very justly concerning the celestial bodies and their motions, vindicates Tycho Brahe from some objections made to his hypothesis, and gives a new theory of the moon: to which are added the Lunar Numbers of Mr. Flamsteed. There are also extracts from several letters between him and Mr. Crabtree, upon various astronomical subjects; with a catalogue of astronomical observations.

There are two things particularly which will perpetuate the memory of this very extraordinary young man. The one is, that he was the first that ever predicted or saw the planet Venus in the sun; for we do not find

that any persons, besides himself and Mr. Crabtree, ever beheld such a phenomenon. Though he was not apprised of the great use that was to be made of it, in discovering the parallax and distance of the sun and planets, yet he made from it many useful observations, corrections, and improvements in the theory of the motions of Venus.—Secondly, his New Theory of Lunar Motions, which Newton himself made the ground work of all his astronomy, relative to the moon, who always spoke of our author as a genius of the first rank.

HORSE-SHOE, in Fortification, is a work sometimes of a round, sometimes of an oval figure, inclosed with a parapet, raised in the ditch of a marshy place, or in low grounds; sometimes also to cover a gate; or to serve as a lodgment for soldiers, to prevent surprises, or relieve an over-tedious defence.

HOSPITAL, or HOPITAL (WILLIAM-FRANCIS-ANTHONY, *marquis of*), a celebrated French mathematician, was born of an ancient family in 1661. He was a mathematician almost from his infancy; for being one day at the duke of Rohan's, where some able mathematicians were speaking of a problem of Pascal's, which appeared to them very difficult, he ventured to say, that he believed he could resolve it. They were surprised at such presumption in a boy of 15, for he was then no more; however, in a few days he sent them the solution.

M. l'Hospital entered early into the army, and was a captain of horse; but being very short-sighted, and on that account exposed to perpetual inconveniences and errors, he at length quitted the army, and applied himself entirely to his favourite amusement.—He contracted a friendship with Malbranche, and took his opinion upon all occasions.—In 1699 he was received an honorary member of the Academy of Sciences at Paris.

He was the first person in France who wrote upon Newton's analysis, and on this account was regarded almost as a prodigy. His work was entitled *l'Analyse des Infinimens Petits*, 1696. He engaged afterwards in another mathematical work, in which he included *Les Sections Coniques, les Lieux Géométriques, la Construction des Equations, et une Théorie des Courbes Mécaniques*: but, a little before he had finished it he was seized with a fever, which carried him off, the 2d of February 1704, at 43 years of age. The work was published after his death, viz, in 1707. There are also six of his pieces inserted in different volumes of the Memoirs of the Academy of Sciences.

HOUR, in Chronology, an aliquot part of a natural day, usually the 24th, but sometimes a 12th part. With us, it is the 24th part of the earth's diurnal rotation, or the time from noon to noon, and therefore it answers to 15 degrees of the whole circle of longitude, or of 360°. The hour is divided by 60ths, viz, first into 60 minutes, then each minute into 60 seconds, &c.

The division of the day into Hours is very ancient; as is shewn by Kircher, *Oedip. Ægypt.* tom. 2, par. 2, class 7, cap. 8. The most ancient Hour is that of the 12th part of the day. Herodotus, lib. 2, observes, that the Greeks learnt from the Egyptians, among other things, the method of dividing the day into 12 parts. And the astronomers of Cathaya, &c, still retain this division.



The division of the day into 24 hours, was not known to the Romans before the Punic war. Till that time they only regulated their days by the rising and setting of the sun. They divided the 12 hours of their day into four; viz, Prime, which commenced at 6 o'clock, Third at 9, Sixth at 12, and None at 3. They also divided the night into four watches, each containing 3 Hours.

There are various kinds of Hours, used by chronologers, astronomers, dialists, &c. Sometimes too,

Hours are divided into Equal and Unequal.

*Equal Hours*, are the 24th parts of a day and night precisely; that is, the time in which the 15 degrees of the equator pass the meridian. These are also called Equinoctial Hours, because measured on the equinoctial; and Astronomical, because used by astronomers.

*Astronomical Hours*, are equal Hours, reckoned from noon to noon, in a continued series of 24.

*Babylonish Hours*, are equal Hours, reckoned from sun-rise in a continued series of 24.

*European Hours*, used in civil computation, are equal Hours, reckoned from midnight; 12 from thence till noon, and 12 more from noon till midnight.

*Jewish, or Planetary, or Ancient Hours*, are 12th parts of the artificial day and night. They are called Ancient or Jewish Hours, because used by the ancients, and still among the Jews. They are called Planetary Hours, because the astrologers pretend, that a new planet comes to predominate every Hour; and that the day takes its denomination from that which predominates the first Hour of it; as Monday from the moon, &c.

*Italian Hours*, are equal Hours, reckoned from sun-set, in a continued series of 24.

*Unequal or Temporary Hours*, are 12th parts of the artificial day and night. The obliquity of the sphere renders these more or less unequal at different times; so that they only agree with the equal Hours at the times of the equinoxes.

*Hour-Circles, or HORARY-Circles*, are great Circles, meeting in the poles of the globe or world, and crossing the equinoctial or equator at right angles; the same as meridians. They are supposed to be drawn through every 15th degree of the equinoctial and equator, each answering to an hour, and dividing them into 24 equal parts; and on both globes they are supplied by the meridian Hour-circle and index.

*Hour-Glass*, a popular kind of chronometer or clepsydra, serving to measure time by the descent or running of sand, water, &c, out of one glass vessel into another. —The best, it is said, are such as, instead of sand, have egg-shells, well dried in the oven, then beaten fine and sifted.

*Hour-Lines*, on a Dial, are lines which arise from the intersections of the plane of the Dial, with the several planes of the Hour-circles of the sphere; and therefore must be all right lines on a plane Dial.

*Hour-Scale*, a divided line on the edge of Collins's quadrant, being only two lines of tangents of 45 degrees each, set together in the middle. Its use, together with the lines of latitude, is to draw the Hour-lines of Dials that have centres, by means of an equilateral triangle, drawn on the dial-planes.

*HOWITZ, or HOWITZER*, in Artillery, a kind of

mortar, or something between a cannon and mortar, partaking of the nature of both, being either a very short gun or a long mortar. It is of German invention, and is mounted upon a carriage like a travelling gun-carriage, with its trunnions placed nearly in the middle. The Howitz is one of the most useful kinds of ordnance, as it can be employed occasionally either as a cannon or mortar, discharging either shells or grape shot, as well as balls, and so doing great execution. They are also very easily travelled about from place to place.

*HUMIDITY*, or moisture, the power or quality of wetting or moistening other bodies, and adhering to them.

Fluids are moist to some bodies, and not to others. Thus, quicksilver is not moist in respect to our hands or clothes, and other things, which it will not stick to; but it may be called Humid in reference to gold, tin, or lead, to the surfaces of which it will presently adhere, and render them soft and moist. Even water itself, which wets almost every thing, and is the great standard of moisture and Humidity, is not capable of wetting all things; for it stands or runs off in globular drops from any thing greased or oiled, or the leaves of cabbages, and many other planets; and it will not wet the feathers of ducks, geese, swans, and other water-fowl.

*HUNDRED*, the number of ten times ten, or the square of 10. The place of Hundreds makes the third in order in the Arabic or modern numeration, being denoted thus 100. In the Roman notation it is denoted by the letter C, being the initial of its name, Centum.

*HUNDRED Weight*, or the great Hundred, contains 112 pounds weight. It is subdivided into 4 quarters, and each quarter into 28 lbs.

*HURTERS*, in Fortification, denote pieces of timber, about 6 inches square, placed at the lower end of the platform, next to the parapet, to prevent the wheels of the gun-carriages from damaging the parapet.

*HUYGENS (CHRISTIAN)*, a very eminent astronomer and mathematician, was born at the Hague in Holland, in 1629, being the son of Constantine Huygens, lord of Zuylichem, who had served three successive princes of Orange in the quality of secretary. He spent his whole life in cultivating the mathematics; and not in the speculative way only, but also in making them subservient to the uses of life. From his infancy he discovered an extraordinary fondness for the mathematics; in a short time made a great progress in them; and perfected himself in those studies under professor Schooten, at Leyden. In 1649 he went to Holstein and Denmark, in the retinue of Henry count of Nassau; and was extremely desirous of going to Sweden, to visit Des Cartes who was then in that country with the queen Christina, but the count's short stay in Denmark would not permit him. —In 1651 he gave the world a specimen of his genius for mathematics, in a treatise intitled, *Theoremata de Quadratura Hyperbolæ, Ellipsis, & Circuli, ex dato Portionum Gravitatis Centro*; in which he clearly shewed what might be expected from him afterwards. —In 1655 he travelled into France, and took the degree of LL.D. at Angers. —In 1658 he published his *Horologium Oscillatorium, sive de Motu Pendulorum*, &c, at the Hague. He had exhibited



hibited in a former work, intituled, *Brevis Institutio de Usu Horologiorum ad inveniendas Longitudines*, a model of a new invented pendulum; but as some persons, envious of his reputation, were labouring to deprive him of the honour of the invention, he wrote this book to explain the construction of it; and to shew that it was very different from the pendulum of astronomers invented by Galileo.—In 1659 he published his *Systema Saturninum*, &c; in which he first of any one explained the ring of Saturn, and discovered also one of the satellites belonging to that planet, which had hitherto escaped the eyes of astronomers: new discoveries, made with glasses of his own forming, which gained him a high rank among the astronomers of his time.

In 1660, he took a second journey into France, and the year after passed over into England, where he communicated his art of polishing glasses for telescopes, and was made Fellow of the Royal Society. About this time the air-pump was invented, which received considerable improvements from him. This year also he discovered the laws of the collision of elastic bodies; as did also about this time Wallis and Wren, with whom he had a dispute about the honour of this discovery. Upon his return to France, in 1663, the minister Colbert, being informed of his great merit, settled a considerable pension upon him, to engage him to fix at Paris; to which Mr. Huygens consented, and staid there from the year 1666 to 1681, where he was admitted a member of the Academy of Sciences. All this time he spent in mathematical pursuits, wrote several books, which were published from time to time, and invented and perfected several useful instruments and machines: particularly he had a dispute, about the year 1668, with Mr. James Gregory, concerning the Quadrature of the Circle and Hyperbola of the latter, then just published, in which Huygens it seems had the better side of the question. But continual application gradually impaired his health; and though he had visited his native country twice, viz, in 1670 and 1675, for the recovery of it, he was now obliged to betake himself to it altogether. Accordingly he left Paris in 1681, and retired to his own country, where he spent the remainder of his life in the same pursuits and employments. He died at the Hague, June 8, 1695, in the 67th year of his age, while his *Cosmotheoros*, or treatise concerning a plurality of worlds, was printing; so that this work did not appear till 1698.

Mr. Huygens loved a quiet and studious manner of life, and frequently retired into the country to avoid interruption, but did not contract that moroseness which is so commonly the effect of solitude and retirement. He was one of the purest and most ingenious mathematicians of his age, and indeed of any other; and made many valuable discoveries. He was the first who discovered Saturn's ring, and a third satellite of that planet, as mentioned above. He invented the means of rendering clocks exact, by applying the pendulum, and of rendering all its vibrations equal, by the cycloid. He brought telescopes to perfection, and made many other useful discoveries.

He was the author of many excellent works. The principal of these are now contained in two collections, of 2 volumes each, printed in 4to, under the care of professor Gravefande. The first was at Leyden

in 1682, under the title of *Opera Varia*; and the second at Amsterdam, in 1728, entitled *Opera Reliqua*.

HYADES, a cluster of 5 stars in the face of the constellation Taurus, or the Bull.

HYALOIDES, the vitreous humour of the eye contained between the tunica-retina and the uvea.

HYBERNAL Occident. See Occident.

HYBERNAL Orient. See Orient.

HYDATOIDES, the watery humour of the eye contained between the cornea and the uvea.

HYDRA, a southern constellation, consisting of a number of stars, imagined to represent a water serpent. The stars in the constellation Hydra, in Ptolemy's catalogue, are 27; in Tycho's, 19; in Hevelius's, 31; and in the Britannic catalogue, 60.

HYDRAULICS, the science of the motion of water and other fluids, with its application in artificial water-works of all sorts.—As to what respects merely the equilibrium of fluids, or their gravitation or action at rest, belongs to Hydrostatics. Upon removing or destroying that equilibrium, motion ensues; and here Hydraulics commence. Hydraulics therefore suppose Hydrostatics; and many writers, from the near relation between them, like mechanics and statics, join the two together, and treat of them conjointly as one science.

The laws of Hydraulics are given under the word Fluid. And the art of raising water, with the several machines employed for that purpose, are described under their several names, Fountain, Hydrocanisterium, Pump, Siphon, Syringe, &c.

The principal writers who have cultivated and improved Hydraulics and Hydrostatics, are Archimedes, in his *Libris de Insidentibus Humido*; Hero of Alexandria, in his *Liber Spiritualium*; Marinus Ghetaldus, in his *Archimedes promotus*; Mr. Oughtred; Jo. Ceva, in his *Geometria Motus*; Jo. Bap. Balianus, *De Motu Naturali Gravium, Solidorum et Liquidorum*; Mariotte, in his treatise of the Motion of Water and other fluids; Boyle, in his *Hydrostatical Paradoxes*; Fran. Tertius de Lanis, in his *Magisterium Naturæ et Artis*; Lamy, in his *Traité de l'Equilibre des Liqueurs*; Rohault; Dr. Wallis, in his *Mechanics*; Dechales; Newton, in his *Principia*; Gulielmeni, in his *Mensura Aquarum Fluentium*; Herman; Wolfius; Gravefande; Musschenbroek; Leopold; Schottus, in his *Mechanica Hydraulico-Pneumatica*; Geo. Andr. Bockler, in his *Architectura Curiosa Germanica*; August. Rammilleis; Lucas Antonius Portius; Sturm, in his treatise on the Construction of Mills; Switzer's *Hydrostatics*; Varignon, in the *Mem. Acad. Sci.*; Jurin; Belidor; Bernoulli; Desaguliers; Clare; Emerson; Ferguson; Ximenes; Bossu; D'Alembert; Buat; &c, &c.

HYDRAULICO-PNEUMATICAL, a term applied by some authors to such engines as raise water by means of the weight or spring of the air.

HYDROGRAPHICAL Charts or Maps, more usually called sea-charts, are projections of some part of the sea, or coast, for the use of navigation.

In these are laid down all the rhumbs or points of the compass, the meridians, parallels, &c, with the coasts, capes, islands, rocks, shoals, shallows, &c, in their proper places, and proportions.

The



The making and selling these charts was for some time the employment of Columbus, the first discoverer of America. The story goes, that happening to be heir to the memoirs and journals of one Alonzo Sanchez de Huelva, a noted pilot and captain of a ship, who by chance had been driven by a storm to the island of St. Domingo, and dying at Columbus's house soon after his return, this gave Columbus the first hint to attempt a discovery of the West Indies.

For the construction and use of the several kinds of Hydrographical Maps, see CHART, and SAILING.

HYDROLOGY, is that part of natural history which examines and explains the nature and properties of water in general.

HYDROMANCY, the art or art of divining or foretelling future events by means of water.

This is one of the four general kinds of divination: the other three respecting the other elements, fire, air, and earth, are denominated respectively pyromancy, aeromancy, and geomancy.

Varro mentions the Persians as the first inventors of Hydromancy, adding, that Numa Pompilius and Pythagoras made use of it.

The writers on optics furnish us with divers Hydromantic machines, vessels, &c. For example,

To construct an Hydromantic machine, by means of which an image or object shall be removed out of the sight of the spectator, and restored again at pleasure, without altering the position, either of the one or the other. Provide two vessels, ABF, CGLK (Plate 12, fig. 3), the uppermost filled with water, and supported by three little pillars, one of which BC is hollow, and furnished with a cock B. Let the lower vessel CL be divided by a partition HI into two parts, the lower of which may be opened or closed by means of a cock at P. Upon the partition place an object, or image, which the spectator at O cannot see by a direct ray GL.

If now the cock B be opened, the water descending into the cavity CI, the ray GL will be refracted from the perpendicular GR to O; so that the spectator will now see the object by the refracted ray OG. And again, shutting the cock B, and opening the other P, the water will descend into the lower cavity HL; where, the refraction ceasing, no rays will now come from the object to the eye: but upon shutting the cock P, and opening the other B, the water will fill the cavity again, and bring the object in sight of O afresh.

To make an Hydromantic vessel, which shall exhibit the images of external objects as if swimming in water. Provide a cylindrical vessel ABCD (fig. 4, pl. 12.) divided into two cavities by a glass EF, not perfectly polished: in G apply a lens convex on both sides; and in H incline a plane mirror, of an elliptic figure, to an angle of 45 degrees; and let IH and HG be something less than the distance of the focus of the lens G; so that the place of the images of an object radiating through the same may fall within the cavity of the upper vessel: let the inner cavity be blacked, and the upper filled with clear water.—If now the vessel be disposed in a dark place, so as the lens be turned towards an object illuminated by the sun, its image will be seen as swimming in the water.

HYDROMETER, an instrument for measuring

the properties and effects of water, as its density, gravity, force, velocity, &c.

That with which the specific gravity of water is determined, is often called an aerometer, or water-poise.

The general principle on which the construction and use of the Hydrometer depends, has been illustrated under the article *Specific Gravity*; where it is shewn that a body specifically lighter than several fluids, will serve to find out their specific gravities; because it will sink deepest in the fluids whose specific gravity is the least. So if AB (fig. 5, pl. 12) be a small even glass tube, hermetically sealed, having a scale of equal divisions marked upon it, with a hollow ball of about an inch in diameter at bottom, and a smaller ball C under it, communicating with the first; into the little ball is put mercury or small shot, before the tube is sealed, so that it may sink in water below the ball, and float or stand upright, the divisions on the stem shewing how far it sinks.—If this instrument be dipped in common water, and sink to D, it will sink only to some lower point E in salt water; but in port wine it will sink to some higher point F, and in brandy perhaps to B.

It is evident that an Hydrometer of this kind will only shew that one liquid is specifically heavier than another; but the true specific weight of any liquid cannot be determined without a calculation for this particular instrument, the tube of which should be truly cylindrical. Besides, these instruments will not serve for fluids whose densities are much different:

Mr. Clarke constructed a new Hydrometer, shewing whether any spirits be proof, or above or below proof, and in what degree. This instrument was made of a ball of copper (because ivory imbibes spirituous liquors, and glass is apt to break), to which is soldered a brass wire about a quarter of an inch thick; upon this wire is marked the point to which it exactly sinks in proof spirits; as also two other marks, one above and one below the former, exactly answering to one-tenth above proof and one-tenth below proof. There are also a number of small weights made to add to it, so as to answer to the other degrees of strength besides those above, and for determining the specific gravities of different fluids. *Philos. Trans. Abr. vol. vi, p. 326.*

Dr. Defaguliers contrived an Hydrometer for determining the specific gravities of different waters, to such a degree of nicety, that it would shew when one kind of water was but the 40,000th part heavier than another. It consists of a hollow glass ball of about 3 inches in diameter, charged with shot to a proper degree, and having fixed in it a long and very slender wire, of only the 40th part of an inch in diameter, and divided into tenths of inches, each tenth answering to the 40,000th part, as above. See his *Exper. Philos. vol. 2, p. 234.*

Mr. Quin and other persons have also constructed Hydrometers, with other and various contrivances, and with different degrees of accuracy; but all nearly on the same general principles.

But there is one circumstance which deserves particular attention in the construction and graduation of Hydrometers, for determining the precise strength of different brandies, and other spirituous liquors. Mr. Reaumur discovered, in making his spirit thermometers, that when rectified spirit and water, or phlegm,



the other constituent part of brandy, are mixed together, there appears to be a mutual penetration of the two liquors, and not merely juxtaposition of parts; so that a part of the one fluid seems to be received into the pores of the other; by which it happens, that if a pint of rectified spirit be added to a pint of water, the mixture will be sensibly less than a quart. The variations hence produced in the bulk of the mixed fluid render the Hydrometer, when graduated in the usual way by equal divisions, an erroneous measure of its strength; because the specific gravity of the compound is found not to correspond to the mean gravity of the two ingredients. M. Montigny constructed a scale for this instrument in the manner before suggested by Dr. Lewis, on actual observation of the sinking or rising of the Hydrometer in various mixtures of alcohol and water, made in certain known proportions. *Hist. de l'Acad. Roy. des Sci.* 1768; also Neumann's *Chem.* by Lewis, p. 450, note r.

M. De Luc has lately published a scheme for the construction of a comparable Hydrometer, so that a workman, after having constructed one upon his principles, may make all others similar to each other, and capable of indicating the same degree on the scale, when immersed in the same liquor of the same temperature. This instrument is proposed to be constructed of a ball of flint glass, communicating with a small hollow cylinder, containing such a quantity of quicksilver for a ballast, that the instrument may sink nearly to the top, in the most spirituous liquor, made as hot as possible; to which is also attached a thin silvered tube, for a scale, &c. The whole description may be seen at large in the *Philos. Trans.* vol. 68, p. 500.

M. Le Roi also published a proposal for constructing comparable Hydrometers. See *Hist. de l'Acad. des Sciens.* for 1770, Mem. 7.

**HYDROMETRIA, HYDROMETRY**, the mensuration of water and other fluid bodies, their gravity, force, velocity, quantity, &c; including both hydrostatics and hydraulics.


**HYDROSCOPE**, an instrument anciently used for the measure of time. It was a kind of water-clock, consisting of a cylindrical tube, conical at bottom: the cylinder was graduated with divisions, to which the top of the water becoming successively contiguous, as it trickled out of the vertex of the cone, pointed out the hour.

**HYDROSTATICAL Balance**, a kind of balance contrived for the exact and easy finding the specific gravities of bodies, both solid and fluid, and thereby of estimating the degree of purity of bodies of all kinds, with the quality and richness of metals, ores, minerals, &c, and the proportions in any mixture, adulteration, or the like.

This is effected by weighing the body both in water, or other fluid, and out of it; and for this purpose one of the scales has usually a hook at the bottom, for suspending the body by some very fine thread. And the use of the instrument is founded on this theorem of Archimedes, that any body weighed in water, loses as much of its weight as is equal to the weight of the same bulk of the water. Thus then is known the proportion of the specific gravities of the solid and fluid, or the proportion of their weights under the same bulk,

viz, the proportion of the weight of the body weighed out of water, to the difference between the same and its weight in water. Hence also, by doing the same thing for several different solids, with the same fluid, or different fluids with the same solid, all their specific gravities become known.

The instrument needs but little description. AB is



a nice balance beam, with its scales C and D, turning with the small part of a grain, the one of them, D, having a hook in the bottom, to receive the loop of a horse hair &c, E, by which the body F is suspended. GH is a jar of water, in which the body is immersed when weighing.

The pieces in the scale C denote the weight of the body out of water; then, upon immersing it, put weights in the scale D to restore the balance again, and they will shew the specific gravity of the body.

There have been various kinds of the Hydrostatical balance, and improvements made on it, by different persons. Thus, Dr. Defaguliers set three screws in the foot of the stand, to move any side higher or lower, till the stem be quite upright, which is known by a plummet hanging over a fixed point in the pedestal. *Defag. Exp. Philos.* vol. 2, p. 196. And for sundry other constructions of this instrument, designed for greater accuracy than the common sort, see Martin's *Phil. Britan.* or Gravesande's *Physices Elem. Math.* tom. 1, lib. 3, cap. 3, &c.

The specific gravities of small weights may be determined by suspending them in loops of horse hair, or fine filken threads, to the hook at the bottom of the scale. Thus, if a guinea suspended in air weigh 129 grains, and upon being immersed in water require  $7\frac{1}{2}$  grains to be put in the scale over it, to restore the equilibrium; we thus find that a quantity of water of equal bulk with the guinea, weighs  $7\frac{1}{2}$  grains, or 7.2; therefore dividing the 129 by the 7.2, the quotient 17.88 shews that the guinea is so many times heavier than its bulk of water. Whence, if any piece of gold be tried, by weighing it first in air, then in water, and if, upon dividing the weight in air by the loss in water, the quotient be 17.88, the gold is good; if the quotient be 18 or more, the gold is more fine; but if it be less than 17.88, the gold is too much alloyed with other metal. If silver be tried in the same manner, and found to be 11 times heavier than water, it is very fine; if it be  $10\frac{1}{2}$  times heavier, it is standard; but if less, it is mixed with some lighter metal, such as tin.

When the body, whose specific gravity is sought, is lighter than water, so that it will not quite sink; annex



to it a piece of another body heavier than water, so that the mass compounded of the two may sink together. Weigh the denser body, and the compound mass, separately, both in water and out of it, thereby finding how much each loses in water; and subtract the loss of these two losses from the greater; then say,

As the remainder

is to the weight of the light body in air,  
so is the specific gravity of water  
to the specific gravity of the light body.


**HYDROSTATICAL Bellows**, a machine for shewing the upward pressure of fluids and the Hydrostatical paradox. It consists of two thick boards, A, D, each about 16 or 18 inches diameter, more or less, covered or connected firmly with leather round the edges, to open and shut like a common bellows, but without valves; only a pipe, B, about 3 feet high is fixed into the bellows at *e*. Now let water be poured into the pipe at C, and it will run into the bellows, gradually separating the boards, by raising the upper one. Then if several weights, as three hundred weights, be laid upon the upper board, by pouring the water in at the pipe till it be full, it will sustain all the weights, though the water in the pipe should not weigh a quarter of a pound; for the pipe or tube may be as small as we please, provided it be but long enough, the whole effect depending upon the height, and not at all on the width of the pipe: for the proportion is always this,

As the area of the orifice of the pipe  
is to the area of the bellows board,  
so is the weight of water in the pipe  
to the weight it will sustain on the board.


Hence if a man stand upon the upper board, and blow into the pipe B, he will raise himself upon the board; and the smaller the pipe, the easier he will be able to raise himself; and then by putting his finger upon the top of the pipe, he can support himself as long as he pleases, provided the bellows be air-tight.

Mr. Ferguson has described another machine, which may be substituted instead of this common Hydrostatical bellows. It is however on the same principle of the Hydrostatical paradox; and may be seen in the Supplement to his Lectures, p. 19.

**HYDROSTATICAL Paradox**, is a principle in Hydrostatics, so called because it has a paradoxical appearance at first view, and it is this; that any quantity of water, or other fluid, how small soever, may be made to balance and support any quantity, or any weight, how great soever. This is partly illustrated in the last article, on the Hydrostatical bellows, where it appears that any weight whatever may be blown up and supported by the breath from a person's mouth. And the principle may be explained as follows: It is well-known that water in a pipe or canal, open at both ends, always rises to the same height at both ends, whether those ends be wide or narrow, equal or unequal. Thus, the small pipe GH being close joined to another open vessel AI, of any size whatever; then




pouring water into the one of these, it will rise up in the other, and stand at the same height, or horizontal line DF in both of them, and that whether they are upright, or inclined in any position. So that all the water that is in the large vessel from A to I, is supported by that which is in the small vessel from D to I only. And as there is no limit to this latter one, but that it may be made as fine even as a hair, it hence evidently appears that any quantity of water may be thus supported by any other the smallest quantity.



Since then the pressure of fluids is directly as their perpendicular heights, without any regard to their quantities, it appears that whatever the figure or size of the vessels may be, if they are but of equal heights, and the areas of their bottoms equal, the pressures of equal heights of water are equal upon the bottoms of these vessels; even though the one should contain a thousand or ten thousand times as much as the other.

Mr. Ferguson confirms and illustrates this paradox by the following experiment.



Let two vessels be prepared of equal heights, but very unequal contents, such as AB and CD; each vessel being open at both ends, and their bottoms E and F of equal widths. Let a brass bottom G and H be exactly fitted to each vessel, not to go into it, but for it to stand upon; and let a piece of wet leather be put between



between each vessel and its brass bottom, for the sake of closeness. Join each bottom to its vessel by a hinge D, so that it may open like the lid of a box; and let each bottom be kept up to its vessel by equal weights W, hung to lines which go over the pulleys P, whose blocks are fixed to the sides of the vessel at *f*, and the lines tied to hooks at *d*, fixed in the brass bottoms opposite to the hinges D. Things being thus prepared and fitted, hold one vessel upright in the hands over a basin on a table, and cause water to be poured slowly into it, till the pressure of the water bears down its bottom at the side *d*, and raises the weight E; and then part of the water will run out at *d*. Mark the height at which the surface H of the water stood in the vessel, when the bottom began to give way at *d*; and then, holding up the other vessel in the same manner, cause water to be poured into it; and it will be seen that when the water rises in this vessel just as high as it did in the former, its bottom will also give way at *d*, and it will lose part of the water.

The natural reason of this surprising phenomenon is, that since all parts of a fluid at equal depths below the surface, are equally pressed in all manner of directions, the water immediately below the fixed part B*f* will be pressed as much upward against its lower surface within the vessel, by the action of the column Ag, as it would be by a column of the same height, and of any diameter whatever; and therefore, since action and reaction are equal and contrary to each other, the water immediately below the surface B*f* will be pressed as much downward by it, as if it were immediately touched and pressed by a column of the height Ag, and of the diameter B*f*; and therefore the water in the cavity BD*df* will be pressed as much downward upon its bottom G, as the bottom of the other vessel is pressed by all the water above it. Lectures, p. 105.

**HYDROSTATICS**, is the science which treats of the nature, gravity, pressure, and equilibrium of fluids; and of the weighing of solids in them.

That part of the science of fluids which treats of their motions, being included under the head of Hydraulics.

Hydrostatics and hydraulics together constitute a branch of philosophy that is justly considered as one of the most curious, ingenious, and useful of any; affording theorems and phenomena not only of the first use and importance, but also surprisingly amusing and pleasant; as appears in the numberless writings upon the subject; the principal points of which may be found under the several particular articles of this work; and the chief writings on this science may be seen under the article Hydraulics.


**HYDRUS**, or *Water Serpent*, one of the new southern constellations, including only ten stars.

**HYEMAL Solstice**, the same with Winter Solstice. See SOLSTICE.

**HYGROMETER**, or **HYGROSCOPE**, or **NOTIOMETER**, an instrument for measuring the degrees of moisture in the air.

There are various kinds of Hygrometers; for whatever body either swells by moisture, or shrinks by dryness, is capable of being formed into an Hygrometer. Such are woods of most kinds, particularly deal, ash, poplar, &c. Such also is catgut, the beard of a wild


oat, and twisted cord, &c. The best and most usual contrivances for this purpose are as follow.



1. Stretch a common cord, or a fiddle-string, ABD along a wall, passing it over a pulley B; fixing it at one end A, and to the other end hanging a weight E, carrying a style or index F. Against the same wall fit a plate of metal HI, graduated, or divided into any number of equal parts; and the Hygrometer is complete.

For it is matter of constant observation, that moisture sensibly shortens cords and strings; and that, as the moisture evaporates, they return to their former length again. The like may be said of a fiddle-string: and from hence it happens that such strings are apt to break in damp weather, if they are not slackened by the screws of the violin. Hence it follows, that the weight E will ascend when the air is more moist, and descend again when it becomes drier. By which means the index F will be carried up and down, and, by pointing to the several divisions on the scale, will shew the degrees of moisture or dryness.

2. Or thus, for a more sensible and accurate Hygro-




meter: strain a whiplcord, or catgut, over several pulleys B, C, D, E, F; and proceed as before for the rest of the construction. Nor does it matter whether the several parts of the cord be parallel to the horizon, as expressed in the annexed figure, or perpendicular to the same, or in any other position; the advantage of this, over the former method, being merely the having a greater length of cord in the same compass; for the longer the cord, the greater is the contraction and dilatation, and consequently the degrees of variation of the index over the scale, for any given change of moisture in the air.

3. Or



3. Or thus: Fasten a twisted cord, or fiddle-string, AB, by one end at A, sustaining a weight at B, carrying an index C round a circular scale DE described on a horizontal board or table.—For a cord or catgut twists itself as it moistens, and untwists again as it dries. Hence, upon an increase or decrease of the humidity of the air, the index will shew the quantity of twisting or untwisting, and consequently the increase or decrease of moisture or dryness.




4. Those Dutch toys, called weather houses, where a small image of a man, and one of a woman, are fixed upon the ends of an index, are constructed upon this principle. For the index, being sustained by a cord or twisted catgut, turns backwards and forwards, bringing out the man in wet weather, and the woman in dry.

5. Or thus: Fasten one end of a cord, or catgut, AB, to a hook at A; and to the other end a ball D of about one pound weight; upon which draw two concentric circles, and divide them into any number of equal parts, for a scale; then fit a style or index EC into a proper support at E, so as the extremity C may almost touch the divisions of the ball.—Here the cord twisting or untwisting, as in the former case, will indicate the change of moisture, by the successive application of the divisions of the circular scale, as the ball turns round, to the index C.



6. Or an Hygrometer may be made of the thin boards of ash or fir, by their swelling or contracting. But this, and all the other kinds of this instrument, above described, become in time sensibly less and less accurate; till at last they lose their effect entirely, and suffer no alteration from the weather. But the following sort is much more durable, serving for many years with tolerable accuracy.




7. Take the Manoscope, described under that article, and instead of the exhausted ball E, substitute a sponge, or other body, that easily imbibes moisture. To prepare the sponge, it may be proper first to wash it in water

very clean; and, when dry again, in water or vinegar in which there has been dissolved sal ammoniac, or salt of tartar; after which let it dry again.—Now, if the air become moist, the sponge will imbibe it and grow heavier, and consequently will preponderate, and turn the index towards C; on the contrary, when the air becomes drier, the sponge becomes lighter, and the index turns towards A; and thus shewing the state of the air.


8. In the last mentioned Hygrometer, Mr. Gould, in the *Philos. Trans.* instead of a sponge, recommends oil of vitriol, which grows sensibly lighter or heavier from the degrees of moisture in the air; so that being saturated in the moistest weather, it afterwards retains or loses its acquired weight, as the air proves more or less moist. The alteration in this liquor is so great, that in the space of 57 days it has been known to change its weight from 3 drachms to 9; and has shifted a tongue or index of a balance 30 degrees. So that in this way a pair of scales may afford a very nice Hygrometer. The same author suggests, that oil of sulphur or campanam, or oil of tartar per deliquium, or the liquor of fixed nitre, might be used instead of the oil of vitriol.

9. This balance may be contrived in two ways; by either having the pin in the middle of the beam, with a slender tongue a foot and a half long, pointing to the divisions on an arched plate, as represented in the last figure above. Or the scale with the liquor may be hung to the point of the beam near the pin, and the other



extremity made so long, as to describe a large arch on a board placed for the purpose; as in the figure here annexed.

10. Mr. Arderon has proposed some improvement in the sponge Hygrometer. He directs the sponge A to



be so cut, as to contain as large a superficies as possible, and to hang by a fine thread of silk upon the beam of a balance.




balance B, and exactly balanced on the other side by another thread of silk at D, strung with the smallest lead shot, at equal distances, so adjusted as to cause an index E to point at G, the middle of a graduated arch FGH, when the air is in a middle state between the greatest moisture and the greatest dryness. Under this silk so strung with shot, is placed a little table or shelf I, for that part of the silk or shot to rest upon which is not suspended. When the moisture imbibed by the sponge increases its weight, it will raise the index, with part of the shot, from the table, and vice versa when the air is dry. *Philos. Transf.* vol. 44, p. 96.

11. From a series of Hygroscoical observations, made with an apparatus of deal wood, described in the *Philos. Transf.* number 480, Mr. Coniers concludes, 1st, That the wood shrinks most in summer, and swells most in winter, but is most liable to change in the spring and fall. 2d, That this motion happens chiefly in the day time, there being scarce any variation in the night. 3d, That there is a motion even in dry weather, the wood swelling in the morning, and shrinking in the afternoon. 4th, That the wood, by night as well as by day, usually shrinks when the wind is in the north, north-east, and east, both in summer and winter. 5th, That by constant observation of the motion and rest of the wood, with the help of a thermometer, the direction of the wind may be told nearly without a weather-cock. He adds, that even the time of the year may be known by it; for in spring it moves more and quicker than in winter; in summer it is more shrunk than in spring; and has less motion in autumn than in summer.

See an account of a method of constructing these and other Hygrometers, in *Phil. Transf. Abr.* vol. 2, p. 30, &c, and plate 1 annexed. See also *Philos. Transf.* vol. 11, p. 647 and 715, vol. 15, p. 1032, vol. 43, p. 6, vol. 44, p. 95, 169 and 184, vol. 54, p. 259, vol. 61, p. 198, vol. 63, p. 404, &c.

12. Dr. Hook's Hygrometer was made of the beard of a wild oat, set in a small box, with a dial plate and an index. See his *Micrographia*, p. 150.

13. The Doctors Hales and Defaguliers both contrived another form of sponge Hygrometer, on this principle. They made an horizontal axis, having a small



part of its length cylindrical, and the remainder tapering conically with a spiral thread cut in it, after the manner of the fuzee of a watch. The sponge is suspended by a fine silk thread to the cylindrical part of the axis, upon which it winds. This is balanced by a small weight W, suspended also by a thread, which

winds upon the spiral fuzee. Then when the sponge grows heavier, in moist weather; it descends and turns the axis, and so draws up the weight, which coming to a thicker part of the axis it becomes a balance to the sponge, and its motion is shewn by an attached scale. And vice versa when the air becomes drier.——Salt of tartar, or any other salt, or pot ashes, may be put into the scale of a balance, and used instead of the sponge. *Defag. Exper. Philos.* vol. 2, p. 300.

14. Mr. Ferguson made an Hygrometer of a thin deal pannel; and to enlarge the scale, and so render its variations more sensible, he employed a wheel and axle, making one cord pass over the axle, which turned a wheel ten times as large, over which passed a line with a weight at the end of it, whose motion was therefore ten times as much as that of the deal pannel. The board should be changed in 3 or 4 years. See *Philos. Transf.* vol. 54, art. 47.

15. Mr. Smeaton gave also an ingenious and elaborate construction of an Hygrometer; which may be seen in the *Philos. Transf.* vol. 61, art. 24.

16. Mr. De Luc's contrivance for an Hygrometer is very ingenious, and on this principle. Finding that even ivory swells with moisture, and contracts with dryness, he made a small and very thin hollow cylinder of ivory, open only at the upper end, into which is fitted the under or open end of a very fine long glass tube, like that of a thermometer. Into these is introduced some quicksilver, filling the ivory cylinder, and a small part of the length up the glass tube. The consequence is this: when moisture swells the ivory cylinder, its bore or capacity grows larger, and consequently the mercury sinks in the fine glass tube; and vice versa, when the air is drier, the ivory contracts, and forces the mercury higher up the tube of glass. It is evident that an instrument thus constructed is in fact also a thermometer, and must necessarily be affected by the vicissitudes of heat and cold, as well as by those of dryness and moisture; or that it must act as a thermometer as well as an Hygrometer. The ingenious contrivances in the structure and mounting of this instrument may be seen in the *Philos. Transf.* vol. 63, art. 38; where it may be seen how the above imperfection is corrected by some simple and ingenious expedients, employed in the original construction and subsequent use of the instrument; in consequence of which, the variations in the temperature of the air, though they produce their full effects on the instrument, as a thermometer, do not interfere with or embarrass its indications as an Hygrometer.

17. In the *Philos. Transf.* for 1791, Mr. De Luc has given a second paper on Hygrometry. This has been chiefly occasioned by a Memoir of M. de Saussure on the same subject, entitled *Essais sur l'Hygrometrie*, in 4to, 1783. In this work M. de S. describes a new Hygrometer of his construction, on the following principle. It is a known fact that a hair will stretch when it is moistened, and contract when dried: and M. de Saussure found, by repeated experiments, that the difference between the greatest extension and contraction, when the hair is properly prepared, and has a weight of about 3 grains suspended by it, is nearly one 40th of its whole length, or one inch in 40. This circumstance suggested the idea of a new Hygrometer. To



render these small variations of the length of the hair perceptible, an apparatus was contrived, in which one of the extremities of the hair is fixed, and the other, bearing the counterpoise abovementioned, surrounds the circumference of a cylinder, which turns upon an axis to which a hand is adapted, marking upon a dial in large divisions the almost insensible motion of this axis. About 12 inches high is recommended as the most convenient and useful: and to render them portable, a contrivance is added, by which the hand and the counterpoise can be occasionally fixed.

But M. de Luc, in his *Idées sur la Meteorologie*, vol. 1, anno 1786, shews that hairs, and all the other animal or vegetable hygroscopic substances, taken lengthwise, or in the direction of their fibres, undergo contrary changes from different variations of humidity; that when immersed in water, they lengthen at first, and afterwards shorten; that when they are near the greatest degree of humidity, if the moisture be increased, they shorten themselves; if it be diminished, they lengthen themselves first before they contract again. These irregularities, which render them incapable of being true measures of humidity, he shews to be the necessary consequence of their organic reticular structure. De Saussure takes his point of extreme moisture from the vapours of water under a glass bell, keeping the sides of the bell continually moistened; and affirms, that the humidity is, there, constantly the same in all temperatures; the vapours even of boiling water having no other effect than those of cold. De Luc, on the contrary, shews that the differences in humidity under the bell are very great, though De Saussure's Hygrometer was not capable of discovering them; and that the real undecomposed vapour of boiling water has the directly opposite effect to that of cold, the effect of extreme dryness; and on this point he mentions an interesting fact, communicated to him by Mr. Watt, viz, that wood cannot be employed in the steam engine, for any of those parts where the vapour of the boiling water is confined, because it dries so as to crack as if exposed to the fire.

To these charges of M. De Luc, a reply is made by M. De Saussure, in his *Defence of the Hair Hygrometer*, in 1788; where he attributes the general disagreement between the two instruments, to irregularities of M. De Luc's; and assigns some aberrations of his own Hygrometer, which could not have proceeded from the above cause, but to its having been out of order; &c.

This has drawn from M. De Luc a second paper on Hygrometry, published in the *Philos. Transf.* for 1791, p. 1, and 389. This author here resumes the four fundamental principles which he had sketched out in the former paper, viz, 1st, That fire is a sure, and the only sure means of obtaining extreme dryness. 2d, That water, in its liquid state, is a sure, and the only sure means of determining the point of extreme moisture. 3d, There is no reason, a priori, to expect, from any hygroscopic substance, that the measurable effects, produced in it by moisture, are proportional to the intensities of that cause.—But, 4th, perhaps the comparative changes of the dimensions of a substance, and of the weight of the same or other substances, by the same variations of moisture, may lead to some discovery in that respect. On these heads M. De Luc expatiates

at large in this paper, shewing the imperfections of M. De Saussure's principles of Hygrometry, and particularly as to a hair, or any such substance when extended lengthwise, being properly used as an Hygrometer. On the other hand, he shews that the expansion of substances across the fibres, or grain, renders them, in that respect, by far the most proper for this purpose. He chooses such as can be made very thin, as ivory, or deal shavings, but over all he finds whalebone to be far the best of any. But, for all the reasonings of these ingenious philosophers on this interesting subject, and complete information, see the publications above quoted, as also the *Monthly Review*, vol. 51, p. 224, vol. 71, p. 213, vol. 76, p. 316, vol. 78, p. 236, and vol. 6, of the new series for the year 1791, p. 133.

HYGROMETRY, the science of the measurement of the moisture of the atmosphere. The chief writings on this science are those of M. De Luc and M. De Saussure, for which see the last article on Hygrometers.

HYGROSCOPE, is commonly used in the same sense with Hygrometer. Wolfius, however, regarding the etymology of the word, makes some difference. According to him, the Hygroscope only shews the alterations of the air in respect of humidity and dryness, but the hygrometer measures them. A Hygroscope therefore is only an indefinite or less accurate hygrometer.

HYPATIA, a very learned and beautiful lady, was born at Alexandria, about the end of the 4th century, as she flourished about the year of Christ 430. She was the daughter of Theon, a celebrated philosopher and mathematician, and president of the famous Alexandrian school. Her father, encouraged by her extraordinary genius, had her not only educated in all the ordinary qualifications of her sex, but instructed in the most abstruse sciences. She made such great progress in philosophy, geometry, astronomy, and the mathematics in general, that she passed for the most learned person of her time. She published commentaries on Apollonius's Conics, on Diophantus's Arithmetic, and other works. At length she was thought worthy to succeed her father in that distinguished and important employment, the government of the school of Alexandria; and to deliver instructions out of that chair where Ammonius, Hierocles, and many other great men, had taught before; and this at a time too when men of great learning abounded both at Alexandria and in many other parts of the Roman empire. Her fame being so extensive, and her worth so universally acknowledged, it was no wonder that she had a crowded auditory. "She explained to her hearers (says Socrates, an ecclesiastical historian of the 5th century, born at Constantinople) the several sciences that go under the general name of philosophy; for which reason there was a confluence to her, from all parts, of those who made philosophy their delight and study."

Her scholars were not less eminent than they were numerous. One of them was the celebrated Synesius, who was afterwards bishop of Ptolomais. This ancient Christian Platonist always expresses the strongest, as well as the most grateful, testimony of the virtue of his tutors; never mentioning her without the most profound respect, and sometimes in terms of affection but little



little short of adoration. But it was not Synesius only, and the disciples of the Alexandrian school, who admired Hypatia for her virtue and learning: never was woman more careffed by the public, and yet never had woman a more unspotted character. She was held as an oracle for her wisdom, for which she was consulted by the magistrates in all important cases; a circumstance which often drew her among the greatest concourse of men, without the least censure of her manners. In short, when Nicephorus intended to pass the highest compliment on the princess Eudocia, he thought he could not do it better than by calling her another Hypatia.

While Hypatia thus reigned the brightest ornament of Alexandria, Orestes was governor of that place for the emperor Theodosius, and Cyril was bishop or patriarch. Orestes, having had a liberal education, could not but admire Hypatia; and as a wise governor often consulted her. This, together with an aversion which Cyril had against Orestes, proved fatal to the lady. About 500 monks assembling, attacked the governor one day, and would have killed him, had he not been rescued by the townsmen; and the respect which Orestes had for Hypatia causing her to be traduced among the Christian multitude, they dragged her from her chair, tore her in pieces, and burnt her limbs.

Cyril is strongly suspected of having fomented this tragedy. Cave indeed endeavours to remove the imputation of so horrid an action from the patriarch; and lays it upon the Alexandrian mob in general, whom he calls "a very trifling inconstant people." But though Cyril should be allowed neither to have been the perpetrator, nor even the contriver, of it, yet it is much to be suspected that he did not discountenance it in the manner he ought to have done: a suspicion which must needs be greatly confirmed by reflecting, that he was so far from blaming the outrage committed by the monks upon Orestes, that he afterwards received the dead body of Ammonius, one of the most forward in that outrage, who had grievously wounded the governor, and who was justly punished with death. Upon this riotous ruffian Cyril made a panegyric in the church where he was laid, in which he extolled his courage and constancy, as one that had contended for the truth; and changing his name to Thaumastus, or the Admirable, ordered him to be considered as a martyr. "However (continues Socrates), the wisest part of Christians did not approve the zeal which Cyril shewed on this man's behalf, being convinced that Ammonius had justly suffered for his desperate attempt."

**HYPERBOLA**, one of the conic sections, being that which is made by a plane cutting a cone so, that, entering one side of the cone, and not being parallel to the opposite side, it may cut the circular base when the opposite side is ever so far produced below the vertex, or shall cut the opposite side of the cone produced above the vertex, or shall make a greater angle with the base than the opposite side of the cone makes; all these three circumstances amounting to the same thing, but in other words.

1. Thus, the figure DAE is an Hyperbola, made by a plane entering the side VQ of a cone PVQ at A, and either cutting the base PEQ when the plane is not parallel to VP, and this is ever so far produced; or when



the angle ARQ is greater than the angle VPQ; or when the plane cuts the opposite side in B above the vertex.

2. By the Hyperbola is sometimes meant the whole plane of the section, and sometimes only the curve line of the section.

3. Hence, the cutting plane meets the opposite cone in B, and there forms another Hyperbola  $dBe$ , equal to the former one, and having the same transverse axis AB; and the same vertices A and B. Also the two are called Opposite Hyperbolas.

4. The centre C is the middle point of the transverse axis.

5. The semi-conjugate axis is CL, a mean proportional between CI and CK, the distances to the sides of the opposite cone, when CI is drawn parallel to the diameter PQ of the base of the cone. Or the whole conjugate axis is a mean proportional between AF and BH, which are drawn parallel to the base of the cone.




6. If DAE and FBG be two opposite Hyperbolas, having the same transverse and conjugate axes AB and  $ab$ , perpendicularly bisecting each other; and if  $d a e$  and  $f b g$  be two other opposite Hyperbolas, having the same axes with the two former, but in the contrary order, viz, having  $ab$  for their first or transverse axis, and AB for their second or conjugate axis: then any two adjacent curves are called Conjugate Hyperbolas, and the whole figure formed by all the four curves, the Figure of the Conjugate Hyperbolas. And if the rectangle HIKL be inscribed within the four conjugate Hyperbolas, touching the vertices A, B,  $a$ ,  $b$ , and having their sides parallel and equal to the two axes; and if then the two diagonals HCK, ICL, of the parallelogram be drawn, these diagonals are the asymptotes of the curves, being lines that continually approach nearer and nearer to the curves, without meeting them, except at an infinite distance, where each asymptote and the two adjacent sides of the two conjugate Hyperbolas may be supposed all to meet; the asymptote being there a common tangent to them both, viz, at that infinite distance.

7. Hence the four Hyperbolas, meeting and running into each other at the infinite distance, may be considered as the four parts of one entire curve, having the same axes, tangents, and other properties.

8. A Diameter in general, is any line, as MN, drawn through the centre C, and meeting, or terminate



nated by the opposite legs of the opposite Hyperbolas. And if parallel to this diameter there be drawn two tangents, at  $m$  and  $n$ , to the opposite legs of the other two opposite Hyperbolas, the line  $mCn$  joining the points of contact, is the conjugate diameter to  $MN$ , and the two mutually conjugates to each other. Or, if to the points  $M$  or  $N$  there be drawn a tangent, and through the centre  $C$  the line  $mn$  parallel to it, that line will be the conjugate to  $MN$ . The points where each of these meet the curves, as  $M$ ,  $N$ ,  $m$ ,  $n$ , are the vertices of the diameters; and the tangents to the curves at the two vertices of any diameter, are parallel to each other, and also to the other or conjugate diameter.



9. Moreover, if those tangents to the four Hyperbolas, at the vertices of two conjugate diameters, be produced till they meet, they will form a parallelogram  $OPQR$ ; and the diagonals  $OQ$  and  $PR$  of the parallelogram will be the asymptotes of the curves; which therefore pass through the opposite angles of all the parallelograms so inscribed between the curves. Also it is a property of these parallelograms, that they are all equal to each other, and therefore equal to the rectangle of the two axes; as will be farther noticed below. Farther, if these diagonals or asymptotes make a right angle between them, or if the inscribed parallelogram be a square, or if the two axes be equal to each other, then the Hyperbola is called a right-angled or an equilateral one.

10. An Ordinate to any diameter, is a line drawn parallel to its conjugate, or to the tangent at its vertex, and terminated by the diameter produced and the curve. So  $MS$  and  $TN$  are ordinates to the axis  $AB$ ; also  $AD$  and  $BG$  are ordinates to the diameter  $MN$  (last fig. but one). Hence the ordinates to the axis are perpendicular to it; but ordinates to the other diameters are oblique to them.

11. An absciss is a part of any diameter, contained between its vertex and an ordinate to it; and every ordinate has two abscisses: as  $AT$  and  $BT$ , or  $MV$  and  $NV$ .

12. The Parameter of any diameter, is a third proportional to the diameter and its conjugate.—The Parameter of the axis is also equal to the line  $AG$  or  $Bg$  (fig. 1), if  $FG$  be drawn to make the angle  $AFG =$  the angle  $BAV$ , or the line  $Hg$  to make the angle  $BHg =$  the angle  $ABV$ .


13. The Focus is the point in the axis where the ordinate is equal to half the parameter of the axis; as

VOL. I.

$S$  and  $T$  (fig. 2) if  $MS$  and  $TN$  be half the parameter, or the 3d proportional to  $CA$  and  $Ca$ . Hence there are two Foci, one on each side the vertex, or one for each of the opposite Hyperbolas. These two points in the axis are called Foci, or burning points, because it is found by opticians that rays of light issuing from one of them, and falling upon the curve of the Hyperbola, are reflected into lines that verge towards the other point or Focus.


To describe an Hyperbola, in various ways.

14. (1st Way by points.)—In the transverse axis  $AF$  produced, take the foci  $F$  and  $f$ , by making  $CF$  and  $Cf = Aa$  or  $Ba$ , assume any point  $I$ : Then with the



radii  $AI$ ,  $BI$ , and centres  $F$ ,  $f$ , describe arcs intersecting in  $E$ , which will give four points in the curves. In like manner, assuming other points  $I$ , as many other points will be found in the curve. Then, with a steady hand, draw the curve line through all the points of intersection  $E$ .—In the same manner are to be constructed the other pair of opposite Hyperbolas, using the axis  $ab$  instead of  $AB$ .

15. (2d Way by points, for a Right-angled Hyperbola only.)—On the axis produced if necessary, take any



point  $I$ , through which draw a perpendicular line, upon which set off  $IM$  and  $IN$  equal to the distance  $Ia$  or  $Ib$  from  $I$  to the extremities of the other axis; and  $M$  and  $N$  will be points in the curve.


16. (3d Way by points, to describe the curve through a given point.)— $CG$  and  $CH$  being the asymptotes, and  $P$  the given point of the curve; through the point  $P$  draw any line  $GPH$  between the asymptotes, upon which take  $GI = PH$ , so shall  $I$  be another point of the curve. And in this manner may any number of points be found, drawing as many lines through the given point  $P$ .

4 K.

17. (4th




17. (4th Way by a continued Motion.)—If one end



of a long ruler  $fMO$  be fastened at the point  $f$ , by a pin on a plane, so as to turn freely about that point as a centre. Then take a thread  $FMO$ , shorter than the ruler, and fix one end of it in  $F$ , and the other to the end  $O$  of the ruler. Then if the ruler  $fMO$  be turned about the fixed point  $f$ , at the same time keeping the thread  $OMF$  always tight, and its part  $MO$  close to the side of the ruler, by means of the pin  $M$ ; the curve line  $AX$  described by the motion of the pin  $M$  is one part of an Hyperbola. And if the ruler be turned, and move on the other side of the fixed point  $F$ , the other part  $AZ$  of the same Hyperbola may be described after the same manner.—But if the end of the ruler be fixed in  $F$ , and that of the thread in  $f$ , the opposite Hyperbola  $xaz$  may be described.

18. (5th Way, by a continued Motion.)—Let  $C$  and  $F$  be the two foci, and  $E$  and  $K$  the two vertices of the Hyperbola. (See the last fig. above.) Take three rulers  $CD$ ,  $DG$ ,  $GF$ , so that  $CD = GF = EK$ , and  $DG = CF$ ; the rulers  $CD$  and  $GF$  being of an indefinite length beyond  $C$  and  $G$ , and having slits in them for a pin to move in; and the rulers having holes in them at  $C$  and  $F$ , to fasten them to the foci  $C$  and  $F$  by means of pins, and at the points  $D$  and  $G$  they are to be joined by the ruler  $DG$ . Then, if a pin be put in the slits, viz, the common intersection of the rulers  $CD$  and  $GF$ , and moved along, causing the two rulers  $GF$ ,  $CD$ , to turn about the foci  $C$  and  $F$ , that pin will describe the portion  $Ee$  of an Hyperbola.—The foregoing are a few among various ways given by several authors.

*Some of the chief Properties of the Hyperbola.*



19. (1st) The squares of the ordinates, of any diameter, are to each other, as the rectangles of their

abscisses; i. e.  $DE^2 : GH^2 :: AD \cdot BD : AG \cdot BG$ .

20. As the square of any diameter, is to the square of its conjugate; so is the rectangle of two abscisses, to the square of their ordinate. That is,  $AB^2 : ab^2 :: AD \cdot BD : DE^2$ .

Or, because the rectangle  $AD \cdot BD$  is = the difference of the squares  $CD^2 - CB^2$ , the same property is,

$$\text{As } AB^2 : ab^2 :: CD^2 - CB^2 : DE^2,$$

$$\text{Or } AB : \frac{ab^2}{AB} :: CD^2 - CB^2 : DE^2,$$

That is  $AB : p :: CD^2 - CB^2 : DE^2$ , where  $p$  is the parameter of the diameter  $AB$ , or the 3d proportional  $\frac{ab^2}{AB}$ .

And hence is deduced the common equation of the Hyperbola, by which its general nature is expressed. Thus, putting  $d$  = the semidiameter  $CA$  or  $CB$ ,

$c$  = its semiconjugate  $Ca$  or  $Cb$ ,

$p$  = its parameter or  $\frac{2d^2}{c}$ ,

$x$  = the absciss  $BD$  from the vertex,

$y$  = the ordinate  $DE$ , and

$v$  = the absciss  $CD$  from the centre :

Then is

$$d^2 : c^2 :: \frac{2d + x}{d} \cdot x : y^2,$$

$$\text{or } d^2 : c^2 :: d + v \cdot x : y^2,$$

$$\text{or } d^2 : c^2 :: v^2 - d^2 : y^2,$$

$$\text{or } d : p :: v^2 - d^2 : y^2; \text{ so that}$$

$$y^2 = \frac{2dx + x^2}{d^2} \cdot c^2 = \frac{v^2 - d^2}{d^2} \cdot c^2 = \frac{v^2 - d^2}{d} \cdot p;$$

any of which equations or proportions express the nature of the curve. And hence arises the name Hyperbola, signifying to exceed, because the ratio of  $d^2$  to  $c^2$ , or of  $d$  to  $p$ , exceeds that of  $2dx$  to  $y^2$ ; that ratio being equal in the parabola, and defective in the ellipse, from which circumstances also these take their names.

21. The distance between the centre and the focus, is equal to the distance between the extremities of the transverse and conjugate axes. That is,  $CF = Aa$  or  $Ab$ , where  $F$  is the focus.

22. The conjugate semi-axis is a mean proportional between the distances of the focus from both vertices of the transverse. That is,  $Ca$  is a mean between  $AF$  and  $BF$ , or  $AF : Ca :: Ca : BF$ , or  $AF \cdot BF = Ca^2$ .

23. The difference of two lines drawn from the foci, to meet in any point of the curve, is equal to the transverse axis. That is,  $fE - FE = AB$ , where  $F$  and  $f$  are the two foci.


24. All the parallelograms inscribed between the four conjugate Hyperbolas are equal to one another, and each equal to the rectangle of the two axes. That is, the parallelogram  $OPQR = AB \cdot ab$  (fig. to art. 9).

25. The difference of the squares of every pair of conjugate diameters, is equal to the same constant quantity, viz, the difference of the squares of the two axes. That is,  $MN^2 - mn^2 = AB^2 - ab^2$ , (fig. to art. 6); where  $MN$  and  $mn$  are any two conjugate diameters.



26. The rectangles of the parts of two parallel lines, terminated by the curve, are to one another, as the rectangles of the parts of any other two parallel lines, any where cutting the former. Or the rectangles of the parts of two intersecting lines, are as the squares of their parallel diameters, or squares of their parallel tangents.

27. All the rectangles are equal which are made of the segments of any parallel lines, cut by the curve, and limited by the asymptotes, and each equal to the square of their parallel diameter. That is,  $HE \cdot EK$  or  $He \cdot eK = CQ^2$  or  $CP^2$ .



28. All the parallelograms are equal, which are formed between the asymptotes and curve, by lines parallel to the asymptotes. That is, the paral.  $CGEK = CPBQ$ .—Hence is obtained another method of expressing the nature of the curve by an equation, involving the absciss taken on one asymptote, and ordinate parallel to the other asymptote. Thus, if  $x = CK$ ,  $y = KE$ ,  $a = CQ$ , and  $b = BQ$  the ordinate at the vertex B of the curve; then, by the property in this article,  $ab = xy$ , or  $a : x :: y : b$ ; that is, the rectangle of the absciss and ordinate is every where of the same magnitude, or any ordinate is reciprocally as its absciss.

29. If the absciss  $CQ, CK, CL$ , &c, taken on the one asymptote, be in geometrical progression increasing; then shall the ordinates  $QB, KE, LM$ , &c, parallel to the other asymptote, be a like geometrical progression in the same ratio, but decreasing; and all the rectangles are equal, under every absciss and its ordinate, viz,  $CQ \cdot QB = CK \cdot KE = CL \cdot LM$ , &c.

30. The absciss  $CQ, CK, CL$ , &c, being taken in geometrical progression; the spaces or asymptotic areas  $BQKE, EKLM$ , &c, will be all equal; or, the spaces  $BQKE, BQLM$ , &c, will be in arithmetical progression; and therefore these spaces are the hyperbolic logarithms of those absciss.

These, and many other curious properties of the Hyperbola, may be seen demonstrated in my Treatise on Conic Sections, and several others. See also CONIC SECTIONS.

*Acute* HYPERBOLA, one whose asymptotes make an acute angle.

*Ambigenal* HYPERBOLA, is that which has one of its infinite legs falling within an angle formed by the asymptotes, and the other falling without that angle. This is one of Newton's triple Hyperbolas of the 2d

order. See his *Enumeratio Lin. tert. Ord.* See also AMBIGENAL.

*Common*, or *Conic* HYPERBOLA, is that which arises from the section of a cone by a plane; called also the Apollonian Hyperbola, being that kind treated on by the first and chief author Apollonius.

*Conjugate* HYPERBOLAS, are those formed or lying together, and having the same axes, but in a contrary order, viz, the transverse of each equal the conjugate of the other; as the two Conjugate Hyperbolas  $Pee$  and  $EEe$  in the last figure but one.

*Equilateral*, or *Rectangular* HYPERBOLA, is that whose two axes are equal to each other, or whose asymptotes make a right angle.—Hence, the property or equation of the equilateral Hyperbola, is  $y^2 = ax + x^2$ , where  $a$  is the axis,  $x$  the absciss, and  $y$  its ordinate; which is similar to the equation of the circle, viz,  $y^2 = ax - x^2$ , differing only in the sign of the second term, and where  $a$  is the diameter of the circle.

*Infinite* HYPERBOLAS, or HYPERBOLAS of the higher kinds, are expressed or defined by general equations similar to that of the conic or common Hyperbola, but having general exponents, instead of the particular numeral ones, but so as that the sum of those on one side of the equation, is equal to the sum of those on the other side. Such as,  $ay^{m+n} = bx^m(d+x)^n$ , where  $x$  and  $y$  are the absciss and ordinate to the axis or diameter of the curve; or  $x^m y^n = a^{m+n}$ , where the absciss  $x$  is taken on one asymptote, and the ordinate  $y$  parallel to the other.


As the Hyperbola of the first kind, or order, viz the conic Hyperbola, has two asymptotes; that of the 2d kind or order has three; that of the 3d kind, four; and so on.

*Obtuse* HYPERBOLA, is that whose asymptotes form an obtuse angle.

*Rectangular* HYPERBOLA, the same as Equilateral Hyperbola.

HYPERBOLIC Arc, is the arc of an Hyperbola.

Put  $a = CA$  the semitransverse axe,  $c = Ca$  the semiconjugate,  $y =$  an ordinate  $PQ$  to the axe drawn from the end  $Q$  of the arc  $AQ$ , beginning at the vertex



$A$ : then putting  $q = \frac{aa + cc}{c^2}$ ,

$A =$  the hyp. log. of  $\frac{y + \sqrt{cc + yy}}{c}$

$= 2.302585093 \times \text{common log. of } \frac{y + \sqrt{cc + yy}}{c}$ ,

$B = \frac{y\sqrt{cc + yy} - ccA}{2}$ ,  $C = \frac{y^3\sqrt{cc + yy} - 3ccB}{4}$ ,

$D = \frac{y^5\sqrt{cc + yy} - 5ccC}{6}$ , &c;

then is the length of the arc  $AQ$  expressed by

$c \times (A + \frac{q}{2} B - \frac{q^2}{2.4} C + \frac{1.3q^3}{2.4.6} D - \frac{1.3.5q^4}{2.4.6.8} E \&c;$   
4 K 2 or



or by  $\frac{15c^2t + (21c^2 + 19t^2)x}{15c^2t + (21c^2 + 9t^2)x} \times y$ , nearly; where  $t$  is the whole transverse axe  $2CA$ ,  $c = 2Ca$  the conjugate,  $x = AP$  the absciss, and  $y = PQ$  the ordinate.

These and other rules may be seen demonstrated in my *Mensuration*, p. 408, &c, 2d edit.

**HYPERBOLIC Area, or Space**, the area or space included by the Hyperbolic curve and other lines.

Putting  $a = CA$  the semitransverse,  $c = Ca$  the semiconjugate,  $y = PQ$  the ordinate, and  $v = CP$  its distance from the centre; then is the

$$\text{area } APQ = \frac{1}{2}vy - \frac{1}{2}ac \times \text{hyp. log. of } \frac{ay + cv}{ac};$$

$$\text{sector } CAQ = \frac{1}{2}ac \times \text{hyp. log. of } \frac{ay + cv}{ac};$$

$$\text{area } APQ = 2xy \times \left( \frac{1}{3} - \frac{z^1}{1.3.5} - \frac{z^2}{3.5.7} - \frac{z^3}{5.7.9} - \frac{z^4}{7.9.11} \text{ \&c} \right),$$

$$\text{where } x = AP, \text{ and } z = \frac{x}{2a + x}; \text{ or}$$

$$APQ = \frac{2cx}{15a} \times (\sqrt{2ax} + 4\sqrt{2ax + \frac{3}{4}xx}) \text{ nearly.}$$

Let  $CT$  and  $CE$  be the two asymptotes, and the ordinates  $DA$ ,  $EF$  parallel to the other asymptote  $CT$ ; then the asymptotic space  $ADEF$  or sector  $CAF$  is

$$= CD \times DA \times \text{hyp. log. of } \frac{CE}{CD} \text{ or}$$

$$= CD \times DA \times \text{hyp. log. } \frac{DA}{EF} \text{ or}$$

$$= CD \times DA \times \frac{DE}{CD} - \frac{DE^2}{2CD^2} + \frac{DE^3}{3CD^3} - \frac{DE^4}{4CD^4} \text{ \&c};$$

and this last series was first given by Mercator in his *Logarithmotechnia*.

See my *Mensuration*, p. 413, &c, 2d edit.



Generally, if  $x^m y^n = a^{m+n}$  be an equation expressing an Hyperbola of any order; then its asymptotic area will be  $\frac{n}{n-m} xy$ ; which space therefore is always

quadrable, in all the orders of Hyperbolas, except the first or common Hyperbola only, in which  $m$  and  $n$  being each 1, the denominator  $n - m$  becomes 0 or nothing.

**HYPERBOLIC Conoid**, a solid formed by the revolution of an Hyperbola about its axis, otherwise called an Hyperboloid.

*To find the Solid Content of an Hyperboloid.*

Let  $AC$  be the semitransverse of the generating Hyperbola, and  $AH$  the height of the solid; then as  $2AC + AH$  is to  $3AC + AH$ , so is the cone of the same base and altitude, to the content of the Conoid.



*To find the Curve Surface of an Hyperboloid.*

Let  $AC$  be the semitransverse, and  $AB$  perpendicular to it, and equal to the semiconjugate of  $ADE$  the generating Hyperbola, or section through the axis of the solid. Join  $CB$ ; make  $CF = CA$ , and on  $CA$  let fall the perpendicular  $FG$ ; then with the semitransverse  $CG$ , and semiconjugate  $GH = AB$ , describe the Hyperbola  $GIK$ ; then as the diameter of a circle is to its circumference, so is the Hyperbolic frustum  $ILAMK$  to the curve surface of the Conoid generated by  $DAE$ . See my *Mensur.* p. 429, &c, 2d edit.

**HYPERBOLIC Cylindroid**, a solid formed by the revolution of an Hyperbola about its conjugate axis, or line through the centre perpendicular to the transverse axis. This solid is treated of in the *Philos. Trans.* by Sir Christopher Wren, where he shews some of its properties, and applies it to the grinding of Hyperbolic Glasses; affirming that they must be formed this way, or not at all. See *Philos. Trans.* vol. 4, pa. 961.

**HYPERBOLIC Leg**, of a curve, is that having an asymptote, or tangent at an infinite distance.—Newton reduces all curves, both of the first and higher kinds, into Hyperbolic and parabolic legs, i. e. such as have asymptotes, and such as have not, or such as have tangents at an infinite distance, and such as have not.

**HYPERBOLIC Line**, is used by some authors for what is more commonly called the Hyperbola itself, being the curve line of that figure; in which sense the surface terminated by it is called the Hyperbola.

**HYPERBOLIC Logarithm**, a logarithm so called as being similar to the asymptotic spaces of the Hyperbola. The Hyperbolic logarithm of a number, is to the common logarithm, as 2.3025850929940457 to 1, or as 1 to .4342944819032518. The first invented logarithms, by Napier, are of the Hyperbolic kind; and so are Kepler's. See **LOGARITHM**.

**HYPERBOLIC Mirror**, is one ground into that shape.

**HYPERBOLIC Space**, that contained by the curve of the Hyperbola, and certain other lines. See **HYPERBOLIC AREA**.

**HYPERBOLICUM Acutum**, a solid made by the revolution of the infinite area or space contained between the curve of the Hyperbola, and its asymptote. This produces a solid, which though infinitely long and generated by an infinite area, is nevertheless equal to a finite solid body; as is demonstrated by Torricelli, who gave it this name.

**HYPERBOLIFORM Figures**, are such curves as approach, in their properties, to the nature of the Hyperbola; called also Hyperboloides.

**HYPERBOLOIDS**, are Hyperbolas of the higher kind, whose nature is expressed by this equation,  $ay^{m+n} = bx^m(a+x)^n$ . See **HYPERBOLA**. It also means the Hyperbolic Conoid. See that article.

**HYPERBOREANS**, the most northern nations, or regions, as dwelling beyond or about the wind Boreas: as the Siberians, Samoeds, &c.

**HYPERTHYRON**, in Architecture, a sort of table, usually placed over gates or doors of the Doric order, above the chambranle, in form of a frieze.

**HYPETHRE**, in Ancient Architecture, two rows of




of pillars surrounding, and ten at each face of a temple, &c, with a peristyle within of six columns.

**HYPOGEUM**, in the ancient Architecture, a name common to all the parts of a building that are under ground; as the cellars, butteries, &c.

**HYPOGEUM**, in Astrology, a name given to the celestial houses that are below the horizon; and especially the *inimæ cœli*, or bottom point of the heavens.

**HYPOMOCHLION**, the fulcrum or prop of a lever; or the point which sustains its pressure, when employed either in raising or lowering bodies. The Hypomochlion is frequently a roller set under the lever; or under stones or pieces of timber, &c, that they may be the more easily lifted up, or removed.

**HYPOTENUSE**, or **HYPOTHENUSE**, in a right-angled triangle, is the side which subtends, or is opposite to the right angle, and is always the longest of the three sides; as the side AC, opposite to the right angle B.



It is a celebrated theorem in Plane Geometry, being the 47th prop. of the 1st book of Euclid, that in every right-angled triangle ABC, the square formed upon the Hypotenuse AC, is equal to both the two squares formed upon the other two sides AB and BC; or that  $AC^2 = AB^2 + BC^2$ . This is particularly called the Pythagorean theorem, from its reputed inventor Pythagoras, who it is said sacrificed a whole hecatomb to the muses, in gratitude for the discovery. But the same thing is true of circles or any other similar figures, viz, that any figure described on the Hypotenuse, is equal to the sum of the two similar figures described on both the other two sides.

**HYPOTHENUSE**. See **HYPOTENUSE**.

**HYPOTHESIS**, in Geometry, or Mathematics, means much the same thing with supposition, being a

supposition or an assumption of something as a condition, upon which to raise a demonstration, or from which to draw an inference.

Dr. Barrow says, Hypotheses, or postulatus, are propositions assuming or affirming some evidently possible mode, action, or motion of a thing, and that there is the same affinity between hypotheses and problems, as between axioms and theorems: a problem shewing the manner, and demonstrating the possibility of some structure, and an Hypothesis assuming some construction which is manifestly possible.

**HYPOTHESIS**, in Philosophy, denotes a kind of system laid down from our own imagination, by which to account for some phenomenon or appearance of nature. Thus there are Hypotheses to account for the tides, for gravity, for magnetism, for the deluge, &c.

The real and scientific causes of natural things generally lie very deep: observation and experiment, the proper means of arriving at them, are in most cases extremely slow; and the human mind is very impatient: hence we are often induced to feign or invent something that may seem like the cause, and which is calculated to answer the several phenomena, so that it may possibly be the true cause.

Philosophers are divided as to the use of such fictions or Hypotheses, which are much less current now than they were formerly. The latest and best writers are for excluding Hypotheses, and standing intirely on observation and experiment. Whatever is not deduced from phenomena, says Newton, is an Hypothesis; and Hypotheses, whether metaphysical, or physical, or mechanical, or of occult qualities, have no place in experimental philosophy. *Phil. Nat. Prin. Math. in Calce.*

**HYPOTHESIS** is more particularly applied, in Astronomy, to the several systems of the heavens; or the divers manners in which different astronomers have supposed the heavenly bodies to be ranged, or moved. The principal Hypotheses are the Ptolomaic, the Tychonic, and the Copernican. This last is now so generally received, and so well established and warranted by observation, that it is thought derogatory to it to call it an Hypothesis.

**HYPOTRACHELION**, in Architecture, is used for a little frieze in the Tuscan and Doric capital, between the astragal and annulets; called also the *colerion* and *gorgerin*.

The word is applied by some authors in a more general sense, to the neck of any column, or that part of its capital below the astragal.



## I. J.


## J A C

**JACK**, in Mechanics, is an instrument in common use for raising heavy timber, or very great weights of any kind; being a certain very powerful combination of teeth and pinions, and the whole inclosed in a strong wooden stock or frame BC, and moved by a winch or handle HP; the outside appearing as in fig. 1, here annexed.

In fig. 6, pl. 12, the wheel or rack-work is shewn, being the view of the inside when the stock is removed. Though it is not drawn in the just proportions and dimensions, for the rack AB must be supposed at least four times as long in proportion to the wheel Q, as the figure represents it; and the teeth, which will be then four times more in number to have about 3 in the inch. Now if the handle HP be 7 inches long, the circumference of this radius will be 44 inches, which is the distance or space the power moves through in one revolution of the handle: but as the pinion of the handle has but 4 leaves, and the wheel Q suppose 20 teeth, or 5 times the number, therefore to make one revolution of the wheel Q, it requires 5 turns of the handle, in which case it passes through 5 times 44 or 220 inches: but the wheel having a pinion R of 3 leaves, these will raise the rack 3 teeth, or one inch, in the same space. Hence then, the handle or power moving 220 times as fast as the weight, will raise or balance a weight of 220 times its own power. And if this be the hand of a man, who can sustain 100 pounds weight, he will, by help of this Jack, be able to raise, or sustain a weight or force of 22000lb, or about 10 tons weight.

This machine is sometimes open behind from the bottom almost up to the wheel Q, to let the lower claw, which in that case is turned up as at B, draw up any weight. When the weight is drawn or pushed sufficiently high, it is kept from going back by hanging the end of the hook S, fixed to a staple, over the curved part of the handle at h.

**JACK** is also the name of a well-known engine in the kitchen, used for turning a spit. Here the weight is the power applied, acting by a set of pulleys; the friction of the parts, and the weight with which the



spit is charged, make the force to be overcome; and a steady uniform motion is maintained by means of the fly.

See the fig. of this machine, pl. 12, fig. 7.

*Smoke JACK*, is an engine used for the same purpose with the common Jack, and is so called from its being moved by means of the smoke, or rarefied air, ascending the chimney, and striking against the sails of the horizontal wheel AB (plate 12, fig. 8), which being inclined to the horizon, is moved about the axis of the wheel, together with the pinion C, which carries the wheels D and E; and E carries the chain F, which turns the spit. The wheel AB should be placed in the narrow part of the chimney, where the motion of the smoke is swiftest, and where also the greatest part of it must strike upon the sails.—The force of this machine depends upon the draught of the chimney, and the strength of the fire.

*JACK-arch*, in Architecture, is an arch of one brick thickness.

*JACK-head*, in Hydraulics, a part sometimes annexed to the forcing pump.

*JACOB's-Staff*, a mathematical instrument for taking heights and distances; the same with the Cross-staff; which see.

**JACOBUS**, a gold coin, worth 25 shillings; so called from king James the first of England, in whose reign it was struck. They distinguished two kinds of the Jacobus, the old and the new; the former valued at 25 shillings, weighing 6 dwts 10 grs; the latter, called also Carolus, valued at 23 shillings, and weighing 5 dwts 20 grains.

**JAMBS**, or **JAUMS**, in Architecture, are the upright sides of chimneys, from the hearth to the mantel-tree. Also door posts, or the upright posts at the ends of the window frames.

*St JAMES's Day*, a festival in the calendar, observed on the 25th of July, in honour of St. James the apostle.

**JANUARY**, the first month of the year, according to the computation now used in the West, and containing 31 days; so called by the Romans from Janus, one of their divinities, to whom they gave two faces; because on the one side, the first day of this month looked towards the new year, and on the other towards the old one. The name may also be derived from Janua, a gate; this month, being the first of the year, may be considered as the gate or entrance of it.

January



January and February were introduced into the year by Numa Pompilius; Romulus's year beginning with the month of March.

JAUMS. See JAMBS.

ICE, a brittle transparent body, formed of some fluid, frozen or fixed by cold. The specific gravity of Ice to water, is various, according to the nature and circumstances of the water, degree of cold, &c. Dr. Irving (Phipps's Voyage towards the North Pole) found the densest Ice he could meet with about a 14th part lighter than water. M. de Mairan found it, at different trials, 1-14th, 18th, or 19th lighter than water; and when the water was previously purged of air, only a 22d part.

The rarefaction of Ice has been supposed owing to the air-bubbles produced in Ice while freezing; these, being considerably large in proportion to the water frozen, render the Ice so much specifically lighter. It is well known that a considerable quantity of air is lodged in the interstices of water, though it has there little or no elastic property, on account of the disunion of its particles; but upon these particles coming closer together, and uniting as the water freezes, light, expansive, and elastic air-bubbles are thus generated, and increase in bulk as the cold grows stronger, and by their elastic force bursts to pieces any vessel in which the water is closely contained. But snow-water, or any water long boiled over the fire, affords an Ice more solid, and with fewer bubbles. Pure water long kept in vacuo and frozen afterwards there, freezes much sooner, on being exposed to the same degree of cold, than water unpurged of its air and set in the open atmosphere. And the Ice made of water thus divested of its air, is much harder, more solid and transparent, and heavier than common Ice.

But M. de Mairan, in a dissertation on Ice, attributes the increase of the bulk of the water under this form, chiefly to a different arrangement of its parts: the icy skin on water being composed of filaments which are found to be joined constantly and regularly at an angle of  $60^\circ$ , and which, by this disposition, occupy a greater volume than if they were parallel. Besides, after Ice is formed, he found it continue to expand by cold; a piece of Ice, which was at first only a 14th part specifically lighter than water, on being exposed some days to the frost, became a 12th part lighter; and thus he accounts for the bursting of Ice in ponds.

It appears from an experiment of Dr. Hooke, in 1663, that Ice refracts the light less than water; whence he infers, that the lightness of Ice, which causes it to swim in water, is not produced merely by the small bubbles which are visible in it, but that it arises from the uniform constitution or general texture of the whole mass: a fact which was afterward confirmed by M. de la Hire. See Hooke's Exper. by Derham, p. 26, Acad. Per. 1693, Mem. p. 25.

Sir Robert Barker thus describes the process of making Ice in the East Indies, in a country where he never saw any natural Ice. On a large plain they dig three or four pits, each about 30 feet square, and 2 feet deep; the bottoms of which are covered, about 8 or 12 inches thick, with sugar-cane, or the stems of the

large Indian corn, dried. On this bed are placed in rows a number of small shallow unglazed earthen pans, formed of a very porous earth, a quarter of an inch thick, and about an inch and a quarter deep; which, at the dusk of the evening, they fill with soft water that has been boiled. In the morning before sunrise the Ice-makers attend at the pits, and collect what has been frozen in baskets, which they convey to the place of preservation. This is usually prepared in some high and dry situation, by sinking a pit 14 or 15 feet deep, which they line first with straw, and then with a coarse kind of blanketing. The Ice is deposited in this pit, and beaten down with rammers, till at length its own accumulated cold again freezes it, and it forms one solid mass. The mouth of the pit is well secured from the exterior air with straw and blankets, and a thatched roof is thrown over the whole. Philos. Trans. vol. 65, p. 252.

ICHOGRAPHY, in Architecture, is a transverse or horizontal section of a building, exhibiting the plot of the whole edifice, and of the several rooms and apartments in any story; together with the thickness of the walls and partitions; the dimensions of the doors, windows, and chimneys; the projectures of the columns and piers, with every thing visible in such a section.

ICHOGRAPHY, in Fortification, is the plan or representation of the length and breadth of a fortress; the distinct parts of which are marked out, either on the ground itself, or upon paper.

ICHOGRAPHY, in Perspective, the view of any thing cut off by a plane parallel to the horizon, just by the base or bottom of it; being the same with what is otherwise called the plan, geometrical plan, or ground-plot, of any thing, and is opposed to Orthography or Elevation.

ICOSAEDRON, or ICOSAHEDRON, one of the five regular bodies or solids, terminated by twenty equilateral and equal triangles. It may be considered as consisting of 20 equal and similar triangular pyramids, whose vertices meet in the centre of a sphere conceived to circumscribe it, and therefore having all their heights and bases equal; therefore the solidity of one of those pyramids multiplied by 20, the number of them, gives the solid content of the Icosaedron.

To form or make the Icosaedron.—Describe upon a card paper, or some other such like substance, 20 equilateral triangles, as in the figure at the article *Regular Body*. Cut it out by the extreme edges, and cut all the other lines half through, then fold the sides up by these edges half cut through, and the solid will be formed.

The linear edge or side of the Icosaedron being  $A$ , then will the surface be  $5A^2\sqrt{3} = 8.6602540A^2$ , and the solidity =

$$\frac{5}{8}A^3\sqrt{\frac{7+3\sqrt{5}}{2}} = 2.1816950A^3.$$

More generally, put  $A$  = the linear edge or side,  $B$  the surface, and  $C$  the solid content of the Icosaedron, also  $r$  the radius of the inscribed, and  $R$  the radius of the circumscribing sphere, then we have these general equations, viz,



$$1^{\text{st}}, A = r\sqrt{42-18\sqrt{5}} = R\sqrt{\frac{10-2\sqrt{5}}{5}} =$$

$$\sqrt{\frac{B\sqrt{3}}{15}} = \sqrt[3]{\frac{6}{5}}C\sqrt{\frac{7-3\sqrt{5}}{2}}.$$

$$2^{\text{d}}, B = 3r^2 \times 7\sqrt{3} - 3\sqrt{15} = 2R^2 \times 5\sqrt{3} - \sqrt{15}$$

$$= 5A^2\sqrt{3} = 3\sqrt[3]{70\sqrt{3} - 30\sqrt{15}}.$$

$$3^{\text{d}}, C = 10r^3 \times 7\sqrt{3} - 3\sqrt{15} = \frac{2}{3}R^3\sqrt{10+2\sqrt{5}}$$

$$= \frac{5}{8}A^3\sqrt{\frac{7+3\sqrt{5}}{2}} = \frac{B}{18}\sqrt{\frac{7\sqrt{3}+3\sqrt{15}}{10}}B.$$

$$4^{\text{th}}, R = r\sqrt{15-6\sqrt{5}} = \frac{1}{2}A\sqrt{\frac{5+\sqrt{5}}{2}} =$$

$$\frac{1}{2}\sqrt{\frac{5\sqrt{3}+\sqrt{15}}{30}}B = \sqrt[3]{\frac{3}{4}}C\sqrt{\frac{5-\sqrt{5}}{10}}.$$

$$5^{\text{th}}, r = R\sqrt{\frac{5+2\sqrt{5}}{15}} = \frac{1}{2}A\sqrt{\frac{7+3\sqrt{5}}{6}}$$

$$= \frac{1}{8}\sqrt{\frac{7\sqrt{3}+3\sqrt{15}}{10}}B = \frac{1}{2}\sqrt[3]{\frac{7\sqrt{3}+3\sqrt{15}}{30}}C.$$

See my Menfuration, p. 258, 2d edit.

IDES, in the Roman Calendar, a name given to a series of 8 days in each month; which, in the full months, March, May, July, and October, commenced on the 15th day; and in the other months, on the 13th day; from thence reckoned backward, so as in those four months to terminate on the 8th day, and in the rest on the 6th. These came between the calends and the nones. And this way of counting is still used in the Roman Chancery, and in the Calendar of the Breviary.

The Ides of May were consecrated to Mercury; the Ides of March were always esteemed unhappy, after the death of Cæsar; the time after the Ides of June was reckoned fortunate for those who entered into matrimony; the Ides of August were consecrated to Diana, and were observed as a feast by the slaves; on the Ides of September, auguries were taken for appointing the magistrates, who formerly entered into their offices on the Ides of May, and afterwards on those of March.

JET D'EAU, a French word, signifying a fountain that throws up water to some height in the air.

A Jet of water is thrown up by the weight of the column of water above its ajutage, or orifice, up to its source or reservoir; and therefore it would rise to the same height as the head or reservoir, if certain causes did not prevent it from rising quite so high. For first, the velocity of the lower particles of the Jet being greater than that of the upper, the lower water strikes that which is next above it; and as fluids press every way, by its impulse it widens, and consequently shortens the column. Secondly, the water at the top of the Jet does not immediately fall off, but forms a kind of ball or head, the weight of which depresses the Jet; but if the Jet be a little inclined, or not quite upright, it will play higher, though it will not be quite so beautiful. Thirdly, the friction against the sides of the pipe and hole of the ajutage, will prevent the Jet from rising quite so high, and a small one will be more impeded than a large one. And the fourth cause is the resistance of the air, which

is proportional to the square of the velocity of the water nearly; and therefore the defect in the height will be nearly in the same proportion, which is also the same as the proportion of the heights of the reservoirs above the ajutage. Hence, and from experience, it is found that a Jet, properly constructed, will rise to different heights according to the height of the reservoir, as in the following table of the heights of reservoirs and the heights of their corresponding Jets; the former in feet, and the latter in feet and tenths of a foot.

*Heights of Reservoirs and their Jets.*

Ref.	Jet.	Ref.	Jet.	Ref.	Jet.	Ref.	Jet.
5	4.9	31	28.3	57	49.0	82	67.0
6	5.9	32	29.2	58	49.7	83	67.7
7	6.8	33	30.0	59	50.5	84	68.4
8	7.8	34	30.8	60	51.2	85	69.1
9	8.7	35	31.6	61	52.0	86	69.8
10	9.7	36	32.5	62	52.7	87	70.5
11	10.6	37	33.3	63	53.5	88	71.1
12	11.6	38	34.1	64	54.2	89	71.8
13	12.5	39	34.9	65	54.9	90	72.5
14	13.4	40	35.7	66	55.7	91	73.2
15	14.3	41	36.6	67	56.4	92	73.8
16	15.2	42	37.4	68	57.1	93	74.5
17	16.1	43	38.1	69	57.8	94	75.2
18	17.0	44	38.9	70	58.6	95	75.8
19	17.9	45	39.8	71	59.3	96	76.5
20	18.8	46	40.5	72	60.0	97	77.2
21	19.7	47	41.3	73	60.7	98	77.8
22	20.6	48	42.1	74	61.4	99	78.5
23	21.5	49	42.9	75	62.1	100	79.1
24	22.3	50	43.7	76	62.8	110	85.6
25	23.2	51	44.4	77	63.5	120	91.9
26	24.1	52	45.2	78	64.2	130	98.0
27	24.9	53	46.0	79	64.9	140	104
28	25.8	54	46.7	80	65.6	150	110
29	26.6	55	47.5	81	66.3	160	116
30	27.5	56	48.2				

By various experiments that have been made by Mariotte, Defaguliers, and others, it has been found, that if the reservoir be 5 feet high, a conduct pipe  $1\frac{3}{4}$  inch diameter will admit a hole in the ajutage from  $\frac{1}{4}$  to  $\frac{3}{8}$  of an inch; and so on as in the following table:

Height Reservoir.	Diam. of the Ajutage.	Diam. of the Conduct Pipe.
5 feet	$\frac{1}{4}$ to $\frac{3}{8}$ inch	$1\frac{3}{4}$ inch
10	$\frac{1}{4}$ to $\frac{1}{2}$	2
15	$\frac{1}{2}$	$2\frac{1}{4}$
20	$\frac{1}{2}$	$2\frac{1}{2}$
25	$\frac{1}{2}$	$2\frac{3}{4}$
30	$\frac{1}{2}$ to $\frac{3}{4}$	$3\frac{1}{4}$
40	$\frac{3}{4}$	$4\frac{1}{2}$
50	$\frac{3}{4}$	$5\frac{1}{4}$
60	1	$5\frac{3}{4}$ or 6
80	$1\frac{1}{4}$	$6\frac{1}{2}$ or 7
100	$1\frac{1}{4}$ to $1\frac{1}{2}$	7 or 8

But



But the size of the pipe will be more or less with the distance.

If it be required to keep any number of Jets of given dimensions playing, by one common conduct-pipe; the diameter of an ajutage must be found that shall be equal to all the small ones that are given, and from this its proper conduct-pipe. Thus, if there be 4 ajutages, each  $\frac{1}{4}$  of an inch diameter; then the square of  $\frac{1}{4}$  is  $\frac{1}{16}$ , which multiplied by 4, the number of them, makes  $\frac{4}{16}$ , the square root of which is  $\frac{1}{2}$  or  $1\frac{1}{2}$ , the diameter of an ajutage equal to all the other four; to which in the table answers a pipe of 8 inches diameter. In general, the diameter of the conduct-pipe should be about 6 times that of the ajutage.

See Mariotte's *Mouvement des Eaux*; Desaguliers's *Exper. Philos.* vol. 2, p. 127, &c; Clare's *Motion of Fluids*, p. 109; &c.

JETTE, the border made round the stilts under a pier, in certain old bridges, being the same with starling, consisting of a strong framing of timber filled with stones, chalk, &c; to preserve the foundations of the piers from injury.

IGNIS FATUUS, a common meteor, chiefly seen in dark nights about meadows, marshes, and other moist places, as also in burying grounds, and near dung-hills. It is known among the people by the appellations, Will with a Wisp, and Jack with a Lantern.

Dr. Shaw describes a remarkable Ignis Fatuus, which he saw in the Holy Land, that was sometimes globular, or in the form of the flame of a candle; and presently afterward it spread itself so much as to involve the whole company in a pale harmless light, and then contract itself again, and suddenly disappear. But in less than a minute it would become visible as before; or, running along from one place to another, with a swift progressive motion, would expand itself at certain intervals over more than 2 or 3 acres of the adjacent mountains. The atmosphere had been thick and hazy, and the dew on the horses' bridles was uncommonly clammy and unctuous. In the same weather he observed those luminous appearances, which skip about the masts and yards of ships at sea, and which the sailors call corpusculæ, by a corruption of the Spanish *cuerposanto*. Shaw's *Travels*, p. 363.

Newton calls it a vapour shining without heat; and supposed it to be of the same nature with the light issuing from putrescent substances. Willughby and Ray were of opinion that it is occasioned by shining insects: but all the appearances of it observed by Derham, Beccaria, and others, sufficiently evince that it must be an ignited vapour. Inflammable air has been found to be the most common of all the factitious airs in nature; and that it is the usual product of the putrefaction and decomposition of vegetable substances in water. Signor Volta writes to Dr. Priestley, that he fires inflammable air by the electric spark, even when the electricity is very moderate: and he supposes that this experiment explains the inflammation of the Ignis Fatui, provided they consist of inflammable air, issuing from marshy ground by help of the electricity of fogs, and by falling stars, which have probably an electrical origin. See Priestley's *Obs. on Air*, vol. 3, p. 382; the *Philos. Trans. Abr.* vol. 7, p. 147 &c.

VOL. I.

ILLUMINATION, the act or effect of a luminous body, or a body that emits light; sometimes also the state of another body that receives it.

Circle of ILLUMINATION. See CIRCLE.

ILLUMINATIVE *Lunar Month*, the space of time that the moon is visible, between one conjunction and another.

IMAGE, in Optics, is the spectre or appearance of an object, made either by reflection or refraction.

In all plane mirrors, the Image is of the same magnitude as the object; and it appears as far behind the mirror as the object is before it. In convex mirrors, the Image appears less than the object; and farther distant from the centre of the convexity, than from the point of reflection. Mr. Molyneux gives the following rule for finding the diameter of an Image, projected in the distinct base of a convex mirror, viz. As the distance of the object from the mirror, is to the distance from the Image to the glass; so is the diameter of the object, to the diameter of the Image. See LENS, MIRROR, REFLECTION, and REFRACTION.

IMAGINARY *Quantities*, or Impossible Quantities, in Algebra, are the even roots of negative quantities; which expressions are Imaginary, or impossible, or opposed to real quantities; as  $\sqrt{-aa}$ , or  $\sqrt[4]{-a^4}$ , &c. For, as every even power of any quantity whatever, whether positive or negative, is necessarily positive, or having the sign +, because + by +, or - by - give equally +; from hence it follows that every even power, as the square for instance, which is negative, or having the sign -, has no possible root; and therefore the even roots of such powers or quantities are said to be impossible or Imaginary. The mixt expressions arising from Imaginary quantities joined to real ones, are also Imaginary; as  $a - \sqrt{-aa}$ , or  $b + \sqrt{-aa}$ .

The roots of negative quantities were, perhaps, first treated of in Cardan's Algebra. As to the uneven roots of such quantities, he shews that they are negative, and he assigns them: but the even roots of them he rejects, observing that they are nothing as to common use, being neither one thing nor another; that is, they are merely Imaginary or impossible. And since his time, it has gradually become a part of Algebra to treat of the roots of negative quantities. Albert Girard, in his *Invention Nouvelle en l'Algebre*, p. 42, gives names to the three sorts of roots of equations, calling them, greater than nothing, less than nothing, and *enveloppée*, as  $\sqrt{-3}$ : but this was soon after called Imaginary or impossible, as appears by Wallis's Algebra, p. 264, &c; where he observes that the square root of a negative quantity, is a mean proportional between a positive and a negative quantity; as  $\sqrt{-bc}$  is the mean proportional between +  $b$  and -  $c$ , or between -  $b$  and +  $c$ ; and this he exemplifies by geometrical constructions. See also p. 313.

The arithmetic of these Imaginary quantities has not yet been generally agreed upon; viz, as to the operations of multiplication, division, and involution; some authors giving the results with +, and others on the contrary with the negative sign -. Thus, Euler, in his Algebra, p. 106 &c, makes the square of  $\sqrt{-3}$  to be -3, of  $\sqrt{-1}$  to be -1, &c; and yet he makes the product of two impossibles, when they are unequal, to be possible and real: as  $\sqrt{-2} \times \sqrt{-3} = \sqrt{6}$ ;

4 L



$= \sqrt{6}$ ; and  $\sqrt{-1} \times \sqrt{-4} = \sqrt{4}$  or 2. But how can the equality or inequality of the factors cause any difference in the signs of the products?

If  $\sqrt{-2} \times \sqrt{-3}$  be  $= \sqrt{+6}$ , how can

$\sqrt{-3} \times \sqrt{-3}$ , which is the square of  $\sqrt{-3}$ , be  $-3$ ? Again, he makes  $\sqrt{-3} \times \sqrt{+5} = \sqrt{-15}$ .

Also in division, he makes  $\sqrt{-4} \div \sqrt{-1}$  to be  $= \sqrt{+4}$  or 2; and  $\sqrt{+3} \div \sqrt{-3} = \sqrt{-1}$ ; also

that 1 or  $\sqrt{+1} \div \sqrt{-1} = \sqrt{\frac{+1}{-1}} = \sqrt{-1}$ ;

consequently, multiplying the quotient root  $\sqrt{-1}$  by the divisor  $\sqrt{-1}$ , must give the dividend  $\sqrt{+1}$ ; and yet, by squaring, he makes the square of  $\sqrt{-1}$ , or the product  $\sqrt{-1} \times \sqrt{-1}$ , equal to  $-1$ .

But Emerson makes the product of Imaginaries to be Imaginary; and for this reason, that "otherwise a real product would be raised from impossible factors, which is absurd. Thus,

$\sqrt{-a} \times \sqrt{-b} = \sqrt{-ab}$ , and  $\sqrt{-a} \times -\sqrt{-b} = -\sqrt{-ab}$ , &c.

Also  $\sqrt{-a} \times \sqrt{-a} = -a$ , and  $\sqrt{-a} \times -\sqrt{-a} = +a$ , &c."

And thus most of the writers on this part of Algebra, are pretty equally divided, some making the product of impossibles real, and others Imaginary.

In the *Philos. Trans.* for 1778, p. 318 &c, Mr. Playfair has given an ingenious dissertation "On the Arithmetic of Impossible Quantities." But this relates chiefly to the applications and uses of them, and not to the algorithm of them, or rules for their products, quotients, squares, &c. From some operations however here performed, we learn that he makes the product of  $\sqrt{-1}$  by  $\sqrt{-1}$ , or the square of  $\sqrt{-1}$ , to be  $-1$ ; and yet in another place he makes the product of  $\sqrt{-1}$  and  $\sqrt{1-z^2}$  to be  $\sqrt{-1+z^2}$ . Mr. Playfair concludes, "that Imaginary expressions are never of use in investigations but when the subject is a property common to the measures both of ratios and of angles; but they never lead to any consequence which might not be drawn from the affinity between those measures; and that they are indeed no more than a particular method of tracing that affinity. The deductions into which they enter are thus reduced to an argument from analogy, but the force of them is not diminished on that account. The laws to which this analogy is subject; the cases in which it is perfect, in which it suffers certain alterations, and in which it is wholly interrupted, are capable of being precisely ascertained. Supported on so sure a foundation, the arithmetic of impossible quantities will always remain an useful instrument in the discovery of truth, and may be of service when a more rigid analysis can hardly be applied. For this reason, many researches concerning it, which in themselves might be deemed absurd, are nevertheless not destitute of utility. M. Bernoulli has found, for example, that if  $r$  be the radius of a circle,

the circumference is  $= \frac{4 \log. \sqrt{-1}}{\sqrt{-1}} r$ . Considered

as a quadrature of the circle, this Imaginary theorem is wholly insignificant, and would deservedly pass for an abuse of calculation; at the same time we learn from

it, that if in any equation the quantity  $\frac{\log. \sqrt{-1}}{\sqrt{-1}}$

should occur, it may be made to disappear, by the substitution of a circular arch, and a property, common to both the circle and hyperbola, may be obtained. The same is to be observed of the rules which have been invented for the transformation and reduction of impossible quantities\*; they facilitate the operations of this imaginary arithmetic, and thereby lead to the knowledge of the most beautiful and extensive analogy which the doctrine of quantity has yet exhibited.

\* The rules chiefly referred to, are those for reducing the impossible roots of an equation to the form  $A + B\sqrt{-1}$ ."

IMAGINARY Roots, of an equation, are those roots or values of the unknown quantity in an equation, which contain some Imaginary quantity. So the roots of the equation  $xx + aa = 0$ , are the two Imaginary quantities  $+\sqrt{-aa}$  and  $-\sqrt{-aa}$ , or  $+a\sqrt{-1}$  and  $-a\sqrt{-1}$ ; also the two roots of the equation  $xx + ax + aa = 0$ , are the Imaginary quantities  $\pm \frac{1}{2}a - \frac{1}{2}a\sqrt{-3}$ ; and the three roots of the equation  $x^3 - 1 = 0$ , or  $x^3 = 1$ , are 1 and  $\frac{-1 + \sqrt{-3}}{2}$  and  $\frac{-1 - \sqrt{-3}}{2}$ , the first real,

and the two latter Imaginary. Sometimes too the real root of an equation may be expressed by Imaginary quantities; as in the irreducible case of cubic equations, when the root is expressed by Cardan's rule; and that happens whenever the equation has no Imaginary roots at all; but when it has two Imaginary roots, then the only real root is expressed by that rule in an Imaginary form. See my paper on Cubic Equations, in the *Philos. Trans.* for 1780, p. 406 &c.

Albert Girard first treated expressly on the impossible or Imaginary roots of equations, and shewed that every equation has as many roots, either real or Imaginary, as the index of the highest power denotes. Thus, the roots of the biquadratic equation  $x^4 = 4x - 3$ , he shews are two real and two Imaginary, viz, 1, 1,  $-1 + \sqrt{-2}$ , and  $-1 - \sqrt{-2}$ ; and he renders the relation general, between all the roots and the coefficients of the terms of the equation. See his *Invention Nouvelle en l'Algebre*, anno 1629, theor. 2, pa. 40 &c.

M. D'Alembert demonstrated, that every Imaginary root of any equation can always be reduced to the form  $e + f\sqrt{-1}$ , where  $e$  and  $f$  are real quantities. And hence it was also shewn, that if one root of an equation be  $e + f\sqrt{-1}$ , another root of it will always be  $e - f\sqrt{-1}$ ; and hence it appears that the number of the Imaginary roots in any equation is always even, if any; i. e. either none, or else two, or four, or six, &c. *Memoirs of the Academy of Berlin*, 1746.

To discover how many impossible roots are contained in any proposed equation, Newton gave this rule, in his *Algebra*, viz, Constitute a series of fractions, whose denominators are the series of natural numbers 1, 2, 3, 4, 5, &c, continued to the number shewing the index or exponent of the highest term of the equations, and their numerators the same series of numbers in the contrary order: and divide each of these fractions by that next before it, and place the resulting quotients over the intermediate terms of the equation; then under each of the intermediate terms, if its square multiplied by



by the fraction over it, be greater than the product of the terms on each side of it, place the sign +; but if not, the sign -; and under the first and last term place the sign +. Then will the equation have as many Imaginary roots as there are changes of the underwritten signs from + to -, and from - to +. So for the equation  $x^3 - 4x^2 + 4x - 6 = 0$ , the series of fractions is  $\frac{3}{1}$ ,  $\frac{2}{1}$ ,  $\frac{1}{1}$ ; then the second divided by the first gives  $\frac{2}{3}$  or  $\frac{1}{1}$ , and the third divided by the second gives  $\frac{1}{2}$  also; hence these quotients placed over the intermediate terms, the whole will stand

$$\begin{array}{cccc} & \frac{1}{3} & & \frac{1}{2} \\ \text{thus, } & x^3 & - & 4x^2 & + & 4x & - & 6. \\ & + & & + & & - & & + \end{array}$$

Now because the square of the 2d term multiplied by its superscribed fraction, is  $\frac{1}{3}x^4$ , which is greater than  $4x^4$  the product of the two adjacent-terms, therefore the sign + is set below the 2d term; and because the square of the 3d term multiplied by its overwritten fraction, is  $\frac{1}{2}x^2$ , which is less than  $24x^2$  the product of the terms on each side of it, therefore the sign - is placed under that term; also the sign + is set under the first and last terms. Hence the two changes of the underwritten signs + + - +, the one from + to -, and the other from - to +, shew that the given equation has two impossible roots.

When two or more terms are wanting together, under the place of the 1st of the deficient terms write the sign -, under the 2d the sign +, under the 3d -, and so on, always varying the signs, except that under the last of the deficient terms must always be set the sign +, when the adjacent terms on both sides of the deficient terms have contrary signs. As in the equation

$$\begin{array}{ccccccccc} x^5 & + & ax^4 & * & * & * & + & a^5 & = & 0, \\ + & & + & - & + & - & & + & & \end{array}$$

which has four Imaginary roots.

The author remarks, that this rule will sometimes fail of discovering all the impossible roots of an equation, for some equations may have more of such roots than can be found by this rule, tho' this seldom happens.

Mr. Maclaurin has given a demonstration of this rule of Newton's, together with one of his own, that will never fail. And the same has also been done by Mr. Campbell. See *Philos. Trans.* vol. 34, p. 104, and vol. 35, p. 515.

The real and imaginary roots of equations may be found from the method of fluxions, applied to the doctrine of maxima and minima, that is, to find such a value of  $x$  in an equation, expressing the nature of a curve, made equal to  $y$ , an abscissa which corresponds to the greatest and least ordinate. But when the equation is above 3 dimensions, the computation is very laborious. See Stirling's treatise on the lines of the 3d order, Schol. pr. 8, pa. 59, &c.

IMBIBE, is commonly used in the same sense as absorb, viz, where a dry porous body takes up another that is moist.

IMMENSE, that whose amplitude or extension cannot be equalled by any measure whatsoever, or how often soever repeated.

IMMERSION, the act of plunging into water, or some other fluid.

IMMERSION, in Astronomy, is when a star, or

planet comes so near the sun, that it cannot be seen; being as it were enveloped, and hid in the rays of that luminary.

IMMERSION also denotes the beginning of an eclipse, or of an occultation, when the body, or any part of it just begins to disappear, either behind the edge of another body, or in its shadow. As, in an eclipse of the moon, when she begins to be darkened by entering into the shadow of the earth: or the beginning of an eclipse of the sun, when the moon's disc just begins to cover him: or the beginning of the eclipses of any of the satellites, as those of Jupiter, by entering into his shadow: or, lastly, the beginning of an occultation of any star or planet, by passing behind the body of the moon or another planet. In all these cases, the darkened body is said to immerge, or to be immersed, or begin to be hid, by dipping as it were into the shade. In like manner, when the darkened body begins to appear again, it is said to emerge, or come out of darkness again.

IMPACT, the simple or single action of one body upon another to put it in motion. Point of Impact, is the place or point where a body acts.

IMPENETRABILITY, a quality by which a thing cannot be pierced or penetrated; or a property of body by which it fills up certain spaces, so that there is no room in them for any other body.

IMPENETRABLE, that cannot be penetrated.

IMPERFECT Number, is that whose aliquot parts, taken all together, do not make a sum that is equal to the number itself, but either exceed it, or fall short of it; being an abundant number in the former case, and a defective number in the latter. Thus, 12 is an abundant Imperfect number, because the sum of all its aliquot parts, 1, 2, 3, 4, 6, makes 16, which exceeds the number 12. And 10 is a defective Imperfect number, because its aliquot parts, 1, 2, 5, taken all together, make only 8, which is less than the number 10 itself.

IMPERIAL Table, is an instrument made of brass, with a box and needle, and staff, &c, used for measuring of land.

IMPERVIOUS, not to be pervaded or entered either because of the closeness of the pores, or the particular configuration of its parts.

IMPETUS, in Mechanics, force, momentum, motion, &c.

IMPOSSIBLE Quantity, or Root, the same as IMAGINARY ones; which see.

IMPOST, in Architecture, a capital or plinth, to a pillar, or pilaster, or pier, that supports an arch, &c.

IMPROPER Fraction, is a fraction whose numerator is either equal to, or greater than, its denominator. As  $\frac{5}{2}$  or  $\frac{3}{2}$  or  $\frac{1}{2}$ . An Improper fraction is reduced to a whole or mixt number, by dividing the numerator by the denominator; the quotient is the integer, and the remainder set over the divisor makes the fractional part of the value of the original Improper fraction. Thus  $\frac{5}{2} = 2\frac{1}{2}$ , and  $\frac{3}{2} = 1\frac{1}{2}$ , and  $\frac{1}{2} = \frac{1}{2}$ . So that when the numerator is just equal to the denominator, the Improper fraction is exactly equal to unity or 1; but when the numerator is the greater, the fraction is greater than 1.



**IMPULSE**, the single or momentary action or force by which a body is impelled; in contradistinction to continued forces; like the blow of a hammer, &c.

**IMPULSIVE**, a term applied to actions by impulse.

**INACCESSIBLE Height or Distance**, is that which cannot be approached, or measured by actual measurement, by reason of some impediment in the way; as water, &c.


See **HEIGHTS and DISTANCES**.

**INCEPTIVE**, of *Magnitude*, a term used by Dr. Wallis, to express such moments, or first principles, as, though of no magnitude themselves, are yet capable of producing such as are. See **INFINITE**, and **INDIVISIBLE**. Thus, a point has no magnitude itself, but is inceptive of a line, which it produces by its motion. Also a line, though it has no breadth, is yet Inceptive of breadth; that is, it is capable, by its motion, of producing a surface, which has breadth.

**INCH**, a common English measure, being the 12th part of a foot, or 3 barley corns in length.

**INCIDENCE**, or *line of INCIDENCE*, in Mechanics, implies the direction or inclination in which one body strikes or acts on another.—In the incursions of two moving bodies, their Incidence is said to be Direct or Oblique, as the directions of their motion make a straight line, or an angle at the point of Impact.

*Angle of INCIDENCE*, by some writers, denotes the angle comprehended between the line of Incidence, and a perpendicular to the body acted on at the point of Incidence. Thus, suppose AB an Incident line, and BF a perpendicular to the plane CB at the incident point B; then ABF is the angle of Incidence, or of inclination.



But, according to Dr. Barrow, and some other writers, the Angle of Incidence is the complement of the former, or the angle made between the incident line, and the plane acted on, or a tangent at the point of Incidence; as the angle ABC.

It is demonstrated by optical writers, 1st, That the Angle of Incidence, of the rays of light, is always equal to the angle of reflection; and that they lie in the same plane. And the same is proved by the writers on Mechanics, concerning the reflection of elastic bodies. That is, the  $\angle ABF = \text{the } \angle FBD$ , or the  $\angle ABC = \text{the } \angle DBE$ .—2d, That the sines of the Angles of Incidence and refraction are to each other, either accurately, or very nearly, in a given or constant ratio.—3dly, That from air into glass, the sine of the Angle of Incidence, is to the sine of the angle of refraction, as 300 to 193, or nearly as 14 to 9: and, on the other hand, that out of glass into air, the sine of the Angle of Incidence, is to the sine of the angle of refraction, as 193 to 300, or as 9 to 14 nearly.

**INCIDENCE of Eclipse**. See **ECLIPSE** and **IMMERSION**.

*Axis of INCIDENCE*, is the line FB perpendicular to the reflecting plane at the point of Incidence B.

*Cathetus of INCIDENCE*. See **CATHETUS**, and **REFLECTION**.

*Line of INCIDENCE*, in Catoptrics, denotes a right line, as AB, in which light is propagated from a radiant point A, to a point B, in the surface of a speculum. The same line is also called an Incident ray.

*Line of INCIDENCE*, in Dioptrics, is a right line, as AB, in which light is propagated unrefracted, in the same medium, from the radiant point to the surface of the refracting body, CBE.


*Point of INCIDENCE*, is the point B on the surface of the reflecting or refracting medium, on which the Incident ray falls.

*Scruples of INCIDENCE*. See **SCRUPLES**.

**INCIDENT Ray**, is the line or ray AB, falling on the surface of any body, at B.

**INCLINATION**, in Geometry, Mechanics, or Physics, denotes the mutual tendency of two lines, planes, or bodies, towards one another; so that their directions make at the point of concurrence some certain angle.

*INCLINATION of the Axis of the Earth*, is the angle it makes with the plane of the ecliptic; or the angle between the planes of the equator and ecliptic.



*INCLINATION of a Line to a plane*, is the acute angle, as CDE, which the line CD makes with another line DE drawn in the plane through the incident point D and the foot of a perpendicular E from any point of the line upon the plane.

*INCLINATION of an Incident ray*, is the angle of inclination, or angle of incidence.

*INCLINATION of the Magnetical needle*. See **DIPPING Needle**.

*INCLINATION of Meridians*, in Dialling, is the angle that the hour-line on the globe, which is perpendicular to the dial-plane, makes with the meridian.

*INCLINATION of the Orbit of a planet*, is the angle formed by the planes of the ecliptic and of the orbit of the planet. The quantity of this Inclination for the several planets, is as follows, viz.

Mercury	-	-	6° 54'
Venus	-	-	3 20
Earth	-	-	0 0
Moon	-	-	5 18
Mars	-	-	1 52
Jupiter	-	-	1 20
Saturn	-	-	2 30
Herschel	-	-	0 48

*INCLINATION of a Plane*, in Dialling, is the arch of a vertical,



a vertical circle, perpendicular both to the plane and the horizon, and intercepted between them.

*INCLINATION of a Planet*, is the arch or angle comprehended between the ecliptic and the place of the planet in its orbit. The greatest Inclination, or declination, is the same as the Inclination of the orbit; which see above.

*INCLINATION of a Reflected ray*, is the angle which a ray after reflection makes with the axis of Inclination; as the angle FBD, in the last fig. but one.


*INCLINATION of Two Planes*, is the angle made by two lines drawn in those planes perpendicular to their common intersection, and meeting in any point of that intersection.

*Angle of INCLINATION*, is the same as what is otherwise called the angle of incidence.

*Argument of INCLINATION*. See ARGUMENT.


*INCLINED Plane*, in Mechanics, is a plane inclined to the horizon, or making an angle with it. It is one of the simple mechanic powers, and the double inclined plane makes the wedge.

1. The power gained by the Inclined plane, is in proportion as the length of the plane is to its height, or as radius to the sine of its inclination; that is, a given weight hanging freely, will balance upon the plane another weight, that shall be greater in that proportion. So, when the greater weight  $W$  on the plane, is balanced by the less weight  $w$  hanging perpendicularly, then is  $w : W :: BC : AC :: \sin. \angle A : \text{radius}$ . Or, in other words, the relative gravity of a body upon the plane, or its force in descending down the plane, is to its absolute gravity or weight, in the same proportion of the height of the plane to its length, or of the sine of inclination to radius.



2. Hence therefore the relative gravities of the same body on different inclined planes, or their forces to descend down the planes, are to each other, as the sines of the angles of inclination, to radius 1, or directly as the heights of the planes, and inversely as their lengths.


3. Hence, if the planes have the same height, and absolute weights of the bodies be directly proportional to the lengths of the planes, then the forces to descend will be equal. Consequently, if the bodies be then connected by a string acting parallel to the planes, they will exactly balance each other; as in the annexed figure.



4. The relative force of gravity upon the plane being in a constant ratio to the absolute weight of the body, viz, as sine of inclination to radius; therefore all the laws relating to the perpendicular free descents of bodies by gravity, hold equally true for the descents on inclined planes; such as, that the motion is a uniformly accelerated one; that the velocities are directly as the times, and the spaces as the square of either of them; using only the relative force upon the plane for the absolute weight of the body, or instead of  $32\frac{1}{2}$  feet, the velocity generated by gravity in the first second of

time, using  $32\frac{1}{2}s$ , where  $s$  is the sine of the inclination to the radius 1.

5. The velocity acquired by a body in descending down an Inclined plane AC, when the body arrives at A, is the same as the velocity acquired by descending freely down the perpendicular altitude BC, when it arrives at B. But the times are very different; for the time of descending down the Inclined plane, is greater than down the perpendicular, in the same proportion as the length of the plane AC, is to the height CB: and so the time of descending from any point C to a horizontal line or plane ABG &c, down any oblique line, or Inclined plane, is directly proportional to the length of that plane, CA, or CD, or CE, or CB, or CF, &c.



6. Hence, if there be drawn AH perpendicular to AC, meeting CB produced in H; then the time of descending down any plane CA, is equal to the time of descending down the perpendicular CH. So that, if upon CH as a diameter a circle be described, the times of descent will be exactly equal, down every chord in the circle, beginning at C, and terminating any where in the circumference, as CI, CA, CK; CH, &c, or beginning any where in the circumference, and terminating at the lowest point of the circle, as CH, IH, AH, KH, &c.

7. When bodies ascend up Inclined planes, their motion is uniformly retarded; and all the former laws for descents, or the generation of motion, hold equally true for ascents, or the destruction of as much motion.

*INCLINED Towers*, are towers inclined, or leaning out of the perpendicular. See TOWERS.

*INCLINERS*, in Dialling, are inclined dials. See DIAL.

*INCOMMENSURABLE*, Lines, or Numbers, or Quantities in general, are such as have no common measure, or no line, number, or quantity of the same kind, that will measure or divide them both without a remainder. Thus, the numbers 15 and 16 are Incommensurable, because, though 15 can be measured by 3 and 5, and 16 by 2, 4, and 8, there is yet no single number that will divide or measure them both.

Euclid demonstrates (prop. 117, lib. 10) that the side of a square and its diagonal are Incommensurable to each other. And Pappus, prop. 17, lib. 4, speaks of Incommensurable angles.

*INCOMMENSURABLE in Power*, is said of quantities whose 2d powers, or squares, are Incommensurable. As  $\sqrt{2}$  and  $\sqrt{3}$ , whose squares are 2 and 3, which are Incommensurable. It is commonly supposed that the diameter and circumference of a circle are Incommensurable to each other; at least their commensurability has never been proved. And Dr. Barrow surmises even that they are infinitely Incommensurable, or that all possible powers of them are Incommensurable.

*INCOMPOSITE Numbers*, are the same with those called by Euclid prime numbers, being such as are not composed by the multiplication together of other numbers. As 3, 5, 7, 11, &c.

INCREMENT,



INCREMENT, is the small increase of a variable quantity. Newton, in his Treatise on Fluxions, calls these by the name Moments, and observes that they are proportional to the velocity or rate of increase of the flowing or variable quantities, in an indefinitely small time; he denotes them by subjoining a cipher 0, to the flowing quantity whose moment or Increment it is; thus  $ao$  the moment of  $a$ . In the doctrine of Increments, by Dr. Brooke Taylor and Mr. Emerson, they are denoted by points below the variable quantities; as  $\dot{x}$ . Some have also denoted them by accents underneath the letter, as  $\acute{x}$ ; but it is now more usual to express them by accents over the same letter; as  $\acute{x}$ .

INCREMENTS, *Method of*, a branch of Analytics, in which a calculus is founded on the properties of the successive values of variable quantities, and their differences, or Increments.

The inventor of the Method of Increments was the learned Dr. Taylor, who, in the year 1715, published a treatise upon it; and afterwards gave some farther account and explication of it in the *Philos. Trans.* as applied to the finding the sums of series. And another ingenious and easy treatise on the same, was published by Mr. Emerson, in the year 1763. The method is nearly allied to Newton's Doctrine of Fluxions, and arises out of it. Also the Differential method of Mr. Stirling, which he applies to the summation and interpolation of series, is of the same nature as the Method of Increments, but not so general and extensive.

From the Method of Increments, Mr. Emerson observes, "The principal foundation of the Method of Fluxions may be easily derived. For as in the Method of Increments, the Increment may be of any magnitude, so in the Method of Fluxions, it must be supposed infinitely small; whence all preceding and successive values of the variable quantity will be equal, from which equality the rules for performing the principal operations of fluxions are immediately deduced. That I may give the reader, continues he, a more perfect idea of the nature of this method: suppose the abscissa of a curve be divided into any number of equal parts, each part of which is called the Increment of the abscissa; and imagine so many parallelograms to be erected thereon; either circumscribing the curvilinear figure, or inscribed in it; then the finding the sum of all these parallelograms is the business of the Method of Increments. But if the parts of the abscissa be taken infinitely small, then these parallelograms degenerate into the curve; and then it is the business of the Method of Fluxions, to find the sum of all, or the area of the curve. So that the Method of Increments finds the sum of any number of finite quantities; and the Method of Fluxions the sum of any infinite number of infinitely small ones: and this is the essential difference between these two methods." Again, "There is such a near relation between the Method of Fluxions, and that of Increments, that many of the rules for the one, with little variation, serve also for the other. And here, as in the Method of Fluxions, some questions may be solved, and the integrals found, in finite terms; whilst in others we are forced to have recourse to infinite series for a solution. And the like difficulties will occur in the Method of Increments, as usually happen

in Fluxions. For whilst some fluxionary quantities have no fluents, but what are expressed by series; so some Increments have no integrals, but what infinite series afford; which will often, as in fluxions, diverge and become useless."

By means of the Method of Increments, many curious and useful problems are easily resolved, which scarcely admit of a solution in any other way. As, suppose several series of quantities be given, whose terms are all formed according to some certain law, which is given; the Method of Increments will find out a general series, which comprehends all particular cases, and from which all of that kind may be found.

The Method of Increments is also of great use in finding any term of a series proposed: for the law being given by which the terms are formed; by means of this general law, the Method of Increments will help us to this term, either expressed in finite quantities, or by an infinite series.

Another use of the Method of Increments, is to find the sums of series; which it will often do in finite terms. And when the sum of a series cannot be had in finite terms, we must have recourse to infinite series; for the integral being expressed by such a series, the sum of a competent number of its terms will give the sum of the series required. This is equivalent to transforming one series into another, converging quicker: and sometimes a very few terms of this series will give the sum of the series sought.

#### *Definitions in the Method of Increments.*

1. When a quantity is considered as increasing, or decreasing, by certain steps or degrees, it is called an Integral.
2. The increase of any quantity from its present value, to the next succeeding value, is called an Increment: or, if it decreases, a Decrement.
3. The increase of any Increment, is the Second Increment; and the increase of the 2d Increment, is the 3d Increment; and so on.
4. Succeeding Values, are the several values of the integral, succeeding one another in regular order, from the present value; and Preceding Values, are such as arise before the present value. All these are called by the general term Factors.
5. A Perfect quantity is such as contains any number of successive values without intermission; and a Defective quantity, is that which wants some of the successive values. Thus  $x^1 x^2 x^3 x^4$  is a Perfect quantity; and  $x^1 x^2 x^3$ , an Imperfect or defective one.

*Notation.* This, according to Mr. Emerson's method, is as follows:

1. Simple Integral quantities are denoted by any letters whatever, as  $z, y, x, u$ , &c.
2. The several values of a simple integral, are denoted by the same letter with small figures under them: so if  $z$  be an integral, then  $z, z^1, z^2, z^3$ , &c. are the present value, and the 1st, 2d, 3d, &c. successive values of it; and the preceding values are denoted by figures with negative signs, thus  $z, z^1, z^2, z^3$ , are the 1st, 2d, 3d, &c. preceding values.



3d, 4th preceding values; and the figure denoting any value, is the characteristic.

3. The Increments are denoted with the same letters, and points under them: thus,  $\dot{x}$  is the Increment of  $x$ , and  $\ddot{z}$  is the increment of  $z$ . Also  $\dot{x}$  is the Increment of  $x$ ; and  $\dot{x}$  of  $x$ , &c.

4. The 2d, 3d, and other Increments, are denoted with two, three, or more points: so  $\ddot{x}$  is the 2d Increment of  $x$ , and  $\dddot{x}$  is the 3d Increment of  $x$ , and so on. And these are denominated Increments of such an order, according to the number of points.

5. If  $x$  be any Increment, then  $[x]$  is the integral of it; also  $^2[x]$  denotes the integral of  $[x]$ , or the 2d integral of  $x$ ; and  $^3[x]$  is the 3d integral of  $x$ , or an integral of the 3d order, &c.

6. Quantities written thus,  $\overset{1}{x} \dots \overset{5}{x}$  mean the same as  $\overset{1}{x} \overset{2}{x} \overset{3}{x} \overset{4}{x} \overset{5}{x}$ , or signify that the quantities are continued from the first to the last, without break or interruption.

To find the Increment of any Integral, or variable quantity.

Rule 1. If the proposed quantity be not fractional, and be a perfect integral, consisting of the successive values of the variable quantity which increases uniformly: Multiply the proposed integral by the number of factors, and change the lowest factor for an Increment. So the Increment of  $a - 3x + 6z$  is  $-3\dot{x} + 6\dot{z}$ ; for the Increment of the constant quantity  $a$  is 0 or nothing. So likewise,

The Increment of  $c \overset{1}{x} \overset{2}{x} \overset{3}{x} \overset{4}{x}$ , is  $4c \overset{1}{x} \overset{2}{x} \overset{3}{x}$ .

The Increment of  $\overset{1}{a} \overset{2}{x} \overset{3}{x} \overset{4}{x}$ , is  $3a \overset{1}{x} \overset{2}{x} \overset{3}{x}$ .

The Increment of  $\overset{1}{x} \dots \overset{n}{x}$ , is  $m + n + 1 \cdot \overset{1}{x} \overset{2}{x} \dots \overset{n}{x}$ .

Rule 2. In fractional quantities, where the denominator is perfect, and the variable quantity increases uniformly: Multiply the proposed integral by the number of factors, and by the constant Increment with a negative sign, and take the next succeeding value into the denominator. Thus,

The Increment of  $\frac{a}{\overset{1}{x} \dots \overset{4}{x}}$ , is  $\frac{-5a\dot{x}}{\overset{1}{x} \dots \overset{5}{x}}$ .

The Increment of  $\frac{c}{\overset{1}{x} \dots \overset{5}{x}}$ , is  $\frac{-9c\dot{x}}{\overset{1}{x} \dots \overset{6}{x}}$ .

Rule 3. The Increment of any power, as  $x^n$  is  $\overset{1}{x} + \overset{2}{x} - x^n$ ; that is, the difference between the present value  $x^n$  and the next succeeding value  $\overset{1}{x} + \overset{2}{x}$ . And generally, the Increment of any quantity whatever, is found by subtracting the present value, or the given quantity, from its next succeeding value. Also by expanding the compound quantity in a series, and subtracting  $x^n$  from it, the Increment will be either

$$= nx^{n-1}\dot{x} + \frac{n \cdot n-1}{2} x^{n-2} \dot{x}^2 + \frac{n \cdot n-1 \cdot n-2}{2 \cdot 3} x^{n-3} \dot{x}^3 \text{ \&c, or}$$

$$= x^n \times : \frac{n\dot{x}}{x} + \frac{n \cdot n-1}{2} \cdot \frac{\dot{x}^2}{x^2} + \frac{n \cdot n-1 \cdot n-2}{2 \cdot 3} \cdot \frac{\dot{x}^3}{x^3} \text{ \&c.}$$

So the Increment of  $x^4$ , is  $\overset{1}{x} + \overset{2}{x} - x^4 = 4x^3\dot{x} + 6x^2\dot{x}^2 + 4x\dot{x}^3 + \dot{x}^4$ .

The Increment of  $\frac{1}{x^3}$  or  $x^{-3}$ .

$$\begin{aligned} \text{is } \frac{1}{\overset{1}{x} + \overset{2}{x}} - \frac{1}{x^3} &= \frac{1}{x + \dot{x}} - \frac{1}{x^3} \\ &= -3x^{-4}\dot{x} + 6x^{-5}\dot{x}^2 - 10x^{-6}\dot{x}^3 \text{ \&c} \\ &= -\frac{3\dot{x}}{x^4} + \frac{6\dot{x}^2}{x^5} - \frac{10\dot{x}^3}{x^6} \text{ \&c.} \end{aligned}$$

The Increment of  $a^x$ ,  $a$  being constant, is  $a^{\overset{1}{x}} - a^x = a^{x+\dot{x}} - a^x = a^x \cdot (a^{\dot{x}} - 1)$ .

$$\begin{aligned} \text{The Increment of } \frac{1}{a^x} \text{ is } \frac{1}{a^{\overset{1}{x}}} - \frac{1}{a^x} \\ &= \frac{a^x - a^{\overset{1}{x}}}{a^x a^{\overset{1}{x}}} = \frac{a^x - a^{x+\dot{x}}}{a^x a^{x+\dot{x}}} = \frac{1 - a^{\dot{x}}}{a^{x+\dot{x}}}. \end{aligned}$$

The Increment of  $xz$  is  $\overset{1}{x} \overset{1}{z} - xz = \overset{1}{x} \overset{1}{z} + \overset{2}{x} \overset{1}{z} - xz = x\dot{z} + \dot{x}z + \dot{x}\dot{z}$ .

And so on for any form of Integral whatever, subtracting the given quantity from its next succeeding value. So,

The Increment of the log. of  $x$  is  $\log. \overset{1}{x} - \log. x = \log. x + \dot{x} - \log. x = \log. \frac{x + \dot{x}}{x}$ , which, by the nature of logarithms, is

$$\frac{\dot{x}}{x} - \frac{\dot{x}^2}{2x^2} + \frac{\dot{x}^3}{3x^3} - \frac{\dot{x}^4}{4x^4} \text{ \&c.}$$

Schol. From hence may be deduced the principles and rules of fluxions; for the method of fluxions is only a particular case of the method of Increments, fluxions being infinitely small Increments; therefore if in any form of Increments the Increment be taken infinitely small, the form or expression will be changed into a fluxional one.

Thus, in  $x\dot{z} + \dot{x}z + \dot{x}\dot{z}$ , which is the Increment of the rectangle  $xz$ , if  $\dot{x}$  and  $\dot{z}$  be changed for  $\dot{x}$  and  $\dot{z}$ , the expression will become  $x\dot{z} + \dot{x}z + \dot{x}\dot{z}$  for the fluxion of  $xz$ , or only  $x\dot{z} + \dot{x}z$ , because  $\dot{x}\dot{z}$  is infinitely less than the rest.

So likewise, if  $\dot{x}$  be changed for  $\dot{x}$  in this

$$nx^{n-1}\dot{x} + \frac{n \cdot n-1}{2} x^{n-2} \dot{x}^2 + \frac{n \cdot n-1 \cdot n-2}{2 \cdot 3} x^{n-3} \dot{x}^3$$

&c, which is the Increment of  $x^n$ , it becomes

$$nx^{n-1}\dot{x} + \frac{n \cdot n-1}{2} x^{n-2} \dot{x}^2 + \frac{n \cdot n-1 \cdot n-2}{2 \cdot 3} x^{n-3} \dot{x}^3$$

&c, or only  $nx^{n-1}\dot{x}$ , for the fluxion of the power  $x^n$ , as all the terms after the first will be nothing, because  $\dot{x}^2$  and  $\dot{x}^3$  &c are infinitely less than  $\dot{x}$ .

And thus may all the other forms of fluxions be derived from the corresponding Increments. And in like manner, the finding of the integrals, is only a more general way of finding fluents, as appears in what follows.

To find out the Integral of any given Increment.

Rule 1. When the variable quantity increases uniformly, and the proposed integral consists of the successive values of it multiplied together, or is a perfect Increment not fractional: Multiply the given Increment



ment by the next preceding value of the variable quantity, then divide by the new number of factors, and by the constant Increment.

Ex. Thus, the integral of  $4cx^2x^3$  is  $cx^3x^3$ .

The integral of  $3ax^2x^3$  is  $ax^3x^3$ .

Rule 2. In a fractional expression, where the variable quantity increases uniformly, and the denominator is perfect, containing the successive values of the variable quantity: Throw out the greatest value of the variable letter, then divide by the new number of factors, and by the constant Increment with a negative sign. So,

The integral of  $\frac{ax}{x \dots x}$  is  $\frac{-a}{5x \dots x}$ .

The integral of  $\frac{-cx}{x \dots x}$  is  $\frac{c}{x \dots x}$ .

Rule 3. Various other particular rules are given, but these and the two foregoing are all best included in the following general table of the most useful forms of Increments and integrals, to be used in the same way as the similar table of fluxions and fluents, to which these correspond.

A Table of Increments and their Integrals.		
Forms	Increments	Integrals
1	$x$ when constant not constant	$x$ , or $x$ , or $x$ , or $x$ &c. $x$ only.
2	$x \dots x$ $x$ constant	$x \dots x$ $m + n$
3	$\frac{ax}{x \dots x}$ $x$ constant	$\frac{-a}{m + n - 2 \dots x}$
4	$zx + xz$	$xz$
5	$\frac{zx - xz}{xz}$	$\frac{x}{z}$
6	$a^x$ $x$ given	$\frac{a^x}{a^x - 1}$

Integrals, when found from given Increments, are corrected in the very same way as fluents when found from given fluxions, viz, instead of every several variable quantity in the integral, substituting such a determinate value of them as they are known to have in some particular case; and then subtracting each side of the resulting equation from the corresponding side of the integral, the remaining equation will be the correct form of the integrals.

For an example of the use of the Method of Increments, suppose it were required to find the sum of

any number of terms of the series  $1.2 + 2.3 + 3.4 + 4.5$  &c.

Let  $x$  be the number of the terms,  
and  $z$  the sum of them.

Then, by the progression of the series, the last or the  $x$  term is  $xx$ , and the next term after that will be  $x^2$ , that is  $z = xx$ , where  $x = 1$ . Hence the integral is  $z = \frac{1}{2}xx = \frac{1}{2}x \cdot x + 1 \cdot x + 2$ , which is the sum of  $x$  terms of the given series. So if the number of terms  $x$  be 10, this becomes  $\frac{1}{2} \cdot 10 \cdot 11 = 55$ , which is the sum of 10 terms of the given series  $1.2 + 2.3 + 3.4$  &c. Or, when  $x = 100$ , the sum of 100 terms of the same series is  $\frac{1}{2} \cdot 100 \cdot 101 = 5050$ .

Again, to find the sum  $z$  of  $n$  terms of the series  $\frac{1}{1.3.5} + \frac{1}{3.5.7} + \frac{1}{5.7.9}$  &c.

Here the  $n$ th term is  $\frac{1}{2n-1.2n+1.2n+3}$ .  
Put  $x = 2n-1$ ; then is  $x = 2n = 2$ , and

the  $n$ th term is  $\frac{1}{x \cdot x + 2 \cdot x + 4} = \frac{1}{x^2 + 2x + 4}$ ;

and the  $n+1$ th term or  $z$  is  $\frac{1}{x^2 + 2x + 4}$ ; the general

integral of which is  $z = \frac{-1}{2x^2 + 2x + 4} = \frac{-1}{4x^2 + 4x + 4}$ .

But this wants a correction; for when  $n = 0$  or no terms, then  $x = -1$ , and the sum  $z = 0$ , and the integral becomes  $z$  or  $0 = \frac{-1}{4 \cdot 1 \cdot 3} = \frac{-1}{12}$ ; that is  $-\frac{1}{12}$  is the correction, and being subtracted, the correct state of the integrals becomes

$z = \frac{1}{12} - \frac{1}{4x^2 + 4x + 4} = \frac{1}{12} - \frac{1}{4 \cdot 2n + 1 \cdot 2n + 3}$ , which

is the sum of  $n$  terms of the proposed series. And when  $n$  is infinite, the latter fraction is nothing, and the sum of the infinite series, or the infinite number of the terms, is accurately  $\frac{1}{12}$ .

When  $n = 100$ , the sum of 100 terms of the series becomes  $z = \frac{1}{12} - \frac{1}{4 \cdot 201 \cdot 203} = \frac{1}{12} - \frac{1}{163212}$ .

For more ample information and application on this science, see Emerson's Increments, Taylor's Methodus Incrementorum, and Stirling's Summatio & Interpolatio Serierum.

INCURVATION of the rays of Light. See LIGHT, and REFRACTION.

INDEFINITE, Indeterminate, that which has no certain bounds, or to which the human mind cannot affix any. Des Cartes uses the word, in his Philosophy, instead of infinite, both in numbers and quantities, to signify an inconceivable number, or number so great, that an unit cannot be added to it; and a quantity so great, as not to be capable of any addition. Thus, he says, the stars, visible and invisible, are in number Indefinite, and not, as the Ancients held, infinite; and that



that quantity may be divided into an Indefinite number of parts, not an infinite number.

Indefinite is now commonly used for indeterminate, number or quantity, that is, a number or quantity in general, in contradistinction from some particular known and given one.

INDETERMINED, or INDETERMINATE, in Geometry, is understood of a quantity, which has no certain or definite bounds.

INDETERMINATE *Problem*, is that which admits of innumerable different solutions, and sometimes perhaps only of a great many different answers; otherwise called an unlimited problem.

In problems of this kind the number of unknown quantities concerned, is greater than the number of the conditions or equations by which they are to be found; from which it happens that generally some other conditions or quantities are assumed, to supply the defect, which being taken at pleasure, give the same number of answers as varieties in those assumptions.

As, if it were required to find two square numbers whose difference shall be a given quantity  $d$ . Here, if  $x^2$  and  $y^2$  denote the two squares, then will  $x^2 - y^2 = d$ , by the question, which is only one equation, for finding two quantities. Now by assuming a third quantity  $z$  so that  $z = x + y$  the sum of the two roots; then

is  $x = \frac{z^2 + d}{2z}$ , and  $y = \frac{z^2 - d}{2z}$ , which are the two

roots having the difference of their squares equal to the given quantity  $d$ , and are expressed by means of an assumed quantity  $z$ ; so that there will be as many answers to the question, as there can be taken values of the Indeterminate quantity  $z$ , that is, innumerable.

Diophantus was the first writer on Indeterminate problems, viz, in his Arithmetic or Algebra, which was first published in 1575 by Xilander, and afterwards in 1621 by Bachet, with a large commentary, and many additions to it. His book is wholly upon this subject; whence it has happened, that such kind of questions have been called by the name of Diophantine problems. Fermat, Des Cartes, Frenicle, in France, and Wallis and others in England, particularly cultivated this branch of Algebra, on which they held a correspondence, proposing difficult questions to each other; an instance of which are those two curious ones, proposed by M. Fermat, as a challenge to all the mathematicians of Europe, viz 1st, To find a cube number which added to all its aliquot parts shall make a square number; and 2d, To find a square number which added to all its aliquot parts shall make a cubic number; which problems were answered after several ways by Dr. Wallis, as well as some others of a different nature. See the Letters that passed between Dr. Wallis, the lord Brouncker, Sir Kenelm Digby, &c, in the Doctor's Works; and the Works of Fermat, which were collected and published by his son. Most authors on Algebra have also treated more or less on this part of it, but more especially Kersey, Prestet, Ozanam, Kirkby, &c. But afterwards, mathematicians seemed to have forgot such questions, if they did not even despise them as useless, when Euler drew their attention by some excellent compositions, demonstrating some general theorems, which had only been

known by induction. M. la Grange has also taken up the subject, having resolved very difficult problems in a general way, and discovered more direct methods than heretofore. The 2d volume of the French translation of Euler's Algebra contains an elementary treatise on this branch, and, with la Grange's additions, an excellent theory of it; treating very generally of Indeterminate problems, of the first and second degree, of solutions in whole numbers, of the method of Indeterminate coefficients, &c.

Finally, Mr. John Leslie has given, in the 2d volume of the Edinburgh Philos. Transactions, an ingenious paper on the resolution of Indeterminate problems, resolving them by a new and general principle. "The doctrine of Indeterminate equations," says Mr. Leslie, "has been seldom treated in a form equally systematic with the other parts of Algebra. The solutions commonly given are devoid of uniformity, and often require a variety of assumptions. The object of this paper is to resolve the complicated expressions which we obtain in the solution of Indeterminate problems, into simple equations, and to do so, without framing a number of assumptions, by help of a single principle, which though extremely simple, admits of a very extensive application."

"Let  $A \times B$  be any compound quantity equal to another,  $C \times D$ , and let  $m$  be any rational number assumed at pleasure; it is manifest that, taking equimultiples,  $A \times mB = C \times mD$ . If, therefore, we suppose that  $A = mD$ , it must follow that  $mB = C$ , or  $B = \frac{C}{m}$ . Thus two equations of a lower dimension are obtained. If these be capable of farther decomposition, we may assume the multiples  $n$  and  $p$ , and form four equations still more simple. By the repeated application of this principle, an higher equation admitting of divisors, will be resolved into those of the first order, the number of which will be one greater than that of the multiples assumed."

For example, resuming the problem at first given, viz, to find two rational numbers, the difference of the squares of which shall be a given number. Let the given number be the product of  $a$  and  $b$ ; then by hypothesis,  $x^2 - y^2 = ab$ ; but these compound quantities admit of an easy resolution, for  $x + y \times x - y = a \times b$ . If therefore we suppose  $x + y = ma$ , we shall obtain  $x - y = \frac{b}{m}$ ; where  $m$  is arbitrary, and if rational,  $x$  and  $y$  must also be rational. Hence the resolution of these two equations gives the values of  $x$  and  $y$ , the numbers sought, in terms of  $m$ ; viz,

$$x = \frac{m^2a + b}{2m}, \text{ and } y = \frac{m^2a - b}{2m}.$$

INDEX, in Arithmetic, is the same with what is otherwise called the characteristic or the exponent of a logarithm; being that which shews of how many places the absolute or natural number belonging to the logarithm consists, and of what nature it is, whether an integer or a fraction; the Index being less by 1 than the number of integer figures in the natural number, and is positive for integer or whole numbers, but negative in fractions, or in the denominator of a fraction;



tion; and in decimals, the negative index is 1 more than the number of ciphers in the decimal, after the point, and before the first significant figure; or, still more generally, the Index shews how far the first figure of the natural number is distant from the place of units, either towards the left hand, as in whole numbers, or towards the right, as in decimals; these opposite cases being marked by the correspondent signs + and -, of opposite affections, the sign - being set over the Index, and not before it, because it is this Index only which is understood as negative, and not the decimal part of the logarithm. Thus, in this logarithm  $2.4234097$ , the figures of whose natural number are 2651, the 2 is the Index, and being positive, it shews that the first figure of the number must be two places removed from the units place, or that there will be three places of integers, the number of these places being always 1 more than the Index; so that the natural number will be 2651. But if the same Index be negative, thus  $\overline{2}.4234097$ , it shews that the natural number is a decimal, and that the first significant figure of it is in the 2d place from units, or that there is one cipher at the beginning of the decimal, being 1 less than the negative Index; and consequently that the natural number of the logarithm in this case is .02651. Hence, by varying the natural number, with respect to the decimal places in it as in the former of the two columns here annexed, the Index of their logarithm will vary as in the 2d column.

Number.	Logarithm.
2651	3.4234097
2651	2.4234097
2651	1.4234097
2651	0.4234097
.2651	$\overline{1}.4234097$
.02651	$\overline{2}.4234097$
.002651	$\overline{3}.4234097$

Mr. Townly introduced a peculiar way of noting these Indices, when they become negative, or express decimal figures, which is now much in use, especially in the log. sines and tangents, &c, viz, by taking, instead of the true Index, its arithmetical complement to 10; so that, in this way, the logarithm  $2.4234097$  is written  $8.4234097$ .

For the addition and subtraction of Indices, see LOGARITHM.

INDEX of a Globe, is a little style fitted on to the north-pole, and turning round with it, pointing out the divisions of the hour-circle.

INDEX of a Quantity, in Arithmetic and Algebra, otherwise called the exponent, is the number that shews to what power it is understood to be raised: as in  $10^3$ , or  $a^3$ , the figure 3 is the Index or exponent of the power, signifying that the root or quantity, 10 or  $a$ , is raised to the 3d power. See this fully treated under EXPONENT.

INDICTION, or Roman INDICTION, a kind of epoch, or manner of counting time, among the Romans; containing a cycle or revolution of 15 years.

The popes have dated their acts by the year of the Indiction, which was fixed to the 1st of January anno Domini 313, ever since Charlemagne made them sovereign; before that time, they dated them by the years of the Emperors.

At the time of reforming the calendar, the year 1582 was reckoned the 10th year of the Indiction; so that beginning to reckon from hence, and dividing the number of years elapsed between that time and this, by 15, the remainder, with the addition of 10, rejecting 15 if the sum be more, will be the year of the Indiction.

But the Indiction will be easier found thus: Add 3 to the given year of Christ; divide the sum by 15, and the remainder after the division, will be the year of the Indiction: if there be no remainder, the Indiction is 15. In either of these ways, the Indiction for the year 1795 is 13.

INDIVISIBLES, are those indefinitely small elements, or principles, into which any body or figure may ultimately be divided.

A line is said to consist of points, a surface of parallel lines, and a solid of parallel surfaces: and because each of these elements is supposed Indivisible, if in any figure a line be drawn perpendicularly through all the elements, the number of points in that line, will be the same as the number of the elements.

Whence it appears, that a parallelogram, or a prism, or a cylinder, is resolvable into elements, as Indivisibles, all equal to each other, parallel, and like or similar to the base; for which reason, one of these elements multiplied by the number of them, that is the base of the figure multiplied by its height, gives the area or content. And a triangle is resolvable into lines parallel to the base, but decreasing in arithmetical progression; so also do the circles, which constitute the parabolic conoid, as well as those which constitute the plane of a circle, or the surface of a cone. In all which cases, as the last or least term of the arithmetic progression is 0, and the length of the figure the same thing as the number of the terms, therefore the greatest term, or base, being multiplied by the length of the figure, half the product is the sum of the whole, or the content of the figure.

And in any other figure or solid, if the law of the decrease of the elements be known, and thence the relation of the sum to the greatest term, which is the base, the whole number of them being the altitude of the figure, then the said sum of the elements is always the content of the figure.


A cylinder may also be resolved into cylindrical curve surfaces, having all the same height, and continually decreasing inwards, as the circles of the base do, on which they insist.

This way of considering magnitudes, is called the Method of Indivisibles, which is only the ancient method of exhaustions, a little disguised and contracted. And it is found of good use, both in computing the contents of figures in a very short and easy way, as above instanced, and in shortening other demonstrations in mathematics; an instance of which may here be given in that celebrated proposition of Archimedes, that a sphere is two-thirds of its circumscribed cylinder. Thus,

Suppose a cylinder, a hemisphere, and an inverted cone, having all the same base and altitude, and cut by an infinite number of planes all parallel to the base, of which EFGH is one; it is evident that the square of EI, the radius of the cylinder, is every where equal



to the square of SF, the radius of the sphere; and also that the square of EI, or of SF, is equal to the sum of the squares of IF and IS, or of IF and IK, because  $IK = IS$ ; that is,  $IE^2 = IF^2 + IK^2$ , in every position; but IE is the radius of the cylinder, IF the corresponding radius of the sphere, and IK that of the cone; and the circular sections of these bodies, are as the squares of their radii; therefore the section of the cylinder is every where equal to the sum of the sections of the hemisphere and cone; and, as the number of all those sections, which is the common height of the figures, is the same, therefore all the sections, or elements, of the cylinder, will be equal to the sum of all those of the hemisphere and cone taken together; that is, the cylinder is equal to both the hemisphere and cone: but as the cone itself is equal to one-third part of the cylinder; therefore the hemisphere is equal to the other two-thirds of it.



The Method of Indivisibles was introduced by Cavalerius, in 1635, in his *Geometria Indivisibilium*. The same was also pursued by Torricelli in his works, printed 1644: and again by Cavalerius himself in another treatise, published in 1647.

INERTIÆ *Vis*. See *Vis Inertiae*.

INFINITE, is applied to quantities which are either greater or less than any assignable ones. In which sense it differs but little from the terms Indefinite and Indeterminate. Thus, an

INFINITE, or *Infinitely great* line, denotes only an indefinite or indeterminate line; or a line to which no certain bounds or limits are prescribed.

INFINITE *Quantities*. Though the idea of magnitude infinitely great, or such as exceeds any assignable quantity, does include a negation of limits, yet all such magnitudes are not equal among themselves; but besides Infinite length, and Infinite area, there are no less than three several sorts of Infinite solidity; all of which are quantities *sui generis*; and those of each species are in given proportions.

Infinite length, or a line infinitely long, may be considered, either as beginning at a point, and so infinitely extended one way; or else both ways from the same point.

As to Infinite surface or area, any right line infinitely extended both ways on a plane infinitely extended every way, divides that plane into two equal parts, one on each side of the line. But if from any point in such a plane, two right lines be infinitely extended, making an angle between them; the Infinite area, intercepted between these Infinite right lines, is to the whole Infinite plane, as that angle is to 4 right angles. And if two Infinite and parallel lines be drawn at a given distance on such an Infinite plane, the area intercepted between them will be likewise Infinite; but yet it will be infinitely less than the whole plane; and even infinitely less than the angular or sectoral space, intercepted between two Infinite lines, that are inclined, though at never so small an angle; because in the one case, the given finite distance of the parallel lines diminishes the Infinity in one of the dimensions; whereas

in a sector, there is Infinity in both dimensions. And thus there are two species of Infinity in surfaces, the one infinitely greater than the other.

In like manner there are species of Infinites in solids, according as only one, or two, or as all their three dimensions, are Infinite; which, though they be all infinitely greater than a finite solid, yet are they in succession infinitely greater than each other.

Some farther properties of Infinite quantities are as follow:

The ratio between a finite and an Infinite quantity, is an Infinite ratio.

If a finite quantity be multiplied by an infinitely small one, the product will be an infinitely small one; but if the former be divided by the latter, the quotient will be infinitely great.

On the contrary, a finite quantity being multiplied by an infinitely great one, the product is infinitely great; but the former divided by the latter, the quotient will be infinitely little.

The product or quotient of an infinitely great or an infinitely little quantity, by a finite one, is respectively infinitely great, or infinitely little.

An infinitely great multiplied by an infinitely little, is a finite quantity; but the former divided by the latter, the quotient is infinitely Infinite.

The mean proportional between infinitely great, and infinitely little, is finite.

*Arithmetic of INFINITES*. See ARITHMETIC. Also Wallis's treatise of this subject; and another by Emerson, at the beginning of his *Conic Sections*; also Bulliald's treatise *Arithmetica Infinitorum*.

INFINITE *Decimals*, such as do not terminate, but go on without end; as  $.333 \text{ \&c} = \frac{1}{3}$ , or  $.142857 \text{ \&c} = \frac{1}{7}$ . See REPETEND.

INFINITELY *Infinite Fractions*, or all the powers of the fractions whose numerator is 1; which are all together equal to unity, as is demonstrated by Dr. Wood, in Hook's *Philos. Coll.* N<sup>o</sup> 3, p. 45; where some curious properties are deduced from the same.

INFINITE *Series*, a series considered as infinitely continued as to the number of its terms. See SERIES.

INFINITESIMALS, are certain infinitely or indefinitely small parts; as also the method of computing by them.

In the method of Infinitesimals, the element by which any quantity increases or decreases, is supposed to be infinitely small, and is generally expressed by two or more terms, some of which are infinitely less than the rest, which being neglected as of no importance, the remaining terms form what is called the *difference* of the proposed quantity. The terms that are neglected in this manner, as infinitely less than the other terms of the element, are the very same which arise in consequence of the acceleration, or retardation, of the generating motion, during the infinitely small time in which the element is generated; so that the remaining terms express the element that would have been produced in that time, if the generating motion had continued uniform. Therefore, those *differences* are accurately in the same ratio to each other, as the generating motions or fluxions. And hence, though in this method, Infinitesimal parts of the elements are neglected, the conclusions are accurately true, without



even an infinitely small error, and agree precisely with those that are deduced by the method of fluxions.

But however safe and convenient this method may be, some will always scruple to admit infinitely little quantities, and infinite orders of Infinitesimals, into a science that boasts of the most evident and accurate principles, as well as of the most rigid demonstrations. In order to avoid such suppositions, Newton considers the simultaneous increments of the flowing quantities as finite, and then investigates the ratio which is the limit of the various proportions which those increments bear to each other, while he supposes them to decrease together till they vanish; which ratio is the same with the ratio of the fluxions. See Maclaurin's Treatise of Fluxions, in the Introduc. p. 39 &c, also art. 495 to 502.

**INFLAMMABILITY**, that property of bodies by which they kindle, or catch fire.

**INFLECTION**, in Optics, called also Diffraction, and Deflection of the rays of light, is a property of them, by reason of which, when they come within a certain distance of any body, they will either be bent from it, or towards it; being a kind of imperfect reflection or refraction.



Some writers ascribe the first discovery of this property to Grimaldi, who first published an account of it, in his Treatise De Lumine, Coloribus, & Irade, printed in 1666. But Dr. Hook also claims the discovery of it, and communicated his observations on this subject to the Royal Society, in 1672. He shews that this property differs both from reflection and refraction; and that it seems to depend on the unequal density of the constituent parts of the ray, by which the light is dispersed from the place of condensation, and rarefied or gradually diverged into a quadrant; and this deflection, he says, is made towards the superficies of the opaque body perpendicularly.

Newton discovered, by experiments, this Inflection of the rays of light; which may be seen in his Optics.

M. De la Hire observed, that when we look at a candle, or any luminous body, with our eyes nearly shut, rays of light are extended from it, in several directions, to a considerable distance, like the tails of comets. The true cause of this phenomenon, which has exercised the sagacity of Des Cartes, Rohault, and others, seems to be, that the light passing among the eyelashes, in this situation of the eye, is inflected by its near approach to them, and therefore enters the eye in a great variety of directions. He also observes, that he found that the beams of the stars being observed, in a deep valley, to pass near the brow of a hill, are always more refracted than if there were no such hill, or the observation was made on the top of it; as if the rays of light were bent down into a curve, by passing near the surface of the mountain.

**Point of INFLECTION**, or of *contrary flexure*, in a curve, is the point or place in the curve where it begins to bend or turn a contrary way; or which separates the concave part from the convex part, and lying between the two; or where the curve changes from concave to convex, or from convex to concave, on the same side of the curve: such as the point E in the annexed figures; where the former of the two is con-

cave towards the axis AD from A to E, and convex from E to F; but, on the contrary, the latter figure is convex from A to E, and concave from E to F.



There are various ways of finding the point of Inflection; but the following, which is new, seems to be the simplest and easiest of all. From the nature of curvature it is evident that, while a curve is concave towards an axis, the fluxion of the ordinate decreases, or is in a decreasing ratio, with regard to the fluxion of the absciss; but, on the contrary, that the said fluxion increases, or is in an increasing ratio to the fluxion of the absciss, where the curve is convex towards the axis; and hence it follows that those two fluxions are in a constant ratio at the point of Inflection, where the curve is neither concave nor convex. That is, if  $x = AD$  the absciss, and  $y = DE$  the ordinate, then  $\dot{x}$  is to  $\dot{y}$  in a constant ratio, or  $\frac{\dot{x}}{\dot{y}}$  or  $\frac{\dot{y}}{\dot{x}}$  is a constant quantity. But constant quantities have no fluxion, or their fluxion is equal to nothing; so that in this case the fluxion of  $\frac{\dot{x}}{\dot{y}}$  or of  $\frac{\dot{y}}{\dot{x}}$  is equal to nothing. And hence we have this general rule: viz,

Put the given equation of the curve into fluxions; from which equation of the fluxions find either  $\frac{\dot{x}}{\dot{y}}$  or  $\frac{\dot{y}}{\dot{x}}$ ; then take the fluxion of this ratio or fraction, and put it equal to 0 or nothing; and from this last equation find also the value of the same  $\frac{\dot{x}}{\dot{y}}$  or  $\frac{\dot{y}}{\dot{x}}$ : then put this latter value equal to the former, which will be an equation from whence, and the first given equation of the curve,  $x$  and  $y$  will be determined, being the absciss or ordinate answering to the point of Inflection in the curve.

Or, putting the fluxion of  $\frac{\dot{x}}{\dot{y}}$  equal to 0, that is,  $\frac{\ddot{x}\dot{y} - \dot{x}\ddot{y}}{\dot{y}^2} = 0$ , or  $\ddot{x}\dot{y} - \dot{x}\ddot{y} = 0$ , or  $\ddot{x}\dot{y} = \dot{x}\ddot{y}$ , or  $\dot{x} : \dot{y} :: \ddot{x} : \ddot{y}$ , that is, the 2d fluxions have the same ratio as the 1st fluxions, which is a constant ratio; and therefore if  $\dot{x}$  be constant, or  $\ddot{x} = 0$ , then shall  $\ddot{y}$  be  $= 0$  also; which gives another rule, viz; Take both the 1st and 2d fluxions of the given equation of the curve, in which make both  $\dot{x}$  and  $\dot{y} = 0$ , and the resulting equations will determine the values of  $x$  and  $y$ , or absciss and ordinate answering to the point of Inflection.

For example, if it be required to find the point of Inflection in the curve whose equation is



$ax^2 = a^2y + x^2y$ . Now the fluxion of this is  $2ax\dot{x} = a^2\dot{y} + 2xy\dot{x} + x^2\dot{y}$ , which gives  $\frac{\dot{x}}{\dot{y}} = \frac{a^2 + x^2}{2ax - 2xy}$ .

Then the fluxion of this again made  $= 0$ , gives

$2x\dot{x} \cdot ax - xy = a^2 + x^2 \cdot a\dot{x} - y\dot{x} - x\dot{y}$ ; and this gives

again  $\frac{\dot{x}}{\dot{y}} = \frac{a^2 + x^2}{a^2 - x^2} \times \frac{x}{a - y}$ . Lastly, this value of  $\frac{\dot{x}}{\dot{y}}$

put  $=$  the former, gives

$\frac{a^2 + x^2}{a^2 - x^2} \cdot \frac{x}{a - y} = \frac{a^2 + x^2}{2x} \cdot \frac{1}{a - y}$ ; and hence

$2x^2 = a^2 - x^2$ , or  $3x^2 = a^2$ , and  $x = a\sqrt{\frac{1}{3}}$ , the absciss.

Hence also, from the original equation,

$y = \frac{ax^2}{a^2 + x^2} = \frac{\frac{2}{3}a^3}{\frac{4}{3}a^2} = \frac{1}{2}a$ , the ordinate to the point of Inflection sought.

When the curve has but one point of Inflection, it will be determined by a simple equation, as above; but when there are several points of Inflection, by the curve bending several times from the one side to the other, the resulting equation will be of a degree corresponding to them, and its roots will determine the absciss or ordinates to the same.

Other methods of determining the points of Inflection in curves, may be seen in most books on the doctrine of fluxions.

To know whether a curve be concave or convex towards any point assigned in the axis; find the value of  $\dot{y}$  at that point; then if this value be positive, the curve will be convex towards the axis, but if it be negative, it will be concave.

**INFORMED Stars**, or **INFORMES Stelle**, are such stars as have not been reduced into any constellation; otherwise called *Sporades*.—There was a great number of this kind left by the ancient astronomers; but Hevelius and some others of the moderns have provided for the greater part of them, by making new constellations.

**ENGINEER**. See **ENGINEER**.

**INGRESS**, in Astronomy, the sun's entrance into one of the signs, especially Aries.

**INNOCENTS Day**, a feast celebrated on the 28th day of December, in commemoration of the infants murdered by Herod.

**INORDINATE Proportion**, is where the order of the terms compared, is disturbed or irregular. As, for example, in two ranks of numbers, three in each rank, viz, in one rank, - - - 2, 3, 9, and in the other rank, - - - 8, 24, 36, which are proportional, the former to the latter, but in a different order, viz, - 2 : 3 :: 24 : 36,

and - 3 : 9 :: 8 : 24. then, casting out the mean terms in each rank, it is concluded that - - - 2 : 9 :: 8 : 36,

that is, the first is to the 3d in the first rank,

as the first is to the 3d in the 2d rank.

**INSCRIBED Figure**, is one that has all its angular points touching the sides of another figure in which the former is said to be inscribed.

**INSCRIBED Hyperbola**, is one that lies wholly within the angle of its asymptotes; as the common or conical hyperbola doth.

**INSTANT**, otherwise called a *Moment*, an infinitely small part of duration, or in which we perceive no succession, or which takes up the time of only one idea in our mind.

It is a maxim in mechanics, that no natural effect can be produced in an Instant, or without some definite time; also that the greater the time, the greater the effect. And hence may appear the reason, why a burthen seems lighter to a person, the faster he carries it; and why, the faster a person slides or scates on the ice, the less liable it is to break, or bend.

**INSULATE**, or **INSULATED**, a term applied to a column or other edifice, which stands alone, or free and detached from any adjacent wall, &c, like an island in the sea.

**INSULATED**, in Electricity, is a term applied to bodies that are supported by electrics, or non-conductors; so that their communication with the earth, by conducting substances, is interrupted.

**INTACTÆ**, are right lines to which curves do continually approach, and yet can never meet them more usually called *Asymptotes*.

**INTEGERS**, denote whole numbers: as contradistinguished from fractions.—Integers may be considered as numbers which refer to unity, as a whole to a part.

**INTEGRAL Number**, an integer; not a fraction.

**INTEGRAL Calculus**, in the New Analysis, is the reverse of the differential calculus, and is the finding the Integral from a given differential; being similar to the inverse method of fluxions, or the finding the fluent to a given fluxion.

**INTEGRANT Parts**, in *Philosophy*, are the similar parts of a body, or parts of the same nature with the whole; as filings of iron are the Integrant parts of iron, having the same nature and properties with the bar or mass they were filed off from.

**INTENSITY**, or **INTENSION**, in Physics, is the degree or rate of the power or energy of any quality; as heat, cold, &c. The Intensity of qualities, as gravity, light, heat, &c, vary in the reciprocal ratio of the squares of the distances from the centre of the radiating quality.

**INTERCALARY Day**, denotes the odd day inserted in the leap-year. See **BISSEXTILE**.

**INTERCEPTED Axis**, in Conic Sections, the same with what is otherwise called the absciss or abscissa.

**INTERCOLUMNATION**, or **INTERCOLUMNIATION**, is the space between column and column.

**INTEREST**, is a sum reckoned for the loan or forbearance of another sum, or principal, lent for, or due at, a certain time, according to some certain rate or proportion; being estimated usually at so much per cent. or by the 100. This forms a particular rule in Arithmetic. The highest legal Interest now allowed in England, is after the rate of 5 per cent. per annum, or the 20th part of the principal for the space of a year, and so in proportion for other times, either greater or less. Except in the case of pawn-brokers, to whom it has lately been made legal to take a higher interest, for one of the worst and most destructive purposes that can be suffered in any state.

Interest.



Another ingenious and general method of computing



puting Interest, is by the following small but comprehensive Table.

A General Interest Table,				
By which the Interest of any Sum, at any Rate, and for any Time, may be readily found.				
Days.	3 per Cent.	3½ per Cent.	4 per Cent.	4½ per Cent.
	l. s. d. q.	l. s. d. q.	l. s. d. q.	l. s. d. q.
1	1 3	2 1	2 2	3 0
2	3 3	4 2	5 1	6 0
3	5 3	6 3	7 3	8 3
4	7 3	9 0	10 2	11 3
5	9 3	11 2	1 1 1	1 2 3
6	11 3	1 1 3	1 3 3	1 5 3
7	1 1 3	1 4 0	1 6 1	1 8 3
8	1 3 3	1 6 1	1 9 0	1 11 3
9	1 5 3	1 8 2	1 11 2	2 2 2
10	1 7 2	1 11 0	2 2 1	2 5 2
20	3 3 1	3 10 0	4 4 2	4 11 1
30	4 11 0	5 9 0	6 6 3	7 4 2
40	6 6 3	7 8 0	8 9 0	9 10 1
50	8 2 2	9 7 0	10 11 2	12 3 3
60	9 10 1	11 6 0	13 1 3	14 9 2
70	11 6 0	13 5 0	15 4 0	17 3 1
80	13 1 3	15 4 0	17 6 1	19 8 3
90	14 9 2	17 3 0	19 8 2	1 2 2 1
100	16 5 1	19 2 0	1 11 0	1 4 8 0
200	1 12 10 2	1 18 4 1	2 3 10 0	2 9 3 3
300	2 9 3 3	2 17 6 1	3 5 9 0	3 13 11 3

N. B. This Table contains the interest of 100l. for all the several days in the 1st column, and at the several rates of 3, 3½, 4, 4½, and 5 per cent. in the other 5 columns.

To find the Interest of 100l. for any other time, as 1 year and 278 days, at 4½ per cent. Take the sums for the several days as here below.

The Int. for 1 year	4	10	0	0
Against 200 ds. is	2	9	3	3
70 ds. -	0	17	3	1
8 ds. -	0	1	11	0

Interest required - 7 18 6 0

For any other Sum than 100l. First find for 100l. as above, and take it so many times or parts as the sum is of 100l. Thus, to find for 355l. at 4½, for 1 year and 278 days.

First, 3 times the above sum,	
(for 300l.) is -	23 15 8 1
½ (for 50l.) is -	3 19 3 1
10 of this (for 5l.)	0 7 11 0

So for 355 it is - 28 2 10 2

When the interest is required for any other rate than those in the Table, it may be easily made out from them. So ½ of 5 is 2½, ½ of 4 is 2, ½ of 3 is 1½, ⅓ of 3 is 1, 1-6th of 3 is ½, and 1-12th of 3 is ¼. And so, by parts, or by adding or subtracting, any rate may be made out.

Compound Interest, called also *Interest-upon-Interest*, is that which is counted not only upon the principal sum lent, but also for its Interest, as it becomes due, at the end of each stated time of payment.

Although it be not lawful to lend money at Compound Interest, yet in purchasing annuities, pensions,

&c, and taking leases in reversion, it is usual to allow Compound Interest to the purchaser for his ready money; and therefore it is very necessary to understand this subject.

Besides the quantities concerned in Simple Interest, viz, the principal  $p$ , the rate or Interest of 1l. for 1 year  $r$ , the amount  $a$ , and the time  $t$ , there is another quantity employed in Compound Interest, viz, the ratio of the rate of Interest, which is the amount of 1l. for 1 time of payment, and which here let be denoted by  $R$ , viz,  $R = 1 + r$ . Then, the particular amounts for the several times may be thus computed, viz, As 1 pound is to its amount for any time, so is any proposed principal sum to its amount for the same time; i. e.

1l. :  $R$  ::  $p$  :  $pR$  the 1st year's amount,

1l. :  $R$  ::  $pR$  :  $pR^2$  the 2d year's amount,

1l. :  $R$  ::  $pR^2$  :  $pR^3$  the 3d year's amount,

and so on.

Therefore in general,  $pR^t = a$  is the amount for the  $t$  year, or  $t$  time of payment. From whence the following general theorems are deduced :

1st,  $a = pR^t$  the amount,

2d,  $p = \frac{a}{R^t}$  the principal,

3d,  $R = \sqrt[t]{\frac{a}{p}}$  the ratio,

4th,  $t = \frac{\log. \text{ of } a - \log. \text{ of } p}{\log. \text{ of } R}$  the time.

From which any one of the quantities may be found, when the rest are given.

For example, suppose it were required to find in how many years any principal sum will double itself, at any rate of Interest. In this case we must employ the 4th theorem, where  $a$  will be  $= 2p$ , and then it

$$\text{is } t = \frac{1. a - 1. p}{\log. R} = \frac{1. 2p - 1. p}{\log. R} = \frac{\log. 2}{\log. R}.$$

So, if the rate of Interest be 5 per cent. per annum; then  $R = 1 + .05 = 1.05$ , and hence

$$t = \frac{\log. 2}{\log. 1.05} = \frac{.3010300}{.0211893} = 14.2067 \text{ nearly;}$$

that is, any sum doubles in 14½ years nearly, at the rate of 5 per cent. per annum Compound Interest.

Hence, and from the like question in Simple Interest, above given, are deduced the times in which any sum doubles itself, at several rates of Interest, both simple and compound : viz,

At	At Simp. Int. Years.	At Comp. Int. Years.
2	50	35.0028
2½	40	28.0701
3	33½	23.4498
3½	28¾	20.1488
4	25	17.6730
4½	22½	15.7473
5	20	14.2067
6	16⅔	11.8957
7	14⅔	10.2448
8	12½	9.0065
9	11⅔	8.0432
10	10	7.2725

The



The following Table will facilitate the calculation of Compound Interest for any sum, and any number of years, at various rates of Interest.

The Amount of 1 <i>l.</i> in any Number of Years.						
Yrs.	3	3½	4	4½	5	6
1	1.0300	1.0350	1.0400	1.0450	1.0500	1.0600
2	1.0609	1.0712	1.0816	1.0920	1.1025	1.1236
3	1.0927	1.1087	1.1249	1.1412	1.1576	1.1910
4	1.1255	1.1475	1.1699	1.1925	1.2155	1.2625
5	1.1593	1.1877	1.2167	1.2462	1.2763	1.3382
6	1.1941	1.2293	1.2653	1.3023	1.3401	1.4185
7	1.2299	1.2723	1.3159	1.3609	1.4071	1.5036
8	1.2668	1.3168	1.3686	1.4221	1.4775	1.5939
9	1.3048	1.3629	1.4233	1.4861	1.5513	1.6895
10	1.3439	1.4106	1.4802	1.5530	1.6289	1.7909
11	1.3842	1.4600	1.5395	1.6229	1.7103	1.8983
12	1.4258	1.5111	1.6010	1.6959	1.7959	2.0122
13	1.4685	1.5640	1.6651	1.7722	1.8856	2.1329
14	1.5126	1.6187	1.7317	1.8519	1.9799	2.2609
15	1.5580	1.6753	1.8009	1.9353	2.0789	2.3966
16	1.6047	1.7340	1.8730	2.0224	2.1829	2.5404
17	1.6528	1.7947	1.9479	2.1134	2.2920	2.6928
18	1.7024	1.8575	2.0258	2.2085	2.4066	2.8543
19	1.7535	1.9225	2.1068	2.3079	2.5270	3.0256
20	1.8061	1.9898	2.1911	2.4117	2.6533	3.2071

The use of this Table, which contains all the powers  $R^t$ , to the 20th power, or the amounts of 1*l.* is chiefly to calculate the Interest, or the amount, of any principal sum, for any time, not more than 20 years. For example, required to find to how much 523*l.* will amount in 15 years, at the rate of 5*l.* per cent. per annum Compound Interest.

In the Table, on the line 15 and column 5 per cent, is the amount of 1*l.* viz. 2.0789,  
this multiplied by the principal 523,

gives the amount 1087.2647  
or 1087*l.* 5*s.* 3¼*d.*  
and therefore the Interest is 564*l.* 5*s.* 3¼*d.*



See Annuities; Discount; Reversion; Smart's Tables of Interest; the Philos. Transf. vol. 6, p. 508; and most books on Arithmetic.

INTERIOR Figure, Angle of. See ANGLE.

INTERIOR Polygon. See POLYGON.

INTERIOR Talus. See TALUS.

INTERNAL Angles, are all angles made within any figure, by the sides of it. In a triangle ABC, the two



angles A and C are peculiarly called Internal and oppo-

site, in respect of the external angle CBD, which is equal to them both together.

INTERNAL Angle is also applied to the two angles formed between two parallels, by a line intersecting those parallels, on each side of the intersecting line. Such are the angles  $a, b, c, d$ , formed between the parallels EF and GH, on each side of the intersecting line.—The two adjacent Internal angles  $a$  and  $b$ , or  $c$  and  $d$ , are together equal to two right angles.

INTERNAL and Opposite Angles, is also applied to the two angles  $a$  and  $b$ , which are respectively equal to the two  $n$  and  $m$ , called the external and opposite angles.

Also the alternate Internal angles are equal to one another; viz,  $a = d$ , and  $b = c$ .

INTERPOLATION, in the modern Algebra, is used for finding an intermediate term of a series, its place in the series being given.

The Method of Interpolation was first invented by Mr. Briggs, and applied by him to the calculation of logarithms, &c, in his Arithmetica Logarithmica, and his Trigonometria Britannica; where he explains, and fully applies the method of Interpolation by differences. His principles were followed by Reginal and Mouton in France, and by Cotes and others in England. Wallis made use of the method of Interpolation in various parts of his works; as his Arithmetic of Infinites, and his Algebra, for quadratures, &c. The same was also happily applied by Newton in various ways: by it he investigated his binomial theorem, and quadratures of the circle, ellipse, and hyperbola: see Wallis's Algebra, chap. 85, &c. Newton also, in lemma 5, lib. 3 Princip. gave a most elegant solution of the problem for drawing a curve line through the extremities of any number of given ordinates; and in the subsequent proposition, applied the solution of this problem to that of finding from certain observed places of a comet, its place at any given intermediate time. And Dr. Waring, who adds, that a solution still more elegant, on some accounts, has been since discovered by Mess. Nichol and Stirling, has also resolved the same problem, and rendered it more general, without having recourse to finding the successive differences. Philos. Transf. vol. 69, part 1, art. 7.

Mr. Stirling indeed pursued this branch as a distinct science, in a separate treatise, viz, Tractatus de Summatione et Interpolatione Serierum Infinitarum, in the year 1730.

When the 1st, 2d, or other successive differences of the terms of a series become at last equal, the Interpolation of any term of such a series may be found by Newton's Differential Method.

When the Algebraic equation of a series is given, the term required, whether it be a primary or intermediate one, may be found by the resolution of affected equations; but when that equation is not given, as it often happens, the value of the term sought must be exhibited by a converging series, or by the quadrature of curves. See Stirling, ut supra, p. 86. Meyer, in Act. Petr. tom. 2, p. 180.

A general theorem for Interpolating any term is as follows: Let  $A$  denote any term of an equidistant series of terms, and  $a, b, c$ , &c, the first of the 1st, 2d, 3d, &c orders of differences; then the term  $x$ , whose distance



distance from A is expressed by  $x$ , will be this, viz,  
Theorem 1,

$$z = A + xa + x \cdot \frac{x-1}{2} b + x \cdot \frac{x-1}{2} \cdot \frac{x-2}{3} c \&c.$$

Hence, if any of the orders of differences become equal to one another, or  $= 0$ , this series for the interpolated term will break off, and terminate, otherwise it will run out in an infinite series.

Ex. To find the 20th term of the series of cubes 1, 8, 27, 64, 125, &c, or  $1^3, 2^3, 3^3, 4^3, 5^3, \&c.$

Set down the series in a column, and take their continual differences as here annexed, where the 4th differences, and all after it become  $= 0$ , also  $A = 1$ ,  $a = 7$ ,  $b = 12$ ,  $c = 6$ , and  $x = 19$ ; therefore the 20th term sought is barely

A	a	b	c	d	e
1	7	12	6	0	
8	19	18	6	0	
27	37	24	6	0	
64	61	24	6	0	
125					

$$z = 1 + 19 \times 7 + 19 \times \frac{18}{2} \times 12 + 19 \cdot \frac{18}{2} \cdot \frac{17}{3} \cdot 6$$

$$= 1 + 133 + 2052 + 5814 = 8000.$$

Theor. 2. In any series of equidistant terms,  $a, b, c, d, \&c$ , whose first differences are small; to find any term wanting in that series, having any number of terms given. Take the equation which stands against the number of given terms, in the following Table; and by reducing the equation, that term will be found.

No.	Equations.
1	$a - b = 0$
2	$a - 2b + c = 0$
3	$a - 3b + 3c - d = 0$
4	$a - 4b + 6c - 4d + e = 0$
5	$a - 5b + 10c - 10d + 5e - f = 0$
6	$a - 6b + 15c - 20d + 15e - 6f + g = 0$
&c.	&c.
n	$a - nb + n \cdot \frac{n-1}{2} c - n \cdot \frac{n-1}{2} \cdot \frac{n-2}{3} d \&c = 0.$

where it is evident that the coefficients in any equation, are the uncia of a binomial  $1 + 1$  raised to the power denoted by the number of the equation.

Ex. Given the logarithms of 101, 102, 104, and 105; to find the log. of 103.

Here are 4 quantities given; therefore we must take the 4th equation  $a - 4b + 6c - 4d + e = 0$ , in which it is the middle quantity or term  $c$  that is to be found, because 103 is in the middle among the numbers 101, 102, 104, 105; then that equation gives the

$$\text{value of } c \text{ as follows, viz } c = \frac{4 \cdot \overline{b + d} - \overline{a + e}}{6}.$$

Now the logs. of the given numbers will be thus:

$$\begin{array}{r} 2.0043214 = a \\ 2.0086002 = b \\ 2.0170333 = d \\ 2.0211893 = e \\ \hline 4.0256335 = b + d \\ \quad \quad 4 \\ \hline 16.1025340 = 4 \cdot \overline{b + d} \\ \text{subtr. } 4.0255107 = a + e \\ \hline 6) 12.0770233 \\ 2.0128372 \text{ the log. of } 103. \end{array}$$

Theor. 3. When the terms  $a, b, c, d, \&c$ , are at unequal distances from each other; to find any intermediate one of these terms, the rest being given.

Let  $p, q, r, s, \&c$ , be the several distances of those terms from each other; then let

$$B = \frac{b-a}{p}, C = \frac{B_1-B}{p+q}, D = \frac{C_1-C}{p+q+r}, E = \frac{D_1-D}{p+q+r+s}$$

$$B_1 = \frac{c-b}{q}, C_1 = \frac{B_2-B_1}{q+r}, D_1 = \frac{C_2-C_1}{q+r+s}$$

$$B_2 = \frac{d-c}{r}, C_2 = \frac{B_3-B_2}{r+s}$$

$$B_3 = \frac{e-d}{s},$$

&c                      &c                      &c

Then the term  $z$ , whose distance from the beginning is  $x$ , will be

$$z = a + Bx + \frac{Cx \cdot \overline{x-p}}{2} + \frac{Dx \cdot \overline{x-p} \cdot \overline{x-p-q}}{6} + \frac{Ex \cdot \overline{x-p} \cdot \overline{x-p-q} \cdot \overline{x-p-q-r}}{24} + \&c;$$

to be continued to as many terms as there are terms in the given series.

By this series may be found the place of a comet, or the sun, or any other object at a given time; by knowing the places of the same for several other given times.

Other methods of Interpolation may be found in the Philos. Transf. number 362; or Stirling's Summation and Interpolation of Series.

INTERSCENDENT, in Algebra, is applied to quantities, when the exponents of their powers are radical quantities. Thus  $x^{\sqrt{2}}, x^{\sqrt{a}}, \&c$ , are interscendent quantities. See FUNCTION.

INTERSECTION, the cutting of one line, or plane, by another; or the point or line in which two lines or two planes cut each other.—The mutual intersection of two planes is a right line. The centre of a circle, or conic section, &c, is in the intersection of two diameters; and the central point of a quadrangle is the Intersection of two diagonals.

INTERSTELLAR, a word used by some authors, to express those parts of the universe, that are without and beyond the limits of our solar system.

In the Interstellar regions, it is supposed there are several other systems of planets moving round the fixed stars, as the centres of their respective motions. And if it be true, as it is not improbable, that each fixed star is thus



thus a sun to some habitable orbs, or earths, that move round it, the Interstellar world will be infinitely the greatest part of the universe.

**INTERTIES**, or **INTERDUCES**, in Architecture, those small pieces of timber which lie horizontally between the summers, or between them and the cell or raising plate.

**INTERVAL**, in Music, the difference between two sounds, in respect of acute and grave. Authors distinguish several divisions of an Interval, as first into Simple and Compound. The

*Simple* INTERVAL is that without parts, or division; such are the octave, and all that are within it; as the 2d, 3d, 4th, 5th, 6th, and 7th, with their varieties.

*Compound* INTERVAL consists of several lesser Intervals; as the 9th, 10th, 11th, 12th, &c, with their varieties.

This Simple Interval was by the ancients called a Diastem, and the Compound they called a System.

An Interval is also divided into Just or True, and into False. The

*Just* or *True* INTERVALS, are such as all those above mentioned, with their varieties, whether major or minor. And the

*False* INTERVALS, are the diminutive or superfluous ones.

An Interval is also divided into the Consonance and Dissonance; which see.

**INTESTINE** *Motion of the parts of fluids*, that which is among its corpuscles or component parts.

When the attracting corpuscles of any fluid are elastic, they must necessarily produce an intestine motion; and that, greater or less, according to the degrees of their elasticity and attractive force. For, two elastic particles, after meeting, will fly from each other, with the same degree of velocity with which they met; abstracting from the resistance of the medium. But when, in leaping back from each other, they approach other particles, their velocity will be increased.

**INTRADOS**, the interior and lower side, or curve, of the arch of a bridge, &c. In contradistinction from the extrados, or exterior curve, or line on the upper side of the arch. See my Treatise on Bridges &c. prop. 5.

**INVERSE**, is applied to a manner of working the rule of three, or proportion, which seems to go backward, i. e. reverse or contrary to the order of the common and direct rule: So that, whereas, in the direct rule, more requires more, or less requires less; in the Inverse rule, on the contrary, more requires less, or less requires more.

For instance, in the direct rule it is said, If 3 yards of cloth cost 20 shillings, how much will 6 yards cost? the answer is 40 shillings: where more yards require more money, and less yards require less money. But in the Inverse rule it is said, If 20 men perform a piece of work in 4 days, in how many days will 40 men perform as much? where the answer is 2 days; and here the more men require the less time, and the fewer men the more time.

**INVERSE Method of Fluxions**, is the method of finding fluents, from the fluxions being given; and is similar to what the foreign mathematicians call the Calculus Integralis. See FLUENTS.

**INVERSE Method of Tangents**, is the method of finding the curve belonging to a given tangent; as opposed to the direct method, or the finding the tangent to a given curve.

As, to find a curve whose subtangent is a third proportional to  $x - y$  and  $y$ , or whose subtangent is equal to the semiordinate, or whose subnormal is a constant quantity.—The solution of this problem depends chiefly on the Inverse method of fluxions. See TANGENT.

**INVERSE Proportion**, or **INVERSE Ratio**, is that in which more requires less, or less requires more. As for instance, in the case of light, or heat from a luminous object, the light received is less at a greater distance, and greater at a less distance; so that here more, as to distance, gives less, as to light, and less distance gives more light. This is usually expressed by the term Inversely, or Reciprocally; as in the case above, where the light is Inversely, or Reciprocally as the square of the distance; or in the Inverse or Reciprocal duplicate ratio of the distance.

**INVERSION**, *Invertendo*, or *by Inversion*, according to the 14th definition of Euclid, lib. 5, is Inverting the terms of a proportion, by changing the antecedents into consequents, and the consequents into antecedents. As in these,  $a : b :: c : d$ , then by Inversion  $b : a :: d : c$ .

**INVESTIGATION**, the searching or finding any thing out, by means of certain steps, traces, or ways.

**INVOLUTE** *Figure* or *Curve*, is that which is traced out by the outer extremity of a string as it is folded or wrapped upon another figure, or as it is unwound from off it.—The Involute of a cycloid, is also a cycloid equal to the former, which was first discovered by Huygens, and by means of which he contrived to make a pendulum vibrate in the curve of a cycloid, and so theoretically at least vibrate always in equal times whether the arch of vibration were great or small, which is a property of that curve.—For the doctrine and nature of Involute and Evolute, see EVOLUTE.

**INVOLUTION**, in Arithmetic and Algebra, is the raising of powers from a given root; as opposed to Evolution, which is the extracting, or developing of roots from given powers. So the Involution of the number 3, or its powers, are thus raised:

3        -        or  $3^1$  or 3 is the 1st power, or root,  
 $3 \times 3$        or  $3^2$  or 9 is the 2d power, or square,  
 $3 \times 3 \times 3$  or  $3^3$  or 27 is the 3d power, or cube,  
 and so on.

And hence, to find any power of a given root, or quantity, let the root be multiplied by itself a number of times which is one less than the number of the index; i. e. once multiplied for the 2d root, twice for the 3d root, thrice for the 4th root, &c.

Thus, to Involve 12 to the 3d power.

12  
 ———  
 0144 square, or 2d power  
 12  
 ———  
 001728 cube, or 3d power.



So also, in Algebra, to Involve the binomial  $a + b$ , or raise its powers.

$$\begin{array}{r} a + b \\ \hline a + b \end{array} \quad \text{1st power, or root}$$

$$\begin{array}{r} a^2 + ab \\ + ab + b^2 \\ \hline \end{array}$$

$$\begin{array}{r} a^2 + 2ab + b^2 \\ \hline a + b \end{array} \quad \text{2d power}$$

$$\begin{array}{r} a^3 + 2a^2b + 2ab^2 \\ + a^2b + ab^2 + b^3 \\ \hline \end{array}$$

$$\begin{array}{r} a^3 + 3a^2b + 3ab^2 + b^3 \\ \hline a + b \end{array} \quad \text{3d power}$$

$$\begin{array}{r} a^4 + 3a^3b + 3a^2b^2 + ab^3 \\ + a^3b + 3a^2b^2 + 3ab^3 + b^4 \\ \hline \end{array}$$

$$a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4 \quad \text{4th power}$$

And in like manner for any other quantities, whatever the number of their terms may be. But compound algebraic quantities are best involved by the BINOMIAL Theorem; which see.

Simple quantities are Involved, by raising the numeral coefficients to the given power, and the literal quantities are raised by multiplying their indices by that of the root; that is, the raising of powers is performed by the multiplication of indices, the same as the multiplication of logarithms. Thus,

The 2d power of  $a$  is  $a^2$ .

The 2d power of  $2a^2$  is  $2^2a^2 \times 2$  or  $4a^4$ .

The 3d power of  $3a^2b^3$  is  $27a^6b^9$ .

The 3d power of  $a^{\frac{1}{2}}b^{\frac{2}{3}}$  is  $a^{\frac{3}{2}}b^2$ .

The  $n$ th power of  $a^m c^p$  is  $a^{mn} c^{pn}$  or  $a^{\overline{m}n} c^{\overline{p}n}$ .

INWARD Flanking Angle, in Fortification, is that made by the curtain and the rasant flanking line of defence.

JOINTS, in Architecture, are the separations between the stones or bricks; which may be filled with mortar, plaster, or cement.

JOINT, in Carpentry, &c, is applied to several manners of assembling, setting, or fixing pieces of wood together. As by a mortise Joint, a dove-tail Joint, &c.

Universal JOINT, in Mechanics, an excellent invention of Dr. Hook, adapted to all kinds of motions and flexures; of which he has given a large account in his Cutlerian Lectures, printed in 1678. This seems to have given occasion to the gimboles used in suspending the sea compasses; the mechanism of which is the same with that of Defaguliers's rolling lamp.

JOINT-Lives, are such as continue during the same time, or that exist together. See LIFE-Annuities.

JOISTS, or JOYSTS, those pieces of timber framed into the girders and summers, and on which the boarding of floors is laid.

"JONES (WILLIAM), F. R. S. a very eminent mathematician, was born at the foot of Bodavon mountain [Mynydd Bodafon] in the parish of Llanfihangel tre'r Bard, in the Isle of Anglesey, North Wales, in the year 1675. His father's name was John George,

his surname being the proper name of his father. For it is a custom in several parts of Wales for the proper name of the father to become the surname of his children. John George the father was commonly called Sion Siors of Llanbabo, to which place he moved, and where his children were brought up. Accordingly our author, whose proper name was William, took the surname of Jones from the proper name of his father, who was a farmer, and of a good family, being descended from Hwfa ap Cynddelw, one of the 15 tribes of North Wales. He gave his two sons the common school education of the country, reading, writing, and accounts, in English, and the Latin Grammar. Harry his second son took to the farming business; but William the eldest, having an extraordinary turn for mathematical studies, determined to try his fortune abroad from a place where the same was but of little service to him. He accordingly came to London, accompanied by a young man, Rowland Williams, afterwards an eminent perfumer in Wych-street. The report in the country is, that Mr. Jones soon got into a merchant's counting house, and so gained the esteem of his master, that he gave him the command of a ship for a West India voyage; and that upon his return he set up a mathematical school, and published his book of Navigation; and that upon the death of the merchant he married his widow: that, Lord Macclesfield's son being his pupil, he was made secretary to the chancellor, and one of the deputy tellers of the exchequer:—and they have a story of an Italian wedding which caused great disturbance in Lord Macclesfield's family, but was compromised by Mr. Jones; which gave rise to a saying, "that Macclesfield was the making of Jones, and Jones the making of Macclesfield." The foregoing account of Mr. Jones, I found among the papers of the late Mr. John Robertson, librarian and clerk to the Royal Society, who had been well known to Mr. Jones, and possessed many of his papers.

Mr. Jones having by his industry acquired a competent fortune, lived upon it as a private gentleman for many years, in the latter part of his life, in habits of intimacy with Sir Isaac Newton and others the most eminent mathematicians and philosophers of his time; and died July 3, 1749, at 74 years of age, being one of the vice-presidents of the Royal Society; leaving at his death one daughter, and his widow with child, which proved a son, who is the present Sir William Jones, now one of the judges in India, and highly esteemed for his great abilities, extensive learning, and eminent patriotism.—Mr. Jones's publications are,

1. *A new Compendium of the Whole Art of Navigation*, &c; in small 8vo, London, 1702. This is a neat little piece, and dedicated to the Rev. Mr. John Harris, the same I believe who was author of the *Lexicon Technicum*, or Universal Dictionary of Arts and Sciences, in whose house Mr. Jones says he composed his book.

2. *Synopsis Palmariorum Matheseos*: Or a New Introduction to the Mathematics, &c; 8vo, London, 1706. Being a very neat and useful compendium of all the mathematical sciences, in about 300 pages.

His papers in the *Philos. Transf.* are the following:

3. *A Compendious Disposition of Equations for exhibiting the relations of Goniometrical Lines*; vol. 44, p. 560.

4 N 2

4. A



4. A Tract on Logarithms ; vol. 61, pa. 455.

5. Account of the person killed by lightning in Tottenham-Court-Chapel, and its effects on the building ; vol. 62, pa. 131.

6. Properties of the Conic Sections, deduced by a compendious method ; vol. 63, pa. 340.

In all these works of Mr. Jones, a remarkable neatness, brevity, and accuracy, every where prevails. He seemed to delight in a very short and comprehensive mode of expression and arrangement ; in so much that sometimes what he has contrived to express in two or three pages, would occupy a little volume in the ordinary style of writing.

Mr. Jones it is said possessed the best mathematical library in England ; scarcely any book of that kind but what was there to be found. He had collected also a great quantity of manuscript papers and letters of former mathematicians, which have often proved useful to writers of their lives, &c. After his death, these were dispersed, and fell into different persons hands ; many of them, as well as of Mr. Jones's own papers, were possessed by the late Mr. John Robertson, before mentioned, at whose death I purchased a considerable quantity of them. From such collections of these it was that Mr. Jones was enabled to give that first and elegant edition, in 4to, 1711, of several of Newton's papers, that might otherwise have been lost, intitled, *Analysis per quantitatum Series, Fluxiones, ac Differentias : cum Enumeratione Linearum Tertii Ordinis*.

IONIC Column, or Order, the 3d of the five orders, or columns, of Architecture. The first idea of this order was given by the people of Ionia ; who, according to Vitruvius, formed it on the model of a young woman, dressed in her hair, and of an easy elegant shape, as the Doric had been formed on the model of a strong robust man.

This column is a medium between the massive and the more delicate orders, the simple and the rich. It is distinguished from the Composite, by having none of the leaves of acanthus in its capital ; and from the Tuscan, Doric, and Corinthian, by the volutes, or rams horns, which adorn its capital ; and from the Tuscan and Doric too, by the channels, or fluting, in its shaft.

The height of this column is 18 modules, or 9 diameters of the column taken at the bottom : indeed at first its height was but 16 modules ; but, to render it more beautiful than the Doric, its height was augmented by adding a base to it, which was unknown in the Doric. M. le Clerc makes its entablature to be 4 modules and 10 minutes, and its pedestal 6 modules ; so that the whole order makes 28 modules 10 minutes.

JOURNAL, in Merchants Accounts, is a book into which every particular article is posted out of the Waste-book, according to the order of time, specifying the debtor and creditor in each account and transaction.

JOURNAL, in Maritime Affairs, is a register kept by the pilot, and others, noticing every thing that happens to the ship, from day to day, and from hour to hour, with regard to the winds, the rhumbs or courses, the knots or rate of running, the rake, soundings, astronomical observations, for the latitudes and longitudes, &c ; to enable them to adjust the reckoning, and determine the place where the ship is.

In all sea Journals, the day, or what is called the 24 hours, is divided into twice 12 hours, those before

noon marked A. M. for ante meridiem, and those from noon to midnight marked P. M. post meridiem, or afternoon.

There are various ways of keeping a sea Journal, according to the different notions of mariners concerning the articles that are to be entered. Some writers direct the keeping such a kind of Journal as is only an abstract of each day's transactions, specifying the weather, what ships or lands were seen, accidents on board, the latitude, longitude, meridional distance, course, and run : these particulars are to be drawn from the ship's log-book, or from that kept by the person himself. Other authors recommend the keeping only of one account, including the log-book, and all the work of each day, with the deductions drawn from it.

JOURNAL is also used for the title of several books which come out at stated times ; and give accounts and abstracts of the new books that are published, with the new improvements daily made in arts and sciences.

The first Journal of this kind was, the Journal des Sçavans, printed at Paris : the design was set on foot for the ease of such as are too busy, or too lazy, to read the entire books themselves. It seems an excellent way of satisfying a man's curiosity, and becoming learned upon easy terms : and so useful has it been found, that it has been executed in most other countries, though under a great variety of titles.

Of this kind are the Acta Eruditorum of Leipzig ; the Nouvelles de la Republique des lettres of Mr. Bayle, &c ; the Bibliothque Universelle, Choisie, et Ancienne et Moderne, of M. le Clerc ; the Memoirs de Trevoux, &c. In 1692, Juncker printed in Latin, An Historical Treatise of the Journals of the Learned, published in the several parts of Europe ; and Wolfius, Struvius, Morhoff, Fabricius, &c, have done something of the same kind.

The Philosophical Transactions of London ; the Memoirs of the Royal Academy of Sciences ; those of the Academy of Belles Lettres ; the Miscellanea Naturæ Curiosorum ; the Experiments of the Academy del Cimento, the Acta Philo-exoticorum Naturæ et Artis, which appeared from March 1686 to April 1687, and which are a history of the Academy of Bresse ; the Miscellanea Berolinensia, or Memoirs of the Academy of Berlin ; the Commentaries of the Academy of Petersburg ; the Memoirs of the Institute at Bologna ; the Acta Literaria Sueciæ ; the Memoirs of the Royal Academy of Stockholm, begun in 1740 ; the Commentarii Societatis Regiæ Gottingensis, begun in 1750, &c, &c, are not so properly Journals, though they are frequently ranked in the number.

Juncker and Wolfius give the honour of the first invention of Journals to Photius. His Bibliotheca, however, is not altogether of the same nature with the modern Journals ; nor was his design the same. It consists of abridgments, and extracts of books which he had read during his embassy in Persia. M. Salo first began the Journal des Sçavans at Paris, in 1665, under the name of the Sieur de Hedonville ; but his death soon after interrupted the work. The abbé Gallois then took it up, and he, in the year 1674, gave way to the abbé de la Roque, who continued it nine years, and was succeeded by M. Cousin, who carried it on till the year 1702, when the abbé Bignon instituted



instituted a new Society, and committed the care of continuing the Journal to them, who improved and published it under a new form. This Society is still continued, and M. de Loyer has had the inspection of the Journal; which is no longer the work of any single author, but of a great number.

The other French Journals are the *Memoirs and Conferences of Arts and Sciences*, by M. Dennis, during the years 1672, 1673, and 1674; *New Discoveries in all the parts of Physic*, by M. de Blegny; the *Journal of Physic*, begun in 1684, and some others, discontinued almost as soon as begun.

Rozier's *Journal de Physique*, begun in July 1771, and continued, till in the year 1780, there were 19 vols. quarto.

The *Nouvelles de la Republique des Lettres*, News from the Republic of Letters, were begun by M. Bayle in 1684, and carried on by him till the year 1687, when M. Bayle being disabled by sickness, his friends, M. Bernard and M. de la Roque, took them up, and continued them till 1699. After an interruption of nine years, M. Bernard resumed the work, and continued it till the year 1710. The *History of the Works of the Learned*, by M. Basnage, was begun in the year 1686, and ended in 1710. The *Universal Historical Library*, by M. le Clerc, was continued to the year 1693, and contained twenty-five volumes. The *Bibliothèque Choisie* of the same author, began in 1703. The *Mercury of France* is one of the most ancient Journals of that country, and is continued by different hands: the *Memoirs of a History of Sciences and Arts*, usually called *Memoires des Trevoux*, from the place where they are printed, began in 1701. The *Essays of Literature* reached but to a twelfth volume in 1702, 1703, and 1704; these only take notice of ancient authors. The *Journal Litteraire*, by Father Hugo, began and ended in 1705. At Hamburgh they have made two attempts for a French Journal, but the design failed: an *Ephemerides Scavantes* has also been undertaken, but that soon disappeared. A *Journal des Scavans*, by M. Dartis, appeared in 1694, and was dropt the year following. That of M. Chauvin, begun at Berlin in 1696, held out three years; and an essay of the same kind was made at Geneva. To these may be added, the *Journal Litteraire* begun at the Hague 1715, and that of Verdun, and the *Memoires Litteraires de la Grande Bretagne* by M. de la Roche; the *Bibliothèque Angloise*, and *Journal Britannique*, which are confined to English books alone. The Italian Journals are, that of abbot Nazari, which lasted from 1668 to 1681, and was printed at Rome. That of Venice began in 1671, and ended at the same time with the other: the authors were Peter Moretti, and Francis Miletto. The *Journal of Parma*, by Roberti and Father Bacchini, was dropped in 1690, and resumed again in 1692. The *Journal of Ferrara*, by the abbé de la Torre, began and ended in 1691. *La Galerio di Minerva*, begun in 1696, is the work of a Society of men of letters. Seignior Apostolo Zeno, Secretary to that Society, began another Journal in 1710, under the protection of the Grand Duke: it is printed at Venice, and several persons of distinction have a hand in it.

The *Fatti Eruditi della Bibliotheca Volante*, were

published at Parma. There has appeared since, in Italy, the *Giornale dei Letterati*.

The principal among the Latin Journals, is that of Leipzig, under the title of *Acta Eruditorum*, begun in 1682: P. P. Manzani began another at Parma. The *Nova Literaria Maris Balthici* lasted from 1698 to 1708. The *Nova Literaria Germaniæ*, collected at Hamburgh, began in 1703. The *Acta Literaria ex Manuscriptis*, and the *Bibliotheca curiosa*, begun in 1705, and ended in 1707, are the work of Struvius. Mess. Kuiter and Sike, in 1697, began a *Bibliotheca Novorum Librorum*, and continued it for two years. Since that time, there have been many Latin Journals; such, besides others, is the *Commentarii de Rebus in Scientia Naturali et Medicina gestis*, by M. Ludwig. The Swiss Journal, called *Nova Literaria Helvetiæ*, was begun in 1702, by M. Scheuchzer; and the *Acta Medica Hafnensia*, published by T. Bartholin, make five volumes from the year 1671 to 1679. There are two Low-Dutch Journals; the one under the title of *Boeckzal van Europe*; it was begun at Rotterdam in 1692, by Peter Rabbus; and continued from 1702 to 1708, by Sewel and Gavern: the other was done by a physician, called Ruiter, who began it in 1710. The German Journals of best note are, the *Monathlichen Unterredungen*, which continued from 1689 to 1698. The *Bibliotheca Curiosa*, began in 1704, and ended in 1707, both by M. Tenzel. The *Magazin d'Hambourg*, begun in 1748: the *Physicalische Belustigungen*, or *Philosophical Amusements*, begun at Berlin in 1751. The *Journal of Hanover* began in 1700, and continued for two years by M. Eccard, under the direction of M. Leibnitz, and afterwards carried on by others. The *Theological Journal*, published by M. Loescher, under the title of *Altes und Neues*, that is, Old and New. A third at Leipzig and Francfort, the authors Mess. Walterck, Krause, and Groschuffius; and a fourth at Hall, by M. Turk.

The English Journals are, The *History of the Works of the Learned*, begun at London in 1699. *Censura Temporum*, in 1708. About the same time there appeared two new ones, the one under the title of *Memoirs of Literature*, containing little more than an English translation of some articles in the foreign Journals, by M. de la Roche; the other a collection of loose tracts, entitled, *Bibliotheca curiosa*, or a Miscellany. These, however, with some others, are now no more, but are succeeded by the *Annual Register*, which began in 1758; the *New Annual Register*, begun in 1780; the *Monthly Review*, which began in the year 1749, and gives a character of all English literary publications, with the most considerable of the foreign ones: the *Critical Review*, which began in 1756, and is nearly on the same plan: as also the *London Review*, by Dr. Kenrick, from 1775 to 1780; *Maty's Review*, from Feb. 1782 to Aug. 1786; the *English Review* begun in Jan. 1783; and the *Analytical Review*, begun in May 1788, and still continues with much reputation. Besides these, we have several monthly pamphlets, called *Magazines*, which, together with a chronological Series of occurrences, contain letters from correspondents, communicating extraordinary discoveries in nature and art, with controversial pieces on all subjects. Of these, the principal are those called, the *Gentleman's Magazine*, which



which began with the year 1731; the London Magazine, which began a few months after, and has lately been discontinued; the Universal Magazine, which is nearly of as old a date.

IRIS, another name for the RAINBOW; which see.

IRIS also denotes the striped variegated circle round the pupil of the eye, formed of a duplicature of the uvea.

In different subjects, the Iris is of several very different colours; whence the eye is called grey, or black &c. In its middle is a perforation, through which appears a small black speck, called the sight, pupil, or apple of the eye, round which the Iris forms a ring.

IRIS is also applied to those changeable colours, which sometimes appear in the glasses of telescopes, microscopes, &c; so called from their similitude to a rainbow.

The same appellation is also given to that coloured spectrum, which a triangular prismatic glass will project on a wall, when placed at a proper angle in the sun-beams.

IRIS *Marina*, the *Sea-Rainbow*. This elegant appearance is generally seen after a violent storm, in which the sea water has been in vast emotions. The celestial rainbow however has great advantage over the marine one, in the brightness and variety of the colours, and in their distinctness one from the other; for in the sea-rainbow, there are scarce any other colours than a dusky yellow on the part next the sun, and a pale green on the opposite side. The other colours are not so bright or so distinct as to be well determined; but the sea-rainbows are more frequent and more numerous than the others: it is not uncommon to see 20 or 30 of them at a time at noon-day.

IRRATIONAL Numbers, or *Quantities*, are the same as surds, or such roots as cannot be accurately extracted, being incommensurable to unity. See SURDS.

IRREDUCIBLE Case, in Algebra, is used for that case of cubic equations where the root, according to Cardan's rule, appears under an impossible or imaginary form, and yet is real. Thus, in the equation  $x^3 - 15x = 4$ , the root, according to Cardan's rule, will be  $x = \sqrt[3]{2 + \sqrt{-121}} + \sqrt[3]{2 - \sqrt{-121}}$ , which is in the form of an impossible expression, and yet it is equal to the quantity 4:

$$\text{for } \sqrt[3]{2 + \sqrt{-121}} = 2 + \sqrt{-1},$$

$$\text{and } \sqrt[3]{2 - \sqrt{-121}} = 2 - \sqrt{-1},$$

therefore their sum is  $x = 4$ .

The other two roots of the equation are also real.

Algebraists, for almost three centuries, have in vain endeavoured to resolve this case, and to bring it under a real form; and the problem is not less celebrated among them, than the squaring of the circle is among geometers.

It is to be observed, that, as in some other cases of cubic equations, the value of the root, though rational, is found under an irrational or surd form; because the root in this case is compounded of two equal surds with contrary signs, which destroy each other; as if  $x = 2 + \sqrt{3} + 2 - \sqrt{3}$ , then  $x = 4$ . In like manner, in the Irreducible case, where the root is rational, there are two equal imaginary quantities, with contrary signs, joined to real quantities; so that the ima-

inary quantities destroy each other; as in the case above of the root of the equation  $x^3 - 15x = 4$ , which was found to be  $2 + \sqrt{-1} + 2 - \sqrt{-1} = 4$ .

It is remarkable that this case always happens, viz one root, by Cardan's rule, in an impossible form, whenever the equation has three real roots, and no impossible ones, but at no time else.

If we were possessed of a general rule for accurately extracting the cube root of a binomial radical quantity, it is evident we might resolve the Irreducible case generally, which consists of two of such cubic binomial roots. But the labours of the algebraists, from Cardan's down to the present time, have not been able to remove this difficulty. Dr. Wallis thought that he had discovered such a rule; but, like most others, it is merely tentative, and can only succeed in certain particular circumstances.

Mr. Maseres, curitor baron of the exchequer, has lately deduced, by a long train of algebraical reasoning, from Newton's celebrated binomial theorem, an infinite series, which will resolve this case, without any mention of either impossible or negative quantities. And I have also discovered several other series which will do the same thing, in all cases whatever; both inserted in the Phil. Trans. See Cardan's Algebra; the articles Algebra, Cubic Equations; Wallis's Algebra, chap. 48; De Moivre in the Appendix to Sanderfon's Algebra, p. 744; Philos. Trans. vol. 68, part 1, art. 42, and vol. 70, p. 387.

IRREGULAR, something that deviates from the common forms or rules. Thus, we say an Irregular fortification, an Irregular building, &c.

IRREGULAR Figure, in Geometry, whether plane or solid, is that whose sides, as well as angles, are not all equal and similar among themselves.

IRREGULARITIES in the Moon's motion. See MOON.

ISAGONE, in Geometry, is sometimes used for a figure consisting of equal angles.

ISLAND, or ISLE, a tract of dry land encompassed by water; whether by the sea, a river, or lake, &c. In which sense Island stands contradistinguished from Continent, or terra firma; like Great Britain, Ireland, Jersey, Sicily, Minorca, &c.

Some naturalists imagine that Islands were formed at the deluge: others think they have been rent and separated from the continent by violent storms, inundations, and earthquakes; while others are thrown up by volcanoes, or otherwise grow or emerge from the sea.

Varenus thinks most of these opinions true in some instances, and believes that there have been Islands produced each of these ways. St. Helena, Ascension, and other steep rocky Islands, he supposes have become so, by the sea's overflowing their neighbouring champaigns. By the heaping up huge quantities of sand &c, he thinks the Islands of Zealand, Japan, &c, were formed: Sumatra and Ceylon, and most of the East India Islands, he rather thinks were rent off from the main land, as England probably was from France. It is also certain that some have emerged from the bottom of the sea; as Santorini formerly, and three other Isles near it lately; the last in 1707, which rose from the bottom of the sea, after an earthquake: the ancients had a tradition that Delos rose from the bottom.



bottom of the sea; and Seneca observes that the Island Therasia rose out of the Ægean sea in his time, of which the mariners were eye witnesses; as they have been within these 10 years, in the sea between Norway and Iceland, where an Island has just emerged. The late circumnavigators too have made it probable, that many of the South-sea Islands have had their foundations, of coral rock, gradually increasing, and growing out of the sea.

ISLES, or rather Ailes, in Architecture, the sides or wings of a building.

ISOCHRONAL, or ISOCHRONOUS, is applied to such vibrations of a pendulum as are performed in equal times. Of which kind are all the vibrations of the same pendulum in a cycloidal curve, and in a circle nearly, whether the arcs it describes be longer or shorter; for when it describes a shorter arc, it moves so much the slower; and when a long one, proportionably faster.

ISOCHRONAL Line, is that in which a heavy body is supposed to descend with a uniform velocity, or without any acceleration.

Leibnitz, in the Act. Erud. Lips. for April 1689, has a discourse on the Linea Isochronea, in which he shews, that a heavy body, with the velocity acquired by its descent from any height, may descend from the same point by an infinite number of Isochronal curves, which are all of the same species, differing from one another only in the magnitude of their parameters (such as are all the quadratocubical paraboloids), and consequently similar to one another. He shews also, how to find a line, in which a heavy body descending, shall recede uniformly from a given point, or approach uniformly to it.

ISOMERIA, in Algebra, a term of Vieta, denoting the freeing an equation from fractions; which is done by reducing all the fractions to one common denominator, and then multiplying each member of the equation by that common denominator, that is rejecting it out of them all.

ISOPERIMETRICAL Figures, are such as have equal perimeters, or circumferences.

It is demonstrated in geometry, that among Isoperimetrical figures, that is always the greatest which contains the most sides or angles. From whence it follows, that the circle is the most capacious of all figures which have the same perimeter with it.

That of two Isoperimetrical triangles, which have the same base, and one of them two sides equal, and the other unequal; that is the greater whose sides are equal.

That of Isoperimetrical figures, whose sides are equal in number, that is the greatest which is equilateral, and equiangular. And hence arises the solution of that popular problem, To make the hedging or walling, which will fence in a certain given quantity of land, also to fence in any other greater quantity of the same. For, let  $x$  be one side of a rectangle that will contain

the quantity  $aa$  of acres; then will  $\frac{aa}{x}$  be its other side,

and double their sum, viz,  $2x + \frac{2aa}{x}$ , will be the

perimeter of the rectangle: let also  $bb$  be any greater number of acres, in the form of a square, then is  $b$  one side of it, and  $4b$  its perimeter, which must be equal to that of the rectangle; and hence the equation  $2x + \frac{2aa}{x} = 4b$ , or  $x^2 + a^2 = 2bx$ , in which qua-

dratic equation the two roots are  $x = b \pm \sqrt{b^2 - a^2}$ , which are the lengths of the two dimensions of the rectangle, viz, whose area  $b^2$  is in any proportion less than the square  $a^2$ , of the same perimeter. As, for example, if one side of a square be 10, and one side of a rectangle be 19, but the other only 1; such square and parallelogram will be Isoperimetrical, viz, each perimeter 40; yet the area of the square is 100, and of the parallelogram only 19.

Isoperimetrical lines and figures have greatly engaged the attention of mathematicians at all times. The 5th book of Pappus's Collections is chiefly upon this subject; where a great variety of curious and important properties are demonstrated, both of planes and solids, some of which were then old in his time, and many new ones of his own. Indeed it seems he has here brought together into this book all the properties relating to Isoperimetrical figures then known, and their different degrees of capacity.

The analysis of the general problem concerning figures that, among all those of the same perimeter, produce maxima and minima, was given by Mr. James Bernoulli, from computations that involve 2d and 3d fluxions. And several enquiries of this nature have been since prosecuted in like manner, but not always with equal success. Mr. Maclaurin, to vindicate the doctrine of fluxions from the imputation of uncertainty, or obscurity, has illustrated this subject, which is considered as one of the most abstruse parts of this doctrine, by giving the resolution and composition of these problems by first fluxions only; and in a manner that suggests a synthetic demonstration, serving to verify the solution. See Maclaurin's Fluxions, p. 486; Analysis Magni problematis Isoperimetrici Act. Erud. Lips. 1701, p. 213; Mem. Acad. Scienc. 1705, 1706, 1718; and the works of John Bernoulli, tom. 1, p. 202, 208, 424, and tom. 2, p. 235; where is contained what he and his brother James published on this problem. Mr. John Bernoulli, in his first paper, considered only two small successive sides of the curve; whereas the true method of resolving this problem in general, requires the considering three such small sides, as may be perceived by examining the two solutions.


M. Euler has also published, on this subject, many profound researches, in the Petersburg commentaries; and there was printed at Lausanne, in 1744, a pretty large work upon it, intitled, Methodus inveniendi lineas curvas, maximi minimive proprietate gaudentes: sive Solutio problematis Isoperimetrici in latissimo sensu accepti.

M. Cramer too, in the Berlin Memoirs for 1752, has given a paper, in which he proposes to demonstrate in general, what can be demonstrated only of regular figures in the elements of geometry, viz, that the circle is the greatest of all Isoperimetrical figures, regular or irregular.



On this head, see also Simpson's *Traëts*, p. 98 ; and the *Philos. Transf.* vol. 49 and 50.

**ISOSCELES Triangle**, is a triangle that has two sides equal. In the 5th prop. of Euclid's 1st book, which prop. is usually called the *Pons Asinorum*, or *Asses bridge*, it is demonstrated, that the angles,  $a$  and  $b$ , at the base of the Isosceles triangle, are equal to each other ; and that if the equal sides be produced, the two angles,  $c$  and  $d$ , below the base, will also be equal. It is also inferred, that every equilateral triangle is also equiangular.



Other properties of this figure are, that the perpendicular AP, from the vertex to the base, bisects the base, the vertical angle, and also the whole triangle. And that if the vertical angles of two Isosceles triangles be equal, the two triangles will be equiangular.

**ISTHMUS**, in Geography, a narrow neck or slip of land, that joins two other large tracts together, and separating two seas, or two parts of the same sea.

The most remarkable Isthmuses are, that of Panama or Straits of Darien, joining north and south America ; that of Suez, which connects Asia and Africa ; that of Corinth, or Peloponnesus, in the Morea ; that of Crim Tartary, otherwise called Taurica Chersonesus ; that of the peninsula Romania and Erisso, or the Isthmus of the Thracian Chersonesus, 12 furlongs broad, and which Xerxes undertook to cut through. The Ancients had several designs of cutting the Isthmus of Corinth, which is a rocky hillock, about 10 miles over ; but without effect, the invention of sluices being not then known. There have also been attempts for cutting the Isthmus of Suez, to make a communication between the Mediterranean and the Red-sea.

**JUDICIAL**, or **JUDICIARY Astrology** ; that relating to the forming of judgments, and making prognostications. See **ASTROLOGY**.

**JULIAN Calendar**, is that depending on, and connected with the Julian year and account of time ; so called from Julius Cæsar, by whom it was established. See **CALENDAR**.

**JULIAN Epoch**, is that of the institution of the Julian reformation of the calendar, which began the 46th year before Christ.

**JULIAN Period**, is a cycle of 7980 consecutive years, invented by Julius Scaliger, from whom it was named ; though some say his name was Joseph Scaliger, and that it was called the Julian Period, because he made use of Julian years. This period is formed by multiplying continually together the three following cycles, viz, that of the sun of 28 years, that of the moon of 19 years, and that of the indiction of 15 years ; so that this epoch, although but artificial or feigned, is yet of good use ; in that every year within the period is distinguishable by a certain peculiar character ; for the year of the sun, moon, and indiction, will not be the same again till the whole 7980 years have revolved. Scaliger fixed the beginning of this period 764 years before the creation, or rather the period naturally reduces to that year, taking the numbers of the three given cycles as he then found them ; and accounting

3950 years from the creation to the birth of Christ, this makes the 1st year of the Christian era answer to the 4714th year of the Julian period ; therefore, to find the year of this period, answering to any proposed year of Christ, to the constant number 4713, add the given year of Christ, and the sum will be the year of the Julian period : thus, to 4713 adding 1791, the sum 6504 is the year of this period for the year of Christ 1791. Hence the first revolution of the Julian period will not be completed till the year of Christ 3267, after which a new revolution of this period will commence.

But the year of this period may be found for any time, from the numbers of the three cycles that compose it, without making use of the given year of Christ, thus : multiply the

numbers  $\left\{ \begin{array}{l} 4845 \\ 4200 \\ 6916 \end{array} \right\}$  respectively by  $\left\{ \begin{array}{l} \text{sun,} \\ \text{moon,} \\ \text{indiction ;} \end{array} \right\}$  the year of the

then add the three products together, and divide the sum by 7980, so shall the remainder after division be the year of the Julian period corresponding to the given years of the other three cycles. Thus, for the year 1791, the years of the solar, lunar, and indiction cycles, are 8, 6, and 9 ; therefore multiplying by these, &c, according to the rule, thus

4845	4200	6916
8	6	9
<hr/>	<hr/>	<hr/>
38760	25200	62244
	38760	
	62244	
	<hr/>	
	7980 ) 126204 ( 15	
	7980	
	<hr/>	
	46404	
	39900	
	<hr/>	

remains 6504 the year of the Jul. period.

**JULIAN Year**, is the old account of the year, established by Julius Cæsar, and consisted of  $365\frac{1}{4}$  days. This year continued in use in all Europe till it was superseded in most parts by the new or Gregorian account, in the year 1582. In England however it continued to be used till the year 1752, when it was abolished by act of parliament, and eleven days added to the account, to bring it up to the new style. In Russia, the old or Julian year and style are still in use.

**JULY**, the 7th month of the year, consisting of 31 days ; about the 21st of which the sun usually enters the sign  $\Omega$  leo. This month was so named by Mark Antony, from Julius Cæsar, who was born in this month.

**JUPITER**,  $\mu$ , one of the superior planets, remarkable for its brightness, being the brightest of all, except sometimes the planet Venus, and is much the largest of all the planets.

Jupiter is situated between Mars and Saturn, being the 5th in order of the primary planets from the sun. His diameter is more than 10 times the diameter of the earth, and therefore his magnitude more than 1000 times.



times. His annual revolution about the sun, is performed in 11 years 314 days 12 hours 20 minutes 9 seconds, going at the rate of more than 25 thousand miles per hour; and he revolves about his own axis in the short space of 9 hours 56 minutes, by which his equatorial parts are carried round at the amazing rate of 26 thousand miles per hour, which is about 25 times faster than the like parts of our earth revolve.

Jupiter is surrounded by faint substances, called zones or belts, in which so many changes appear, that they are generally ascribed to clouds: for some of them have been first interrupted and broken, and then have vanished entirely. They have sometimes been observed of different breadths, and afterwards have all become nearly of the same breadth. Large spots have been seen in these belts; and when a belt vanishes, the contiguous spots disappear with it. The broken ends of some belts have often been observed to revolve in the same time with the spots: only those nearer the equator in somewhat less time than those nearer the poles; perhaps on account of the sun's greater heat near the equator, which is parallel to the belts and course of the spots. Several large spots, which appear round at one time, grow oblong by degrees, and then divide into two or three round spots. The periodical time of the spots near the equator is 9 hours 50 minutes, but of those near the poles 9 hours 56 minutes. See Dr. Smith's Optics, § 1105 and 1109.

The axis of Jupiter is so nearly perpendicular to his orbit, that he has no sensible change of seasons; which is a great advantage, and wisely ordered by the Author of Nature. For, if the axis of this planet were inclined any considerable number of degrees, just so many degrees round each pole would in their turn be almost 6 years together in darkness. And, as each degree of a great circle on Jupiter contains about 706 miles, it is easy to judge what vast tracts of land would be rendered uninhabitable by any considerable inclination of his axis.

The difference between the equatorial and polar diameters of Jupiter, is upwards of 6000 miles; the former being to the latter as 13 to 12: so that his poles are more than 3000 miles nearer his centre than the equator is. This happens from his quick motion round his axis; for the fluids, together with the light particles, which they can carry or wash away with them, recede from the poles which are at rest, towards the equator where the motion is quickest, until there be a sufficient number accumulated to make up the deficiency of gravity lost by the centrifugal force, which always arises from a quick motion round an axis: and when the deficiency of weight or gravity of the particles is made up by a sufficient accumulation, there is then an equilibrium, and the equatorial parts rise no higher.

Jupiter's orbit is  $1^{\circ} 20'$  inclined to the ecliptic. The place of his aphelion  $9^{\circ} 10'$  of  $\text{♈}$ , the place of his ascending node  $7^{\circ} 29'$  of  $\text{♊}$ , and that of his south or descending node  $7^{\circ} 29'$  of  $\text{♋}$ . The eccentricity of his orbit is  $\frac{1}{20}$  of his mean distance from the sun.

The sun appears to Jupiter but the 48th part so large as to us; and his light and heat are in the same

small proportion, but compensated by the quick returns of them, and by 4 moons, some of them larger than our earth, which revolve about him; so that there is scarce any part of this huge planet but what is, during the whole night, enlightened by one or more of these moons, except his poles, whence only the farthest moons can be seen, and where their light is not wanted, because the sun constantly circulates in or near the horizon, and is very probably kept in view of both poles by the refraction of Jupiter's atmosphere, which, if it be like ours, has certainly refractive power enough for that purpose. This planet seen from its nearest moon, appears 1000 times as large as our moon does to us; increasing and waning in all her monthly shapes, every  $42\frac{1}{2}$  hours. The periods, distances, in semidiameters of Jupiter, and angles of the orbits of these moons, seen from the earth are as follow:

No.	Periods round Jupiter.	Distances.	Angles of orbits.
1	$1^d 18^h 36^m$	$5\frac{2}{3}$	$3' 55''$
2	3 13 15	9	6 14
3	7 3 59	$14\frac{1}{3}$	9 58
4	16 18 30	$25\frac{1}{3}$	17 30

The three nearest moons of Jupiter fall into his shadow, and are eclipsed in every revolution: but the orbit of the 4th satellite is so much inclined, that it passeth by its opposition to Jupiter, without falling into his shadow, two years in every fix. By these eclipses, astronomers have not only discovered that the sun's light takes up 8 minutes of time in coming to us; but have also by them determined the longitudes of places on this earth, with greater certainty and facility, than by any other method yet known. The outermost of these satellites will appear nearly as large as the moon does to us. See M. De la Place's Theory of Jupiter's Satellites, in the Memoires de l'Acad. and in the Connoissance des Temps for 1792, pa. 273.

Though there be 4 primary planets below Jupiter, yet an eye placed on his surface would never perceive any of them; unless perhaps as spots passing over the sun's disc, when they happen to come between the eye and the sun.—The parallax of the sun, viewed from Jupiter, will scarce be sensible, being not much above 20 seconds; and the sun's apparent diameter in Jupiter, but about 6 minutes.—Dr. Gregory adds, that an astronomer in Jupiter would easily distinguish two kinds of planets, four nearer him, viz his satellites, and two more remote, viz the sun and Saturn: the former however will fall vastly short of the sun in brightness, notwithstanding the great disproportion in the distances and apparent magnitude.

JURIN (Dr. JAMES), a very distinguished person in several walks of literature, particularly medicine, mathematics, and philosophy, which he cultivated with equal success. He was secretary of the Royal Society in London, as well as president of the College of Physicians there, at the time of his death, which happened March 22, 1750.

Doctor Jurin was author of several ingenious compositions; particularly "an Essay upon Distinct and In-



distinct Vision ;" printed at the end of the 2d volume of Dr. Smith's System of Optics ; also several controversial papers ; against Michellotti, upon the momentum of running waters ; against Robins, upon distinct vision ; and against the partisans of Leibnitz, upon the forces of moving bodies ; &c. His papers inserted in the Philos. Transf. are the following :

1. On the Suspension of Water in Capillary Tubes : vol. 30, p. 739.
2. Observations on the Motion of Running Water : p. 748.
3. On an old Roman Inscription : p. 813.
4. A Discourse on the Power of the Heart : p. 863 and 929.
5. On the Specific Gravity of Human Blood : p. 1000.
6. Defence of his Doctrine of the Power of the Heart against the Objections of Dr. Keill : p. 1039.

7. On the Action of Glass-Tubes upon Water and Quicksilver : p. 1083.

8. On the Specific Gravity of Solids when weighed in Water : vol. 31, p. 223.

9. On the Motion of Running Water, against Michellotti : vol. 32, p. 179.

10. Remarkable Instance of the Small-pox : vol. 32, p. 191.

11. Inoculated and Natural Small-pox compared : vol. 32, p. 213.

12. On Meteorological Diaries : vol. 32, p. 422.

13. On the Measure and Motion of Running Water : vol. 41, p. 5 and 65.

14. Meteorological Observations in Charles Town : vol. 42, p. 491.

15. On the Action of Springs : vol. 43, p. 46.

16. On the Force of Bodies in Motion : p. 423.

17. Dynamic Principles, or Meteorological Principles of Mechanics : vol. 66, p. 103.

END OF VOLUME I.



DIALS.




Fig. 2




Fig. 4





Fig. 5



Halley's DIVING BELL.

Triewald's.

Fig. 6





Fig. 7



Profile of a FORTIFICATION.










Fig. 8















ELECTRICAL MACHINES.

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 7




Fig. 8





Fig. 6









General View of FORTIFICATION.









FOUNTAINS.



Fig. 1




Fig. 2



Fig. 3



Fig. 4




Fig. 5




Fig. 6




Fig. 7

Fountain Pen



Fig. 8








Artificial Globe




Fig. 1


Fig. 2



Profile of the Body of the Place and the Ravelin with Revetement.



Profile of the Body of the Place and Ravelin with demi Revetement.



HYDROMANTIC




Fig. 3

Fig. 4



Hydrometer




Fig. 5




Fig. 6

Smoke JACK




Fig. 8

Kitchen or Fly JACK




Fig. 7



























